Title: Nutritional status and fecundity are synchronised by muscular exopheresis

Authors: Michał Turek[1*], Małgorzata Piechota[2†], Katarzyna Banasiak[2†], Nilesh

Shanmugam[2†], Matylda Macias[3], Małgorzata Alicja Śliwińska[4], Marta Niklewicz[2], Konrad

Kowalski[2], Natalia Nowak[4], Agnieszka Chacińska[1], Wojciech Pokrzywa[2*]

Affiliations:

[1]*Laboratory of Mitochondrial Biogenesis, ReMedy International Research Agenda Unit,*

University of Warsaw, Warsaw, Poland.

[2]*Laboratory of Protein Metabolism in Development and Aging, International Institute of*

Molecular and Cell Biology, Warsaw, Poland.

[3]*Core Facility, International Institute of Molecular and Cell Biology in Warsaw, Poland*

[4]*Laboratory of Imaging Tissue Structure and Function, Nencki Institute of Experimental*

Biology Polish Academy of Sciences, Warsaw, Poland

* Co-corresponding authors

[†] These authors contributed equally to this work

Correspondence should be directed to:

WP: wpokrzywa@iimcb.gov.pl

MT: m.turek@ibb.waw.pl

Abstract

Organismal functionality and reproduction depend on metabolic rewiring and balanced energy resources. However, the crosstalk between organismal homeostasis and fecundity, and the associated paracrine signaling mechanisms are still poorly understood. Using the *Caenorhabditis elegans* we discovered that large extracellular vesicles termed exophers, attributed in neurons and cardiomyocytes to the removal of damaged subcellular components, are released by body wall muscles to support embryonic growth. We found that exopher formation (exopheresis) is a non-cell autonomous process regulated by egg formation in the uterus. Our data suggest that exophers serve as transporters for muscle-generated yolk proteins used for nourishing and improving the growth rate of the next generation. We propose that the primary role of muscular exopheresis is to stimulate the reproductive capacity, thereby influencing the adaptation of worm populations to the current environmental conditions.

Introduction

The proper cellular function relies on the removal of unwanted contents by proteolysis or degradation mainly via the ubiquitin-proteasome system (UPS) and autophagy (Dikic, 2017). Recently a complementary mechanism was described in *Caenorhabditis elegans*. Under proteotoxic stress, worms' neurons can remove protein aggregates and damaged mitochondria via large membrane-surrounded vesicles called exophers. This process is beneficial for neurons functionality as cells that generate exophers perform better than those devoid of this mechanism (Melentijevic *et al*, 2017). However, not only neurons can remove cellular content via exophers. Murine cardiomyocytes can eject the subcellular content along with the mitochondria through exopher-like structures that are ultimately uptaken and eliminated by cardiac macrophages (Nicolás-Ávila *et al*, 2020). This extrusion phenomenon constitutes a

significant but still poorly explored metabolic waste management pathway. Here, we report

that *C. elegans* body wall muscle cells (BWM) can produce exophers in a sex-specific

manner. Furthermore, the generation of muscular exophers depends on developing embryos

in worms' uterus and positively correlates with the number of retained eggs. Finally, we

show that exophers serve as transporters for muscle-generated yolk proteins, which support

offspring development. We identified a new role for exopheresis, which goes beyond the

removal of proteotoxic cellular components, and has a transgenerational effect on the animal

population.

Results and Discussion

C. elegans **muscles remove cellular content via exophers**

Using worms expressing fluorescent reporters in the body wall muscle cells, we identified

muscle-derived vesicles resembling exophers. These reporters allow simultaneous tracking of

mitochondria, which can be extruded in neuronal and cardiac exophers (Melentijevic *et al.*,

2017) and proteasomes, a key component of the proteostasis network. Similar to *C. elegans*

neurons, muscles generate a small fraction (approximately 12%) of the vesicles that contain

the mitochondria, but all of those we examined were filled with proteasomal subunits that are

components of the 19S and 20S proteasome (**Figure 1A**, **S1A**). The size of the described

vesicles extruded by BWM also corresponds to the neural exophers (Melentijevic *et al.*,

2017). However, those of muscular origin are generally more abundant and prominent (with

diameters ranging from 2 to 15 µm) (**Figure 1B**). As with neural, exopheresis in muscles is

not limited to a particular reporter or transgenic line (**Figures 1A-C, 2C, 4H**). Moreover, our

electron microscopy (EM) studies visualized the membrane surrounded exophers adjacent to

the BWM. Vesicles analyzed by EM show similar compactness, with the characteristic

presence of mitochondria (**Figure 1C, S1C-D**). Most of them showed marked morphology

changes, characterized by increased area and disturbed cristae organization as in cardiac

exophers. However, we noted the presence of probably not compromised mitochondria in

BWM-derived structures, suggesting that they are not only used to transport defective

organelles. To finally confirm that the identified vesicles are the counterpart of exophers, we

tracked their formation in response to RNAi knockdown of genes that regulate neuronal

exopheresis (Melentijevic et al., 2017). Indeed, the appearance of muscle-derived exophers

was also affected by depletion of EMB-8 cytochrome reductase and actin-binding protein

POD-1, both linked to the generation of cell polarity (**Figure S1B**).

Exophers are formed in the muscle cell and expelled outside via a pinching off mechanism

(**Figure 1D**), and the majority are generated by adult hermaphrodite mid-body muscles

(**Figure 1E**). Some exophers that bud off remain connected with the extruding BWM via a

thin but elastic tube that permits the continued transfer of a large amount of cellular material

into the extruded vesicle (**Figure 1G, Video S1**), similar to neuronal exophers (Melentijevic

et al., 2017). We also observed that large exophers could fragment into smaller vesicles

(**Figure 1F**). Proteostasis impairment significantly increases neuronal exopher output (Hualin

et al, 2019; Melentijevic *et al.*, 2017). In contrast, the number of muscular exophers did not

change in response to depletion of the central proteostasis transcription factor HSF-1 (via *hsf-*

1 RNAi) or heat stress, and the number of exophers increased slightly under conditions of

oxidative stress (**Figure 1H-I**). These observations suggest that proteostasis regulation might

not be the core function of muscle exopheresis.

Muscular exopheresis is a sex-specific process regulated in a non-cell autonomous

manner

Next, we assessed the number of exophers at different time points of the *C.*

elegans hermaphrodite life cycle. Reminiscent of neuronal exophers, muscular exophers are

not produced during the larval stages, and their maximum level is reached around the second

and third days of hermaphrodite adulthood (**Figure 2A**). Because this time point coincides with the worm's maximum reproductive rate, we wondered if reproduction could influence exopher formation. To examine this possibility, we followed exopheresis in males. For the first three days of adulthood, males did not produce any exophers (**Figure 2A**). This finding suggests that germ cell maturation in the reproductive system of hermaphrodite worms, the process of oocyte fertilization, or embryonic development might regulate muscle exopheresis. To test these hypotheses, we took advantage of a thermosensitive *C. elegans fem-1* mutant strain that does not produce viable sperm at the restrictive temperature of 25 °C. At the permissive temperature of 15 °C, some animals can reproduce like wild type hermaphrodites, whereas the rest of the population is sterile (Nelson *et al*, 1978). The offspring-producing *fem-1* mutant animals grown at 15 °C generated a high number of muscular exophers. Contrarily, animals raised either at 15 °C or 25 °C that were unable to fertilize oocytes did not activate muscular exopheresis (**Figure 2B-C**), indicating that neither the female gonad nor the temperature itself was sufficient to trigger exopher release. However, re-establishing fertility of *fem-1* mutants by mating them to *him*-5 mutant males restored exopher production at the restrictive temperature (**Figure 2B**). Moreover, hermaphrodites sterilized via fluorodeoxyuridine (FUdR) treatment (Hosono, 1978) extruded no exophers or only a few per animal (**Figure 2D**), suggesting that the occurrence of developing embryos could indeed stimulate muscular exopheresis. We also found that FUdR-treated hermaphrodites often contained exopher-like structures in their BWM (**Figure 2E**, middle and right panels). Interestingly, we detected objects resembling non-extruded exophers in the BWM of males (**Figure 2E**, left panel), indicating that males are devoid of mechanisms triggering their expulsion.

The above results suggest that the occurrence of developing embryos could induce muscular exopheresis. Consistently, we observed a positive correlation between the number of

exophers released and the number of eggs present in the worm uterus (**Figure 3A**). To further

explore this link, we depleted the mRNA of genes responsible for various processes

associated with egg-laying. RNAi depletion of the G-protein signalling gene *goa-1* leads to

hyperactive egg-laying behaviour, resulting in the presence of fewer early-stage eggs within

the uterus (Bany *et al*, 2003). As expected, *goa-1* knockdown caused a significant drop in the

level of exopher release. In contrast, the egg-laying defects induced by *egl-1* and *egl-4* RNAi,

which lead to egg retention in the uterus (Hirose *et al*, 2003), increased exopher formation by

muscle cells (**Figure 3B**). In the absence of food, worms halt egg-laying and retain fertilized

eggs in the uterus (Daniels *et al*, 2000; Dong *et al*, 2000); therefore, we anticipated that

worms would generate more exophers when experiencing a food shortage. Indeed, the

accumulation of developing eggs in the uterus caused by the transfer of adult worms to food-

free plates resulted in a significant increase in muscle exopher secretion in contrast to L4

larvae grown under conditions of food shortage (**Figure 3C**).

Next, we tested whether worm embryos could directly induce exopher production. We

exposed young adult worms to extract from developing eggs or L2/L3 larvae of the wild-type

strain (**Figure 3D**). Intriguingly, worms placed in contact with material derived only from

lysed embryos increased exopheresis significantly (**Figure 3E**). This observation suggests

that molecules that diffuse from embryos *in utero* are responsible for exopheresis induction.

Finally, we decided to confirm that even in the face of disturbed proteostasis specifically in

the BWM, a signal associated with the developing embryos in the uterus would be the

primary regulator of muscular exopheresis. To this end, we knocked down the myosin-

directed chaperone UNC-45, which results in severe defects in the sarcomeric proteostasis, as

well as embryonic defects in cytokinesis and polarity determination (Gazda *et al*, 2013;

Kachur *et al*, 2004; Pokrzywa & Hoppe, 2013). We initiated UNC-45 depletion in L4 larvae,

which first leads to a disturbance of proteostasis in BWM, and later, in young gravid adults,

inhibiting embryonic development. Despite the dysfunction of a myosin-chaperone network

in muscle cells, which was indicated by complete paralysis of worms, we observed a

dramatic inhibition of exopher formation (**Figure 3F**), highlighting the predominant role of

maturating eggs in muscular exopheresis.

Exophers transport muscle-synthesized yolk proteins and supports offspring

development

The high number of exophers produced during the *C. elegans* lifespan by BWM leads to

removing a rather significant portion of subcellular components by a single tissue. Thus, we

hypothesized that this process should have substantial effects on worm muscle functionality

and healthspan. To this end, we selected three types of worms from the synchronized

population of gravid adults based on the number of extruded exophers, i.e., few (< 2), many

(> 20), and control (6-8) animals (**Figure 4A**) and analyzed their locomotion. In neurons,

exophers production is correlated with improved cell functionality; however, worms with

intensified exopheresis did not show enhanced mobility (**Figure 4B, S2**). On the contrary,

these animals displayed a reduction in exploratory locomotion state (**Figure 4C-D**). Since the

production of exopher does not seem to have a positive effect on the functionality of the

muscles, we next wondered whether, in connection with the role of embryos in exopheresis,

this process benefited the offspring. Previous work showed that neuronal exopher content

could be transported through the worm body to reach distant scavenger cells (coelomocytes)

(Melentijevic *et al.*, 2017). However, the fate of muscle exophers may be different from

neuronal counterparts, as their processing is not dependent on the apoptotic engulfment

pathway proteins CED-1 and CED-6, the depletion of which did not change the muscle

exopheresis level (**Figure S3A**). Muscle-specific transcriptomic analysis revealed the

presence of significant levels of vitellogenin mRNAs (i.e., *vit-2*, *-5* and *-6*) (Blazie *et al*,

2015). Therefore, we hypothesized that the yolk components from BWM are transported

through exophers to be used as a source of raw materials for maturating eggs. To address this

possibility, we used RNAi knockdown of *vit-1* (a vitellogenin-coding gene) to reduce the

level of the principal intestinal yolk protein (Perez & Lehner, 2019) and investigated whether

it led to an increase in exopher biogenesis as a possible compensatory mechanism. Indeed,

the number of accumulated muscle-released exophers nearly doubled in response to yolk

protein depletion (**Figure 4E**). Moreover, the intensification of exopheresis in the mother

increased the amount of vitellogenin in developing eggs (**Figure 4F**). In contrast, depletion of

the RME-2 yolk receptor and subsequent inhibition of yolk uptake by oocytes led to a drastic

reduction in exopheresis (**Figure 4G**), which cannot be explained by the sterility of worms,

as the embryos' viability is not entirely compromised even in the *rme-2* deletion mutant

(b1008) (more than 20% of embryos are alive) (Grant & Hirsh, 1999). Next, we followed the

localization of endogenous vitellogenin-2 (VIT-2) fused to GFP (CRISPR/Cas9 knock-in

strain) and muscle exopher markers in the hermaphrodite. We detected the presence of

endogenous VIT-2::GFP in the BWM of day-2 adult worms, as well as a significant

accumulation in many muscle exophers (**Figure 4H, S3B**). These results suggest that

exophers can mediate the transport of additional portions of muscle-produced vitellogenin,

which is ultimately deposited in oocytes from the body cavity (Hall *et al*, 1999). Finally, we

followed the growth of the progeny of hermaphrodites exhibiting different levels of

exopheresis (**Figure 4A**). We found that offspring from mothers with a high number of

muscle-released exophers grew faster (**Figure 4I**). This is in line with previous reports

showing that yolk-reach eggs support animals post-embryonic survival and larvae

development (Perez *et al*, 2017; Perez & Lehner, 2019; Van Rompay *et al*, 2015).

Exophers result from the activation of non-canonical mechanisms that allow the neuronal and

muscle cells to release the sub-cellular components together with organelles such as

mitochondria or nano-organelles such as proteasomes. Accordingly, based on their content

and participation of the autophagy machinery in exophers formation, this vesicle can be

considered part of the proteostasis network. Here we show that exophers are not only a

storage compartment of cellular waste, but the muscular exopheresis in *C. elegans* represents

a previously uncharacterized nutrient management program associated with nourishing the

next generation of worms. Activation of exopheresis occurs with the first appearance of

developing eggs in the uterus and intensifies in environmental food depletion situations.

Hermaphrodites utilize a narrow time window between 2-4 days of adulthood to

simultaneously expel probably parts of the dysfunctional cell elements out of the muscles, as

well as material that can be used by the embryos. Likewise, the disturbance of yolk synthesis

increases exopher biosynthesis. Consequently, in mothers with highly active exopheresis, the

volume of yolk content in eggs increased. Our results show that yolk protein produced in

BWM is transferred to exophers and ultimately delivered to oocytes. Therefore, the use of

exophers for the transport of vitellogenin represents an elegant mechanism by which remote

cells can enrich the nourishment of developing embryos. This process leads to the production

of larvae better prepared to thrive in the current environmental conditions. Thus, muscular

exopheresis is likely an adaptive mechanism that affects the dynamics of population growth.

The impact of exopheresis on early reproduction may be particularly crucial for wild worms,

given the significantly shortened life expectancy observed under more natural conditions

(Van Voorhies *et al.*, 2005). In addition, yolk protein can be synthesized in the muscles of

oviparous animals like zebrafish (Zhong *et al*, 2014), and is supplemented from the mother to

the intraovarian embryo in viviparous animals (Iida *et al*, 2019). Hence, it is tempting to

speculate that the role of muscular exopheresis in supporting progeny development could be

evolutionarily conserved. However, the exact mechanisms by which oocyte fertilization and

subsequent embryonic development initiate exopher formation and how this exopheresis is

executed at the molecular level require further studies.

References

Bany IA, Dong MQ, Koelle MR (2003) Genetic and cellular basis for acetylcholine inhibition
of Caenorhabditis elegans egg-laying behavior. *J Neurosci* 23: 8060-8069
Bishop D, Nikić I, Brinkoetter M, Knecht S, Potz S, Kerschensteiner M, Misgeld T (2011)
Near-infrared branding efficiently correlates light and electron microscopy. *Nat Methods* 8:
568-570
Blazie SM, Babb C, Wilky H, Rawls A, Park JG, Mangone M (2015) Comparative RNA-Seq
analysis reveals pervasive tissue-specific alternative polyadenylation in Caenorhabditis
elegans intestine and muscles. *BMC Biol* 13: 4
Brenner S (1974) The genetics of Caenorhabditis elegans. *Genetics* 77: 71-94
Daniels SA, Ailion M, Thomas JH, Sengupta P (2000) egl-4 acts through a transforming
growth factor-beta/SMAD pathway in Caenorhabditis elegans to regulate multiple neuronal
circuits in response to sensory cues. *Genetics* 156: 123-141
Deerinck TJ, Bushong EA, Thor A, Ellisman MH, 2010. *NCMIR methods for 3D EM: A new*
protocol for preparation of biological specimens for serial block face scanning electron
microscopy. National Center for Microscopy and Imaging Research, University of
California San Diego, La Jolla, CA92093.
Dikic I (2017) Proteasomal and Autophagic Degradation Systems. *Annu Rev Biochem* 86:
193-224
Dong MQ, Chase D, Patikoglou GA, Koelle MR (2000) Multiple RGS proteins alter neural G
protein signaling to allow C. elegans to rapidly change behavior when fed. *Genes Dev* 14:
2003-2014
El Mouridi S, Lecroisey C, Tardy P, Mercier M, Leclercq-Blondel A, Zariohi N, Boulin T
(2017) Reliable CRISPR/Cas9 Genome Engineering in. *G3 (Bethesda)* 7: 1429-1437
Gazda L, Pokrzywa W, Hellerschmied D, Lowe T, Forne I, Mueller-Planitz F, Hoppe T,
Clausen T (2013) The myosin chaperone UNC-45 is organized in tandem modules to support
myofilament formation in C. elegans. *Cell* 152: 183-195
Grant B, Hirsh D (1999) Receptor-mediated endocytosis in the Caenorhabditis elegans
oocyte. *Mol Biol Cell* 10: 4311-4326
Hall DH, Winfrey VP, Blaeuer G, Hoffman LH, Furuta T, Rose KL, Hobert O, Greenstein D
(1999) Ultrastructural features of the adult hermaphrodite gonad of Caenorhabditis elegans:
relations between the germ line and soma. *Dev Biol* 212: 101-123
Hanson HH, Reilly JE, Lee R, Janssen WG, Phillips GR (2010) Streamlined embedding of
cell monolayers on gridded glass-bottom imaging dishes for correlative light and electron
microscopy. *Microsc Microanal* 16: 747-754
Hirose T, Nakano Y, Nagamatsu Y, Misumi T, Ohta H, Ohshima Y (2003) Cyclic GMP-
dependent protein kinase EGL-4 controls body size and lifespan in C elegans. *Development*
130: 1089-1099
Hosono R (1978) Sterilization and growth inhibition of Caenorhabditis elegans by 5-
fluorodeoxyuridine. *Exp Gerontol* 13: 369-374
Hualin F, Jilong L, Peng D, Weilin J, Daxiang C, 2019. Metabolic wastes are extracellularly
disposed by excretosomes, nanotubes and exophers in mouse HT22 cells through an
autophagic vesicle clustering mechanism. bioRxiv.
Iida A, Arai HN, Someya Y, Inokuchi M, Onuma TA, Yokoi H, Suzuki T, Hondo E, Sano K
(2019) Mother-to-embryo vitellogenin transport in a viviparous teleost. *Proc Natl Acad Sci U*
S A 116: 22359-22365

Kachur T, Ao W, Berger J, Pilgrim D (2004) Maternal UNC-45 is involved in cytokinesis
and colocalizes with non-muscle myosin in the early Caenorhabditis elegans embryo. *J Cell*
Sci 117: 5313-5321

Kamath RS, Ahringer J (2003) Genome-wide RNAi screening in Caenorhabditis elegans.
Methods 30: 313-321

Melentijevic I, Toth ML, Arnold ML, Guasp RJ, Harinath G, Nguyen KC, Taub D, Parker
JA, Neri C, Gabel CV *et al* (2017) C. elegans neurons jettison protein aggregates and
mitochondria under neurotoxic stress. *Nature* 542: 367-371

Mitchell DH, Stiles JW, Santelli J, Sanadi DR (1979) Synchronous growth and aging of
Caenorhabditis elegans in the presence of fluorodeoxyuridine. *J Gerontol* 34: 28-36

Nelson GA, Lew KK, Ward S (1978) Intersex, a temperature-sensitive mutant of the
nematode Caenorhabditis elegans. *Dev Biol* 66: 386-409

Nicolás-Ávila JA, Lechuga-Vieco AV, Esteban-Martínez L, Sánchez-Díaz M, Díaz-García E,
Santiago DJ, Rubio-Ponce A, Li JL, Balachander A, Quintana JA *et al* (2020) A Network of
Macrophages Supports Mitochondrial Homeostasis in the Heart. *Cell* 183: 94-109.e123

Paul-Gilloteaux P, Heiligenstein X, Belle M, Domart MC, Larijani B, Collinson L, Raposo
G, Salamero J (2017) eC-CLEM: flexible multidimensional registration software for
correlative microscopies. *Nat Methods* 14: 102-103

Perez MF, Francesconi M, Hidalgo-Carcedo C, Lehner B (2017) Maternal age generates
phenotypic variation in Caenorhabditis elegans. *Nature* 552: 106-109

Perez MF, Lehner B (2019) Vitellogenins - Yolk Gene Function and Regulation in. *Front*
Physiol 10: 1067

Pokrzywa W, Hoppe T (2013) Chaperoning myosin assembly in muscle formation and aging.
Worm 2: e25644

Porta-de-la-Riva M, Fontrodona L, Villanueva A, Cerón J (2012) Basic Caenorhabditis
elegans methods: synchronization and observation. *J Vis Exp*: e4019

Praitis V, Casey E, Collar D, Austin J (2001) Creation of low-copy integrated transgenic lines
in Caenorhabditis elegans. *Genetics* 157: 1217-1226

Redemann S, Schloissnig S, Ernst S, Pozniakowsky A, Ayloo S, Hyman AA, Bringmann H
(2011) Codon adaptation-based control of protein expression in C. elegans. *Nat Methods* 8:
250-252

Van Rompay L, Borghgraef C, Beets I, Caers J, Temmerman L (2015) New genetic
regulators question relevance of abundant yolk protein production in C. elegans. *Sci Rep* 5:
16381

Van Voorhies WA, Fuchs J, Thomas S (2005) The longevity of Caenorhabditis elegans in
soil. *Biol Lett* 1: 247-249

Watanabe S, Punge A, Hollopeter G, Willig KI, Hobson RJ, Davis MW, Hell SW, Jorgensen
EM (2011) Protein localization in electron micrographs using fluorescence nanoscopy. *Nat*
Methods 8: 80-84

Zhang Y, Werling U, Edelmann W (2012) SLiCE: a novel bacterial cell extract-based DNA
cloning method. *Nucleic Acids Res* 40: e55

Zhong L, Yuan L, Rao Y, Li Z, Zhang X, Liao T, Xu Y, Dai H (2014) Distribution of
vitellogenin in zebrafish (Danio rerio) tissues for biomarker analysis. *Aquat Toxicol* 149: 1-7

Śliwińska MA, Cały A, Borczyk M, Ziółkowska M, Skonieczna E, Chilimoniuk M, Bernaś
T, Giese KP, Radwanska K (2020) Long-term Memory Upscales Volume of Postsynaptic
Densities in the Process that Requires Autophosphorylation of αCaMKII. *Cereb Cortex* 30:
2573-2585

 Figures

Figure 1. *C. elegans* muscles expel cellular content via exophers. (A) Muscular exophers
contain organelles and large protein complexes. Arrows: white – exopher, blue – muscle cell,
green – mitochondria, red – proteasome foci, yellow – unidentified vesicle. MOM –
mitochondrial outer membrane. (B) BMW actively releases significant amounts of exopher.
The image shows the middle part of the worm's body with muscles marked with dashed lines.
Arrows indicate representative exophers, and the asterisk indicates the position of the vulva.
(C) The ultrastructure of the muscular exopher and its schematic view. Arrows: red –
morphologically changed mitochondria inside the exopher, yellow – normal, elongated
mitochondria inside muscle cell. c – cuticle, m – muscle, e – exopher. (D) Exophers are
formed via a pinching-off mechanism. Arrows: white – exopher, blue – muscle cell, red –
distorted muscle cell membrane during exopher formation. (E) Production of muscular
exophers is not evenly distributed across all muscle cells. The highest number of exophers is
produced by the muscles adjacent to the vulva. n = 46 animals; three biological replicates. (F)
Large exophers may spontaneously disintegrate into smaller exophers. Arrows: white –
exopher, blue – muscle cell. (G) Exophers may remain connected to the sending BWM cells
via thin elastic tubes that allow further transfer of cellular material. Arrows: white – exopher,
blue – muscle cell, red – cellular material transferred to exopher via elastic tube. (H)

Proteostasis disruption by *hsf-1* knockdown does not increase exopher production. n = 60 and

55 animals, two biological replicates. (I) Proteostasis induced via oxidative stress but not via

heat stress increases exopher production. n = 30 animals; three biological replicates.

Scale bars are 10 µm (A, B, F, G) and 1 µm (C). Data are shown as mean ± SEM. ns – not

significant, ** P < 0.01, **** P < 0.0001; (E, H, I) Mann-Whitney test.

Figure 2. Exopher formation is sex-specific and fertility-dependent. (A) The highest number of exophers is produced during the hermaphrodite reproductive period and in aging animals. Males do not produce exophers during the first days of adulthood and begin to generate a small number of exophers later in life. Starting n = 90 hermaphrodites and 150 males; three biological replicates. (B) Feminized hermaphrodites of thermosensitive a *fem-1* mutant strain do not produce exophers regardless of growth temperature. This phenotype can be partially rescued by mating *fem-1* mutants with males. n = 10 - 26 animals; one biological replicate. (C) Representative images of the middle part of the worm body in panel B. (D) Hermaphrodites sterilized via FUdR treatment produce no exophers or only a few per animal. n = 17 and 20 animals; two biological replicates. (E) Males and sterile hermaphrodites (via FUdR treatment) show the formation of spherical structures in the BWM that resemble mature exophers. MOM – mitochondrial outer membrane.

Scale bars are 10 µm. Data are shown as mean ± SEM; ** P < 0.01, **** P < 0.0001; (B) Mann-Whitney test.

Figure 3. Muscular exopheresis is a non-cell autonomous process regulated by *in utero* developing embryos. (A) The number of produced exophers positively correlates with the number of *in utero* embryos. The violet line is a linear regression line, and each orange point represents one animal. All animals were 1 - 3 days old; n = 54 animals; three biological replicates. (B) RNAi knockdown of genes regulating the egg laying rate and their presence in the uterus influences exopher production. n = 50 - 60 animals; two biological replicates. (C) Eggs retention in the uterus caused by starvation during worm's reproductive period increases exopher production. n = 81 - 90 animals; three biological replicates. (D) Schematic representation of the experimental setup for investigating the influence of egg lysate on exopher production. (E) Exposure of young-adult worms to embryo-derived substances increases exopher production. n = 24 - 77 animals; two biological replicates. (F) RNAi knockdown of the essential muscular co-chaperone *unc-45* reduces exopher production. n = 28 and 17 animals; two biological replicates.

Data are shown as mean ± SEM; ns – not significant, * P < 0.05, **** P < 0.0001; (B, C, F) Mann-Whitney test; (E) two-tailed Welch's t-test.

Figure 4. Muscular exopheresis benefits offspring development. (A) Schematic representation of the experimental setup for investigating the influence of overactive exopheresis on F0 and F1 worm generation. (B) Exopher production does not improve muscle functionality. n = 11 - 38; three biological replicates. (C-D) Animals with overactive exopheresis have reduced exploratory behaviour presented as a reduction in straight line distance travelled (C) and by representative worms traces on the plate (D). n = 11 - 34; three biological replicates. (E) RNAi knockdown of the egg yolk precursor protein VIT-1 increases the number of muscular exophers. n = 20 animals; two biological replicates. (F) Embryos from hermaphrodite mothers that produce a high number of exophers contain more egg yolk precursor protein VIT-2. Representative images of embryos with endogenous VIT-2::GFP levels from mothers with different exopheresis activity are shown above the graph. n = 15 and 14 eggs, five and six biological replicates (animals). (G) RNAi knockdown of RME-2 yolk receptor abolish exopher production. n = 110 – 123 animals; three biological replicates. (H) Muscle-produced VIT-2 is released from muscles via exophers. The image shows the midbody of worms expressing the proteasome subunit RPN-5 tagged with RFP in BWM and VIT-2::GFP endogenous expression. Arrows: white – exopher, blue – muscle cell. Dashed lines mark eggs present in the uterus. Scale bar is 10 µm. (I) Offspring of worms with overactive exopheresis develop faster. YA – young adult stage, L4 – last larval stage. n = 317 – 372 animals; two biological replicates.

Data are shown as mean ± SEM. * P < 0.05, ** P < 0.01, **** P < 0.0001; (E, G) Mann-Whitney test; (F) two-tailed Welch's t-test; (I) Fisher's exact test.

Figure S1. Ultrastructure of muscle exophers, the formation of which depends on cell polarity regulating proteins. (A) Small fraction of muscular exophers contain mitochondria. n = 60; six biological replicates. (B) The formation of muscle-derived exophers is affected by

depletion of proteins linked to the generation of cell polarity. n = 80 – 120; *pod-1* – two biological replicates, *emb-8* RNAi – three biological replicates. Data are shown as mean ± SEM. **** P < 0.0001, Mann-Whitney test. (C-D) Exophers visualization using correlative light electron microscopy (CLEM). Exopher's red fluorescence (red arrow). (C'-D') Exopher (red arrow) surrounded by marks (asterisk) after NIRB, autofluorescence of branded frame edges is visible. (C''-D'') CLEM tracking the exopher - LM image, EM image and their overlay. (C'''-D''') Image shows exopher's (coloured in red) ultrastructure. Thick, grey arrows show order of steps done for the final exopher detection under the electron microscope. Scale bars are 1 µm.

Figure S2. Active exophoresis affects worm mobility. Graph shows the track lengths of animals exhibiting different exopheresis activities. n = 11 - 38; three biological replicates. Data are shown as mean ± SEM.

Figure S3. Exophers contain muscle-produced vitellogenin. (A) Muscular exophers formation does not depend on the apoptotic engulfment pathway proteins CED-1 and CED-6. n = 114 – 120; three biological replicates. Data are shown as mean ± SEM. ns – not significant, Mann-Whitney test. (B) Images show the formation of an exopher filled with the proteasome and vitellogenin. Images were captured before and after RPN-5::RFP

photobleaching, confirming that the high signal from endogenous VIT-2::GFP in forming

exophers is not an imaging artefact. The white circle marks the area that was bleached using a

555 nm laser and the position of developing exopher. Dashed lines mark different tissue

borders: m – muscle, i – intestine, g – gonad. Scale bar is 10 µm.

Methods

Data reporting

No statistical methods were used to predetermine the sample size. The experiments were not

randomized. The investigators were not blinded to allocation during experiments and

outcome assessment except for exploratory behaviour experiments.

Worm maintenance and strains

Worms were maintained on Nematode Growing Medium (NGM) plates seeded with

OP50 *Escherichia coli* bacteria at 20 °C unless otherwise stated (Brenner, 1974). A list of all

strains used in the study, together with the information in which experiments they were used

is provided in Supplementary Table 1.

Generation of plasmids

All constructs were cloned using the SLiCE method (Zhang *et al*, 2012) and were sequenced

for verification. Construct for the expression of GFP on the mitochondrial outer membrane in

the worm's body wall muscles was prepared as follows. First, destination vector pMT26

containing pCG150 vector backbone with the *myo-3* promoter and *unc-54* 3'UTR sequences

separated by KspAI restriction site was made. Next, codon-optimized sequences (Redemann

et al, 2011) for GFP, linker containing attB5 sequence, and the sequence of the first 55 amino

acids from TOMM-20 protein (Watanabe *et al*, 2011) (from a plasmid that was synthesized

for this study) were PCR amplified and inserted into linearized pMT26 vector. To generate

construct for expression of RPN-5 or PAS-7 tagged with RFP in worm's body wall muscles,

sequences encoding respective proteins were PCR amplified and inserted into pMT26

linearized destination vector. As a template for the *rpn-5*-coding sequence, we used plasmid

bearing codon-optimized *rpn-5* cDNA with 3 artificial introns, which was synthesized for this

study. The sequence of *pas-7* gene was directly amplified from N2 worms genomic DNA,

and the sequence of RFP (wrmScarlet) was amplified from pSEM89 plasmid (El Mouridi *et*

al, 2017).

Transgenic strains generation

Worms transgenic strains were created by microparticle bombardment using *unc-*

119(ed3) rescue as a selection marker (Praitis *et al*, 2001). After phenotypic confirmation of

successful plasmid insertion, transformants were backcrossed two times against N2 strain to

remove the *unc-119(ed3)* background.

Scoring exophers and fluorescence microscopy

For scoring of exophers, a confocal microscope or a stereomicroscope was used. When using

the confocal microscope, animals were transferred onto 3 % agarose pads prepared in H_2O

and formed on a microscope slide. Next, animals were immobilized on the pad using 6 μL of

PolySciences 0.05 μm polystyrene microspheres or 25 μM tetramisole and covered with a

glass coverslip. Immediately afterward, animals were imaged using an inverted Zeiss

LSM800 laser-scanning confocal microscope with 40x or 63x oil immersion objectives. The

488 nm and 561 nm lasers were used for excitation of GFP and RFP fluorescent proteins,

respectively. Z-stacks, which covered the whole animal, were taken and the number of

exophers released by muscles was counted. Finally, data were normalized to the average

number of exophers present in control animals and compared between conditions.

For scoring exophers with the stereomicroscope, we used Leica M165FC stereomicroscope

equipped with Leica EL6000 lamp and standard Texas Red and GFP filter sets. Age

synchronized, freely moving, day-2 adult animals were directly visualized on culturing NGM

plates and the number of visible exophers released by muscles were counted. Finally, data

were normalized to the average number of exophers present in control animals and compared

between conditions.

Representative pictures of exophers presented in the manuscript were taken with the usage of

an inverted Zeiss 700 laser-scanning confocal microscope equipped with a 40x oil objective.

The 488 nm and 555 nm lasers were used for excitation of GFP and RFP fluorescent proteins,

respectively. To investigate the presence and distribution of exophers, z-stacks were taken

and processed with ZEN software.

Electron microscopy

EM sample preparation

C. elegans day 2 adult worms were fixed with 2% paraformaldehyde (Sigma Aldrich, P6148)

and 1% glutaraldehyde (Sigma Aldrich, EM grade) in 0.2M Hepes pH 7.3 overnight. Next,

the samples were washed three times in 0.2M Hepes pH 7.3, followed by overnight

incubation with 0.1% ruthenium red solution in distilled water. After staining, worms were

mounted with 2% gelatin on gridded 35mm dishes (MatTek 35mm dish, No. 1.5 Gridded

Coverslip 14 mm Glass Diameter, P35G-1.5-14-CGRD) and designated for branding.

Near infra-red branding (NIRB)

Worms were localised on a gridded dish in bright field mode. Next, the exophers' red

fluorescence was imaged using multiphoton Examiner.Z1 LSM 7MP microscope equipped

with LSM NDD detectors and water 20x NA 1.0 objective (Zeiss, Oberkochen, Germany).

For the imaging, pulsed laser at 1020 nm (Coherent Chameleon) and 600-700 nm detection

filter were used. Z-stacks of the samples were acquired using 4 times averaging in the line

mode, pixel size of 138 x 138 nm and 500 nm interval in the Z axis. For each sample a Z-

stack was acquired before and after the branding.

The ROI was marked by NIRB (Bishop *et al*, 2011). The frame contour was bleached below

the place of interest with a pulsed laser at 910 nm (Coherent Chameleon, 2 W at 910 nm),

using 20-40 iterations with 20% of the laser power and pixel dwell time of 6-25 µs. The final

parameters (number of iteration and pixel dwell time) were adjusted separately for each

sample. The branding was repeated until the edges of the frame were highly fluorescent at

910 nm.

EM processing

Branded samples were prepared for electron microscopy according to a published

protocol(Deerinck *et al*, 2010) with minor changes (Śliwińska *et al*, 2020). Briefly, animals

were post-fixed with 1% aqueous solution of osmium tetroxide (Polysciences Europe GmbH

0972B-5) and 1.5% potassium ferrocyanide (Sigma Aldrich, St. Louis, MO, USA, P3289) in

phosphate buffer for 30 min on ice. Next, samples were immersed in 1% aqueous

thiocarbohydrazide (Sigma Aldrich, St. Louis, MO, USA, #88535) for 40 min, post-fixed

with 2% aqueous solution of osmium tetroxide for 60 min (all at room temperature) and

incubated in 1% aqueous uranyl acetate at 4ºC overnight. The next day, samples were

immersed in 0.66% lead aspartate for 60 min at 60ºC, dehydrated with increasing dilutions of

ethanol, infiltrated with Durcupan resin (Sigma Aldrich, St. Louis, MO, USA, #44610),

embedded using BEEM capsules according to a published protocol (Hanson *et al*, 2010) and

cured at 70ºC for 72 h.

The resin blocks were trimmed and cut with an ultramicrotome (ultracut R or EM UC7,

Leica) and ultrathin sections (70 nm thick) were collected on formvar-coated copper grids,

mesh 100 (Agar Scientific, AGS138-1 or EMS FF100-CU-50) or copper slot grids (EMS

FCF 2010-CU-SB-50). For serial block face scanning electron microscopy trimmed pyramids

were cut off with a razor blade, mounted to aluminium pins (Gatan system pins, Micro to

Nano, Netherlands, 10-006003-50) with cyanoacrylate glue, trimmed, block face polished

and grounded with silver paint (Ted Pella, Redding, USA, 16 062-15).

Transmission Electron Microscopy Imaging

Specimen grids were examined with a transmission electron microscope JEM 1400 (JEOL

Co., Tokyo, Japan, 2008), equipped with a 11 megapixel TEM camera MORADA G2

(EMSIS GmbH, Münster, Germany) at Nencki Institute of Experimental Biology PAS or

with a Tecnai T12 BioTwin electron microscope (FEI, Hillsboro, OR, USA) equipped with a

16 megapixel TemCam-F416 (R) camera (TVIPS GmbH) at International Institute of

Molecular and Cell Biology in Warsaw.

Exopher tracking

Identification of branded expoher was performed with eC-CLEM open source software (Paul-

Gilloteaux *et al*, 2017). The NIRB frame is visible on both FM and EM image. The marks on

EM image are placed on the frame's edges and respectively its position is adjusted on FM

image allowing for correlation of both images. These coordinates allow to set out exopher

position on the TEM image.

RNA interference

RNA interference in *C. elegans* was performed using the standard RNAi feeding method and

RNAi clone (Kamath & Ahringer, 2003). For experiments, we used NGM plates

supplemented with 1 mM IPTG and 25 µg/µl carbenicillin seeded with HT115 *E.*

coli bacteria expressing double-stranded RNA (dsRNA) against the gene of interest or, as a

control, with bacteria without a vector. Worms were placed on freshly prepared RNAi plates,

either as age-synchronized pretzel-stage eggs, L1 larvae, or L4 larvae. The number of

exophers was counted on day 2 adult worms using a confocal microscope or a

stereomicroscope.

Stress influence on exophers production

Worms were age-synchronized using alkaline hypochlorite solution (bleaching procedure), as

previously described (Porta-de-la-Riva *et al*, 2012). The harvested eggs were incubated

overnight at 16 °C for hatching. Approx. 1000 L1 larvae were transferred on NGM plates and

incubated at 20 °C till they reached day-2 of adulthood. The worms were further channelled

to the respective stress treatments.

Oxidative stress

Approx. 100, day-2 adult worms were washed-off from the NGM plates and rinsed 3 times

with M9 buffer. The worms to be stressed were suspended in 1 ml of 5 mM hydrogen

peroxide solution prepared in M9 buffer, whereas control worms were suspended in M9

buffer. The tubes were incubated on a shaker at 20 °C for 60 minutes.

Heat stress

Similarly, approx. 100, day-2 adult worms were washed-off form NGM plates and rinsed 3

times with M9 buffer. The worms were further suspended in a 1 ml M9 buffer. The worms to

be heat stressed were incubated in a shaker at 33 °C for 60 minutes, whereas the control

animals were incubated at 20 °C for 60 minutes.

Exophers quantification

From each stress/control treatment, 30 worms were picked onto agarose pad slides

individually for the quantification of exophers. Exophers were quantified using a confocal

microscope. Obtained data were normalized to the average number of exophers present in

control animals and compared between conditions.

Number of exophers on consecutive days

For each biological replicate, 30 L4 larvae hermaphrodites or 50 L4 larvae males were

transferred to fresh NGM plates (5 hermaphrodites or 10 males per plate). For the next

consecutive 15 days, the number of exophers in each animal was counted using a

stereomicroscope. Hermaphrodites were transferred to fresh plates every 2 – 3 days. Males

were kept on the same plate until the end of the experiment. All animals that died during the

experiment time course were removed from the plate.

Measuring exophers in *fem-1* mutant

Approx. 50 L1 larvae from *fem-1(hc17)ts* mutant strain expressing GFP in BWM were

transferred to 2 fresh NGM plates. During the time course of the experiment, one of the

plates was kept at 15 °C while the second one was kept at 25 °C. When worms reached L4

larval stage, each worm was transferred to a separate plate and grown at the same

temperature as before. After 48 hours at 25 °C or 72 hours at 15 °C, using a

stereomicroscope, a number of exophers released from worms muscles was counted and

plates were scored for F1 offspring to assign all worms as fertile or infertile.

Additionally, a phenotype rescue experiment was performed for sterile animals grown at 25

°C. 24 hours after they reached the adulthood, single animals were placed on separate plates

and mated with 6-8 *him-5* mutant day-1 males. After next 48 hours, a number of exophers

released from worms muscles was counted.

FUdR assay

Age-synchronized animals were placed on NGM plates seeded with OP50 *E. coli* bacteria as

a food source until they reached young adulthood (day 0). Next, animals were selected and

moved to test plates containing 25 μM fluorodeoxyuridine (FUdR) to prevent embryonic

development and egg hatching (Mitchell *et al*, 1979) or control plates without FUdR. The

number of exophers was counted on adult day-2 using confocal microscopy.

Exophers and *in utero* eggs correlation

For correlating the number of exophers with the number of eggs present in worms uterus, 1 to

3-days old animals were used. First, muscular exophers for a single worm were counted using

the stereomicroscope. Next, worm was transferred to a 10 µl drop of 1.8 % hypochlorite

solution put on a microscope slide. Finally, after approx. 5 minutes when the hermaphrodite

mother was fully bleached, the number of eggs released from worm's uterus was counted.

Data analysis was performed using GraphPad Prism 8 software.

Starvation assay

To assess the influence of worms starvation on the exophers production, day-2 adult worms

and L4 larvae were moved to bacteria-free NGM plates. After 24 hours of food deprivation,

the number of exophers was counted using a stereomicroscope. As a control, day-3 adult

worms and day-1 adult worms grown for the whole time on NGM plates seeded with bacteria

were used. Obtained data were normalized to the average number of exophers present in

control animals and compared between conditions.

Eggs lysate assay

To obtain eggs, age-synchronized N2 gravid hermaphrodites were bleached using an alkaline

hypochlorite solution. Harvested eggs were suspended in appropriate volume of M9 buffer to

reach the concentration of approx. 200 eggs per µl. As a control, age-synchronized L2/L3

larvae were collected and suspended in M9 buffer to reach the concentration of approx. 25

larvae per µl. Next, eggs and larvae were flash-frozen in liquid nitrogen, thawed on ice, and

sonicated 3 times for 10 seconds to obtain eggs and larvae lysate. 150 µl of eggs or larvae

lysate was mixed with 150 µl of concentrated OP50 *E. coli* bacteria and placed in a 0.5 ml

Eppendorf tube. Approx. 30 day-1 young adult hermaphrodites were transferred to Eppendorf

tube containing a mixture of eggs or larvae lysate and bacteria. Next, the Eppendorf tube was

placed on the rotator for 1 hour at room temperature. After 1 hour, the content of the tube was

placed on the fresh NGM plate seeded with OP50 bacteria. 18 hours later the number of

exophers in each worm was counted. For mock control animals, the whole protocol was the

same except that for bleaching step no worms were used.

Worm exploratory behaviour

Age-synchronized, day-2 adult worms which had an average (6-8), few (< 2), or many (> 20)

exophers were sorted to separate NGM plates. Approximately 10 worms per replicate were

brought onto NGM plates and the worm movement was recorded for 2 minutes using the

WormLab system (MBF Bioscience). The frame rate, exposure time, and gain were set to 7.5

frames per second, 0.0031 s, and 1, respectively. The track length, straight-line distance,

center point speed, and the overall track pattern of individual worms were analyzed using the

WormLab software (MBF Bioscience).

Vitellogenin levels in embryos

Vitellogenin levels in embryos were measured based on the GFP signal from fluorescently

tagged VIT-2 protein. Approx. 200 late L4 larvae were transferred to fresh NGM plate. On

day 2 of worms adulthood, the number of exophers in each animal was counted and worms

with less than 2 or more than 20 visible exophers were transferred to new plates. Next,

animals from each group were individually transferred to a 10 µl drop of M9 buffer which

was placed on a microscope slide. Using a sharp injection needle, worm was cut open to

release eggs from the uterus. Fluorescent signal from 2-cell embryo stage was captured using

Leica M165FC stereomicroscope equipped with Leica EL6000 lamp, standard GFP filter set,

and Leica DFC365 FX CCD camera. Magnification used for recording pictures was set to

192x. Exposure time and gain were set to 600 ms and 2, respectively. The fluorescent signal

was quantified using Leica Las X software and was normalized to the average signal from

eggs, which were obtained from animals with less than 2 exophers.

Worms development assay

25 – 30 age-synchronized, day-2 adult worms which had an average (6-8), few (< 2), or many

(> 20) exophers were sorted to separate NGM plates. Gravid adults were allowed to lay eggs

for 4 hours, afterward removed from the plates, and the development of their offspring was

followed. 46 hours later, using the stereomicroscope, developmental stage of each animal was

checked, and the proportion between L4 larvae stage worms and young adult worms was

calculated.

Statistical analysis

Statistical tests used in this study were two-tailed Welch's t-test and Fisher's exact test. P-

value < 0.05 was considered significant.

Acknowledgements

We thank the Caenorhabditis Genetics Center (funded by the NIH National Center for

Research Resources, P40 OD010440) for strains; T. Hoppe for expert advice and Carl

Kutzner for RNAi clones; T. Wegierski for confocal microscopy assistance; B. Uszczyńska-

Ratajczak, K. Szczepanowska, H. Bringmann, and members of Chacińska and Pokrzywa

laboratories for discussions and comments on the manuscript. Multiphoton microscopy

imaging, near infrared branding, and serial block face scanning electron microscopy imaging
were performed at the Laboratory of Imaging Tissue Structure and Function at Nencki
Institute of Experimental Biology PAS. Transmission electron microscopy imaging and
ultramicrotomy was performed either at Core Facility of International Institute of Molecular
and Cell Biology in Warsaw or using the equipment of Laboratory of Electron Microscopy at
Nencki Institute of Experimental Biology PAS.

Funding
Work in the W.P. laboratory was supported by the Foundation for Polish Science
co financed by the European Union under the European Regional Development Fund (grant
POIR.04.04.00 00 5EAB/18 00 to K.B., K.K. and W.P.), the European Molecular
Biology Organization (EMBO Installation Grant No. 3916 to M.P. and W.P.), the Polish
National Science Center (grant UMO 2016/23/B/NZ3/00753 to N.S. and W.P.), and the
Deutsche Forschungsgemeinschaft (DFG FOR 2743 to W.P.). Work in the A.C. laboratory
was funded by "Regenerative Mechanisms for Health" project MAB/2017/2 (M.T. and A.C.)
carried out within the International Research Agendas programme of the Foundation for
Polish Science co-financed by the European Union under the European Regional
Development Fund and was supported by a POLONEZ Fellowship of National Science
Centre, Poland, 2016/21/P/NZ3/03891 (M.T.), within European Union's Horizon 2020
research and innovation programme under the Marie Skłodowska-Curie grant agreement no.
665778.

Author contributions
M.T., W.P., M.P., K.B., N.S., M.M., M.A.Ś., M.N., K.K. and N.N. conducted and designed
experiments, M.T. and W.P. conceived the project and supervised the study. W.P., A.C. and

M.T. secured the funding. W.P. and M.T. wrote the manuscript with input from A.C. and

M.P.

Author Information

The authors declare no competing financial interests. Further information and requests for

resources and reagents should be directed to and will be fulfilled by the Lead Contact,

Wojciech Pokrzywa (wpokrzywa@iimcb.gov.pl).

Supplementary Information

Supplementary Table 1

List of *C. elegans* strains used in this study

Supplementary Video 1

The video shows the cellular content being transfer from BWM cell to exopher via elastic

tube.