Biological invasions alter environmental microbiomes: a meta-analysis

Antonino Malacrinò[1,*], Victoria A. Sadowski[1], Tvisha K. Martin[2], Nathalia Cavichiolli de Oliveira[3], Ian J. Brackett[1], James D. Feller[1], Kristian J. Harris[1], Orlando Combita Heredia[1], Rosa Vescio[1,4], Alison E. Bennett[1]

[1] Department of Evolution, Ecology and Organismal Biology, The Ohio State University, Columbus (OH)
[2] College of Food, Agricultural and Environmental Sciences, The Ohio State University, Columbus (OH)
[3] Department of Entomology, University of Sao Paulo, Piracicaba (SP), Brazil
[4] Dipartimento AGRARIA, Università degli Studi Mediterranea, Reggio Calabria, Italy

*Corresponding author.
Email: malacrino.1@osu.edu; antonino.malacrino@gmail.com (AM)

Abstract

Biological invasions impact both agricultural and natural systems. The damage can be quantified in terms of both economic loss and reduction of biodiversity. Although the literature is quite rich about the impact of invasive species on plant and animal communities, their impact on environmental microbiomes is underexplored. Here, we re-analyze publicly available data using a common framework to create a global synthesis of the effects of biological invasions on environmental microbial communities. Our findings suggest that non-native species are responsible for the loss of microbial diversity and shifts in the structure of microbial populations. Therefore, the impact of biological invasions on native ecosystems might be more pervasive than previously thought, influencing both macro- and micro-biomes. We also identified gaps in the literature which encourage research on a wider variety of environments and invaders, and the influence of invaders across seasons and geographical ranges.

Introduction

Biological invasions have severe impacts on biodiversity, community composition and ecosystem functions [1–5]. Invasive plants can alter many important ecosystem functions including the nitrogen cycle [6], carbon cycle, and decomposition. For example, invasion by the plant Amur Honeysuckle altered the decomposition rate in the invaded environment likely through changes in litter quality [7]. Exotic snails have been found to alter carbon and nitrogen fluxes in freshwater systems through their consumption/excretion activity [8]. These functions are provided by environmental microbiomes. Yet, despite the implications for ecosystem functioning, we are still learning the consequences of biological invasions on environmental microbiomes.

Previous studies have shown biological invasions can impact the diversity and taxonomical structure of environmental microbiomes. For example, we often see a shift in soil microbiota following invasion by non-native plant species [9–19]. Removal of feral pigs increased the diversity of soil bacterial communities and shifted their structure [20], and invasive crustaceans [21], mussels [22] and jellyfish [23] produce changes in the structure of water microbiomes. However, shifts in environmental microbiome as consequence of biological invasions do not always occur. For example, invasion by the plants *Robinia pseudoacacia* [24], *Eucalyptus* sp. [25], *Vincetoxicum rossicum* [26] did not alter the structure of soil microbial communities. Also, microcosms exposed to the simultaneous invasion of multiple plant species [27–29] did not alter soil microbiomes. Similarly, soil microbiome structure in microcosms did not change with the addition of the invasive earthworm *Aporrectodea trapezoides* [30]. Several of these studies used techniques (e.g. DGGE, PLFA, t-RFLP) that limit fine scale investigations of biological invasions on environmental microbiome diversity and taxonomical composition. Among the studies using high-throughput amplicon-sequencing techniques, most did not find changes in microbiome diversity [13,15,18,19,21,25,27,28,30], few reported a decrease of microbial diversity in response to invasion [16,20,29], and fewer still reported an increase [11,14]. Thus, there is still little consensus on the effects of biological invasions on the diversity and taxonomical structure of the environmental microbiomes, both tied to the stability and function of microbial communities [31,32].

Our ability to draw broad conclusions from published studies is limited, because
individual studies have occurred within a limited geographical range or with a limited group of
species. Meta-analyses of published biological means have long enabled more robust
conclusions than individual studies [33–36]. However, the meta-analytic approach has less
frequently been applied to amplicon-sequencing data that represent environmental
microbiome community composition. The majority of meta-analytic metabarcoding studies
have occurred in the medical sciences [37–45]. This approach can be successfully used to
address ecological questions. For example, meta-analytic metabarcoding studies have found
common patterns in the structure of indoor microbiomes [46] and freshwater eukaryotes [47].
Shade et al. [48] also used a meta-analysis of metabarcoding datasets from different
environments highlighting a time-dependent structure of microbiomes. A meta-analytic
approach has also been used to test the effects of stressors (e.g. water availability,
temperature, heavy metals) on environmental microbiomes [49]. Thus meta-analyses on
microbiome data have a striking potential to address global-scale questions, generate new
hypotheses and model common patterns [50], because they provide across study comparisons
[39,51,52].
Here, we aim to test whether the effect of biological invasion on environmental
microbiomes can be generalized or is idiosyncratic. To do so, we collected publicly available
data and re-analyzed this data under a common framework. We tested the effect of invasive
species on the diversity and structure of environmental microbiomes with the hypothesis that
the presence of invasive species will decrease microbial diversity, and alter the composition of
the environmental microbiome. We then investigated whether certain taxonomical groups are
more responsive to biological invasions.

Methods

Data collection

We searched for metabarcoding studies that evaluated the effect of biological invasions
on environmental microbiomes, and compared invaded and non-invaded habitats. Our
literature search for this study was conducted using Web of Science Core Collection (accessed
on March 6[th], 2020) using the keywords "Invasive speci*" and "microbio*" published between
2010-2020, and found 1,471 studies. Two additional studies were added by searching the same
keywords on Google Scholar (Fig S1). Records were manually filtered based on the study design
appropriate for our research question. This step yielded 22 studies, and we further filtered
these studies based on data availability in public repositories. When data was not available, we
attempted to contact the corresponding author. Finally we selected only studies that used the
16S rRNA marker gene, primer pair 515F/806R [53] or 341F/785R [54], and Illumina MiSeq
sequencing platform. After discarding studies that failed quality checks (see below), we were
able to include a total of five studies (Tab. 1), summing up to a total of 356 samples. Three
studies focused on invasive plants, and the remaining studies focused on a mammal and a
mussel (Tab. 1). The invasive mussel study [22] was the only study performed in an aquatic
environment, while the remaining studies examined soil environments.
101
102

103
104   **Table 1**. Summary of studies included in the meta-analysis

Study ID	Invasive organism	Species	Invaded environment	Reference
MPG13011	Plant	*Agropyron cristatum, Bromus tectorum, Sisymbrium altissimum, Erodium cicutarium* and *Poa bulbosa*	Soil	Gibbons et al. [28]
MPG87547	Mammal	*Sus scrofa*	Soil	Wehr et al. [20]
PRJNA296487	Plant	*Microstegium vimineum, Rhamnus davurica* and *Ailanthus altissima*	Soil	Rodrigues et al. [19]
PRJNA320310	Plant	*Artemisia rothrockii*	Soil	Collins et al. [11]
PRJNA385848	Mussel	*Dreissena bugensis*	Water	Denef et al. [22]

105
We took the following steps to alleviate some of the potential sources of bias due to
studies performed in different labs, using different protocols and sequenced on different
instruments. First, all studies included were performed using the Illumina MiSeq platform, in
order to reduce the potential bias that might be generated by directly comparing data obtained
from different platforms. Second, all studies targeted the same region of the 16S rRNA, as
several primer pairs targeting different regions are currently published and widely used. Three
out of five papers we considered in our analysis used the 515F/806R primer pair [53], while two
used the 341F/785R [54]. Although there might be small differences between them, they
overlap in the V4 region of 16S gene (Fig. S2), so we feel confident that the chance of including
spurious OTUs in our analysis is quite negligible. To account for study-specific variances due to
small differences in sampling procedures and lab protocols, we also included the study itself,
the environment where the study was performed (i.e., soil or water) and the identity of the
invasive species as stratification variables in the PERMANOVA and as random factors in our
linear model. This allowed us to ensure that our results are not biased by study-specific
features.
Once the papers were selected, we assigned each a "Study ID" and collected meta-data
from each sample in each study (invasive species, type of organism, invaded environment). We
then downloaded data from repositories using SRA Toolkit 2.10.4 for data on the SRA
databases, or by directly downloading files from the MG-RAST database.
125

Data processing and analysis

Paired-end reads were merged using FLASH 1.2.11 [55] and data were processed using
QIIME 1.9.1 [56]. Quality-filtering of reads was performed using default parameters, binning
OTUs and discarding chimeric sequences identified with VSEARCH 2.14.2 [57]. Taxonomy for
representative sequences was determined by querying against the SILVA database v132 [58]

using the BLAST method. A phylogeny was obtained by aligning representative sequences using
MAFFT v7.464 [59] and reconstructing a phylogenetic tree using FastTree [60].

Data analysis was performed using R statistical software 3.5 [61] with the packages
phyloseq [62] and *vegan* [63]. Read counts were normalized using DESeq2 v1.22.2 [64] prior to
data analysis. Singletons and sequences classified as chloroplast were excluded, as well as
samples which had less than 5000 sequence counts. Shannon diversity was fit to a linear mixed-
effects model specifying *sample type* (invaded or control), *organism* (plant, mammal, mussel),
and their interactions as fixed factors with *studyID* and *environment* (soil or water) included as
random factors. Models were fit using the *lmer*() function under the *lme4* package [65] and the
package *emmeans* was used to infer pairwise contrasts (corrected using False Discovery Rate,
FDR). Furthermore, we explored the effects of *sample type* and *organism* on the structure of
the microbial communities using a multivariate approach. Distances between pairs of samples,
in terms of community composition, were calculated using a Unifrac matrix, and then visualized
using an RDA procedure. Differences between sample groups were inferred through
PERMANOVA multivariate analysis (999 permutations stratified at level of *studyID*, *environment*
and *identity of invasive species*). Pairwise contrasts from PERMANOVA were subjected to FDR
correction. Finally, the relative abundance of each bacterial family was fit using the *lmer*()
function to test the effects of *sample type* (invaded or control) on single taxa, with *studyID*,
organism (plant, mammal, mussel) and *environment* included as random factors.

Results

Our search yielded 5 studies with an appropriate experimental design and available
data, for a total of 356 samples. A few samples failed quality checks and we further considered
335 samples for downstream analyses. Sequences clustered into 22831 OTUs, after quality
checks, removal of singletons and "chloroplast" reads, with an average of 61776.92 reads per
sample. Although the number of OTUs might seem high, it is important to consider that we are
analyzing samples across multiple environments (soil and water) and from different
geographical regions which we would expect to increase richness.

Biological invasions led to a reduction in Shannon diversity (χ^2= 3.85, df=1, *P*=0.04, Fig
1A). We also found biological invasions altered microbiome community composition in the
invaded environment compared to the control (Tab. 2, Fig 1B). The type of invasive organism
(plant, mammal, or mussel) produced a different community structure (pairwise *P*<0.01, FDR
corrected). A deeper analysis of bacterial families (Fig. 1C, Table S2) revealed that some
taxonomic groups are significantly more abundant in invaded environments (Blastocatellaceae,
Chitinophagaceae, Nitrosomonadaceae, Pirellulaceae, Sphingomonadaceae), while others are
more abundant in non-invaded samples (Acetobacteraceae, Beijerinckiaceae, Gemmataceae,
Micromonosporaceae, Pedosphaeraceae, Solibacteraceae, Solirubrobacteraceae).

Table 2. Results from PERMANOVA analysis testing the effects of *sample type*
(invaded/control), *organism group* (plant, mammal, mussel) and their interaction on microbial
community composition. The factors *studyID* (unique for each study) and *environment* (soil or
water) were used as strata to constrain permutations.

Factor	df	R^2	F	P
Sample type (Invaded/Control)	1	0.011	6.68	<0.001
Organism group (plant, mammal, mussel)	3	0.411	118.86	<0.001
Sample type * Organism group	3	0.007	2.1	0.02

173
174

175   **Fig 1**. (A) Comparison of Shannon index between control and invaded environments. (B)
176   RDA ordination using a Bray-Curtis distance matrix of samples. (C) Comparison of the relative
177   abundance of microbial classes between control and invaded environments. * *P*<0.05, **
178   *P*<0.01, *** *P*<0.001

Discussion

We show biological invasions decrease the diversity of environmental microbiomes.
Thus, understanding the impact of biological invasions on environmental microbiomes is of high
priority if we want to preserve ecosystem functions [66]. While several studies have
investigated the effects of species invasions on environmental microbiomes, we still lack a
generalized consensus across different environmental microbiomes and systems. Previous
studies have found that invasive species increased environmental microbial diversity [11,14],
while others reported a decrease [16,20,29]. But the majority of studies did not analyze the
microbial diversity, because they used techniques that did not allow such analysis, or reported
no changes [9,10,12,13,15,17–19,21–28,30]. Within the studies included in our analysis,
invasion by feral pigs decreased soil microbial diversity, while invasion by *Artemisia rothrockii*
increased soil microbial diversity. The remaining three studies in our analysis reported no
effects of biological invasions on environmental microbiome diversity. Microbial diversity is tied
to the function of microbiomes, and changes in diversity can reflect changes in function [67–
69]. Changes in microbial diversity and function do not always have the same direction [70] and
this might explain the discrepancy between our results and other studies.

In contrast to diversity, our report of changes in community composition was relatively
consistent with the published literature and the individual results of the studies we analyzed.
Most studies of the influence of biological invasions on environmental microbiomes found that
biological invasions alter environmental microbial community composition. However, some
previous reports did not report changes [24–30], including the study by Gibbons et al. [28]
considered in our analysis. This variation may be due to individual effects of organisms on the
environment. For example, invasive plants may alter soil microbiome composition through root
exudates [5], and invasive mussels may alter water microbiome composition via bacterial
removal through their feeding activity [22]. Thus, reported influences on community
composition are more consistent. Alternatively changes in community composition might due
to the response of some bacterial groups to environmental disturbance. The bacterial families
that we found to be differentially abundant between the invaded and control environments

have diverse ecological functions ranging from nitrogen fixation and carbohydrate metabolism
to antimicrobial properties. Many of the families that showed a significant difference between
invaded and control environments have species members that play important roles at various
points during nitrogen and carbon cycling (i.e. Nitrosomonadaceae, Acetobacteraceae,
Chitinophagaceae, Micromonosporaceae, Gemmataceae, Beijerinckiaceae, Pirellulaceae) [71–
81]. However, nitrogen fixing and carbohydrate degrading bacteria did not have a unified
response to invaded environments as some increased and others decreased in abundance in
invaded environments. Many nitrogen fixing bacteria have been shown to respond to
environmental disturbance, such as Acidobacteria abundances during forest to pasture
conversions or Pirellulaceae's response to the presence of microplastics [82,83]. Thus, changes
in environmental microbiome community composition appear to be linked to changes in
ecosystem functions, although this pattern is not yet predictable for all functions and taxa.

Few previous studies on biological invasions have reported details on the differential
abundance of taxa, and among these we found limited general consensus. For example, some
studies report a decrease in abundance of bacteria associated with nitrogen cycling (e.g.
Nitrosphaeria, Nitrospira, Nitrosomonadales) [13,14,19], while others report an increase of
Nitrosomonadaceae following invasion [30]. In our study some groups associated with the
nitrogen cycle were positively associated with biological invasions (i.e. Nitrosomonadaceae,
Pirellulaceae, Chitinophagaceae) while others were negatively associated (Beijerinckiaceae,
Micromonosporaceae). Unfortunately, amplicon-based sequencing has a limited power to infer
changes in the functions of microbiomes. Future metagenomic and metatranscriptomic studies
are needed to investigate whether biological invasions alter gene content or gene expression of
environmental microbiomes, and whether this reflects changes in biogeochemical cycling.

Meta-analyses are also useful to highlight gaps in the literature, and here we highlight
some aspects that warrant further investigation. We identified a large gap in the availability of
sequencing data from multiple types of environments and types of invasive species. For our
analysis almost all available data came from two environments: four sets of data came from soil
and one came from freshwater. Greater effort is needed for sample collection from invasions in
both freshwater and marine environments. Without sufficient diversity of sample
environments, it is impossible to tell whether microbial shifts following an invasion are unique
to an invaded environment. Second, in our analysis the majority of data came from one type of
invasive species: plants. Noticeably absent from our dataset were invasions by insects, fish, and
amphibians. Sequencing data is needed from a larger number of invasive species to allow us to
broadly assess shifts in microbial community structure. A third gap we identified was the lack of
spatial and temporal resolution. Almost all of the initially identified 22 studies we assessed
were also restricted to one season of sampling and were conducted in the Northern
Hemisphere. Thus, it is impossible to validate the influence of latitude with existing datasets or
explore how seasonality and biological invasions interact to modulate microbial communities.
Thus, there are a number of opportunities for future research on how biological invasions alter
environmental microbial communities.

Here we analyzed 16S amplicon sequencing data from five studies and show that
biological invasions influence both the diversity and the structure of environmental
microbiomes. We identified a number of gaps in our knowledge, including the need to assess a
wider range of environments, invasive species, temporal variation, and latitudinal variation. We

also demonstrate the power of re-analysis of publicly available datasets using a common
pipeline which benefited from open-data initiatives.

Acknowldgements
254   We would like to thank Kali Mattingly (Ohio State University, USA) and Davide Rassati
255 (Università degli Studi di Padova, Italy) for their helpful comments on the manuscript.

Data accessibility
257   Raw data from single studies is publicly available. The code used to perform our
258 analyses has been deposited on GitHub: https://github.com/amalacrino/
259 Biol_Invasion_Microb_MA

References

1. Chapin III FS, Zavaleta ES, Eviner VT, Naylor RL, Vitousek PM, Reynolds HL, et al. Consequences of changing biodiversity. Nature. 2000;405: 234–242. doi:10.1038/35012241

2. Pejchar L, Mooney HA. Invasive species, ecosystem services and human well-being. Trends Ecol Evol. 2009;24: 497–504. doi:10.1016/j.tree.2009.03.016

3. Shackleton RT, Shackleton CM, Kull CA. The role of invasive alien species in shaping local livelihoods and human well-being: A review. J Environ Manage. 2019;229: 145–157. doi:10.1016/j.jenvman.2018.05.007

4. Charles H, Dukes JS. Impacts of Invasive Species on Ecosystem Services. Biological Invasions. Berlin, Heidelberg: Springer Berlin Heidelberg; 2008. pp. 217–237. doi:10.1007/978-3-540-36920-2_13

5. Weidenhamer JD, Callaway RM. Direct and Indirect Effects of Invasive Plants on Soil Chemistry and Ecosystem Function. J Chem Ecol. 2010;36: 59–69. doi:10.1007/s10886-009-9735-0

6. Lee MR, Bernhardt ES, van Bodegom PM, Cornelissen JHC, Kattge J, Laughlin DC, et al. Invasive species' leaf traits and dissimilarity from natives shape their impact on nitrogen cycling: a meta-analysis. New Phytol. 2017;213: 128–139. doi:10.1111/nph.14115

7. Arthur MA, Bray SR, Kuchle CR, McEwan RW. The influence of the invasive shrub, Lonicera maackii, on leaf decomposition and microbial community dynamics. Plant Ecol. 2012;213: 1571–1582. doi:10.1007/s11258-012-0112-7

8. Hall Jr RO, Tank JL, Dybdahl MF. Exotic snails dominate nitrogen and carbon cycling in a highly productive stream. Front Ecol Environ. 2003;1: 407–411. doi:10.1890/1540-9295(2003)001[0407:ESDNAC]2.0.CO;2

9. Lazzaro L, Giuliani C, Fabiani A, Agnelli AE, Pastorelli R, Lagomarsino A, et al. Soil and plant changing after invasion: The case of Acacia dealbata in a Mediterranean ecosystem. Sci Total Environ. 2014;497–498: 491–498. doi:10.1016/j.scitotenv.2014.08.014

10. Malinich E, Lynn-Bell N, Kourtev PS. The effect of the invasive Elaeagnus umbellata on soil microbial communities depends on proximity of soils to plants. Ecosphere. 2017;8: e01827. doi:10.1002/ecs2.1827

11. Collins CG, Carey CJ, Aronson EL, Kopp CW, Diez JM. Direct and indirect effects of native range expansion on soil microbial community structure and function. Austin A, editor. J Ecol. 2016;104: 1271–1283. doi:10.1111/1365-2745.12616

12. Lorenzo P, Pereira CS, Rodríguez-Echeverría S. Differential impact on soil microbes of allelopathic compounds released by the invasive Acacia dealbata Link. Soil Biol Biochem. 2013;57: 156–163. doi:10.1016/j.soilbio.2012.08.018

13. Parsons LS, Sayre J, Ender C, Rodrigues JLM, Barberán A. Soil microbial communities in restored and unrestored coastal dune ecosystems in California. Restor Ecol. 2020; rec.13101. doi:10.1111/rec.13101

14. Piper CL, Siciliano SD, Winsley T, Lamb EG. Smooth brome invasion increases rare soil bacterial species prevalence, bacterial species richness and evenness. De Deyn G, editor. J Ecol. 2015;103: 386–396. doi:10.1111/1365-2745.12356

15. Zhang H-Y, Goncalves P, Copeland E, Qi S-S, Dai Z-C, Li G-L, et al. Invasion by the weed

Conyza canadensis alters soil nutrient supply and shifts microbiota structure. Soil Biol
Biochem. 2020;143: 107739. doi:10.1016/j.soilbio.2020.107739

16. Le Roux JJ, Ellis AG, van Zyl L-M, Hosking ND, Keet J-H, Yannelli FA. Importance of soil
legacy effects and successful mutualistic interactions during Australian acacia invasions in
nutrient-poor environments. Aerts R, editor. J Ecol. 2018;106: 2071–2081.
doi:10.1111/1365-2745.12965

17. Brunel C, Beifen Y, Pouteau R, Li J, van Kleunen M. Responses of Rhizospheric Microbial
Communities of Native and Alien Plant Species to Cuscuta Parasitism. Microb Ecol.
2020;79: 617–630. doi:10.1007/s00248-019-01438-z

18. Pickett B, Irvine IC, Bullock E, Arogyaswamy K, Aronson E. Legacy effects of invasive grass
impact soil microbes and native shrub growth. Invasive Plant Sci Manag. 2019;12: 22–35.
doi:10.1017/inp.2018.32

19. Rodrigues RR, Pineda RP, Barney JN, Nilsen ET, Barrett JE, Williams MA. Plant Invasions
Associated with Change in Root-Zone Microbial Community Structure and Diversity. Liu J,
editor. PLoS One. 2015;10: e0141424. doi:10.1371/journal.pone.0141424

20. Wehr NH, Kinney KM, Nguyen NH, Giardina CP, Litton CM. Changes in soil bacterial
community diversity following the removal of invasive feral pigs from a Hawaiian tropical
montane wet forest. Sci Rep. 2019;9: 14681. doi:10.1038/s41598-019-48922-7

21. Boeker C, Geist J. Effects of invasive and indigenous amphipods on physico-chemical and
microbial properties in freshwater substrates. Aquat Ecol. 2015;49: 467–480.
doi:10.1007/s10452-015-9539-y

22. Denef VJ, Carrick HJ, Cavaletto J, Chiang E, Johengen TH, Vanderploeg HA. Lake Bacterial
Assemblage Composition Is Sensitive to Biological Disturbance Caused by an Invasive
Filter Feeder. Kent AD, editor. mSphere. 2017;2. doi:10.1128/mSphere.00189-17

23. Manzari C, Fosso B, Marzano M, Annese A, Caprioli R, D'Erchia AM, et al. The influence of
invasive jellyfish blooms on the aquatic microbiome in a coastal lagoon (Varano, SE Italy)
detected by an Illumina-based deep sequencing strategy. Biol Invasions. 2015;17: 923–
940. doi:10.1007/s10530-014-0810-2

24. Lazzaro L, Mazza G, D'Errico G, Fabiani A, Giuliani C, Inghilesi AF, et al. How ecosystems
change following invasion by Robinia pseudoacacia: Insights from soil chemical
properties and soil microbial, nematode, microarthropod and plant communities. Sci
Total Environ. 2018;622–623: 1509–1518. doi:10.1016/j.scitotenv.2017.10.017

25. Hernández-Gómez O, Byrne AQ, Gunderson AR, Jenkinson TS, Noss CF, Rothstein AP, et
al. Invasive vegetation affects amphibian skin microbiota and body condition. PeerJ.
2020;8: e8549. doi:10.7717/peerj.8549

26. Thompson GL, Bell TH, Kao-Kniffin J. Rethinking Invasion Impacts across Multiple Field
Sites Using European Swallowwort (Vincetoxicum rossicum) as a Model Invader.
Invasive Plant Sci Manag. 2018;11: 109–116. doi:10.1017/inp.2018.22

27. Carey CJ, Beman JM, Eviner VT, Malmstrom CM, Hart SC. Soil microbial community
structure is unaltered by plant invasion, vegetation clipping, and nitrogen fertilization in
experimental semi-arid grasslands. Front Microbiol. 2015;6.
doi:10.3389/fmicb.2015.00466

28. Gibbons SM, Lekberg Y, Mummey DL, Sangwan N, Ramsey PW, Gilbert JA. Invasive Plants
Rapidly Reshape Soil Properties in a Grassland Ecosystem. Shade A, editor. mSystems.

2017;2. doi:10.1128/mSystems.00178-16

29. Gomes SIF, Merckx VSFT, Hynson NA. Biological invasions increase the richness of
arbuscular mycorrhizal fungi from a Hawaiian subtropical ecosystem. Biol Invasions.
2018;20: 2421–2437. doi:10.1007/s10530-018-1710-7

30. de Menezes AB, Prendergast-Miller MT, Macdonald LM, Toscas P, Baker G, Farrell M, et
al. Earthworm-induced shifts in microbial diversity in soils with rare versus established
invasive earthworm populations. FEMS Microbiol Ecol. 2018;94.
doi:10.1093/femsec/fiy051

31. Bier RL, Bernhardt ES, Boot CM, Graham EB, Hall EK, Lennon JT, et al. Linking microbial
community structure and microbial processes: an empirical and conceptual overview.
Muyzer G, editor. FEMS Microbiol Ecol. 2015;91: fiv113. doi:10.1093/femsec/fiv113

32. Gibbons SM, Gilbert JA. Microbial diversity — exploration of natural ecosystems and
microbiomes. Curr Opin Genet Dev. 2015;35: 66–72. doi:10.1016/j.gde.2015.10.003

33. Gurevitch J, Koricheva J, Nakagawa S, Stewart G. Meta-analysis and the science of
research synthesis. Nature. 2018;555: 175–182. doi:10.1038/nature25753

34. Nakagawa S, Poulin R. Meta-analytic insights into evolutionary ecology: an introduction
and synthesis. Evol Ecol. 2012;26: 1085–1099. doi:10.1007/s10682-012-9593-z

35. Nakagawa S, Santos ESA. Methodological issues and advances in biological meta-analysis.
Evol Ecol. 2012;26: 1253–1274. doi:10.1007/s10682-012-9555-5

36. Senior AM, Grueber CE, Kamiya T, Lagisz M, O'Dwyer K, Santos ESA, et al. Heterogeneity
in ecological and evolutionary meta-analyses: its magnitude and implications. Ecology.
2016;97: 3293–3299. doi:10.1002/ecy.1591

37. Lozupone CA, Stombaugh J, Gonzalez A, Ackermann G, Wendel D, Vazquez-Baeza Y, et al.
Meta-analyses of studies of the human microbiota. Genome Res. 2013;23: 1704–1714.
doi:10.1101/gr.151803.112

38. Walters WA, Xu Z, Knight R. Meta-analyses of human gut microbes associated with
obesity and IBD. FEBS Lett. 2014;588: 4223–4233. doi:10.1016/j.febslet.2014.09.039

39. Duvallet C, Gibbons SM, Gurry T, Irizarry RA, Alm EJ. Meta-analysis of gut microbiome
studies identifies disease-specific and shared responses. Nat Commun. 2017;8: 1784.
doi:10.1038/s41467-017-01973-8

40. Koren O, Knights D, Gonzalez A, Waldron L, Segata N, Knight R, et al. A Guide to
Enterotypes across the Human Body: Meta-Analysis of Microbial Community Structures
in Human Microbiome Datasets. Eisen JA, editor. PLoS Comput Biol. 2013;9: e1002863.
doi:10.1371/journal.pcbi.1002863

41. Krych L, Hansen CHF, Hansen AK, van den Berg FWJ, Nielsen DS. Quantitatively Different,
yet Qualitatively Alike: A Meta-Analysis of the Mouse Core Gut Microbiome with a View
towards the Human Gut Microbiome. Bereswill S, editor. PLoS One. 2013;8: e62578.
doi:10.1371/journal.pone.0062578

42. Wang J, Kurilshikov A, Radjabzadeh D, Turpin W, Croitoru K, Bonder MJ, et al. Meta-
analysis of human genome-microbiome association studies: the MiBioGen consortium
initiative. Microbiome. 2018;6: 101. doi:10.1186/s40168-018-0479-3

43. Jiao N, Baker SS, Nugent CA, Tsompana M, Cai L, Wang Y, et al. Gut microbiome may
contribute to insulin resistance and systemic inflammation in obese rodents: a meta-
analysis. Physiol Genomics. 2018;50: 244–254. doi:10.1152/physiolgenomics.00114.2017

44.   Wirbel J, Pyl PT, Kartal E, Zych K, Kashani A, Milanese A, et al. Meta-analysis of fecal
393   metagenomes reveals global microbial signatures that are specific for colorectal cancer.
394   Nat Med. 2019;25: 679–689. doi:10.1038/s41591-019-0406-6

395   45.   Bisanz JE, Upadhyay V, Turnbaugh JA, Ly K, Turnbaugh PJ. Meta-Analysis Reveals
396   Reproducible Gut Microbiome Alterations in Response to a High-Fat Diet. Cell Host
397   Microbe. 2019;26: 265-272.e4. doi:10.1016/j.chom.2019.06.013

398   46.   Adams RI, Bateman AC, Bik HM, Meadow JF. Microbiota of the indoor environment: a
399   meta-analysis. Microbiome. 2015;3: 49. doi:10.1186/s40168-015-0108-3

400   47.   Debroas D, Domaizon I, Humbert J-F, Jardillier L, Lepère C, Oudart A, et al. Overview of
401   freshwater microbial eukaryotes diversity: a first analysis of publicly available
402   metabarcoding data. FEMS Microbiol Ecol. 2017;93. doi:10.1093/femsec/fix023

403   48.   Shade A, Gregory Caporaso J, Handelsman J, Knight R, Fierer N. A meta-analysis of
404   changes in bacterial and archaeal communities with time. ISME J. 2013;7: 1493–1506.
405   doi:10.1038/ismej.2013.54

406   49.   Rocca JD, Simonin M, Blaszczak JR, Ernakovich JG, Gibbons SM, Midani FS, et al. The
407   Microbiome Stress Project: Toward a Global Meta-Analysis of Environmental Stressors
408   and Their Effects on Microbial Communities. Front Microbiol. 2019;9.
409   doi:10.3389/fmicb.2018.03272

410   50.   Duvallet C. Meta-analysis generates and prioritizes hypotheses for translational
411   microbiome research. Microb Biotechnol. 2018;11: 273–276. doi:10.1111/1751-
412   7915.13047

413   51.   Drewes JL, White JR, Dejea CM, Fathi P, Iyadorai T, Vadivelu J, et al. High-resolution
414   bacterial 16S rRNA gene profile meta-analysis and biofilm status reveal common
415   colorectal cancer consortia. npj Biofilms Microbiomes. 2017;3: 34. doi:10.1038/s41522-
416   017-0040-3

417   52.   Ramirez KS, Knight CG, de Hollander M, Brearley FQ, Constantinides B, Cotton A, et al.
418   Detecting macroecological patterns in bacterial communities across independent studies
419   of global soils. Nat Microbiol. 2018;3: 189–196. doi:10.1038/s41564-017-0062-x

420   53.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al.
421   Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.
422   Proc Natl Acad Sci. 2011;108: 4516–4522. doi:10.1073/pnas.1000080107

423   54.   Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of
424   general 16S ribosomal RNA gene PCR primers for classical and next-generation
425   sequencing-based diversity studies. Nucleic Acids Res. 2013;41: e1–e1.
426   doi:10.1093/nar/gks808

427   55.   Magoc T, Salzberg SL. FLASH: fast length adjustment of short reads to improve genome
428   assemblies. Bioinformatics. 2011;27: 2957–2963. doi:10.1093/bioinformatics/btr507

429   56.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-
430   throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms.
431   ISME J. 2012;6: 1621–1624. doi:10.1038/ismej.2012.8

432   57.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source tool
433   for metagenomics. PeerJ. 2016;4: e2584. doi:10.7717/peerj.2584

434   58.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal
435   RNA gene database project: improved data processing and web-based tools. Nucleic

Acids Res. 2012;41: D590–D596. doi:10.1093/nar/gks1219

59.   Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol Biol Evol. 2013;30: 772–780.
doi:10.1093/molbev/mst010

60.   Price MN, Dehal PS, Arkin AP. FastTree: Computing Large Minimum Evolution Trees with
Profiles instead of a Distance Matrix. Mol Biol Evol. 2009;26: 1641–1650.
doi:10.1093/molbev/msp077

61.   R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2016. 2017.

62.   McMurdie PJ, Holmes S. phyloseq: An R package for reproducible interactive analysis and
graphics of microbiome census data. Watson M, editor. PLoS One. 2013;8: e61217.
doi:10.1371/journal.pone.0061217

63.   Dixon P. VEGAN, a package of R functions for community ecology. J Veg Sci. 2003;14:
927–930. doi:10.1111/j.1654-1103.2003.tb02228.x

64.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15: 550. doi:10.1186/s13059-014-0550-8

65.   Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. J
Stat Softw. 2015;67: 1–48. doi:10.18637/jss.v067.i01

66.   Doblas-Miranda E, Martínez-Vilalta J, Lloret F, Álvarez A, Ávila A, Bonet FJ, et al.
Reassessing global change research priorities in mediterranean terrestrial ecosystems:
how far have we come and where do we go from here? Glob Ecol Biogeogr. 2015;24: 25–
43. doi:10.1111/geb.12224

67.   Delgado-Baquerizo M, Maestre FT, Reich PB, Jeffries TC, Gaitan JJ, Encinar D, et al.
Microbial diversity drives multifunctionality in terrestrial ecosystems. Nat Commun.
2016;7: 10541. doi:10.1038/ncomms10541

68.   Miki T, Yokokawa T, Matsui K. Biodiversity and multifunctionality in a microbial
community: a novel theoretical approach to quantify functional redundancy. Proc R Soc B
Biol Sci. 2014;281: 20132498. doi:10.1098/rspb.2013.2498

69.   Philippot L, Spor A, Hénault C, Bru D, Bizouard F, Jones CM, et al. Loss in microbial
diversity affects nitrogen cycling in soil. ISME J. 2013;7: 1609–1619.
doi:10.1038/ismej.2013.34

70.   Mendes LW, Tsai SM, Navarrete AA, de Hollander M, van Veen JA, Kuramae EE. Soil-
Borne Microbiome: Linking Diversity to Function. Microb Ecol. 2015;70: 255–265.
doi:10.1007/s00248-014-0559-2

71.   Chen J, Mo L, Zhang Z, Nan J, Xu D, Chao L, et al. Evaluation of the ecological restoration
of a coal mine dump by exploring the characteristics of microbial communities. Appl Soil
Ecol. 2020;147: 103430. doi:10.1016/j.apsoil.2019.103430

72.   Cheng J, Chen Y, He T, Liao R, Liu R, Yi M, et al. Soil nitrogen leaching decreases as biogas
slurry DOC/N ratio increases. Appl Soil Ecol. 2017;111: 105–113.
doi:10.1016/j.apsoil.2016.12.001

73.   Warneke C, Karl T, Judmaier H, Hansel A, Jordan A, Lindinger W, et al. Acetone,
methanol, and other partially oxidized volatile organic emissions from dead plant matter
by abiological processes: Significance for atmospheric HO x chemistry. Global
Biogeochem Cycles. 1999;13: 9–17. doi:10.1029/98GB02428

74.   Kämpfer P, Lodders N, Falsen E. Hydrotalea flava gen. nov., sp. nov., a new member of
the phylum Bacteroidetes and allocation of the genera Chitinophaga, Sediminibacterium,
Lacibacter, Flavihumibacter, Flavisolibacter, Niabella, Niastella, Segetibacter,
Parasegetibacter, Terrimonas, Fer. Int J Syst Evol Microbiol. 2011;61: 518–523.
doi:10.1099/ijs.0.023002-0

75.   Madhaiyan M, Poonguzhali S, Senthilkumar M, Pragatheswari D, Lee J-S, Lee K-C.
Arachidicoccus rhizosphaerae gen. nov., sp. nov., a plant-growth-promoting bacterium in
the family Chitinophagaceae isolated from rhizosphere soil. Int J Syst Evol Microbiol.
2015;65: 578–586. doi:10.1099/ijs.0.069377-0

76.   Reis VM, Teixeira KR dos S. Nitrogen fixing bacteria in the family Acetobacteraceae and
their role in agriculture. J Basic Microbiol. 2015;55: 931–949.
doi:10.1002/jobm.201400898

77.   Trujillo ME, Bacigalupe R, Pujic P, Igarashi Y, Benito P, Riesco R, et al. Genome Features of
the Endophytic Actinobacterium Micromonospora lupini Strain Lupac 08: On the Process
of Adaptation to an Endophytic Life Style? Brüggemann H, editor. PLoS One. 2014;9:
e108522. doi:10.1371/journal.pone.0108522

78.   Carro L, Nouioui I, Sangal V, Meier-Kolthoff JP, Trujillo ME, Montero-Calasanz M del C, et
al. Genome-based classification of micromonosporae with a focus on their
biotechnological and ecological potential. Sci Rep. 2018;8: 525. doi:10.1038/s41598-017-
17392-0

79.   Dedysh SN, Ivanova AA. Planctomycetes in boreal and subarctic wetlands: diversity
patterns and potential ecological functions. FEMS Microbiol Ecol. 2019;95.
doi:10.1093/femsec/fiy227

80.   Wegner C-E, Gorniak L, Riedel S, Westermann M, Küsel K. Lanthanide-Dependent
Methylotrophs of the Family Beijerinckiaceae: Physiological and Genomic Insights.
Stams AJM, editor. Appl Environ Microbiol. 2019;86. doi:10.1128/AEM.01830-19

81.   Neufeld JD, Schäfer H, Cox MJ, Boden R, McDonald IR, Murrell JC. Stable-isotope probing
implicates Methylophaga spp and novel Gammaproteobacteria in marine methanol and
methylamine metabolism. ISME J. 2007;1: 480–491. doi:10.1038/ismej.2007.65

82.   Miao L, Wang P, Hou J, Yao Y, Liu Z, Liu S, et al. Distinct community structure and
microbial functions of biofilms colonizing microplastics. Sci Total Environ. 2019;650:
2395–2402. doi:10.1016/j.scitotenv.2018.09.378

83.   Navarrete AA, Venturini AM, Meyer KM, Klein AM, Tiedje JM, Bohannan BJM, et al.
Differential Response of Acidobacteria Subgroups to Forest-to-Pasture Conversion and
Their Biogeographic Patterns in the Western Brazilian Amazon. Front Microbiol. 2015;6.
doi:10.3389/fmicb.2015.01443