Biological invasions alter environmental microbiomes: a

meta-analysis

Antonino Malacrinò[1],*, Victoria A. Sadowski[1], Tvisha K. Martin[2], Nathalia Cavichiolli de Oliveira[3],

Ian J. Brackett[1], James D. Feller[1], Kristian J. Harris[1], Orlando Combita Heredia[1], Rosa Vescio[1,4],

Alison E. Bennett[1]

[1] Department of Evolution, Ecology and Organismal Biology, The Ohio State University,

Columbus (OH)

[2] College of Food, Agricultural and Environmental Sciences, The Ohio State University,

Columbus (OH)

[3] Department of Entomology, University of Sao Paulo, Piracicaba (SP), Brazil

[4] Dipartimento AGRARIA, Università degli Studi Mediterranea, Reggio Calabria, Italy

*Corresponding author.

Email: malacrino.1@osu.edu; antonino.malacrino@gmail.com (AM)

Abstract

Biological invasions impact both agricultural and natural systems. The damage can be quantified in terms of both economic loss and reduction of biodiversity. Although the literature is quite rich about the impact of invasive species on plant and animal communities, their impact on environmental microbiomes is underexplored. Here, we re-analyze publicly available data using a common framework to create a global synthesis of the effects of biological invasions on environmental microbial communities. Our findings suggest that non-native species are responsible for the loss of microbial diversity and shifts in the structure of microbial populations. Therefore, the impact of biological invasions on native ecosystems might be more pervasive than previously thought, influencing both macro- and micro-biomes. We also identified gaps in the literature which encourage research on a wider variety of environments and invaders, and the influence of invaders across seasons and geographical ranges.

Introduction

Biological invasions have severe impacts on biodiversity, community composition and ecosystem functions [1–5]. Invasive plants can alter many important ecosystem functions including the nitrogen cycle [6], carbon cycle, and decomposition. For example, invasion by the plant Amur Honeysuckle altered the decomposition rate in the invaded environment likely through changes in litter quality [7]. Exotic snails have been found to alter carbon and nitrogen fluxes in freshwater systems through their consumption/excretion activity [8]. These functions are provided by environmental microbiomes. Yet, despite the implications for ecosystem

functioning, we are still learning the consequences of biological invasions on environmental

microbiomes.

Previous studies have shown biological invasions can impact the diversity and

taxonomical structure of environmental microbiomes. For example, we often see a shift in soil

microbiota following invasion by non-native plant species [9,10,19,11–18]. Removal of feral pigs

increased the diversity of soil bacterial communities and shifted their structure [20], and

invasive crustaceans [21], mussels [22] and jellyfish [23] produce changes in the structure of

water microbiomes. However, shifts in environmental microbiome as consequence of biological

invasions do not always occur. For example, invasion by the plants *Robinia pseudoacacia* [24],

Eucalyptus sp. [25], and *Vincetoxicum rossicum* [26] did not alter the structure of soil microbial

communities. Also, some microcosms exposed to the simultaneous invasion of multiple plant

species [27–29] did not alter soil microbiomes. Similarly, soil microbiome structure in

microcosms did not change with the addition of the invasive earthworm *Aporrectodea*

trapezoides [30]. Several of these studies used techniques (e.g. DGGE, PLFA, t-RFLP) that limit

fine scale investigations of biological invasions on environmental microbiome diversity and

taxonomical composition. Among the studies using high-throughput amplicon-sequencing

techniques, most did not find changes in microbiome diversity [12,14,17,18,21,25,27,28,30],

few reported a decrease of microbial diversity in response to invasion [15,20,29], and fewer still

reported an increase [13,19]. Thus, there is still little consensus on the effects of biological

invasions on the diversity and taxonomical structure of the environmental microbiomes, both

tied to the stability and function of microbial communities [31,32].

Our ability to draw broad conclusions from published studies is limited, because

individual studies have occurred within a limited geographical range or with a limited group of

species. Meta-analyses of published biological means have long enabled more robust

conclusions than individual studies [33–36]. However, the meta-analytic approach has less

frequently been applied to amplicon-sequencing data that represent environmental

microbiome community composition. The majority of meta-analytic metabarcoding studies

have occurred in the medical sciences [37–45]. This approach can be successfully used to

address ecological questions. For example, meta-analytic metabarcoding studies have found

common patterns in the structure of indoor microbiomes [46] and freshwater eukaryotes [47].

Shade et al. [48] also used a meta-analysis of metabarcoding datasets from different

environments highlighting a time-dependent structure of microbiomes. A meta-analytic

approach has also been used to test the effects of stressors (e.g. water availability,

temperature, heavy metals) on environmental microbiomes [49]. Thus meta-analyses on

microbiome data have a striking potential to address global-scale questions, generate new

hypotheses and model common patterns [50], because they provide across study comparisons

[39,51,52].

Here, we aim to test whether the effect of biological invasion on environmental

microbiomes can be generalized or is idiosyncratic. To do so, we collected publicly available

data and re-analyzed this data under a common framework. We tested the effect of invasive

species on the diversity and structure of environmental microbiomes, with the hypothesis that

the presence of invasive species will decrease microbial diversity and alter the composition of

the environmental microbiome. We then investigated whether certain taxonomical groups are

more responsive to biological invasions.

Methods

Data collection

We searched for metabarcoding studies that evaluated the effect of biological invasions on

environmental microbiomes, and compared invaded and non-invaded habitats. Our literature

search for this study was conducted using Web of Science Core Collection (accessed on March

6th, 2020) using the keywords "Invasive speci*" and "microbio*" published between 2010-2020,

and found 1,471 studies. Two additional studies were added by searching the same keywords

on Google Scholar (S1 Figure). Records were manually filtered based on the study design

appropriate for our research question. This step yielded 22 studies, and we further filtered

these studies based on data availability in public repositories. When data was not available, we

attempted to contact the corresponding author. Finally we selected only studies that used the

16S rRNA marker gene, primer pair 515F/806R [53] or 341F/785R [54], and Illumina MiSeq

sequencing platform. After discarding studies that failed quality checks (see below), we were

able to include a total of five studies (Tab. 1), summing up to a total of 356 samples. The study

by Gibbons et al. [28] tested the impact of five invasive plant species (*Agropyron cristatum,*

Bromus tectorum, Sisymbrium altissimum, Erodium cicutarium and *Poa bulbosa*) on soil

microbiome using microcosms, comparing monocultures of each one of them towards a

mixture of eight native plant species. A similar question was tested in Rodrigues et al. [18] in

field condition. They identified three locations invaded by three different exotic plant species

(*Microstegium vimineum*, *Rhamnus davurica* and *Ailanthus altissima*) and, within each location,

they sampled soil form an invaded area and a non-invaded area for comparisons. Similarly,

Collins et al. [19] compared the soil microbial community of field sites invaded by *Artemisia*

rothrockii to non-invaded sites. The study by Wehr et al. [20] focused on the effects of feral pig

(*Sus scrofa*) invasion on soil microbiome comparing invaded areas to those where pigs were

removed over a ~25 year chronosequence. Finally, the only study performed in an aquatic

environment [22] compared water samples collected in lake areas invaded by the exotic mussel

Dreissena bugensis to non-invaded sites. Three studies focused on invasive plants, and the

remaining studies focused on a mammal and a mussel (Tab. 1).

Table 1. Summary of studies included in the meta-analysis

Study ID	Invasive organism	Species	Invaded environment	Reference
MPG13011	Plant	*Agropyron cristatum, Bromus tectorum, Sisymbrium altissimum, Erodium cicutarium* and *Poa bulbosa*	Soil	Gibbons et al. [28]
MPG87547	Mammal	*Sus scrofa*	Soil	Wehr et al. [20]
PRJNA296487	Plant	*Microstegium vimineum, Rhamnus davurica* and *Ailanthus altissima*	Soil	Rodrigues et al. [18]
PRJNA320310	Plant	*Artemisia rothrockii*	Soil	Collins et al.

				[19]
PRJNA385848	Mussel	*Dreissena bugensis*	Water	Denef et al. [22]

We took the following steps to alleviate some of the potential sources of bias due to

studies performed in different labs, using different protocols and sequenced on different

instruments. First, all studies included were performed using the Illumina MiSeq platform, in

order to reduce the potential bias that might be generated by directly comparing data obtained

from different platforms. Second, all studies targeted the same region of the 16S rRNA, as

several primer pairs targeting different regions are currently published and widely used. Three

out of five papers we considered in our analysis used the 515F/806R primer pair [53], while two

used the 341F/785R [54]. Because these primer pairs overlap in the V4 region of 16S rRNA we

feel confident that the chance of including spurious OTUs in our analysis is quite negligible. To

account for study-specific variances due to small differences in sampling procedures and lab

protocols, we also included the study itself, the environment where the study was performed

(i.e., soil or water) and the identity of the invasive species as stratification variables in the

PERMANOVA and as random factors in our linear model. This allowed us to ensure that our

results are not biased by study-specific features.

Once the papers were selected, we assigned each a "Study ID" and collected meta-data

from each sample in each study (invasive species, type of organism, invaded environment). We

then downloaded data from repositories using SRA Toolkit 2.10.4 for data on the SRA

databases, or by manually downloading files from the MG-RAST database.

Data processing and analysis

Paired-end reads were merged using FLASH 1.2.11 [55] and data were processed using

QIIME 1.9.1 [56]. Quality-filtering of reads was performed using default parameters, binning

OTUs and discarding chimeric sequences identified with VSEARCH 2.14.2 [57]. Taxonomy for

representative sequences was determined by querying against the SILVA database v132 [58]

using the BLAST method. A phylogeny was obtained by aligning representative sequences using

MAFFT v7.464 [59] and reconstructing a phylogenetic tree using FastTree [60].

Data analysis was performed using R statistical software 3.5 [61] with the packages

phyloseq [62] and *vegan* [63]. Read counts were normalized using DESeq2 v1.22.2 [64] prior to

data analysis. Singletons and sequences classified as chloroplast were excluded, as well as

samples which had less than 5000 sequence counts. Shannon diversity was fit to a linear mixed-

effects model specifying *sample type* (invaded or control), *organism* (plant, mammal, mussel),

and their interactions as fixed factors. We included *studyID* or both *studyID* and *environment*

(soil or water) as random factors, and both models reported similar results (S2 Table). We

focused on the one with only *studyID* as random effect due to the lower AIC value. Models

were fit using the *lmer*() function under the *lme4* package [65] and the package *emmeans* was

used to infer pairwise contrasts (corrected using False Discovery Rate, FDR). Furthermore, we

explored the effects of *sample type* and *organism* on the structure of the microbial

communities using a multivariate approach. Distances between pairs of samples, in terms of

community composition, were calculated using a Unifrac matrix, and then visualized using an

RDA procedure. Differences between sample groups were inferred through PERMANOVA

multivariate analysis (999 permutations). We ran two different PERMANOVA models obtaining

similar results: in one we stratified permutations at level of *studyID* and *identity of invasive*

species, and in the other we stratified permutations at level of *studyID*, *environment* and

identity of invasive species, obtaining similar results (S3 Table). Pairwise contrasts from

PERMANOVA were subjected to FDR correction. Finally, the relative abundance of each

bacterial family was fit using the *lmer*() function to test the effects of *sample type* (invaded or

control) on individual taxa. We ran two different linear mixed-effects models, one including

studyID, *organism* (plant, mammal, mussel) and *environment* as random factors, and another

with *studyID* and *organism* as random factors, obtaining similar results (S4 Table).

Results

Our search yielded 5 studies with an appropriate experimental design and available

data, for a total of 356 samples. A few samples failed quality checks and we further considered

335 samples for downstream analyses. Sequences clustered into 22831 OTUs, after quality

checks, removal of singletons and "chloroplast" reads, with an average of 61776.92 reads per

sample. This high OTU count is likely a result of increased richness from analyzing samples

across multiple environments (soil and water) and from different geographical regions.

Biological invasions led to a reduction in Shannon diversity (χ^2= 3.85, df=1, *P*=0.04, Fig

1A). We also found biological invasions altered microbiome community composition in the

invaded environment compared to the control (Tab. 2, Fig 1B). The type of invasive organism

(plant, mammal, or mussel) produced a different community structure (pairwise *P*<0.01, FDR

corrected). A deeper analysis of bacterial families (Fig. 1C, S4 Table) revealed that some

taxonomic groups are significantly more abundant in invaded environments (Blastocatellaceae,

Chitinophagaceae, Nitrosomonadaceae, Pirellulaceae, Sphingomonadaceae), while others are

more abundant in non-invaded samples (Acetobacteraceae, Beijerinckiaceae, Gemmataceae,

Micromonosporaceae, Pedosphaeraceae, Solibacteraceae, Solirubrobacteraceae).

Table 2. Results from PERMANOVA analysis testing the effects of *sample type*

(invaded/control), *organism group* (plant, mammal, mussel) and their interaction on microbial

community composition. The factors *studyID* (unique for each study) was used as strata to

constrain permutations.

Factor	df	R^2	F	P
Sample type (Invaded/Control)	1	0.011	6.68	<0.001
Organism group (plant, mammal, mussel)	3	0.411	118.86	<0.001
Sample type * Organism group	3	0.007	2.1	0.01

Fig 1. (A) Comparison of Shannon index between control and invaded environments.

Data is represented as violin plot (showing the probability distribution of Shannon index for

each group) with a boxplot inside each violin plot indicating median and interquartile range. (B)

RDA ordination using a Bray-Curtis distance matrix of samples. (C) Comparison of the relative

abundance of microbial classes between control and invaded environments. * *P*<0.05, **

P<0.01, *** *P*<0.001

Discussion

Here we show biological invasions decrease the diversity of environmental microbiomes. While several studies have investigated the effects of species invasions on environmental microbiomes, we still lack a generalized consensus across different environmental microbiomes and systems. Previous studies have found that invasive species increased environmental microbial diversity [13,19], while others reported a decrease [15,20,29]. However, the majority of studies did not analyze the microbial diversity, as they used techniques that did not allow for such analysis, or reported no changes [9,10,23–28,30,11,12,14,16–18,21,22]. Within the studies included in our analysis, invasion by feral pigs decreased soil microbial diversity, while invasion by *Artemisia rothrockii* increased soil microbial diversity. The remaining three studies in our analysis reported no effects of biological invasions on environmental microbiome diversity. Our analysis was constrained in terms of sampled environment (soil) and invasive organism (plants), and an expanded dataset would be beneficial to generalize our results. Microbial diversity is tied to the function of microbiomes, and changes in diversity can reflect changes in function [66–68]. Changes in microbial diversity and function do not always have the same direction [69], which might explain the discrepancy between our results and other studies.

In contrast to diversity, our report of changes in community composition was relatively consistent with the published literature and the individual results of the studies we analyzed. Most studies of the influence of biological invasions on environmental microbiomes found that biological invasions alter environmental microbial community composition. However, some previous reports did not report changes [24–30], including the study by Gibbons et al. [28]

considered in our analysis. This variation may be due to individual effects of organisms on the

environment. For example, invasive plants may alter soil microbiome composition through root

exudates [5], and invasive mussels may alter water microbiome composition via bacterial

removal through their feeding activity [22]. Thus, reported influences on community

composition are more consistent. Alternatively changes in community composition might be

due to the response of some bacterial groups to environmental disturbance. The bacterial

families that we found to be differentially abundant between the invaded and control

environments have diverse ecological functions ranging from nitrogen fixation and

carbohydrate metabolism to antimicrobial properties. Many of the families that showed a

significant difference in abundance between invaded and control environments include taxa

that play important roles at various points during nitrogen and carbon cycling (i.e.

Nitrosomonadaceae, Acetobacteraceae, Chitinophagaceae, Micromonosporaceae,

Gemmataceae, Beijerinckiaceae, Pirellulaceae) [70,71,80,72–79]. However, nitrogen-fixing and

carbohydrate-degrading bacteria did not have a unified response to invaded environments as

some increased and others decreased in abundance in invaded environments. Many nitrogen-

fixing bacteria have been shown to respond to environmental disturbance, such as

Acidobacteria abundances during forest to pasture conversions or Pirellulaceae's response to

the presence of microplastics [81,82]. Thus, changes in environmental microbiome community

composition appear to be linked to changes in ecosystem functions, although this pattern is not

yet predictable across functions and taxa.

Few previous studies on biological invasions have reported details on the differential

abundance of taxa, and among these we found limited general consensus. For example, some

studies report a decrease in abundance of bacteria associated with nitrogen cycling (e.g.

Nitrosphaeria, Nitrospira, Nitrosomonadales) [12,13,18], while others report an increase of

Nitrosomonadaceae following invasion [30]. In our study some groups associated with the

nitrogen cycle were positively associated with biological invasions (i.e. Nitrosomonadaceae,

Pirellulaceae, Chitinophagaceae) while others were negatively associated (Beijerinckiaceae,

Micromonosporaceae). Unfortunately, amplicon-based sequencing has a limited power to infer

changes in the functions of microbiomes. Future metagenomic and metatranscriptomic studies

are needed to investigate whether biological invasions alter gene content or gene expression of

environmental microbiomes, and whether this reflects changes in biogeochemical cycling.

Meta-analyses are also useful to highlight gaps in the literature, and here we highlight

some aspects that warrant further investigation. First, we identified a large gap in the

availability of sequencing data from multiple types of environments and types of invasive

species. For our analysis almost all available data came from two environments: four sets of

data came from soil and one came from freshwater. Greater effort is needed for sample

collection from invasions in both freshwater and marine environments. Without sufficient

diversity of sample environments, it is impossible to tell whether microbial shifts following an

invasion are unique to an invaded environment. Second, in our analysis the majority of data

came from one type of invasive species: plants. Noticeably absent from our dataset were

invasions by insects, fish, and amphibians. Sequencing data is needed from a larger number of

invasive species to allow us to broadly assess shifts in microbial community structure. A third

gap we identified was the lack of spatial and temporal resolution. Almost all of the initially

identified 22 studies we assessed were also restricted to one season of sampling and were

conducted in the Northern Hemisphere. Thus, it is infeasible with existing datasets to validate

the influence of latitude or explore how seasonality and biological invasions interact to

modulate microbial communities. Thus, there are a number of opportunities for future research

on how biological invasions alter environmental microbial communities.

Here we analyzed 16S amplicon sequencing data from five studies and show that

biological invasions influence both the diversity and the structure of environmental

microbiomes. Understanding the impact of biological invasions on environmental microbiomes

is of high priority to preserve ecosystem functions [83]. We identified a number of gaps in our

knowledge, including the need to assess a wider range of environments, invasive species,

temporal variation, and latitudinal variation. We also demonstrate the power of re-analysis of

publicly available datasets using a common pipeline which benefited from open-data initiatives.

Acknowledgements

268 We would like to thank Kali Mattingly (Ohio State University, USA) and Davide Rassati

269 (Università degli Studi di Padova, Italy) for their helpful comments on the manuscript.

Data accessibility

271 Raw data from single studies is publicly available. The code used to perform our

272 analyses has been deposited on GitHub:

273 https://github.com/amalacrino/Biol_Invasion_Microb_MA

Supporting information

S1 Figure. PRISMA workflow. From: Moher D, Liberati A, Tetzlaff J, Altman DG, The

PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses:

The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

S2 Figure. Primers pairs considered in our analysis mapped to the structure of the 16S

rRNA gene that includes here just the hypervariable regions V1–V4.

S1 Table. PRIMA checklist. From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA

Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The

PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

S2 Table. Comparison of two different linear mixed-effects models testing the effect of

biological *sample type* (invaded or control), *organism* (plant, mammal, mussel), and their

interactions, on Shannon diversity index of the environmental microbiome. In Model 1 we

included *studyID* as random factor, while in Model 2 we included both *studyID* and *environment*

(soil or water) as random factors

S3 Table. Comparison of two different PERMANOVA models testing the effect of

biological *sample type* (invaded or control), *organism* (plant, mammal, mussel), and their

interactions, on the structure of the environmental microbiome. In Model 1 we included

studyID and *speciesID* as stratification factor, while in Model 2 we included *studyID*, *speciesID*

and *environment* (soil or water) as stratification factors.

S4 Table. Results from two different the linear mixed-effects model testing the

abundance of each bacterial family against *sample type* (invaded or control), *organism* (plant,

mammal, mussel), and their interactions. In Model 1 we included *studyID* as random factor,

while in Model 2 we included both *studyID* and *environment* (soil or water) as random factors.

References

1. Chapin III FS, Zavaleta ES, Eviner VT, Naylor RL, Vitousek PM, Reynolds HL, et al. Consequences of changing biodiversity. Nature. 2000;405: 234–242. doi:10.1038/35012241

2. Pejchar L, Mooney HA. Invasive species, ecosystem services and human well-being. Trends Ecol Evol. 2009;24: 497–504. doi:10.1016/j.tree.2009.03.016

3. Shackleton RT, Shackleton CM, Kull CA. The role of invasive alien species in shaping local livelihoods and human well-being: A review. J Environ Manage. 2019;229: 145–157. doi:10.1016/j.jenvman.2018.05.007

4. Charles H, Dukes JS. Impacts of Invasive Species on Ecosystem Services. Biological Invasions. Berlin, Heidelberg: Springer Berlin Heidelberg; 2008. pp. 217–237. doi:10.1007/978-3-540-36920-2_13

5. Weidenhamer JD, Callaway RM. Direct and Indirect Effects of Invasive Plants on Soil Chemistry and Ecosystem Function. J Chem Ecol. 2010;36: 59–69. doi:10.1007/s10886-009-9735-0

6. Lee MR, Bernhardt ES, van Bodegom PM, Cornelissen JHC, Kattge J, Laughlin DC, et al. Invasive species' leaf traits and dissimilarity from natives shape their impact on nitrogen cycling: a meta-analysis. New Phytol. 2017;213: 128–139. doi:10.1111/nph.14115

7. Arthur MA, Bray SR, Kuchle CR, McEwan RW. The influence of the invasive shrub, Lonicera maackii, on leaf decomposition and microbial community dynamics. Plant Ecol. 2012;213: 1571–1582. doi:10.1007/s11258-012-0112-7

8. Hall Jr RO, Tank JL, Dybdahl MF. Exotic snails dominate nitrogen and carbon cycling in a

highly productive stream. Front Ecol Environ. 2003;1: 407–411. doi:10.1890/1540-

9295(2003)001[0407:ESDNAC]2.0.CO;2

9.   Lazzaro L, Giuliani C, Fabiani A, Agnelli AE, Pastorelli R, Lagomarsino A, et al. Soil and

plant changing after invasion: The case of Acacia dealbata in a Mediterranean ecosystem.

Sci Total Environ. 2014;497–498: 491–498. doi:10.1016/j.scitotenv.2014.08.014

10.   Malinich E, Lynn-Bell N, Kourtev PS. The effect of the invasive Elaeagnus umbellata on

soil microbial communities depends on proximity of soils to plants. Ecosphere. 2017;8:

e01827. doi:10.1002/ecs2.1827

11.   Lorenzo P, Pereira CS, Rodríguez-Echeverría S. Differential impact on soil microbes of

allelopathic compounds released by the invasive Acacia dealbata Link. Soil Biol Biochem.

2013;57: 156–163. doi:10.1016/j.soilbio.2012.08.018

12.   Parsons LS, Sayre J, Ender C, Rodrigues JLM, Barberán A. Soil microbial communities in

restored and unrestored coastal dune ecosystems in California. Restor Ecol. 2020;

rec.13101. doi:10.1111/rec.13101

13.   Piper CL, Siciliano SD, Winsley T, Lamb EG. Smooth brome invasion increases rare soil

bacterial species prevalence, bacterial species richness and evenness. De Deyn G, editor.

J Ecol. 2015;103: 386–396. doi:10.1111/1365-2745.12356

14.   Zhang H-Y, Goncalves P, Copeland E, Qi S-S, Dai Z-C, Li G-L, et al. Invasion by the weed

Conyza canadensis alters soil nutrient supply and shifts microbiota structure. Soil Biol

Biochem. 2020;143: 107739. doi:10.1016/j.soilbio.2020.107739

15.   Le Roux JJ, Ellis AG, van Zyl L-M, Hosking ND, Keet J-H, Yannelli FA. Importance of soil

legacy effects and successful mutualistic interactions during Australian acacia invasions in

nutrient-poor environments. Aerts R, editor. J Ecol. 2018;106: 2071–2081.

doi:10.1111/1365-2745.12965

16. Brunel C, Beifen Y, Pouteau R, Li J, van Kleunen M. Responses of Rhizospheric Microbial

Communities of Native and Alien Plant Species to Cuscuta Parasitism. Microb Ecol.

2020;79: 617–630. doi:10.1007/s00248-019-01438-z

17. Pickett B, Irvine IC, Bullock E, Arogyaswamy K, Aronson E. Legacy effects of invasive grass

impact soil microbes and native shrub growth. Invasive Plant Sci Manag. 2019;12: 22–35.

doi:10.1017/inp.2018.32

18. Rodrigues RR, Pineda RP, Barney JN, Nilsen ET, Barrett JE, Williams MA. Plant Invasions

Associated with Change in Root-Zone Microbial Community Structure and Diversity. Liu J,

editor. PLoS One. 2015;10: e0141424. doi:10.1371/journal.pone.0141424

19. Collins CG, Carey CJ, Aronson EL, Kopp CW, Diez JM. Direct and indirect effects of native

range expansion on soil microbial community structure and function. Austin A, editor. J

Ecol. 2016;104: 1271–1283. doi:10.1111/1365-2745.12616

20. Wehr NH, Kinney KM, Nguyen NH, Giardina CP, Litton CM. Changes in soil bacterial

community diversity following the removal of invasive feral pigs from a Hawaiian tropical

montane wet forest. Sci Rep. 2019;9: 14681. doi:10.1038/s41598-019-48922-7

21. Boeker C, Geist J. Effects of invasive and indigenous amphipods on physico-chemical and

microbial properties in freshwater substrates. Aquat Ecol. 2015;49: 467–480.

doi:10.1007/s10452-015-9539-y

22. Denef VJ, Carrick HJ, Cavaletto J, Chiang E, Johengen TH, Vanderploeg HA. Lake Bacterial

Assemblage Composition Is Sensitive to Biological Disturbance Caused by an Invasive

Filter Feeder. Kent AD, editor. mSphere. 2017;2. doi:10.1128/mSphere.00189-17

23.   Manzari C, Fosso B, Marzano M, Annese A, Caprioli R, D'Erchia AM, et al. The influence of

370   invasive jellyfish blooms on the aquatic microbiome in a coastal lagoon (Varano, SE Italy)

371   detected by an Illumina-based deep sequencing strategy. Biol Invasions. 2015;17: 923–

372   940. doi:10.1007/s10530-014-0810-2

373   24.   Lazzaro L, Mazza G, D'Errico G, Fabiani A, Giuliani C, Inghilesi AF, et al. How ecosystems

374   change following invasion by Robinia pseudoacacia: Insights from soil chemical

375   properties and soil microbial, nematode, microarthropod and plant communities. Sci

376   Total Environ. 2018;622–623: 1509–1518. doi:10.1016/j.scitotenv.2017.10.017

377   25.   Hernández-Gómez O, Byrne AQ, Gunderson AR, Jenkinson TS, Noss CF, Rothstein AP, et

378   al. Invasive vegetation affects amphibian skin microbiota and body condition. PeerJ.

379   2020;8: e8549. doi:10.7717/peerj.8549

380   26.   Thompson GL, Bell TH, Kao-Kniffin J. Rethinking Invasion Impacts across Multiple Field

381   Sites Using European Swallowwort (Vincetoxicum rossicum) as a Model Invader.

382   Invasive Plant Sci Manag. 2018;11: 109–116. doi:10.1017/inp.2018.22

383   27.   Carey CJ, Beman JM, Eviner VT, Malmstrom CM, Hart SC. Soil microbial community

384   structure is unaltered by plant invasion, vegetation clipping, and nitrogen fertilization in

385   experimental   semi-arid   grasslands.   Front   Microbiol.   2015;6.

386   doi:10.3389/fmicb.2015.00466

387   28.   Gibbons SM, Lekberg Y, Mummey DL, Sangwan N, Ramsey PW, Gilbert JA. Invasive Plants

388   Rapidly Reshape Soil Properties in a Grassland Ecosystem. Shade A, editor. mSystems.

389   2017;2. doi:10.1128/mSystems.00178-16

29. Gomes SIF, Merckx VSFT, Hynson NA. Biological invasions increase the richness of
arbuscular mycorrhizal fungi from a Hawaiian subtropical ecosystem. Biol Invasions.
2018;20: 2421–2437. doi:10.1007/s10530-018-1710-7

30. de Menezes AB, Prendergast-Miller MT, Macdonald LM, Toscas P, Baker G, Farrell M, et
al. Earthworm-induced shifts in microbial diversity in soils with rare versus established
invasive   earthworm   populations.   FEMS   Microbiol   Ecol.   2018;94.
doi:10.1093/femsec/fiy051

31. Bier RL, Bernhardt ES, Boot CM, Graham EB, Hall EK, Lennon JT, et al. Linking microbial
community structure and microbial processes: an empirical and conceptual overview.
Muyzer G, editor. FEMS Microbiol Ecol. 2015;91: fiv113. doi:10.1093/femsec/fiv113

32. Gibbons SM, Gilbert JA. Microbial diversity — exploration of natural ecosystems and
microbiomes. Curr Opin Genet Dev. 2015;35: 66–72. doi:10.1016/j.gde.2015.10.003

33. Gurevitch J, Koricheva J, Nakagawa S, Stewart G. Meta-analysis and the science of
research synthesis. Nature. 2018;555: 175–182. doi:10.1038/nature25753

34. Nakagawa S, Poulin R. Meta-analytic insights into evolutionary ecology: an introduction
and synthesis. Evol Ecol. 2012;26: 1085–1099. doi:10.1007/s10682-012-9593-z

35. Nakagawa S, Santos ESA. Methodological issues and advances in biological meta-analysis.
Evol Ecol. 2012;26: 1253–1274. doi:10.1007/s10682-012-9555-5

36. Senior AM, Grueber CE, Kamiya T, Lagisz M, O'Dwyer K, Santos ESA, et al. Heterogeneity
in ecological and evolutionary meta-analyses: its magnitude and implications. Ecology.
2016;97: 3293–3299. doi:10.1002/ecy.1591

37. Lozupone CA, Stombaugh J, Gonzalez A, Ackermann G, Wendel D, Vazquez-Baeza Y, et al.

Meta-analyses of studies of the human microbiota. Genome Res. 2013;23: 1704–1714.

doi:10.1101/gr.151803.112

38. Walters WA, Xu Z, Knight R. Meta-analyses of human gut microbes associated with

obesity and IBD. FEBS Lett. 2014;588: 4223–4233. doi:10.1016/j.febslet.2014.09.039

39. Duvallet C, Gibbons SM, Gurry T, Irizarry RA, Alm EJ. Meta-analysis of gut microbiome

studies identifies disease-specific and shared responses. Nat Commun. 2017;8: 1784.

doi:10.1038/s41467-017-01973-8

40. Koren O, Knights D, Gonzalez A, Waldron L, Segata N, Knight R, et al. A Guide to

Enterotypes across the Human Body: Meta-Analysis of Microbial Community Structures

in Human Microbiome Datasets. Eisen JA, editor. PLoS Comput Biol. 2013;9: e1002863.

doi:10.1371/journal.pcbi.1002863

41. Krych L, Hansen CHF, Hansen AK, van den Berg FWJ, Nielsen DS. Quantitatively Different,

yet Qualitatively Alike: A Meta-Analysis of the Mouse Core Gut Microbiome with a View

towards the Human Gut Microbiome. Bereswill S, editor. PLoS One. 2013;8: e62578.

doi:10.1371/journal.pone.0062578

42. Wang J, Kurilshikov A, Radjabzadeh D, Turpin W, Croitoru K, Bonder MJ, et al. Meta-

analysis of human genome-microbiome association studies: the MiBioGen consortium

initiative. Microbiome. 2018;6: 101. doi:10.1186/s40168-018-0479-3

43. Jiao N, Baker SS, Nugent CA, Tsompana M, Cai L, Wang Y, et al. Gut microbiome may

contribute to insulin resistance and systemic inflammation in obese rodents: a meta-

analysis. Physiol Genomics. 2018;50: 244–254. doi:10.1152/physiolgenomics.00114.2017

44. Wirbel J, Pyl PT, Kartal E, Zych K, Kashani A, Milanese A, et al. Meta-analysis of fecal

metagenomes reveals global microbial signatures that are specific for colorectal cancer.

Nat Med. 2019;25: 679–689. doi:10.1038/s41591-019-0406-6

45.   Bisanz JE, Upadhyay V, Turnbaugh JA, Ly K, Turnbaugh PJ. Meta-Analysis Reveals

Reproducible Gut Microbiome Alterations in Response to a High-Fat Diet. Cell Host

Microbe. 2019;26: 265-272.e4. doi:10.1016/j.chom.2019.06.013

46.   Adams RI, Bateman AC, Bik HM, Meadow JF. Microbiota of the indoor environment: a

meta-analysis. Microbiome. 2015;3: 49. doi:10.1186/s40168-015-0108-3

47.   Debroas D, Domaizon I, Humbert J-F, Jardillier L, Lepère C, Oudart A, et al. Overview of

freshwater microbial eukaryotes diversity: a first analysis of publicly available

metabarcoding data. FEMS Microbiol Ecol. 2017;93. doi:10.1093/femsec/fix023

48.   Shade A, Gregory Caporaso J, Handelsman J, Knight R, Fierer N. A meta-analysis of

changes in bacterial and archaeal communities with time. ISME J. 2013;7: 1493–1506.

doi:10.1038/ismej.2013.54

49.   Rocca JD, Simonin M, Blaszczak JR, Ernakovich JG, Gibbons SM, Midani FS, et al. The

Microbiome Stress Project: Toward a Global Meta-Analysis of Environmental Stressors

and Their Effects on Microbial Communities. Front Microbiol. 2019;9.

doi:10.3389/fmicb.2018.03272

50.   Duvallet C. Meta-analysis generates and prioritizes hypotheses for translational

microbiome research. Microb Biotechnol. 2018;11: 273–276. doi:10.1111/1751-

7915.13047

51.   Drewes JL, White JR, Dejea CM, Fathi P, Iyadorai T, Vadivelu J, et al. High-resolution

bacterial 16S rRNA gene profile meta-analysis and biofilm status reveal common

colorectal cancer consortia. npj Biofilms Microbiomes. 2017;3: 34. doi:10.1038/s41522-

017-0040-3

52.   Ramirez KS, Knight CG, de Hollander M, Brearley FQ, Constantinides B, Cotton A, et al.

Detecting macroecological patterns in bacterial communities across independent studies

of global soils. Nat Microbiol. 2018;3: 189–196. doi:10.1038/s41564-017-0062-x

53.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al.

Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.

Proc Natl Acad Sci. 2011;108: 4516–4522. doi:10.1073/pnas.1000080107

54.   Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of

general 16S ribosomal RNA gene PCR primers for classical and next-generation

sequencing-based diversity studies. Nucleic Acids Res. 2013;41: e1–e1.

doi:10.1093/nar/gks808

55.   Magoc T, Salzberg SL. FLASH: fast length adjustment of short reads to improve genome

assemblies. Bioinformatics. 2011;27: 2957–2963. doi:10.1093/bioinformatics/btr507

56.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-

throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms.

ISME J. 2012;6: 1621–1624. doi:10.1038/ismej.2012.8

57.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source tool

for metagenomics. PeerJ. 2016;4: e2584. doi:10.7717/peerj.2584

58.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal

RNA gene database project: improved data processing and web-based tools. Nucleic

Acids Res. 2012;41: D590–D596. doi:10.1093/nar/gks1219

59.   Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Mol Biol Evol. 2013;30: 772–780.

doi:10.1093/molbev/mst010

60.   Price MN, Dehal PS, Arkin AP. FastTree: Computing Large Minimum Evolution Trees with

Profiles instead of a Distance Matrix. Mol Biol Evol. 2009;26: 1641–1650.

doi:10.1093/molbev/msp077

61.   R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R

Foundation for Statistical Computing; 2016. 2017.

62.   McMurdie PJ, Holmes S. phyloseq: An R package for reproducible interactive analysis and

graphics of microbiome census data. Watson M, editor. PLoS One. 2013;8: e61217.

doi:10.1371/journal.pone.0061217

63.   Dixon P. VEGAN, a package of R functions for community ecology. J Veg Sci. 2003;14:

927–930. doi:10.1111/j.1654-1103.2003.tb02228.x

64.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014;15: 550. doi:10.1186/s13059-014-0550-8

65.   Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. J

Stat Softw. 2015;67: 1–48. doi:10.18637/jss.v067.i01

66.   Delgado-Baquerizo M, Maestre FT, Reich PB, Jeffries TC, Gaitan JJ, Encinar D, et al.

Microbial diversity drives multifunctionality in terrestrial ecosystems. Nat Commun.

2016;7: 10541. doi:10.1038/ncomms10541

67.   Miki T, Yokokawa T, Matsui K. Biodiversity and multifunctionality in a microbial

community: a novel theoretical approach to quantify functional redundancy. Proc R Soc B

Biol Sci. 2014;281: 20132498. doi:10.1098/rspb.2013.2498

68.   Philippot L, Spor A, Hénault C, Bru D, Bizouard F, Jones CM, et al. Loss in microbial

diversity affects nitrogen cycling in soil. ISME J. 2013;7: 1609–1619.

doi:10.1038/ismej.2013.34

69.   Mendes LW, Tsai SM, Navarrete AA, de Hollander M, van Veen JA, Kuramae EE. Soil-

Borne Microbiome: Linking Diversity to Function. Microb Ecol. 2015;70: 255–265.

doi:10.1007/s00248-014-0559-2

70.   Chen J, Mo L, Zhang Z, Nan J, Xu D, Chao L, et al. Evaluation of the ecological restoration

of a coal mine dump by exploring the characteristics of microbial communities. Appl Soil

Ecol. 2020;147: 103430. doi:10.1016/j.apsoil.2019.103430

71.   Cheng J, Chen Y, He T, Liao R, Liu R, Yi M, et al. Soil nitrogen leaching decreases as biogas

slurry DOC/N ratio increases. Appl Soil Ecol. 2017;111: 105–113.

doi:10.1016/j.apsoil.2016.12.001

72.   Kämpfer P, Lodders N, Falsen E. Hydrotalea flava gen. nov., sp. nov., a new member of

the phylum Bacteroidetes and allocation of the genera Chitinophaga, Sediminibacterium,

Lacibacter, Flavihumibacter, Flavisolibacter, Niabella, Niastella, Segetibacter,

Parasegetibacter, Terrimonas, Fer. Int J Syst Evol Microbiol. 2011;61: 518–523.

doi:10.1099/ijs.0.023002-0

73.   Madhaiyan M, Poonguzhali S, Senthilkumar M, Pragatheswari D, Lee J-S, Lee K-C.

Arachidicoccus rhizosphaerae gen. nov., sp. nov., a plant-growth-promoting bacterium in

the family Chitinophagaceae isolated from rhizosphere soil. Int J Syst Evol Microbiol.

2015;65: 578–586. doi:10.1099/ijs.0.069377-0

74. Reis VM, Teixeira KR dos S. Nitrogen fixing bacteria in the family Acetobacteraceae and

523    their role in agriculture. J Basic Microbiol. 2015;55: 931–949.

524    doi:10.1002/jobm.201400898

525 75. Trujillo ME, Bacigalupe R, Pujic P, Igarashi Y, Benito P, Riesco R, et al. Genome Features of

526    the Endophytic Actinobacterium Micromonospora lupini Strain Lupac 08: On the Process

527    of Adaptation to an Endophytic Life Style? Brüggemann H, editor. PLoS One. 2014;9:

528    e108522. doi:10.1371/journal.pone.0108522

529 76. Carro L, Nouioui I, Sangal V, Meier-Kolthoff JP, Trujillo ME, Montero-Calasanz M del C, et

530    al. Genome-based classification of micromonosporae with a focus on their

531    biotechnological and ecological potential. Sci Rep. 2018;8: 525. doi:10.1038/s41598-017-

532    17392-0

533 77. Dedysh SN, Ivanova AA. Planctomycetes in boreal and subarctic wetlands: diversity

534    patterns and potential ecological functions. FEMS Microbiol Ecol. 2019;95.

535    doi:10.1093/femsec/fiy227

536 78. Wegner C-E, Gorniak L, Riedel S, Westermann M, Küsel K. Lanthanide-Dependent

537    Methylotrophs of the Family Beijerinckiaceae : Physiological and Genomic Insights.

538    Stams AJM, editor. Appl Environ Microbiol. 2019;86. doi:10.1128/AEM.01830-19

539 79. Neufeld JD, Schäfer H, Cox MJ, Boden R, McDonald IR, Murrell JC. Stable-isotope probing

540    implicates Methylophaga spp and novel Gammaproteobacteria in marine methanol and

541    methylamine metabolism. ISME J. 2007;1: 480–491. doi:10.1038/ismej.2007.65

542 80. Warneke C, Karl T, Judmaier H, Hansel A, Jordan A, Lindinger W, et al. Acetone,

543    methanol, and other partially oxidized volatile organic emissions from dead plant matter

by abiological processes: Significance for atmospheric HO x chemistry. Global

Biogeochem Cycles. 1999;13: 9–17. doi:10.1029/98GB02428

81.   Miao L, Wang P, Hou J, Yao Y, Liu Z, Liu S, et al. Distinct community structure and

microbial functions of biofilms colonizing microplastics. Sci Total Environ. 2019;650:

2395–2402. doi:10.1016/j.scitotenv.2018.09.378

82.   Navarrete AA, Venturini AM, Meyer KM, Klein AM, Tiedje JM, Bohannan BJM, et al.

Differential Response of Acidobacteria Subgroups to Forest-to-Pasture Conversion and

Their Biogeographic Patterns in the Western Brazilian Amazon. Front Microbiol. 2015;6.

doi:10.3389/fmicb.2015.01443

83.   Doblas-Miranda E, Martínez-Vilalta J, Lloret F, Álvarez A, Ávila A, Bonet FJ, et al.

Reassessing global change research priorities in mediterranean terrestrial ecosystems:

how far have we come and where do we go from here? Glob Ecol Biogeogr. 2015;24: 25–

43. doi:10.1111/geb.12224