Ten Genetic Loci Identified for Milk Yield, Fat, and Protein in Holstein Cattle

Liyuan Liu[1,2#], Jinghang Zhou[1,2#], Chunpeng James Chen[2#], Juan Zhang[1], Wan Wen[3], Jia Tian[3], Zhiwu Zhang[2], Yaling Gu[1*]*

[1]School of Agriculture, Ningxia University, Yinchuan, Ningxia, China, [2]Department of Crop and Soil Sciences, Washington State University, Pullman, Washington, USA, [3]Animal Husbandry of Extension Station, Yinchuan, Ningxia, China.

[#]These authors contributed equally to this work.

[*]Correspondence should be addressed to Zhiwu Zhang (email: zhiwu.zhang@wsu.edu) or Yaling Gu (email: guyl@nxu.edu.cn)

Abstract

The production of high-yield and high-quality milk is the direct economic goals of dairy industry breeding. Understanding the genetic architecture underlying milk-related traits is beneficial so that genetic variants can be targeted toward the overall improvement. In this study, we measured five traits related to milk production and components in Holstein cattle population from China. These traits included milk yield, protein yield, and fat yields; fat percentage and protein percentages. We used the estimated breeding values as dependent variables to conduct the genome-wide association study (GWAS). Breeding values were estimated through pedigree relationships by using a mixed linear model for individuals with and without phenotypic data. Genotyping was carried out on the individuals with phenotypes by using the Illumina BovineSNP150 BeadChip. The association analyses were conducted by using the Fixed and

random model Circulating Probability Unification (FarmCPU) method. A total of 10 SNPs was detected above the genome-wide significant threshold, six were located in previously reported QTL regions. We also found 8 candidate genes each is within a distance of 120 kb upstream and downstream to an associated SNPs, the most significant SNP affecting milk fat and protein percentage is in the *DGAT1* gene. These genetic variants and potential candidate genes would be valuable resources to enhance dairy cattle breeding.

Keywords: Dairy cattle, GWAS, milk traits, *DGAT1*

Introduction

Traditional dairy breeding mainly depends on pedigree and phenotype information that is expensive, labor-intensive, and time-consuming. For example, milk is not available until a cow turns three years old. In addition, dairy cows have a long generation interval and give birth only once a year. It was demonstrated that implementation of genomic selection for young increased genetic gain [1] and marker-assisted selection also had the potential to enhance genetic improvement [2,3]. The production of high-yield and high-quality milk is the direct economic goals of dairy breeding. The premise for the potential success of molecular breeding is the ability to identify major genes associated with important dairy milk-related traits.

Genome-wide association studies (GWAS) have proven to be a powerful method for identifying potential genetic variants, especially single-nucleotide polymorphisms (SNPs) associated with complex diseases in humans and important economic traits in animals [4,5]. A large number of association studies reveals numerous QTLs for milk-related traits in Holstein population around the world (CatttleQTLdb: https://www.animalgenome.org/cgi-bin/QTLdb/BT/index), Pausch et

al identified a total of 25 QTL for milk fat and protein using meta-analysis across three cattle breeds [6]. Although in 2010, Jiang et al revealed 105 SNPs associated with milk production traits using Illumina BovineSNP50 BeadChip in Chinese Holstein population [7], what's different from this study is that we used a higher density chips and another Holstein population. Genotype at high density provide markers close to the QTL and therefore in stronger linkage disequilibrium (LD) with QTL [8], and help in fine mapping of candidate QTLs.

In this study, we conducted GWAS using Illumina BovineSNP150 BeadChip and phenotypic data from Holstein cows raised in the Ningxia region of northwest China. Our objectives were to identify new genetic variants associated with five milk traits (milk yield, fat yield, protein yield, fat percentage, and protein percentage), We expect the newly identified genetic variants and potential candidate genes would become valuable resources for breeding.

Material and methods

Population structure and phenotypic data

The study population is Holstein cows that were raised on 21 dairy farms in the Ningxia area of China. We used estimated breeding values (EBVs) as phenotypes implement association analysis, Milk yield (MY), fat yield (FY), protein yield (PY), fat percentage (FP), protein percentage (PP) measurements were recorded once a month for each cow after calving and the start of lactation. The milk yield was automatically recorded by the milking system in each farm, the milk components are tested by the Dairy Herd Improvement lab at Animal Husbandry of Extension Station in Ningxia, using spectrometers. FY was calculated as (FP*MY)/100; PY was calculated as (PP*MY)/100. The distribution of phenotypes and correlations between the different phenotypic traits are illustrated in **Figures** S1, S2.

Estimated breeding values

The DMU package (MUItivariate analysis based on a Derivative-free approach) [9] was used to estimate breeding values using Random Regression Test-Day Model [10,11]. Totally about 452,920 test-day records from 61,600 cows spanning a 9-year period (2011-2019) at their first lactation stage, all recorded data reflect the first lactation for each cow. We considered herd-test-day and calving year-season as fixed effects, calving month-age as a fixed regression effect, and individual additive effect and permanent environment effect as random regression effects. Fixed regression using a fourth order Legendre polynomial. The model equation is as follows:

$$y_{ijklm} = HTD_i + cays_j + \sum_{m=1}^{5} b_{km}X_{bm}(\omega) + \sum_{m=1}^{r} a_{lm}Z_{am}(\omega) + \sum_{m=1}^{s} p_{lm}W_{pm}(\omega) + e_{ijklm}$$

Where y_{ijklm} is the test-day records m of cow ; HTD_i is the fixed effect of the i th herd-test day; $cays_j$ is the fixed effect of the j th combined of calving years and season; b_{km} is the m th fixed regression coefficient of the k th calving year-season , X_{bm} is the Legendre polynomial m th covariable; a_{lm} is the m th random regression coefficient under additive genetic effects, corresponds to the l th cow in the pedigree; p_{lm} is the m th random regression coefficient under the permanent environment genetic effects, correspond to the l th cow in the data file; Z_{am} and W_{pm} are the m th covariance elements for additive effects and permanent environment effects; ω is the days of lactation after standardization; and e_{ijklm} is the residual effects.

Genotypic data

Blood samples from the 1,220 cows were collected by cattle farm staff in this study. DNA was extracted and Genotyping was carried out by Compass Biotechnology (http://www.kangpusen.com/) using the Illumina BovineSNP150 BeadChip. We used genome

reference the Bos_taurus_UMD_3.1. In total, there were 124,743 variants for the association

analysis after conducted quality control by Plink software [12]. Markers were removed if (1) the

call rate of an individual genotype was less than 95%, (2) the call rate of a single SNP genotype

was less than 90%, and (3) if the MAF of a SNP was less than 0.05 and deviated from Hardy-

Weinberg equilibrium ($p < 1.0e-6$). We calculated marker intervals and linkage disequilibrium

(LD) to estimate R square for all markers and plotted the marker distribution in **Figure** 1.

Association analysis

We performed the single-SNP analysis for the markers using the FarmCPU (Fixed and random

model Circuitous Probability Unification) [13]. FarmCPU can consider a population structure

matrix as a covariate variable to correct false positive results caused by population stratification

[13]. Considering the stratification in the dairy cow population, we fit the first three principal

components as covariate variables in the GWAS models.

Annotation of candidate gene and pathway analysis

The genome reference Bos_taurus_UMD_3.1 was used to search candidate gene. The online

website "https://oct2018.archive.ensembl.org/Bos_taurus/Info/Index",

"https://www.ncbi.nlm.nih.gov/gene/ ", "https://www.genome.jp/kegg/pathway.html" were used

for functional analysis and pathway analysis of the candidate gene by GWAS.

Results

Marker information

We conducted GWAS analyses with 1,220 Holstein dairy cows and 124,743 markers after quality control (QC). Markers covered all 29 autosomes plus the X sex chromosome (**Figure 1**). After the QC filtering, we re-calculated the minor allele frequency (MAF) for all SNPs. The minimum MAF was 0.038. Marker density was high when marker interval was less than 20 kb and LD was strong within a distance of 10 kb.

Population structure

The 1,220 individuals included in the study population came from 21 different farms. To determine the level of population stratification, we plotted the population structure by principal component analysis (PCA). The population stratified into two unevenly sized groups (**Figure 2**). To adjust for the population stratification, we fit the principal components (PCs) as covariate variables in the association analysis. We also made a scatter plot between the first three principal components and the ten traits, there are no correlation observed between the PCA and these phenotypes (**Figure** S3).

Phenotypic and estimated genetic parameters

Phenotype distributions and correlations among phenotypic traits, estimated breeding values, and residuals are shown in the supplementary material (**Figure** S1). We found strong positive phenotypic correlations between MY and PY ($r_p = 0.90$), MY and FY ($r_p = 0.70$), and PY and FY ($r_p = 0.74$). We also found strong positive genetic correlations between MY and PY ($r_g = 0.92$), MY and FY ($r_g = 0.84$), and PY and FY ($r_g = 0.88$). In contrast, there were weak negative phenotypic and genetic correlations between MY and FP ($r_p = -0.15$, $r_g = -0.32$), MY and PP ($r_p = -0.20$, $r_g = -0.44$).

In this study, we used EBVs as the dependent variables for GWAS, **Figure** S2 showed that the heatmap of EBVs for five milk traits. We used test-day records to estimate the heritability and breeding values of MY, FY, and PY over a 305-day lactation period. For the heritability and breeding values of FP and PP, the 305-day averages were estimated. The heritability estimates for MY, FY, PY, FP and PP were 0.12, 0.21, 0.23, 0.30 and 0.32, respectively (**Table** 1).

Results of the genome-wide associations with milk traits

By drawing the Quantile-Quantile (QQ) plots, we found that the model for GWAS analysis in this study was reasonable, and the point at the upper right corner also shown that some significant markers were found that associated with four milk traits (**Figure** 3). We found a total of ten highly significant SNPs associated with FP (3SNPs), FY (4 SNPs), PP (3 SNPs) and PY (1 SNP) (**Table** 2, **Figure** 3). We did not find threshold significant SNPs associated with MY. We used $P < 4.0e-07$ as threshold, which was corresponding to 1% of type I error after Bonferroni multiple test correction. The ten significant SNPs are located on Bos_taurus autosomes (BTAs) 1, 5, 7, 8, 11, 14, 17, 21 and X chromosome (**Table** 3, **Figure** 3). The three SNPs (rs42295213, rs136949224 and rs109421300) associated with FP are located on BTA1, 8 and 14, four SNPs (rs43526055, rs137676276, rs109528658 and rs135780687) associated with FY are located on BTA7, 11, 17, and X, respectively. The three SNPs (rs109875012, rs109421300 and rs108996837) associated with PP are located on BTA5, 14, and 21, respectively. One SNP associated with PY is located on BTA5. Four of these ten significant SNPs are located inside genes *EPHA6* (EPH receptor A6), *SLCO1A2* (solute carrier organic anion transporter family member 1A2), *DGAT1* (diacylglycerol O-acyltransferase 1) *and EP400* (E1A binding protein p400). The SNP (rs109875012) on BTA5 located close to the *ZNF384* (zinc finger protein 384).

The SNP (rs10705865) on BTA8 located close to *SCARA5* (scavenger receptor class A member 5) gene. The SNP (rs137676276) on BTA 11 located close to *VIT* (vitrin). The SNP (rs108996837) on BTA 21 located close to *EXOC3L4* (exocyst complex component 3 like 4), and the SNP (rs135780687) on X chromosome located close to *GRPR* (gastrin releasing peptide receptor). The most significant SNP (rs109421300) associated with both FP and PP located in the *DGAT1* gene. Two SNPs (rs137676276, rs108996837) exhibit notably smaller MAFs compared to other SNPs, 0.11 and 0.12, respectively (**Table** 2).

Pleiotropic QTLs for milk production traits

We used the SNPs with $P < 0.0005$ to make heatmap to look for markers associated with two or more milk production traits (**Figure** S4). We expected to find pleiotropic QTLs because these milk traits are moderately or highly correlated (**Figure** S1, S2). QTL on BTA1 at 0.4Mb, on BTA2 at 14.8Mb, on BTA2 at 134.3Mb, on BTA3 at 64.4Mb, on BTA11 at 39 Mb, on BTA17 at 29−32.9Mb, on BTA20 at 67.8 Mb, BTA22 at 54.6Mb associated with three milk production traits (MY, PY, and FY). QTL on BTA1 at 40.3-41.1Mb, on BTA6 at 37.7Mb, on BTA14 at 73-77Mb, on BTA17 at 29-32Mb associated with MY, PY. QTL on BTA20 at 24Mb associated with FY and FY.

Discussion

Population structure

Population stratification is an important confounding factor due to systematic ancestry differences that can cause false positives in GWAS [14]. Partially in cattle population, not all the registered cattle are the full blood. This point can be explained by looking at the cattle breed

registration requirements in different countries. Here we take the Holstein cattle as an example, one of the requirements to register a Holstein cattle in China is that the cattle at least has 87.5% blood of Holstein (Chinese Holstein, GB/T 3157 2008), all the animals with Holstein genetics can be registered in Canadian (https://www.holstein.ca/Public/en/Services/Registration/Registration_Eligibilities) and there are the similar clause in USA (http://www.holsteinusa.com/animal_id/register.html). According to the above standards we can see the common ground to register a Holstein cattle between different countries is that not all the Holstein cattle are the full blood. Even most of the Holstein cattle are pure breed, some of registered cattle could still contains a little other blood in the long-term breeding progress. That's why the population structure analysis is necessary in this study. By the principle component analysis, the PCA scatter plot showed that there are two groups in the population in this study; that is, a small number of individuals divided into the other group (**Figure** 2). There are probably two reasons caused the deviation of the population structure, the first reason is the semen the cattle farm used are different, as we know most of the farms in this study participated in a dairy breeding project that introduce of some Holstein frozen semen overseas annually from 2013 to 2018. Another reason is that some cows are introduced from different countries there might has some various background.

As we observed the population structure, the principle components were fitted as covariance do association analysis to correcting population stratification. After adjusted PCA factors, there are four SNPs are overlapped with association model not fitted PCs, will discuss that later.

GWAS for milk production traits

Milk yield and composition are very important economic traits in dairy cattle because good milk production performance can bring greater economic benefits. As early as 1994, a research

identified a QTL significantly associated with FY was linked to kappa-casein and a QTL for PY was linked to beta-lactoglobulin [15]. Subsequently, a growing number of studies detected thousands and tens of QTLs through the 30 chromosomes associated with 646 different traits in cattle (Cattle QTLdb). In this study, there were no significant SNPs passed the Bonferroni correction threshold for MY, but ten SNPs were significantly associated with the other four milk traits (FP, FY, PP and PY).

Six of these significant SNPs are located within previously reported QTL regions, and four SNPs (rs42295213, rs134480235, rs109421300, and rs109528658) located within four genes, *EPHA6*, *SLCO1A2*, *DGAT1*, *EP400*. More precisely, three SNPs (rs42295213, rs136949224, rs109421300) on BTA1, 8, 14 were associated with FP, they are within reported QTLs [16–19]. The SNP on BTA1 is in the *EPHA6* gene, which have functions of ATP binding (GO:0005524), protein binding (GO: 0005515), protein tyrosine kinase activity (GO: 0004713), and has been proposed to participate in Axon guidance pathway (KEGG: bta04360). The SNP on BTA8 located close to *SCARA5* gene, this gene is a member of the scavenger receptor (SR) family, which is broad expression in fat tissue in human individuals, research showed that scavenger receptors involved in lipid accumulation and inflammation [20], some studies reported the gene plays a critical role in progression and metastasis of breast cancer [21], and involved in breast carcinogenesis [22]. *SCARA5* gene may be a potential candidate gene involved in the biosynthesis of milk fat. The SNP on BTA14 associated both with FP and PP, which is in the *DGAT1* gene. As we know, *DGAT1* gene is widely reported associated with milk yield and composition, especially K232A polymorphism affecting on milk fat and protein [17,23,24]. In this study, the SNP (rs109421300) we identified is within an intron of *DGAT1* gene is the most significantly affecting FP ($P = 9.92e-25$), and also affecting PP ($P = 7.05e-9$), and a lot of

studies have reported for milk yield and composition in Holstein, Jersey and Holstein-Friesian cattle [25–27]. *DGAT1* gene encodes an multipass transmembrane protein that functions as a key metabolic enzyme, involved in triacylglycerol biosynthesis, and glycerolipid metabolism (KEGG: bta00561), retinol metabolism (KEGG: bta00830), metabolic pathways (KEGG: bta01100) and fat digestion and absorption (KEGG: bta04975). The PP-associated SNP (rs109875012) on BTA5 is within previously reported milk fatty acid content QTL [18], this SNP is close to the *ZNF384*, this gene encodes a C2H2-type zinc finger protein, which may function as a transcription factor and have DNA binding and metal ion binding functions. Another PP-associated SNP (rs108996837) on BTA21. Four SNPs (rs43526055, rs137676276, rs109528658, rs135780687) on BTA7, 11, 17 and X chromosome are associated with FY. The SNP on BTA11 at 19Mb located near the VIT, this gene encodes an extracellular matrix (ECM) protein, which has function of glycosaminoglycan binding (GO:0005539), is has been proposed participate in growth plate cartilage chondrocyte morphogenesis pathway (GO:0003429) and nervous system development pathway (GO:0007399). The SNP on BTA17 is within previously reported QTL for milk protein composition [28], and it also located in the *EP400*, this gene has been participate in histone H2A acetylation, histone H4 acetylation, and has the function of ATP binding, DNA binding, protein binding and helicase activity. The FY-associated SNP on chromosome X was located nearby the *GRPR*, this gene is gastrin releasing peptide receptor, gastrin-releasing peptide regulates numerous functions of gastrointestinal and central nervous system, which involved in calcium signaling pathway and neuroactive ligand-receptor interaction. Song reviewed a lot of studies on calcium signaling pathway participated in the effect of the sympathetic never in regulating adipose metabolism [29]. The PY-associated SNP (rs134480235) on BTA5 is within previously reported QTL region, which is associated with milk fatty acid

content, this SNP is in the *SLCO1A2* gene, which is encoding solute carrier anion transporter family, member 1A2, the gene participate in digestive system, organic anion transport process and bile secretion pathway (KEGG: 04976). Furthermore, when the association model not fitted PCs as covariance, four SNPs are overlapped with model with fitted PCs：rs109875012 (*ZNF384*), rs136949224 (*SCARA5*), rs109421300 (*DGAT1*), and rs109528658 (*EP400*), it is suggested that these significant SNPs could more stable and reliable.

Correlations among milk traits and pleiotropic QTLs

Estimates of heritability and genetic correlations are essential population genetic parameters in animal breeding research and application of animal breeding programs [30]. Genetic correlation can be useful for indirect selection, selection in different environment and selecting multiple traits simultaneously. We found there were relatively high genetic correlation between milk yield and protein yield, is consistent with the results in Holsteins [31] and the results in Jersey dairy cattle [30]. We expect that there are overlapping region of genetic variation between different traits that are relatively high correlation. A heatmap using *P*-values helped identify pleiotropic QTLs (**Figure** S4). The QTL on BTA1 (0.4 Mb) associated with three milk production traits (MY, FY, and PY). Another QTL on BTA1 (40.3-41.06Mb) is associated with MY, PY, and FP. The QTL on BTA2 at 14.8Mb showed that affecting MY, PY and FY. The QTL on BTA2 at 134Mb associated with MY, PY and FY, and also reported associated with the milk composition in another Holstein study [32]. The MY- and PY-associated QTL is adjacent to the *ABCG2* gene on BTA 6 at 37Mb, previously studies reported a missense mutations in the *ABCG2* gene is associated with milk yield and composition in Holstein, Braunvieh, and Fleckvieh cattle [6,33] , a intron variant affecting milk fatty acids in Chinese Holstein [34], and also reported for body

weight, calving ease direct in US cattle breed [35,36]. The QTL on BTA14 at 1.8Mb in the *DGAT1* gene associated with milk yield, fat and protein percentage. The QTL on BTA17 at 29-32Mb associated with MY, PY and FY, some study have pointed to QTL on this region for milk fatty acid, body weight [32,37]. The QTL on BTA 20 (67Mb) is associated with MY, PY and FY, previous study reported for PP and body weight in this region [38,39]. The QTL on BTA24 at 24Mb is pleiotropic and is associated with MY and FY, Ibeagha-Awemu and Peter reported it's associated with milk fatty acid [40], Kiser and Clancey reported for conception rate in Holstein cows [41].

Conclusions

We performed genome-wide association studies using test-day records for five milk production traits (milk yield, fat yield, protein yield, fat percentage, and protein percentage) in Holstein cows. In total of 10 significantly SNPs associated with fat and protein percentage, fat and protein yield, five SNPs are located within previously reported QTLs. An important top variant in the *DGAT1* gene affecting milk fat and protein, and we also found *SCARA5* and *GRPR* gene may be potential candidate gene involved in the biosynthesis of milk fat. In addition, some pleiotropic QTLs on BTA1, 2, 3, 6, 11, 14, 17, 20, and 24 associated with more than two milk traits, these results could provide some basis for molecular breeding and targets for gene cloning.

Acknowledgements

This study was partially supported by the High-Level Academic Papers in Ningxia University, the project of High-yield and High-quality Dairy Cattle Breeding (2013NYYZ0501), the National Science Foundation (Award # DBI 1661348), and the USDA National Institute of Food

and Agriculture Hatch project (1014919). The authors would like to thank the DHI measurement center of Ningxia Animal Husbandry of Extension Station and 21 dairy farms for their phenotypic data. We also thank Dr. Linda R. Klein for valuable writing advice and editing the manuscript.

Author contributions

Conceived experiment: Zhiwu Zhang and Yaling Gu;

Data analyses: Liyuan Liu, Jinghang Zhou, and James Chen;

Data collection: Juan Zhang, Jia Tian, and Wan Wen;

Wrote manuscript: Liyuan Liu, Jinghang Zhou, James Chen, Zhiwu Zhang, and Yaling Gu

Disclosures

The authors declare no conflict of interest.

Ethics Statement

No applicable. This study only did statistical analysis based on the former database which come from the Holstein breeding project in Ningxia province, China. All the authors in this study didn't participated in any sample collection process. All the farms involved were consent and agree to take part in this research.

Literature Cited

1. Hayes, B.J.; Bowman, P.J.; Chamberlain, A.J.; Goddard, M.E. Invited review: Genomic selection in dairy cattle: progress and challenges. *J. Dairy Sci.* **2009**, *92*, 433–443, doi:10.3168/jds.2008-1646.
2. Moniruzzaman, M.; Khatun, R.; Mintoo, A. Application of marker assisted selection for livestock improvement in Bangladesh. *Bangl. vet.* **2015**, *31*, 1–11, doi:10.3329/bvet.v31i1.22837.
3. Wakchaure, R.; Ganguly, S. Marker Assisted Selection (MAS) in Animal Breeding: A Review. *J Drug Metab Toxicol* **2015**, *06*, doi:10.4172/2157-7609.1000e127.

4. Kristensen, V.N.; Børresen-Dale, A.-L. SNPs associated with molecular subtypes of breast cancer: On the usefulness of stratified Genome-wide Association Studies (GWAS) in the identification of novel susceptibility loci. *Mol Oncol* **2008**, *2*, 12–15, doi:10.1016/j.molonc.2008.02.003.

5. Yang, W.; Gu, C.C. A whole-genome simulator capable of modeling high-order epistasis for complex disease. *Genet. Epidemiol.* **2013**, *37*, 686–694, doi:10.1002/gepi.21761.

6. Pausch, H.; Emmerling, R.; Gredler-Grandl, B.; Fries, R.; Daetwyler, H.D.; Goddard, M.E. Meta-analysis of sequence-based association studies across three cattle breeds reveals 25 QTL for fat and protein percentages in milk at nucleotide resolution. *BMC Genomics* **2017**, *18*, 853, doi:10.1186/s12864-017-4263-8.

7. Jiang, L.; Liu, J.; Sun, D.; Ma, P.; Ding, X.; Yu, Y.; Zhang, Q. Genome Wide Association Studies for Milk Production Traits in Chinese Holstein Population. *PLoS ONE* **2010**, *5*, e13661, doi:10.1371/journal.pone.0013661.

8. Pryce, J.E.; Johnston, J.; Hayes, B.J.; Sahana, G.; Weigel, K.A.; McParland, S.; Spurlock, D.; Krattenmacher, N.; Spelman, R.J.; Wall, E.; et al. Imputation of genotypes from low density (50,000 markers) to high density (700,000 markers) of cows from research herds in Europe, North America, and Australasia using 2 reference populations. *Journal of Dairy Science* **2014**, *97*, 1799–1811, doi:10.3168/jds.2013-7368.

9. Madsen, P.; Jensen, J. A Package for Analysing Multivariate Mixed Models. Version 6, release 5. 31.

10. Schaeffer, L.R. Application of random regression models in animal breeding. *Livestock Production Science* **2004**, *86*, 35–45, doi:10.1016/S0301-6226(03)00151-9.

11. Cobuci, J.A.; Euclydes, R.F.; Lopes, P.S.; Costa, C.N.; Torres, R. de A.; Pereira, C.S. Estimation of genetic parameters for test-day milk yield in Holstein cows using a random regression model. *Genetics and Molecular Biology* **2005**, *28*, 75–83, doi:10.1590/S1415-47572005000100013.

12. Purcell, S.; Neale, B.; Todd-Brown, K.; Thomas, L.; Ferreira, M.A.R.; Bender, D.; Maller, J.; Sklar, P.; de Bakker, P.I.W.; Daly, M.J.; et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **2007**, *81*, 559–575, doi:10.1086/519795.

13. Liu, X.; Huang, M.; Fan, B.; Buckler, E.S.; Zhang, Z. Iterative Usage of Fixed and Random Effect Models for Powerful and Efficient Genome-Wide Association Studies. *PLOS Genetics* **2016**, *12*, e1005767, doi:10.1371/journal.pgen.1005767.

14. Price, A.L.; Patterson, N.J.; Plenge, R.M.; Weinblatt, M.E.; Shadick, N.A.; Reich, D. Principal components analysis corrects for stratification in genome-wide association studies. *Nat. Genet.* **2006**, *38*, 904–909, doi:10.1038/ng1847.

15. Bovenhuis, H.; Weller, J.I. Mapping and Analysis of Dairy Cattle Quantitative Trait Loci by Maximum Likelihood Methodology Using Milk Protein Genes as Genetic Markers. *Genetics* **1994**, *137*, 267–280.

16. Thaller, G.; Krämer, W.; Winter, A.; Kaupe, B.; Erhardt, G.; Fries, R. Effects of DGAT1 variants on milk production traits in German cattle breeds1. *Journal of Animal Science* **2003**, *81*, 1911–1918, doi:10.2527/2003.8181911x.

17. Kühn, C.; Thaller, G.; Winter, A.; Bininda-Emonds, O.R.P.; Kaupe, B.; Erhardt, G.; Bennewitz, J.; Schwerin, M.; Fries, R. Evidence for Multiple Alleles at the *DGAT1* Locus Better Explains a Quantitative Trait Locus With Major Effect on Milk Fat Content in Cattle. *Genetics* **2004**, *167*, 1873–1881, doi:10.1534/genetics.103.022749.

18. Bouwman, A.C.; Bovenhuis, H.; Visker, M.H.; van Arendonk, J.A. Genome-wide association of milk fatty acids in Dutch dairy cattle. *BMC Genet* **2011**, *12*, 43, doi:10.1186/1471-2156-12-43.

19. Sanchez, M.P.; Govignon-Gion, A.; Ferrand, M.; Gelé, M.; Pourchet, D.; Amigues, Y.; Fritz, S.; Boussaha, M.; Capitan, A.; Rocha, D.; et al. Whole-genome scan to detect quantitative trait loci associated with milk protein composition in 3 French dairy cattle breeds. *Journal of Dairy Science* **2016**, *99*, 8203–8215, doi:10.3168/jds.2016-11437.

20. Nayeri, S.; Sargolzaei, M.; Abo-Ismail, M.K.; May, N.; Miller, S.P.; Schenkel, F.; Moore, S.S.; Stothard, P. Genome-wide association for milk production and female fertility traits in Canadian dairy Holstein cattle. *BMC Genet.* **2016**, *17*, 75, doi:10.1186/s12863-016-0386-1.

21. You, K.; Su, F.; Liu, L.; Lv, X.; Zhang, J.; Zhang, Y.; Liu, B. SCARA5 plays a critical role in the progression and metastasis of breast cancer by inactivating the ERK1/2, STAT3, and AKT signaling pathways. *Mol Cell Biochem* **2017**, *435*, 47–58, doi:10.1007/s11010-017-3055-4.

22. Ulker, D.; Ersoy, Y.E.; Gucin, Z.; Muslumanoglu, M.; Buyru, N. Downregulation of SCARA5 may contribute to breast cancer via promoter hypermethylation. *Gene* **2018**, *673*, 102–106, doi:10.1016/j.gene.2018.06.036.

23. Grisart, B.; Farnir, F.; Karim, L.; Cambisano, N.; Kim, J.-J.; Kvasz, A.; Mni, M.; Simon, P.; Frere, J.-M.; Coppieters, W.; et al. Genetic and functional confirmation of the causality of the DGAT1 K232A quantitative trait nucleotide in affecting milk yield and composition. *Proceedings of the National Academy of Sciences* **2004**, *101*, 2398–2403, doi:10.1073/pnas.0308518100.

24. Bovenhuis, H.; Visker, M.H.P.W.; Poulsen, N.A.; Sehested, J.; van Valenberg, H.J.F.; van Arendonk, J.A.M.; Larsen, L.B.; Buitenhuis, A.J. Effects of the diacylglycerol o-acyltransferase 1 (DGAT1) K232A polymorphism on fatty acid, protein, and mineral composition of dairy cattle milk. *Journal of Dairy Science* **2016**, *99*, 3113–3123, doi:10.3168/jds.2015-10462.

25. Meredith, B.K.; Kearney, F.J.; Finlay, E.K.; Bradley, D.G.; Fahey, A.G.; Berry, D.P.; Lynn, D.J. Genome-wide associations for milk production and somatic cell score in Holstein-Friesian cattle in Ireland. *BMC Genet* **2012**, *13*, 21, doi:10.1186/1471-2156-13-21.

26. Aliloo, H.; Pryce, J.E.; González-Recio, O.; Cocks, B.G.; Hayes, B.J. Validation of markers with non-additive effects on milk yield and fertility in Holstein and Jersey cows. *BMC Genet* **2015**, *16*, 89, doi:10.1186/s12863-015-0241-9.

27. Frąszczak, M.; Szyda, J. Comparison of significant single nucleotide polymorphisms selections in GWAS for complex traits. *J Appl Genetics* **2016**, *57*, 207–213, doi:10.1007/s13353-015-0305-6.

28. Schopen, G.C.B.; Koks, P.D.; van Arendonk, J.A.M.; Bovenhuis, H.; Visker, M.H.P.W. Whole genome scan to detect quantitative trait loci for bovine milk protein composition. *Animal Genetics* **2009**, *40*, 524–537, doi:10.1111/j.1365-2052.2009.01880.x.

29. Song, Z.; Wang, Y.; Zhang, F.; Yao, F.; Sun, C. Calcium Signaling Pathways: Key Pathways in the Regulation of Obesity. *IJMS* **2019**, *20*, 2768, doi:10.3390/ijms20112768.

30. Missanjo, E.; Imbayarwo-Chikosi, V.; Halimani, T. Estimation of Genetic and Phenotypic Parameters for Production Traits and Somatic Cell Count for Jersey Dairy Cattle in Zimbabwe. *ISRN Veterinary Science* **2013**, *2013*, 1–5, doi:10.1155/2013/470585.

31. Zaabza, H.B.; Gara, A.B.; Rekik, B. Genetic analysis of milk production traits of Tunisian Holsteins using random regression test-day model with Legendre polynomials. *ajas* **2018**, *31*, 636–642, doi:https://doi.org/10.5713/ajas.17.0332.

32. Gebreyesus, G.; Buitenhuis, A.J.; Poulsen, N.A.; Visker, M.H.P.W.; Zhang, Q.; van Valenberg, H.J.F.; Sun, D.; Bovenhuis, H. Multi-population GWAS and enrichment analyses reveal novel genomic regions and promising candidate genes underlying bovine milk fatty acid composition. *BMC Genomics* **2019**, *20*, 178, doi:10.1186/s12864-019-5573-9.

33. Cohen-Zinder, M. Identification of a missense mutation in the bovine ABCG2 gene with a major effect on the QTL on chromosome 6 affecting milk yield and composition in Holstein cattle. *Genome Research* **2005**, *15*, 936–944, doi:10.1101/gr.3806705.

34. Li, C.; Sun, D.; Zhang, S.; Yang, S.; Alim, M.A.; Zhang, Q.; Li, Y.; Liu, L. Genetic effects of FASN, PPARGC1A, ABCG2 and IGF1 revealing the association with milk fatty acids in a Chinese Holstein cattle population based on a post genome-wide association study. *BMC Genet* **2016**, *17*, 110, doi:10.1186/s12863-016-0418-x.

35. Saatchi, M.; Schnabel, R.D.; Taylor, J.F.; Garrick, D.J. Large-effect pleiotropic or closely linked QTL segregate within and across ten US cattle breeds. *BMC Genomics* **2014**, *15*, 442, doi:10.1186/1471-2164-15-442.

36. Smith, J.L.; Wilson, M.L.; Nilson, S.M.; Rowan, T.N.; Oldeschulte, D.L.; Schnabel, R.D.; Decker, J.E.; Seabury, C.M. Genome-wide association and genotype by environment interactions for growth traits in U.S. Gelbvieh cattle. *BMC Genomics* **2019**, *20*, 926, doi:10.1186/s12864-019-6231-y.

37. McClure, M.C.; Morsci, N.S.; Schnabel, R.D.; Kim, J.W.; Yao, P.; Rolf, M.M.; McKay, S.D.; Gregg, S.J.; Chapple, R.H.; Northcutt, S.L.; et al. A genome scan for quantitative trait loci influencing carcass, post-natal growth and reproductive traits in commercial Angus cattle. *Anim. Genet.* **2010**, *41*, 597–607, doi:10.1111/j.1365-2052.2010.02063.x.

38. Snelling, W.M.; Allan, M.F.; Keele, J.W.; Kuehn, L.A.; McDaneld, T.; Smith, T.P.L.; Sonstegard, T.S.; Thallman, R.M.; Bennett, G.L. Genome-wide association study of growth in crossbred beef cattle12. *Journal of Animal Science* **2010**, *88*, 837–848, doi:10.2527/jas.2009-2257.

39. Jiang, J.; Ma, L.; Prakapenka, D.; VanRaden, P.M.; Cole, J.B.; Da, Y. A Large-Scale Genome-Wide Association Study in U.S. Holstein Cattle. *Front. Genet.* **2019**, *10*, 412, doi:10.3389/fgene.2019.00412.

40. Ibeagha-Awemu, E.M.; Peters, S.O.; Akwanji, K.A.; Imumorin, I.G.; Zhao, X. High density genome wide genotyping-by-sequencing and association identifies common and low frequency SNPs, and novel candidate genes influencing cow milk traits. *Sci Rep* **2016**, *6*, 31109, doi:10.1038/srep31109.

41. Kiser, J.N.; Clancey, E.; Moraes, J.G.N.; Dalton, J.; Burns, G.W.; Spencer, T.E.; Neibergs, H.L. Identification of loci associated with conception rate in primiparous Holstein cows. *BMC Genomics* **2019**, *20*, 840, doi:10.1186/s12864-019-6203-2.

Figures and Tables

Figure. 1 Properties of Single Nucleotide Polymorphisms (SNPs). A total of 1,220 cows were genotyped by the Illumina Bovine 150 k BeadChip. After conducting quality control on both minor allele frequency (MAF, above 5%) and missing rate (< 10%), 1,220 individuals and 124,743 SNP remained. The distribution of the filtered SNPs is displayed over the 30 bovine chromosomes except for Y chromosome (a). The MAFs of SNPs were re-calculated after the filtering and were displayed by a heat map. Consequently, there SNPs with MAF < 5% remained, as demonstrated by the histogram (c). The density of SNPs is displayed by the frequency of the distance between adjacent SNPs (b). The distances over 60 kb clustered into one group. The maximum distance was 100.06 kb. Pairwise Linkage Disequilibrium (LD) was calculated as the R square for SNPs within the 100 kb window. The decay of LD over distance (red line) is displayed by the pairwise LD and moving average (d).

Figure. 2 Population structure demonstrated by principal component analysis. Principal Component Analysis (PCA) was conducted with the 124,743 SNPs covering the whole bovine genome. The population structure is demonstrated by the pairwise scatter plots (a, b, and c) and the 3D plot (d) of the first three principal components (PCs).

Figure. 3 Associations between 124,743 SNPs and milk traits. Milk traits include fat yield (FY), protein yield (PY), fat percentage (FP), protein percentage (PP). The association analyses were conducted by the FarmCPU R package. Manhattan plots display the negative logarithms of the observed *P* values for SNPs across 30 bovine chromosomes (left panel). The green line indicates the Bonferroni multiple test threshold at *P* = 4.0e-07. The Quantile-Quantile (QQ) plots represent the negative logarithms of the expected *P* values (X-axis) and observed *P* values (Y-axis) (right panel).

Table. 1 Variance components and heritability of milk traits

Variance component	MY	FP	FY	PP	PY
Genetic variance	1592.62	4.77	1.48	1.65	1.20
Permanent environmental variance	4267.93	10.58	5.46	3.45	3.96
Residual variance	7413.63.63	0.51	0.08	0.07	0.03
h^2	0.12	0.30	0.21	0.32	0.23

*MY, milk yield; FP, fat percentage; FY, fat yield; PP, protein percentage, PY, protein yield.

Table. 2 Genome-wide significant SNPs associated with milk traits

Milk traits	SNP	Chr._Mb	Position(bp)	MAF	Nearest gene	Distance(kb)	*P*-value
FP	rs42295213	1_41	41061715	0.36	*EPHA6*	within	1.50E-07
PP	rs109875012	5_104	104120905	0.45	*ZNF384*	2.6	4.03E-08
PY	rs134480235	5_89	89267320	0.49	*SLCO1A2*	within	1.57E-08
FY	rs43526055	7_73	73431219	0.31	*ADRA1B*	179.8	4.48E-09
FP	rs136949224	8_10	10705865	0.14	*SCARA5*	43.5	3.57E-08
FY	rs137676276	11_19	19277448	0.11	*VIT*	25	8.58E-09
FP, PP	rs109421300	14_1.8	1801116	0.23	*DGAT1*	within	9.92E-25
FY	rs109528658	17_46	46090458	0.40	*EP400*	within	7.05E-09
PP	rs108996837	21_69	69386346	0.12	*EXOC3L4*	32.6	2.36E-08
FY	rs135780687	X_134	134726985	0.42	*GRPR*	83	1.63E-10

*FP, fat percentage; PP, protein percentage; PY, protein yield; FY, fat yield.