Evolution and spread of SARS-CoV-2 likely to be affected by climate

Authors: Priyanka Bajaj, Prakash Chandra Arya

Correspondence: priyankaba@iisc.ac.in, prakasha@iisc.ac.in

Graphical Abstract

*Climate types adapted by SARS-CoV-2

In Brief:

The authors elucidate adaptation of SARS-CoV-2 to different climates by studying phylogenetics and the distribution of strains on Koppen's climate map.

Highlights:

- Phylogenetic network divides SARS-CoV-2 strains into two variant groups, G1 and G2.
- G1 strains is restricted to Koppen's *"temperate"* climate (mainly *Cfa-Cfb*).
- G2 strains has evolved from G1 to sustain in other climates mainly "*humid-continental" (Dfa-Dfb)* and "*tropical-savannah" (Aw)* climate.

Evolution and spread of SARS-CoV-2 likely to be affected by climate

Priyanka Bajaj[1#*] and Prakash Chandra Arya[2#**]

[1]Molecular Biophysics Unit, Indian Institute of Science, Bangalore-560012, India
[2]Centre for Earth Sciences, Indian Institute of Science, Bangalore-560012, India
# Both authors have equally contributed to this work
*Correspondence: priyankaba@iisc.ac.in
**Correspondence: prakasha@iisc.ac.in

Abstract

COVID-19 pandemic has been extensively studied by many researchers. However, it is still unclear why it was restricted to higher latitudes during the initial days and later cascaded in the tropics. Here, we analyzed 176 SARS-CoV-2 genomes across different latitudes and climate (Koppen's climate) that provided insights about within species virus evolution and its relation to abiotic factors. Two genetically variant groups, named as G1 and G2 were identified, well defined by four mutations. The G1 group (ancestor), is mainly restricted to warm and moist, temperate climate (Koppen's C climate) while its descendent G2 group surpasses the climatic restrictions of G1, initially cascading into neighboring cold climate (D) of higher latitudes and later into hot climate of the tropics (A). It appears that the gradation of temperate climate (Cfa-Cfb) to cold climate (Dfa-Dfb) drives the evolution of G1 into G2 variant group which later adapted to tropical climate (A) as well. It seems this virus followed inverse latitudinal gradient in the beginning due to its preference towards temperate (C) and cold climate (D). Our work elucidates virus evolutionary studies combined with climatic studies can provide crucial information about the pathogenesis and natural spreading pathways in such

outbreaks which is hard to achieve through individual studies. Mutational insights gained may help design an efficacious vaccine.

Keywords: SARS-CoV-2, molecular phylogeny, virus cluster SNPs, climate zones, Koppen's climate

Introduction

The first case of Corona Virus Disease-19 (COVID-19) pandemic caused by (Severe Acute Respiratory Syndrome-2) SARS-CoV-2 pathogen was first reported from Wuhan, China[1]. In spite of various precautions such as lockdown, social distancing, wearing mask, and sanitization, the disease was able to reach almost every part of the world[2]. This zoonotic virus is like SARS (Severe Acute Respiratory Syndrome) coronavirus (79% similarity) and MERS (50% similarity) and is closely related to bat derived coronaviruses[1]. The SARS-CoV-2 can survive up to 3, 4, and 24 hours on aerosols, copper, and cardboard respectively, and up to 2-3 days on stainless-steel or plastic[1] and spreads faster than its ancestors SARS-CoV-1 and Middle East Respiratory Syndrome (MERS-CoV)[1,3]. This led it concur a larger geographical region by infecting larger population. Since the social behaviour and travelling of humans have not changed much, what makes few respiratory viruses confined locally and others spread globally is still unclear. COVID-19 outbreak led to a big discussion, that does climate have a role in the spread of the disease? The ancestor SARS-CoV-1 losses its viability at higher temperature (38°C) and relatively higher humidity (>95%)[4]. Experiments support that SARS-CoV-2 is highly stable at 4°C but is sensitive to heat[5]. Studies both in favour and against have been published but the topic is still debatable[6,7]. A recent review compiles 61 studies relating COVID-19 with climate[1]. Several climatic factors such as humidity, precipitation, radiation, temperature, and wind speed affecting this virus spread

have been incorporated. Both positive and negative association of COVID-19 with temperature and humidity have been published[1]. Recently, Carlson et al. mentioned that COVID-19 transmission could be affected by climate but discouraged the use of SDMs (Species Distribution Models) for COVID-19 transmission due to their limitations, which generally takes only climatic parameters as input, as they may not be appropriate tools[7]. This study was challenged by Araujo et al. mentioning strengths and limitations of the tools and reasoned that R_0 of COVID-19 depends on several factors, it may also be affected by climate[6]. Since only climatic parameters are insufficient to capture climatic signatures of COVID-19 spread, such patterns can be recognized by combining phylogenetic and climatic studies[8]. Such approach enables to probe the similarities or differences in virus genome across similar or different climate types present all over the world. To understand such a behaviour, we have attempted to study the genomic sequence of the SARS-CoV-2 across different latitudes and climates.

Till now, pathogens are poorly mapped and little is known about their underlying ecological and evolutionary causes[9]. Nucleotide substitution has been proposed to be one of the most important mechanisms of viral evolution in nature[10]. However, factors responsible for the generation of these mutations are not well understood. One of the possible factors is adaptation to new environments, dictated by natural selection that discriminates among genetic variations and favours survival of the fittest[11]. Virus evolution as a consequence of climate change is poorly understood. SARS-CoV-2 consists of large single-stranded ~30kb long positive-sense RNA. These viruses majorly have a conserved genomic organization, consisting of a unique 265bp long leader sequence, ORF1ab polyprotein, and structural proteins like S (spike glycoprotein), E (Envelope), M (Membrane), and N (Nucleocapsid).

ORF1ab encodes replicase, transcriptase and helicase, essential enzymes required for replication, along with non-structural and accessory proteins. Expression of non-structural proteins is facilitated by ribosomal frameshifting[12]. All coronaviruses express structural proteins S, E, M, N; spike glycoprotein being the most immunogenic to T-cell response[13]. Spike glycoprotein of coronaviruses binds to human angiotensin-converting enzyme 2 (hACE2) receptor for viral fusion and entry and is the main target for neutralizing antibodies and development of vaccines[14]. Membrane protein is also antigenic as it stimulates a humoral immune response[15]. E protein is responsible for virus assembly and release of virion particles[16]. Nucleocapsid protein packages RNA genome into a helical ribonucleocapsid protein (RNP) complex during virion assembly and is capable of eliciting an immune response[17]. Since it is still not clear whether SARS-CoV-2 evolution and spread have relation with climate, our study may act as a missing link between genomic sequence, climate and COVID-19 severity. If SARS-CoV-2 is responding towards external climate it can be delineated by superimposing its genomic variants across different latitudes and Koppen's climate[8]. The earliest and the most simple classification of Earth's climate is based on latitudes which divide the Earth's climate into seven climate zones, North Frigid Zone (NFZ), North Temperate Zone (NTZ), North Subtropical Zone (NSTZ), Tropical Zone (TZ), South Subtropical Zone (SSTZ), South Temperate Zone (STZ) and South Frigid Zone (SFZ) lying between 90°N to 66.5°N, 66.5°N to 30°N, 30°N to 23.5°N, 23.5°N to 23.5°S, 23.5°S to 30°S, 30°S to 66.5°S and 66.5°S to 90°S, respectively[18]. Based on temperature and precipitation, Wladimir Koppen divided Earth's climate into five major climates, A (Tropical), B (Arid), C (Temperate), D (Cold or Continental) and E (Polar) which are further subdivided into 30 climate types[19]. To understand the

effect of climate on SARS-CoV-2 evolution, the present study comprises of two parts: (1) Sequence analysis of SARS-CoV-2 strains, (2) Mapping SARS-CoV-2 strains across different climates. These combined studies can provide insights on within species evolution and preferential distribution of SARS-CoV-2 across different climatic zones which might be difficult to probe through individual studies.

Methodology

Molecular phylogenetic analysis

Approximately, 11,000 full-length genome sequences of SARS-CoV-2 were available in Global Initiative on Sharing Avian Influenza Data (GISAID) database, accessed till 2[nd] May 2020. 185 full-length SARS-CoV-2 genomic sequences from countries across the globe, with genome length more than 29 kb and high coverage were obtained from GISAID database and the reference genome was retrieved from GenBank24 (Supplementary Table S1). This sample size was selected by taking 95% confidence interval, 0.5 standard deviation and 7% margin of error. To avoid bias related to the geographical area covered by a country, genomic sequence of strains isolated from different locations from each country/ climate type was retrieved, depending on the availability of data. The corresponding location, latitude, Koppen's climate, Koppen's Climate type, SARS-CoV-2 variant group, Environment/ Region, Climate zone, temperature, and precipitation of each strain is provided in Supplementary Table S2. These sequences were aligned to the full reference genome[20] by using Biomanager and Seqinr packages of R (version 3.6.3). Among 185 genomes, some partial genomes were discarded. Thus, finally 176 genomes were analysed. NC_045512 genome sequence was used as reference and the genomic coordinate in this study is based on this

reference genome. Based on protein annotations, nucleotide level variants were

converted into amino acid codon variants for alignments when its location within a

gene was identified. The amino acid position numbering is according to its position

within the specified gene (Coding Sequence) as annotated in reference sequence

(NC_045512, NCBI)[20]. To ensure comparability, we trimmed the flanks of all

sequences. The aligned sequences were used to construct a phylogenetic tree

using MEGA X[21]. The evolutionary history was inferred using the Neighbor-Joining

method (500 bootstrap tests)[22]. The optimal tree with the sum of branch length =

0.01116462 is shown. The tree is drawn to scale, with branch lengths in the same

units as those of the evolutionary distances used to infer the phylogenetic tree. The

evolutionary distances were computed using the Maximum Composite Likelihood

method[23] and are in the units of the number of base substitutions per site. All

ambiguous positions were removed for each sequence pair (pairwise deletion

option). A total of 29408 positions were present in the final dataset. The results are

presented in the form of DNA sequencing i.e., U (uracil) is read as T (thymine). We

have labeled each virus strain by the GISAID Accession ID and the location from

which it was isolated in the format "Location|EPI ISL Accession ID", in the

constructed phylogenetic tree. For ease of visualization, we have marked a new

Strain ID (1 to 176) against each SARS-CoV-2 isolate in the phylogenetic tree

(Figure 1). The same Strain ID is used for the climatic studies in this article. High-

frequency SNPs (Single Nucleotide Polymorphisms) distinguishing one virus

cluster from the others is referred to as "virus cluster SNPs" throughout this paper.

Mapping virus strain on the Koppen's climate map

The location of each SARS-CoV-2 strain is obtained from the METADATA file

provided in GISAID database for each viral isolate (Supplementary Table S1). The

coordinates of the locations were taken from the official website of USGS Earth

Explorer[24]. The Gieger-Koppen's climate map is used for climatic studies[19]. The

Koppen's climate type, temperature, precipitation of each strain is assessed from

weatherbase[25] and CLIMATE.ORG[26]. The above information for each strain is

tabulated in Supplementary Table S2. The map is georeferenced by using 'Arc-

GIS 10.1'[27]. The locations of all strains (n=176) were transferred to the

georeferenced map[27]. On the map, the G1 strains were symbolized as 'Yellow-

circle', and G2 as 'Square' (Figure 3). Each strain in the map is labelled as per their

Strain ID (1 to 176) (Figure 3), the map combines information of the phylogeny,

climate, and global distribution of SARS-CoV-2. These locations were classified

into coastal and continental region, we define the coastal region as land region <

500 km from the ocean/sea and the continental region as land lying >500 km from

the coastline measured through google maps.

Statistical analysis

Chi-square were performed in Microsoft Excel (2016) to test our null hypothesis.

Null hypothesis for these tests are mentioned in the text and its corresponding

contingency table is mentioned in Supplementary Table S2. Histograms depicting

the distribution of coronavirus in coastal region, continental region, Koppen's

climate and climate type were plotted using R (version 3.6.3). SigmaPlot10 was

used to generate box plot, regression plot, and mesh plot to statistically compare

frequency distribution of latitude, temperature, and precipitation of G1 and G2

strains. We performed one-way ANOVA to estimate statistical differences in the

latitude, temperature and precipitation between G1 and G2 virus populations.

Various scatterplots between latitude, temperature, and precipitation of G1 and G2

strains were plotted in R (version 3.6.3). Values were considered statistically

significant for P values below 0.05. Exact P values are provided in appropriate

Figures.

Data accessibility

The full-length genomic sequences were downloaded from GISAID website

(https://www.gisaid.org/), an open-source database for influenza viruses. The data

is downloaded as FASTA file along with the acknowledgement. The location of

each strain is accessed from its METADATA file. The Koppen's Climate map is

taken from the published article[19]. The Koppen's climate type, temperature and

precipitation for each strain is taken from weatherbase

(https://www.weatherbase.com/) and CLIMATEDATA.ORG (https://en.climate-

data.org/). Refer Supplementary Tables S1-S5. The code is available from the

corresponding authors on request.

Results

Molecular phylogeny analysis to infer genomic similarities and their

distribution in different climates

To probe genomic similarities between SARS-CoV-2 virus isolates, a phylogenetic

tree was constructed by aligning 176 virus genomes to the reference genome[20]

retrieved from GISAID. Interestingly, our Multiple Sequence Alignment (MSA)

results reveal sixty virus cluster Single Nucleotide Polymorphisms (SNPs) (see

methodology). Table1 comprises of SNPs of virus clusters across different climatic

zones, Koppen's climate and climate type. Climatic parameters (temperature and

precipitation) for each virus strain is mentioned in Supplementary Table S2. Based

on phylogenetic clustering, 176 SARS-CoV-2 strains are majorly divided into two

groups, we named them as G1 (1-58) and G2 (59-176) (Figure 1). Predominantly

four mutations distinguish G2 from G1 group, i.e., 1) a synonymous mutation
(C241T) appeared in the unique leader sequence, 2) F924 (C3037T) appeared in
nsp3, encoding for papain-like proteinase[28], 3) a non-synonymous mutation,
P214L (C14408T) arose in ORF1b, that codes for four putative non-structural
proteins (nsp13, nsp14, nsp15 and nsp16), functionally involved in replication-
transcription complex[29], and 4) D614G (A23403G) arose in S gene, encoding spike
glycoprotein[13] (Figure 2a). Among four mutations in G2, the D614G mutation, lying
in spike glycoprotein was widely studied due to its higher infectivity and
involvement in entering the host cell through hACE2 receptors[30–33]. The other three
mutations in G2 have co-evolved with D614G making it distinguishable from G1.
We explored the extent of genome-wide divergence of G1 and G2 group across
different climate zones and Koppen's climate (Figure 2b). 59% of G1 viruses fall in
NTZ, 14% in NSTZ, 12% in TZ, 10% in SSTZ and 5% in STZ. 76% of the virus
isolates in G2 group are present in the NTZ, 13.5% in TZ, 7.6% in STZ and
remaining 2% is equally distributed in NSTZ and SSTZ, showing G2 strain variants
evolved to adapt to temperate zones as their population decreased drastically in
the subtropical zones. These results show both G1 and G2 strains have a strong
preference towards higher latitudes i.e., NTZ (Figure 2c). Mapping viral strains on
Koppen's map (thoroughly discussed in the next section) reveal their prevalence
majorly in the C and D climate (Figure 2d). 71% of G1 lie in C climate, 17% in D
and the remaining is equally distributed in the A and B climate. 54% of G2 lie in C
climate, 36% in D, 9% in A and 1% in B climate pointing towards a preferential shift
of the novel coronavirus towards D climate (Figure 2b), alluding G2 is climatically
and genomically more diverse than G1. The analysis suggests that the G1 group
is mostly restricted to temperate climate (C) and G2 is climatically and

geographically widely distributed, it is possible that these mutations were acquired by G1 to stabilize itself in different climates hence allowing it to spread globally. Similar climatic concordance with the temperate climate (C) was also observed for SARS-CoV-1 that was responsible for 2002-2004 epidemic as it prevailed in regions of Australia, Europe, Canada and, China[34], having Koppen's C climate. Such similar occurrence of SARS-CoV-1 and G1 group of SARS-CoV-2 hints towards, why initially G1 variant group (consisting of the reference genome NC_045512[20]) that has 79% similarity to SARS-CoV-1[1] was majorly located in the temperate climate (C), and latter it evolved into G2 variant group that allowed it to extend its climatic boundaries into temperate, cold and tropical climate. These results suggest these four SNPs could be the key factors in increasing the virulence, transmission and sustainability of the virus in humans.

We further analyzed the order in which the phylogenetic clusters evolved from the ancestor 45-57 cluster (containing the reference genome, Strain ID: 50) based on nodes, mutational branches and branch length. The order in which the virus evolved is 44-47 (G1440A, G392D; G2891A, A876T), 1-22 (C8782T, S2839; T28144C, L84S), 33-43 (G26144T, G251V), 23-32, 58-61 (C15324T, N519), 80-115 (G28881A, G28882A, R203K; G28883C, G204R), 116-125 (A20268G, L2167), 126-176 (G25563T, Q57H) and 62-79 (cluster, acquired genomic mutation and its corresponding amino acid mutation). In Figure 2e, looking at the distribution of the viruses in different climate zones, no such preference was observed as the virus evolved. Virus cluster 58-61, linking G1 and G2 has an equal distribution of virus strains in C and D climate. The virus cluster 80-115 of G2 more closely related to G1, is widely distributed in A, C and D climates. Within 80-115 virus cluster, 106-115 cluster shows distribution in C and A climate. A trend was observed that virus

clusters in G2 group gradually evolved to sustain in Koppen's D climate (80-115 to

116-125 to 126-176). Within these major virus clusters, small clusters also exist as

shown in Table1 with their mutations along with their climatic distribution.

We have examined whether climatic conditions exhibit any selective pressure on

each gene (Figure 2f). Since, the present picture of the data shows that SARS-

CoV-2 spreads widely in NTZ, as expected all genes are having mutations in NTZ,

suggesting the virus is probably using varied mechanisms to adapt to the two main

climates of NTZ i.e., temperate (C) and cold or continental (D). Mutations in the M

gene are only pertaining to NTZ and NSTZ and are present in C and D climate. In

particular, there is a surge in the virus strains carrying SNPs in ORF8 in the NSTZ

(20%). 77% of the SNPs in ORF8 lie in the C and 20% in the D climate. Overall,

the distribution of virus cluster SNPs of ORF1ab, S, ORF3a, and N gene follows a

similar pattern across all the climatic zones and Koppen's climate, implying no

difference in selective pressure of the climate in generating mutations in these

genes. S, M, and N proteins are immunogenic[13,15,17], implicating virus evades

immune response by introducing these substitutions.

Apart from non-synonymous mutations, synonymous mutations within the gene

can also significantly affect protein function due to codon usage bias[35] and through

mechanisms such as ribosome stalling[36] and mRNA secondary structure

formation[37]. We probed the frequency of derived synonymous versus non-

synonymous mutations and observed a very similar distribution pattern of the

derived synonymous versus missense mutations across all climate zones and

Koppen's climate (Figure 2g). These analyses suggest novel coronavirus is using

varied mechanisms both at the transcriptional as well as translational level to

adapt, survive, and increase infectivity in all types of climates. These findings

unequivocally bolster a requirement for further prompt, comprehensive studies that

join genomic information, epidemiological information, and climatic distribution with

COVID-19 severity.

Distribution of strains across Koppen's climate

To probe the relation between climate and SARS-CoV-2 strains, we superimposed

genomic information along with their geolocations on the climate map of Wladimir

Koppen (Figure 3). We carefully examined the distribution of strains on Koppen's

map and an overview of the map shows, the distribution of 176 strains are mainly

concentrated in the western coasts of Europe and North America, and eastern

coasts of China, North America, Australia and South America (Figure 3).

Throughout the text Koppen's climate type is marked within quotations and its

standard symbol is written within brackets e.g., "*humid-subtropical" (Cfa)*.   List of

Koppen's symbol of each climate type is given in Supplementary Table S3 and its

criteria for classification is given in Supplementary Table S4. Mostly the SARS-

CoV-2 strains are distributed in the "*humid-subtropical" (Cfa)* and "*marine-*

temperate" (Cfb) and "*humid-continental" (Dfa-Dfb)* climate and, two strains from

virus clusters (80-115 and 126-176) belonging to South America, are found in

"tropical-savanna" *(Aw)* of 'A' climate (Supplementary Table S5). The map displays

~86% (151/176) of virus isolates are distributed in the coastal regions and the

remaining in continental region (Chi-square test, *P*<.001 considering null

hypothesis that equal number of virus isolates are found in coastal and continental

region, Figure 4a, Supplementary Table S6). Around ~74% (130/176) of the total

strains are distributed in "humid-subtropical" (Cfa) and "marine-temperate" (Cfb)

climate type of C climate and "humid-continental" (Dfa-Dfb) climate type of D

climate. The remaining ~26% (46/176) strains are distributed in other climate types

of Koppen's climate including non 'Cfa-Cfb' of C climate and non 'Dfa-Dfb' of D

climate (Figure 4b). It seems that spread of COVID-19 is maximally in areas with

'Cfa' and 'Cfb' climate type. The climatic parameters (temperature and

precipitation) in which these strains lie, were analyzed. Statistically, significant

difference was found in the latitudes of G1 ~24.14±3.5 (mean±s.e.) and G2

~34.03±2.7 (one-way ANOVA, P=.03251, Figure 5a). Statistically, significant

difference was observed in the temperatures of G1 (15.82±0.75 ^{o}C (mean±s.e.)

and G2 (11.67±0.68 ^{o}C) strains (one-way ANOVA, P<.001, Figure 5b). However,

the difference in precipitation for G1 (1046.95±80 mm) and G2 (896.64±35.48 mm)

strains is statistically not significant (one-way ANOVA, P=.06118, Figure 5c). The

latitudes and temperature are inversely related to each other (r = -0.6649,

Supplementary Figure S1a), which explains the occurrence of G1 strains in lower

and G2 strains in higher latitudes (Figure 5d). Such relation between latitude and

precipitation has not been observed (r = -0.3064, Supplementary Figure S1b and

Figure 5e). A mesh plot simultaneously evaluates all climatic parameters for both

G1 and G2 strains, the results agree to the limited temperature and wider

precipitation range of G1 group and interestingly the G2 group appears in a wider

temperature and shows a preferential shift towards lower temperature which is

evident from the fact that it initially appeared more in higher latitudes

(Supplementary Figure S2). A complete description of the distribution of G1 and

G2 strains lying in different countries and/or continents of the world is provided in

Supplementary Figure S3 and Supplementary Material.

Discussion

A pattern observed and discussed by several authors is that SARS-CoV-2

predominantly affected higher latitude countries (eg., China, Europe, USA and

Australia, Japan, Turkey etc.) in its preliminary stage (November 2019 to March

2020). This led many researchers predict a reduced spread in regions with higher

temperature such as tropical countries[38,39]. Our phylogenetic analysis provides one

more dimension to the above observations in which we have identified two groups

G1 and G2, similar observations were also reflected in other studies[40,41]. We have

further integrated the genomic information with climatic data and analyzed the

spread of these two variant groups across the globe and its climate.

There can be many reasons behind the observed pattern of spread. First, it could

be because of less testing in tropical countries. However, if this would be true, local

media would have reported over-crowding of patients in hospitals. For example,

Italy escalated its testing when media reported over-crowding of hospitals with

patients having COVID symptoms. But such incidences were not reported from

tropical countries. Second, it could be higher human mobility between China US,

and China Europe. But mobility between China and South-East Asia is also very

high. Thus, no outbreak in tropical countries initially is perplexing (till March 2020).

Third, most countries implemented lock-down measures from mid-March. Until

March people were working in normal mode. Hence, it is not possible that stringent

measures in containing the virus evaded it from cascading into the tropical

countries. Fourth, the population density of the tropics could be lower than

temperate countries. But this is also not true, in fact most of world population

resides in tropical countries and are mostly developing with limited proper health

and hygiene facilities compared to temperate countries. In such situation the

disease should have equally spread in tropics like that of temperate countries as

positive cases of COVID-19 were identified from many tropical countries in January

itself. Fifth, our analyses show that G1 group spread largely in temperate climate.

A possible reason behind this could be that host-immune response of people residing in the tropical countries resisted the infection. However, no supporting evidence have been found till now and needs detailed investigation.

Hence, these reasons negate the fact that the observed patterns could solely be due to community structure, social dynamics, government policies, global connectivity, population density and number of reported cases. Similar observations i.e. more COVID cases in higher latitudes were reflected in COVID-19 growth curves of Yusuf and Bukhari until March 2020 for different countries, which led them interpret that natural factors (temperature and humidity) are responsible for less number of COVID cases in tropical countries[42]. Thus, we are inclined to say that environmental factors, directly or indirectly play a role to explain the observed pattern of spread in cold countries during the initial stages. Since, in winters mostly people stay indoors, thus climate can indirectly be involved in increased growth rate of virus in cold countries. However, if climate indirectly had a role to play in the spread, spread of the disease would have been limited in summers in tropical countries. On the contrary, the COVID-19 outbreak in tropical countries started in summers, soon after emergence of G2 strains.

Results of this study depict that in the initial stages (until March 2020) of the pandemic the G1 group was restricted to temperate climate (Koppen's C type) whereas G2 group spread all over the world. Our analyses suggest that same regions were affected by G1 variant group and SARS-CoV-1. It is known that SARS-CoV-1 was not able to survive in regions with warm climate as its lipid bilayer was prone to degradation at higher temperature[43] and it loses its viability at higher temperature[4]. A recent study shows the structural stability of SARS-CoV-2 Virus

Like Particles (VLPs) degrades on increasing temperature[44], supporting our

observations.

Our analyses favor that evolution of G1 into G2 helps to sustain this virus from

temperate to cold and tropical climate successively, mainly due to four mutations

i.e., in leader sequence, ORF1ab, and S gene. The leader sequence and ORF1ab

is involved in replication and transcription, and the S gene is involved in binding to

the host cell through hACE2 receptors. Substitutions in the ORF1ab gene may

increase the synthesis of replicase-transcriptase complex, thus, increasing the

replication rate of the virus and blocking the host innate-immune response. 614

position in spike glycoprotein lies near the S1/S2 subunit junction where the furin-

cleavage site is present (R667) that enhances virion cell-cell fusion[45]. This

suggests, aspartate to glycine substitution in the vicinity of the furin-recognition site

may result in a conformational change of the spike glycoprotein that favors higher

affinity of the Receptor Binding Domain (RBD) to hACE2. A recent article showed

retroviruses pseudotyped with Glycine at 614 position infected ACE2-expressing

cells markedly more efficiently than those with Aspartic acid due to less S1

shedding and greater incorporation of the S protein into the pseudovirion[30]. Several

studies reported D614G mutation is increasing at an alarming rate[31,32]. Few

observed that this alteration correlated with increased viral loads in COVID-19

patients[31]. This is consistent with the epidemiological data showing proportion of

viruses bearing G614 is correlated to increased case fatality rate on a country by

country basis[33]. This substitution coevolved with substitution in the leader

sequence, nsp3 and RdRp (RNA dependent RNA polymerase) proteins,

suggesting these mutations allow the virus to transmit more efficiently. This

explains these mutations have not emerged merely because of founder's effect but

this virus under selection pressure has made itself more stable and infective. Also,

Forster et al. observed in his phylogenetic analysis, the preferential geographical

spread of SARS-CoV-2 and provided a plausible cause which could be founders

effect or immunological or environmental effect[41]. Although there is a possibility

that the stable variant might have appeared because of host innate immune

response or some unknown reason, in such a case it would not show any close

association with climate. Through our analysis, we are inclined to say that climate

has an effect of SARS-CoV-2 evolution. However, in particular, selective pressure

of climate on each gene of SARS-CoV-2 is not visible. Our genomic analysis of

virus strains show that the novel coronavirus undergoes both synonymous as well

as non-synonymous mutations throughout its genome in various climates,

suggesting the novel coronavirus uses multiple mechanisms both at the

transcriptional and translational level for evading the immune response, developing

drug resistance and increasing pathogenesis. However, the actual role of these

mutations is not yet determined, and these studies need to be further enlightened

by biophysical and biochemical studies. Such mutational insights will aid the design

of efficacious vaccines that can be stored and transported in a wide range of

temperature and conditions, thereby minimizing cold storage costs.

To delineate the signatures of underlying abiotic factors (temperature,

precipitation, and latitude) responsible for evolution of SARS-CoV-2 (n=176),

spreading patterns of G1 and G2 strains were carefully examined on Koppen's

climate map. Figure 3 shows an elevated spread of COVID-19 in the western and

eastern coasts of the continents and a diminished spread in the hot and cold

deserts. The G1 strains are majorly present in the eastern and western coasts of

the continents and G2 strains lie in both the coastal regions and continent's interior.

On a closer inspection, the eastern coasts of continents consist of "humid-subtropical" (Cfa) climate while the western coasts of continents consist of "marine-temperate" (Cfb), commonly known as east and west coast climate, respectively. These two climates are very similar to each other and belongs to temperate climate also known as C type climate of Koppen's classification scheme. A very large portion (~94%) of habitable China consists of temperate climate (C), i.e., humid subtropical climate (Cfa), which explains presence of only G1 strains in China and one strain of G1 is present in cold climate (D) present near the transition of temperate (C) to cold climate (D), thus probably temperate climate was suitable for G1. A similar association of G1 with temperate climate (C) was found in eastern and western coast of North America, eastern coast of South America, western coast of Europe and eastern and western coast of Australia. Statistically, distribution of G1 strains all over the globe is in concordance with the temperate climate and strongly favor C climate (Chi-square test, $P<.01$ for null hypothesis that G1 strains are equally distributed in all climates, Supplementary Table S6) as compared to any other climate. If climate does not have any role in the evolution and preferential spread of coronavirus, in such a case G1 would have been evenly distributed in all climate types which is not the case. Few exceptions of G1 seen in other climate types are most probably because of travel as they remained subsided in that climate, implying their inability to sustain in other climate types. It appears that the G1 strains existed in temperate climate all over the world but could not extend their geographical territories beyond temperate climate. Contrastingly, the evolved G2 strains can sustain in temperate (C), cold (D) and tropical (A) climate surpassing the climatic restrictions of G1. Map interpretation suggests that G2 strains enters the continent's interior through D climate (e.g., North America

and Russia). Temperate climate (C) generally grades into cold climate (D) and deserts (B) in the northern hemisphere (e.g., C to D: Europe to Russia, and USA to Canada; C to B: China, and USA). In southern hemisphere, gradation of temperate climate (C) into tropical climate (A) and deserts (B) exists (e.g., C to A, Brazil; C to B, Australia), C to A transition is identified by virus cluster 105-115 in phylogenetic tree. In Russia, 91.3% (21/23) of the strains belong to G2 (Figure 3), are mainly present in the ~8500 km long and 600-1700 km wide D climate belt ('Dfa-Dfb-Dw'), suggesting the G2 strains might have adapted to the D climate (Chi-square test, P<.001 for null hypothesis that both G1 and G2 strains are equally found in D climate of Russia, Supplementary Table S6). Similar observations are seen for North America, South America, and Australia. The eastern and western coasts of North America have temperate climate and are connected by cold climate along USA-Canada boundary (i.e., having humid subtropical (Cfa) in eastern coast and marine temperate climate (Cfb) in western coast) (Figure 3). The G2 strains follows this cold climate (Dfa-Dfb) belt which is ~3800 km long and ~600 to 1000 km wide. The dominance of G2 and nearly absence of G1 population in cold climate of North America is similar to the observations of Russia. Our analyses suggests that a fall of temperature from temperate to cold climate might have dictated the evolution of G1 into G2 variant group. Similarly, a change in climate from C to A probably made the strains stable in tropical regions. Overall, our analyses suggest that SARS-CoV-2 has likely evolved to sustain in different climate, thereby increasing its spread. Studies combining genetic information with climate can provides useful information about virus evolution and possible climatic pathways during an outbreak.

Conclusion

It is reasonable to assume COVID-19 transmission pathway and evolution is influenced by climate. Phylogenetic network classified 176 SARS-CoV-2 strains into two variant groups G1 and G2. The G1 strains were habituated to C climate that evolved into G2 by undergoing significant mutations (C241T in leader sequence, F924 in ORF1a, P214L in ORF1b and D614G in S gene), plausibly extended its climatic boundaries from C to D climate, displaying role of natural selection on virus evolution. In our analysis SARS-CoV-2, were found resistive to desert climate (B). Gradually, strains are adapting to A climate in South America. The strains adapted to *"tropical-savannah" (Aw)* climate are a threat to all the tropical countries, which were initially less affected by COVID-19. Nevertheless, due to the uncertainty of COVID-19 data, the results should be carefully interpreted and should not be extrapolated to climate types and climatic conditions other than those analyzed here for the early evolution period. The study agrees that viruses are sensitive to their environment and respond towards naturally occurring abiotic factors such as temperature, latitude and humidity to sustain in different climate of the Earth, which also provides insights about seasonal variations possibly being a strong reason for the spread of other viral diseases as well. Here we showed a more refined description of genes based on phylogenetics and their distribution across different climates. This finer-grained analyses led to highly relevant insights on evolutionary dynamics of poorly understood SARS-CoV-2 genome and provides vital information about the direction of the spread and highlights vulnerable regions of Earth. Such inter-disciplinary studies will play an imperative role in designing antiviral strategies and taking pre-emptive precautionary measures to combat such a pandemic.

References

1. Briz-Redón, Á. & Serrano-Aroca, Á. The effect of climate on the spread of the COVID-19 pandemic: A review of findings, and statistical and modelling techniques. *Prog. Phys. Geogr.* (2020). doi:10.1177/0309133320946302

2. WHO. *Coronavirus disease (COVID-19).* (2020).

3. Vellingiri, B. *et al.* COVID-19: A promising cure for the global panic. *Science of the Total Environment* (2020). doi:10.1016/j.scitotenv.2020.138277

4. Chan, K. H. *et al.* The effects of temperature and relative humidity on the viability of the SARS coronavirus. *Adv. Virol.* (2011). doi:10.1155/2011/734690

5. Chin, A. W. H. *et al.* Stability of SARS-CoV-2 in different environmental conditions. *The Lancet Microbe* (2020). doi:10.1016/s2666-5247(20)30003-3

6. Araújo, M. B., Mestre, F. & Naimi, B. Ecological and epidemiological models are both useful for SARS-CoV-2. *Nature Ecology and Evolution* (2020). doi:10.1038/s41559-020-1246-y

7. Carlson, C. J., Chipperfield, J. D., Benito, B. M., Telford, R. J. & O'Hara, R. B. Species distribution models are inappropriate for COVID-19. *Nature Ecology and Evolution* (2020). doi:10.1038/s41559-020-1212-8

8. Chandra Arya, P. & Bajaj, P. Mini-Review Evolutionary Studies Combined with Species Distribution Models are Useful to Understand COVID-19 Spread and Aid Vaccine Design. *J. Vaccines Vaccin. 1 J Vaccines Vaccin* **11**, 1000438 (2020).

9. Guernier, V., Hochberg, M. E. & Guégan, J. F. Ecology drives the worldwide distribution of human diseases. *PLoS Biol.* **2**, 740–746 (2004).

10. Lauring, A. S. & Andino, R. Quasispecies theory and the behavior of RNA viruses. *PLoS Pathogens* (2010). doi:10.1371/journal.ppat.1001005

11. Racevska, E. Natural Selection. in *Encyclopedia of Animal Cognition and Behavior* (eds. Vonk, J. & Shackelford, T.) 1–14 (Springer International Publishing, 2018). doi:10.1007/978-3-319-47829-6_542-1

12. Fehr, A. R. & Perlman, S. Coronaviruses: An overview of their replication and pathogenesis. in *Coronaviruses: Methods and Protocols* 1–23 (2015). doi:10.1007/978-1-4939-2438-7_1

13. Li, C. K. *et al.* T Cell Responses to Whole SARS Coronavirus in Humans. *J. Immunol.* **181**, 5490–5500 (2008).

14. Li, F., Li, W., Farzan, M. & Harrison, S. C. Structural biology: Structure of SARS coronavirus spike receptor-binding domain complexed with receptor. *Science (80-.).* **309**, 1864–1868 (2005).

15. Liu, J. *et al.* The Membrane Protein of Severe Acute Respiratory Syndrome Coronavirus Acts as a Dominant Immunogen Revealed by a Clustering Region of Novel Functionally and Structurally Defined Cytotoxic T-Lymphocyte Epitopes. *J. Infect. Dis.* **202**, 1171–1180 (2010).

16. Ruch, T. R. & Machamer, C. E. The coronavirus E protein: Assembly and beyond. *Viruses* (2012). doi:10.3390/v4030363

17. Chang, C. K., Hou, M. H., Chang, C. F., Hsiao, C. D. & Huang, T. H. The SARS coronavirus nucleocapsid protein - Forms and functions. *Antiviral Research* (2014). doi:10.1016/j.antiviral.2013.12.009

18. Allaby, M. *Atmosphere. A scientific history of air, weather, and climate. Advances in Space Research* (2009).

19. Peel, M. C., Finlayson, B. L. & McMahon, T. A. Updated world map of the Köppen-Geiger climate classification. *Hydrol. Earth Syst. Sci.* **11**, 1633–1644 (2007).

20. Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* (2020). doi:10.1038/s41586-020-2008-3

21. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**, 1547–1549 (2018).

22. Nei, M. & Saitou, N. The neighbor-joining method: a new method for reco... [Mol Biol Evol. 1987] - PubMed result. *Mol Biol Evol* 406–425 (1987).

23. Tamura, K., Nei, M. & Kumar, S. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 11030–11035 (2004).

24. EarthExplorer.

25. Travel Weather Averages (Weatherbase).

26. Climate data for cities worldwide - Climate-Data.org.

27. Herbei, M., Ciolac, V., Smuleac, A. & Ciolac, L. Georeferencing of Topographical Maps Using the Software ArcGIS. *Res. J. Agric. Sci.* **42**, 595–606 (2010).

28. Harcourt, B. H. *et al.* Identification of Severe Acute Respiratory Syndrome Coronavirus Replicase Products and Characterization of Papain-Like Protease Activity. *J. Virol.* (2004). doi:10.1128/jvi.78.24.13600-13612.2004

29. Snijder, E. J., Decroly, E. & Ziebuhr, J. The Nonstructural Proteins Directing Coronavirus RNA Synthesis and Processing. in *Advances in Virus Research* (2016). doi:10.1016/bs.aivir.2016.08.008

30. Zhang, L. *et al.* The D614G mutation in the SARS-CoV-2 spike protein reduces S1 shedding and increases infectivity. *bioRxiv* 2020.06.12.148726 (2020). doi:10.1101/2020.06.12.148726

31. Korber, B. *et al.* Spike mutation pipeline reveals the emergence of a more transmissible form of SARS-CoV-2. *bioRxiv* (2020). doi:10.1101/2020.04.29.069054

32. Junior, I. J. M. *et al.* The global population of SARS-CoV-2 is composed of six major subtypes. *bioRxiv* (2020). doi:10.1101/2020.04.14.040782

33. Becerra-Flores, M. & Cardozo, T. SARS-CoV-2 viral spike G614 mutation
exhibits higher case fatality rate. *Int. J. Clin. Pract.* (2020).
doi:10.1111/ijcp.13525

34. WHO | SARS (Severe Acute Respiratory Syndrome). Available at:
https://www.who.int/ith/diseases/sars/en/. (Accessed: 10th September 2020)

35. Boël, G. *et al.* Codon influence on protein expression in E. coli correlates with
mRNA levels. *Nature* **529**, 358–363 (2016).

36. Tsai, C. J. *et al.* Synonymous Mutations and Ribosome Stalling Can Lead to
Altered Folding Pathways and Distinct Minima. *Journal of Molecular Biology*
383, 281–291 (2008).

37. Shabalina, S. A., Ogurtsov, A. Y. & Spiridonov, N. A. A periodic pattern of
mRNA secondary structure created by the genetic code. *Nucleic Acids Res.*
(2006). doi:10.1093/nar/gkl287

38. Jia, J. *et al.* Modeling the control of COVID-19: Impact of policy interventions
and meteorological factors. *Electron. J. Differ. Equations* (2020).

39. Bannister-Tyrrell, M., Meyer, A., Faverjon, C. & Cameron, A. Preliminary
evidence that higher temperatures are associated with lower incidence of
COVID-19, for cases reported globally up to 29th February 2020. *medRxiv*
(2020). doi:10.1101/2020.03.18.20036731

40. Nie, Q. *et al.* Phylogenetic and phylodynamic analyses of SARS-CoV-2. *Virus
Res.* (2020). doi:10.1016/j.virusres.2020.198098

41. Forster, P., Forster, L., Renfrew, C. & Forster, M. Phylogenetic network
analysis of SARS-CoV-2 genomes. *Proc. Natl. Acad. Sci. U. S. A.* (2020).
doi:10.1073/pnas.2004999117

42. Bukhari, Q. & Jameel, Y. Will Coronavirus Pandemic Diminish by Summer?
SSRN Electron. J. (2020). doi:10.2139/ssrn.3556998

43. Raamsman, M. J. B. *et al.* Characterization of the Coronavirus Mouse
Hepatitis Virus Strain A59 Small Membrane Protein E. *J. Virol.* (2000).
doi:10.1128/jvi.74.5.2333-2342.2000

44. Sharma, A. *et al.* Structural stability of SARS-CoV-2 degrades with
temperature. *bioRxiv* (2020).

45. Follis, K. E., York, J. & Nunberg, J. H. Furin cleavage of the SARS coronavirus
spike glycoprotein enhances cell-cell fusion but does not affect virion entry.
Virology **350**, 358–369 (2006).

Potential caveats

635 We acknowledge that there are few caveats due to uncertainty in the COVID-19

636 data. The data from the tropical regions is limited because at the time of data

collection (SARS-CoV-2 strains) from all over the world, the strains from the tropical countries were very limited, from few tropical regions strains were available (e.g., Ghana (Africa); India, Mexico, Nepal, Pakistan) but the data has been discarded due to the travel history of the strains, a large fraction of strains without travel history have large gaps in genomic sequences which were not suitable for the present study. Also, case history of each patient is not reported in the METADATA file as collecting all information from each patient is time-consuming. Hence, there are chances patients from whom these strains were isolated may have a migratory history. Data from different individual locations without travel history and large gaps in genomic sequences have been incorporated in this study. Due to these reasons, our analyses should be carefully interpreted and should not be extrapolated to climate types and climatic conditions other than those analyzed here for the early evolution period.

Authors Contribution

Phylogenetic study and genomic analysis is carried out by P.B. GIS study and Koppen's climate map interpretations is done by P.C.A. Statistical analysis is carried out by both the authors. Sequence alignment script was written by both the authors. Both authors have written, reviewed and edited the manuscript.

Acknowledgement

We gratefully acknowledge the authors and originating and submitting laboratories of the sequences from GISAID's EpiFlu (TM) Database on which this research is based. A table of the contributors is available in, Supplementary Table S1. We thank Prof. Raghavan Varadarajan, Prof. Raman Sukumar, Dr. Teena Jangid, and Chetankumar Jalihal of Indian Institute of Science for proofreading the article.

661   **Conflict of Interest**

662   Authors declare no conflict of interest.

664   **TABLES and FIGURES**

666   **Table 1: SNPs representing virus cluster and their distribution across varied**
667   **climates.**

Virus cluster	Nucleotide mutation	Amino acid mutation	Gene	Climate Zone	KCT	KC
1-22	C8782T T28144C	S2839 L84S	ORF1a ORF8	NTZ	Cfa	C
5-6	C29095T	F274	N	NTZ	Cfa	C
8-9	T9477A G25979T C28657T C28863T	F3071Y G196V D128 S197L	ORF1a ORF3a N N	NTZ, TZ	Mix	C-A
10-17	C18060T	L1431	ORF1b	NTZ	Cfa-Cfb	C
12-17	A17858G	Y1364C	ORF1b	NTZ	Cfa-Cfb	C
13-17	C17747T	P1327L	ORF1b	NTZ	Cfa	C
20-22	C24034T T26729C G28077C	N824 A69 V62L	S M ORF8	NTZ	Cfa	C
21-22	T490A C3177T T18736C	D75 P971L F1657L	ORF1a ORF1a ORF1b	NTZ, NTSZ	Cfa	C
23-25	C6312A C13730T C23929T C28311T	T2016K L4489 Y789 P13L	ORF1a ORF1a S N	NTZ, TZ, SSTZ	Mix	Mix
28-32	G1397A T28688C	D392G L139	ORF1a N	NTZ	Mix	Mix
33-43	G26144T	G251V	ORF3a	NTZ	Cfa-Cfb	C
37-39	A2480G C2558T	I739V P765S	ORF1a ORF1a	NTZ	Mix	Mix
37-43	C14805T	Y346	ORF1b	NTZ	Cfa	C
42-43	T17247C	R1160	ORF1b	NTZ	Cfb	B
44-47	G1440A G2891A	G392D A876T	ORF1a ORF1a	NTZ	Cfb	C
58-61	C15324T	N519	ORF1b	NTZ	Cfa-Dfb	C-D
59-176	C3037T A23403G C14408T	F924 D614G P214L	ORF1a S ORF1b	NTZ	Cfa-Cfb-Dfb-Aw	C-D - A

Position	SNP	AA	Gene	Zone	Cluster	KC
59-125,	C241T		Leader seq.	NTZ	Cfa-Cfb-Dfa-Dfb	C-D-A
127-176	C241T					C-D
66-68	A26530G	D3G	M	NTZ	Cfc-Dfb	C-D
70-71	G4201T	M1312I	ORF1a	NTZ	Cfa-Dwc	C-D
	C26527T	A2V	M			
80-115	G28881A	R203K	N	NTZ		
	G28882A	R203K	N			
	G28883C	G204R	N		Cfa-Cfb-Dfb-Aw	C-D-A
86-87	C27046T	T175M	M	NTZ	Cfa-Dfb	C-D
88-89	C3373A	D1036E	ORF1a	NTZ	Dfb-Cfb	C-D
105-107	T29148C	I292T	N	TZ, STZ	Cfa-Aw	C-A
106-107	A27299C	I33T	ORF6	NTZ, TZ	Cfa-Aw	C-A
108-111	C313T	L16	ORF1a	NTZ, TZ	Cfa-Cfb-Aw	C
113-115	C4002T	T1246I	ORF1a	STZ	Cfa-Cfb-Am	C-A
	G10097A	G3278S	ORF1a			
	C13536T	T4424I	ORF1a			
	C23731T	T723	S			
116-125	A20268G	L2167	ORF1b	NTZ	Cfa-Cfb-Dfa-Dfb	C-D
126-176	G25563T	Q57H	ORF3a	NTZ	Cfa-Cfb-Dfa-Dfb	C-D
126-130	C18877T	L1704	ORF1b	NTZ	Cfa-Dfa-Dcb	C
131-135	C2416T	Y717	ORF1a	NTZ	Cfa-Dfa-Aw	D
136-176	C1059T	T265I	ORF1a	NTZ	Cfa-Cfb-Dfa-Dfb	C-D
138-139	C18998T	A1744V	ORF1b	NTZ, TZ	Cfa-Am	C-A
	G29540A					
138-141	C11916T	S3884L	ORF1a	NTZ	Cfa-Csb-Am	C
143-147	C27964T	S24L	ORF8	NTZ	Cfa-Cfb-Dfa-Dfb	C-D
148-149	C11224T	V3653	ORF1a	NTZ	Dfa-Dfb	D
157-159	G29553A			NTZ	Cfa	C

NOTE: Virus clusters are named by Strain ID as depicted on the tree (Supplementary Table
S1 and S2). Genomic coordinates in this study is based on reference genome[20]. The SNP
positions are based on the reference genome. Nucleotide T represents nucleotide U in the
SARS-CoV-2 RNA genome. Mutation at the protein level is not mentioned for the SNPs arising
in the non-coding region. The amino acid position numbering is according to its position within
the specified gene (CDS). In Climate zone column we have mentioned the major climate zone
for the corresponding virus cluster[18]. KCT is Koppen's Climate Type and KC is Koppen's
Climate columns display the main Koppen's climate in which most of the virus isolates of the
corresponding virus cluster lie. 'Mix' implies no particular climate type is favored[19].

Figure 1: Phylogenetic network divides 176 SARS-CoV-2 strains into two variant
groups. Largely, the left side of the tree (1 to 58) constitute the G1 group and the right side
of the tree constitutes the G2 group (59 to 176). Branch length is proportional to the genomic
relatedness of the viral isolates. Closely related virus isolates comprise the same SNP with
respect to the reference genome (Strain ID: 50) and form a cluster. The evolutionary history
of 176 taxa was inferred using the Neighbor-Joining method[22] (500 bootstrap tests). A total of
29408 positions were analysed with nucleotide position numbering according to the reference
sequence[20].

(a)

(b)

(e)

(f)

(g)

SARS CoV-2 genes

Mutant type

NTZ
NSTZ
TZ
SSTZ
STZ

A
B
C
D

Figure 2: Molecular phylogeny analysis to infer genomic similarities of SARS-CoV-2 and their distribution across different climate zones[18] and Koppen's climate types[19]. (a) Genomic architecture of SARS-CoV-2 genome highlighting four positions, substitutions on these positions probably enabled evolution of G1 into G2 variant group. (b, e-g) Strains found within a virus cluster (as shown in the phylogenetic tree and mentioned in Table 1) were analysed for significant mutations that may have arisen due to climatic pressure. Hence, percentage of such virus strains is plotted according to the geographical location of the climate zone from where they were isolated. The height of the bar is proportional to percent virus strain occurring in the specified condition i.e., labelled on the x-axis. Box in the left panel consist of color code for each climate zone and box in the right panel consist of color code for Koppen's climate. Left panel shows distribution of percent virus strains in different climate zones and right panel shows distribution of percent virus strain in Koppen's climate (b) Percent virus strains prevailing in different climate zones, stratified by SARS-CoV-2 variant groups. Width of curves of violin plot is proportional to the number of SARS-CoV-2 strains (n=176) in varied (c) climate zones and (d) Koppen's climate. (e) Abiotic factors influencing evolutionary dynamics of phylogenetic virus clusters. (f) Percent of virus strains with high frequency SNPs in each gene. (g) Type of mutation i.e., non-synonymous (Non-syn) or synonymous (Syn) exhibited by viruses.

Figure 3: Global distribution of SARS-CoV-2 strains on the Gieger-Koppen's map displaying different climate types[19]. Each strain is labelled as per the strain ID (1 to 176) within parenthesis. The G1 strains were symbolized as 'Yellow-circle', and G2 as 'Square', pink square denotes strain clusters (80-115) stable across C, D and A climate, purple square represents strain cluster (126-176) stable majorly in D climate, the remaining G2 strains (blue squares) are stable across C and D climate. Standard Koppen's climate-type symbols are mentioned in the legend, the criteria for distinguishing these climate types, is mentioned in Table S3. Table S4 contains full form of these symbols. All symbols with initials 'A' (Af, Am, Aw) are of tropical climate, initials with 'B' belong to desert climate, 'C' to temperate and 'D' to cold and 'E' to polar climate. The shades of blue on the map, in North America and Russia belongs to D climate. The shades of blue in South America, Africa, and South Asia belong to tropical climate. Shades of yellow and green belongs to C climate. Shades of red, orange and pink belong to Desert climate.

(a)   **(b)**

Figure 4: Global distribution of SARS-CoV-2 strains (n=176) (a) in the coastal and continental region (b) and in different Koppen's climate types[19]. Number of virus strains in G1 population is represented by light grey color and of virus strains in G2 population is represented by dark grey color.

Figure 5: Comparative analysis of different climatic parameters such as latitude, temperature and precipitation between G1 and G2 variant groups. (a) Positive values represent the latitude range falling in Northern Hemisphere and negative values represent latitude range falling in Southern Hemisphere. The G2 strains preferentially occur towards the higher latitudes than G1 (P=.032; 95% CI 17.12-31.12 for G1; 95% CI 28.67-68.06 for G2). (b) The mean annual temperature of G2 is significantly lower than the G1 strains (P<.001; 95% CI 17.32-14.32 for G1; 95% CI 13.02-10.33 for G2) (c) Mean annual precipitation of G1 and G2 strains is nearly same (P=.061; 95% CI 1207.16-886.75 for G1; 95% CI 966.91-826.37 for G2). (a-c) Black horizontal line in the middle of the box is median, upper and lower limits of the box indicate first and third quartile. Black dots represent outliers. P values is based on one-way ANOVA. Scatter plot for (d) latitude and annual temperature and (e) latitude and precipitation for each SARS-CoV-2 strain (n=176) belonging to G1 group (n=58, shown in light grey) and G2 group (n=118, shown in dark grey).