Nonlinear ridge regression improves robustness of cell-type-specific differential expression analysis

Fumihiko Takeuchi and Norihiro Kato

Department of Gene Diagnostics and Therapeutics, Research Institute, National Center for Global Health and Medicine (NCGM), Tokyo, Japan

Correspondence

Fumihiko Takeuchi

Department of Gene Diagnostics and Therapeutics, Research Institute, National Center for Global Health and Medicine (NCGM)

1-21-1 Toyama, Shinjuku-ku, Tokyo, 162-8655, Japan

Email: fumihiko@takeuchi.name

Abstract

Background: Epigenome-wide association studies (EWAS) and differential

gene expression analyses are generally performed on tissue samples, which

consist of multiple cell types. Cell-type-specific effects of a trait, such as

disease, on the omics expression are of interest but difficult or costly to

measure experimentally. By measuring omics data for the bulk tissue, cell

type composition of a sample can be inferred statistically. Subsequently, cell-

type-specific effects are estimated by linear regression that includes terms

representing the interaction between the cell type proportions and the trait.

This approach involves two issues, scaling and multicollinearity.

Results: First, although cell composition is analyzed in linear scale,

differential methylation/expression is analyzed suitably in the logit/log scale.

To simultaneously analyze two scales, we applied nonlinear regression.

Second, we show that the interaction terms are highly collinear, which is

obstructive to ordinary regression. To cope with the multicollinearity, we

applied ridge regularization. In simulated data, nonlinear ridge regression

attained well-balanced sensitivity, specificity and precision. In real data,

nonlinear ridge regression detected signals consistently over the examined

cases.

Conclusion: Nonlinear ridge regression performed cell-type-specific

association test on bulk omics data more robustly than previous methods.

The omicwas package for R implements nonlinear ridge regression for cell-

type-specific EWAS, differential gene expression and QTL analyses. The

software is freely available from https://github.com/fumi-github/omicwas

Keywords

Epigenome-wide association study, Differential gene expression analysis, Cell

type, Nonlinear regression, Ridge regression, mQTL, eQTL

Background

Epigenome-wide association studies (EWAS) and differential gene expression analyses elucidate the association of disease traits (or conditions) with the level of omics expression, namely DNA methylation and gene expression. Thus far, tissue samples, which consist of heterogeneous cell types, have mainly been examined, because cell sorting is not feasible in most tissues and single-cell assay is still expensive. Nevertheless, the cell type composition of a sample can be quantified statistically by comparing omics measurement of the target sample with reference data obtained from sorted or single cells [1,2]. By utilizing the composition, the disease association specific to a cell type was statistically inferred for gene expression [3-10] and DNA methylation [11-14].

For the imputation of cell type composition, omics markers are usually analyzed in the original linear scale, which measures the proportion of mRNA molecules from a specific gene or the proportion of methylated cytosine molecules among all cytosines at a specific CpG site [15]. The proportion can differ between cell types, and the weighted average of cell-type-specific proportions becomes the proportion in a bulk tissue sample. Using the fact that the weight equals the cell type composition, the cell type composition of a sample is imputed. In contrast, gene expression analyses are performed in the log-transformed scale because the signal and noise are normally distributed after log-transformation [16]. In DNA methylation analysis, the logit-transformed scale, which is called the M-value, is statistically valid [17]. Consequently, the optimal scales for analyzing differential gene expression or methylation can differ from the optimal scale for analyzing cell type composition.

Aiming to perform cell-type-specific EWAS or differential gene expression analyses by using unsorted tissue samples, we study two issues that have been overlooked. Whereas previous studies were performed in linear scale, we develop a nonlinear regression, which simultaneously analyzes cell type composition in linear scale and differential expression/methylation in log/logit scale. The second issue is multicollinearity. Cell-type-specific effects of a trait, such as disease, on omics expression are usually estimated by linear

63 regression that includes terms representing the interaction between the cell
64 type proportions and the trait. We show that the interaction terms can
65 mutually be highly correlated, which obstructs ordinary regression. To cope
66 with the multicollinearity, we implement ridge regularization. Our methods
67 and previous ones are compared in simulated and real data.
68

Results

Multicollinearity of interaction terms

Typically, cell-type-specific effects of a trait on omics marker expression is
analyzed by the linear regression in equation (2). For each omics marker, the
goal is to estimate $\beta_{h,k}$, the effect of trait k on the expression level in cell type
h. This is estimated based on the relation between the bulk expression level
Y_i of sample i and the regressor $W_{h,i}X_{i,k}$, which is an interaction term defined
as the product of the cell type proportion $W_{h,i}$ and the trait value $X_{i,k}$ of the
sample. We assume that Y_i, $W_{h,i}$ and $X_{i,k}$ are given as input data.
The variable $W_{h,i}$ for cell type composition cannot be mean-centered for
our purpose. If $W_{h,i}$ were centered, we would obtain, instead of $\beta_{h,k}$, the
deviation of $\beta_{h,k}$ from the average across cell types. In general, interaction
terms involving uncentered variables can become collinear [18]. We first
survey the extent of multicollinearity in real data for cell-type-specific
association.
In peripheral blood leukocyte data from a rheumatoid arthritis study
(GSE42861), the proportion of cell types ranged from 0.59 for neutrophils to
0.01 for eosinophils (Table 1A). The proportion of neutrophils was negatively
correlated with the proportion of other cell types (apart from monocytes) with
correlation coefficient of −0.68 to −0.46, whereas the correlation was weaker
for other pairs (Table 1B). Rheumatoid arthritis status was modestly
correlated with proportions of cell types. The product of the disease status
X_k, centered to have zero mean, and the proportion of a cell type becomes
an interaction term. The correlation coefficients between the interaction
terms were mostly >0.8, apart from eosinophils (Table 1C). The coefficient

of variation (CV), which is the ratio of standard deviation to mean, of the
proportion was low for all cell types apart from eosinophils (Table 1A). The
interaction terms for low-CV cell types were strongly correlated with X_k,
which in turn caused strong correlation between the relevant interaction
terms.

The situation was the same for the interaction with age in GTEx data. The
granulocytes (which include neutrophils and eosinophils) were the most
abundant (Table 2A). The proportion of granulocytes was negatively
correlated with other cell types (apart from monocytes) with correlation
coefficient of −0.89 to −0.41, and the correlation between other pairs was
generally weaker (Table 2B). Age was modestly correlated with proportions
of cell types. In this dataset, the CV of the proportion was low in all cell types
(Table 2A), which caused strong mutual correlation between interaction
terms (Table 2C).

In the above empirical data, multicollinearity between interaction terms
seemed to arise not due to the correlation between cell type proportions or
X_k, but due to the low CV in the cell type proportions. Subsequently, this
property was derived mathematically. As we derived in equation (17), the
correlation between interaction terms $W_h X_k$ and $W_{h'} X_k$ approaches one when
$CV[W_h]$ and $CV[W_{h'}]$ are low, irrespective of $Cor[W_h, W_{h'}]$ (Fig. 1). The CV was
0.2 to 0.6 (apart from eosinophils) in the rheumatoid arthritis dataset and
0.1 to 0.2 in the GTEx dataset. We looked up datasets of several ethnicities
and found the CV to be ≤0.6 in majority of blood cell types (Additional file 1:
Table S1). Thus, multicollinearity can be a common problem for cell-type-
specific association analyses. Biologically, in tissues where cell type
composition is tightly controlled, the CV of cell type proportion becomes low
and the multicollinearity is exacerbated.

Evaluation in simulated data

By using simulated data, we evaluated previous methods and new
approaches of the omicwas package. In order to simultaneously analyze two
scales, the linear scale for heterogeneous cell mixing and the log/logit scale
for trait effects, we applied nonlinear regression (equations (4) and (5)). To

cope with the multicollinearity of interaction terms, we applied ridge
regularization (formula (8)).

Previous regression type methods are based either on the full model of
linear regression (equation (2)) or the marginal model (equation (3)). The
full model fits and tests cell-type-specific effects for all cell types
simultaneously, and its variations include TOAST, csSAM.lm and CellDMC. The
marginal model fits and tests cell-type-specific effect for one cell type at a
time, and its variations include csSAM.monovariate and TCA. We also
examined a hybrid of the two models (Marginal.Full005), which becomes
positive when the models agree.

The simulation data was generated from real datasets of DNA methylation
(658 samples; 451,725 CpG sites) and gene expression (389 samples;
14,038 genes). The original cell type composition was retained for all samples,
and the case-control status was randomly assigned. Ninety-five percent of
omics markers were set to be unassociated with disease status, 2.5% were
up-regulated in cases at one cell type, and 2.5% were similarly down-
regulated. The cell-type-specific effect-size was fixed in a simulation trial,
either to methylation odds ratio of 1.3, 1.6 or 1.9 or to gene expression fold
change of 1.7, 3.0 or 5.0. The significance level was set to $P < 2.4 \times 10^{-7}$ for
DNA methylation and false discovery rate <5% for gene expression. In each
simulation trial, the sensitivity, specificity and precision for detecting cell-
type-specific association was calculated for each cell type. To compare
algorithms, the performance measures for the same effect-size and cell type
were averaged over the simulation trials.

Overall, in the simulation for DNA methylation the sensitivity (Fig. 2) was
higher under large effect-size (bottom row of panels) and in abundant cell
types (left columns of panels). The average specificity (Fig. 3) was high
across the effect-size settings and across cell types, being >0.97 for the
Marginal model, >0.98 for TCA and >0.999 for other algorithms. The
precision (Fig. 4) was higher in abundant cell types within each effect-size
setting. As effect-size increased, the precision decreased in neutrophils but
increased in the other minor cell types. Excluding the cases where all
algorithms lacked sensitivity (monocytes and B cells under methOR=1.3 and

eosinophils), the average precision of omicwas.logit.ridge was >0.78 and was
the highest in 13/16 of the cases.

There was trade-off between sensitivity and precision. Among the
algorithms, the Marginal model attained the highest sensitivity and the lowest
precision. TCA, which is a variation of the marginal model, had relatively high
sensitivity and relatively low precision. Marginal.Full005 attained the second
highest sensitivity and moderate precision. The ridge regressions
(omicwas.logit.ridge and omicwas.identity.ridge) attained moderate
sensitivity and high precision. The full models without ridge regularization
(omicwas.logit, omicwas.identity, Full, TOAST and CellDMC) had the lowest
sensitivity.

The overall tendency was similar in the simulation for gene expression.
The sensitivity (Fig. 5) was higher under large effect-size and in abundant
cell types. The average specificity (Fig. 6) was high across the effect-size
settings and across cell types, being >0.96 for the marginal models (Marginal
and csSAM.monovariate), >0.98 for the nonlinear and ridge regressions
(omicwas.log.ridge, omicwas.identity.ridge, omicwas.log) and >0.996 for
other algorithms. The precision (Fig. 7) was higher in abundant cell types
within each effect-size setting. As effect-size increased, the precision
decreased in granulocytes but increased in the other minor cell types.
Excluding the full models (omicwas.identity, Full, TOAST, csSAM.lm) that
lacked sensitivity, the algorithms that were frequently top in average
precision were omicwas.identity.ridge (5 cases), omicwas.log (5 cases),
omicwas.log.ridge (3 cases) and Marginal.Full005 (3 cases).

There again was trade-off between sensitivity and precision. Among the
algorithms, the marginal models attained the highest sensitivity but relatively
low precision. The nonlinear and ridge regressions and Marginal.Full005
attained moderate sensitivity and highest precision. The full models had very
low average sensitivity of <0.01.

For gene expression, we also simulated a scenario where cell-type-specific
disease effect occurred in cell type "marker" genes (Additional files 2, 3, 4:
Figs. S1, S2, S3). In other words, the expression level in the target cell type
differed between cases and controls, and the expression level in other cell

types was zero (in linear scale). Thus, non-target cell types did not introduce
noise to bulk expression level. The Marginal model attained the highest
average sensitivity of >0.93 but relatively low average precision of ~0.16. As
equal number of differentially expressed genes were generated in the six cell
types, picking up signals for all such genes, including those not for the tested
cell type, would result in precision of 1/6 = 0.16. The full models had low
sensitivity. The nonlinear and ridge regressions and Marginal.Full005 attained
moderate sensitivity and moderate precision. With regards to the frequency
of being the top in average precision, the algorithms were ordered
Marginal.Full005   (9   cases),   omicwas.log.ridge   (6   cases),
omicwas.identity.ridge (2 cases) and omicwas.log (1 case), excluding the full
models that lacked sensitivity.

204 Cell-type-specific association with rheumatoid arthritis and age

The detection of cell-type-specific association in bulk tissue was evaluated by
using physically sorted cells. In principle, sorted cells should serve as genuine
verification, however, due to the relatively small sample size (94 or 203 for
rheumatoid arthritis and 214 or 1202 for age) the available datasets were
underpowered to generate a gold standard list of differentially expressed
omics markers [19]. Instead, we generated a benchmark set of differentially
expressed markers by imposing a relaxed significance level of $P < 0.05$; the
set would be enriched for true differentially expressed markers yet also
include unassociated markers. The benchmark set was cross-checked with
the prediction by each algorithm; the prediction cutoff value was varied and
the performance was measured by the area under the curve (AUC) of receiver
operating characteristic (ROC). In other words, we examined if the prediction
in bulk tissue by an algorithm could be replicated in sorted cells under $P <$
0.05.

The cell-type-specific association of DNA methylation with rheumatoid
arthritis was predicted using bulk peripheral blood leukocyte data and was
evaluated in sorted monocytes (Fig. 8A) and B cells (Fig. 8B). Whereas the
full model (and its variations) performed the best and the marginal model
(and its variations) performed the worst in monocytes, the performance

ranking was opposite in B cells. A robust algorithm would consistently achieve
high performance relative to the best algorithm in each instance. Nonlinear
ridge regression (omicwas.logit.ridge) was the most robust, performing 65%
to 93% relative to the best method. The hybrid Marginal.Full005 did not
perform well.

The cell-type-specific association of gene expression with age was
predicted using whole blood data and was evaluated in sorted CD4$^+$ T cells
(Fig. 8C) and monocytes (Fig. 8D). All algorithms performed poorly in CD4$^+$
T cells, and the marginal model performed the best in monocytes. Overall,
nonlinear ridge regression (omicwas.log.ridge) was next to the marginal
model, performing 21% to 47% to the marginal. The dataset of sorted CD4$^+$
T cells (214 samples) is smaller than the monocyte dataset (1202 samples)
thus could be underpowered to pick enough true differentially expressed
genes into the benchmark set.

For DNA methylation dataset GSE42861 and for GTEx gene expression
dataset, the omicwas.logit.ridge and omicwas.log.ridge models of the
omicwas package was computed in 8.1 and 0.7 hours respectively, using 8
cores of a 2.5 GHz Xeon CPU Linux server.
242

Discussion
Aiming to elucidate cell-type-specific trait association in DNA methylation and
gene expression, this article explored two aspects, multicollinearity and scale.
We observed multicollinearity in real data and derived mathematically how it
emerges. To cope with the multicollinearity, we applied ridge regularization.
To properly handle multiple scales simultaneously, we applied nonlinear
regression. Among the examined algorithms, nonlinear ridge regression
attained moderate sensitivity and highest precision in simulated data, and
achieved robust performance in real data. We also developed an algorithm
that combines full and marginal models, which attained balanced sensitivity
and precision in simulation.

The statistical methods discussed in this article are applicable, in principle,
to any tissue. For validation of the methods, we need datasets for bulk tissue

as well as sorted cells, ideally of several hundred samples. Currently, the
publicly available data is limited to peripheral blood. By no means, we claim
the rheumatoid arthritis EWAS datasets [20-22] or the datasets for age
association of gene expression [23,24] to be representative. Nevertheless,
we think verification in real data is valuable.

By the performance in simulated and real data, we can roughly divide
algorithms into three groups: full (and its variations), marginal (and its
variations) and the third group that includes ridge regressions and the hybrid
Marginal.Full005. In marginal models, we test one cell type at a time. If we
knew in advance that one particular cell type is associated with the trait,
which would be a rare situation, testing that cell type with the marginal model
is the most simple and correct approach. However, when the test target cell
type is not associated, but instead another cell type is associated, the
marginal models can pick up false signals due to the collinearity between
regressor variables. Indeed, marginal models attained highest sensitivity
(Figs. 2, 5) and relatively low precision (Figs. 4, 7), which could lead to
unstable performance. In real data, the marginal models were the most
powerful for detecting B cell specific association with rheumatoid arthritis (Fig.
8B) but were the least powerful for monocytes (Fig. 8A). The full model fits
and tests all cell types simultaneously, by which it adjusts for the effects of
other cell types. Due to the simultaneous inclusion of collinear predictors, the
sensitivity was low (Figs. 2, 5). The ridge regressions (omicwas.identity.ridge,
omicwas.logit.ridge and omicwas.log.ridge) were in the middle between full
and marginal models with regards to the sensitivity (Figs. 2, 5), while
attaining the highest specificity (Figs. 4, 7). The nonlinear ridge regressions
consistently achieved high performance relative to the best algorithm in real
data (Fig. 8). The hybrid Marginal.Full005 algorithm is intended to gain
sensitivity by the marginal model while keeping precision by incorporating
the full model. It attained moderate sensitivity (Figs. 2, 5) and moderate
precision (Figs. 4, 7) in simulation but did not perform well in real data (Fig.
8).

Conclusions

For cell-type-specific differential expression analysis by using unsorted tissue samples, we recommend trying the nonlinear ridge regression as a first choice because it balances sensitivity and precision and was robust in experimental data. Although marginal models can be powerful when the tested cell type actually is the only one associated with the trait, caution is needed due to its low precision. Under the idea of first scanning by the marginal model and then reanalyzing in full model, we developed the hybrid Marginal.Full005 algorithm, which attained balanced sensitivity and precision but was not corroborated in experimental data. Ridge regression is preferable compared to the full model without ridge regularization because ridge estimator of the effect-size has smaller mean squared error (equation (13)). We do not claim the ridge regression to substitute previous algorithms. Indeed, we think none of the current algorithms is superior to others in all aspects, indicating possibility for future improvement.

Methods

Linear regression

We begin by describing the linear regressions used in previous studies. Let the indexes be h for a cell type, i for a sample, j for an omics marker (CpG site or gene), k for a trait that has cell-type-specific effects on marker expression, and l for a trait that has a uniform effect across cell types. The input data is given in four matrices. The matrix $W_{h,i}$ represents cell type composition. The matrices $X_{i,k}$ and $C_{i,l}$ represent the values of the traits that have cell-type-specific and uniform effects, respectively. We assume the two matrices are centered: $\sum_i X_{i,k} = \sum_i C_{i,l} = 0$. For example, $X_{i,k} = 0.5$ for disease cases and $X_{i,k} = -0.5$ for controls when the number of cases and controls are equal. The matrix $Y_{i,j}$ represents the omics marker expression level in tissue samples.

The parameters we estimate are the cell-type-specific trait effect $\beta_{h,j,k}$, tissue-uniform trait effect $\gamma_{j,l}$, and basal marker level $\alpha_{h,j}$ in each cell type.

320 For the remaining of the first five sections (up to "Multicollinearity of
321 interaction terms"), we focus on one marker j, and omit the index for
322 readability. For cell type h, the marker level of sample i is

$$\alpha_h + \sum_k \beta_{h,k} X_{i,k}. \quad (1)$$

324 This is a representative value rather than a mean because we do not model
325 a probability distribution for cell-type-specific expression. By averaging the
326 value over cell types with weight $W_{h,i}$, and combining with the tissue-uniform
327 trait effects, we obtain the mean marker level in bulk tissue of sample i,

$$\mu_i = \sum_h \alpha_h W_{h,i} + \sum_{h,k} \beta_{h,k} W_{h,i} X_{i,k} + \sum_l \gamma_l C_{i,l}.$$

329 With regards to the statistical model, we assume the error of the marker
330 level to be normally distributed with variance σ^2, independently among
331 samples, as

$$Y_i = \mu_i + \varepsilon_i,$$

$$\varepsilon_i \sim N(0, \sigma^2).$$

334 The statistical significance of all parameters is tested under the *full* model of
335 linear regression,

$$Y_i = \sum_h \alpha_h W_{h,i} + \sum_{h,k} \beta_{h,k} W_{h,i} X_{i,k} + \sum_l \gamma_l C_{i,l} + \varepsilon_i, \quad (2)$$

337 or its variations [5,9,13]. Alternatively, the cell-type-specific effects of traits
338 can be fitted and tested for one cell type h at a time by the *marginal* model,

$$Y_i = \sum_{h'} \alpha_{h'} W_{h',i} + \sum_k \beta_{h,k} W_{h,i} X_{i,k} + \sum_l \gamma_l C_{i,l} + \varepsilon_i, \quad (3)$$

or its variations [7,8,10,11,14].

Nonlinear regression

Aiming to simultaneously analyze cell type composition in linear scale and
differential expression/methylation in log/logit scale, we develop a nonlinear
regression model. The differential analyses are performed after applying
normalizing transformation. The normalizing function is the natural logarithm
f = log for gene expression, and f = logit for methylation (see Background).
Conventional linear regression can be formulated by defining f as the identity

function. We denote the inverse function of f by g; g = exp for gene

expression, and g = logistic for methylation. Thus, f converts from the linear

scale to the normalized scale, and g does the opposite.

The marker level in a specific cell type (formula (1)) is modeled in the

normalized scale. The level is linearized by applying function g, then averaged

over cell types with weight $W_{h,i}$, and normalized by applying function f.

Combined with the tissue-uniform trait effects, the mean normalized marker

level in bulk tissue of sample i becomes

356
$$\mu_i = f\left(\sum_h W_{h,i}\, g\left(\alpha_h + \sum_k \beta_{h,k} X_{i,k}\right)\right) + \sum_l \gamma_l C_{i,l}. \quad (4)$$

357   We assume the normalized marker level to have an error that is normally

358 distributed with variance σ^2, independently among samples, as

359
$$f(Y_i) = \mu_i + \varepsilon_i, \quad (5)$$

360
$$\varepsilon_i \sim N(0, \sigma^2).$$

361 We obtain the ordinary least squares (OLS) estimator of the parameters by

362 minimizing the residual sum of squares,

363
$$\text{RSS} = \sum_i (f(Y_i) - \mu_i)^2, \quad (6)$$

364 and then estimate the error variance as

365
$$\widehat{\sigma^2} = \frac{1}{n - p}\text{RSS}, \quad (7)$$

where n is the number of samples and p is the number of parameters [[25],

section 6.3.1].

Ridge regression

The parameters $\beta_{h,k}$ for cell-type-specific effect cannot be estimated

accurately by ordinary linear regression because the regressors $W_{h,i} X_{i,k}$ in

equation (2) are highly correlated between cell types (see below).

Multicollinearity also occurs to the nonlinear case in formula (4) because of

local linearity. To cope with the multicollinearity, we apply ridge regression

with a regularization parameter $\lambda \geq 0$, and obtain the ridge estimator of the

parameters that minimizes

$$376 \qquad \text{RSS} + \lambda \sum_{h,k} {\beta_{h,k}}^2 , \quad (8)$$

377   where the second term penalizes $\beta_{h,k}$ for taking large absolute values. The

378   ridge estimator $\hat{\boldsymbol{\theta}}(\lambda)$ is asymptotically normally distributed (see Additional file

379   5: Supplementary note) with

$$380 \qquad \text{Mean}\big[\hat{\boldsymbol{\theta}}(\lambda)\big] = Q(\lambda)^{-1}\, Q(0)\, \boldsymbol{\theta}, \quad (9)$$

$$381 \qquad \text{Var}\big[\hat{\boldsymbol{\theta}}(\lambda)\big] = \sigma^2\, Q(\lambda)^{-1} \left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}\right)^{T} \left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}\right) Q(\lambda)^{-1}, \quad (10)$$

$$382 \qquad Q(\lambda) = \left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}\right)^{T} \left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}\right) + \lambda \begin{pmatrix} O & O & O \\ O & I & O \\ O & O & O \end{pmatrix} - (f(Y) - \boldsymbol{\mu}(\boldsymbol{\theta}))^{T} \cdot \left(\frac{\partial^2 \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^{T}}\right),$$

where $\boldsymbol{\mu}$ is the vector form of μ_i, $\boldsymbol{\theta}$ is the vector form of the parameters α_h,

$\beta_{h,k}$ and γ_l combined, $(\partial \boldsymbol{\mu}/\partial \boldsymbol{\theta})$ is the Jacobian matrix, $(\partial^2 \boldsymbol{\mu}/\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^{T})$ is the

array of Hessian matrices for μ_i taken over samples, and superscript T

indicates matrix transposition. The dot product of $(f(Y) - \boldsymbol{\mu}(\boldsymbol{\theta}))^{T}$ and the

array of Hessians is taken by multiplying for each sample and then summing

up over samples. The matrix after λ has one only in the diagonal

corresponding to $\beta_{h,k}$. The assigned value $\boldsymbol{\theta}$ is the true parameter value. By

taking the expectation of Q, we obtain a rougher approximation [26] as

$$391 \qquad \text{Mean}\big[\hat{\boldsymbol{\theta}}(\lambda)\big] = Q^{*}(\lambda)^{-1}\, Q^{*}(0)\, \boldsymbol{\theta}, \quad (11)$$

$$392 \qquad \text{Var}\big[\hat{\boldsymbol{\theta}}(\lambda)\big] = \sigma^2\, Q^{*}(\lambda)^{-1} \left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}\right)^{T} \left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}\right) Q^{*}(\lambda)^{-1}, \quad (12)$$

$$393 \qquad Q^{*}(\lambda) = \text{E}[Q(\lambda)] = \left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}\right)^{T} \left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}\right) + \lambda \begin{pmatrix} O & O & O \\ O & I & O \\ O & O & O \end{pmatrix}.$$

The matrices Q and Q^{*} are the observed and expected Fisher matrices

multiplied by σ^2 and adapted to ridge regression, respectively.

Since our objective is to predict the cell-type-specific trait effects, we

choose the regularization parameter λ that can minimize the mean squared

error (MSE) of $\beta_{h,k}$. Our methodology is based on [27]. To simplify the

explanation, we assume the Jacobian matrices $(\partial \boldsymbol{\mu}(\boldsymbol{\theta})/\partial \boldsymbol{\alpha})$, $(\partial \boldsymbol{\mu}(\boldsymbol{\theta})/\partial \boldsymbol{\beta})$ and

$(\partial \boldsymbol{\mu}(\boldsymbol{\theta})/\partial \boldsymbol{\gamma})$ to be mutually orthogonal, where $\boldsymbol{\alpha}$, $\boldsymbol{\beta}$ and $\boldsymbol{\gamma}$ are the vector

forms of α_h, $\beta_{h,k}$ and γ_l, respectively. Then, from formulae (11) and (12), the

ridge estimator $\hat{\boldsymbol{\beta}}(\lambda)$ is asymptotically normally distributed with

$$403 \qquad \text{Mean}[\hat{\boldsymbol{\beta}}(\lambda)] = \left[\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right) + \lambda I\right]^{-1} \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)\boldsymbol{\beta},$$

$$404 \qquad \text{Var}[\hat{\boldsymbol{\beta}}(\lambda)] = \sigma^2 \left[\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right) + \lambda I\right]^{-1} \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)$$

$$405 \qquad \left[\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right) + \lambda I\right]^{-1},$$

406 where the assigned values $\boldsymbol{\theta}$ and $\boldsymbol{\beta}$ are the true parameter values. We apply
407 singular value decomposition

$$408 \qquad \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right) = UDV^T,$$

409 where U and V are orthogonal matrices, the columns of V are $\boldsymbol{v}_1, \cdots, \boldsymbol{v}_M$, and
410 the diagonals of diagonal matrix D are sorted $d_1 \geq \cdots \geq d_M \geq 0$. The bias,
411 variance and MSE of the ridge estimator are decomposed as

$$412 \qquad \text{Bias}[\hat{\boldsymbol{\beta}}(\lambda)] = \text{E}[\hat{\boldsymbol{\beta}}(\lambda) - \boldsymbol{\beta}]$$

$$413 \qquad = -\lambda \left[\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right) + \lambda I\right]^{-1} \boldsymbol{\beta}$$

$$414 \qquad = \left\{\sum_{m=1}^{M} \boldsymbol{v}_m \frac{-\lambda}{d_m^2 + \lambda} \boldsymbol{v}_m^T\right\}\boldsymbol{\beta},$$

$$415 \qquad \text{Var}[\hat{\boldsymbol{\beta}}(\lambda)] = \sigma^2 \sum_{m=1}^{M} \boldsymbol{v}_m \frac{d_m^2}{\left(d_m^2 + \lambda\right)^2} \boldsymbol{v}_m^T,$$

$$416 \qquad \text{MSE}[\hat{\boldsymbol{\beta}}(\lambda)] = \text{E}\left[\|\hat{\boldsymbol{\beta}}(\lambda) - \boldsymbol{\beta}\|^2\right]$$

$$417 \qquad = \|\text{Bias}[\hat{\boldsymbol{\beta}}(\lambda)]\|^2 + \text{tr}\left(\text{Var}[\hat{\boldsymbol{\beta}}(\lambda)]\right)$$

$$418 \qquad = \sum_{m=1}^{M} \left(\frac{\lambda}{d_m^2 + \lambda}\right)^2 (\boldsymbol{v}_m^T\boldsymbol{\beta})^2 + \left(\frac{d_m^2}{d_m^2 + \lambda}\right)^2 \left(\frac{\sigma^2}{d_m^2}\right). \quad (13)$$

For each m in the summation of (13), the minimum of the summand is
attained at $\lambda_m = \sigma^2/(\boldsymbol{v}_m^T\boldsymbol{\beta})^2$. To minimize MSE, we need to find some
"average" of the optimal λ_m over the range of m. Hoerl et al. [28] proposed
to use the harmonic mean $\lambda = M\sigma^2/\|\boldsymbol{\beta}\|^2$. However, if an OLS estimator $\hat{\boldsymbol{\beta}}(0)$
is actually plugged into $\|\boldsymbol{\beta}\|^2$, the denominator is biased upwards, and the
computed mean is biased downwards. Indeed, with regards to the estimator
of $1/\sqrt{\lambda_m}$, we notice that

$$426 \qquad \frac{1}{\sigma}\boldsymbol{v}_m^T\hat{\boldsymbol{\beta}}(0) \sim N\left(\frac{1}{\sigma}\boldsymbol{v}_m^T\boldsymbol{\beta}, \frac{1}{d_m^2}\right),$$

427   where the terms with larger m have larger variance. Thus, we take the

428   average of $(v_m^T\hat{\beta}(0))^2/\sigma^2$, weighted by $d_m^2/\sum_{m=1}^M d_m^2$, and also subtract the

429   upward bias as,

430
$$\kappa = \frac{1}{\sum_{m=1}^M d_m^2} \sum_{m=1}^M \left\{ \frac{d_m^2\left(v_m^T\hat{\beta}(0)\right)^2}{\sigma^2} - 1 \right\}. \quad (14)$$

431   The weighting and subtraction were mentioned in [27], where the subtraction

432   term was dismissed, under the assumption of large effect-size β. Since the

433   effect-size could be small in our application, we keep the subtraction term.

434   The statistic κ can be nonpositive, and is unbiased in the sense that

435
$$\mathrm{E}[\kappa] = \frac{1}{\sum_{m=1}^M d_m^2} \sum_{m=1}^M \frac{d_m^2(v_m^T\beta)^2}{\sigma^2} = \frac{1}{\sum_{m=1}^M d_m^2} \sum_{m=1}^M \frac{d_m^2}{\lambda_m}$$

436   equals the weighted sum of $1/\lambda_m$. Our choice of regularization parameter is

437
$$\lambda = \begin{cases} 1/\kappa & \text{if } \kappa > 0, \\ d_1^2 & \text{otherwise}, \end{cases} \quad (15)$$

where d_1^2 is taken instead of positive infinity.

Implementation of omicwas package

For each omics marker, the parameters α, β and γ (denoted in combination

by θ) are estimated and tested by nonlinear ridge regression in the following

steps. As we assume the magnitude of trait effects β and γ to be much

smaller than that of basal marker level α, we first fit α alone for numerical

stability.

1. Compute OLS estimator $\hat{\alpha}(0)$ by minimizing formula (6) under $\beta = \gamma = 0$.

Apply Wald test.

2. Calculate $\widehat{\sigma^2}$ by formula (7). Use it as a substitute for σ^2. The residual

degrees of freedom $n - p$ is the number of samples minus the number of

parameters in α.

3. Compute OLS estimators $\hat{\beta}(0)$ and $\hat{\gamma}(0)$ by minimizing formula (6) under

$\alpha = \hat{\alpha}(0)$. Let $\hat{\theta}(0) = (\hat{\alpha}(0)^T, \hat{\beta}(0)^T, \hat{\gamma}(0)^T)^T$.

4. Apply singular value decomposition $(\partial\mu(\hat{\theta}(0))/\partial\beta) = UDV^T$.

5. Calculate κ and then the regularization parameter λ by formulae (14)

and (15).

6. Compute ridge estimators $\widehat{\boldsymbol{\beta}}(\lambda)$ and $\widehat{\boldsymbol{\gamma}}(\lambda)$ by minimizing formula (8) under

$\boldsymbol{\alpha} = \widehat{\boldsymbol{\alpha}}(0)$. Let $\widehat{\boldsymbol{\theta}}(\lambda) = (\widehat{\boldsymbol{\alpha}}(0)^T, \widehat{\boldsymbol{\beta}}(\lambda)^T, \widehat{\boldsymbol{\gamma}}(\lambda)^T)^T$.

7. Approximate the variance of ridge estimator, according to formula (10),

by

459
$$\text{Var}\left[\begin{pmatrix}\widehat{\boldsymbol{\beta}}(\lambda)\\\widehat{\boldsymbol{\gamma}}(\lambda)\end{pmatrix}\right] = \widehat{\sigma^2}\, Q(\lambda)^{-1} \left(\frac{\partial \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}}\right)^T \left(\frac{\partial \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}}\right) Q(\lambda)^{-1},$$

460
$$Q(\lambda) = \left(\frac{\partial \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}}\right)^T \left(\frac{\partial \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}}\right) + \lambda \begin{pmatrix}I & O\\O & O\end{pmatrix}$$

461
$$-\left(f(Y) - \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)\right)^T \cdot \left(\frac{\partial^2 \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix} \partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}^T}\right).$$

462   8. Apply the "non-exact" t-type test [29]. For the s-th coordinate,

464
$$\frac{\begin{pmatrix}\widehat{\boldsymbol{\beta}}(\lambda)\\\widehat{\boldsymbol{\gamma}}(\lambda)\end{pmatrix}_s}{\sqrt{\text{Var}\left[\begin{pmatrix}\widehat{\boldsymbol{\beta}}(\lambda)\\\widehat{\boldsymbol{\gamma}}(\lambda)\end{pmatrix}\right]_{s,s}}} \sim t_{n-p}, \quad (16)$$

under the null hypothesis $\begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}_s = 0$.

The formula (16) is the same as a Wald test, but the test differs, because the

ridge estimators are not maximum-likelihood estimators. The algorithm was

implemented as a package for the R statistical language. We used the NL2SOL

algorithm of the PORT library [30] for minimization.

In analyses of quantitative trait locus (QTL), such as methylation QTL

(mQTL) and expression QTL (eQTL), an association analysis that takes the

genotypes of a single nucleotide polymorphism (SNP) as $X_{i,k}$ is repeated for

many SNPs. In order to speed up the computation, we perform rounds of

linear regression. First, the parameters $\widehat{\boldsymbol{\alpha}}(0)$ and $\widehat{\boldsymbol{\gamma}}(0)$ are fit by ordinary

linear regression under $\boldsymbol{\beta} = \mathbf{0}$, which does not depend on $X_{i,k}$. By taking the

residuals, we practically dispense with $\widehat{\boldsymbol{\alpha}}(0)$ and $\widehat{\boldsymbol{\gamma}}(0)$ in the remaining steps.

Next, for $X_{i,k}$ of each SNP, $\widehat{\boldsymbol{\beta}}(0)$ is fit by ordinary linear regression under $\boldsymbol{\alpha} =$

$\widehat{\boldsymbol{\alpha}}(0)$, $\boldsymbol{\gamma} = \widehat{\boldsymbol{\gamma}}(0)$. The regularization parameter λ is computed according to

steps 4 and 5 above. Finally, $\widehat{\boldsymbol{\beta}}(\lambda)$ is fitted and tested by linear ridge
regression under $\boldsymbol{\alpha} = \widehat{\boldsymbol{\alpha}}(0)$, $\boldsymbol{\gamma} = \widehat{\boldsymbol{\gamma}}(0)$.

Multicollinearity of interaction terms

The regressors for cell-type-specific trait effects in the full model (equation
(2)) are the interaction terms $W_{h,i}X_{i,k}$. To assess multicollinearity, we
mathematically derive the correlation coefficient between two interaction
terms $W_{h,i}X_{i,k}$ and $W_{h',i}X_{i,k}$. In this section, we treat $W_{h,i}$, $W_{h',i}$ and $X_{i,k}$ as
sampled instances of random variables W_h, $W_{h'}$ and X_k, respectively; note
that the sample index i is omitted. For simplicity, we assume W_h and $W_{h'}$ are
independent of X_k. Let $E[\bullet]$, $Var[\bullet]$, $Cov[\bullet]$, $Cor[\bullet]$ and $CV[\bullet]$ denote the
expectation, variance, covariance, correlation and coefficient of variation,
respectively. Since X_k is centered, $E[W_hX_k] = E[W_{h'}X_k] = 0$. The correlation
coefficient between interaction terms becomes

491
$$\text{Cor}[W_hX_k, W_{h'}X_k] = \frac{E[W_hX_kW_{h'}X_k]}{\sqrt{E[W_h{}^2X_k{}^2]}\sqrt{E[W_{h'}{}^2X_k{}^2]}}$$

492
$$= \frac{E[W_hW_{h'}]}{\sqrt{E[W_h{}^2]}\sqrt{E[W_{h'}{}^2]}}$$

493
$$= \frac{\text{Cov}[W_h, W_{h'}] + E[W_h]\,E[W_{h'}]}{\sqrt{\text{Var}[W_h] + E[W_h]^2}\sqrt{\text{Var}[W_{h'}] + E[W_{h'}]^2}}$$

494
$$= \frac{\frac{\text{Cor}[W_h, W_{h'}]\sqrt{\text{Var}[W_h]}\sqrt{\text{Var}[W_{h'}]}}{E[W_h]E[W_{h'}]} + 1}{\sqrt{\frac{\text{Var}[W_h]}{E[W_h]^2} + 1}\sqrt{\frac{\text{Var}[W_{h'}]}{E[W_{h'}]^2} + 1}}$$

495
$$= \frac{\text{Cor}[W_h, W_{h'}]\,\text{CV}[W_h]\,\text{CV}[W_{h'}] + 1}{\sqrt{\text{CV}[W_h]^2 + 1}\sqrt{\text{CV}[W_{h'}]^2 + 1}}. \quad (17)$$

496 If $\text{CV}[W_h]$ and $\text{CV}[W_{h'}]$ approach zero, the correlation of interaction terms
497 approaches one, irrespective of $\text{Cor}[W_h, W_{h'}]$.

EWAS of rheumatoid arthritis

EWAS datasets for rheumatoid arthritis were downloaded from the Gene Expression Omnibus. Using the RnBeads package (version 2.2.0) [31] of R, IDAT files of HumanMethylation450 array were preprocessed by removing low quality samples and markers, by normalizing methylation level, and by removing markers on sex chromosomes and outlier samples. The association of methylation level with disease status was tested with adjustment for sex, age, smoking status and experiment batch; the covariates were assumed to have uniform effects across cell types. After quality control, dataset GSE42861 included bulk peripheral blood leukocyte data for 336 cases and 322 controls [21].

The cell type composition of bulk samples was imputed using the Houseman algorithm [32] in the GLINT software (version 1.0.4) [33]. The reference data of GLINT software characterizes seven cell types [34] by 300 CpG sites [35], of which 284 were measured in our data. We used prediction results for the seven cell types (Table 1).

Dataset GSE131989 included sorted CD14$^+$ monocyte data for 63 cases and 31 controls [22]. By meta-analysis of GSE131989 and GSE87095 [20], we obtained sorted CD19$^+$ B cell data for 108 cases and 95 controls. Under the nominal significance level $P < 0.05$ (two-sided), the number of CpG sites up- or down-regulated in cases were 20,545 (5%) and 14,742 (3%), respectively, in CD14$^+$ monocyte and 28,216 (6%) and 26,447 (6%) in CD19$^+$ B cell.

Differential gene expression by age

Whole blood RNA-seq data of GTEx v7 was downloaded from the GTEx website [23]. Genes of low quality or on sex chromosomes were removed, expression level was normalized, outlier samples were removed, and 389 samples were retained. The association of read count with age was tested with adjustment for sex.

The cell type composition of bulk samples was imputed using the DeconCell package (version 0.1.0) [10] of R. The reference data of DeconCell

characterizes 33 cell types by two to 217 signature genes. Our data measured

39% of the signature genes. We used prediction results for six main cell types

(Table 2), for which the prediction performance was 44.4 to 90.9.

From the Gene Expression Omnibus dataset GSE56047 [24], we obtained

sorted CD14$^+$ monocyte data for 1202 samples and sorted CD4$^+$ T cell data

for 214 samples. Under the nominal significance level P < 0.05 (two-sided),

the number of genes up- or down-regulated by higher age were 2081 (16%)

and 1984 (15%), respectively, in CD14$^+$ monocyte and 693 (5%) and 842

(6%) in CD4$^+$ T cell.

Simulation of cell-type-specific disease association

Bulk tissue sample data for case-control comparison were simulated based

on the above-mentioned EWAS dataset GSE42861 and GTEx gene expression

dataset. We randomly assigned the case-control status to the samples.

Among the omics markers, 2.5% were set to be up-regulated in cases in

single cell type, 2.5% were similarly down-regulated, and 95% were

unrelated to case-control status. The cell-type-specific effect-size of the

differentially expressed markers was fixed within a simulation trial, and was

chosen from methylation odds ratio (OR) of 1.3, 1.6 or 1.9 for EWAS [19]

and fold-change of 1.7, 3.0 or 5.0 for gene expression analysis; the effect-

sizes correspond to log(1.3), log(1.6) and log(1.9) or log(1.7), log(3.0) and

log(5.0) in normalized scale.

For each effect-size, we performed 20 simulation trials for the EWAS

dataset and 50 trials for the gene expression dataset. In each simulation trial,

we randomly assigned half of the samples as cases ($X_{i,k} = 0.5$) and the other

half as controls ($X_{i,k} = -0.5$). We retained the covariates matrix $C_{i,l}$ and the

cell type composition matrix $W_{h,i}$ from the original data. From the original

bulk expression level matrix $Y_{i,j}$, 95% of the markers were randomly chosen

and retained; these markers had no association with disease because the

case-control status was randomized. Cell-type-specific association was

introduced into the remaining 5% of markers, such that an equal number of

markers were up- or down-regulated in each cell type. For example, in the

EWAS dataset, $451{,}725 \times 0.05 \times 0.5 \div 7 = 1613$ CpG sites were up-regulated in

neutrophils of cases.

The bulk expression level of a marker j with normalized-scale effect-size

β specific to a cell type h was generated as follows. First, the average μ and

the variance σ^2 of the normalized bulk expression level $f(Y_{i,j})$ in the original

data was measured. Next, we generated normalized expression level in each

cell type. For cases, the expression level in cell type h was randomly sampled

from the normal distribution $N(\mu + \beta, \sigma^2)$ and the expression level in each of

the other cell types was sampled from $N(\mu, \sigma^2)$. For controls, the expression

level in each cell type was sampled from $N(\mu, \sigma^2)$. Finally, for each individual,

the expression levels in cell types were converted to the linear scale,

multiplied by the cell type composition and added, to obtain the bulk

expression level in linear scale.

For gene expression, we also simulated a scenario where cell-type-specific

disease effect occurs in cell type marker genes. The simulation procedure is

same as above except that the expression level was set to zero (in linear

scale) in all cell types other than the target cell type h, for both cases and

controls.

Evaluation of statistical methods

Cell-type-specific effects of traits was statistically tested by using bulk tissue

data as input. We applied the omicwas package with the normalizing function

f = log, logit, identity without ridge regularization (omicwas.log,

omicwas.logit,   omicwas.identity)   or   under   ridge   regression

(omicwas.log.ridge,   omicwas.logit.ridge,   omicwas.identity.ridge).   The

omicwas package was used also for conventional linear regression under the

full and marginal models. We also developed a hybrid of marginal and full

models (Marginal.Full005): if the effect direction agreed in two models and if

$P < 0.05$ in the full model, we adopted the Z-score of the marginal model;

otherwise, the Z-score was set to zero.

Among previous methods, we evaluated those that accept cell type

composition as input and compute test statistics for cell-type-specific

association. For DNA methylation data, we applied TOAST (version 1.2.0) [9],

CellDMC (version 2.0.2) [13] and TCA (version 1.0.0) [14]. For gene

expression data, we applied TOAST and csSAM (version 1.4) [5]. For csSAM,
we either fitted all cell types together or one cell type at a time, and denoted
the results as csSAM.lm and csSAM.monovariate, respectively. The csSAM
method is applicable to binomial traits but not to quantitative traits.

For simulated data of EWAS dataset GSE42861, we adopted the
significance level $P < 2.4 \times 10^{-7}$, which accounts for the correlation among
the probes on HumanMethylation450 array [36]. For the GTEx gene
expression dataset, multiple testing was controlled by the Benjamini-
Hochberg procedure with the false discovery rate <5% in each cell type [37].

For the association with rheumatoid arthritis and age, "true" association
was determined from the measurements in physically sorted blood cells,
under the nominal significance level $P < 0.05$ (two-sided). The obtained
benchmark set of differentially expressed markers were "up-regulated" (in
rheumatoid arthritis cases or elders) or "down-regulated." For a set of
differentially expressed markers in a cell type (e.g., up-regulated in
monocytes), the prediction performance of an algorithm was measured by
the AUC of ROC. Standard error of AUC was computed by the jackknife
estimator by splitting the markers into 100 groups by chromosomal position.
The relative performance of an algorithm was evaluated by its AUC − 0.5
divided by that for the best algorithm in each setting.
613

Supplementary information

Supplementary information
Additional file 1: Table S1. Blood cell type proportion in Tsimane
Amerindians, Caucasians and Hispanics.
Additional file 2: Fig. S1. Sensitivity for detecting cell-type-specific
association in simulated data for gene expression of marker genes.
Additional file 3: Fig. S2. Specificity for detecting cell-type-specific
association in simulated data for gene expression of marker genes.
Additional file 4: Fig. S3. Precision (positive predictive value) for detecting
cell-type-specific association in simulated data for gene expression of marker
genes.
Additional file 5: Supplementary note. Asymptotic distribution of ridge

estimator.

Abbreviations

AUC: area under the curve, CV: coefficient of variation, eQTL: expression QTL,

EWAS: epigenome-wide association study, mQTL: methylation QTL, MSE:

mean squared error, OLS: ordinary least squares, OR: odds ratio, QTL:

quantitative trait locus, ROC: receiver operating characteristic, SNP: single

nucleotide polymorphism

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated and analyzed during the current study are available

in the figshare repository, https://dx.doi.org/10.6084/m9.figshare.10718282

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by JSPS KAKENHI [grant number JP16K07218] and

by the NCGM Intramural Research Fund [grant numbers 19A2004, 20A1013].

The funding body had no role in the design and collection of the study,

experiments, analyses and interpretations of data, and in writing the

manuscript.

Author's contributions

FT developed the methodology, wrote the software, implemented the study,

and wrote the manuscript. NK revised the manuscript. All authors read and
approved the final manuscript.

Acknowledgements

Not applicable.

References

1. Teschendorff AE, Zheng SC. Cell-type deconvolution in epigenome-wide
association studies: a review and recommendations. Epigenomics.
2017;9:757–68.

2. Sturm G, Finotello F, Petitprez F, Zhang JD, Baumbach J, Fridman WH, et
al. Comprehensive evaluation of transcriptome-based cell-type
quantification methods for immuno-oncology. Bioinformatics.
2019;35:i436–45.

3. Ghosh D. Mixture models for assessing differential expression in complex
tissues using microarray data. Bioinformatics. 2004;20:1663–9.

4. Stuart RO, Wachsman W, Berry CC, Wang-Rodriguez J, Wasserman L,
Klacansky I, et al. In silico dissection of cell-type-associated patterns of
gene expression in prostate cancer. Proc Natl Acad Sci USA. National
Academy of Sciences; 2004;101:615–20.

5. Shen-Orr SS, Tibshirani R, Khatri P, Bodian DL, Staedtler F, Perry NM, et
al. Cell type–specific gene expression differences in complex tissues. Nat
Meth. Nature Publishing Group; 2010;7:287–9.

6. Erkkilä T, Lehmusvaara S, Ruusuvuori P, Visakorpi T, Shmulevich I,
Lähdesmäki H. Probabilistic analysis of gene expression measurements from
heterogeneous tissues. Bioinformatics. 2010;26:2571–7.

7. Kuhn A, Thu D, Waldvogel HJ, Faull RLM, Luthi-Carter R. Population-
specific expression analysis (PSEA) reveals molecular changes in diseased

brain. Nat Meth. Nature Publishing Group; 2011;8:945–7.

8. Westra H-J, Arends D, Esko T, Peters MJ, Schurmann C, Schramm K, et
al. Cell Specific eQTL Analysis without Sorting Cells. Pastinen T, editor. PLoS
Genet. Public Library of Science; 2015;11:e1005223–17.

9. Li Z, Wu Z, Jin P, Wu H. Dissecting differential signals in high-throughput
data from complex tissues. Hancock J, editor. Bioinformatics.
2019;35:3898–905.

10. Aguirre-Gamboa R, de Klein N, di Tommaso J, Claringbould A, van der
Wijst MG, de Vries D, et al. Deconvolution of bulk blood eQTL effects into
immune cell subpopulations. BMC Bioinformatics. 2020;21:243.

11. Montaño CM, Irizarry RA, Kaufmann WE, Talbot K, Gur RE, Feinberg AP,
et al. Measuring cell-type specific differential methylation in human brain
tissue. Genome Biol. BioMed Central; 2013;14:R94–9.

12. White N, Benton M, Kennedy D, Fox A, Griffiths L, Lea R, et al.
Accounting for cell lineage and sex effects in the identification of cell-
specific DNA methylation using a Bayesian model selection algorithm.
Sawalha AH, editor. PLoS ONE. 2017;12:e0182455–18.

13. Zheng SC, Breeze CE, Beck S, Teschendorff AE. Identification of
differentially methylated cell types in epigenome-wide association studies.
Nat Meth. Nature Publishing Group; 2018;15:1059–66.

14. Rahmani E, Schweiger R, Rhead B, Criswell LA, Barcellos LF, Eskin E, et
al. Cell-type-specific resolution epigenetics without the need for cell sorting
or single-cell biology. Nature Communications. Nature Publishing Group;
2019;10:3417–11.

15. Cobos FA, Vandesompele J, Mestdagh P. Computational deconvolution
of transcriptomics data from mixed cell populations. Bioinformatics.
2018;34:1969–79.

16. Hoyle DC, Rattray M, Jupp R, Brass A. Making sense of microarray data
distributions. Bioinformatics. 2002;18:576–84.

17. Du P, Zhang X, Huang C-C, Jafari N, Kibbe WA, Hou L, et al.
Comparison of Beta-value and M-value methods for quantifying methylation
levels by microarray analysis. BMC Bioinformatics. BioMed Central;
2010;11:1–9.

18. Aiken LS, West SG. Multiple Regression: Testing and Interpreting
Interactions. Sage Publications; 1991.

19. Rakyan VK, Down TA, Balding DJ, Beck S. Epigenome-wide association
studies for common human diseases. Nat Rev Genet. 2011;12:529–41.

20. Julià A, Absher D, López-Lasanta M, Palau N, Pluma A, Waite Jones L, et
al. Epigenome-wide association study of rheumatoid arthritis identifies
differentially methylated loci in B cells. Hum Mol Genet. 2017;26:2803–11.

21. Liu Y, Aryee MJ, Padyukov L, Fallin MD, Hesselberg E, Runarsson A, et
al. Epigenome-wide association data implicate DNA methylation as an
intermediary of genetic risk in rheumatoid arthritis. Nat Biotechnol. Nature
Publishing Group; 2013;31:142–7.

22. Rhead B, Holingue C, Cole M, Shao X, Quach HL, Quach D, et al.
Rheumatoid Arthritis Naive T Cells Share Hypermethylation Sites With
Synoviocytes. Arthritis & Rheumatology. John Wiley & Sons, Ltd;
2017;69:550–9.

23. GTEx Consortium. Genetic effects on gene expression across human
tissues. Nature. Nature Publishing Group; 2017;550:204–13.

24. Reynolds LM, Taylor JR, Ding J, Lohman K, Johnson C, Siscovick D, et
al. Age-related variations in the methylome associated with gene expression
in human monocytes and T cells. Nature Communications. 2014;5:5366.

25. Riazoshams H, Midi H, Ghilagaber G. Robust Nonlinear Regression: with

Applications using R. John Wiley & Sons; 2019.

26. Lim C. Robust ridge regression estimators for nonlinear models with
applications to high throughput screening assay data. Statist. Med.
2014;34:1185–98.

27. Lawless JF, Wang P. A simulation study of ridge and other regression
estimators. Communications in Statistics - Theory and Methods.
1976;5:307–23.

28. Hoerl AE, Kannard RW, Baldwin KF. Ridge regression: some simulations.
Communications in Statistics - Theory and Methods. 1975;4:105–23.

29. Halawa AM, Bassiouni El MY. Tests of regression coefficients under ridge
regression models. Journal of Statistical Computation and Simulation.
2000;65:341–56.

30. Dennis JE, Gay DM, Welsch RE. An adaptive nonlinear least-squares
algorithm. ACM Transactions on Mathematical Software. 1981;7:348–68.

31. Müller F, Scherer M, Assenov Y, Lutsik P, Walter J, Lengauer T, et al.
RnBeads 2.0: comprehensive analysis of DNA methylation data. Genome
Biol. BioMed Central; 2019;20:55–12.

32. Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ,
Nelson HH, et al. DNA methylation arrays as surrogate measures of cell
mixture distribution. BMC Bioinformatics. 2012;13:86.

33. Rahmani E, Yedidim R, Shenhav L, Schweiger R, Weissbrod O, Zaitlen
N, et al. GLINT: a user-friendly toolset for the analysis of high-throughput
DNA-methylation array data. Hancock JM, editor. Bioinformatics.
2017;33:1870–2.

34. Reinius LE, Acevedo N, Joerink M, Pershagen G, Dahlén S-E, Greco D, et
al. Differential DNA Methylation in Purified Human Blood Cells: Implications
for Cell Lineage and Studies on Disease Susceptibility. Ting AH, editor. PLoS

ONE. Public Library of Science; 2012;7:e41361–13.

35. Koestler D. Improving Cell Mixture Deconvolution by Identifying Optimal
DNA methylation Libraries (IDOL). BMC Bioinformatics. BMC Bioinformatics;
2016;:1–21.

36. Saffari A, Silver MJ, Zavattari P, Moi L, Columbano A, Meaburn EL, et al.
Estimation of a significance threshold for epigenome-wide association
studies. Genet Epidemiol. John Wiley & Sons, Ltd; 2017;42:20–33.

37. Anders S, McCarthy DJ, Chen Y, Okoniewski M, Smyth GK, Huber W, et
al. Count-based differential expression analysis of RNA sequencing data using
R and Bioconductor. Nat Protoc. 2013;8:1765–86.
768

 TABLES

Table 1A Blood cell type proportion in rheumatoid arthritis dataset

Cell type	Neu	CD4$^+$T	CD8$^+$T	NK	Mono	Bcells	Eos
Mean	0.59	0.10	0.08	0.08	0.07	0.07	0.01
SD	0.11	0.06	0.05	0.04	0.02	0.03	0.02
CV	0.2	0.6	0.6	0.5	0.3	0.4	2.7

SD, standard deviation; CV, coefficient of variation

Table 1B Correlation between blood cell type proportion and rheumatoid arthritis (X_k)

r	Neu	CD4$^+$T	CD8$^+$T	NK	Mono	Bcells	Eos	X_k=Disease
Neu	1	-0.68	-0.60	-0.46	-0.06	-0.49	-0.48	0.44
CD4$^+$T	-0.68	1	0.14	0.05	-0.17	0.38	0.26	-0.33
CD8$^+$T	-0.60	0.14	1	0.08	-0.05	0.19	0.13	-0.27
NK	-0.46	0.05	0.08	1	-0.04	0.01	0.11	-0.27
Mono	-0.06	-0.17	-0.05	-0.04	1	-0.17	0.05	0.10
Bcells	-0.49	0.38	0.19	0.01	-0.17	1	0.11	-0.22
Eos	-0.48	0.26	0.13	0.11	0.05	0.11	1	-0.10

Table 1C Correlation between interaction terms

r	Neu*X_k	CD4$^+$T*X_k	CD8$^+$T*X_k	NK*X_k	Mono*X_k	Bcells*X_k	Eos*X_k
Neu*X_k	1	0.83	0.80	0.85	0.93	0.90	0.27
CD4$^+$T*X_k	0.83	1	0.78	0.78	0.83	0.88	0.42
CD8$^+$T*X_k	0.80	0.78	1	0.77	0.82	0.83	0.35
NK*X_k	0.85	0.78	0.77	1	0.85	0.83	0.35
Mono*X_k	0.93	0.83	0.82	0.85	1	0.88	0.35
Bcells*X_k	0.90	0.88	0.83	0.83	0.88	1	0.36
Eos*X_k	0.27	0.42	0.35	0.35	0.35	0.36	1

Neu, neutrophils; Mono, monocytes; Eos, eosinophils.

Table 2A Blood cell type proportion in GTEx dataset

Cell type	Gran	CD4$^+$T	CD8$^+$T	Mono	NK	Bcells
Mean	0.53	0.22	0.10	0.07	0.05	0.03
SD	0.037	0.020	0.013	0.004	0.012	0.003
CV	0.1	0.1	0.1	0.1	0.2	0.1

SD, standard deviation; CV, coefficient of variation

Table 2B Correlation between blood cell type proportion and age (X_k)

r	Gran	CD4$^+$T	CD8$^+$T	Mono	NK	Bcells	X_k=Age
Gran	1	-0.89	-0.83	0.56	-0.76	-0.41	-0.23
CD4$^+$T	-0.89	1	0.59	-0.64	0.50	0.51	0.14
CD8$^+$T	-0.83	0.59	1	-0.40	0.59	0.15	0.15
Mono	0.56	-0.64	-0.40	1	-0.44	-0.42	0.02
NK	-0.76	0.50	0.59	-0.44	1	0.13	0.31
Bcells	-0.41	0.51	0.15	-0.42	0.13	1	-0.03

Table 2C Correlation between interaction terms

r	Gran*X_k	CD4$^+$T*X_k	CD8$^+$T*X_k	Mono*X_k	NK*X_k	Bcells*X_k
Gran*X_k	1	0.99	0.98	1.00	0.96	0.99
CD4$^+$T*X_k	0.99	1	1.00	0.99	0.98	1.00
CD8$^+$T*X_k	0.98	1.00	1	0.99	0.98	0.99
Mono*X_k	1.00	0.99	0.99	1	0.96	0.99
NK*X_k	0.96	0.98	0.98	0.96	1	0.97
Bcells*X_k	0.99	1.00	0.99	0.99	0.97	1

Gra, granulocytes; Mono, monocytes.

 FIGURE LEGENDS

Figure 1

Contour plot of the correlation coefficient between interaction terms $W_h X_k$

and $W_{h'} X_k$. W_h and $W_{h'}$ represent proportions of cell types h and h', and X_k

represents the value of trait k. For this plot, we assume the coefficient of

variation $\text{CV}[W_h]$ and $\text{CV}[W_{h'}]$ to be equal. As the CV decreases 0.6, 0.4 to

0.2, the correlation coefficient raises >0.5, >0.7 to >0.9, over most range of

$\text{Cor}[W_h, W_{h'}]$.

Figure 2

Sensitivity for detecting cell-type-specific association in simulated data for

DNA methylation. Panels are aligned in rows according to the simulation

settings with the methylation odds ratio of 1.3, 1.6 or 1.9. In each row, panels

for different cell types are aligned in decreasing order of proportion. The

vertical axis indicates sensitivity. In each panel, results from different

algorithms are aligned horizontally in different colors. Results from 20

simulation trials are summarized in a box plot. The middle bar of the box plot

indicates the median, and the lower and upper hinges correspond to the first

and third quartiles. The whiskers extend to the value no further than 1.5 $\times$

inter-quartile range from the hinges. MethOR, methylation odds ratio; Neu,

neutrophils; Mono, monocytes; Eos, eosinophils.

Figure 3

Specificity for detecting cell-type-specific association in simulated data for

DNA methylation. The figure format is same as Fig. 3.

Figure 4

Precision (positive predictive value) for detecting cell-type-specific

association in simulated data for DNA methylation. The figure format is same

as Fig. 3.

Figure 5

Sensitivity for detecting cell-type-specific association in simulated data for
gene expression. Panels are aligned in rows according to the simulation
settings with the gene expression fold change of 1.7, 3.0 or 5.0. In each row,
panels for different cell types are aligned in decreasing order of proportion.
The vertical axis indicates sensitivity. In each panel, results from different
algorithms are aligned horizontally in different colors. Results from 50
simulation trials are summarized in a box plot. The middle bar of the box plot
indicates the median, and the lower and upper hinges correspond to the first
and third quartiles. The whiskers extend to the value no further than $1.5 \times$
inter-quartile range from the hinges. FC, fold change; Gran, granulocytes;
Mono, monocytes.

Figure 6

Specificity for detecting cell-type-specific association in simulated data for
gene expression. The figure format is same as Fig. 5.

Figure 7

Precision (positive predictive value) for detecting cell-type-specific
association in simulated data for gene expression. The figure format is same
as Fig. 5.

Figure 8

Performance of cell-type-specific association prediction. For rheumatoid
arthritis association of DNA methylation in monocytes (A) and B cells (B);
Age association of gene expression in CD4$^+$ T cells (C) and monocytes (D).
The prediction is evaluated separately for up-regulated and down-regulated
markers. The AUC of ROC and its 95% confidence interval are plotted for each
statistical algorithm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

(A) Rheumatoid arthritis association in Monocytes

(B) Rheumatoid arthritis association in B cells

(C) Age association in CD4+ T cells

(D) Age association in Monocytes