Nonlinear ridge regression improves cell-type-specific differential expression analysis

Fumihiko Takeuchi and Norihiro Kato

Department of Gene Diagnostics and Therapeutics, Research Institute, National Center for Global Health and Medicine (NCGM), Tokyo, Japan

Correspondence

Fumihiko Takeuchi

Department of Gene Diagnostics and Therapeutics, Research Institute, National Center for Global Health and Medicine (NCGM)

1-21-1 Toyama, Shinjuku-ku, Tokyo, 162-8655, Japan

Email: fumihiko@takeuchi.name

Abstract

Background: Epigenome-wide association studies (EWAS) and differential

gene expression analyses are generally performed on tissue samples, which

consist of multiple cell types. Cell-type-specific effects of a trait, such as

disease, on the omics expression are of interest but difficult or costly to

measure experimentally. By measuring omics data for the bulk tissue, cell

type composition of a sample can be inferred statistically. Subsequently, cell-

type-specific effects are estimated by linear regression that includes terms

representing the interaction between the cell type proportions and the trait.

This approach involves two issues, scaling and multicollinearity.

Results: First, although cell composition is analyzed in linear scale,

differential methylation/expression is analyzed suitably in the logit/log scale.

To simultaneously analyze two scales, we applied nonlinear regression.

Second, we show that the interaction terms are highly collinear, which is

obstructive to ordinary regression. To cope with the multicollinearity, we

applied ridge regularization. In simulated data, nonlinear ridge regression

attained well-balanced sensitivity, specificity and precision. Marginal model

attained the lowest precision and highest sensitivity and was the only

algorithm to detect weak signal in real data.

Conclusion: Nonlinear ridge regression performed cell-type-specific

association test on bulk omics data with well-balanced performance. The

omicwas package for R implements nonlinear ridge regression for cell-type-

specific EWAS, differential gene expression and QTL analyses. The software

is freely available from https://github.com/fumi-github/omicwas

Keywords

Epigenome-wide association study, Differential gene expression analysis, Cell

type, Nonlinear regression, Ridge regression, mQTL, eQTL

Background

Epigenome-wide association studies (EWAS) and differential gene expression analyses elucidate the association of disease traits (or conditions) with the level of omics expression, namely DNA methylation and gene expression. Thus far, tissue samples, which consist of heterogeneous cell types, have mainly been examined, because cell sorting is not feasible in most tissues and single-cell assay is still expensive. Nevertheless, the cell type composition of a sample can be quantified statistically by comparing omics measurement of the target sample with reference data obtained from sorted or single cells [1,2]. By utilizing the composition, the disease association specific to a cell type was statistically inferred for gene expression [3-10] and DNA methylation [11-14].

For the imputation of cell type composition, omics markers are usually analyzed in the original linear scale, which measures the proportion of mRNA molecules from a specific gene or the proportion of methylated cytosine molecules among all cytosines at a specific CpG site [15]. The proportion can differ between cell types, and the weighted average of cell-type-specific proportions becomes the proportion in a bulk tissue sample. Using the fact that the weight equals the cell type composition, the cell type composition of a sample is imputed. In contrast, gene expression analyses are performed in the log-transformed scale because the signal and noise are normally distributed after log-transformation [16]. In DNA methylation analysis, the logit-transformed scale, which is called the M-value, is statistically valid [17], although the linear scale could yield comparable performance under large sample size [18]. Consequently, the optimal scales for analyzing differential gene expression or methylation can differ from the optimal scale for analyzing cell type composition.

Aiming to perform cell-type-specific EWAS or differential gene expression analyses by using unsorted tissue samples, we study two issues that have been overlooked. Whereas previous studies were performed in linear scale, we develop a nonlinear regression, which simultaneously analyzes cell type composition in linear scale and differential expression/methylation in log/logit scale. The second issue is multicollinearity. Cell-type-specific effects of a trait,

such as disease, on omics expression are usually estimated by linear

regression that includes terms representing the interaction between the cell

type proportions and the trait. We show that the interaction terms can

mutually be highly correlated, which obstructs ordinary regression. To cope

with the multicollinearity, we implement ridge regularization. Our methods

and previous ones are compared in simulated and real data.

Results

Multicollinearity of interaction terms

Typically, cell-type-specific effects of a trait on omics marker expression is

analyzed by the linear regression in equation (2). For each omics marker, the

goal is to estimate $\beta_{h,k}$, the effect of trait *k* on the expression level in cell type

h. This is estimated based on the relation between the bulk expression level

Y_i of sample *i* and the regressor $W_{h,i}X_{i,k}$, which is an interaction term defined

as the product of the cell type proportion $W_{h,i}$ and the trait value $X_{i,k}$ of the

sample. We assume that Y_i, $W_{h,i}$ and $X_{i,k}$ are given as input data.

The variable $W_{h,i}$ for cell type composition cannot be mean-centered for

our purpose. If $W_{h,i}$ were centered, we would obtain, instead of $\beta_{h,k}$, the

deviation of $\beta_{h,k}$ from the average across cell types. In general, interaction

terms involving uncentered variables can become collinear [19]. We first

survey the extent of multicollinearity in real data for cell-type-specific

association.

In peripheral blood leukocyte data from a rheumatoid arthritis study

(GSE42861), the proportion of cell types ranged from 0.59 for neutrophils to

0.01 for eosinophils (Table 1A). The proportion of neutrophils was negatively

correlated with the proportion of other cell types (apart from monocytes) with

correlation coefficient of −0.68 to −0.46, whereas the correlation was weaker

for other pairs (Table 1B). Rheumatoid arthritis status was modestly

correlated with proportions of cell types. The product of the disease status

X_k, centered to have zero mean, and the proportion of a cell type becomes

an interaction term. The correlation coefficients between the interaction

terms were mostly >0.8, apart from eosinophils (Table 1C). The coefficient

of variation (CV), which is the ratio of standard deviation to mean, of the

proportion was low for all cell types apart from eosinophils (Table 1A). The

interaction terms for low-CV cell types were strongly correlated with X_k,

which in turn caused strong correlation between the relevant interaction

terms.

The situation was the same for the interaction with age in GTEx data. The

granulocytes (which include neutrophils and eosinophils) were the most

abundant (Table 2A). The proportion of granulocytes was negatively

correlated with other cell types (apart from monocytes) with correlation

coefficient of −0.89 to −0.41, and the correlation between other pairs was

generally weaker (Table 2B). Age was modestly correlated with proportions

of cell types. In this dataset, the CV of the proportion was low in all cell types

(Table 2A), which caused strong mutual correlation between interaction

terms (Table 2C).

In the above empirical data, multicollinearity between interaction terms

seemed to arise not due to the correlation between cell type proportions or

X_k, but due to the low CV in the cell type proportions. Subsequently, this

property was derived mathematically. As we derived in equation (19), the

correlation between interaction terms $W_h X_k$ and $W_{h'} X_k$ approaches one when

$\mathrm{CV}[W_h]$ and $\mathrm{CV}[W_{h'}]$ are low, irrespective of $\mathrm{Cor}[W_h, W_{h'}]$ (Fig. 1). The CV was

0.2 to 0.6 (apart from eosinophils) in the rheumatoid arthritis dataset and

0.1 to 0.2 in the GTEx dataset. We looked up datasets of several ethnicities

and found the CV to be ≤0.6 in majority of blood cell types (Additional file 1:

Table S1). Thus, multicollinearity can be a common problem for cell-type-

specific association analyses. Biologically, in tissues where cell type

composition is tightly controlled, the CV of cell type proportion becomes low

and the multicollinearity is exacerbated.

Evaluation in simulated data

By using simulated data, we evaluated previous methods and new

approaches of the omicwas package. In order to simultaneously analyze two

scales, the linear scale for heterogeneous cell mixing and the log/logit scale

for trait effects, we applied nonlinear regression (equations (4) and (5)). To

cope with the multicollinearity of interaction terms, we applied ridge

regularization (formula (10)).

Previous regression type methods are based either on the full model of

linear regression (equation (2)) or the marginal model (equation (3)). The

full model fits and tests cell-type-specific effects for all cell types

simultaneously, and its variations include TOAST, csSAM.lm and CellDMC. The

marginal model fits and tests cell-type-specific effect for one cell type at a

time, and its variations include csSAM.monovariate and TCA. We also

examined a hybrid of the two models (Marginal.Full005), which becomes

positive when the models agree.

The simulation data was generated from real datasets of DNA methylation

(658 samples; 451,725 CpG sites) and gene expression (389 samples;

14,038 genes). The original cell type composition was retained for all samples,

and the case-control status was randomly assigned. Ninety-five percent of

omics markers were set to be unassociated with disease status, 2.5% were

up-regulated in cases at one cell type, and 2.5% were similarly down-

regulated. The cell-type-specific effect-size was fixed in a simulation trial,

either to methylation odds ratio (OR) of 1.3, 1.6 or 1.9 or to gene expression

fold change of 1.7, 3.0 or 5.0. The significance level was set to $P < 2.4 \times 10^{-7}$

for DNA methylation and false discovery rate <5% for gene expression. In

each simulation trial, the sensitivity, specificity and precision for detecting

cell-type-specific association was calculated for each cell type. To compare

algorithms, the performance measures for the same effect-size and cell type

were averaged over the simulation trials.

Overall, in the simulation for DNA methylation the sensitivity (Fig. 2) was

higher under large effect-size (bottom row of panels) and in abundant cell

types (left columns of panels). The average specificity (Fig. 3) was high

across the effect-size settings and across cell types, being >0.97 for the

Marginal model, >0.98 for TCA and >0.999 for other algorithms. The

precision (Fig. 4) was higher in abundant cell types within each effect-size

setting. As effect-size increased, the precision decreased in neutrophils but

increased in the other minor cell types. Excluding the cases where all

algorithms lacked sensitivity (monocytes and B cells under methOR=1.3 and
eosinophils), the average precision of omicwas.logit.ridge was >0.79 and was
the highest in 13/16 of the cases.

There was trade-off between sensitivity and precision. Among the
algorithms, the Marginal model attained the highest sensitivity and the lowest
precision. TCA, which is a variation of the marginal model, had relatively high
sensitivity and relatively low precision. Marginal.Full005 attained the second
highest sensitivity and moderate precision. The ridge regressions
(omicwas.logit.ridge and omicwas.identity.ridge) attained moderate
sensitivity and high precision. The full models without ridge regularization
(omicwas.logit, omicwas.identity, Full, TOAST and CellDMC) had the lowest
sensitivity.

The overall tendency was similar in the simulation for gene expression.
The sensitivity (Fig. 5) was higher under large effect-size and in abundant
cell types. The average specificity (Fig. 6) was high across the effect-size
settings and across cell types, being >0.96 for the marginal models (Marginal
and csSAM.monovariate), >0.98 for the nonlinear and ridge regressions
(omicwas.log.ridge, omicwas.identity.ridge, omicwas.log) and >0.996 for
other algorithms. The precision (Fig. 7) was higher in abundant cell types
within each effect-size setting. As effect-size increased, the precision
decreased in granulocytes but increased in the other minor cell types.
Excluding the full models (omicwas.identity, Full, TOAST, csSAM.lm) that
lacked sensitivity, the algorithms that were frequently top in average
precision were omicwas.identity.ridge (5 cases), omicwas.log (5 cases),
omicwas.log.ridge (3 cases) and Marginal.Full005 (3 cases).

There again was trade-off between sensitivity and precision. Among the
algorithms, the marginal models attained the highest sensitivity but relatively
low precision. The nonlinear and ridge regressions and Marginal.Full005
attained moderate sensitivity and highest precision. The full models had very
low average sensitivity of <0.01.

For gene expression, we also simulated a scenario where cell-type-specific
disease effect occurred in cell type "marker" genes (Additional files 2, 3, 4:
Figs. S1, S2, S3). In other words, the expression level in the target cell type

differed between cases and controls, and the expression level in other cell
types was zero (in linear scale). Thus, non-target cell types did not introduce
noise to bulk expression level. The Marginal model attained the highest
average sensitivity of >0.93 but relatively low average precision of ~0.16. As
equal number of differentially expressed genes were generated in the six cell
types, picking up signals for all such genes, including those not for the tested
cell type, would result in precision of 1/6 = 0.16. The full models had low
sensitivity. The nonlinear and ridge regressions and Marginal.Full005 attained
moderate sensitivity and moderate precision. With regards to the frequency
of being the top in average precision, the algorithms were ordered
Marginal.Full005   (9   cases),   omicwas.log.ridge   (6   cases),
omicwas.identity.ridge (2 cases) and omicwas.log (1 case), excluding the full
models that lacked sensitivity.

205 Cell-type-specific association with rheumatoid arthritis and age

The detection of cell-type-specific association in bulk tissue was evaluated by
using physically sorted cells. In principle, sorted cells should serve as genuine
verification, however, due to the relatively small sample size (94 or 203 for
rheumatoid arthritis and 214 or 1202 for age) the available datasets were
underpowered to generate a gold standard list of differentially expressed
omics markers [20]. Instead, we generated a benchmark set of differentially
expressed markers by imposing a relaxed significance level of $P < 0.05$; the
set would be enriched for true differentially expressed markers yet also
include unassociated markers. The benchmark set was cross-checked with
the prediction by each algorithm; in the same manner as the simulation
analysis, we assessed the sensitivity, specificity and precision.

The cell-type-specific association of DNA methylation with rheumatoid
arthritis was predicted using bulk peripheral blood leukocyte data and was
evaluated in sorted monocytes and B cells (Fig. 8). The input bulk methylation
data   was   normalized   by   applying   the   logit-transformation   for   the
Marginal.logit algorithm, which otherwise was the same as Marginal. Although
the sensitivity was extremely low for all algorithms, it was positive in both
cell types for Marginal ($0.8-1.1 \times 10^{-4}$), Marginal.logit ($1.2-1.4 \times 10^{-4}$), TCA

(0.5–1.2 $\times$ 10^{-4}) and omicwas.logit (0.5 $\times$ 10^{-4}). The cell-type-specific
association of DNA methylation with age was predicted using the same bulk
dataset and was evaluated in sorted CD4+T cells and monocytes (Fig. 8). The
Marginal and Marginal.logit models attained by far the highest sensitivity
(both 0.15–0.27) in both cell types, and moderate precision (0.59–0.68 and
0.60–0.68 respectively).

The cell-type-specific association of gene expression with age was
predicted using whole blood data and was evaluated in sorted CD4+ T cells
and monocytes (Fig. 9). The input bulk gene expression data was normalized
by applying the log-transformation for the Marginal.log algorithm, which
otherwise was the same as Marginal. Although the sensitivity was low for all
algorithms, it was positive in both cell types for Marginal (0.02–0.07),
Marginal.log (0.07–0.11), omicwas.identity.ridge (0.01–0.22) and
omicwas.log (0.03–0.05). The precision was modest for Marginal (0.06–0.31),
Marginal.log (0.07–0.28), omicwas.identity.ridge (0.03–0.21) and
omicwas.log (0.04–0.17). The dataset of sorted CD4+ T cells (214 samples)
is smaller than the monocyte dataset (1202 samples) thus could be
underpowered to pick enough true differentially expressed genes into the
benchmark set.

For DNA methylation dataset GSE42861 and for GTEx gene expression
dataset, the omicwas.logit.ridge and omicwas.log.ridge models of the
omicwas package was computed in 8.1 and 0.7 hours respectively, using 8
cores of a 2.5 GHz Xeon CPU Linux server.

Discussion
Aiming to elucidate cell-type-specific trait association in DNA methylation and
gene expression, this article explored two aspects, multicollinearity and scale.
We observed multicollinearity in real data and derived mathematically how it
emerges. To cope with the multicollinearity, we applied ridge regularization.
To properly handle multiple scales simultaneously, we applied nonlinear
regression. Among the examined algorithms, nonlinear ridge regression
attained moderate sensitivity and highest precision in simulated data. We also

developed an algorithm that combines full and marginal models, which
attained balanced sensitivity and precision in simulation. In real benchmark
data, all algorithms performed poorly yet the marginal models tended to
attain the highest sensitivity.

The statistical methods discussed in this article are applicable, in principle,
to any tissue. For validation of the methods, we need datasets for bulk tissue
as well as sorted cells, ideally of several hundred samples. Currently, the
publicly available data is limited to peripheral blood. By no means, the
rheumatoid arthritis EWAS datasets [21-23] or the datasets for age
association of gene expression [24,25] are representative. Nevertheless, we
think verification in real data is valuable.

By the performance in simulated and real data, we can roughly divide
algorithms into three groups: full (and its variations), marginal (and its
variations) and the third group that includes ridge regressions and the hybrid
Marginal.Full005. In marginal models, we test one cell type at a time. If we
knew in advance that one particular cell type is associated with the trait,
which would be a rare situation, testing that cell type with the marginal model
is the most simple and correct approach. However, when the test target cell
type is not associated, but instead another cell type is associated, the
marginal models can pick up false signals due to the collinearity between
regressor variables. Indeed, marginal models attained highest sensitivity
(Figs. 2, 5) and relatively low precision (Figs. 4, 7), which could lead to
unstable performance. The full model fits and tests all cell types
simultaneously, by which it adjusts for the effects of other cell types. Due to
the simultaneous inclusion of collinear predictors, the sensitivity was low (Figs.
2, 5). The ridge regressions (omicwas.identity.ridge, omicwas.logit.ridge and
omicwas.log.ridge) were in the middle between full and marginal models with
regards to the sensitivity (Figs. 2, 5), while attaining the highest specificity
(Figs. 4, 7). The hybrid Marginal.Full005 algorithm is intended to gain
sensitivity by the marginal model while keeping precision by incorporating
the full model. It attained moderate sensitivity (Figs. 2, 5) and moderate
precision (Figs. 4, 7) in simulation. In real data, all algorithms performed
poorly yet Marginal, Marginal.logit and Marginal.log tended to attain the

highest sensitivity. With regards to the performance measures of all
algorithms, the association of DNA methylation with age (Fig. 8) was roughly
similar to the simulation setting of methylation OR = 1.6 for B cells (Figs. 2–
4), and the association of gene expression with age (Fig. 9) was roughly
similar to the simulation setting of fold change = 1.7 for CD8$^+$T cells (Figs.
5–7). For the respective simulation settings, the median coefficient of
determination for the Marginal model was 0.020 and 0.007, indicating weak
association.

A limitation of our simulation is that only one cell type was assumed to be
associated with disease status at each marker. In reality, two or more cell
types can be associated with disease under homogeneous or heterogeneous
effect. In the physically sorted cells, the association of DNA methylation with
rheumatoid arthritis tended to be consistent between monocytes and B cells;
the association statistics across CpG sites were positively correlated with
Spearman's rank correlation coefficient of 0.20 (P-value < 2.2 × 10^{-16}).
Similarly, the association with age tended to be consistent between CD4$^+$T
cells and monocytes with correlation coefficient of 0.27 and 0.07 (P-value <
2.2 × 10^{-16}), respectively, for DNA methylation and gene expression. The
consistency suggests that multiple cell types tend to be associated under
homogeneous effect. If the association is completely consistent, the effect-
size is uniform across cell types. As there is no cell-type-specific effect, a
simple regression by disease (or relevant trait), ignoring the cell type
composition, becomes the appropriate modeling (formula (8)). Moreover,
when cell type composition has low CV (as observed in Tables 1 and 2), the
marginal model with normalized input (formula (9)) becomes almost identical
to the simple regression. In other words, the marginal model can pick up
signal in cases where effect-size is homogeneous across cell types.
Correspondingly, in real data of DNA methylation Marginal.logit performed
the best and was slightly better than Marginal (Fig. 8), and in gene expression
Marginal.log performed mostly the best and was better than Marginal (Fig.
9).

Conclusions

For cell-type-specific differential expression analysis by using unsorted tissue samples, we recommend trying the nonlinear ridge regression as a first choice because it balances sensitivity and precision. Although marginal models can be powerful when the tested cell type actually is the only one associated with the trait, caution is needed in its low precision. Under the idea of first scanning by the marginal model and then reanalyzing in full model, we developed the hybrid Marginal.Full005 algorithm, which attained balanced sensitivity and precision but was not corroborated in experimental data. Ridge regression is preferable compared to the full model without ridge regularization because ridge estimator of the effect-size has smaller mean squared error (equation (15)). The number of cell types associated with disease at each marker was restricted to one in our simulation but could be two or more with homogeneous or heterogeneous effect. If the effect-size is uniform across all cell types, a simple regression by disease status is suitable, which can be substituted with the marginal model that takes normalized input. We do not claim the ridge regression to substitute previous algorithms. Indeed, we think none of the current algorithms is superior to others in all aspects, indicating possibility for future improvement.

Methods

Linear regression

We begin by describing the linear regressions used in previous studies. Let the indexes be h for a cell type, i for a sample, j for an omics marker (CpG site or gene), k for a trait that has cell-type-specific effects on marker expression, and l for a trait that has a uniform effect across cell types. The input data is given in four matrices. The matrix $W_{h,i}$ represents cell type composition. The matrices $X_{i,k}$ and $C_{i,l}$ represent the values of the traits that have cell-type-specific and uniform effects, respectively. We assume the two matrices are centered: $\sum_i X_{i,k} = \sum_i C_{i,l} = 0$. For example, $X_{i,k} = 0.5$ for disease cases and $X_{i,k} = -0.5$ for controls when the number of cases and controls are

equal. The matrix $Y_{i,j}$ represents the omics marker expression level in tissue
samples.

The parameters we estimate are the cell-type-specific trait effect $\beta_{h,j,k}$,
tissue-uniform trait effect $\gamma_{j,l}$, and basal marker level $\alpha_{h,j}$ in each cell type.
For the remaining of the first five sections (up to "Multicollinearity of
interaction terms"), we focus on one marker j, and omit the index for
readability. For cell type h, the marker level of sample i is

359
$$\alpha_h + \sum_k \beta_{h,k} X_{i,k}. \quad (1)$$

360 This is a representative value rather than a mean because we do not model
361 a probability distribution for cell-type-specific expression. By averaging the
362 value over cell types with weight $W_{h,i}$, and combining with the tissue-uniform
363 trait effects, we obtain the mean marker level in bulk tissue of sample i,

364
$$\mu_i = \sum_h \alpha_h W_{h,i} + \sum_{h,k} \beta_{h,k} W_{h,i} X_{i,k} + \sum_l \gamma_l C_{i,l}.$$

365   With regards to the statistical model, we assume the error of the marker
366 level to be normally distributed with variance σ^2, independently among
367 samples, as

368
$$Y_i = \mu_i + \varepsilon_i,$$

369
$$\varepsilon_i \sim N(0, \sigma^2).$$

370 The statistical significance of all parameters is tested under the *full* model of
371 linear regression,

372
$$Y_i = \sum_h \alpha_h W_{h,i} + \sum_{h,k} \beta_{h,k} W_{h,i} X_{i,k} + \sum_l \gamma_l C_{i,l} + \varepsilon_i, \quad (2)$$

373 or its variations [5,9,13]. Alternatively, the cell-type-specific effects of traits
374 can be fitted and tested for one cell type h at a time by the *marginal* model,

375
$$Y_i = \sum_{h'} \alpha_{h'} W_{h',i} + \sum_k \beta_{h,k} W_{h,i} X_{i,k} + \sum_l \gamma_l C_{i,l} + \varepsilon_i, \quad (3)$$

376 or its variations [7,8,10,11,14].

Nonlinear regression

Nonlinear regression
Aiming to simultaneously analyze cell type composition in linear scale and
differential expression/methylation in log/logit scale, we develop a nonlinear

regression model. The differential analyses are performed after applying

normalizing transformation. The normalizing function is the natural logarithm

f = log for gene expression, and *f* = logit for methylation (see Background).

Conventional linear regression can be formulated by defining *f* as the identity

function. We denote the inverse function of *f* by *g*; *g* = exp for gene

expression, and *g* = logistic for methylation. Thus, *f* converts from the linear

scale to the normalized scale, and *g* does the opposite.

The marker level in a specific cell type (formula (1)) is modeled in the

normalized scale. The level is linearized by applying function *g*, then averaged

over cell types with weight $W_{h,i}$, and normalized by applying function *f*.

Combined with the tissue-uniform trait effects, the mean normalized marker

level in bulk tissue of sample *i* becomes

392
$$\mu_i = f\left(\sum_h W_{h,i}\, g\left(\alpha_h + \sum_k \beta_{h,k} X_{i,k}\right)\right) + \sum_l \gamma_l C_{i,l}. \quad (4)$$

393   We assume the normalized marker level to have an error that is normally

394   distributed with variance σ^2, independently among samples, as

395
$$f(Y_i) = \mu_i + \varepsilon_i, \quad (5)$$

396
$$\varepsilon_i \sim N(0, \sigma^2).$$

397   We obtain the ordinary least squares (OLS) estimator of the parameters by

398   minimizing the residual sum of squares,

399
$$\mathrm{RSS} = \sum_i (f(Y_i) - \mu_i)^2, \quad (6)$$

400   and then estimate the error variance as

401
$$\widehat{\sigma^2} = \frac{1}{n-p}\mathrm{RSS}, \quad (7)$$

where *n* is the number of samples and *p* is the number of parameters [[26],

section 6.3.1].

In the special case where the marker expression is homogeneous across

cell types, the formulae become simple. Suppose that α_h regardless of cell

type *h* equals α and that $\beta_{h,k}$ equals β_k. The regression formulae (4) and (5)

of sample *i* reduces to

408
$$f(Y_i) = \alpha + \sum_k \beta_k X_{i,k} + \sum_l \gamma_l C_{i,l} + \varepsilon_i. \quad (8)$$

409 On the other hand, the marginal model for cell type h in formula (3) reduces
410 to

$$Y_i = \alpha + \sum_k \beta_k W_{h,i} X_{i,k} + \sum_l \gamma_l C_{i,l} + \varepsilon_i.$$

412 Moreover, when the CV for cell type composition is low, the cell type
413 proportion $W_{h,i}$ of sample i approximately equals the average $\overline{W_h}$ taken over
414 samples. Thus, the formula reduces further to

$$Y_i = \alpha + \overline{W_h} \cdot \sum_k \beta_k X_{i,k} + \sum_l \gamma_l C_{i,l} + \varepsilon_i.$$

416 If we replace the input bulk expression level Y_i with the normalized value
417 $f(Y_i)$, the model becomes

$$f(Y_i) = \alpha + \overline{W_h} \cdot \sum_k \beta_k X_{i,k} + \sum_l \gamma_l C_{i,l} + \varepsilon_i. \quad (9)$$

Under the special case of cell-type-homogeneous expression and low-CV cell
type composition, formula (8) for nonlinear regression and formula (9) for
the marginal model with normalized input become almost identical. The
difference is the multiplication by constant $\overline{W_h}$, which does not change the
test statistics for β_k.

Ridge regression

The parameters $\beta_{h,k}$ for cell-type-specific effect cannot be estimated
accurately by ordinary linear regression because the regressors $W_{h,i} X_{i,k}$ in
equation (2) are highly correlated between cell types (see below).
Multicollinearity also occurs to the nonlinear case in formula (4) because of
local linearity. To cope with the multicollinearity, we apply ridge regression
with a regularization parameter $\lambda \geq 0$, and obtain the ridge estimator of the
parameters that minimizes

$$\mathrm{RSS} + \lambda \sum_{h,k} \beta_{h,k}^2, \quad (10)$$

433 where the second term penalizes $\beta_{h,k}$ for taking large absolute values. The
434 ridge estimator $\hat{\boldsymbol{\theta}}(\lambda)$ is asymptotically normally distributed (see Additional file
435 5: Supplementary note) with

$$436 \qquad \mathrm{Mean}\big[\widehat{\boldsymbol{\theta}}(\lambda)\big] = Q(\lambda)^{-1}\,Q(0)\,\boldsymbol{\theta}, \quad (11)$$

$$437 \qquad \mathrm{Var}\big[\widehat{\boldsymbol{\theta}}(\lambda)\big] = \sigma^2\,Q(\lambda)^{-1}\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\theta}}\right)^T\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\theta}}\right)Q(\lambda)^{-1}, \quad (12)$$

$$438 \qquad Q(\lambda) = \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\theta}}\right)^T\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\theta}}\right) + \lambda\begin{pmatrix} O & O & O \\ O & I & O \\ O & O & O \end{pmatrix} - (f(Y) - \boldsymbol{\mu}(\boldsymbol{\theta}))^T\cdot\left(\frac{\partial^2\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\theta}\partial\boldsymbol{\theta}^T}\right),$$

where $\boldsymbol{\mu}$ is the vector form of μ_i, $\boldsymbol{\theta}$ is the vector form of the parameters α_h,

$\beta_{h,k}$ and γ_l combined, $(\partial\boldsymbol{\mu}/\partial\boldsymbol{\theta})$ is the Jacobian matrix, $(\partial^2\boldsymbol{\mu}/\partial\boldsymbol{\theta}\partial\boldsymbol{\theta}^T)$ is the

array of Hessian matrices for μ_i taken over samples, and superscript T

indicates matrix transposition. The dot product of $(f(Y) - \boldsymbol{\mu}(\boldsymbol{\theta}))^T$ and the

array of Hessians is taken by multiplying for each sample and then summing

up over samples. The matrix after λ has one only in the diagonal

corresponding to $\beta_{h,k}$. The assigned value $\boldsymbol{\theta}$ is the true parameter value. By

taking the expectation of Q, we obtain a rougher approximation [27] as

$$447 \qquad \mathrm{Mean}\big[\widehat{\boldsymbol{\theta}}(\lambda)\big] = Q^*(\lambda)^{-1}\,Q^*(0)\,\boldsymbol{\theta}, \quad (13)$$

$$448 \qquad \mathrm{Var}\big[\widehat{\boldsymbol{\theta}}(\lambda)\big] = \sigma^2\,Q^*(\lambda)^{-1}\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\theta}}\right)^T\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\theta}}\right)Q^*(\lambda)^{-1}, \quad (14)$$

$$449 \qquad Q^*(\lambda) = \mathrm{E}[Q(\lambda)] = \left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\theta}}\right)^T\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\theta}}\right) + \lambda\begin{pmatrix} O & O & O \\ O & I & O \\ O & O & O \end{pmatrix}.$$

The matrices Q and Q^* are the observed and expected Fisher matrices

multiplied by σ^2 and adapted to ridge regression, respectively.

Since our objective is to predict the cell-type-specific trait effects, we

choose the regularization parameter λ that can minimize the mean squared

error (MSE) of $\beta_{h,k}$. Our methodology is based on [28]. To simplify the

explanation, we assume the Jacobian matrices $(\partial\boldsymbol{\mu}(\boldsymbol{\theta})/\partial\boldsymbol{\alpha})$, $(\partial\boldsymbol{\mu}(\boldsymbol{\theta})/\partial\boldsymbol{\beta})$ and

$(\partial\boldsymbol{\mu}(\boldsymbol{\theta})/\partial\boldsymbol{\gamma})$ to be mutually orthogonal, where $\boldsymbol{\alpha}$, $\boldsymbol{\beta}$ and $\boldsymbol{\gamma}$ are the vector

forms of α_h, $\beta_{h,k}$ and γ_l, respectively. Then, from formulae (13) and (14), the

ridge estimator $\widehat{\boldsymbol{\beta}}(\lambda)$ is asymptotically normally distributed with

$$459 \qquad \mathrm{Mean}\big[\widehat{\boldsymbol{\beta}}(\lambda)\big] = \left[\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right) + \lambda I\right]^{-1}\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)\boldsymbol{\beta},$$

$$460 \qquad \mathrm{Var}\big[\widehat{\boldsymbol{\beta}}(\lambda)\big] = \sigma^2\left[\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right) + \lambda I\right]^{-1}\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)$$

$$461 \qquad \left[\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right)^T\left(\frac{\partial\boldsymbol{\mu}(\boldsymbol{\theta})}{\partial\boldsymbol{\beta}}\right) + \lambda I\right]^{-1},$$

462 where the assigned values $\boldsymbol{\theta}$ and $\boldsymbol{\beta}$ are the true parameter values. We apply

463 singular value decomposition

464
$$\left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\beta}}\right) = UDV^T,$$

465 where U and V are orthogonal matrices, the columns of V are $\boldsymbol{v}_1, \cdots, \boldsymbol{v}_M$, and

466 the diagonals of diagonal matrix D are sorted $d_1 \geq \cdots \geq d_M \geq 0$. The bias,

467 variance and MSE of the ridge estimator are decomposed as

468
$$\mathrm{Bias}[\widehat{\boldsymbol{\beta}}(\lambda)] = \mathrm{E}[\widehat{\boldsymbol{\beta}}(\lambda) - \boldsymbol{\beta}]$$

469
$$= -\lambda \left[\left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\beta}}\right)^T \left(\frac{\partial \boldsymbol{\mu}(\boldsymbol{\theta})}{\partial \boldsymbol{\beta}}\right) + \lambda I\right]^{-1} \boldsymbol{\beta}$$

470
$$= \left\{\sum_{m=1}^{M} \boldsymbol{v}_m \frac{-\lambda}{d_m^2 + \lambda} \boldsymbol{v}_m^T\right\} \boldsymbol{\beta},$$

471
$$\mathrm{Var}[\widehat{\boldsymbol{\beta}}(\lambda)] = \sigma^2 \sum_{m=1}^{M} \boldsymbol{v}_m \frac{d_m^2}{\left(d_m^2 + \lambda\right)^2} \boldsymbol{v}_m^T,$$

472
$$\mathrm{MSE}[\widehat{\boldsymbol{\beta}}(\lambda)] = \mathrm{E}\left[\|\widehat{\boldsymbol{\beta}}(\lambda) - \boldsymbol{\beta}\|^2\right]$$

473
$$= \|\mathrm{Bias}[\widehat{\boldsymbol{\beta}}(\lambda)]\|^2 + \mathrm{tr}\left(\mathrm{Var}[\widehat{\boldsymbol{\beta}}(\lambda)]\right)$$

474
$$= \sum_{m=1}^{M} \left(\frac{\lambda}{d_m^2 + \lambda}\right)^2 (\boldsymbol{v}_m^T \boldsymbol{\beta})^2 + \left(\frac{d_m^2}{d_m^2 + \lambda}\right)^2 \left(\frac{\sigma^2}{d_m^2}\right). \quad (15)$$

For each m in the summation of (13), the minimum of the summand is

attained at $\lambda_m = \sigma^2/(\boldsymbol{v}_m^T \boldsymbol{\beta})^2$. To minimize MSE, we need to find some

"average" of the optimal λ_m over the range of m. Hoerl et al. [29] proposed

to use the harmonic mean $\lambda = M\sigma^2/\|\boldsymbol{\beta}\|^2$. However, if an OLS estimator $\widehat{\boldsymbol{\beta}}(0)$

is actually plugged into $\|\boldsymbol{\beta}\|^2$, the denominator is biased upwards, and the

computed mean is biased downwards. Indeed, with regards to the estimator

of $1/\sqrt{\lambda_m}$, we notice that

482
$$\frac{1}{\sigma} \boldsymbol{v}_m^T \widehat{\boldsymbol{\beta}}(0) \sim N\left(\frac{1}{\sigma} \boldsymbol{v}_m^T \boldsymbol{\beta}, \frac{1}{d_m^2}\right),$$

483 where the terms with larger m have larger variance. Thus, we take the

484 average of $(\boldsymbol{v}_m^T \widehat{\boldsymbol{\beta}}(0))^2/\sigma^2$, weighted by $d_m^2/\sum_{m=1}^{M} d_m^2$, and also subtract the

485 upward bias as,

486
$$\kappa = \frac{1}{\sum_{m=1}^{M} d_m^2} \sum_{m=1}^{M} \left\{\frac{d_m^2 \left(\boldsymbol{v}_m^T \widehat{\boldsymbol{\beta}}(0)\right)^2}{\sigma^2} - 1\right\}. \quad (16)$$

487   The weighting and subtraction were mentioned in [28], where the subtraction

488   term was dismissed, under the assumption of large effect-size $\boldsymbol{\beta}$. Since the

489   effect-size could be small in our application, we keep the subtraction term.

490   The statistic κ can be nonpositive, and is unbiased in the sense that

$$\mathrm{E}[\kappa] = \frac{1}{\sum_{m=1}^{M} d_m{}^2} \sum_{m=1}^{M} \frac{d_m{}^2 (\boldsymbol{v}_m{}^T \boldsymbol{\beta})^2}{\sigma^2} = \frac{1}{\sum_{m=1}^{M} d_m{}^2} \sum_{m=1}^{M} \frac{d_m{}^2}{\lambda_m}$$

492   equals the weighted sum of $1/\lambda_m$. Our choice of regularization parameter is

$$\lambda = \begin{cases} 1/\kappa & \text{if } \kappa > 0, \\ d_1{}^2 & \text{otherwise,} \end{cases} \quad (17)$$

494   where $d_1{}^2$ is taken instead of positive infinity.

Implementation of omicwas package

For each omics marker, the parameters $\boldsymbol{\alpha}$, $\boldsymbol{\beta}$ and $\boldsymbol{\gamma}$ (denoted in combination

by $\boldsymbol{\theta}$) are estimated and tested by nonlinear ridge regression in the following

steps. As we assume the magnitude of trait effects $\boldsymbol{\beta}$ and $\boldsymbol{\gamma}$ to be much

smaller than that of basal marker level $\boldsymbol{\alpha}$, we first fit $\boldsymbol{\alpha}$ alone for numerical

stability.

1. Compute OLS estimator $\widehat{\boldsymbol{\alpha}}(0)$ by minimizing formula (6) under $\boldsymbol{\beta} = \boldsymbol{\gamma} = \mathbf{0}$.

Apply Wald test.

2. Calculate $\widehat{\sigma^2}$ by formula (7). Use it as a substitute for σ^2. The residual

degrees of freedom $n - p$ is the number of samples minus the number of

parameters in $\boldsymbol{\alpha}$.

3. Compute OLS estimators $\widehat{\boldsymbol{\beta}}(0)$ and $\widehat{\boldsymbol{\gamma}}(0)$ by minimizing formula (6) under

$\boldsymbol{\alpha} = \widehat{\boldsymbol{\alpha}}(0)$. Let $\widehat{\boldsymbol{\theta}}(0) = (\widehat{\boldsymbol{\alpha}}(0)^T, \widehat{\boldsymbol{\beta}}(0)^T, \widehat{\boldsymbol{\gamma}}(0)^T)^T$.

4. Apply singular value decomposition $\left(\partial \boldsymbol{\mu}(\widehat{\boldsymbol{\theta}}(0)) / \partial \boldsymbol{\beta} \right) = UDV^T$.

5. Calculate κ and then the regularization parameter λ by formulae (16)

and (17).

6. Compute ridge estimators $\widehat{\boldsymbol{\beta}}(\lambda)$ and $\widehat{\boldsymbol{\gamma}}(\lambda)$ by minimizing formula (10)

under $\boldsymbol{\alpha} = \widehat{\boldsymbol{\alpha}}(0)$. Let $\widehat{\boldsymbol{\theta}}(\lambda) = (\widehat{\boldsymbol{\alpha}}(0)^T, \widehat{\boldsymbol{\beta}}(\lambda)^T, \widehat{\boldsymbol{\gamma}}(\lambda)^T)^T$.

7. Approximate the variance of ridge estimator, according to formula (12),

by

$$515 \qquad \mathrm{Var}\left[\begin{pmatrix}\widehat{\boldsymbol{\beta}}(\lambda)\\\widehat{\boldsymbol{\gamma}}(\lambda)\end{pmatrix}\right] = \widehat{\sigma^2}\, Q(\lambda)^{-1} \left(\frac{\partial \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}}\right)^T \left(\frac{\partial \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}}\right) Q(\lambda)^{-1},$$

$$516 \qquad Q(\lambda) = \left(\frac{\partial \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}}\right)^T \left(\frac{\partial \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}}\right) + \lambda \begin{pmatrix} I & O \\ O & O \end{pmatrix}$$

$$517 \qquad - \left(f(Y) - \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)\right)^T \cdot \left(\frac{\partial^2 \boldsymbol{\mu}\left(\widehat{\boldsymbol{\theta}}(\lambda)\right)}{\partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix} \partial \begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}^T}\right).$$

518   8. Apply the "non-exact" t-type test [30]. For the s-th coordinate,

$$520 \qquad \frac{\begin{pmatrix}\widehat{\boldsymbol{\beta}}(\lambda)\\\widehat{\boldsymbol{\gamma}}(\lambda)\end{pmatrix}_s}{\sqrt{\mathrm{Var}\left[\begin{pmatrix}\widehat{\boldsymbol{\beta}}(\lambda)\\\widehat{\boldsymbol{\gamma}}(\lambda)\end{pmatrix}\right]_{s,s}}} \sim t_{n-p}, \quad (18)$$

under the null hypothesis $\begin{pmatrix}\boldsymbol{\beta}\\\boldsymbol{\gamma}\end{pmatrix}_s = 0$.

The formula (18) is the same as a Wald test, but the test differs, because the

ridge estimators are not maximum-likelihood estimators. The algorithm was

implemented as a package for the R statistical language. We used the NL2SOL

algorithm of the PORT library [31] for minimization.

In analyses of quantitative trait locus (QTL), such as methylation QTL

(mQTL) and expression QTL (eQTL), an association analysis that takes the

genotypes of a single nucleotide polymorphism (SNP) as $X_{i,k}$ is repeated for

many SNPs. In order to speed up the computation, we perform rounds of

linear regression. First, the parameters $\widehat{\boldsymbol{\alpha}}(0)$ and $\widehat{\boldsymbol{\gamma}}(0)$ are fit by ordinary

linear regression under $\boldsymbol{\beta} = \mathbf{0}$, which does not depend on $X_{i,k}$. By taking the

residuals, we practically dispense with $\widehat{\boldsymbol{\alpha}}(0)$ and $\widehat{\boldsymbol{\gamma}}(0)$ in the remaining steps.

Next, for $X_{i,k}$ of each SNP, $\widehat{\boldsymbol{\beta}}(0)$ is fit by ordinary linear regression under $\boldsymbol{\alpha} =$

$\widehat{\boldsymbol{\alpha}}(0)$, $\boldsymbol{\gamma} = \widehat{\boldsymbol{\gamma}}(0)$. The regularization parameter λ is computed according to

steps 4 and 5 above. Finally, $\widehat{\boldsymbol{\beta}}(\lambda)$ is fitted and tested by linear ridge

regression under $\boldsymbol{\alpha} = \widehat{\boldsymbol{\alpha}}(0)$, $\boldsymbol{\gamma} = \widehat{\boldsymbol{\gamma}}(0)$.

Multicollinearity of interaction terms

The regressors for cell-type-specific trait effects in the full model (equation (2)) are the interaction terms $W_{h,i}X_{i,k}$. To assess multicollinearity, we mathematically derive the correlation coefficient between two interaction terms $W_{h,i}X_{i,k}$ and $W_{h',i}X_{i,k}$. In this section, we treat $W_{h,i}$, $W_{h',i}$ and $X_{i,k}$ as sampled instances of random variables W_h, $W_{h'}$ and X_k, respectively; note that the sample index i is omitted. For simplicity, we assume W_h and $W_{h'}$ are independent of X_k. Let $\mathrm{E}[\bullet]$, $\mathrm{Var}[\bullet]$, $\mathrm{Cov}[\bullet]$, $\mathrm{Cor}[\bullet]$ and $\mathrm{CV}[\bullet]$ denote the expectation, variance, covariance, correlation and coefficient of variation, respectively. Since X_k is centered, $\mathrm{E}[W_hX_k] = \mathrm{E}[W_{h'}X_k] = 0$. The correlation coefficient between interaction terms becomes

$$\mathrm{Cor}[W_hX_k, W_{h'}X_k] = \frac{\mathrm{E}[W_hX_kW_{h'}X_k]}{\sqrt{\mathrm{E}\left[{W_h}^2{X_k}^2\right]}\sqrt{\mathrm{E}\left[{W_{h'}}^2{X_k}^2\right]}}$$

$$= \frac{\mathrm{E}[W_hW_{h'}]}{\sqrt{\mathrm{E}\left[{W_h}^2\right]}\sqrt{\mathrm{E}\left[{W_{h'}}^2\right]}}$$

$$= \frac{\mathrm{Cov}[W_h, W_{h'}] + \mathrm{E}[W_h]\,\mathrm{E}[W_{h'}]}{\sqrt{\mathrm{Var}[W_h] + \mathrm{E}[W_h]^2}\sqrt{\mathrm{Var}[W_{h'}] + \mathrm{E}[W_{h'}]^2}}$$

$$= \frac{\dfrac{\mathrm{Cor}[W_h, W_{h'}]\sqrt{\mathrm{Var}[W_h]}\sqrt{\mathrm{Var}[W_{h'}]}}{\mathrm{E}[W_h]\mathrm{E}[W_{h'}]} + 1}{\sqrt{\dfrac{\mathrm{Var}[W_h]}{\mathrm{E}[W_h]^2} + 1}\sqrt{\dfrac{\mathrm{Var}[W_{h'}]}{\mathrm{E}[W_{h'}]^2} + 1}}$$

$$= \frac{\mathrm{Cor}[W_h, W_{h'}]\,\mathrm{CV}[W_h]\,\mathrm{CV}[W_{h'}] + 1}{\sqrt{\mathrm{CV}[W_h]^2 + 1}\sqrt{\mathrm{CV}[W_{h'}]^2 + 1}}. \quad (19)$$

If $\mathrm{CV}[W_h]$ and $\mathrm{CV}[W_{h'}]$ approach zero, the correlation of interaction terms approaches one, irrespective of $\mathrm{Cor}[W_h, W_{h'}]$.

EWAS of rheumatoid arthritis and age

EWAS datasets for rheumatoid arthritis were downloaded from the Gene Expression Omnibus. Using the RnBeads package (version 2.2.0) [32] of R, IDAT files of HumanMethylation450 array were preprocessed by removing

low quality samples and markers, by normalizing methylation level, and by
removing markers on sex chromosomes and outlier samples. The association
of methylation level with disease status was tested with adjustment for sex,
age, smoking status and experiment batch; the covariates were assumed to
have uniform effects across cell types. Alternatively, the association of
methylation level with age was tested with adjustment for disease status ,
sex, smoking status and experiment batch. After quality control, dataset
GSE42861 included bulk peripheral blood leukocyte data for 336 cases and
322 controls [22].

The cell type composition of bulk samples was imputed using the
Houseman algorithm [33] in the GLINT software (version 1.0.4) [34]. The
reference data of GLINT software characterizes seven cell types [35] by 300
CpG sites [36], of which 284 were measured in our data. We used prediction
results for the seven cell types (Table 1).

Dataset GSE131989 included sorted CD14$^+$ monocyte data for 63 cases
and 31 controls [23]. By meta-analysis of GSE131989 and GSE87095 [21],
we obtained sorted CD19$^+$ B cell data for 108 cases and 95 controls. Under
the nominal significance level P < 0.05 (two-sided), the number of CpG sites
up- or down-regulated in cases were 20,869 (5%) and 14,911 (3%),
respectively, in CD14$^+$ monocyte and 28,004 (6%) and 26,582 (6%) in CD19$^+$
B cell.

From the Gene Expression Omnibus dataset GSE56047 [25], we obtained
sorted CD14$^+$ monocyte data for 1200 samples and sorted CD4$^+$ T cell data
for 214 samples. Under the nominal significance level P < 0.05 (two-sided),
the number of CpG sites up- or down-regulated by higher age were 45,283
(10%) and 80,871 (18%), respectively, in CD14$^+$ monocyte and 35,822 (8%)
and 25,020 (5%) in CD4$^+$ T cell.

Differential gene expression by age

Whole blood RNA-seq data of GTEx v7 was downloaded from the GTEx
website [24]. Genes of low quality or on sex chromosomes were removed,
expression level was normalized, outlier samples were removed, and 389

samples were retained. The association of read count with age was tested
with adjustment for sex.

The cell type composition of bulk samples was imputed using the
DeconCell package (version 0.1.0) [10] of R. The reference data of DeconCell
characterizes 33 cell types by two to 217 signature genes. Our data measured
39% of the signature genes. We used prediction results for six main cell types
(Table 2), for which the prediction performance was 44.4 to 90.9.

From the Gene Expression Omnibus dataset GSE56047 [25], we obtained
sorted CD14$^+$ monocyte data for 1202 samples and sorted CD4$^+$ T cell data
for 214 samples. Under the nominal significance level P < 0.05 (two-sided),
the number of genes up- or down-regulated by higher age were 2715 (11%)
and 3240 (13%), respectively, in CD14$^+$ monocyte and 1082 (4%) and 1246
(5%) in CD4$^+$ T cell.

Simulation of cell-type-specific disease association
Bulk tissue sample data for case-control comparison were simulated based
on the above-mentioned EWAS dataset GSE42861 and GTEx gene expression
dataset. We randomly assigned the case-control status to the samples.
Among the omics markers, 2.5% were set to be up-regulated in cases in
single cell type, 2.5% were similarly down-regulated, and 95% were
unrelated to case-control status. The cell-type-specific effect-size of the
differentially expressed markers was fixed within a simulation trial, and was
chosen from methylation OR of 1.3, 1.6 or 1.9 for EWAS [20] and fold-change
of 1.7, 3.0 or 5.0 for gene expression analysis; the effect-sizes correspond to
log(1.3), log(1.6) and log(1.9) or log(1.7), log(3.0) and log(5.0) in
normalized scale. If the mean methylation level of a CpG site in cases and
controls are μ_{case} and μ_{control}, respectively, the methylation odds become
$\mu_{\text{case}}/(1 - \mu_{\text{case}})$ and $\mu_{\text{control}}/(1 - \mu_{\text{control}})$. The methylation OR represents the
case-control contrast of methylation level by the ratio of odds,
$\{\mu_{\text{case}}/(1 - \mu_{\text{case}})\}/\{\mu_{\text{control}}/(1 - \mu_{\text{control}})\}$ (see [20]).

For each effect-size, we performed 50 simulation trials. In each simulation
trial, we randomly assigned half of the samples as cases ($X_{i,k} = 0.5$) and the

other half as controls ($X_{i,k} = -0.5$). We retained the covariates matrix $C_{i,l}$ and

the cell type composition matrix $W_{h,i}$ from the original data. From the original

bulk expression level matrix $Y_{i,j}$, 95% of the markers were randomly chosen

and retained; these markers had no association with disease because the

case-control status was randomized. Cell-type-specific association was

introduced into the remaining 5% of markers, such that an equal number of

markers were up- or down-regulated in each cell type. For example, in the

EWAS dataset, $451,725 \times 0.05 \times 0.5 \div 7 = 1613$ CpG sites were up-regulated in

neutrophils of cases.

The bulk expression level of a marker j with normalized-scale effect-size

β specific to a cell type h was generated as follows. First, the average μ and

the variance σ^2 of the normalized bulk expression level $f(Y_{i,j})$ in the original

data was measured. Next, we generated normalized expression level in each

cell type. For cases, the expression level in cell type h was randomly sampled

from the normal distribution $N(\mu + \beta, \sigma^2)$ and the expression level in each of

the other cell types was sampled from $N(\mu, \sigma^2)$. For controls, the expression

level in each cell type was sampled from $N(\mu, \sigma^2)$. Finally, for each individual,

the expression levels in cell types were converted to the linear scale,

multiplied by the cell type composition and added, to obtain the bulk

expression level in linear scale.

In the truly disease-associated cell type h, we introduced signal β and

noise σ^2. The signal level was fixed in a simulation trial, for example to

methylation OR = 1.3. Since the noise level was taken from real data, the

level varied between markers. In the process of obtaining bulk expression the

expression of all cell types was mixed, which dilutes the signal. The signal

dilution becomes stronger if h is a minor cell type. The mixing process adds

noise from other cell types, which becomes stronger if h is a minor cell type.

Consequently, minor cell types tend to manifest weaker association in bulk

tissue. We empirically measured the strength of association by the coefficient

of determination, R^2, for the marginal model. The coefficient of determination

is defined as the proportion of variance explained by the model, and

$R^2/(1 - R^2)$ equals the signal-to-noise ratio. Under methylation OR of 1.3, 1.6,

1.9 for EWAS simulation, the median R^2 was 0.322, 0.589, 0.712 for

neutrophils, 0.010, 0.033, 0.057 for NK cells, and 0.001, 0.003, 0.005 for

eosinophils. Under fold-change of 1.7, 3.0 or 5.0 for gene expression

simulation, the median R^2 was 0.135, 0.331, 0.434 for granulocytes, 0.007,

0.026, 0.049 for CD8$^+$ T cells, and 0.001, 0.003, 0.007 for B cells.

For gene expression, we also simulated a scenario where cell-type-specific

disease effect occurs in cell type marker genes. The simulation procedure is

same as above except that the expression level was set to zero (in linear

scale) in all cell types other than the target cell type *h*, for both cases and

controls.

Evaluation of statistical methods

Cell-type-specific effects of traits was statistically tested by using bulk tissue

data as input. We applied the omicwas package with the normalizing function

f = log, logit, identity without ridge regularization (omicwas.log,

omicwas.logit, omicwas.identity) or under ridge regression

(omicwas.log.ridge, omicwas.logit.ridge, omicwas.identity.ridge). The

omicwas package was used also for conventional linear regression under the

full and marginal models. We also developed a hybrid of marginal and full

models (Marginal.Full005): if the effect direction agreed in two models and if

P < 0.05 in the full model, we adopted the Z-score of the marginal model;

otherwise, the Z-score was set to zero.

Among previous methods, we evaluated those that accept cell type

composition as input and compute test statistics for cell-type-specific

association. For DNA methylation data, we applied TOAST (version 1.2.0) [9],

CellDMC (version 2.0.2) [13] and TCA (version 1.0.0) [14]. For gene

expression data, we applied TOAST and csSAM (version 1.4) [5]. For csSAM,

we either fitted all cell types together or one cell type at a time, and denoted

the results as csSAM.lm and csSAM.monovariate, respectively. The csSAM

method is applicable to binomial traits but not to quantitative traits.

For simulated data of EWAS dataset GSE42861, we adopted the

significance level P < 2.4 × 10^{-7}, which accounts for the correlation among

the probes on HumanMethylation450 array [37]. For the GTEx gene

expression dataset, multiple testing was controlled by the Benjamini-
Hochberg procedure with the false discovery rate <5% in each cell type [38].
The performance of an algorithm for the simulated data was assessed by
sensitivity, specificity and precision. The performance measures were
obtained from each simulation trial. For a target cell type h, we counted the
four possible outcomes, true positives (TP_h), true negatives (TN_h), false
positives (FP_h) and false negatives (FN_h). The sum $\text{TP}_h + \text{TN}_h + \text{FP}_h + \text{FN}_h$
equals the total number of omics markers (which was 451,725 CpG sites for
DNA methylation and 14,038 genes for gene expression). For 5% of the
markers, one randomly selected cell type h^* was set to be truly associated
with disease status at data generation. The remaining 95% of the markers
were null cases with no truly associated cell types. The outcome counts can
be subtotaled according to the truly associated cell type, which is denoted in
superscript,

698
$$\text{TP}_h = \text{TP}_h^{h^*=h},$$

699
$$\text{TN}_h = \sum_{h^* \neq h} \text{TN}_h^{h^*} + \text{TN}_h^{\text{Null}},$$

700
$$\text{FP}_h = \sum_{h^*} \text{FP}_h^{h^*} + \text{FP}_h^{\text{Null}},$$

701
$$\text{FN}_h = \text{FN}_h^{h^*=h}.$$

702 Remark that FP_h can occur when in cell type h a marker is truly up-regulated
703 in disease cases but an algorithm predicts the marker to be down-regulated
704 in h. The performance measures can be represented as

705
$$\text{sensitivity}_h = \frac{\text{TP}_h}{\text{TP}_h + \text{FN}_h} = \frac{\text{TP}_h^{h^*=h}}{\text{TP}_h^{h^*=h} + \text{FN}_h^{h^*=h}},$$

706
$$\text{specificity}_h = \frac{\text{TN}_h}{\text{TN}_h + \text{FP}_h} = \frac{\sum_{h^* \neq h} \text{TN}_h^{h^*} + \text{TN}_h^{\text{Null}}}{\text{FP}_h^{h^*=h} + \sum_{h^* \neq h}(\text{TN}_h^{h^*} + \text{FP}_h^{h^*}) + (\text{TN}_h^{\text{Null}} + \text{FP}_h^{\text{Null}})},$$

707
$$\text{precision}_h = \frac{\text{TP}_h}{\text{TP}_h + \text{FP}_h} = \frac{\text{TP}_h^{h^*=h}}{(\text{TP}_h^{h^*=h} + \text{FP}_h^{h^*=h}) + \sum_{h^* \neq h} \text{FP}_h^{h^*} + \text{FP}_h^{\text{Null}}}.$$

Whereas sensitivity is obtained solely from markers that are truly associated
in the target cell type h, the specificity and precision are obtained by
aggregating with the markers associated in other cell types and the null
markers.

For the association with rheumatoid arthritis and age, "true" association
was determined from the measurements in physically sorted blood cells,
under the nominal significance level $P < 0.05$ (two-sided). In the same
manner as the simulation analysis, we assessed the sensitivity, specificity
and precision.
717

Supplementary information
Additional file 1: Table S1. Blood cell type proportion in Tsimane
Amerindians, Caucasians and Hispanics.
Additional file 2: Fig. S1. Sensitivity for detecting cell-type-specific
association in simulated data for gene expression of marker genes.
Additional file 3: Fig. S2. Specificity for detecting cell-type-specific
association in simulated data for gene expression of marker genes.
Additional file 4: Fig. S3. Precision (positive predictive value) for detecting
cell-type-specific association in simulated data for gene expression of marker
genes.
Additional file 5: Supplementary note. Asymptotic distribution of ridge
estimator.

Abbreviations
CV: coefficient of variation, eQTL: expression QTL, EWAS: epigenome-wide
association study, mQTL: methylation QTL, MSE: mean squared error, OLS:
ordinary least squares, OR: odds ratio, QTL: quantitative trait locus, SNP:
single nucleotide polymorphism

Declarations

Ethics approval and consent to participate
Not applicable.

Consent for publication
Not applicable.

Availability of data and materials

740 The datasets generated and analyzed during the current study are available
742 in the figshare repository, https://dx.doi.org/10.6084/m9.figshare.10718282

Competing interests

744 The authors declare that they have no competing interests.

Funding

746 This work was supported by JSPS KAKENHI [grant number JP16K07218] and
747 by the NCGM Intramural Research Fund [grant numbers 19A2004, 20A1013].
748 The funding body had no role in the design and collection of the study,
749 experiments, analyses and interpretations of data, and in writing the
750 manuscript.

Author's contributions

752 FT developed the methodology, wrote the software, implemented the study,
753 and wrote the manuscript. NK revised the manuscript. All authors read and
754 approved the final manuscript.

Acknowledgements

756 Not applicable.

757
758

References

References

1. Teschendorff AE, Zheng SC. Cell-type deconvolution in epigenome-wide

association studies: a review and recommendations. Epigenomics.

2017;9:757–68.

2. Sturm G, Finotello F, Petitprez F, Zhang JD, Baumbach J, Fridman WH, et

al. Comprehensive evaluation of transcriptome-based cell-type

quantification methods for immuno-oncology. Bioinformatics.

2019;35:i436–45.

3. Ghosh D. Mixture models for assessing differential expression in complex

tissues using microarray data. Bioinformatics. 2004;20:1663–9.

4. Stuart RO, Wachsman W, Berry CC, Wang-Rodriguez J, Wasserman L,

Klacansky I, et al. In silico dissection of cell-type-associated patterns of

gene expression in prostate cancer. Proc Natl Acad Sci USA. National

Academy of Sciences; 2004;101:615–20.

5. Shen-Orr SS, Tibshirani R, Khatri P, Bodian DL, Staedtler F, Perry NM, et

al. Cell type–specific gene expression differences in complex tissues. Nat

Meth. Nature Publishing Group; 2010;7:287–9.

6. Erkkilä T, Lehmusvaara S, Ruusuvuori P, Visakorpi T, Shmulevich I,

Lähdesmäki H. Probabilistic analysis of gene expression measurements from

heterogeneous tissues. Bioinformatics. 2010;26:2571–7.

7. Kuhn A, Thu D, Waldvogel HJ, Faull RLM, Luthi-Carter R. Population-

specific expression analysis (PSEA) reveals molecular changes in diseased

brain. Nat Meth. Nature Publishing Group; 2011;8:945–7.

8. Westra H-J, Arends D, Esko T, Peters MJ, Schurmann C, Schramm K, et

al. Cell Specific eQTL Analysis without Sorting Cells. Pastinen T, editor. PLoS

Genet. Public Library of Science; 2015;11:e1005223–17.

9. Li Z, Wu Z, Jin P, Wu H. Dissecting differential signals in high-throughput

data from complex tissues. Hancock J, editor. Bioinformatics.

2019;35:3898–905.

10. Aguirre-Gamboa R, de Klein N, di Tommaso J, Claringbould A, van der
Wijst MG, de Vries D, et al. Deconvolution of bulk blood eQTL effects into
immune cell subpopulations. BMC Bioinformatics. 2020;21:243.

11. Montaño CM, Irizarry RA, Kaufmann WE, Talbot K, Gur RE, Feinberg AP,
et al. Measuring cell-type specific differential methylation in human brain
tissue. Genome Biol. BioMed Central; 2013;14:R94–9.

12. White N, Benton M, Kennedy D, Fox A, Griffiths L, Lea R, et al.
Accounting for cell lineage and sex effects in the identification of cell-
specific DNA methylation using a Bayesian model selection algorithm.
Sawalha AH, editor. PLoS ONE. 2017;12:e0182455–18.

13. Zheng SC, Breeze CE, Beck S, Teschendorff AE. Identification of
differentially methylated cell types in epigenome-wide association studies.
Nat Meth. Nature Publishing Group; 2018;15:1059–66.

14. Rahmani E, Schweiger R, Rhead B, Criswell LA, Barcellos LF, Eskin E, et
al. Cell-type-specific resolution epigenetics without the need for cell sorting
or single-cell biology. Nature Communications. Nature Publishing Group;
2019;10:3417–11.

15. Cobos FA, Vandesompele J, Mestdagh P. Computational deconvolution
of transcriptomics data from mixed cell populations. Bioinformatics.
2018;34:1969–79.

16. Hoyle DC, Rattray M, Jupp R, Brass A. Making sense of microarray data
distributions. Bioinformatics. 2002;18:576–84.

17. Du P, Zhang X, Huang C-C, Jafari N, Kibbe WA, Hou L, et al.
Comparison of Beta-value and M-value methods for quantifying methylation
levels by microarray analysis. BMC Bioinformatics. BioMed Central;
2010;11:1–9.

18. Zhuang J, Widschwendter M, Teschendorff AE. A comparison of feature

selection and classification methods in DNA methylation studies using the

Illumina Infinium platform. BMC Bioinformatics. BioMed Central;

2012;13:1–14.

19. Aiken LS, West SG. Multiple Regression: Testing and Interpreting

Interactions. Sage Publications; 1991.

20. Rakyan VK, Down TA, Balding DJ, Beck S. Epigenome-wide association

studies for common human diseases. Nat Rev Genet. 2011;12:529–41.

21. Julià A, Absher D, López-Lasanta M, Palau N, Pluma A, Waite Jones L, et

al. Epigenome-wide association study of rheumatoid arthritis identifies

differentially methylated loci in B cells. Hum Mol Genet. 2017;26:2803–11.

22. Liu Y, Aryee MJ, Padyukov L, Fallin MD, Hesselberg E, Runarsson A, et

al. Epigenome-wide association data implicate DNA methylation as an

intermediary of genetic risk in rheumatoid arthritis. Nat Biotechnol. Nature

Publishing Group; 2013;31:142–7.

23. Rhead B, Holingue C, Cole M, Shao X, Quach HL, Quach D, et al.

Rheumatoid Arthritis Naive T Cells Share Hypermethylation Sites With

Synoviocytes. Arthritis & Rheumatology. John Wiley & Sons, Ltd;

2017;69:550–9.

24. GTEx Consortium. Genetic effects on gene expression across human

tissues. Nature. Nature Publishing Group; 2017;550:204–13.

25. Reynolds LM, Taylor JR, Ding J, Lohman K, Johnson C, Siscovick D, et

al. Age-related variations in the methylome associated with gene expression

in human monocytes and T cells. Nature Communications. 2014;5:5366.

26. Riazoshams H, Midi H, Ghilagaber G. Robust Nonlinear Regression: with

Applications using R. John Wiley & Sons; 2019.

27. Lim C. Robust ridge regression estimators for nonlinear models with

applications to high throughput screening assay data. Statist. Med.
2014;34:1185–98.

28. Lawless JF, Wang P. A simulation study of ridge and other regression
estimators. Communications in Statistics - Theory and Methods.
1976;5:307–23.

29. Hoerl AE, Kannard RW, Baldwin KF. Ridge regression: some simulations.
Communications in Statistics - Theory and Methods. 1975;4:105–23.

30. Halawa AM, Bassiouni El MY. Tests of regression coefficients under ridge
regression models. Journal of Statistical Computation and Simulation.
2000;65:341–56.

31. Dennis JE, Gay DM, Welsch RE. An adaptive nonlinear least-squares
algorithm. ACM Transactions on Mathematical Software. 1981;7:348–68.

32. Müller F, Scherer M, Assenov Y, Lutsik P, Walter J, Lengauer T, et al.
RnBeads 2.0: comprehensive analysis of DNA methylation data. Genome
Biol. BioMed Central; 2019;20:55–12.

33. Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ,
Nelson HH, et al. DNA methylation arrays as surrogate measures of cell
mixture distribution. BMC Bioinformatics. 2012;13:86.

34. Rahmani E, Yedidim R, Shenhav L, Schweiger R, Weissbrod O, Zaitlen
N, et al. GLINT: a user-friendly toolset for the analysis of high-throughput
DNA-methylation array data. Hancock JM, editor. Bioinformatics.
2017;33:1870–2.

35. Reinius LE, Acevedo N, Joerink M, Pershagen G, Dahlén S-E, Greco D, et
al. Differential DNA Methylation in Purified Human Blood Cells: Implications
for Cell Lineage and Studies on Disease Susceptibility. Ting AH, editor. PLoS
ONE. Public Library of Science; 2012;7:e41361–13.

36. Koestler D. Improving Cell Mixture Deconvolution by Identifying Optimal

DNA methylation Libraries (IDOL). BMC Bioinformatics. BMC Bioinformatics;

2016;:1–21.

37. Saffari A, Silver MJ, Zavattari P, Moi L, Columbano A, Meaburn EL, et al.

Estimation of a significance threshold for epigenome-wide association

studies. Genet Epidemiol. John Wiley & Sons, Ltd; 2017;42:20–33.

38. Anders S, McCarthy DJ, Chen Y, Okoniewski M, Smyth GK, Huber W, et

al. Count-based differential expression analysis of RNA sequencing data

using R and Bioconductor. Nat Protoc. 2013;8:1765–86.

TABLES

Table 1A Blood cell type proportion in rheumatoid arthritis dataset

Cell type	Neu	CD4$^+$T	CD8$^+$T	NK	Mono	Bcells	Eos
Mean	0.59	0.10	0.08	0.08	0.07	0.07	0.01
SD	0.11	0.06	0.05	0.04	0.02	0.03	0.02
CV	0.2	0.6	0.6	0.5	0.3	0.4	2.7

SD, standard deviation; CV, coefficient of variation

Table 1B Correlation between blood cell type proportion and rheumatoid arthritis (X_k)

r	Neu	CD4$^+$T	CD8$^+$T	NK	Mono	Bcells	Eos	X_k=Disease
Neu	1	-0.68	-0.60	-0.46	-0.06	-0.49	-0.48	0.44
CD4$^+$T	-0.68	1	0.14	0.05	-0.17	0.38	0.26	-0.33
CD8$^+$T	-0.60	0.14	1	0.08	-0.05	0.19	0.13	-0.27
NK	-0.46	0.05	0.08	1	-0.04	0.01	0.11	-0.27
Mono	-0.06	-0.17	-0.05	-0.04	1	-0.17	0.05	0.10
Bcells	-0.49	0.38	0.19	0.01	-0.17	1	0.11	-0.22
Eos	-0.48	0.26	0.13	0.11	0.05	0.11	1	-0.10

Table 1C Correlation between interaction terms

r	Neu*X_k	CD4$^+$T*X_k	CD8$^+$T*X_k	NK*X_k	Mono*X_k	Bcells*X_k	Eos*X_k
Neu*X_k	1	0.83	0.80	0.85	0.93	0.90	0.27
CD4$^+$T*X_k	0.83	1	0.78	0.78	0.83	0.88	0.42
CD8$^+$T*X_k	0.80	0.78	1	0.77	0.82	0.83	0.35
NK*X_k	0.85	0.78	0.77	1	0.85	0.83	0.35
Mono*X_k	0.93	0.83	0.82	0.85	1	0.88	0.35
Bcells*X_k	0.90	0.88	0.83	0.83	0.88	1	0.36
Eos*X_k	0.27	0.42	0.35	0.35	0.35	0.36	1

Neu, neutrophils; Mono, monocytes; Eos, eosinophils.

Table 2A Blood cell type proportion in GTEx dataset

Cell type	Gran	CD4$^+$T	CD8$^+$T	Mono	NK	Bcells
Mean	0.53	0.22	0.10	0.07	0.05	0.03
SD	0.037	0.020	0.013	0.004	0.012	0.003
CV	0.1	0.1	0.1	0.1	0.2	0.1

SD, standard deviation; CV, coefficient of variation

Table 2B Correlation between blood cell type proportion and age (X_k)

r	Gran	CD4$^+$T	CD8$^+$T	Mono	NK	Bcells	X_k=Age
Gran	1	-0.89	-0.83	0.56	-0.76	-0.41	-0.23
CD4$^+$T	-0.89	1	0.59	-0.64	0.50	0.51	0.14
CD8$^+$T	-0.83	0.59	1	-0.40	0.59	0.15	0.15
Mono	0.56	-0.64	-0.40	1	-0.44	-0.42	0.02
NK	-0.76	0.50	0.59	-0.44	1	0.13	0.31
Bcells	-0.41	0.51	0.15	-0.42	0.13	1	-0.03

Table 2C Correlation between interaction terms

r	Gran*X_k	CD4$^+$T*X_k	CD8$^+$T*X_k	Mono*X_k	NK*X_k	Bcells*X_k
Gran*X_k	1	0.99	0.98	1.00	0.96	0.99
CD4$^+$T*X_k	0.99	1	1.00	0.99	0.98	1.00
CD8$^+$T*X_k	0.98	1.00	1	0.99	0.98	0.99
Mono*X_k	1.00	0.99	0.99	1	0.96	0.99
NK*X_k	0.96	0.98	0.98	0.96	1	0.97
Bcells*X_k	0.99	1.00	0.99	0.99	0.97	1

Gra, granulocytes; Mono, monocytes.

FIGURE LEGENDS

Figure 1

Contour plot of the correlation coefficient between interaction terms $W_h X_k$

and $W_{h'} X_k$. W_h and $W_{h'}$ represent proportions of cell types h and h', and X_k

represents the value of trait k. For this plot, we assume the coefficient of

variation $\mathrm{CV}[W_h]$ and $\mathrm{CV}[W_{h'}]$ to be equal. As the CV decreases 0.6, 0.4 to

0.2, the correlation coefficient raises >0.5, >0.7 to >0.9, over most range of

$\mathrm{Cor}[W_h, W_{h'}]$.

Figure 2

Sensitivity for detecting cell-type-specific association in simulated data for

DNA methylation. Panels are aligned in rows according to the simulation

settings with the methylation odds ratio of 1.3, 1.6 or 1.9. In each row, panels

for different cell types are aligned in decreasing order of proportion. The

vertical axis indicates sensitivity. In each panel, results from different

algorithms are aligned horizontally in different colors. Results from 20

simulation trials are summarized in a box plot. The middle bar of the box plot

indicates the median, and the lower and upper hinges correspond to the first

and third quartiles. The whiskers extend to the value no further than 1.5 ×

inter-quartile range from the hinges. MethOR, methylation odds ratio; Neu,

neutrophils; Mono, monocytes; Eos, eosinophils.

Figure 3

Specificity for detecting cell-type-specific association in simulated data for

DNA methylation. The figure format is same as Fig. 3.

Figure 4

Precision (positive predictive value) for detecting cell-type-specific

association in simulated data for DNA methylation. The figure format is same

as Fig. 3.

Figure 5

Sensitivity for detecting cell-type-specific association in simulated data for
gene expression. Panels are aligned in rows according to the simulation
settings with the gene expression fold change of 1.7, 3.0 or 5.0. In each row,
panels for different cell types are aligned in decreasing order of proportion.
The vertical axis indicates sensitivity. In each panel, results from different
algorithms are aligned horizontally in different colors. Results from 50
simulation trials are summarized in a box plot. The middle bar of the box plot
indicates the median, and the lower and upper hinges correspond to the first
and third quartiles. The whiskers extend to the value no further than 1.5 $\times$
inter-quartile range from the hinges. FC, fold change; Gran, granulocytes;
Mono, monocytes.

Figure 6

Specificity for detecting cell-type-specific association in simulated data for
gene expression. The figure format is same as Fig. 5.

Figure 7

Precision (positive predictive value) for detecting cell-type-specific
association in simulated data for gene expression. The figure format is same
as Fig. 5.

Figure 8

Performance of the predictions for cell-type-specific association of DNA
methylation. For the association with rheumatoid arthritis in monocytes and
B cells and the association with age in CD4$^+$ T cells and monocytes, sensitivity
(top), specificity (middle) and precision (bottom) are plotted. In each panel,
results from different algorithms are aligned horizontally in different colors.
Precision is not plotted when there were no positive CpG sites. RA,
rheumatoid arthritis; Mono, monocytes.

Figure 9

Performance of the predictions for cell-type-specific association of gene

expression. For the association with age in CD4$^+$ T cells and monocytes,

sensitivity (top), specificity (middle) and precision (bottom) are plotted. In

each panel, results from different algorithms are aligned horizontally in

different colors. Precision is not plotted when there were no positive genes.

Mono, monocytes.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9