A regularized functional regression model enabling transcriptome-wide dosage-dependent association study of cancer drug response

Biomarker detection for revealing anticancer drug dynamics

Evanthia Koukouli[1*], Dennis Wang [2,3], Frank Dondelinger[4◗], Juhyun Park[1◗]

1 Department of Mathematics and Statistics, Fylde College, Lancaster University, Bailrigg, Lancaster, UK

2 Sheffield Institute for Translational Neuroscience, University of Sheffield, Sheffield, UK

3 Department of Computer Science, University of Sheffield, Sheffield, UK

4 Centre for Health Informatics and Statistics, Lancaster Medical School, Lancaster University, Bailrigg, Lancaster, UK

◗These authors contributed equally to this work.

*e.koukouli@lancaster.ac.uk (EK)

Abstract

Cancer treatments can be highly toxic and frequently only a subset of the patient population will benefit from a given treatment. Tumour genetic makeup plays an important role in cancer drug sensitivity. We suspect that gene expression markers could be used as a decision aid for treatment selection or dosage tuning. Using *in vitro* cancer cell line dose-response and gene expression data from the Genomics of Drug Sensitivity in Cancer (GDSC) project, we build a dose-varying regression model. Unlike existing approaches, this allows us to estimate dosage-dependent associations with gene expression. We include the transcriptomic profiles as dose-invariant covariates into the regression model and assume that their effect varies smoothly over the dosage levels. A two-stage variable selection algorithm (variable screening followed by penalised regression) is used to identify genetic factors that are associated with drug response over

the varying dosages. We evaluate the effectiveness of our method using simulation studies focusing on the choice of tuning parameters and cross-validation for predictive accuracy assessment. We further apply the model to data from five *BRAF* targeted compounds applied to different cancer cell lines under different dosage levels. We highlight the dosage-dependent dynamics of the associations between the selected genes and drug response, and we perform pathway enrichment analysis to show that the selected genes play an important role in pathways related to tumourgenesis and DNA damage response.

Author Summary

Tumour cell lines allow scientists to test anticancer drugs in a laboratory environment. Cells are exposed to the drug in increasing concentrations, and the drug response, or amount of surviving cells, is measured. Generally, drug response is summarized via a single number such as the concentration at which 50% of the cells have died (IC50). To avoid relying on such summary measures, we adopted a functional regression approach that takes the dose-response curves as inputs, and uses them to find biomarkers of drug response. One major advantage of our approach is that it describes how the effect of a biomarker on the drug response changes with the drug dosage. This is useful for determining optimal treatment dosages and predicting drug response curves for unseen drug-cell line combinations. Our method scales to large numbers of biomarkers by using regularisation and, in contrast with existing literature, selects the most informative genes by accounting for drug response at untested dosages. We demonstrate its value using data from the Genomics of Drug Sensitivity in Cancer project to identify genes whose expression is associated with drug response. We show that the selected genes recapitulate prior biological knowledge, and belong to known cancer pathways.

Introduction

Cancer is a heterogeneous disease, with individual tumours showing sometimes very different mutational and molecular profiles. The genetic makeup of a tumour influences how it reacts to a given anti-cancer drug. However, due to lack of predictive markers of

tumour response, often patients with very different tumour genetic makeup will receive   5
the same therapy, resulting in high rates of treatment failure [1]. Large clinical trials in   6
rapidly lethal diseases are expensive, complex and often lead to failure due to lack of   7
efficacy at a given dosage [2]. One major issue for some cancer treatments, e.g.   8
chemotherapies, are cytotoxic effects that result in collateral damage of the healthy host   9
tissue [3]. Patient remission depends not only on the selection of the right drug but also   10
on the determination of the optimal dosage, especially when drugs with small   11
therapeutic range, high toxicity levels or both are administered. Genetic factors can   12
help fine-tune the dosage for individual patients, so that the minimal effective dosage   13
can be delivered [4].   14

Treatment response in patients with specific cancers had been intensely examined in   15
relation to the molecular characteristics of the tumours [5]. However, cellular   16
heterogeneity within the tumour and the lack of standard metrics for quantifying drug   17
response in patients can make it difficult to computationally model response as a   18
function of molecular features. Cancer cell line drug screens can provide valuable   19
information about the effect of genetic features on drug dose-response in a controlled   20
setting. During the last decade, there have been several systematic studies that   21
examined the relationship between genetic variants and drug response in cell   22
lines [6–10]. There have also been studies that measured transcriptional profiles [11,12]   23
and drug response in cancer cells after administering anticancer drugs at various   24
dosages [13,14]. By comparing multiple genomic features of cell lines to drug response,   25
the investigators were able to identify gene signatures for drug responsiveness in specific   26
cancer types. However, these signatures were selected based on a single summary   27
statistic of response, usually IC50, that may not always be the most useful metric for   28
differentiating drugs [15], and only provides information on one dose concentration.   29
While these existing signatures of drug response provide a way towards selecting the   30
right drug for a patient, none of them characterise gene-dose relationships that may   31
ultimately identify the optimal dose for a drug to use in the clinic.   32

With regards to the high-dimensional nature of genomic datasets, it is worth noting   33
that highly-complex data sets with non-stationary trends are not easily amenable to   34
analysis by classic parametric or semi-parametric mixed models. Such effects, e.g. the   35
effect of genes on drug response over different drug dosages (dose-varying effect), can be   36

examined using varying coefficient models which allow for the covariate effect to be ₃₇ varying instead of constant [16]. Methods to estimate varying covariate effects include ₃₈ global and local smoothing, e.g. kernel estimators [17,18], basis approximation [19] or ₃₉ penalised splines [20]. Although non-parametric techniques can reduce modelling ₄₀ biases [21], they often suffer from the "curse of dimensionality" [22]. Inference in these ₄₁ models becomes impossible as the number of predictors increases, and often selecting a ₄₂ smaller number of important variables for inclusion into the model is clinically ₄₃ beneficial. Sparse regression has enabled a more flexible and computationally ₄₄ "inexpensive" way of choosing the best subset of predictors [23]. However, these ₄₅ methods cannot handle ultra-high dimensional problems without losing statistical ₄₆ accuracy and algorithmic stability, since they handle all of the predictors jointly. ₄₇ Consequently, there is a need of prior univariate tests focused on filtering out the ₄₈ unimportant predictors by estimating the association of each predictor to the outcome ₄₉ variable separately [21, 24, 25]. The advantage of using varying coefficient models along ₅₀ with a variable screening algorithm on genomic data sets was first introduced to explore ₅₁ the effect of genetic mutations on lung function [24]. Recently, Tansey et al. [26] ₅₂ proposed a method for modelling drug-response curves via Gaussian processes and ₅₃ linking them to biomarkers using a neural network prediction model. The authors did ₅₄ not use their model for dosage-dependent inference of biomarker effects, and the highly ₅₅ non-linear neural network model makes interpretation of biomarker effects challenging. ₅₆

Here, we extended the methodology of Chu et al. [24] to the objective of assessing ₅₇ the transcriptomic effect on anti-cancer drug response, where our coefficient functions ₅₈ were allowed to vary with dosage. We developed a functional regression framework to ₅₉ study the effectiveness of multiple anticancer agents applied in different cancer cell lines ₆₀ under different dosage levels, adjusting for the transcriptomic profiles of the cell lines ₆₁ under treatment. We considered a dose-varying coefficient model, along with a ₆₂ two-stage variable selection method in order to detect and evaluate drug-gene ₆₃ relationships, and then applied this method to data extracted from the Genomics for ₆₄ Drug Sensitivity in Cancer (GDSC) project [7]. To compare and differentiate similar ₆₅ treatments, we examined a case study of five BRAF targeted compounds under different ₆₆ dosages to almost 1000 cancer cell lines. We used baseline gene expression ₆₇ measurements for the cancer cell lines to investigate gene-drug response relationships for ₆₈

almost 18000 genes. Gene rankings were obtained based on the estimated effects of the   69
genes on the drug response. The resulting model describes the whole dose-response   70
curve, rather than a summary statistic of drug response (e.g. IC50), which allowed us to   71
identify trends in the gene-drug association at untested dose concentrations.   72

Materials and Methods   73

The Genomic of Drug Sensitivity in Cancer data   74

Drug sensitivity data and molecular measures derived from 951 cancer cell lines used for   75
the screening of 138 anticancer compounds were downloaded from the GDSC database   76
(https://www.cancerrxgene.org/). We specifically focused on cell lines of cancers of   77
epithelial, mesenchymal and haematopoietic origin treated by five BRAF targeted   78
inhibitors (PLX-4720, Dabrafenib, HG6-64-1, SB590885 and AZ628; GDSC1 data). The   79
maximum screening concentration for each different drug was: 10.00 uM for PLX-4720   80
and Dabrafenib, 5.12 uM for HG6-64-1, 5.00 uM for SB590885 and 4.00 uM for AZ628.   81
Additionally, we used the independently generated GDSC2 dataset to validate our   82
approach on drugs targeting *MEK1*, *MEK2* genes (Trametinib−1.00 uM;   83
Selumetinib−10.00 uM, and; PD0325901−0.250 uM) and the PI3K/MTOR signalling   84
pathway (Alpelisib−10.00 uM; AMG-319−10.00 uM, and; AZD8186−10.00 uM). The   85
drug sensitivity measurement was obtained via fluorescence-based cell viability assays   86
72 hours after drug administration [7]. Approximately 66% of drug sensitivity responses   87
were measured over nine dose concentrations (2-fold dilutions) and 34% were measured   88
over five drug concentrations (4-fold dilutions). In total, we considered 3805 cancer cell   89
line-drug combinations (experimental units). The distribution of different tissues of   90
origin treated were similar across the different drugs tested (for additional information   91
see Fig S2). Paired microarray gene expression data (17737 genes) was available   92
together with the drug response dataset   93
(https://www.cancerrxgene.org/gdsc1000/GDSC1000_WebResources/Home.html).   94

The dose-response dataset also included a blank response for wells on the   95
experimental plate that had not been seeded with cells or treated with a drug. Blank   96
responses have been used to adjust for the magnitude of the observation error while   97

measuring the amount of cells in each well. We used an affine transformation to the reported responses in order to normalise them within the drug concentration interval, 0 (0% of the maximum dosage) to 1 (100% of the maximum dosage). In particular, for the normalising procedure, we have used the formula:

$$NR_{ij} = \frac{R_{ij} - BR_i}{CR_i - BR_i} \tag{1}$$

where R_{ij} is the response of the i^{th} subject at the j^{th} dosage level, CR_i is the response under no drug administration (zero dose, $n_i = 1$), BR_i is the blank response of the i^{th} subject as described above and NR_{ij} is the new score taken from the transformation, $i = 1, \ldots, 3805, j = 1, \ldots, n_i$.

A two-stage algorithm for identification of gene-drug associations

Non-parametric techniques have been a great tool for reducing modelling bias and producing data driven inference. Flexible modelling techniques applied on high-dimensional genomic data sets can cause real problems in inference making. Sparse regression techniques, such as the LASSO, can be used as dimensionality reduction techniques, but cannot handle ultra-high dimensional problems without introducing statistical inaccuracies, algorithmic instability and a huge computational burden. Hence, the need for a feature screening algorithm which will marginally filter unimportant variables becomes essential [23]. Below we further explain the two-stage algorithm that has been built in order to detect and explore dose-dependent gene-response associations.

Let the repeated measures data $\{(d_{ij}, y_{ij}, \boldsymbol{z}_i, \boldsymbol{x}_i) : j = 1, \ldots, n_i, \ i = 1, \ldots, n\}$, where y_{ij} is the response of the ith experimental unit (corresponds to a drug sensitivity assay of a specific drug on a specific cell line) at the jth drug dosage level d_{ij} and $\boldsymbol{z}_i$ along with $\boldsymbol{x}_i$ are the corresponding vectors of scalar (dose-invariant) covariates. The covariate vector $\boldsymbol{z}_i = (1, z_{i1}, \ldots, z_{ip})^T$ is a low-dimensional vector of predictors that should be included in the model, whereas $\boldsymbol{x}_i = (x_{i1}, x_{i2}, \ldots, z_{iG})^T$ is a high-dimensional vector, i.e. 17737 gene expression measurements, that needs to be screened. We assumed that only a small number of x-variables (in our case, genes) are truly associated with the response while most of them are expected to be irrelevant; i.e. we

make a sparsity assumption.

To explore potential dose-varying effects between the covariates and the drug response, we consider the following varying coefficient model:

$$y_{ij} = \sum_{k=0}^{p} z_{ik} \beta_k(d_{ij}) + \sum_{g=1}^{G} x_{ig} \gamma_g(d_{ij}) + \varepsilon_{ij} \tag{2}$$

where $\{\beta_k(\cdot), k = 0, \ldots, p\}$ and $\{\gamma_g(\cdot), g = 1, \ldots, G\}$ are smooth functions of dosage level $d \in \mathscr{D}$, where $\mathscr{D}$ is a closed and bounded interval of $\mathbb{R}$. The errors ε_{ij} were assumed to be independent across subjects and potentially dependent within the same subject with conditional mean equal to zero and variance $\mathrm{Var}(\varepsilon) = \sigma^2(d) = V(d)$.

Methods for estimating the coefficient functions in Eq (2) include local and global smoothing methods, such as kernel smoothing, local polynomial smoothing, basis approximation smoothing etc. For computational convenience, in this application we used basis approximation smoothing via B-splines.

Let the sets of basis functions $\{B_{lk}(\cdot) : l = 1, \ldots, L_k\}$ and $\{B'_{lg}(\cdot) : l = 1, \ldots, L_g\}$ and constants $\{\zeta_{lk} : l = 1, \ldots, L_k\}$ and $\{\eta_{lg} : l = 1, \ldots, L_g\}$ where $k = 0, \ldots, p$ and $g = 1, \ldots, G$ such that, $\forall d \in \mathscr{D}$, $\beta_k(d)$ and $\gamma_g(d)$ can be approximated by the expansion

$$\beta_k(\cdot) \approx \sum_{l=1}^{L_k} \zeta_{lk} B_{lk}(\cdot) \quad \text{for} \quad k = 0, \ldots, p \tag{3}$$

$$\gamma_g(\cdot) \approx \sum_{l=1}^{L_g} \eta_{lg} B'_{lg}(\cdot) \quad \text{for} \quad g = 1, \ldots, G. \tag{4}$$

Substituting $\beta_k(\cdot)$ and $\gamma_g(\cdot)$ of Eq (2) with Eq (3) and Eq (4), we approximated Eq (2) by

$$y_{ij} \approx \sum_{k=0}^{p} z_{ik} \sum_{l=1}^{L_k} \zeta_{lk} B_{lk}(d_{ij}) + \sum_{g=1}^{G} x_{ig} \sum_{l=1}^{L_g} \eta_{lg} B'_{lg}(d_{ij}) + \varepsilon_{ij} \tag{5}$$

If $B_k(\cdot)$ and $B'_g(\cdot)$ are groups of B-spline basis functions of degree q_k and q_g respectively, and $\delta_0 < \delta_1 < \ldots < \delta_{K_k} < \delta_{K_k+1}$ and $\delta_0 < \delta_1 < \ldots < \delta_{K_g} < \delta_{K_g+1}$ are the corresponding knots, then $L_k = K_k + q_k$ and $L_g = K_g + q_g$.

Using the approximation Eq (5), the coefficients $\boldsymbol{\zeta} = (\zeta_0, \zeta_1, \ldots, \zeta_p)^T$ and

$\boldsymbol{\eta} = (\eta_1, \eta_2, \ldots, \eta_G)^T$ can be estimated by minimizing the squared error

$$\ell_w((\boldsymbol{\zeta}, \boldsymbol{\eta})^T) = \sum_{i=1}^{n} \sum_{j=1}^{n_i} w_{ij} \left[y_{ij} - \sum_{k=0}^{p} z_{ik} \sum_{l=1}^{K_k} \zeta_{lk} B_{lk}(d_{ij}) - \sum_{g=1}^{G} x_{ig} \sum_{l=1}^{L_g} \eta_{lg} B'_{lg}(d_{ij}) \right] \qquad (6)$$

where w_{ij} are known non-negative weights.

In cases where $p + G >> n$ though, minimisation of Eq (6) is infeasible. Our aim was to identify factors of the covariate vector $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_G)^T$ (genes) that are truly associated with the response (cancer cell line sensitivity to the drug). In addition, we wanted to explore potential dose-varying effects on the drug response.

We make the following sparsity assumption: any valid solution $\hat{\boldsymbol{\gamma}}(d)$ will have $\hat{\gamma}_g(d) = 0, \forall d \in \mathscr{D}$ for the majority of components g. To detect non-zero coefficient functions we applied a two-stage approach which incorporated a variable screening step and a further variable selection step.

Screening

The sparsity assumption applies only to components of $\mathbf{x}$, the high-dimensional covariate vector in Eq (2).

Let the set of indices

$$\mathcal{M}_0 = \{1 \le g \le G : ||\gamma_g(\cdot)||_2 > 0\} \qquad (7)$$

where $|| \cdot ||_2$ is the L_2-norm. In order to rank the different components of $\boldsymbol{x}$, we fitted the marginal non-parametric regression model for the gth x-predictor:

$$y_{ij} \approx \sum_{k=0}^{p} z_{ik} \sum_{l=1}^{K_k} \zeta_{lk}^{(g)} B_{lk}^{(g)}(d_{ij}) + x_{ig} \sum_{l=1}^{L_g} \eta_{lg}^{(g)} B_{lg}^{(g)\prime}(d_{ij}) + \varepsilon_{ij}^{(g)} \qquad (8)$$

where: $\{B_{lk}^{(g)}(\cdot) : l = 1, \ldots, L_k\}$ and $\{B_{lg}^{(g)\prime}(\cdot) : l = 1, \ldots, L_g\}$ are sets of coefficient functions; $\{\zeta_{lk}^{(g)} : l = 1, \ldots, L_k\}$ and $\{\eta_{lg}^{(g)} : l = 1, \ldots, L_g\}$ are constants to be estimated, $k = 0, \ldots, p$; and, $\varepsilon^{(g)}$ is the error term similar to Eq (5). We then computed the

following weighted mean squared error for each $g \in \{1, \ldots, G\}$,

$$\hat{u}_g = \frac{1}{n} \sum_{i=1}^{n} (\mathbf{y}_i - \hat{\mathbf{y}}_i^{(g)})^T \mathbf{W}_i (\mathbf{y}_i - \hat{\mathbf{y}}_i^{(g)}) \tag{9}$$

to quantify the importance of the gth x-variable. Here,

$$\boldsymbol{W}_i = \frac{1}{n_i} \hat{\boldsymbol{V}}_i^{-\frac{1}{2}} \boldsymbol{R}_i^{-1}(\hat{\boldsymbol{\phi}}) \hat{\boldsymbol{V}}_i^{-\frac{1}{2}} \tag{10}$$

where $\hat{\boldsymbol{V}}_i$ is the $n_i \times n_i$ diagonal matrix consisting of the dose-varying variance

$$\hat{\boldsymbol{V}}_i = \begin{bmatrix} \hat{V}(d_{i1}) & 0 & \ldots & 0 \\ 0 & \hat{V}(d_{i2}) & \ldots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \ldots & \hat{V}(d_{in_i}) \end{bmatrix} \tag{11}$$

and $\boldsymbol{R}_i(\boldsymbol{\phi}) = (R_{jk})$ the $n_i \times n_i$ working correlation matrix for the i^{th} subject. By $\boldsymbol{\phi}$, we denoted the $s \times 1$ vector that fully characterises the correlation structure. The estimate of $\boldsymbol{\phi}$, $\hat{\boldsymbol{\phi}}$, was obtained by taking the moment estimators for the parameters $\boldsymbol{\phi}$ in the correlation structure based on the residuals obtained from fitting the following model

$$y_{ij} = \sum_{k=0}^{p} z_{ik} \beta_k(d_{ij}) + \varepsilon_{ij} \quad \text{where} \quad i = 1, \ldots, n, j = 1, \ldots, n_i. \tag{12}$$

The variance function $V(d)$ in Eq (11) was estimated using techniques described in [24].

After having obtained $\{\hat{u}_g : g = 1, \ldots, G\}$, we sorted gene utilities in an increasing order, where smaller $\hat{u}_g$ values indicate stronger marginal associations. The x-predictors included in the screened submodel are, then, given by

$$\widehat{\mathscr{M}_{\tau_n}} = \{1 \leq g \leq G : \hat{u}_g \text{ ranks among the first } \tau_n(\nu)\} \tag{13}$$

where $\tau_n(\nu)$ corresponds to the size of the submodel which is chosen to be smaller than the sample size n.

Variable selection using a group SCAD (gSCAD) penalty

Screening algorithms aim to discard all unimportant variables but tend to be
conservative. In order to preserve only the most important x-predictors in the final
model, we considered a model including the first $\tau_n(\nu)$ outranked genes and we applied
a gSCAD penalty by minimising the following criterion:

$$\frac{1}{2}\sum_{i=1}^{n}\sum_{j=1}^{n_i} w_{ij}\Bigg\{ y_{ij} - \sum_{k=0}^{p} z_{ik}\sum_{l=1}^{L_k}\zeta_{lk}B_{lk}(d_{ij}) - \tag{14}$$

$$\sum_{g\in\widehat{\mathscr{M}}_{\tau_n}} x_{ig}\sum_{l=1}^{L_g}\eta_{lg}B'_{lg}(d_{ij})\Bigg\}^2 + \sum_{g\in\widehat{\mathscr{M}}_{\tau_n}} p_{\lambda,\alpha}(||\boldsymbol{\eta}_g||) \tag{15}$$

where

$$p_{\lambda,\alpha}(u) = \begin{cases} \lambda u & \text{if } 0 \le u \le \lambda \\ -\frac{(u^2 - 2\alpha\lambda u + \lambda^2)}{2(\alpha-1)} & \text{if } \lambda \le u \le \alpha\lambda \\ \frac{(\alpha+1)\lambda^2}{2} & \text{if } u \ge \alpha\lambda, \end{cases}$$

α is a scale parameter, λ controls for the penalty size and $||\cdot||$ is the Euclidean
L_2-norm. At this point, note that grouping is applied for the coefficients $\boldsymbol{\eta}_g$ that
correspond to the same coefficient function. In addition, in order to reduce the bias
introduced when applying a LASSO penalty, we alternatively chose the SCAD, which
coincides with the LASSO until $u = \lambda$, then transits to a quadratic function until
$u = \alpha\lambda$ and then it remains constant $\forall u > \alpha\lambda$, meaning that retains the penalisation
and bias rates of the LASSO for small coefficients but at the same time relaxes the rate
of penalisation as the absolute value of the coefficients increases. In Fig 1 the reader can
find a brief overview of the employed methodology.

Tuning parameters selection

We used knots placed at the median of the observed data values along with cubic
B-splines with 1 interior knot, resulting from calculating the number of interior knots
suitable using the formula $N_n = [n^{\frac{1}{2p+3}}]$ proposed and applied by [19, 27, 28]. Due to the

Fig 1. The two-stage algorithm for identifying dose-dependent associations between genes and drugs. Gene expression and drug response data from a drug screening study (e.g. GDSC) are used to fit our dose-varying coefficients model to estimate the dose-varying effect between covariates and drug response. A two-stage variable screening and selection algorithm is applied to rank gene-drug associations. The selected genes can then be used to predict dose-dependent response for drugs of interest.

computational burden this would add, we did not apply cross-validation.   198

    As for the screening threshold τ_n, its magnitude could be determined by the fraction   199

$\nu[\frac{n}{log(n)}]$, $\nu \in \{1, 2, 3, \ldots\}$. We conducted a pilot simulation study in order to decide the   200

most appropriate size (for further details see Text S1). We also considered an   201

automated algorithm for its selection (Greedy Iterative Non-parametric Independence   202

Screening-Greedy INIS, [29]). Finally, the penalty size for the gSCAD step λ was   203

determined using a 5-fold cross-validation.   204

Simulation study

Monte Carlo simulations were conducted to examine the ability of our model to detect the genes that are truly associated with the drug response. This had a key role in tuning model parameters and simultaneously assess model goodness-of-fit using a fraction of the original data set in order to reduce the computational burden of conducting a simulation study under the original dimensions of the data. Responses over different dosage levels were generated based on a subset of genes, the corresponding low-dimensional GDSC data covariates (drug and cancer type) and some specified smooth coefficient functions (see Text S1). Due to the computational burden associated with a simulation of the same scale as the data set, we conducted a simulation study using smaller random fragments of the original GDSC data set. In particular, we repeatedly sampled without replacement 190 experimental units and 886 genes based on which the simulated responses have been generated. The performance of the employed methodology has been assessed based on 1000 simulations using three screening thresholds ($\tau_n(\nu) = [\frac{n}{log(n)}]$, $\tau_n(\nu) = [\frac{2n}{log(n)}]$ and $\tau_n(\nu)$ chosen using the greedy-INIS algorithm [29]) and two estimated covariance structure scenarios (independence and rational quadratic covariance structure). Cubic B-splines and knots placed at the median of the observed data values have been used for estimating the coefficient functions.

To evaluate the performance of the proposed procedure we used the following summary measures: TP−number of genes correctly identified as active; FP−number of the genes incorrectly identified as active; TN−number of the genes correctly identified as inactive; FN−number of the genes incorrectly identified as inactive.

Simulation results suggested that our method accurately detects the drug associated genes from the simulated responses under most of the examined scenarios (Fig A in Text S1). A screening threshold of size $[\frac{2n}{log(n)}]$ and regression weights adjusted for the covariance structure of the data have been identified as the scenario where our method reached its maximum accuracy. Consequently, for the GDSC application, we chose the screening threshold to be the maximum possible, i.e. 923 genes derived from the formula $[\frac{2n}{log(n)}]$, and weights derived by assuming a rational quadratic covariance structure for the repeated measures.

Data and software availability

The analysis has been conducted using R version 3.6.3. Code for applying the two-stage variable selection algorithm is available online as an R package at `https://github.com/koukoulEv/fbioSelect`. The data is available online at the Genomics of Drug Sensitivity in Cancer website `https://www.cancerrxgene.org/`.

Results and Discussion

Dose-dependent associations with gene expression in a large-scale drug sensitivity assay

We applied the two stage variable selection algorithm under the dose-varying coefficient model framework described above. Gene rankings and predicted mean drug effects over different dosage levels were obtained. Our algorithm identified 230 candidate genes associated with drug response. The effect of each of those genes was assessed with respect to:

1. the area under the estimated coefficient curve (AUC) and its corresponding standard deviation (estimated using bootstrapping);

2. the effect on cell survival (overall positive, overall negative, mixed);

3. Spearman correlation between the coefficient function value and the dosage level;

4. the mean fold change of the expression of cell lines carrying *BRAF* mutations with respect to wild type; and,

5. the protein-protein interaction network distance between the *BRAF* gene and the selected genes using the Omnipath database [30].

The 230 genes were ranked based on the estimated AUC value (Table S4), and the top 30 genes were highlighted for further analysis (Table 1). The higher the AUC, the larger the effect of the gene on the drug response. The overall effect on cell survival can be either positive, negative or vary over the different dosage levels as determined by the range of the estimated coefficient function. Spearman's rank correlation was used as an indicator of the coefficient function's monotonicity by characterising the progress of the

genetic effect over different dosage levels. For instance, high expression of the *C3orf58* gene at baseline has a positive effect on cancer cell survival, which becomes stronger as the dosage increases (Spearman's correlation=0.922). In other words, high expression of this gene can be an indicator of drug resistant cell lines. On the other hand, the *DLC1* gene has a decreasing (Spearman's correlation=-0.928) and negative effect on cancer cell survival which suggested that as the dosage increases, higher baseline expression of this gene can indicate higher drug sensitivity at higher dosage. Elevated expression of DLC1 has been observed in melanoma and is a well known tumour suppressor that could be a novel marker of BRAF inhibition [31]. Finally, in cases where the overall effect varies (changes between positive and negative), the effect of gene expression on the drug response depends on the drug dosage. In particular, the effect of *DLX6* increases and then decreases at higher dosages (Fig 2). Given the biological and technical variation in drug screens, we should treat the mean effect estimates with caution and consider the confidence intervals of the coefficient functions in order to derive conclusions about the exact effect of the selected genes on the dose response (Fig 2).

Coefficient function estimates provide a lot of information about the dosage, cancer type and genetic effects on drug response. Fig 2 illustrates the estimated coefficient functions for different drugs, cancer types and three genes in relation to the model's intercept, Dabrafenib response in *BRAF* mutant cell lines originating from the skin (melanoma). Except from HG6-64-1, all other BRAF inhibitors (AZ628, SB590885 and PLX4720) showed no addition effect compared to this intercept. Similar patterns can be observed for cancer cell lines coming from most of the tissues examined. This result indicates that the examined drugs may have similar or worse behaviour over the different dosages for most of the examined cancer types. Interestingly, we observed greater efficacy (negative values of the coefficient function) for cell lines originating from the endocrine system, autonomic ganglia and heamatopoietic and lymphoid tissues at lower dosages. The observed effect in endocrine system cell lines reflects the Dabrafenib responses observed in anaplastic thyroid cancer patients [32]. Interestingly, the drug, Trametinib, taken in combination with Dabrafenib is a MEK inhibitor, and genes interacting with MEK (*MAP2K1*) were selected features from our model (Fig 3). Together these results provide important insights into the effectiveness of the five *BRAF* targeted drugs examined on different cancer types, highlighting the potential for

Table 1. Top 30 gene rankings based on the estimated area under the coefficient function curve.

Gene Name	Area	SD	Sign	Spearman's Correlation	Mean fold change in *BRAF* mutant vs wild-type cell lines	Protein-protein interaction network distance to *BRAF*
KIR3DL1	0.370	0.107	-	-0.874	0.978	3
CHST11	0.257	0.092	-	-0.817	0.899	NI
APOC1P1	0.247	0.09	-	-0.918	1.190	NI
PLEKHA6	0.239	0.086	-	-0.908	1.037	3
PPM1F	0.223	0.068	+	0.910	0.883	3
BFSP1	0.222	0.074	-	-0.800	1.217	NI
PPP1R3A	0.217	0.082	+	0.774	1.078	3
C16orf87	0.207	0.087	+	0.851	0.977	NI
PARVA	0.203	0.081	+	0.890	0.984	2
SLC39A13	0.202	0.079	-	-0.461	1.055	NI
UCN2	0.198	0.07	-	-0.928	0.979	NI
STMN3	0.198	0.087	+	0.834	1.201	2
RNF130	0.197	0.083	-	-0.927	1.153	NI
C3orf58	0.196	0.076	+	0.922	1.133	NI
CXXC4	0.188	0.079	+	0.866	0.995	NI
THBD	0.179	0.093	0	-0.967	1.231	4
SIRT3	0.173	0.066	-	-0.760	1.013	3
PLAT	0.172	0.092	-	-0.878	1.322	4
MPPED1	0.168	0.066	+	0.430	0.978	NI
INSL3	0.162	0.068	-	-0.973	0.965	NI
FAM163A	0.159	0.078	-	-0.983	1.106	NI
CNIH3	0.153	0.08	-	-0.918	0.938	NI
GJA3	0.153	0.067	0	-0.940	0.933	NI
BTG2	0.152	0.078	+	0.959	1.035	2
DLX6	0.152	0.059	0	0.686	0.987	NI
DLC1	0.151	0.053	-	-0.928	0.974	3
GAPDHS	0.150	0.077	+	0.886	1.232	NI
JAG2	0.149	0.069	-	-0.994	0.981	3
SMOX	0.146	0.057	0	0.816	1.070	NI
ZMYND8	0.145	0.091	+	0.907	1.020	3

Gene rankings of the top 30 selected genes based on the magnitude of the genetic effect on drug response. A positive (+) sign translates to a positive effect on cells survival after drug administration, a negative (-) sign translates to a negative effect on cells survival and a mixed (0) effect translates to a varying effect on cells survival which depends on drug dosage. Spearman's correlation is calculated between drug dosage and gene estimated coefficient function values as an indicator of the magnitude change of the gene effect over the increasing dosage. Area corresponds to the area under the estimated coefficient curve and the SD corresponds to the standard deviation of the area based on bootstrapping. Mean fold change is calculated between the selected gene expression values of the cell lines carrying *BRAF* mutations with respect to wild type. Protein-protein interaction network distance is computed based on the shortest interaction path between the *BRAF* gene and each of the selected genes. Here, NI denotes absence of any interaction.

effective treatment of a wide range of cancers given the tumour genetic characteristics.

Since the *BRAF* gene is the target of the drugs, mean fold change and protein-protein interaction network distance were used to examine whether and how the selected genes are related to inhibitors' target. From the selected genes, 120 genes had a mean fold change greater than 1 whereas the rest had a mean fold change between 1

Fig 2. Estimated coefficient functions for the low-dimensional predictors and three of the selected genes. Estimated coefficient functions for intercept, different drugs, tissue of origin and three of the selected genes along with 95% bootstrap confidence intervals. Baseline corresponds to *BRAF* mutant cell lines treated with Dabrafenib in skin tumours.

and 0.792. Some of the genes with the highest mean fold change of *BRAF* mutation were *PSMC3IP*, *KIF3C*, *UBE2Q2*, *SERPIND1* and *PLAT*, however only *PLAT* is displayed in Table 1. From the genes identified through the two-stage algorithm, 35% of them encode proteins interacting with the *BRAF* gene, though none of them directly. Most of the selected genes interact with the *BRAF* gene via pathways mediated by

Fig 3. Protein-protein interaction network for the genes selected from the two-stage variable selection algorithm. (A) Undirected protein-protein interaction network between the 230 selected (blue) and the *BRAF* (red) genes (full scale analysis). (B) Undirected protein-protein interaction network between the 65 genes selected from the two-stage variable selection algorithm for the cell lines resistant to BRAF inhibitors (blue) and the *BRAF* (red) gene. In both panels genes depicted with black are the interaction mediators. Common mediators include the *HRAS*, *MAPK1*, *MAP2K1* and *BAD* genes.

HRAS, *MAPK1* (*ERK*), *MAP2K1* (*MEK*) and *BAD* (Fig 3). 305

Since *HRAS* mutations are frequent in patients receiving *BRAF* targeted 306

therapies [33], we examined the mean estimated trajectory over different dosages under 307

treatment with *BRAF* inhibitors tested in six cancer cell lines with and without *BRAF* 308

and *HRAS* mutations (Fig 4). As stated previously, we observed that in most cases 309

HG6-64-1 seems to be the most effective drug. The estimated coefficient functions 310

facilitate drug examination and response prediction under the different dosages. In 311

some instances, we observed different drugs having similar behaviour for lower drug 312

dosages and larger divergence for higher dosages. In most cases, regardless of the cell 313

line origin, our method successfully estimates the expected survival rates of the cancer 314

cell lines for the different drugs given their gene expression information. 315

For validation purposes, we performed the same analysis using the 316

independently-generated GDSC2 dataset, with a different set of drugs. Note that the 317

drug set in GDSC2 only partially overlaps with the one used in GDSC1. The results are 318

reported in supplementary Fig S11, and show similar properties to the analysis of the ₃₁₉
BRAF drugs in Fig 4. As the GDSC2 dose-responses are produced from ₃₂₀
independently-generated experiments, the measured drug response is different for some ₃₂₁
of the drug-cell line combinations. We observe some divergences between GDSC1 and ₃₂₂
GDCS2 estimated trajectories for Dabrafenib and PLX-4720, which can be explained ₃₂₃
due to measurement error in the GDSC experiments themselves. ₃₂₄

**Fig 4. Estimated mean drug response trajectories for six cancer cell lines
with *BRAF* and *HRAS* mutations.** Observed responses (points) and estimated
mean trajectory (lines) of cells' concentration for cancer cell lines with and without
BRAF and *HRAS* mutations after treatment with the five anticancer compounds
examined.

Variable selection algorithm identifies cancer pathways ₃₂₅
associated with BRAF inhibitor response ₃₂₆

Using our functional regression approach, we identified 230 genes that were selected via ₃₂₇
the SCAD step (observed gene set). We used the Enrichr [34, 35], WikiPathways [36] ₃₂₈
databases to see if the selected genes can be grouped into common functional classes or ₃₂₉

pathways. In total, 183 pathways identified, from which 11 were statistically significant ₃₃₀ at 5% level, including apoptosis modulation, NOTCH1 regulation, and MAPK signaling ₃₃₁ (Table S6). The model identified genes (*IKBKB*, *RASGRF1*, *DUSP16*, *DUSP8*, *DUSP6*, ₃₃₂ *MAPT* and *IL1R2*) downstream of the MAPK signaling pathway targeted by BRAF ₃₃₃ inhibitors. ₃₃₄

Previous studies of these pathways have found associations with tumourgenesis and ₃₃₅ cancer treatment [37–40]. Genes in more than one of these pathways include *IKBKB*, ₃₃₆ *PLAT*, *IL1R2* and *PDPK1*. The IKB kinase composed of IKBKB had previously been ₃₃₇ suggested as a marker of sensitivity for combination therapy with BRAF inhibitors [41]. ₃₃₈ Taken together, these results suggest that the identified associations between the drug ₃₃₉ response and the observed genes may reveal new predictive markers of tumour response ₃₄₀ to the examined BRAF inhibitors. ₃₄₁

In addition to the pathway enrichment analysis, we used the Molecular Signatures ₃₄₂ Database (MSigDB database v7.0 updated August 2019: [42]) to compute overlaps ₃₄₃ between the observed gene set and known oncogenic gene sets. Fig 5 displays the 29 ₃₄₄ overlaps found. Interestingly, we identified three instances where the observed gene set ₃₄₅ significantly overlapped with gene sets over-expressing an oncogenic form of the *KRAS* ₃₄₆ gene. ₃₄₇

We further explored potential biologically relevant pathways using the Reactome ₃₄₈ database [44, 45]. More than 40 enriched pathways were identified at a 5% significance ₃₄₉ level. The top 40 pathways are depicted in Fig 6 along with the pathway-gene network ₃₅₀ of the top 5 pathways (for the full list, see Table S12). Interestingly, axon guidance and ₃₅₁ VEGF signalling were among the enriched pathways, confirming relevance of the ₃₅₂ selected genes to the intended role of the examined compounds, since BRAF kinase ₃₅₃ activity drives axon growth in the central nervous system [46] and VEGF blockade has ₃₅₄ potential anti-tumour effects when combined with BRAF inhibitors [47]. Note that ₃₅₅ axon growth has not thus far been directly implicated in BRAF inhibitor activity, and ₃₅₆ thus our analysis provides important evidence towards this theory. ₃₅₇

Fig 5. Overlaps between the observed gene set and oncogenic signatures in the Molecular Signatures Database (full data analysis); signalling pathways enriched for genes predictive of *BRAF* inhibitor response (resistant cell lines). (A) Full gene set names can be found in Table S8. Overlaps have been detected using gene set enrichment analysis performed using a hypergeometric distribution. The false discovery rate analog of the hypergeometric p-value is displayed after correction for multiple hypothesis testing according to Benjamini and Hochberg [43]. (B) Top 20 enriched signalling pathways along with the adjusted p-values and the number of overlapping genes obtained after pathway enrichment analysis to the resistant cell line analysis results (for full list of the pathways identified see Table S7).

Identifying dose-dependent genes in drug-resistance conditions

Acquired resistance to BRAF inhibitors is often observed in the clinic [48]. To further examine the utility of the employed methodology, we applied the variable selection algorithm to a data subset containing only cell lines with mutations activating resistant mechanisms to BRAF inhibitors [49]. Out of the 951 cell lines in the data, 191 had some mutation in any of the following: *RAC1* gene, *NRAS* gene, *cnaPANCAN44* or *cnaPANCAN315*. We identified 65 genes associated with dose-response, though none of

Fig 6. Pathway enrichment analysis using the Reactome database. (A) Top 40 enriched signalling pathways along with the adjusted p-values. For the full list, see Table S12. (B) Pathway-gene network of the top 5 enriched signalling pathways as found using Reactome [44, 45].

them were directly associated with the MAPK/ERK pathway. However, from these, 25 365
genes have been found to indirectly interact with the *BRAF* gene (Fig 3) and 21 to 366
overlap with three oncogenic gene sets in the Molecular Signatures Database (genes 367
down-regulated in NCI-60 panel of cell lines with mutated *TP53*; genes up-regulated in 368
Sez-4 cells (T lymphocyte) that were first starved of *IL2* and then stimulated with *IL21*, 369
and; genes down-regulated in mouse fibroblasts over-expressing *E2F1* gene; Table S9). 370
Finally, we found 34 pathways enriched for genes predicting drug response of the 371
mutated cell lines to the examined BRAF inhibitors, of which the top 20 are depicted in 372
Fig 5(B). 373

Table 2 presents gene rankings based on the AUC and the overall coefficient function 374
effect (sign) for the 42 genes in either the enriched pathways, the three oncogenic gene 375

sets discussed above or the protein-protein interaction network with the *BRAF* gene $\qquad$ 376
(full list available in Table S5). Eight of the selected genes in the current $\qquad$ 377
implementation were also selected from the algorithm implemented on the full data: $\qquad$ 378
ASB9, *PRSS33*, *GJA3*, *PLAT*, *KLF9*, *BFSP1*, *MTARC1* and *UCN2*. $\qquad$ 379

Predictive performance of dose-dependent models $\qquad$ 380

As discussed above, the employed methodology gives a good overview of the baseline $\qquad$ 381
genetic effect on drug response. We assessed the overall predictive performance of our $\qquad$ 382
method using 10-fold cross validation under two different scenarios. For the first, we $\qquad$ 383
split the data into training and test set holding out the experimental units (cancer cell $\qquad$ 384
line-drug combinations) and for the second, holding out cancer cell lines. The absolute $\qquad$ 385
mean error for both cases was around 0.12. Our analysis shows robust cross-validated $\qquad$ 386
performance when it comes to predicting sensitivity to the administered drugs, as shown $\qquad$ 387
in Fig S3 which displays the correlation between predicted and true response. This $\qquad$ 388
result was further validated by repeating the analysis on the independently-generated $\qquad$ 389
GDSC2 dataset, using a different set of drugs (Fig S10), which demonstrated $\qquad$ 390
comparable predictive performance. $\qquad$ 391

Predictive accuracy for the dose-response curves was evaluated under four different $\qquad$ 392
sub-scenarios: prediction of the most effective drug-dosage combination for the 951 cell $\qquad$ 393
lines in the data set; prediction of the most effective drug given a cell line; prediction of $\qquad$ 394
the most effective dosage given treatment with a particular drug and prediction of the $\qquad$ 395
most effective dosage range given treatment with a drug (Table 3). The proposed model $\qquad$ 396
performs well when it comes to predicting the most effective drug or dosage range $\qquad$ 397
($\approx$79% in both scenarios). Results are less reliable when it comes to prediction of the $\qquad$ 398
exact dosage or drug-dosage combination ($\approx$48-49% and $\approx$57-58% in both scenarios) $\qquad$ 399
but this can be due to either the large variability observed in the observed responses or $\qquad$ 400
due to the small number of cell lines for some predictor level combinations. Results were $\qquad$ 401
similar for both cross-validation scenarios (differences range from 0 to $<$2%, Table 3), $\qquad$ 402
meaning that as long as a cell line has similar genetic characteristics to those observed, $\qquad$ 403
the model can be reliable in predicting the outcome after anticancer drug $\qquad$ 404
administration. $\qquad$ 405

Table 2. Rankings of the genes identified from the pathway and oncogenic gene set enrichment analysis.

Gene Name	Area	SD	Sign	Spearman's Correlation	Mean fold change in *BRAF* mutant vs wild-type cell lines	Protein-protein interaction network distance to BRAF
MYO5A	0.531	0.261	+	0.955	1.358	4
S100A1	0.488	0.189	+	0.812	1.263	NI
GPNMB	0.424	0.196	+	1	1.169	3
ACP5	0.359	0.149	-	-0.998	1.039	NI
FCGR2A	0.341	0.158	-	-0.588	1.25	3
CITED1	0.28	0.348	0	-0.603	1.63	3
SPRY4	0.274	0.127	-	-0.611	1.228	2
CD44	0.239	0.164	+	0.868	1.413	3
RAP2B	0.236	0.179	0	0.927	1.254	NI
KCNJ13	0.205	0.094	0	-0.604	1.101	3
ALX1	0.202	0.099	-	-1	1.104	NI
PLAT	0.201	0.121	-	-0.405	1.312	4
RETSAT	0.201	0.142	0	0.689	1.127	NI
GSN	0.196	0.109	+	0.588	1.079	4
CDH19	0.185	0.102	0	0.943	0.933	NI
ATP1B3	0.178	0.115	-	-1	1.063	NI
BAZ1A	0.173	0.105	+	-0.29	1.109	4
SLC16A4	0.166	0.117	-	-0.298	1.234	NI
ST6GALNAC2	0.164	0.102	0	-0.815	1.264	NI
MFSD12	0.16	0.148	0	-0.788	1.13	NI
GJA3	0.157	0.075	0	-0.85	1.071	NI
CYP27A1	0.156	0.09	-	-0.743	1.373	NI
EGLN1	0.15	0.119	-	-0.442	1.053	3
TRPV2	0.147	0.118	0	0.769	1.074	NI
MITF	0.146	0.106	+	1	0.743	2
TBC1D7	0.146	0.118	0	-0.603	1.304	NI
SLC6A8	0.144	0.111	0	-0.263	0.941	NI
PTPRZ1	0.139	0.138	-	-0.808	1.074	4
PLOD3	0.132	0.135	0	0.696	1.166	NI
ANKRD7	0.131	0.12	+	0.92	1.241	NI
KANK1	0.107	0.113	0	-0.493	1.345	NI
GYPC	0.105	0.092	+	-0.3	1.072	NI
TYR	0.1	0.098	-	0.467	1.11	4
TYRP1	0.1	0.097	0	0.457	1.326	3
IGSF8	0.09	0.129	0	-0.668	1.313	5
SPRED1	0.067	0.116	0	-0.556	1.239	4
ITGA9	0.056	0.111	0	0.785	1.154	4
KREMEN1	0.053	0.086	0	-0.555	1.123	4
LAMA4	0.038	0.083	-	0.344	1.151	4
MLANA	0.037	0.097	0	0.534	1.147	NI
KLF9	0.011	0.074	0	0.932	1.064	NI

Table notes rankings of the genes found to have some biological importance. A positive (+) sign translates to a positive effect on cells survival after drug administration, a negative (-) sign translates to a negative effect on cells survival and a neutral (0) effect translates to a varying effect on cells survival which depends on drug dosage. Spearman's correlation is calculated between drug dosage and gene estimated coefficient function values as an indicator of the magnitude change of the gene effect over the increasing dosage. Area corresponds to the area under the estimated coefficient curve and the SD corresponds to the standard deviation of the area based on bootstrapping. Pearson's correlation is calculated between the selected gene microarray expression values and the *BRAF* expression across all the cell lines. Protein-protein interaction network distance is computed based on the shortest interaction path between the *BRAF* gene and each of the selected genes. Here, NI denotes absence of interaction.

We additionally compared the performance of our two-stage algorithm approach to a penalized linear (LASSO) regression for predicting the IC50 and area under the dose-response curve (AUC). Note that our functional regression model is not directly predicting either of these values, but rather predicts the full drug-response curve. As this is a harder problem, we would expect the LASSO to have a natural advantage; however, our method has the added benefit of being able to detect dose-dependent associations, which is not possible when predicting summary statistics of the dose-response curve directly. We employed 10-fold cross-validation to evaluate the predictive error in terms of root mean squared error (RMSE), and we used a sigmoid curve fit for estimating the IC50 values from the predicted dose-response curves with our two-stage method. Our method outperformed standard LASSO in terms of predicting the AUC ($\text{RMSE}_{2-stage} = 0.176$; $\text{RMSE}_{LASSO} = 0.347$) and performed well on predicting the IC50, although the LASSO performed better ($\text{RMSE}_{2-stage} = 1.969$; $\text{RMSE}_{LASSO} = 1.134$). This could be expected, as estimating the IC50 from the predicted dose-response curve adds a further level of complexity, compared to directly predicting this value using the LASSO.

Table 3. Predictive performance of the employed model (mean absolute error=0.121).

Scenario	Accuracy EU	Accuracy CL
Model predicts the more effective drug-dosage combination	57.85%	57.42%
Model predicts the more effective drug given a cell-line	78.21%	78.21%
Model predicts the more effective dosage given a drug	48.44%	48.65%
Model predicts the most effective dosage range ($>$ or $\leq$ 31.25% of the maximum dosage)	79.47%	79.28%

Table notes the predictive performance of the model based on the percentages for correctly identifying the most effective drug, dosage or drug-dosage combinations. Results obtained based on 10-fold cross-validation of the final model (based on holding out either experimental units−EU− or cancer cell lines−CL−).

Conclusion

Genetic alternations and gene expression in tumours are known to affect disease progression and response to treatment. Here, we studied dosage-dependent associations between gene expression and drug response, using a functional regression approach which adjusts for genetic factors. We analysed data from the Genomics of Drug Sensitivity in Cancer project relating to drug effectiveness for suspending cancer cell proliferation under different dosages, and examined five *BRAF* targeted inhibitors, each applied in a number of common and rare types of cancer cell lines. Our implementation

of a two-stage screening algorithm revealed a number of genes that are potentially 430 associated with drug response. Gene, drug and cancer type trajectories have been 431 modelled using a varying coefficient modelling framework. The proposed methodology 432 allows for dose-dependent analysis of genetic associations with drug response data. It 433 enables us to study the effect of different drugs simultaneously, which results in high 434 accuracy of drug response prediction. Drug comparisons using the proposed 435 methodology could support drug repositioning, especially in disease indications where 436 existing treatment options are limited. In addition, our methodology can help to reveal 437 unknown potential relationships between genetic characteristics and drug efficacy. 438 Hence, the good predictive performance of our method could be due to the fact that 439 some genes may act as proxies for unmeasured phenotypes that are directly relevant to 440 drug sensitivity. 441

Our work relies on two major assumptions. First, that out of tens of thousands 442 genes regulating protein composition only a small proportion is actually associated with 443 cancer cell survival in a dosage-dependent manner. In other words, transcriptomic 444 profiles exert influence on disease progress after drug administration in a sparse and 445 dynamic way. However, if a large number of genes are associated with the drug response, 446 our method may produce biased results, and some important information about the 447 biological mechanisms can be lost. Secondly, we assume that the different drugs are 448 comparable on the scale of maximum dosage percentage level for our joint model. 449 However, we acknowledge that different drugs have different chemical structure and 450 maximum screening concentrations. Our focus is to identify genetic components that 451 could be informative for dose response given drugs that belong to a particular family, 452 for example $BRAF$ targeted therapies. However, our methodology is flexible enough to 453 allow each drug to be examined separately if it appears to be clinically appropriate. 454

Drug response prediction from gene expression data has been widely studied in the 455 literature. Sparse regression methods, gene selection algorithms such as the Ping-pong 456 algorithm [50], or a combination of network analysis and penalised regression, e.g. the 457 sparse network-regularized partial least squares method [51], have all been employed to 458 simultaneously predict drug response and select genetic factors that seem to be 459 associated with the drug response. However, none of these methods are able to quantify 460 the effect of drug dosage on the response, which is one of the main contributions of this 461

work. Employing the proposed dose-varying model gives a detailed picture of different drug effects and can be extremely valuable in predicting drug response for agents with small therapeutic range and high toxicity levels. Our algorithm showed moderate predictive performance due to the complexity of predicting whole drug-response curves. Methods for further enhancing the performance of the proposed methodology, such as judicious use of prior information and leveraging information sharing across multiple data sources should be explored in the future in order to overcome this issue and make good use of its full potentials.

To conclude, the main purpose of this paper is to examine the dose-dependent associations between genes and drugs. The proposed methodology, by using the raw data to infer the effects of interest, allows to obtain a more comprehensive picture of the biological mechanisms that undergo cancer treatment and the role of drug dose on that. In addition, due to its simple structure, it allows extension to different types of molecular data (e.g. RNA-seq gene expression, methylation or mutational profiles) and enrichment with further information, such as drug chemical composition.

References

1. Chang JC, Wooten EC, Tsimelzon A, Hilsenbeck SG, Gutierrez MC, Elledge R, et al. Gene expression profiling for the prediction of therapeutic response to docetaxel in patients with breast cancer. The Lancet. 2003;362(9381):362–369.

2. Cook D, Brown D, Alexander R, March R, Morgan P, Satterthwaite G, et al. Lessons learned from the fate of AstraZeneca's drug pipeline: a five-dimensional framework. Nature reviews Drug discovery. 2014;13(6):419–431.

3. Corrie PG. Cytotoxic chemotherapy: clinical aspects. Medicine. 2008;36(1):24–28.

4. Relling MV, Dervieux T. Pharmacogenetics and cancer therapy. Nature reviews cancer. 2001;1(2):99.

5. Zhang W, Liu Y, Sun N, Wang D, Boyd-Kirkup J, Dou X, et al. Integrating Genomic, Epigenomic, and Transcriptomic Features Reveals Modular Signatures Underlying Poor Prognosis in Ovarian Cancer. Cell reports. 2013;4(3):542–553.

6. Iorio F, Knijnenburg TA, Vis DJ, Bignell GR, Menden MP, Schubert M, et al. A landscape of pharmacogenomic interactions in cancer. Cell. 2016;166(3):740–754.

7. Yang W, Soares J, Greninger P, Edelman EJ, Lightfoot H, Forbes S, et al. Genomics of Drug Sensitivity in Cancer (GDSC): a resource for therapeutic biomarker discovery in cancer cells. Nucleic acids research. 2012;41(D1):D955–D961.

8. Hyman DM, Taylor BS, Baselga J. Implementing genome-driven oncology. Cell. 2017;168(4):584–599.

9. Ben-David U, Siranosian B, Ha G, Tang H, Oren Y, Hinohara K, et al. Genetic and transcriptional evolution alters cancer cell line drug response. Nature. 2018;560:325–330.

10. Tavassoly I, Hu Y, Zhao S, Mariottini C, Boran A, Chen Y, et al. Genomic signatures defining responsiveness to allopurinol and combination therapy for lung cancer identified by systems therapeutics analyses. Molecular Oncology. 2019;13(8):1725–1743.

11. Ji RR, de Silva H, Jin Y, Bruccoleri RE, Cao J, He A, et al. Transcriptional profiling of the dose response: a more powerful approach for characterizing drug activities. PLoS computational biology. 2009;5(9).

12. Delpuech O, Rooney C, Mooney L, Baker D, Shaw R, Dymond M, et al. Identification of Pharmacodynamic Transcript Biomarkers in Response to FGFR Inhibition by AZD4547. Molecular Cancer Therapeutics. 2016;15(11):2802–2813.

13. Falcetta F, Lupi M, Colombo V, Ubezio P. Dynamic rendering of the heterogeneous cell response to anticancer treatments. PLoS computational biology. 2013;9(10):e1003293.

14. Silverbush D, Grosskurth S, Wang D, Powell F, Gottgens B, Dry J, et al. Cell-specific computational modeling of the PIM pathway in acute myeloid leukemia. Cancer research. 2017;77(4):827–838.

15. Keshava N, Toh TS, Yuan H, Yang B, Menden MP, Wang D. Defining subpopulations of differential drug response to reveal novel target populations. npj Systems Biology and Applications. 2019;5:36.

16. Hastie T, Tibshirani R. Varying-coefficient models. Journal of the Royal Statistical Society: Series B (Methodological). 1993;55(4):757–779.

17. Wu CO, Chiang CT, Hoover DR. Asymptotic confidence regions for kernel smoothing of a varying-coefficient model with longitudinal data. Journal of the American statistical Association. 1998;93(444):1388–1402.

18. Wu CO, Chiang CT. Kernel smoothing on varying coefficient models with longitudinal dependent variable. Statistica Sinica. 2000; p. 433–456.

19. Huang JZ, Wu CO, Zhou L. Polynomial spline estimation and inference for varying coefficient models with longitudinal data. Statistica Sinica. 2004; p. 763–788.

20. Qu A, Li R. Quadratic inference functions for varying-coefficient models with longitudinal data. Biometrics. 2006;62(2):379–391.

21. Fan J, Feng Y, Song R. Nonparametric independence screening in sparse ultra-high-dimensional additive models. Journal of the American Statistical Association. 2011;106(494):544–557.

22. Geenens G, et al. Curse of dimensionality and related issues in nonparametric functional regression. Statistics Surveys. 2011;5:30–43.

23. Song R, Yi F, Zou H. On varying-coefficient independence screening for high-dimensional varying-coefficient models. Statistica Sinica. 2014;24(4):1735.

24. Chu W, Li R, Reimherr M. Feature screening for time-varying coefficient models with ultrahigh dimensional longitudinal data. The annals of applied statistics. 2016;10(2):596.

25. Fan J, Ma Y, Dai W. Nonparametric independence screening in sparse ultra-high-dimensional varying coefficient models. Journal of the American Statistical Association. 2014;109(507):1270–1284.

26. Tansey W, Li K, Zhang H, Linderman SW, Rabadan R, Blei DM, et al. 545
Dose-response modeling in high-throughput cancer drug screenings: An 546
end-to-end approach. arXiv preprint arXiv:181205691. 2018;. 547

27. Xue L, Qu A, Zhou J. Consistent model selection for marginal generalized 548
additive model for correlated data. Journal of the American Statistical 549
Association. 2010;105(492):1518–1530. 550

28. Xue L, Qu A. Variable selection in high-dimensional varying-coefficient models 551
with global optimality. Journal of Machine Learning Research. 552
2012;13(Jun):1973–1998. 553

29. Fan J, Feng Y, Song R. Nonparametric Independence Screening in Sparse 554
Ultra-High-Dimensional Additive Models. Journal of the American Statistical 555
Association. 2011;106(494):544–557. 556

30. Türei D, Korcsmáros T, Saez-Rodriguez J. OmniPath: guidelines and gateway for 557
literature-curated signaling pathway resources. Nature methods. 2016;13(12):966. 558

31. Yang X, Hu F, Liu JA, Yu S, Cheung MPL, Liu X, et al. Nuclear DLC1 exerts 559
oncogenic function through association with FOXK1 for cooperative activation of 560
MMP9 expression in melanoma. Oncogene. 2020;39(20):4061–4076. 561

32. Subbiah V, Kreitman RJ, Wainberg ZA, Cho JY, Schellens JH, Soria JC, et al. 562
Dabrafenib and trametinib treatment in patients with locally advanced or 563
metastatic BRAF V600–mutant anaplastic thyroid cancer. Journal of Clinical 564
Oncology. 2018;36(1):7. 565

33. Sharma SP. RAS mutations and the development of secondary tumours in 566
patients given BRAF inhibitors. The Lancet Oncology. 2012;13(3):e91. 567

34. Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, et al. Enrichr: 568
interactive and collaborative HTML5 gene list enrichment analysis tool. BMC 569
bioinformatics. 2013;14(1):128. 570

35. Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. 571
Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. 572
Nucleic acids research. 2016;44(W1):W90–W97. 573

36. Slenter DN, Kutmon M, Hanspers K, Riutta A, Windsor J, Nunes N, et al. WikiPathways: a multifaceted pathway database bridging metabolomics to other omics research. Nucleic Acids Research. 2017;46(D1):D661–D667. doi:10.1093/nar/gkx1064.

37. Rangaswami H, Bulbule A, Kundu GC. Osteopontin: role in cell signaling and cancer progression. Trends in cell biology. 2006;16(2):79–87.

38. Sharma N, Jha S. NLR-regulated pathways in cancer: opportunities and obstacles for therapeutic interventions. Cellular and molecular life sciences. 2016;73(9):1741–1764.

39. Whyte J, Bergin O, Bianchi A, McNally S, Martin F. Key signalling nodes in mammary gland development and cancer. Mitogen-activated protein kinase signalling in experimental models of breast cancer progression and in mammary gland development. Breast Cancer Research. 2009;11(5):209.

40. Mortezaee K, Salehi E, Mirtavoos-mahyari H, Motevaseli E, Najafi M, Farhood B, et al. Mechanisms of apoptosis modulation by curcumin: Implications for cancer therapy. Journal of cellular physiology. 2019;234(8):12537–12550.

41. Colomer C, Margalef P, Villanueva A, Vert A, Pecharroman I, Solé L, et al. IKKα Kinase Regulates the DNA Damage Response and Drives Chemo-resistance in Cancer. Molecular cell. 2019;75(4):669–682.

42. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proceedings of the National Academy of Sciences. 2005;102(43):15545–15550.

43. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal statistical society: series B (Methodological). 1995;57(1):289–300.

44. Wilke CO. Bringing molecules back into molecular evolution. PLoS Computational Biology. 2012;8(6):e1002572.

45. Jassal B, Matthews L, Viteri G, Gong C, Lorente P, Fabregat A, et al. The reactome pathway knowledgebase. Nucleic acids research. 2020;48(D1):D498–D503.

46. O'Donovan KJ, Ma K, Guo H, Wang C, Sun F, Han SB, et al. B-RAF kinase drives developmental axon growth and promotes axon regeneration in the injured mature CNS. Journal of Experimental Medicine. 2014;211(5):801–814.

47. Comunanza V, Corà D, Orso F, Consonni FM, Middonti E, Di Nicolantonio F, et al. VEGF blockade enhances the antitumor effect of BRAFV 600E inhibition. EMBO molecular medicine. 2017;9(2):219–237.

48. Solit DB, Rosen N. Resistance to BRAF inhibition in melanomas. New England Journal of Medicine. 2011;364(8):772–774.

49. Manzano JL, Layos L, Bugés C, de los Llanos Gil M, Vila L, Martinez-Balibrea E, et al. Resistant mechanisms to BRAF inhibitors in melanoma. Annals of translational medicine. 2016;4(12).

50. Kutalik Z, Beckmann JS, Bergmann S. A modular approach for integrative analysis of large-scale gene-expression and drug-response data. Nature biotechnology. 2008;26(5):531.

51. Chen J, Zhang S. Integrative analysis for identifying joint modular patterns of gene-expression and drug-response data. Bioinformatics. 2016;32(11):1724–1732.

Supporting information captions

S1 Text. Accurate detection of drug associated genes from simulated responses. Simulated responses have been generated to examine the accuracy of the employed method in detecting the genes that are truly associated to drug response. Three screening thresholds, three active gene sets and two covariance structure scenarios for the repeated measurements simulation have been considered. This text includes all the details of the simulation study that we conducted.

S2 Fig. Distribution of tissue of origin across the five BRAF compounds used for cell line screening in the Genomics of Drug Sensitivity in Cancer

data. Overall, similar proportion of cell lines have been treated with all of the   630
compounds examined with smaller number of cell lines been treated with AZ628,   631
Dabrafenib and PLX-4720. Larger number of cell lines in the data set were originated   632
from the lungs, the gastrointestinal tract and the haematopoietic and lymphoid tissues.   633

S3 Fig. Prediction accuracy for each different drug and scenario. Pearson   634
correlation was estimated across observed and predicted AUC values. AUC values have   635
been computed by calculating the area under the coefficient function curve (both   636
observed and predicted). Training and test sets have been considered based on either   637
the experimental units or on cancer cell lines only.   638

S4 Table. Full gene rankings based on the estimated area under the   639
coefficient function curve (analysis on the full data set). Gene rankings of all   640
selected genes based on the magnitude of the genetic effect on drug response. A positive   641
(+) sign translates to a positive effect on cells survival after drug administration, a   642
negative (-) sign translates to a negative effect on cells survival and a mixed (0) effect   643
translates to a varying effect on cells survival which depends on drug dosage.   644
Spearman's correlation is calculated between drug dosage and gene estimated coefficient   645
function values as an indicator of the magnitude change of the gene effect over the   646
increasing dosage. Area corresponds to the area under the estimated coefficient curve   647
and the SD corresponds to the standard deviation of the area based on bootstrapping.   648
Mean fold change is calculated between the selected gene expression values of the cell   649
lines carrying BRAF mutations with respect to wild type. Protein-protein interaction   650
network distance is computed based on the shortest interaction path between the BRAF   651
gene and each of the selected genes. Here, NI denotes absence of any interaction.   652

S5 Table. Full gene rankings based on the estimated area under the   653
coefficient function curve (analysis on resistant cell lines). Gene rankings of all   654
selected genes based on the magnitude of the genetic effect on drug response. A positive   655
(+) sign translates to a positive effect on cells survival after drug administration, a   656
negative (-) sign translates to a negative effect on cells survival and a mixed (0) effect   657
translates to a varying effect on cells survival which depends on drug dosage.   658
Spearman's correlation is calculated between drug dosage and gene estimated coefficient   659
function values as an indicator of the magnitude change of the gene effect over the   660
increasing dosage. Area corresponds to the area under the estimated coefficient curve   661

and the SD corresponds to the standard deviation of the area based on bootstrapping. Mean fold change is calculated between the selected gene expression values of the cell lines carrying *BRAF* mutations with respect to wild type. Protein-protein interaction network distance is computed based on the shortest interaction path between the BRAF gene and each of the selected genes. Here, NI denotes absence of any interaction.

S6 Table. Signalling pathways linked to genes predictive of BRAF inhibitor response (analysis on the full data set). Signalling pathways along with the adjusted p-values and the number of overlapping genes obtained after pathway enrichment analysis to the full scale analysis results.

S7 Table. Signalling pathways linked to genes predictive of BRAF inhibitor response (analysis on resistant cell lines). Signalling pathways along with the adjusted p-values and the number of overlapping genes obtained after pathway enrichment analysis to the resistant cell line analysis results. **S8 Table. Overlaps between the observed gene set and oncogenic signatures in the Molecular Signatures Database (analysis on the full data set).** Overlaps have been detected using gene set enrichment analysis performed using a hypergeometric distribution. The false discovery rate analog of the hypergeometric p-value is displayed after correction for multiple hypothesis testing according to Benjamini and Hochberg.

S9 Table. Overlaps between the observed gene set and oncogenic signatures in the Molecular Signatures Database (resistant cell lines analysis). Overlaps have been detected using gene set enrichment analysis performed using a hypergeometric distribution. The false discovery rate analog of the hypergeometric p-value is displayed after correction for multiple hypothesis testing according to Benjamini and Hochberg. **S10 Fig. Prediction accuracy for each different drug and scenario: analysis performed on GDSC2 data.** Pearson correlation across observed and predicted AUC values. AUC values have been computed by calculating the area under the coefficient function curve (both observed and predicted) using the GDSC2 data. Training and test sets have been considered based on either the experimental units or on cancer cell lines only.

S11 Fig. Estimated mean drug response trajectories for *BRAF* and *HRAS* mutated and non-mutated cancer cell lines: analysis performed on GDSC2 data. Observed responses (points) and estimated mean trajectory (lines) of cells'

concentration for cancer cell lines with and without *BRAF* and *HRAS* mutations after treatment with the eight anticancer compounds examined using data from GDSC2.

S12 Table. Signalling pathways linked to genes predictive of BRAF inhibitor response (analysis on the full data set)-Reactome. Signalling pathways along with the adjusted p-values and the number of overlapping genes obtained after pathway enrichment analysis to the full scale analysis results using the Reactome database.

Acknowledgments

We acknowledge Emily Chambers and the Sheffield Bioinformatics Core Facility for their assistance in GDSC2 data pre-processing. Their input to this study is greatly appreciated.