Two opposite voltage-dependent currents control the unusual early development pattern of embryonic Renshaw cell electrical activity

Juliette Boeri[1¶], Claude Meunier[2 ¶], Hervé Le Corronc[1,3 ¶], Pascal Branchereau[4], Yulia Timofeeva[2,5,6], François Xavier Lejeune[7], Christine Mouffle[1], Hervé Arulkandarajah[1], Jean Marie Mangin[1], Pascal Legendre[1&*], Antonny Czarnecki[1,4&*]

Affiliation

[1] INSERM, UMR_S 1130, CNRS, UMR 8246, Neuroscience Paris Seine, Institute of Biology Paris Seine, Sorbonne Univ, Paris, France.
[2] Centre de Neurosciences Intégratives et Cognition, CNRS UMR 8002, Institut Neurosciences et Cognition, Université de Paris, Paris, France.
[3] Univ Angers, Angers, France.
[4] Univ. Bordeaux, CNRS, EPHE, INCIA, UMR 5287 F-33000 Bordeaux, France
[5] Department of Computer Science and Centre for Complexity Science, University of Warwick, Coventry, UK.
[6] Department of Clinical and Experimental Epilepsy, UCL Queen Square Institute of Neurology, University College London, London, UK.
[7] Institut du Cerveau et de la Moelle Epinière, Centre de Recherche CHU Pitié-Salpétrière, INSERM, U975, CNRS, UMR 7225, Sorbonne Univ, Paris, France.

* Corresponding authors
Email: pascal.legendre@inserm.fr (PL)
Email: antonny.czarnecki@u-bordeaux.fr (AC)

[¶]These authors contributed equally to this work.

[&]These authors also contributed equally to this work.

Abstract

Renshaw cells (V1R) are excitable as soon as they reach their final location next to the spinal motoneurons and display heterogeneous activity patterns at the onset of the synaptogenesis (12 days post-fertilization; E12.5). Using multiple experimental approaches, in combination with biophysical modeling and dynamical systems theory, we analyzed, for the first time, the mechanisms underlying the electrophysiological properties of V1R during early embryonic development of the spinal cord locomotor networks (E11.5-E16.5). We found that these interneurons are subdivided into several functional clusters from E11.5 and then display an unexpected transitory involution process during which they lose their ability to sustain tonic firing. V1R begin to recover this capability when a locomotor-like activity emerges (after E15.5). Our computational analysis, closely based on experimental data, demonstrated that the essential factor controlling the diversity of the discharge pattern of embryonic V1R is the ratio of a persistent sodium conductance to a delayed rectifier potassium conductance. We also found that V1R involution is due to a decrease of the persistent sodium conductance as V1R develop and their size increases. Taken together, our results reveal how a simple mechanism, based on the synergy of two voltage-dependent conductances that are ubiquitous in neurons, can produce functional diversity in V1R and control their early developmental trajectory. Our findings also indicate that homeostatic regulation of channel density or channel degeneracy known to maintain homogeneous firing pattern within specific populations of mature neurons, are unlikely to occur at these early developmental stages.

Keywords: development, spinal cord, embryo, Renshaw cell, firing pattern, functional involution, electrophysiology, biophysical modeling.

Introduction

The development of the central nervous system (CNS) follows complex steps, which depend on genetic and environmental factors and involve interactions between multiple elements of the neural tissue. Remarkably, emergent neurons begin to synchronize soon after the onset of synapse formation, generating long episodes of low frequency (<0.01 Hz) correlated spontaneous network activity (SNA) [1-8]. In the mouse embryonic spinal cord (SC), SNA is driven by an excitatory cholinergic-GABAergic loop between motoneurons (MNs) and interneurons (INs), GABA being depolarizing before embryonic day 16.5 (E16.5) [9]. SNA emerges around E12.5 [4, 6, 10-12], at a time when functional neuromuscular junctions are not yet established [13], and sensory and supraspinal inputs have not yet reached the spinal motor networks [14-17].

Several studies pointed out that SNA is an essential component in neuronal networks formation. [18-21]. In the SC, pharmacologically-induced disturbances of SNA between E12.5 and E14.5 induce defects in the formation of motor pools, in motor axon guidance to their target muscles and in the development of motor networks [4, 21-23]. During SNA episodes, long lasting giant depolarization potentials (GDPs) are evoked in the SC, mainly by the massive release of GABA onto MNs [12]. Immature Renshaw cells (V1[R]) are likely the first GABAergic partners of MNs in the mouse embryo [24, 25], and the massive release of GABA during SNA probably requires that many of them display repetitive action potential firing or plateau potential activity [25].

However, little is known about the firing pattern of embryonic V1[R] and the maturation of their intrinsic properties. We recently found that V1[R] exhibit heterogeneous excitability properties when SNA emerges in the SC [25] in contrast to adult Renshaw cells that constitute a functionally homogeneous population [26, 27]. Whether this early functional

diversity really reflects distinct functional classes of V1^R, how this diversity evolves during

development, and what are the underlying biophysical mechanisms remain open questions.

The present study addresses these issues using multiple approaches, including patch-clamp

recordings, cluster analysis, biophysical modeling and dynamical systems theory. The firing

patterns of V1^R and the mechanisms underlying their functional diversity are analyzed during

a developmental period covering the initial phase of development of SC activity in the

mouse embryo (E11.5-E14.5), when SNA is present, and during the critical period (E14.5-

E16.5), when GABAergic neurotransmission gradually shifts from excitation to inhibition [28]

and locomotor-like activity emerges [4, 10, 11].

We discover that the balance between the slowly inactivating subthreshold persistent

sodium inward current (I_{Nap}) [29] and the delayed rectifier potassium outward current

(I_{Kdr}), accounts for the heterogeneity of embryonic V1^R and the changes in firing pattern

during development. The heterogeneity of V1^R at E12.5 arises from the existence of distinct

functional groups. Surprisingly, and in opposition to the classically accepted development

scheme [30-35], we show that the embryonic V1^R population loses its ability to support tonic

firing from E13.5 to E15.5, exhibiting a transient functional involution during its

development. Our experimental and theoretical results provide a global view of the

developmental trajectories of embryonic V1^R. They demonstrate that a simple mechanism,

based on the synergy of only two major opposing voltage-dependent currents, accounts for

functional diversity in these immature neurons.

Results

The delayed rectifier potassium current I_{Kdr} **is a key partner of the persistent sodium**

current I_{Nap} **in controlling embryonic V1^R firing patterns during development**

We previously highlighted that V1^R are spontaneously active at E12.5. Their response to a 2 s

suprathreshold depolarizing current steps revealed four main patterns, depending of the

recorded interneuron [25]: i) single spiking (SS) V1^R that fires only 1-3 APs at the onset of the

depolarizing pulse, ii) repetitive spiking (RS) V1^R, iii) mixed events (ME) V1^R that shows an

alternation of action potentials (APs) and plateau potentials or, iv) V1^R that displays a long-

lasting sodium-dependent plateau potential (PP) (*Figure 1A1–A4*).

We also uncovered a relationship between I_{Nap} and the ability of embryonic V1^R to sustain

repetitive firing [25]. However, the heterogeneous firing patterns of V1^R observed at E12.5

could not be fully explained by variations in I_{Nap} [25], suggesting the involvement of other

voltage-gated channels in the control of the firing pattern of V1^R, in particular potassium

channels, known to control firing and AP repolarization. Our voltage clamp protocol,

performed in the presence of TTX (1 μM), did not disclose any inward rectifying current

(hyperpolarizing voltage steps to -100 mV from V_H = -20 mV, data not shown), but revealed

two voltage-dependent outward potassium currents, a delayed rectifier current (I_{Kdr}) and a

transient potassium current (I_A) in all embryonic V1^R, whatever the firing pattern (*Figure

1B1–B4*). These currents are known to control AP duration (I_{Kdr}) or firing rate (I_A),

respectively [36]. The activation threshold of I_{Kdr} lied between -30 mV and -20 mV and the

threshold of I_A between -60 mV and -50 mV, (n = 27; N = 27 embryos) (*Figure 1C1–C4*).

Removing external calcium had no effect on potassium current I/V curves (data not shown),

suggesting that calcium-dependent potassium currents are not yet present at E12.5.

It was unlikely that the heterogeneity of V1^R firing patterns resulted from variations in the

intensity of I_A. Indeed, its voltage-dependent inactivation (time constant: 23.3 ± 2.6 ms, n =

8; N = 8), which occurs during the depolarizing phase of an AP, makes it ineffective to control

AP or plateau potential durations. This was confirmed by our theoretical analysis (*see Figure

7—figure supplement 1). We thus focused our study on I_{Kdr}. At E12.5, PP V1^R had a

significantly lower G_{Kdr} (2.12 ± 0.44 nS, n = 6; N = 6) than SS V1^R (5.57 ± 0.56 nS, n = 9; N = 9)

and RS V1^R (6.39 ± 0.83 nS, n = 7; N = 7) (*Figure 1D*). However, there was no significant

difference in G_{Kdr} between SS V1^R and RS V1^R at E12.5 (*Figure 1D*), which indicated that

variations in G_{Kdr} alone could not explain all the firing patterns observed at E12.5. Similarly,

there was no significant difference in G_{Nap} between RS V1^R (0.91 ± 0.21nS, n = 8; N = 8) and

PP V1^R (1.24 ± 0.19 nS, n = 6; N = 6) at E12.5 (*Figure 1E*), indicating that variations

in G_{Nap} alone could not explain all the firing patterns of V1^R at E12.5 [25]. In contrast

G_{Nap} measured in SS V1^R at E12.5 (0.21 ± 0.20 nS, n = 9; N = 9) were significantly lower

compared to G_{Nap} measured in RS V1^R and in PP V1^R at E12.5 (*Figure 1E*).

Mature neurons often display multiple stable firing patterns [37-39]. This usually depends on

the combination of several outward and inward voltage- or calcium-dependent

conductances and on their spatial localization [37-39]. In contrast, immature V1^R have a

limited repertoire of voltage-dependent currents (I_{Nat} and I_{Nap}, I_{Kdr} and I_A) at E12.5, and

we did not find any evidence of voltage-dependent calcium currents at this age [25].

Blocking I_{Nap} prevented plateau potential activity, PP-V1^R becoming unexcitable, and turned

repetitive spiking V1^R into single spiking V1^R [25]. Therefore, we hypothesized that the

different firing patterns of V1^R observed at E12.5 were related to the G_{Nap} / G_{Kdr} ratio only,

with variations in the intensity of I_A being unlikely to account for the heterogeneity of firing

pattern. We found that this ratio was significantly lower for SS V1^R recorded at E12.5

(G_{Nap} / G_{Kdr} = 0.043 ± 0.015, n = 9) compared to RS V1^R (0.154 ± 0.022, n = 8) and PP V1^R

(0.66 ± 0.132, n = 6) (*Figure 1F*). We also found that the G_{Nap} / G_{Kdr} ratio was significantly

lower for RS V1^R compared to PP V1^R (*Figure 1F*).

Altogether, these results strongly suggest that, although the presence of I_{Nap} is required for

embryonic V1^R to fire repetitively or to generate plateau potentials [25], the heterogeneity

of the firing pattern observed between E12.5 is not determined by I_{Nap} *per se* but likely by

the balance between I_{Nap} and I_{Kdr}.

Manipulating the balance between G$_{Nap}$ and G$_{Kdr}$ changes embryonic V1^R firing patterns

We previously showed that blocking I_{Nap} with riluzole converted PP V1^R or RS V1^R into SS

V1^R [25]. To confirm further that the balance between G_{Nap} and G_{Kdr} was the key factor in

the heterogeneity of V1^R firing patterns, we assessed to what extent a given E12.5 SS V1^R cell

could change its firing pattern when I_{Kdr} was gradually blocked by 4-aminopiridine (4-AP).

We found that I_{Kdr} could be blocked by micromolar concentrations of 4-AP without

affecting I_A (*Figure 2—figure supplement 1*). 4-AP, applied at concentrations ranging from

0.3 μM to 300 μM, specifically inhibited I_{Kdr} with an IC$_{50}$ of 2.9 μM (*Figure 2—figure*

supplement 1C1).

We then determined to what extent increasing the concentration of 4-AP modified the firing

pattern of V1^R at E12.5. Applying 4-AP at concentrations ranging from 3 μM to 300 μM

changed the firing pattern of SS V1^R (n = 10; N = 10) in a concentration-dependent manner

(*Figure 2A1–A3*). In 50% of the recorded V1^R, increasing 4-AP concentrations successfully

transformed SS V1^R into PP V1^R with the following sequence: SS → RS → ME → PP (*Figure*

2A1). In a second group of embryonic V1^R (25%), 4-AP application only evoked mixed activity,

with the same sequence as aforementioned (SS → RS → ME) (data not shown). In the

remaining SS V1^R (25%), increasing 4-AP concentration only led to sustained AP firing (*Figure*

2A2). Application of 300 μM 4-AP on RS V1^R at E12.5 evoked mixed events or plateau

potentials (*Figure 2—figure supplement 2*). Plateau potentials and repetitive spiking evoked

in the presence of 300 μM 4-AP were fully blocked by 0.5-1 μM TTX, indicating that they

were generated by voltage-gated Na$^+$ channels (*Figure 2B,C* and *Figure 2—figure*

supplement 2). It should be noted that the application of 300 μM of 4-AP induced a

significant 30.5 ± 12.4 % increase (P = 0.0137; Wilcoxon test) of the input resistance (1.11 ±

0.08 GΩ versus 1.41 ± 0.12 GΩ; n = 11; N = 11).

These results show that, in addition to I_{Nap}, I_{Kdr} is also a major determinant of the firing

pattern of embryonic V1^R. The above suggests that the firing patterns depend on a synergy

between I_{Nap} and I_{Kdr} and that the different patterns can be ordered along the following

sequence SS → RS → ME → PP when the ratio G_{Nap}/ G_{Kdr} is increased.

The heterogeneity of the V1^R **firing patterns decreases during embryonic development**

It was initially unclear whether these different firing patterns corresponded to well

separated classes within the E12.5 V1^R population or not. To address this question, we

performed a hierarchical cluster analysis on 163 embryonic V1^R, based on three quantitative

parameters describing the firing pattern elicited by the depolarizing pulse: the mean

duration of evoked APs or plateau potentials measured at half-amplitude (mean ½Ad), the

variability of the event duration during repetitive firing (coefficient of variation of ½Ad: CV

½Ad) and the total duration of all events, expressed in percentage of the pulse duration

(depolarizing duration ratio: ddr) (*Figure 3A inserts*). In view of the large dispersion of mean

½Ad and ddr values, cluster analysis was performed using the (decimal) logarithm of these

two quantities [40]. The analysis of the distribution of log mean ½Ad, CV ½Ad and log ddr

revealed multimodal histograms that could be fitted with several Gaussians (*Figure 3—*

figure supplement 1A1–C1). Cluster analysis based on these three parameters showed that

the most likely number of clusters was 5 (*Figure 3A,B*), as determined by the silhouette

width measurement (*Figure 3B*). Two clearly separated embryonic V1^R groups with CV ½Ad =

0 stood out, as shown in the 3D plot in *Figure 5C*. The cluster with the largest ½Ad (mean

½Ad = 833.5 ± 89.99 ms) and the largest ddr (0.441 ± 0.044) contained all PP V1^R (n = 35; N =

29) (*Figure 3C-D* and *Figure 3—figure supplement 1A2,C2*). Similarly, the cluster with the

shortest ½Ad (9.73 ± 0.66 ms) and the lowest ddr (0.0051 ± 0.0004) contained all SS V1^R (n =

46; N = 37) (*Figure 3C-D* and *Figure 3—figure supplement 1A2,C2*).

The three other clusters corresponded to V1^R with nonzero values of CV ½Ad (*Figure 3C*). A

first cluster regrouping all RS V1^R (n = 69; N = 61) was characterized by smaller values of ½Ad

(23.91 ± 1.43 ms), CV ½Ad (27.36 ± 1.64%) and ddr (0.11 ± 0.01) (*Figure 3C-D* and *Figure 3—*

figure supplement 1A2,C2). The last two clusters corresponded to ME V1^R (*Figure 3C,D*). The

smaller cluster, characterized by a larger CV ½Ad (170.9 ± 8.9%; n= 4; N = 4), displayed a mix

of APs and short plateau potentials, while the second cluster, with smaller CV ½Ad (87.61 ±

7.37%; n = 9; N = 9), displayed a mix of APs and long-lasting plateau potentials (*Figure 3D*

and *Figure 3—figure supplement 1B2*). Their ½Ad and ddr values were not significantly

different (*Figure 3—figure supplement 1A2,C2*).

It must be noted that three embryonic V1^R (1.8%) were apparently misclassified since they

were aggregated within the RS cluster although having zero CV ½Ad (*Figure 3C; arrows*).

Examination of their firing pattern revealed that this was because they generated only two

APs, although their ddr (0.16 to 0.2) and ½ Ad values (31.6 to 40.3 ms) were well in the range

corresponding of the RS cluster.

These different firing patterns of V1^R might reflect different states of neuronal development

[31, 41-43]. Single spiking and/or plateau potentials are generally believed to be the most

immature forms of firing pattern, repetitive spiking constituting the most mature form [19,

44]. If it were so, the firing patterns of embryonic V1^R would evolve during embryonic

development from single spiking or plateau potential to repetitive spiking, this latter firing pattern becoming the only one in neonates [26] and at early postnatal stages [27]. However, RS neurons already represent 41% of V1^R at E12.5. We therefore analyzed the development of firing patterns from E11.5, when V1^R terminate their migration and reach their final position [45], to E16.5. This developmental period covers a first phase of development (E11.5-E14.5), where lumbar spinal networks exhibit SNA, and a second phase (E14.5-E16.5), where locomotor-like activity emerges [4, 11, 46, 47]. We first analyzed changes in the intrinsic properties (input capacitance C_{in}, input resistance $R_{in} = 1/G_{in}$ and spike voltage threshold) of V1^R. C_{in} did not change significantly from E11.5 to E13.5 (*Figure 4A1*), remaining of the order of 12 pF, in agreement with our previous work [25]. However, it increased significantly at the transition between the two developmental periods (E13.5-E15.5) to reach about 23.5 pF at E15.5 (*Figure 4A1*). A similar developmental pattern was observed for R_{in}, which remained stable during the first phase from E11.5 to E14.5 ($R_{in} \approx 1$-1.2 GΩ) but decreased significantly after E14.5 to reach about 0.7 GΩ at E15.5 (*Figure 4A2*). Spike threshold also decreased significantly between the first and the second developmental phases, dropping from about -34 mV at E12.5 to about -41 mV at E16.5 (*Figure 4A3*). Interestingly, this developmental transition around E14.5 correspond to the critical stage at which SNA gives way to a locomotor-like activity [11, 46, 47] and rhythmic activity becomes dominated by glutamate release rather than acetylcholine release [4].

This led us to hypothesize that this developmental transition could be also critical for the maturation of V1^R firing patterns. The distinct firing patterns observed at E12.5 were already present at E11.5 (*Figure 4B1,C*), but the percentage of RS V1^R strongly increased from E11.5 to E12.5, while the percentage of ME V1^R decreased significantly (*Figure 4C*). The heterogeneity of V1^R firing patterns then substantially diminished. Plateau potentials were

no longer observed at E13.5 (*Figure 4B2,C*), and ME V1^R disappeared at E14.5 (*Figure 4B3,C*).

Interestingly, the proportion of SS V1^R remained high from E13.5 to E15.5 and even slightly

increased (91.23% at E14.5 and 93.33% at E15.5; *Figure 4C*). This trend was partially

reversed at E16.5, as the percentage of RS V1^R increased at the expense of SS V1^R (67.86% SS

V1^R and 32.34% RS V1^R; *Figure 4B5,C*). This decrease in repetitive firing capability after E13.5

was surprising in view of what is classically admitted on the developmental pattern of

neuronal excitability [18, 48]. Therefore, we verified that it did not reflect the death of some

V1^R after E13.5. Our data did not reveal any activated caspase3 (aCaspase3) staining in V1^R

(FoxD3 staining) at E14.5 (n = 10 SCs; N = 10) (*Figure 5*), in agreement with previous reports

showing that developmental cell death of V1^R does not occur before birth [49].

To determine whether G_{Nap} and G_{Kdr} also controlled the firing pattern of V1^R at E14.5 (see

Figure 4B3,C), we assessed the presence of I_{Nap} and I_{Kdr} in single spiking V1^R at this

embryonic age. Both I_{Nap} and I_{Kdr} were present in V1^R at E14.5 (*Figure 6—figure

supplement 1* **and** *Figure 6—figure supplement 2*) whereas, as in V1^R at E12.5, no calcium-

dependent potassium current was detected at this developmental age (not shown). In SS V1^R,

G_{Kdr} was significantly higher at E14.5 (11.11 ± 1.12 nS, n = 10; N = 10) than at E12.5 (*Figure

1D*). In contrast, G_{Nap} was similar at E14.5 (0.13 ± 0.14 nS, n = 10; N = 10) and E12.5 (*Figure

1E*). We also found that the G_{Nap} / G_{Kdr} ratio was significantly lower for SS V1^R recorded at

E14.5 (0.012 ± 0.004, n = 10) compared to RS V1^R (0.154 ± 0.022, n = 8) and PP V1^R (0.66 ±

0.132, n = 6) recorded at E12.5 (*Figure 1F*).

We tested the effect of 4-AP in SS V1^R at E14.5. At this embryonic age, 300 μM 4-AP

inhibited only 59.2% of I_{Kdr}. Increasing 4-AP concentration to 600 μM did not inhibit I_{Kdr}

significantly more (60.2%) (*Figure 6—figure supplement 2*), indicating that inhibition of I_{Kdr}

by 4-AP reached a plateau at around 300 μM. 600 μM 4-AP application had no significant

effect on I_A (*Figure 6—figure supplement 2*). The application of the maximal concentration

of 4-AP tested (600 μM) converted SS V1^R (n = 13; N = 13) to PP V1^R (23.1%; *Figure 6A1,B*),

RS V1^R (38.5%; *Figure 6A2,B*) or ME V1^R (38.4%; *Figure 6B*), as was observed at E12.5, thus

indicating that the firing pattern of V1^R depends on the balance between I_{Nap} and

I_{Kdr} also at E14.5. Plateau potential and repetitive spiking recorded in the presence of 4-AP

at E14.5 were fully blocked by 0.5-1 μM TTX indicating that they were generated by voltage-

gated sodium channels (*Figure 6A1,A2*), as observed at E12.5.

Theoretical analysis: the basic model

As shown in *Figure 7A* for 26 cells, in which both G_{Nap} and G_{Kdr} were measured, the three

largest clusters revealed by the hierarchical clustering analysis (SS, RS and PP, which account

together for the discharge of more than 95% of cells, see *Figure 5*) correspond to well

defined regions of the G_{Nap} - G_{Kdr} plane. Single spiking is observed only when G_{Nap} is

smaller than 0.6 nS. For larger values of G_{Nap}, repetitive spiking occurs when G_{Kdr} is larger

than 3.5 nS, and V1^R display plateau potentials when G_{Kdr} is smaller than 3.5 nS. Mixed

events (ME, 4.5% of the 163 cells used in the cluster analysis), where plateaus and spiking

episodes alternate, are observed at the boundary of RS and PP clusters. This suggested to us

that a conductance-based model incorporating only the leak current, I_{Nat}, I_{Nap} and I_{Kdr} (see

Materials and Methods) could account for most experimental observations, the observed

zonation being explained in terms of bifurcations between the different stable states of the

model. Therefore, we first investigated a simplified version of the model without I_A and slow

inactivation of I_{Nap}.

A one-parameter bifurcation diagram of this "basic" model is shown in *Figure 7B* for two

values of G_{Kdr} (2.5 nS and 10 nS) and a constant injected current I = 20 pA. In both cases,

the steady-state membrane voltage (stable or unstable) and the peak and trough voltages of

stable and unstable periodic solutions are shown as the function of the maximal

conductance G_{Nap} of the I_{Nap} current, all other parameters being kept constant. For G_{Kdr} =

10 nS, the steady-state membrane voltage progressively increases (in gray) with G_{Nap}, but

repetitive spiking (in red, see voltage trace for G_{Nap} = 1.5 nS) is not achieved until G_{Nap}

reaches point SN$_1$, where a saddle node (SN) bifurcation of limit cycles occurs. This fits with

the experimental data, where a minimal value of G_{Nap} is required for repetitive spiking (see

also [25], and is in agreement with the known role of I_{Nap} in promoting repetitive discharge

[50, 51]. Below SN$_1$, the model responds to the onset of a current pulse by firing only one

spike before returning to quiescence (see voltage trace for G_{Nap}= 0.2 nS), or a few spikes

when close to SN$_1$ (not shown) before returning to quiescence. The quiescent state becomes

unstable through a subcritical Hopf bifurcation (HB) at point HB$_1$, with bistability between

quiescence and spiking occurring between SN$_1$ and HB$_1$ points. Repetitive firing persists

when G_{Nap} is increased further and eventually disappears at point SN$_2$. The firing rate does

not increase much throughout the RS range (*Figure 7—figure supplement 1C*), remaining

between 20.1 Hz (at SN$_1$) and 28.7 Hz (at SN$_2$). A stable plateau appears at point HB$_2$ through

a subcritical HB. The model is bistable between HB$_2$ and SN$_2$, with plateau and large

amplitude APs coexisting in this range.

The model behaves very differently when G_{Kdr} is reduced to 2.5 nS (gray-blue curve in

Figure 7B). It exhibits a unique stable fixed point whatever the value of G_{Nap} is, and the

transition from quiescence to plateau is gradual as G_{Nap} is increased. No repetitive spiking is

ever observed. This indicates that the activity pattern is controlled not only by G_{Nap} but also

by G_{Kdr}. This is demonstrated further in *Figure 7C*, where G_{Nap} was fixed at 1.2 nS while

G_{Kdr} was increased from 0 to 25 nS. The model exhibits a plateau potential until G_{Kdr} is

increased past point the subcritical HB point HB_2, repetitive spiking sets in before at point

SN_2 *via* a SN of limit cycles bifurcation. When G_{Kdr} is further increased, repetitive firing

eventually disappears through a SN bifurcation of limit cycles at point SN_1, the quiescent

state becomes stable through a subcritical HB at point HB_1, and bistability occurs between

these two points. This behavior is in agreement with *Figure 7A*.

Since both conductances G_{Nap} and G_{Kdr} control the firing pattern of embryonic V1[R] cells, we

computed a two-parameters bifurcation diagram (*Figure 7D*), where the stability regions of

the different possible activity states and the transition lines between them are plotted in the

G_{Nap} - G_{Kdr} plane. The black curves correspond to the bifurcations HB_1 and HB_2 and delimit

a region where only repetitive firing occurs. The red curves correspond to the SN

bifurcations of periodic orbits associated with the transition from quiescence to firing (SN_1)

and the transition from plateau to firing (SN_2). They encompass a region (shaded area)

where repetitive firing can be achieved but may coexist with quiescence (between the HB_1

and SN_1 lines) or plateau potential (in the narrow region between the HB_2 and SN_2 lines).

Some important features of the diagram must be emphasized: 1) minimal values of both

G_{Nap} (to ensure sufficient excitability) and G_{Kdr} (to ensure proper spike repolarization) are

required for repetitive spiking, 2) quiescence and plateau can be clearly distinguished only

when they are separated by a region of repetitive spiking (see also *Figure 7B* for G_{Kdr} = 10

nS), otherwise the transition is gradual (*Figure 7B* for G_{Kdr}= 2.5 nS), 3) only oblique lines

with an intermediate slope cross the bifurcation curve and enter the RS region (see, for

example, the red line in *Figure 7D*). This means that repetitive spiking requires an

appropriate balance between I_{Nap} and I_{Kdr}. If the ratio G_{Nap}/G_{Kdr} is too large (blue line) or

too small (gray line), only plateau potentials or quiescence will be observed at steady state.

This is exactly what is observed in experiments, as shown by the cumulative distribution

function of the ratio G_{Nap}/G_{Kdr} for the different clusters of embryonic V1[R] in *Figure 7E*

(same cells as in *Figure 7A*). The ratio increases according to the sequence SS → RS → ME →

PP, with an overlap of the distributions for SS V1[R] and RS V1[R]. Note also that the ratio for ME

cells (around 0.25) corresponds to the transition between repetitive spiking and plateau

potentials (more on this below).

Embryonic V1[R] cells display voltage fluctuations that may exceed 5 mV and are presumably

due to channel noise. The relatively low number of sodium and potassium channels (of the

order of a few thousands) led to voltage fluctuations in the stochastic version of our model

comparable to those seen experimentally when the cell was quiescent (top voltage trace in

Figure 7D) or when a voltage plateau occurred (bottom trace). Channel noise caused some

jitter during repetitive spiking (middle trace), and induced clearly visible variations in the

amplitude of APs. However, repetitive firing proved to be very robust and was not disrupted

by voltage fluctuations. Altogether, channel noise little alters the dynamics (compare the

deterministic voltage traces in *Figure 7B* and the noisy traces in *Figure 7D*). This is likely

because channel noise has a broad power spectrum and displays no resonance with the

deterministic solutions of the model.

The one-parameter bifurcation diagram of the model was not substantially modified when

we took I_A into account, as shown in *Figure 6—figure supplement 1*. It just elicited a slight

membrane hyperpolarization, an increase in the minimal value of G_{Nap} required for firing,

and a decrease of the firing frequency. The transition from repetitive firing to plateau was

not affected because I_A is then inactivated by depolarization.

The bifurcation diagram of *Figure 7D* accounts *qualitatively* for the physiological data on V1[R]

at E12.5 presented in *Figure 7A*, as shown in *Figure 7F* where the conductance data of

Figure 7A were superimposed on it. However, one must beware of making a more

quantitative comparison because the theoretical bifurcation diagram was established for a

constant injected current of 20 pA, whereas the current injected in experiments data varied

from neuron to neuron and ranged from 10 to 30 pA in the sample shown in *Figure 7A*. The

position of bifurcation lines in the G_{Nap} - G_{Kdr} plane depends not only on the value of the

injected current, but on the values chosen for the other parameters, which also vary from

cell to cell but were kept at fixed values in the model (Ori et al. 2018). For instance, the

diagrams were computed in *Figure 7D,F* for G_{in}= 1 nS and C_{in}= 13 pF, the median values of

the input conductance and capacitance at E12.5, taking no account of the cell-to-cell

variations of these quantities. Between E12.5 and E14.5, C_{in} which provides an estimate of

the cell size, increases by 38% in average, whereas G_{in} is not significantly modified (see

Figure 4). As illustrated in *Figure 7G* the two-parameters bifurcation diagram is then shifted

upward and rightward compared to *Figure 7F*, because larger conductances are required to

obtain the same firing pattern. The observed regression of excitability from E12.5 to E14.5-

E15.5 (see *Figure 4C*) thus comes from a decrease in G_{Nap} density (see presumable

developmental trajectories indicated by arrows in *Figure 7F*) together with a shift of the RS

region as cell size increases. As a result, all 10 cells shown in *Figure 7G* are deeply inside the

SS region at E14.5.

It is less straightforward to explain on the basis of our model the experiments where 4-AP

changed the firing pattern of SS V1[R] (*Figure 2*). Indeed, the decrease of G_{Kdr} (*Figure 6—*

figure supplement 2), although it may exceed 70% at the higher concentrations of 4-AP we

used, is not sufficient by itself to account for the change in the firing pattern of V1[R] (*Figure*

6—figure supplement 2) because data points in the SS cluster will not cross the bifurcation

lines between SS and RS (SN1) and between RS and PP (SN2) when displaced downward in

the G_{Nap}- G_{Kdr} plane. However, 4-AP at a 300 μM concentration also decreases G_{in} (by 23%

in average and up to 50% in some neurons), the rheobase current with it, and the current

that was injected in cells during experiments was reduced accordingly. hen we take into

account this reduction of both G_{in} and I the two parameters bifurcation diagram pf the

model remains qualitatively the same, but it is shifted leftward and downward in the G_{Nap}-

G_{Kdr} plane (*Figure 6—figure supplement 2*). As a consequence, the bifurcation lines

between SS and RS and between RS and PP (SN2) are then successively crossed when G_{Kdr} is

reduced, in accordance with experimental results.

Theoretical analysis: slow inactivation of I_{Nap} and bursting

Our basic model accounts for the firing pattern of 73% of the 163 cells used in the cluster

analysis. However, bursting, under the form of recurring plateaus separated by brief

repolarization episodes (see a typical trace in *Figure 8A left*), was experimentally observed in

half of PP V1[R] (24 out of 46), and plateaus intertwined with spiking episodes were recorded

in the 13 cells of the ME cluster (8% of the total sample, see *Figure 8A right* for a typical

example). Recurrent plateaus indicate membrane bistability and require that the $I - V$ curve

be S-shaped. This occurs when G_{Nap} is large and G_{Kdr} small. (*Figure 8B1,B2*). However, our

basic model lacks a mechanism for switching between quiescent state and plateau, even in

this case. Channel noise might induce such transitions, but our numerical simulations

showed that this is too infrequent to account for bursting (see voltage trace in *Figure 8B1*

where the plateau state is maintained despite channel noise).

To explain recurrent plateaus during a constant current pulse, we have to incorporate in our

model an additional slow dynamical process. Therefore, we took into account the slow

inactivation of I_{Nap} that is observed in experiments. I_{Kdr} also inactivates slowly but over

times that are much longer than the timescale of bursting, which is why we did not take its

slow inactivation into account. The one-parameter bifurcation diagram of the basic model

without slow inactivation of I_{Nap} is shown in *Figure 8C* for G_{Kdr} = 5 nS and an injected

current reduced to 10 pA (as compared to 20 pA in the previous section), so as to allow for

bistability (see *Figure 8B2*). The $G_{Nap} - V$ curve is then S-shaped, as shown in *Figure 8B1*,

with a bistability region for G_{Nap} between 1.36 and 1.85 nS. This is in contrast with *Figure 7B*

where the $G_{Nap} - V$ curve was monotonic. Adding the slow (de)inactivation of I_{Nap} then

caused periodic transitions between up (plateau) and down (quiescent) states, and the

model displayed a stable limit cycle (shown in black in *Figure 8C*), as illustrated by the top

voltage trace on the right. This mechanism is known as pseudo-plateau or plateau-like

bursting (a.k.a. fold-subcritical HB bursting) [52]. It is akin to square wave bursting [53-56],

but the up-state is a stable fixed point rather than a limit cycle [57-59], which is why

recurrent plateaus are obtained rather than bursts of APs. The duration of the plateaus and

repolarization episodes depends on the values of G_{Nap} and G_{Kdr}. A voltage-independent

time constant $\tau_s = 2\,ms$ leads to up and down states of comparable durations (see top left

voltage trace in *Figure 8C*). In agreement with the bifurcation diagram of *Figure 8C*, the

persistent sodium current inactivates during plateaus (phase 1, see bottom right trace in in

Figure 8C) and de-inactivates during quiescent episodes (phase 3, see bottom right trace).

Transitions from the down-state to the up-state occurs when inactivation has reached its

maximal value (phase 2) and transition from the up-state to the down state when it has

reached its maximum (phase 4).

Increasing G_{Nap} by 50% (or decreasing G_{Kdr}) makes plateaus much longer than quiescent

episodes (see top right voltage trace in *Figure 8C*). This again points out to the fact that the

ratio is an important control parameter. Adding channel noise preserves bursting but

introduces substantial randomness in the duration of plateaus and repolarization episodes

(bottom right voltage trace in *Figure 8C*). Moreover, it substantially decreases the duration

of both plateaus and quiescent episodes by making transition between the two states easier

(compare the two top voltage traces in *Figure 8C*, both computed for $\tau_s = 2\ ms$). We also

noted that adding the I_A current lengthened the quiescence episodes (*Figure 6—figure*

supplement 1).

Slow inactivation of I_{Nap} also provides an explanation for mixed patterns, where plateaus

alternate with spiking episodes (*Figure 8A, right*). They take place in our model near the

transition between repetitive spiking and plateau, as in experiments (see *Figure 8A*). Slow

inactivation can lead to elliptic bursting, notably when the bifurcation HB$_2$ is subcritical [60,

61], which is the case here (*Figure 8D*). The model then displays a stable limit cycle with

alternating plateaus and spiking episodes, arising from crossing the bifurcation points HB$_2$

and SN$_2$ back and forth (see bifurcation diagram in *Figure 8D* and top voltage trace). We

note that sufficient de-inactivation of I_{Nap} for triggering a new plateau (phase 3 in the

bottom trace of *Figure 8D*) may be difficult to achieve during spiking episodes, because

voltage oscillates over a large range, which tends to average out the variations of the

inactivation level. If de-inactivation is not sufficient, the model keeps on spiking repetitively

without returning to the plateau state. This is what occurs for cells well within the RS region,

far away from the RS-PP transition. It also probably explains why it was difficult in many

recorded cells to elicit plateaus by increasing the injected current, inactivation balancing

then the increase of I_{Nap} induced by the larger current.

Altogether, our study shows that a model incorporating the slow inactivation of

I_{Nap} accounts for all the firing patterns displayed by cells from the PP and ME clusters.

Discussion

V1[R] constitute a homogeneous population when referring to their transcription factor

V1[R] constitute a homogeneous population when referring to their transcription factor

program during development [24, 62], their physiological function [63] and their firing

pattern at postnatal stages [27]. Surprisingly, our electrophysiological recordings and our

cluster analysis clearly indicate that distinct functional classes of V1[R] are transiently present

during development at the onset of the SNA (E11.5-E12.5). Five different groups of

embryonic V1[R] were defined using cluster analysis, according to their firing properties.

Development of the firing pattern of embryonic V1[R] during SNA

It is generally assumed that, during early development, newborn neurons cannot sustain

repetitive firing [35, 48]. Later on, neurons progressively acquire the ability to fire

repetitively, APs become sharper, and neurons eventually reach their mature firing pattern,

due to the progressive appearance of a panoply of voltage-gated channels with different

kinetics [18, 35, 48]. Our results challenge the general view that single spiking is a more

primitive form of excitability [35]. Indeed, we show that repetitive firing and plateau

potentials dominated at early stages (E11.5-E12.5), while single spiking was prevailing only

later (E13.5- E16.5).

The different V1[R] firing patterns observed at E11.5-E12.5 might reflect variability in the

maturation level between V1[R] at a given developmental stage, as suggested for developing

MNs [64, 65]. However, this is unlikely since V1[R] transiently lose their ability to sustain tonic

firing or plateau potential after E13.5. The heterogeneous discharge patterns of V1[R]

observed before E13.5 contrasts with the unique firing pattern of V1[R] at postnatal age [27].

Accordingly, the transient functional heterogeneity of V1[R] rather reflects an early initial

developmental stage (E11.5-E13.5) of intrinsic excitability.

The physiological meaning of the transient functional involution of V1[R] that follows, after

E12.5, is puzzling. To our knowledge, such a phenomenon was never described in

vertebrates during CNS development. So far, a functional involution was described only for

inner hair cells between E16 and P12 [66, 67] and cultured oligodendrocytes [68], and it was

irreversible. Because most V1[R] cannot sustain tonic firing after E12.5, it is likely that their

participation to SNA is limited to the developmental period before other GABAergic

interneuron subtypes mature and start to produce GABA and glycine [69]. Interestingly,

embryonic V1[R] begin to recover their capability to sustain tonic firing when locomotor-like

activity emerges [4, 11], a few days before they form their recurrent synaptic loop with MNs

(around E18.5 in the mouse embryos, [70]). One possible function of the transient involution

between E12.5 and E15.5 could be to regulate the growth of V1[R] axons toward their targets.

It is indeed known that low calcium fluctuations within growth cones are required for axon

growth while high calcium fluctuations stop axon growth and promote growth cone

differentiation [71].

Ion channels mechanisms underlying the functional heterogeneity of embryonic V1[R]

Blockade of I_{Nap} leads to single spiking [25], which emphasizes the importance of this

current for the occurrence of repetitive firing and plateau potentials in V1[R] at early

developmental stages. But these neurons can also switch from one firing pattern to another,

when G_{Kdr} is decreased by 4-AP, which emphasizes the importance of I_{Kdr}. We found that

the main determinant of embryonic V1[R] firing pattern is the balance between G_{Nap} and G_{Kdr}.

A Hodgkin-Huxley-type model incorporating a persistent sodium current I_{Nap} provided a

parsimonious explanation of all the firing patterns recorded in the V1[R] population at E12.5. It

provided a mathematical interpretation for the clustering of embryonic V1[R] shown by the

hierarchical analysis and accounted for the effect of 4-AP and riluzole [25] on the discharge.

Remarkably, it highlighted how a simple mechanism involving only the two opposing

currents I_{Nap} and I_{Kdr} , but not I_A, could produce functional diversity in a population of

developing neurons. The model explained why minimal G_{Nap} and G_{Kdr} are required for firing,

and how a synergy between G_{Nap} and G_{Kdr} controls the firing pattern and accounts for the

zonation of the $G_{Nap} - G_{Kdr}$ plane that is observed experimentally.

Taking into account the slow inactivation of I_{Nap} to the model allowed us to explain the

bursting patterns displayed by cells of the PP and ME clusters. We showed, in particular, that

mixed events arose from elliptic bursting at the repetitive spiking-plateau transition and that

smooth repetitive plateaus could be explained by a pseudo-plateau bursting mechanism [52,

58]. Such bursting scenario has been previously studied in models of endocrine cells [57, 72,

73] and adult neurons [74], but rarely observed in experiments.

Heterogeneity of the discharge pattern of pacemaker neurons has also been observed in the

embryonic pre-Bötzinger network [75]. However, it was related there to the gradual change

of balance between two inward currents, I_{Nap} and the calcium-activated nonselective

cationic current I_{CAN} during neuronal maturation, which led to the progressive replacement

of pseudo-plateau bursting by elliptic bursting. Such a scenario cannot account for the

variety of discharge patterns observed in embryonic V1[R] at the E11.5-12.5 stage of

development [25]. Our theoretical analysis and our experimental data clearly indicate that

the interplay between two opposing currents is necessary to explain all the firing patterns of

V1[R]. Our model is of course not restricted to embryonic V1[R], but may also apply to any

electrically compact cell, the firing activity of which is dominated by I_{Nap} and delayed

rectifier potassium currents. This is the case of many classes of embryonic cells in mammals

at an early stage of their development. It can also apply to the axon initial segment, where

G_{Nap} and G_{Kdr} are known to play the major role in the occurrence of repetitive firing [76].

Altogether our experimental and theoretical results provide a global view of the

developmental trajectories of embryonic V1^R (see *Figure 7F,G*). At E12.5, conductances of

embryonic V1^R are widely spread in the $G_{Nap} - G_{Kdr}$ plane, which explains the

heterogeneity of their firing patterns. This likely results from the random and uncorrelated

expression of sodium and potassium channels from cell to cell at this early stage. Between

E12.5 and E14.5-15.5 cell size increases, and G_{Kdr} with it, while the density of sodium

channels decreases (see *Figure 1* **and** *4*). The functional involution displayed by V1^R between

E12.5 and E15.5 thus mainly results from a decrease of G_{Nap} coordinated with an increase of

G_{Kdr}. How these synergistic processes are controlled during this developmental period

remains an open issue.

It is important to note that the presence of I_{Nap} is required for the functional diversity of V1^R.

Indeed, in the absence of I_{Nap}, V1^R lose their ability to generate plateau potentials or to fire

repetitively. More generally, when the diversity of voltage-gated channels is limited, as

observed in embryonic neurons [18], changes in the balance between I_{Kdr} and non (or

poorly) inactivating inward current can modify the firing pattern. This can be achieved not

only by I_{Nap}, but also by other slowly or non-inactivating inward conductances, such as I_{CAN}

[75]. Our work also clearly indicates that a change in the firing pattern can only occur if a

change in inward conductances cannot be counterbalanced by a corresponding change in

outward conductances. This implies that there is no homeostatic regulation of channel

density to ensure the robustness of V1^R excitability during its early development, contrarily

to the mature CNS [37]. In addition, the poor repertoire of voltage-gated channels at this

developmental stage precludes channel degeneracy, which is also known to ensure the

robustness of excitability in mature neurons [37].

In conclusion, our results demonstrate that a simple mechanism involving only two slowly

inactivating voltage-gated channels with opposite effects is sufficient to produce functional

diversity in immature neurons having a limited repertoire of voltage-gated channels.

Materials and Methods

Isolated spinal cord preparation

Experiments were performed in accordance with European Community guiding principles on

the care and use of animals (86/609/CEE, CE Off J no. L358, 18 December 1986), French

decree no. 97/748 of October 19, 1987 (J Off République Française, 20 October 1987, pp.

12245-12248). All procedures were carried out in accordance with the local ethics

committee of local Universities and recommendations from the CNRS. We used GAD67eGFP

knock-in mice to visualize putative GABAergic INs [77], as in our previous study [25]. To

obtain E12.5-E16.5 transgenic GAD67-eGFP embryos, 8 to 12 weeks old wild-type Swiss

female mice were crossed with GAD67-eGFP Swiss male mice.

Isolated mouse SCs from 420 embryos were used in this work and obtained as previously

described [28, 78]. Briefly, pregnant mice were anesthetized by intramuscular injection of a

mix of ketamine and xylazine and sacrificed using a lethal dose of CO_2 after embryos of

either sex were removed. Whole SCs were isolated from eGFP-positive embryos and

maintained in an artificial cerebrospinal fluid (ACSF) containing 135 mM NaCl, 25 mM

$NaHCO_3$, 1 mM NaH_2PO_4, 3 mM KCl, 11 mM glucose, 2 mM $CaCl_2$, and 1 mM $MgCl_2$ (307

mOsm/kg H_2O), continuously bubbled with a 95% O_2-5% CO_2 gas mixture.

In the lumbar SC of GAD67eGFP mouse embryos, eGFP neurons were detected using 488 nm

UV light. They were localized in the ventro-lateral marginal zone between the motor

columns and the ventral funiculi [62]. Embryonic $V1^R$ identity was confirmed by the

expression of the forkhead transcription factor Foxd3 [25].

Whole-cell recordings and analysis

The isolated SC was placed in a recording chamber and was continuously perfused (2 ml/min)

at room temperature (22-26°C) with oxygenated ACSF. Whole-cell patch-clamp recordings of

lumbar spinal embryonic $V1^R$ were carried out under direct visualization using an infrared-

sensitive CCD video camera. Whole-cell patch-clamp electrodes with a resistance of 4-7 MΩ

were pulled from thick-wall borosilicate glass using a P-97 horizontal puller (Sutter

Instrument Co., USA). They were filled with a solution containing (in mM): 96.4 K

methanesulfonate, 33.6 KCl, 4 $MgCl_2$, 4 Na_2ATP, 0.3 Na_3GTP, 10 EGTA, and 10 HEPES (pH 7.2;

290 mOsm/kg-H_2O). This intracellular solution led to an equilibrium potential of chloride ions,

E_{Cl}, of about -30 mV, close to the physiological values measured at E12.5 in spinal MNs [28].

The junction potential (6.6 mV) was systematically corrected offline.

Signals were recorded using Multiclamp 700B amplifiers (Molecular Devices, USA). Data

were low-pass filtered (2 kHz), digitized (20 kHz) online using Digidata 1440A or 1550B

interfaces and acquired using pCLAMP 10.5 software (Molecular Devices, USA). Analyses

were performed off-line using pCLAMP 10.5 and Axograph 1.7.2 (Molecular devices;

RRID:SCR_014284) software packages.

In voltage-clamp mode, voltage-dependent K^+ currents (I_{Kv}) were elicited in the presence of

1 μM TTX by 500 ms depolarizing voltage steps (10 mV increments, 10 s interval) after a

prepulse of 300 ms at V_H = -100 mV. To isolate I_{Kdr}, voltage steps were applied after a 300

ms prepulse at V_H = -30 mV that inactivated the low threshold transient potassium current

I_A. I_A was then obtained by subtracting offline I_{Kdr} from the total potassium current I_{Kv}.

Capacitance and leak current were subtracted using on-line P/4 protocol provided by

pCLAMP 10.5.

In current-clamp mode, V1^R discharge was elicited using 2 s depolarizing current steps (from

0 to ≈ 50 pA in 5-10 pA increments, depending on the input resistance of the cell) with an 8 s

interval to ensure that the membrane potential returned to V_H. When a cell generated a

sustained discharge, the intensity of the depolarizing pulse was reduced to the minimal

value compatible with repetitive firing.

I_{Nap} was measured in voltage-clamp mode using a 70 mV/s depolarizing voltage ramp [79].

This speed was slow enough to preclude substantial contamination by the inactivating

transient current and fast enough to avoid substantial inactivation of I_{Nap}. Subtraction of

the current evoked by the voltage ramp in the presence of 1 μM tetrodotoxin (TTX) from the

control voltage ramp-evoked current revealed I_{Nap}.

Pharmacological reagents

During patch-clamp recordings, bath application of TTX (1 μM, Alomone, Israel) or 4-AP (0.3 -

600 μM, Sigma) was done using 0.5 mm diameter quartz tubing positioned, under direct

visual control, 50 μm away from the recording area. The quartz tubing was connected to 6

solenoid valves linked with 6 reservoirs *via* a manifold. Solutions were gravity-fed into the

quartz tubing. Their application was controlled using a VC-8 valve controller (Warner

Instruments, USA).

4-aminopyridine (4-AP; Sigma Aldrich, USA) was used to block I_{Kdr}. To determine the

concentration–response curve, $I - V$ curves of I_{Kdr} for different concentrations of 4-AP (0.3

to 300 µM) were compared to the control curve obtained in the absence of 4-AP. The

percentage of inhibition for a given concentration was calculated by dividing the peak

intensity of I_{Kdr} by the peak value obtained in control condition. The obtained normalized

concentration–response curves were fitted using the Hill equation:

$\frac{100 - I_{min}}{1 + ([4-AP]/IC_{50})^{n_H}} + I_{min}$,

where [4-AP] is the 4-AP concentration, I_{min} is the residual current (in percentage of the

peak I_{Kdr}), $100 - I_{min}$ is the maximal inhibition achieved for saturating concentration of 4-

AP, IC_{50} is the 4-AP concentration producing half of the maximal inhibition, and n_H is the

Hill coefficient. Curve fitting was performed using KaleidaGraph 4.5 (Synergy Software, USA).

Immunohistochemistry and confocal microscopy

E14.5 embryos were collected from pregnant females. Once dissected out of their yolk sac,

SCs were dissected and immediately immersion-fixed in phosphate buffer (PB 0.1 M)

containing 4% paraformaldehyde (PFA; freshly prepared in PB, pH 7.4) for 1 h at 4°C. Whole

SCs were then rinsed out in 0.12 M PB at 4°C, thawed at room temperature, washed in PBS,

incubated in NH_4Cl (50 mM), diluted in PBS for 20 min and then permeabilized for 30 min in

a blocking solution (10% goat serum in PBS) with 0.2% Triton X-100. They were incubated for

48 h at 4°C in the presence of the following primary antibodies: guinea pig anti-FoxD3

(1:5000, gift from Carmen Birchmeier and Thomas Müller of the Max Delbrück Center for

Molecular Medicine in Berlin) and rabbit anti-cleaved Caspase-3 (1:1000, Cell Signaling

Technology Cat# 9661, RRID:AB_2341188). SCs were then washed in PBS and incubated for 2

h at RT with secondary fluorescent antibodies (goat anti-rabbit-conjugated 649; donkey anti-

guinea pig-conjugated Alexa Fluor 405 [1:1000, ThermoFisher]) diluted in 0.2% Triton X-100

blocking solution. After washing in PBS, SCs were dried and mounted in Mowiol medium

(Millipore, Molsheim, France). Preparations were then imaged using a Leica SP5 confocal

microscope. Immunostaining was observed using a 40X oil-immersion objective with a

numerical aperture of 1.25, as well as with a 63X oil-immersion objective with a numerical

aperture of 1.32. Serial optical sections were obtained with a Z-step of 1 μm (40X) and 0.2-

0.3 μm (63X). Images (1024x1024; 12-bit color scale) were stored using Leica software LAS-

AF and analyzed using ImageJ 1.5 (N.I.H., USA; http//rsb.info.nih.gov/ij/) and Adobe

Photoshop CS6 (Adobe, USA) software.

Cluster analysis

To classify the firing patterns of embryonic V1^R, we performed a hierarchical cluster analysis

on a population of 163 cells. Each cell was characterized by three quantitative measures of

its firing pattern (see legend of Figure 5). After normalizing these quantities to zero mean

and unit variance, we performed a hierarchical cluster analysis using the hclust function in R

software (version 3.3.2, https://cran.r-project.org/) that implements the complete linkage

method. The intercluster distance was defined as the maximum Euclidean distance between

the points of two clusters, and, at each step of the process, the two closest clusters were

merged into a single one, thus constructing progressively a dendrogram. Clusters were then

displayed in data space using the dendromat function in the R package 'squash' dedicated to

color-based visualization of multivariate data. The best clustering was determined using the

silhouette measure of clustering consistency [80]. The silhouette of a data point, based on

the comparison of its distance to other points in the same cluster and to points in the closest

cluster, ranges from -1 to 1. A value near 1 indicates that the point is well assigned to its

cluster, a value near 0 indicates that it is close to the decision boundary between two

neighboring clusters, and negative values may indicate incorrect assignment to the cluster.

This allowed us to identify an optimal number k of clusters by maximizing the overall

average silhouette over a range of possible values for k [80], using the silhouette function in

the R package 'cluster'.

Biophysical modeling

To understand the relationship between the voltage-dependent membrane conductances

and the firing patterns of embryonic V1$^{\text{R}}$, we relied on a single compartment conductance-

based model that included the leak current, the transient and persistent components of the

sodium current, I_{Nat} and I_{Nap}, a delayed rectifier potassium current I_{Kdr} and the

inactivating potassium current I_A revealed by experiments. Voltage evolution then followed

the equation

$C_{in}\frac{dV}{dt} = G_{in}(V_r - V) + G_{Nat}m^3h(E_{Na} - V) + G_{Nap}m_p^3s(V_{Na} - V) + G_{Kdr}n^3(E_K - V) +$

$G_A m_A h_A(E_K - V) + I$ (1),

,

where C_{in} was the input capacitance; G_{in} the input conductance; G_{Nat}, G_{Nap}, G_{Kdr} and

G_A the maximal conductances of the aforementioned currents; m, m_p, n and m_A their

activation variables; h the inactivation variable of I_{Nat}, s the slow inactivation variable of

I_{Nap}, and m_A the inactivation variable of I_A. V_r is the baseline potential imposed by *ad hoc*

current injection in current-clamp experiments; E_{Na} and E_K are the Nernst potentials of

sodium and potassium ions, and I the injected current. All gating variables satisfied

equations of the form:

$\tau_x\frac{dx}{dt} = x_\infty(V) - x,$

where the (in)activation curves were modeled by a sigmoid function of the form:

$x_\infty = \dfrac{1}{1+\exp\left(-(V-V_x)/k_x\right)}$

with k_x being positive for activation and negative for inactivation. The time constant τ_x was

voltage-independent except for the inactivation variables h and s. The activation variable

m_A of I_A was assumed to follow instantaneously voltage changes.

The effect of channel noise was investigated with a stochastic realization of the model,

where channels kinetics were described by Markov-type models, assuming a unitary channel

conductance of 10 pS for all channels.

Choice of model parameters

Most model parameters were chosen on the basis of experimental measurements

performed in the present study or already reported [25]. Parameters that could not be

constrained from our experimental data were chosen from an experimentally realistic range

of values. V_r was set at -60 mV as in experiments (see *table 1*). C_{in} (average 13.15 pF, 50%

between 11.9 and 15.1 pF, only 18 cells out of 246 in the first quartile below 7.2 pF or in the

fourth quartile above 19 pF) and G_{in} (50% of cells between 0.71 and 1.18 nS, only 7 out of

242 with input conductance above 2 nS) were not spread much in the cells recorded at E12.5,

which showed that most embryonic V1[R] were of comparable size. Interestingly, C_{in} and G_{in}

were not correlated, which indicated that the input conductance was determined by the

density of leak channels rather than by the sheer size of the cell. Moreover, no correlation

was observed between the passive properties and the firing pattern [25]. Therefore, we

always set G_{in} and C_{in} to 1 nS and 13 pF in the model (except in *Figure 6—figure*

supplement 2), close to the experimental medians (0.96 nS and 13.15 pF, respectively). The

membrane time constant C_{in}/G_{in} was then equal to 13 ms, which was also close to the

experimental median (13.9 ms, N=241).

E_{Na} was set to 60 mV (see [25]). The activation curve of I_{Nap} was obtained by fitting

experimental data, leading to an average mid-activation of -36 mV and an average steepness

of 9.5 mV. The experimentally measured values of G_{Nap} were in the range 0-2.2 nS. We

assumed that the activation curve of I_{Nat} was shifted rightward by 10 mV in comparison to

I_{Nap}. No experimental data was available for the inactivation of I_{Nat}. We chose a mid-

inactivation voltage V_h= -45 mV and a steepness k_h=-5 mV. We also assumed that the

activation time constant of both I_{Nat} and I_{Nap} was 1.5 ms, and that the inactivation time

constant was voltage-dependent: $\tau_h(V) = 16.5 - 13.5\tanh((V+20)/15)$, decreasing by

one order of magnitude (from 30 ms down to 3 ms) with the voltage increase. This enabled

us to account for the shape of the action potentials recorded in experiments, showing a slow

rise time and rather long duration. The conductance G_{Nat} was not measured experimentally.

When choosing a reasonable value of 20 nS for G_{Nat}, the model behaved very much as

recorded embryonic V1[R]: with similar current threshold (typically 10-20 pA) and stable

plateau potential obtained for the largest values of G_{Nap}.

When taking into account slow inactivation of I_{Nap} (see *Figure 8*), we chose V_s = -30 mV for

the mid-inactivation voltage and set the steepness k_s at -5 mV (as for the inactivation of

I_{Nat}). For simplicity, we assumed that the inactivation time constant was voltage-

independent and set it at a value of 2 s.

E_K was set to the experimentally measured value of -96 mV [25]. The activation parameters

of I_{Kdr} were obtained by fitting the experimental data: V_n = -20 mV, k_n = 15 mV, τ_n = 10 ms

and an activation exponent of 3. The activation and inactivation properties of I_A were also

chosen based on experimental measurements. Accordingly, V_{m_A} = -30 mV, k_{m_A} = -12 mV,

V_{h_A} = -70 mV, k_{h_A} = -7 mV, and τ_{h_A}= 23 ms. When I_A was taken into account, we assumed

that $G_A = G_{Kdr}$, consistently with experimental data (see *Figure 6—figure supplement 1*).

Numerical simulations and dynamical systems analysis

We integrated numerically the deterministic model using the freeware XPPAUT [81] and a

standard fourth-order Runge-Kutta algorithm. XPPAUT was also used to compute one-

parameter and two-parameters bifurcation diagrams. The stochastic version of the model

was also implemented in XPPAUT and computed with a Gillespie's algorithm [82].

To investigate the dynamics of the model with slow inactivation of I_{Nap}, we relied on

numerical simulations together with fast/slow dynamics analysis [83]. In this approach, one

distinguishes slow dynamical variables (here only s) and fast dynamical variables. Slow

variables vary little at the time scale of fast variables and may therefore be considered as

constant parameters of the fast dynamics in first approximation. In contrast, slow variables

are essentially sensitive to the time average of the fast variables, much more than to their

instantaneous values. This separation of time scales allows one to conduct a phase plane

analysis of the full dynamics.

Statistics

Samples sizes (n) were determined based on previous experience. The number of embryos

(N) is indicated in the main text and figure captions. No power analysis was employed, but

sample sizes are comparable to those typically used in the field. All values were expressed as

mean with standard error of mean (SEM). Statistical significance was assessed by non-

parametric Kruskal-Wallis test with Dunn's post hoc test for multiple comparisons, Mann-

Whitney test for unpaired data or Wilcoxon matched pairs test for paired data using

GraphPad Prism 7.0e Software (USA). Significant changes in the proportions of firing

patterns with age were assessed by chi-square test for large sample and by Fisher's exact

test for small sample using GraphPad Prism 7.0e Software. Significance was determined as

$p < 0.05$ (*), $p < 0.01$ (**) or $p < 0.001$ (***). The exact p value was mentioned in the result

section or in the figure captions.

Acknowledgments

We thank Susanne Bolte, Jean-François Gilles and France Lam for assistance with confocal imaging (IBPS imaging facility) and IBPS rodent facility team for animal care and production. We thank University Paris Descartes for hosting Yulia Timofeeva as an invited professor. This work was supported by INSERM, CNRS, Sorbonne Université (Paris), Université de Bordeaux, Université Paris Descartes and Fondation pour la Recherche Médicale.

Additional information

Competing interests

The authors declare no competing interests

Figure captions

Figure 1. G$_{Kdr}$ and G$_{Nap}$ in embryonic V1^R at E12.5 and E14.5

 (A) Representative traces of voltage responses showing single-spiking activity in E12.5 SS V1^R (A1), repetitive action potential firing in RS V1^R (A2), Mixed of plateau potential activity and repetitive action potential firing in ME V1^R (A3) and plateau potential activity in PP V1^R (A4). (B) Representative examples of the total outward K$^+$ currents (IK$_V$ total) obtained from V_H = -100 mV (left traces), of I_{Kdr} (V_H = -30 mV, middle traces) and of isolated I_A (left traces) recorded at E12.5 in SS V1^R (B1), RS V1^R (B2), ME V1^R (B3) and PP V1^R (B4). Voltage-dependent potassium currents were evoked in response to 10 mV membrane potential steps (200 ms) from -100 or from -30 mV to +40 mV (10 s interval between pulses). V1^R were voltage clamped at V_H = -60 mV. A prepulse of -40 mV (300 ms) was applied to activate both I_A and I_{Kdr}. I_{Kdr} was isolated by applying a prepulse of 30 mV (300 ms) to inactivate I_A (B1

insert). I_A was isolated by subtracting step-by-step the currents obtained using a pre-pulse of

30 mV (V_H = - 30 mV) from the currents obtained using a pre-pulse of -40 mV (V_H = -100 mV).

(C) Current-voltage relationship ($I - V$ curves) of I_{Kdr} (filled circles) and of I_A (open circles)

recorded in SS V1^R (C1), RS V1^R (C2), ME V1^R (C3) and PP V1^R (C4). $I - V$ curves were

obtained from currents shown in B1, B2, B3 and B4. Note that $I - V$ curves are similar

between SS V1^R, RS V1^R, ME V1^R and PP V1^R. (D) Bar graph showing maximal G_{Kdr} value

(Max G_{Kdr}) in SS V1^R at E12.5 (n = 9; N = 9; gray bar) and at E14.5 (n = 10; N = 10 gray bar),

and in RS V1^R (n = 7; N = 7; red bar), ME V1^R (n = 3; N = 3 purple bar) and PP V1^R at E12.5 (n =

7; N = 7 blue bar). G_{Kdr} was calculated from I_{Kdr} at V_H = + 20 mV, assuming a K$^+$ equilibrium

potential of -96 mV. There is no significant difference in G_{Kdr} between SS V1^R and RS V1^R,

while G_{Kdr} is significantly smaller in PP V1^R as compared to SS V1^R and RS V1^R. G_{Kdr} was

significantly higher in SS V1^R at E14.5 than in SS V1^R, RS V1^R and PP V1^R at E12.5. (Kruskall-

Wallis test P < 0.0001; SS V1^R versus RS V1^R at E12.5, P = 0.5864; SS V1^R versus PP V1^R at

E12.5, P = 0.0243; RS V1^R versus PP V1^R at E12.5, P = 0.0086; E14.5 SS V1^R versus E12.5 SS V1^R,

P = 0.0048; E14.5 SS V1^R versus E12.5 RS V1^R, P = 0.0384, E14.5 SS V1^R versus E12.5 PP V1^R, P

< 0.0001). The increase in G_{Kdr} between E12.5 and E14.5 is likely to be due to the increase in

neuronal size (input capacitance; Figure 2A). Indeed, there was no significant difference

(Mann Whitney test, P = 0.133) in G_{Kdr} density between SS V1^R at E12.5 (n = 9; N = 9 gray

bar) and at E14.5 (n = 10; N = 10 gray bar). (E) Bar graph showing the maximal G_{Nap} value

(Max G_{Nap}) in SS V1^R at E12.5 (n = 9; N = 9 gray bar) and E14.5 (n = 10; N = 10 gray bar), and

in RS V1^R (n = 8; N = 8 red bar), ME V1^R (n = 3; N = 3 purple bar) and PP V1^R (n = 6; N = 6 blue

bar) at E12.5. Max G_{Nap} was calculated from maximal I_{Nap} value measured on current

evoked by assuming a Na$^+$ equilibrium potential of +60 mV. There was no difference

in G_{Nap} between RS V1^R and PP V1^R. On the contrary, G_{Nap} measured in SS V1^R at E12.5 or at

E14.5 was significantly smaller as compared to G_{Nap} measured at E12.5 in RS V1^R or in PP

V1^R. G_{Nap} measured at E12.5 and E14.5 in SS V1^R were not significantly different (Kruskall-

Wallis test $P < 0.0001$; E12.5 SS V1^R versus E12.5 RS V1^R, $P = 0.0034$; E12.5 SS V1^R versus

E12.5 PP V1^R, $P = 0.0006$; E12.5 RS V1^R versus E12.5 PP V1^R, $P = 0.5494$; E14.5 SS V1^R versus

E12.5 SS V1^R, $P = 0.5896$; E14.5 SS V1^R versus E12.5 RS V1^R, $P = 0.0005$; E14.5 SS V1^R versus

E12.5 PP V1^R, $P < 0.0001$). (F) Histograms showing the G_{Nap} / G_{Kdr} ratio in SS V1^R at E12.5 (n

= 9; gray bar) and E14.5 (n = 10; green bar) and in RS V1^R (n = 8; red bar), ME V1^R (n = 3;

purple bar) and PP V1^R (n = 6; blue bar) at E12.5. Note that the G_{Nap} / G_{Kdr} ratio differs

significantly between SS V1^R, RS V1^R and PP V1^R at E12.5, while it is not different between SS

V1^R recorded at E12.5 and at E14.5 (Kruskall-Wallis test $P < 0.0001$; SS V1^R versus RS V1^R at

E12.5, $P = 0.0367$; SS V1^R versus PP V1^R at E12.5, $P < 0.0001$; RS V1^R versus PP V1^R at E12.5, P

= 0.0159; E14.5 SS V1^R versus E12.5 SS V1^R, $P = 0.2319$; E14.5 SS V1^R versus E12.5 RS V1^R, $P =$

0.0017; E14.5 SS V1^R versus E12.5 PP V1^R $P < 0.0001$). Data shown in A and B were used to

calculate G_{Nap} / G_{Kdr} ratio shown in C. (*$P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

Figure 2. Increasing 4-AP concentration changed the firing pattern of single spiking

embryonic V1^R recorded at E12.5

The firing pattern of embryonic V1^R was evoked by 2 s suprathreshold depolarizing current

steps. (A) Representative traces showing examples of the effect of increasing concentration

of 4-AP (from 3 to 300 µM) on the firing pattern of a SS V1^R recorded at E12.5. Note that in

A1, increasing 4-AP concentration converted single spiking (gray trace) to repetitive spiking

(red trace), repetitive spiking to a mixed event pattern (purple trace) and mixed events to

plateau potential (blue trace). (A2) Example of SS V1^R in which increasing 4-AP concentration

converted single spiking to repetitive spiking only. (A3) Bar plots showing the change in the

firing pattern of SS V1^R according to 4-AP concentrations (control n = 10; N = 10, 3 µM 4-AP n

= 8; N = 8, 10 µM 4-AP n = 10; N = 10, 30 µM 4-AP n = 10; N = 10, 100 µM 4-AP n = 10; N = 10,

300 µM 4-AP n = 8; N = 8). (B) Representative traces showing the effect of 0.5 µM TTX on a

plateau potential evoked in a SS V1^R in the presence of 300 µM 4-AP. (C) Representative

traces showing the effect of 0.5 µM TTX on repetitive AP firing evoked in a SS V1^R in the

presence of 300 µM 4-AP. In both cases, the application of TTX fully blocked the responses

evoked in the presence of 4-AP, indicating that they were underlain by the activation of

voltage-gated Na$^+$ channels.

Figure 3. Cluster analysis of V1^R firing pattern at E12.5

(A, inserts) Cluster analysis of embryonic V1^R firing pattern was performed using three

parameters that describe the firing pattern during a 2 s suprathreshold depolarizing pulses:

the mean of the half-amplitude event duration (mean ½Ad), the coefficient of variation of ½

Ad (CV ½Ad) allowing to quantify the AP variation within a train (CV was set to 0 when the

number of spikes evoked by a depolarizing pulse was ≤ 3) and the duration ratio ddr = Σ½

Ad/Pw, obtained by dividing the sum of ½ Ad by the pulse duration Pw, that indicates the

total time spent in the depolarized state. For example, ddr = 1 when a plateau potential lasts

as long as the depolarizing pulse. Conversely, its value is low when the depolarizing pulse

evokes a single AP only. (A) Dendrogram for complete linkage hierarchical clustering of 164

embryonic V1^R (N = 140) according to the values of log mean ½Ad, of CV ½Ad and of log ddr.

The colored matrix below the dendrogram shows the variations of these three parameters

for all the cells in the clusters (colored trees) extracted from the dendrogram. (B) The

number of clusters was determined by analyzing the distribution of silhouette width values

(see Material and Methods). The boxplots show the distribution of silhouette width values

when the number of clusters k varies from 2 to 12. The mean silhouette width values (red

diamond shaped points) attained their maximum when the estimated cluster number was 5.

(C) 3D plot showing cluster distribution of embryonic V1[R] according to log mean ½Ad, CV

½Ad and log ddr. Each cluster corresponds to a particular firing pattern as illustrated in D.

V1[R] that cannot sustain repetitive firing of APs (1 to 3 AP/pulse only, gray, Single spiking, SS),

V1[R] that can fire tonically (red, Repetitive spiking, RS), V1[R] with a firing pattern characterized

by a mix of APs and relatively short plateau potentials (dark purple, Mixed event short PP,

ME short PP), V1[R] with a firing pattern characterized by a mix of APs and relatively long

plateau potentials (light purple, Mixed event long PP, ME long PP) and V1[R] with evoked

plateau potentials only (blue, Plateau potential, PP). The arrow in C indicates 3 misclassified

V1[R] that could not sustain repetitive firing although they were assigned to the cluster of

repetitively firing V1[R] (see text).

Figure 4. Developmental changes of embryonic V1[R] firing patterns from E11.5 to E16.5

(A1) Graph showing how the input capacitance C_{in} of V1[R] changes with embryonic age. C_{in}

significantly increased between E12.5 or E13.5 and E14.5 (Kruskall-Wallis test P < 0.0001;

E12.5 versus E11.5 P = 0.258, E12.5 versus E13.5 P = 0.904, E12.5 versus E14.5 P < 0.0001,

E12.5 versus E15.5 P < 0.0001, E12.5 versus E16.5 P < 0.0001, E13.5 versus E14.5 P < 0.0001,

E13.5 versus E15.5 P < 0.0001, E13.5 versus E16.5 P < 0.0001; E11.5 n = 31; N = 27, E12.5 n =

267; N = 152, E13.5 n = 43; N = 40, E14.5 n = 61; N = 49, E15.5 n = 16; N = 4, E16.5 n = 30; N =

9). (A2) Graph showing how the input resistance R_{in} of V1[R] changes with embryonic age. R_{in}

significantly decreased between E12.5 or E14.5 and E15.5 (Kruskall-Wallis test P < 0.0001;

E12.5 versus E11.5 P > 0.999, E12.5 versus E13.5 P = 0.724, E12.5 versus E14.5 P > 0.999,

E12.5 versus E15.5 P = 0.0004, E12.5 versus E16.5 P = 0.0005, E14.5 versus E15.5 P = 0.0019,

E14.5 versus E16.5 $P < 0.0058$; E11.5 n = 31, E12.5 n = 261; N = 146, E13.5 n = 43; N = 40,

E14.5 n = 60; N = 48, E15.5 n = 16; N = 4, E16.5 n = 30; N = 9). (A3) Graph showing how the

threshold of regenerative events (APs and plateau potentials) of V1^R changes with

embryonic age. The average threshold became significantly more hyperpolarized after E12.5

(Kruskall-Wallis test $P < 0.0001$; E12.5 versus E11.5 $P = 0.676$, E12.5 versus E13.5 $P = 0.0039$,

E12.5 versus E14.5 $P < 0.0001$, E12.5 versus E15.5 $P < 0.0001$, E12.5 versus E16.5 $P < 0.0001$,

E13.5 versus E14.5 $P > 0.999$, E13.5 versus E15.5 $P = 0.1398$, E13.5 versus E16.5 $P = 0.0013$;

E14.5 versus E15.5 $P > 0.999$, E14.5 versus E16.5 $P = 0.0634$, E15.5 versus E16.5 $P > 0.999$;

E11.5 n = 20; N = 16, E12.5 n = 162; N = 139, E13.5 n = 31; N = 28, E14.5 n = 30; N = 26, E15.5

n = 16; N = 4, E16.5 n = 30; N = 9). Yellow and purple bars below the graphs indicate the two

important phases of the functional development of spinal cord networks. The first one is

characterized by synchronized neuronal activity (SNA) and the second one is characterized

by the emergence of a locomotor-like activity (see text). Note that changes in C_{in} and

R_{in} occurred at the end of the first developmental phase. (*$P < 0.05$, ** $P < 0.01$, *** $P <$

0.001; control, E12.5). The intrinsic activation properties were analyzed using 2 s

suprathreshold depolarizing current steps. (B) Representative traces of voltage responses

showing Single Spiking (SS) V1^R (gray), Repetitive Spiking (RS) V1^R (red), ME V1^R (purple) and

Plateau Potential (PP) V1^R (blue) at E11.5 (B1), E13.5 (B2), E14.5 (B3) E15.5 (B4) and E16.5

(B5). (C) Bar graph showing how the proportions of the different firing patterns change from

E11.5 to E16.5 (E11.5 n = 22; N = 18, E12.5 n = 163; N = 140, E13.5 n = 32; N = 29, E14.5 n =

57; N = 45, E15.5 n = 15; N = 4, E16.5 n = 28; N = 9). Yellow and purple bars below the graphs

indicate the first and the second phase of functional embryonic spinal cord networks. The

proportions of the different firing patterns significantly changed between E11.5 to E12.5

(Fisher's exact test, $P = 0.0052$) with a significant increase in the proportion of RS V1^R

(Fisher's exact test, P = 0.0336) and a significant decrease in the proportion of ME V1[R]

(Fisher's exact test, P = 0.01071) at E12.5. Only two firing patterns (SS and RS) were observed

after E13.5 and most embryonic V1[R] lost their ability to sustain tonic firing after E13.5.

However, at E16.5 the proportion of RS V1[R] significantly increased at the expense of SS V1[R]

when compared to E14.5 (Fisher's exact test, P = 0.0112), indicating that embryonic V1[R]

began to recover the ability to sustain tonic firing after E15.5.

Figure 5. Activated caspase-3 is not observed in embryonic V1[R] at E14.5

Representative confocal image of the ventral part of an isolated lumbar spinal cord of E14.5

GAD67-eGFP mouse embryo showing immunostainings using antibodies against eGFP (A),

FoxD3 (B) and activated Caspase 3 (aCaspace 3) (C). (D) Superimposition of the three

stainings shows that embryonic V1[R] (eGFP+ and FoxD3+) were not aCaspase 3

immunoreactive. (A1, B1, C1 and D1). Enlarged images from A, B and C showing that

aCaspase 3 staining is localized in areas where eGFP and Foxd3 staining were absent. (A2, B2,

C2 and D2) Enlarged images from A, B and C showing that aCaspase 3 staining is absent in

the area where V1[R] (eGFP+ and FoxD3+) are located. aCaspase 3 staining that did not co-

localize with GAD67eGFP likely indicates MN developmental cell death.

Figure 6. 600 μM 4-AP changed the firing pattern of single spiking embryonic V1[R] recorded

at E14.5

The firing pattern of embryonic V1[R] was evoked by 2 s suprathreshold depolarizing current

steps. (A) Representative traces showing the effect of 4-AP application (600 μM) on the

firing pattern of single spiking (SS) V1[R] recorded at E14.5. Note that the applications of 600

μM 4-AP evoked either a plateau potential (A1) or repetitive spiking (A2), both fully blocked

by TTX. (B) Bar plots showing the proportions of the different firing patterns observed in the

presence of 600 μM 4-AP versus control recorded in SS V1^R at E14.5 (n = 14; N = 14). Single

Spiking (SS) V1^R (gray), Repetitive Spiking (RS) V1^R (red), Mixed Events (ME) V1^R (purple),

Plateau Potential (PP) V1^R (blue).

Figure 7. Embryonic V1^R firing patterns predicted by computational modeling

(A) Firing patterns of 27 recorded cells, in which both G_{Nap} and G_{Kdr} were measured. Gray:

SS, red: RS, blue: PP. The three purple points located at the boundary between the RS and PP

regions correspond to mixed events (ME) where plateau potentials alternate with spiking

episodes. Note that no cell exhibited low values of both G_{Nap} and G_{Kdr} (lower left), or large

values of both conductances (upper right). (B) Bifurcation diagram of the deterministic

model when G_{Kdr} is kept fixed to 2.5 nS or 10 nS while G_{Nap} is varied between 0 and 2.5 nS.

G_{in} = 1 nS and I = 20 pA. For G_{Kdr} = 10 nS (i.e., in the top experimental range), the red

curves indicate the maximal and minimal voltages achieved on the stable limit cycle

associated with repetitive firing (solid lines) and on the unstable limit cycle (dashed lines).

The fixed point of the model is indicated by a gray solid line when it corresponds to the

stable quiescent state, a gray dashed line when it is unstable and a solid blue line when it

corresponds to a stable plateau potential. The two HB corresponding to the change of

stability of the quiescence state (HB$_1$, G_{Nap}= 0.81 nS) and of the voltage plateau (HB$_2$, G_{Nap}=

2.13 nS) are indicated, as well as the two SN bifurcations of limit cycles associated with the

onset (SN$_1$, G_{Nap} = 0.65 nS) and offset (SN$_2$, G_{Nap}= 2.42 nS) of repetitive spiking as G_{Nap} is

increased. For G_{Kdr} = 2.5 nS, the model does not display repetitive firing; it possesses a

unique fixed point, which is always stable (blue-gray curve). The transition from quiescence

to plateau is gradual with no intervening bifurcation. Representative voltage traces of the

three different activity patterns are shown: single spiking in response to a 2 s current pulse

(gray, G_{Nap} = 0.2 nS, G_{Kdr} = 10 nS), repetitive spiking (red, G_{Nap} = 1.2 nS, G_{Kdr} = 10 nS) and

plateau potential (blue, G_{Nap} = 1.2 nS, G_{Kdr} = 2.5 nS). Note that the plateau never outlasts

the current pulse. (C) Bifurcation diagram when G_{Nap} is kept fixed at 1.2 nS and G_{Kdr} is

varied between 0 and 25 nS (I = 20 pA). Same conventions as in B. Plateau potential is stable

until the subcritical Hopf bifurcation HB$_2$ (G_{Kdr} = 6.34 nS) is reached, repetitive firing can be

observed between SN$_2$ (G_{Kdr} = 5.93 nS) and SN$_1$ (G_{Kdr} = 22.65 nS). The quiescent state is

stable from point HB$_1$ (G_{Kdr} = 17.59 nS) onward. (D) Two-parameters bifurcation diagram of

the model in the G_{Nap} - G_{Kdr} plane (I = 20 pA). The black merged curves indicate the

bifurcations HB$_1$ and HB$_2$. The red curves indicate the SN bifurcations of limit cycles SN$_1$ and

SN$_2$. The shaded area indicates the region where repetitive firing can occur. The oblique lines

through the points labeled 1, 2 and 3, the same as in B, correspond to three different values

of the ratio G_{Nap}/G_{Kdr} : 0.02 (gray), 0.12 (red) and 0.48 (blue). Voltage traces on the right

display the response to a 2 s current pulse when channel noise is taken into account for the

three regimes: quiescence (top, gray trace and dot in the diagram), repetitive firing (middle,

red) and plateau potential (bottom, blue). They correspond to the three deterministic

voltage traces shown in B. Note that the one-parameter bifurcation diagrams shown in B

correspond to horizontal lines through points 1 and 2 (G_{Kdr} = 10 nS) and through point 3

(G_{Kdr} = 2.5 nS), respectively. The bifurcation diagram in C corresponds to a vertical line

through point 2 and 3 (G_{Nap} = 1.2 nS). (E) Cumulative distribution function of the ratio

G_{Nap}/G_{Kdr} for the four clusters in A, showing the sequencing SS (gray) → RS (red) → ME

(purple, 3 cells only) → PP (blue) predicted by the two-parameters bifurcation diagram in D.

The wide PP range, as compared to SS and RS, merely comes from the fact that G_{Kdr} is small

for cells in this cluster. The three colored points indicate the slopes of the oblique lines

displayed in D (0.02, 0.12 and 0.48, respectively). (F) The data points in A are superimposed

on the two-parameters bifurcation diagram shown in D, demonstrating a good agreement

between our basic model and experimental data (the same color code as in A for the

different clusters). The bifurcation diagram is simplified compared to A, only the region

where repetitive spiking is possible (i.e. between the lines SN_1 and SN_2 in A) being displayed

(shaded area). Notice that 3 ME cells (purple dots) are located close to the transition

between the RS and PP regions. The four arrows indicate the presumable evolution of G_{Nap}

and G_{Kdr} for SS, RS, ME and PP cells between E12.5 and E14.5-15.5. G_{Nap} eventually

decreases while G_{Kdr} keeps on increasing. (G) Distribution of a sample of cells in the G_{Kdr} -

G_{Kdr} plane at E14.5. All the cells are located well within the SS region far from bifurcation

lines because of the decreased G_{Nap} compared to E12.5, the increased G_{Kdr}, and the shift of

the RS region (shaded) due to capacitance increase (18 versus 13 pF).

1001

Figure 8. Effects of the slow inactivation of I_{Nap} on firing patterns predicted by

computational modeling

(A) Examples of repetitive plateaus (left) and mixed events (right) recorded in V1[R] at E12.5

during a 2 s current pulse. (B1) Current-voltage curve of the basic model (without slow

inactivation of I_{Nap}, and without I_A or channel noise) for G_{Kdr} = 5 nS and for G_{Nap} = 1.65 nS

(lower curve) and 2 nS (upper curve). Solid lines denote stable fixed points and dashed lines

unstable ones. For G_{Nap} =1.65 nS, bistability between quiescence and plateau occurs

between 1.39 and 10.48 pA. When G_{Nap} is increased to 2 nS, the bistability region ranges

from -10.84 to 9.70 pA, thus extending into the negative current range. This implies that

once a plateau has been elicited, the model will stay in that stable state and not return to

the resting state, even though current injection is switched off (see insert). B1 Insert. Voltage

response to a 2 s current pulse of 15 pA for G_{Nap} = 2 nS. The resting state (gray dot on the

lower curve in B1 is destabilized at pulse onset and a plateau is elicited (blue dot on the

upper curve in B1). At pulse offset, the plateau is maintained, even though the injected

current is brought back to zero, and channel noise is not sufficient to go back to the resting

state. (B2) Domain of bistability between quiescence and plateau (shaded) in the I-

G_{Nap} plane for G_{Kdr}= 5 nS. It is delimited by the line SN$_2$ where a SN bifurcation of fixed

points occurs and by the subcritical Hopf bifurcation line HB where the plateau becomes

unstable. Bistability requires that G_{Nap} exceeds 1.35 nS, and the domain of bistability

enlarges as G_{Nap} is increased further. The two horizontal lines correspond to the two cases

shown in B1: G_{Nap} =1.65 nS and 2 nS. (C) Behavior of the model when slow inactivation is

incorporated. The bifurcation diagram of the basic model (without slow inactivation) for I =

10 pA and G_{Kdr} = 5 nS (same conventions as in Fig 7B) and the stable limit cycle (black solid

curve) obtained when slow inactivation is added are superimposed. The limit cycle is

comprised of four successive phases (see labels): 1) long plateau during which I_{Nap} slowly

inactivates, 2) fast transition to the quiescent state, 3) repolarization episode during which

I_{Nap} slowly deinactivates, 4) fast transition back to the plateau. Each plateau starts with a

full-blown action potential followed by rapidly decaying spikelets. Note that the bifurcation

HB is subcritical here (unstable limit cycle shown by dashed red curve), at variance with

square wave bursting (supercritical bifurcation and stable limit cycle); this is a characteristic

feature of pseudo-plateau bursting. Note also that the plateau extends beyond the

bifurcation HB because it is only weakly unstable then. Responses to a 15 s current pulse are

shown on the right side. Top left: voltage response, Top right: behavior of the "effective"

conductance of I_{Nap} channel, i.e., the maximal conductance G_{Nap} multiplied by the slow

inactivation variable s. Bottom left: voltage trace when channel noise is added to fast and

slow gating variables, Bottom right: Voltage trace when G_{Nap} is increased by 50%. (D) Mixed

events. The bifurcation diagram of the basic model for G_{Kdr} = 5 nS and I = 12 pA and the

stable limit cycle obtained in the presence of slow inactivation (G_{Nap} = 2.5 nS) are

superimposed. Here again, the limit cycle is comprised of four successive phases (see labels):

1) slow inactivation of I_{Nap} that leads to the crossing of the bifurcation point HB_2 and then

to the destabilization of the plateau potential, 2) fast transition to the spiking regime, 3)

repetitive spiking during which I_{Nap} slowly de-inactivates, which leads to the crossing of the

bifurcation point SN_2 and terminates the spiking episode, 4) fast transition back to the stable

plateau potential. Response to a 15 s current pulse of 12 pA is shown on the right in the

absence of any channel noise. Top: Voltage trace (same labels as in the bifurcation diagram

on the left), Bottom: Variations of the "effective" conductance $G_{Nap}s$ (same labels as in the

voltage trace). Note that de-inactivation sufficient to trigger a new plateau occurs over a

series of successive spikes, hence the small oscillations are visible on the trace. Note also

that in C and D the first plateau lasts longer than the following ones, as in

electrophysiological recordings of embryonic V1[R] cells displaying repetitive plateaus. This

form of adaptation is caused by the slow inactivation of the persistent sodium current.

1053

References

1054

1. O'Donovan MJ. The origin of spontaneous activity in developing networks of the

vertebrate nervous system. Curr Opin Neurobiol. 1999;9(1):94-104. PubMed PMID:

10072366.

2. Saint-Amant L. Development of motor rhythms in zebrafish embryos. Progress in

brain research. 2010;187:47-61. doi: 10.1016/B978-0-444-53613-6.00004-6. PubMed PMID:

21111200.

3. Blankenship AG, Feller MB. Mechanisms underlying spontaneous patterned activity

in developing neural circuits. Nature reviews Neuroscience. 2010;11(1):18-29. doi:

10.1038/nrn2759. PubMed PMID: 19953103; PubMed Central PMCID: PMC2902252.

4. Myers CP, Lewcock JW, Hanson MG, Gosgnach S, Aimone JB, Gage FH, et al.

Cholinergic input is required during embryonic development to mediate proper assembly of

spinal locomotor circuits. Neuron. 2005;46(1):37-49. Epub 2005/04/12. doi: S0896-

6273(05)00165-0 [pii]

10.1016/j.neuron.2005.02.022. PubMed PMID: 15820692.

5. Milner LD, Landmesser LT. Cholinergic and GABAergic inputs drive patterned

spontaneous motoneuron activity before target contact. J Neurosci. 1999;19(8):3007-22.

PubMed PMID: 10191318.

6. Hanson MG, Landmesser LT. Characterization of the circuits that generate

spontaneous episodes of activity in the early embryonic mouse spinal cord. J Neurosci.

2003;23(2):587-600. PubMed PMID: 12533619.

7. Momose-Sato Y, Sato K. Large-scale synchronized activity in the embryonic

brainstem and spinal cord. Frontiers in cellular neuroscience. 2013;7:36. doi:

10.3389/fncel.2013.00036. PubMed PMID: 23596392; PubMed Central PMCID:

PMC3625830.

8. Khazipov R, Luhmann HJ. Early patterns of electrical activity in the developing

cerebral cortex of humans and rodents. Trends in neurosciences. 2006;29(7):414-8. doi:

10.1016/j.tins.2006.05.007. PubMed PMID: 16713634.

9. Allain AE, Le Corronc H, Delpy A, Cazenave W, Meyrand P, Legendre P, et al.

Maturation of the GABAergic transmission in normal and pathologic motoneurons. Neural

plasticity. 2011;2011:905624. doi: 10.1155/2011/905624. PubMed PMID: 21785735;

PubMed Central PMCID: PMC3140191.

10. Branchereau P, Chapron J, Meyrand P. Descending 5-hydroxytryptamine raphe inputs

repress the expression of serotonergic neurons and slow the maturation of inhibitory systems

in mouse embryonic spinal cord. J Neurosci. 2002;22(7):2598-606. doi: 20026199. PubMed
PMID: 11923425.

11.   Yvert B, Branchereau P, Meyrand P. Multiple spontaneous rhythmic activity patterns
generated by the embryonic mouse spinal cord occur within a specific developmental time
window. Journal of neurophysiology. 2004;91(5):2101-9. doi: 10.1152/jn.01095.2003.
PubMed PMID: 14724265.

12.   Czarnecki A, Le Corronc H, Rigato C, Le Bras B, Couraud F, Scain AL, et al.
Acetylcholine controls GABA-, glutamate-, and glycine-dependent giant depolarizing
potentials that govern spontaneous motoneuron activity at the onset of synaptogenesis in the
mouse   embryonic   spinal   cord.   J   Neurosci.   2014;34(18):6389-404.   doi:
10.1523/JNEUROSCI.2664-13.2014. PubMed PMID: 24790209.

13.   Pun S, Sigrist M, Santos AF, Ruegg MA, Sanes JR, Jessell TM, et al. An intrinsic
distinction in neuromuscular junction assembly and maintenance in different skeletal muscles.
Neuron.   2002;34(3):357-70.   doi: 10.1016/s0896-6273(02)00670-0.   PubMed   PMID:
11988168.

14.   Angelim M, Maia L, Mouffle C, Ginhoux F, Low D, Amancio-Dos-Santos A, et al.
Embryonic macrophages and microglia ablation alter the development of dorsal root ganglion
sensory neurons in mouse embryos. Glia. 2018;66(11):2470-86. doi: 10.1002/glia.23499.
PubMed PMID: 30252950.

15.   Marmigere F, Ernfors P. Specification and connectivity of neuronal subtypes in the
sensory lineage. Nature reviews Neuroscience. 2007;8(2):114-27. doi: 10.1038/nrn2057.
PubMed PMID: 17237804.

16.   Ozaki S, Snider WD. Initial trajectories of sensory axons toward laminar targets in the
developing mouse spinal cord. The Journal of comparative neurology. 1997;380(2):215-29.
PubMed PMID: 9100133.

17.   Ballion B, Branchereau P, Chapron J, Viala D. Ontogeny of descending serotonergic
innervation and evidence for intraspinal 5-HT neurons in the mouse spinal cord. Brain
research   Developmental   brain   research.   2002;137(1):81-8.   doi: 10.1016/s0165-
3806(02)00414-5. PubMed PMID: 12128257.

18.   Moody WJ, Bosma MM. Ion channel development, spontaneous activity, and activity-
dependent development in nerve and muscle cells. Physiological reviews. 2005;85(3):883-941.
doi: 10.1152/physrev.00017.2004. PubMed PMID: 15987798.

19.   Spitzer   NC.   Electrical   activity   in   early   neuronal   development.   Nature.
2006;444(7120):707-12. doi: 10.1038/nature05300. PubMed PMID: 17151658.

20. Katz LC, Shatz CJ. Synaptic activity and the construction of cortical circuits. Science.
1996;274(5290):1133-8. PubMed PMID: 8895456.

21. Hanson MG, Milner LD, Landmesser LT. Spontaneous rhythmic activity in early
chick spinal cord influences distinct motor axon pathfinding decisions. Brain Res Rev.
2008;57(1):77-85. Epub 2007/10/09. doi: S0165-0173(07)00127-0 [pii]
10.1016/j.brainresrev.2007.06.021. PubMed PMID: 17920131.

22. Hanson MG, Landmesser LT. Normal patterns of spontaneous activity are required for
correct motor axon guidance and the expression of specific guidance molecules. Neuron.
2004;43(5):687-701. doi: 10.1016/j.neuron.2004.08.018. PubMed PMID: 15339650.

23. Hanson MG, Landmesser LT. Increasing the frequency of spontaneous rhythmic
activity disrupts pool-specific axon fasciculation and pathfinding of embryonic spinal
motoneurons. J Neurosci. 2006;26(49):12769-80. doi: 10.1523/JNEUROSCI.4170-06.2006.
PubMed PMID: 17151280.

24. Benito-Gonzalez A, Alvarez FJ. Renshaw cells and Ia inhibitory interneurons are
generated at different times from p1 progenitors and differentiate shortly after exiting the cell
cycle. J Neurosci. 2012;32(4):1156-70. doi: 10.1523/JNEUROSCI.3630-12.2012. PubMed
PMID: 22279202; PubMed Central PMCID: PMC3276112.

25. Boeri J, Le Corronc H, Lejeune FX, Le Bras B, Mouffle C, Angelim M, et al.
Persistent Sodium Current Drives Excitability of Immature Renshaw Cells in Early
Embryonic Spinal Networks. J Neurosci. 2018;38(35):7667-82. doi:
10.1523/JNEUROSCI.3203-17.2018. PubMed PMID: 30012693.

26. Perry S, Gezelius H, Larhammar M, Hilscher MM, Lamotte d'Incamps B, Leao KE, et
al. Firing properties of Renshaw cells defined by Chrna2 are modulated by hyperpolarizing
and small conductance ion currents Ih and ISK. The European journal of neuroscience.
2015;41(7):889-900. doi: 10.1111/ejn.12852. PubMed PMID: 25712471.

27. Bikoff JB, Gabitto MI, Rivard AF, Drobac E, Machado TA, Miri A, et al. Spinal
Inhibitory Interneuron Diversity Delineates Variant Motor Microcircuits. Cell.
2016;165(1):207-19. doi: 10.1016/j.cell.2016.01.027. PubMed PMID: 26949184; PubMed
Central PMCID: PMC4808435.

28. Delpy A, Allain AE, Meyrand P, Branchereau P. NKCC1 cotransporter inactivation
underlies embryonic development of chloride-mediated inhibition in mouse spinal
motoneuron. J Physiol. 2008;586(4):1059-75. doi: 10.1113/jphysiol.2007.146993. PubMed
PMID: 18096599; PubMed Central PMCID: PMC2375629.

29. Crill WE. Persistent sodium current in mammalian central neurons. Annual review of
physiology. 1996;58:349-62. doi: 10.1146/annurev.ph.58.030196.002025. PubMed PMID:
8815799.

30. Sillar KT, Simmers AJ, Wedderburn JF. The post-embryonic development of cell
properties and synaptic drive underlying locomotor rhythm generation in Xenopus larvae.
Proceedings Biological sciences. 1992;249(1324):65-70. doi: 10.1098/rspb.1992.0084.
PubMed PMID: 1359549.

31. Gao BX, Ziskind-Conhaim L. Development of ionic currents underlying changes in
action potential waveforms in rat spinal motoneurons. Journal of neurophysiology.
1998;80(6):3047-61. PubMed PMID: 9862905.

32. Gao H, Lu Y. Early development of intrinsic and synaptic properties of chicken
nucleus laminaris neurons. Neuroscience. 2008;153(1):131-43. doi:
10.1016/j.neuroscience.2008.01.059. PubMed PMID: 18355968.

33. McKay BE, Turner RW. Physiological and morphological development of the rat
cerebellar Purkinje cell. J Physiol. 2005;567(Pt 3):829-50. doi: 10.1113/jphysiol.2005.089383.
PubMed PMID: 16002452; PubMed Central PMCID: PMC1474219.

34. Liu X, Pfaff DW, Calderon DP, Tabansky I, Wang X, Wang Y, et al. Development of
Electrophysiological Properties of Nucleus Gigantocellularis Neurons Correlated with
Increased CNS Arousal. Developmental neuroscience. 2016;38(4):295-310. doi:
10.1159/000449035. PubMed PMID: 27788521; PubMed Central PMCID: PMC5127753.

35. Pineda R, Ribera A. Evolution of the Action Potential. In: Kaas JH, editor. Evolution
of Nervous Systems. 1: Elsevier Ltd; 2010. p. 211-38.

36. Coetzee WA, Amarillo Y, Chiu J, Chow A, Lau D, McCormack T, et al. Molecular
diversity of K+ channels. Annals of the New York Academy of Sciences. 1999;868:233-85.
doi: 10.1111/j.1749-6632.1999.tb11293.x. PubMed PMID: 10414301.

37. O'Leary T, Williams AH, Caplan JS, Marder E. Correlations in ion channel expression
emerge from homeostatic tuning rules. Proceedings of the National Academy of Sciences of
the United States of America. 2013;110(28):E2645-54. Epub 2013/06/27. doi:
10.1073/pnas.1309966110. PubMed PMID: 23798391; PubMed Central PMCID:
PMCPMC3710808.

38. Taylor AL, Goaillard JM, Marder E. How multiple conductances determine
electrophysiological properties in a multicompartment model. J Neurosci. 2009;29(17):5573-
86. Epub 2009/05/01. doi: 10.1523/jneurosci.4438-08.2009. PubMed PMID: 19403824;
PubMed Central PMCID: PMCPMC2821064.

39. Alonso LM, Marder E. Visualization of currents in neural models with similar
behavior and different conductance densities. eLife. 2019;8. Epub 2019/02/01. doi:
10.7554/eLife.42722. PubMed PMID: 30702427; PubMed Central PMCID:
PMCPMC6395073.

40. Sigworth FJ, Sine SM. Data transformations for improved display and fitting of
single-channel dwell time histograms. Biophysical journal. 1987;52(6):1047-54. doi:
10.1016/S0006-3495(87)83298-8. PubMed PMID: 2447968; PubMed Central PMCID:
PMC1330104.

41. Ramoa AS, McCormick DA. Developmental changes in electrophysiological
properties of LGNd neurons during reorganization of retinogeniculate connections. J Neurosci.
1994;14(4):2089-97. PubMed PMID: 8158259; PubMed Central PMCID: PMC6577110.

42. Belleau ML, Warren RA. Postnatal development of electrophysiological properties of
nucleus accumbens neurons. Journal of neurophysiology. 2000;84(5):2204-16. doi:
10.1152/jn.2000.84.5.2204. PubMed PMID: 11067966.

43. Picken Bahrey HL, Moody WJ. Early development of voltage-gated ion currents and
firing properties in neurons of the mouse cerebral cortex. Journal of neurophysiology.
2003;89(4):1761-73. doi: 10.1152/jn.00972.2002. PubMed PMID: 12611962.

44. Tong H, McDearmid JR. Pacemaker and plateau potentials shape output of a
developing locomotor network. Current biology : CB. 2012;22(24):2285-93. doi:
10.1016/j.cub.2012.10.025. PubMed PMID: 23142042; PubMed Central PMCID:
PMC3525839.

45. Alvarez FJ, Benito-Gonzalez A, Siembab VC. Principles of interneuron development
learned from Renshaw cells and the motoneuron recurrent inhibitory circuit. Annals of the
New York Academy of Sciences. 2013;1279:22-31. doi: 10.1111/nyas.12084. PubMed PMID:
23530999; PubMed Central PMCID: PMC3870136.

46. Allain AE, Segu L, Meyrand P, Branchereau P. Serotonin controls the maturation of
the GABA phenotype in the ventral spinal cord via 5-HT1b receptors. Annals of the New
York Academy of Sciences. 2010;1198:208-19. doi: 10.1111/j.1749-6632.2010.05433.x.
PubMed PMID: 20536936.

47. Branchereau P, Morin D, Bonnot A, Ballion B, Chapron J, Viala D. Development of
lumbar rhythmic networks: from embryonic to neonate locomotor-like patterns in the mouse.
Brain research bulletin. 2000;53(5):711-8. doi: 10.1016/s0361-9230(00)00403-2. PubMed
PMID: 11165805.

48.   Spitzer NC, Vincent A, Lautermilch NJ. Differentiation of electrical excitability in
motoneurons. Brain research bulletin. 2000;53(5):547-52. PubMed PMID: 11165790.

49.   Prasad T, Wang X, Gray PA, Weiner JA. A differential developmental pattern of
spinal interneuron apoptosis during synaptogenesis: insights from genetic analyses of the
protocadherin-gamma gene cluster. Development. 2008;135(24):4153-64. doi:
10.1242/dev.026807. PubMed PMID: 19029045; PubMed Central PMCID: PMC2755264.

50.   Taddese A, Bean BP. Subthreshold sodium current from rapidly inactivating sodium
channels drives spontaneous firing of tuberomammillary neurons. Neuron. 2002;33(4):587-
600. doi: 10.1016/s0896-6273(02)00574-3. PubMed PMID: 11856532.

51.   Kuo JJ, Lee RH, Zhang L, Heckman CJ. Essential role of the persistent sodium current
in spike initiation during slowly rising inputs in mouse spinal neurones. J Physiol.
2006;574(Pt 3):819-34. doi: 10.1113/jphysiol.2006.107094. PubMed PMID: 16728453;
PubMed Central PMCID: PMC1817738.

52.   Teka W, Tsaneva-Atanasova K, Bertram R, Tabak J. From plateau to pseudo-plateau
bursting: making the transition. Bulletin of mathematical biology. 2011;73(6):1292-311. doi:
10.1007/s11538-010-9559-7. PubMed PMID: 20658200; PubMed Central PMCID:
PMC3152987.

53.   Bertram R, Butte MJ, Kiemel T, Sherman A. Topological and phenomenological
classification of bursting oscillations. Bulletin of mathematical biology. 1995;57(3):413-39.
doi: 10.1007/BF02460633. PubMed PMID: 7728115.

54.   Izhikevich EM. Neural excitability, spiking and bursting. Int J Bifurcation Chaos.
2000;10(06):1171-266.

55.   Borisyuk A, Rinzel J. Understanding neuronal dynamics by geometrical dissection of
minimal models. In: Chow C, Gutkin B, Hansel D, Meunier C, Dalibard J, editors. Models
and Methods in Neurophysics. Proc Les Houches Summer School 2003, (Session LXXX):
Elsevier; 2005. p. 19-72.

56.   Rinzel J. Bursting oscillations in an excitable membrane model. In: Sleeman B, Jarvis
R, editors. Ordinary and Partial Differential Equations Lecture Notes in Mathematics. 1151.
Berlin, Heidelberg: Springer; 1985. p. 304–16.

57.   Stern JV, Osinga HM, LeBeau A, Sherman A. Resetting behavior in a model of
bursting in secretory pituitary cells: distinguishing plateaus from pseudo-plateaus. Bulletin of
mathematical biology. 2008;70(1):68-88. doi: 10.1007/s11538-007-9241-x. PubMed PMID:
17703340.

58.   Osinga HM, Tsaneva-Atanasova KT. Dynamics of plateau bursting depending on the

location of its equilibrium. Journal of neuroendocrinology. 2010;22(12):1301-14. doi:

10.1111/j.1365-2826.2010.02083.x. PubMed PMID: 20955345.

59.   Osinga HM, Sherman A, Tsaneva-Atanasova K. Cross-Currents between Biology and

Mathematics: The Codimension of Pseudo-Plateau Bursting. Discrete and continuous

dynamical systems Series A. 2012;32(8):2853-77. doi: 10.3934/dcds.2012.32.2853. PubMed

PMID: 22984340; PubMed Central PMCID: PMC3439852.

60.   Izhikevich EM. Subcritical Elliptic Bursting of Bautin Type. SIAM Journal on

Applied Mathematics. 2000;60(2):503-35. doi: https://doi.org/10.1137/S003613999833263X.

61.   Su J, Rubin J, Terman D. Effects of noise on elliptic bursters. Nonlinearity.

2004;17(1):133-57. doi: https://doi.org/10.1088/0951-7715/17/1/009.

62.   Stam FJ, Hendricks TJ, Zhang J, Geiman EJ, Francius C, Labosky PA, et al. Renshaw

cell interneuron specialization is controlled by a temporally restricted transcription factor

program. Development. 2012;139(1):179-90. doi: 10.1242/dev.071134. PubMed PMID:

22115757; PubMed Central PMCID: PMC3231776.

63.   Eccles JC, Fatt P, Landgren S. The inhibitory pathway to motoneurones. Progress in

neurobiology. 1956;(2):72-82. PubMed PMID: 13441782.

64.   Vinay L, Brocard F, Clarac F. Differential maturation of motoneurons innervating

ankle flexor and extensor muscles in the neonatal rat. The European journal of neuroscience.

2000;12(12):4562-6. doi: 10.1046/j.0953-816x.2000.01321.x. PubMed PMID: 11122369.

65.   Durand J, Filipchuk A, Pambo-Pambo A, Amendola J, Borisovna Kulagina I,

Gueritaud JP. Developing electrical properties of postnatal mouse lumbar motoneurons.

Frontiers in cellular neuroscience. 2015;9:349. doi: 10.3389/fncel.2015.00349. PubMed

PMID: 26388736; PubMed Central PMCID: PMC4557103.

66.   Marcotti W, Johnson SL, Holley MC, Kros CJ. Developmental changes in the

expression of potassium currents of embryonic, neonatal and mature mouse inner hair cells. J

Physiol. 2003;548(Pt 2):383-400. doi: 10.1113/jphysiol.2002.034801. PubMed PMID:

12588897; PubMed Central PMCID: PMC2342842.

67.   Marcotti W, Johnson SL, Rusch A, Kros CJ. Sodium and calcium currents shape

action potentials in immature mouse inner hair cells. J Physiol. 2003;552(Pt 3):743-61. doi:

10.1113/jphysiol.2003.043612. PubMed PMID: 12937295; PubMed Central PMCID:

PMC2343463.

68.   Sontheimer H, Trotter J, Schachner M, Kettenmann H. Channel expression correlates

with differentiation stage during the development of oligodendrocytes from their precursor

cells in culture. Neuron. 1989;2(2):1135-45. doi: 10.1016/0896-6273(89)90180-3. PubMed PMID: 2560386.

69.   Allain AE, Bairi A, Meyrand P, Branchereau P. Ontogenic changes of the GABAergic system in the embryonic mouse spinal cord. Brain research. 2004;1000(1-2):134-47. doi: 10.1016/j.brainres.2003.11.071. PubMed PMID: 15053961.

70.   Sapir T, Geiman EJ, Wang Z, Velasquez T, Mitsui S, Yoshihara Y, et al. Pax6 and engrailed 1 regulate two distinct aspects of renshaw cell development. J Neurosci. 2004;24(5):1255-64. doi: 10.1523/JNEUROSCI.3187-03.2004. PubMed PMID: 14762144; PubMed Central PMCID: PMC2997484.

71.   Henley J, Poo MM. Guiding neuronal growth cones using Ca2+ signals. Trends in cell biology. 2004;14(6):320-30. doi: 10.1016/j.tcb.2004.04.006. PubMed PMID: 15183189; PubMed Central PMCID: PMC3115711.

72.   Tsaneva-Atanasova K, Osinga HM, Riess T, Sherman A. Full system bifurcation analysis of endocrine bursting models. Journal of theoretical biology. 2010;264(4):1133-46. doi: 10.1016/j.jtbi.2010.03.030. PubMed PMID: 20307553; PubMed Central PMCID: PMC3128456.

73.   Tagliavini A, Tabak J, Bertram R, Pedersen MG. Is bursting more effective than spiking in evoking pituitary hormone secretion? A spatiotemporal simulation study of calcium and granule dynamics. American journal of physiology Endocrinology and metabolism. 2016;310(7):E515-25. doi: 10.1152/ajpendo.00500.2015. PubMed PMID: 26786781.

74.   Oster A, Faure P, Gutkin BS. Mechanisms for multiple activity modes of VTA dopamine neurons. Frontiers in computational neuroscience. 2015;9:95. doi: 10.3389/fncom.2015.00095. PubMed PMID: 26283955; PubMed Central PMCID: PMC4516885.

75.   Chevalier M, Toporikova N, Simmers J, Thoby-Brisson M. Development of pacemaker properties and rhythmogenic mechanisms in the mouse embryonic respiratory network. eLife. 2016;5. doi: 10.7554/eLife.16125. PubMed PMID: 27434668; PubMed Central PMCID: PMC4990420.

76.   Kole MH, Stuart GJ. Signal processing in the axon initial segment. Neuron. 2012;73(2):235-47. doi: 10.1016/j.neuron.2012.01.007. PubMed PMID: 22284179.

77.   Tamamaki N, Yanagawa Y, Tomioka R, Miyazaki J, Obata K, Kaneko T. Green fluorescent protein expression and colocalization with calretinin, parvalbumin, and somatostatin in the GAD67-GFP knock-in mouse. The Journal of comparative neurology. 2003;467(1):60-79. doi: 10.1002/cne.10905. PubMed PMID: 14574680.

78.   Scain AL, Le Corronc H, Allain AE, Muller E, Rigo JM, Meyrand P, et al. Glycine

release from radial cells modulates the spontaneous activity and its propagation during early

spinal cord development. J Neurosci. 2010;30(1):390-403. Epub 2010/01/08. doi: 30/1/390

[pii]

10.1523/JNEUROSCI.2115-09.2010. PubMed PMID: 20053920.

79.   Huang H, Trussell LO. Control of presynaptic function by a persistent Na(+) current.

Neuron. 2008;60(6):975-9. doi: 10.1016/j.neuron.2008.10.052. PubMed PMID: 19109905;

PubMed Central PMCID: PMC2657474.

80.   Rousseeuw PJ. Silhouettes - a Graphical Aid to the Interpretation and Validation of

Cluster-Analysis. J Comput Appl Math. 1987;20:53-65. doi: Doi 10.1016/0377-

0427(87)90125-7. PubMed PMID: WOS:A1987L111800005.

81.   Ermentrout B. Simulating, Analyzing, and Animating Dynamical Systems: A Guide to

XPPAUT for Researchers and Students. Philadelphia: Society for Industrial and Applied

Mathematics; 2002.

82.   Gillespie DT. A general method for numerically simulating the stochastic time

evolution of coupled chemical reactions. Journal of Computational Physics. 1976;22(4):403-

34. doi: https://doi.org/10.1016/0021-9991(76)90041-3.

83.   Witelski T, Bowen M. Fast/slow Dynamical Systems. In: Methods of Mathematical

Modelling: Springer, Cham; 2015.

1342

1343

Table 1

Parameter	Basic model	Model with slow inactivation of I_{Nap}
Passive parameters		
Input conductance G_{in}	1 nS	same
Input capacitance C_{in}	13 pF (E12.5, Figs. 7B, C, D and F and 8B to D) or 18 pF (E14.5, Fig. 7G)	13 pF
Resting potential V_r	-60 mV	same
Injected current I	20 pA (Fig. 7B to G)	10 pA (Fig. 8C) or 12 pA (Fig. 8D) variable in Fig. 8B
Transient sodium current I_{nat}		
Maximal conductance G_{Nat}	20 nS	same
Reversal potential E_{Na}	60 mV	
Activation exponent	3	
Mid-activation V_m	-26 mV	
Steepness of activation k_m	9.5 mV	
Activation time constant	1.5 ms	
Mid-inactivation V_h	-45 mV	
Steepness of inactivation K_h	-5 mV	
Inactivation time constant τ_m	Voltage-dependent (see Material and Methods)	
Persistent sodium current I_{Nap}		
Maximal conductance	variable (see text and figure captions)	same
Mid-activation voltage	-36 mV	same
Mid-inactivation V_s		-30 mV
Steepness of inactivation k_s		-5 mV
Inactivation time constant	Slow inactivation not included	2 s
Delayed rectifier potassium current I_{Kdr}		
Maximal conductance G_{Kdr}	variable (see text and figure captions)	same
Reversal potential E_K	-96 mV	
Activation exponent	3	
Mid-activation V_n	-20 mV	
Steepness of activation k_n	15 mV	
Activation time constant τ_m	10 ms	
Potassium A current I_A (when included in the basic model)		
Maximal conductance G_A	Equal to G_{Kdr}	never included
Mid-activation V_{mA}	-30 mV	
Steepness of activation k_{mA}	12 mV	
Activation time constant	Instantaneous activation	
Mid-inactivation V_{hA}	-70 mV	
Steepness of inactivation k_{hA}	-7 mV	
Inactivation time constant τ_{hA}	23 ms	

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7

Fig 8

Supplementary figures

Figure 2—figure supplement 1. Effect of 4-AP on I_{Kdr} and I_A in embryonic V1[R]
(A1) Example of voltage-dependent potassium currents evoked in response to 10 mV membrane potential steps (200 ms) from -100 mV or from -30 mV to +40 mV (10 s interval between pulses). V1[R] were voltage clamped at V_H = -60 mV. A prepulse of -40 mV (300 ms) was applied to activate both I_A and I_{Kdr}. I_{Kdr} was evoked in response to 10 mV membrane potential steps (200 ms) from -100 mV to +40 mV. V1[R] were voltage clamped at V_H = -60 mV. A prepulse of 30 mV (V_H = - 30 mV) was applied to isolate I_{Kdr}. (A1) Representative example of the effect of 300 µM 4-AP application on I_{Kdr} recorded from embryonic V1[R] at E12.5. (B1) Curves showing current-voltage relationships of I_{Kdr} in control and in the presence of 300 µM 4-AP. Measurements were performed on traces shown in A1. (C1) Dose-response relationship of 4-AP-evoked I_{Kdr} inhibition (mean + SE). Data were normalized to I_{Kdr} amplitude measured in the absence of 4-AP (V_H = 40mV) and fitted as explained in Materials and Methods. Note that 4-AP IC$_{50}$ is in µM range (2.9 µM). 0.3 µM 4-AP n = 3; N = 3, 1 µM 4-AP n = 3; N = 3, 3 µM 4-AP n = 9; N = 9, 10 µM 4-AP n = 13; N = 13, 30 µM 4-AP n = 7; N = 7, 100 µM 4-AP n = 7; N = 7, 300 µM 4-AP n = 7; N = 7. (A2) I_A was obtained as the difference between currents evoked from V_H = -100 mV and currents evoked from V_H = -30 mV (10 mV voltage step). (A2) Representative example of the effect of 300 µM 4-AP on I_A in V1[R] recorded at E12.5. (B2) I_A Current-voltage ($I - V$) relationship in control conditions and in the presence of 300 µM 4-AP. The $I - V$ curves were obtained from the traces shown in A1. (C2) Bar graph showing the percentage of I_A block elicited by 4-AP. Note that 4-AP did not significantly block I_A (Wilcoxon test P = 0.065, n = 10).

Figure 2—figure supplement 2. Relates to Fig 2. Effect of 4-AP application in repetitively spiking V1[R] at E12.5

(A) Representative traces showing the effect of 4-AP application (300 µM) on Repetitive Spiking (RS) V1[R] at E12.5. Note that plateau potential activity evoked in the presence of 4-AP (middle trace) was blocked by 0.5 µM TTX (right trace). (B) Bar plots showing the changes in the firing pattern of RS V1[R] evoked by 300 µM 4-AP application (n = 14). 4-AP application evoked a plateau potential in 71.4 % of the recorded neurons (10/14) and mixed events in 14.3% of the recorded neurons (2/14). The excitability pattern was not modified in 2 neurons. Repetitive Spiking (RS) V1[R] (red), Mixed events (ME) V1[R] (purple), Plateau Potential (PP) V1[R] (blue).

Figure 3—figure supplement 1. Distributions of log ½Ad, CV ½Ad and log ddr values related to the cluster analysis of embryonic V1[R] firing patterns

(A1) Histogram of log mean ½Ad (mean half amplitude event duration) for the whole V1[R] population at E12.5 (n= 164; bin width 0.1). The histogram was well fitted by the sum of three Gaussian curves with means and SDs of 1.135, 2.046 & 2.84, and 0.316, 0.181 & 0.21, respectively. (A2) Histogram of the values of log mean ½Ad sorted after cluster analysis showing single spiking (SS) V1[R] (gray), repetitive spiking (RS) V1[R] (red), mixed events (ME) V1[R] with short plateau potentials (ME short PP V1[R], light purple), ME V1[R] with long plateau potentials (ME long PP V1[R], dark purple) and plateau potential (PP) V1[R] (blue). log mean ½Ad was significantly different between SS V1[R], PP V1[R], the whole ME V1[R] population (ME$_s$ and ME$_l$ V1[R]) and PP V1[R] (Kruskall-Wallis test $P < 0.0001$; SS V1[R] versus RS V1[R], $P < 0.0001$; SS V1[R] versus ME V1[R], $P < 0.0001$; SS V1[R] versus PP V1[R], $P < 0.0001$; RS V1[R] versus ME V1[R], $P = 0.0004$; RS V1[R] versus PP V1[R], $P < 0.0001$; ME V1[R] versus PP V1[R], $P = 0.018$; SS V1[R] n = 46, RS V1[R] n = 69, ME$_s$ V1[R] n = 9, ME$_l$ V1[R] n = 4, PP V1[R] n = 35). (B1) Histogram of CV ½Ad for the whole V1[R] population at E12.5 (n= 164; bin width 5%). Note that a large population of V1[R] had zero CV ½Ad (n = 83). The histogram for CV ½Ad ≠ 0 was fitted by the sum of three Gaussian curves with means and SDs of 23.4, 68.4 & 117 (%) and 8.9, 6.8 & 4.1, respectively. (B2) Histograms of the values of CV ½Ad sorted after cluster analysis showing SS V1[R] (black), RS V1[R] (red), ME$_s$ V1[R] (light purple), ME$_l$ V1[R] (dark purple) and PP V1[R]. CV ½Ad was not significantly different between SS V1[R] and PP V1[R] (CV ½Ad of SS V1[R] and PP V1[R] = 0.682 % and 0% respectively: only one of the 46 SS V1[R] displayed 3 PA and had a CV ½Ad of 31.37). CV ½Ad was significantly different between RS V1[R] and the whole ME V1[R] population and

also between SS V1^R or PP V1^R and RS V1^R or ME V1^R (Kruskall-Wallis test $P < 0.0001$; SS V1^R versus RS V1^R $P < 0.0001$, SS V1^R versus ME V1^R $P < 0.0001$, SS V1^R versus PP V1^R $P = 0.846$, RS V1^R versus ME V1^R $P = 0.0003$, RS V1^R versus PP V1^R $P < 0.0001$, ME V1^R versus PP V1^R $P < 0.0001$). (C1) Histogram of log ddr (sum of ½Ad divided by pulse duration) for the whole V1^R population at E12.5 (n= 164; bin width 0.2). The histogram was fitted by the sum of two Gaussian curves with means and SDs of -2.51 & -0.851, and 0.2 & 0.46, respectively. (C2) Histograms of the values of log ddr sorted after cluster analysis showing SS V1^R (black), RS V1^R (red), ME$_s$ V1^R (light purple), ME$_l$ V1^R (dark purple) and PP V1^R. log (ddr) was not significantly different between ME V1^R and PP V1^R, while it was significantly different between SS V1^R and RS V1^R, SS V1^R and the whole ME V1^R population, SS V1^R and PP V1^R, RS V1^R and the whole ME V1^R population, RS V1^R and PP V1^R (Kruskall-Wallis test $P < 0.0001$; SS V1^R versus RS V1^R, $P < 0.0001$; SS V1^R versus ME V1^R, $P < 0.0001$; SS V1^R versus PP V1^R, $P < 0.0001$; RS V1^R versus ME V1^R, $P < 0.0001$; RS V1^R versus PP V1^R, $P < 0.0001$; ME V1^R versus PP V1^R, $P = 0.977$). ME$_s$ V1^R and ME$_l$ V1^R differed only by their CV ½Ad (Mann-Whitney test, log mean ½Ad for ME$_s$ V1^R versus log mean ½Ad for ME$_l$ V1^R, $P = 0.26$; CV ½Ad for ME$_s$ V1^R versus CV ½Ad ME$_l$ V1^R, $P = 0.0028$ and log ddr for ME$_s$ V1^R versus log ddr for ME$_l$ V1^R, $P = 0.1483$). It is noteworthy that the distribution of the values of each metric was multimodal thus indicating that each of them could partially discriminate different groups of embryonic V1^R according to their firing pattern.

Figure 6—figure supplement 1. I_{Nap} **is present in embryonic V1^R recorded at E14.5**

(A) Representative trace of I_{Nap} evoked by a slow depolarizing voltage ramp (70 mV/s, upper insert) in SS embryonic V1^R (lower insert). I_{Nap} was isolated by subtracting currents evoked by depolarizing ramps in the presence of 1 µM TTX to the control current evoked in the absence of TTX (upper insert). (B) Voltage dependence of G_{Nap} conductance calculated from the trace shown in A. The activation curve was obtained by transforming the current evoked by a depolarizing voltage ramp from -100 mV to 20 mV (70 mV/s) using the following equation: G_{NaP} = $-I_{Nap}/((-Vh)+E_{Na})$ where Vh is the holding potential at time t during a depolarizing voltage ramp and E_{Na} is the equilibrium potential for sodium (E_{Na} = 60 mV). The G_{NaP}/Vh curve was fitted with the following Boltzmann function: G = $G_{MAX}/(1+exp(-(V-V_{HALF})/k)))$ (Boeri et al. 2018), where V_{half} is the Vh value for G_{Nap} half activation, k the slope factor of the curve and G_{max} the maximum conductance. We found no significant difference between the values of V_{half} (Mann-Whitney test: P = 0.8518) and of k (Mann-Whitney test: P = 0.7546) obtained at E12.5 (Boeri et al. 2018) and those obtained at E14.5. At E14.5 V_{half} = -27 ± 5.1 mV and k = 7.73 ± 0.78 (n = 6).

Figure 6—figure supplement 2. I_{Kdr} was inhibited by 4-AP in V1^R recorded at E14.5
(A1) Representative examples of the total outward K$^+$ currents obtained from V_H = -100 mV (left traces), of I_{Kdr} (V_H = -30 mV, middle traces) and of isolated I_A (left traces) recorded in single spiking (SS) V1^R at E14.5. (A2) Current-voltage relationship of I_{Kdr} (filled circle) and of I_A (open circle) in SS V1^R at E14.5. $I - V$ curves were obtained from currents shown in A1. (B1) Representative example of the effect of 4-AP at 600 μM in V1^R at E14.5. (B2) Current-voltage curves in control condition and in the presence of 600 μM 4-AP. (B3) Bar plots showing the percentage of I_{Kdr} inhibition evoked by 300 μM 4-AP application (n = 8) and by 600 μM 4-AP application (n = 7). The percentages of I_{Kdr} inhibition evoked by 300 μM 4-AP and by 600 μM 4-AP applications were not significantly different (*P* = 0.574). (C1) Representative example of the effect of 600 μM 4-AP on I_A in V1^R recorded at E14.5. (C2) $I - V$ curves in control conditions and in the presence of 600 μM 4-AP. These curves were obtained from the traces shown in B1. (C3) Bar graph showing the percentage of I_A block elicited by 4-AP. 4-AP did not significantly block I_A (Wilcoxon test *P* = 0.11, n = 6).

Figure 7—figure supplement 1. Effects of I_A on embryonic V1^R firing patterns predicted by computational modeling

(A) The maximal conductances of I_{Kdr} and I_A at E12.5 are linearly correlated. Best fit: G_A = 1.09 G_{Kdr} (R^2 = 0.81, N=44). (B) Effect of I_A on the dynamics of the basic model. The one-parameter bifurcation diagrams in control condition (black, I = 20 pA, G_{Kdr} = 10 nS, no I_A, same as in Fig 7B) and with I_A added (orange, G_A = 10 nS) are superimposed. The I_A current shifts the firing threshold SN$_1$ to the right by 0.18 nS (see also C) as indicated by the orange arrow, with little effect on the amplitude of action potentials (see also insert in C). In contrast, I_A shifts SN$_2$ by only 0.03 nS because it is inactivated by depolarization. (C) I_A also slows down the discharge frequency, as shown by comparing the $G_{Nap} - V$ curves without I_A (black) and with I_A (orange). For G_{Nap} = 1 nS, for instance, the firing frequency is reduced by 31%, from 15 to 10.4 Hz. Here again, the effect of I_A progressively decreases as G_{Nap} increases because of the membrane depolarization elicited by I_{Nap}. For G_{Nap} = 2.4 nS, for instance, the firing frequency is reduced by 11% only, from 19.1 to 17 Hz. This frequency reduction elicited by I_A does not merely result from the increased firing threshold. Note also that the latency of the first spike is increased (see voltage trace in insert), which is a classical effect of I_A. (D) I_A reduces the frequency of pseudo-plateau bursting by lengthening quiescent episodes (doubling their duration in the example shown) without affecting the duration of plateaus much (here a mere 5% increase), as shown by the comparison of the voltage traces obtained without I_A (control, G_{Kdr} = 2.5 nS, black) and with I_A ($G_{Kdr} = G_A$ = 2.5 nS, orange). This is because I_A is activated near rest but inactivated during voltage plateaus. Note that increasing G_{Kdr}, in the absence of I_A has not the same effect; it shortens both plateaus and quiescent episodes (see Fig 8C, where G_{Kdr} = 5 nS). Again, this is because I_{Kdr} does not inactivate (or does it only very slowly), in contrast to I_A.

Figure 7—figure supplement 2. Explaining the effect of 4-AP on the firing pattern

The RS region of the basic model, where repetitive firing may occur, is displayed in the G_{Nap} $- G_{Kdr}$ plane in control condition for E12.5 V1^R (C_{in} = 13 pF, G_{in} = 1 nS, I = 20 pA, shaded area), and when G_{in} and I were both reduced by 25% (middle curve) or by 50% (left curve). The reduced I accounts for the decrease in rheobase, and thus in the current injected in the experiments, following the decrease in G_{in}. If 4-AP reduced only G_{Kdr} (as indicated by the downward arrow) the firing pattern of SS V1^R would not change, the RS region being too far to the right to be visited. In contrast, when the effects of 4-AP on the input conductance and rheobase are taken into account, the bifurcation diagram moves leftwards and downwards, as indicated by the oblique black arrow, and the RS and PP regions are then successively entered as G_{Kdr} is reduced. The same explanation holds at E14.5.