Reduced cue-induced reinstatement of cocaine-seeking behavior in

Plcb1+/- mice

Running title: *Plcb1* **is involved in cocaine-seeking behavior.**

Judit Cabana-Domínguez*[1,2,3,4], Elena Martín-García*[5], Ana Gallego-Roman[5], Rafael Maldonado[#5,6],

Noèlia Fernàndez-Castillo[#1,2,3,4] and Bru Cormand[#1,2,3,4]

[1] Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de

Barcelona, Catalonia, Spain

[2] Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), Spain

[3] Institut de Biomedicina de la Universitat de Barcelona (IBUB), Catalonia, Spain

[4] Institut de Recerca Sant Joan de Déu (IR-SJD), Esplugues de Llobregat, Barcelona, Catalonia Spain

[5] Laboratory of Neuropharmacology-Neurophar, Department of Experimental and Health Sciences,

Universitat Pompeu Fabra (UPF), Barcelona, Spain

[6] Hospital del Mar Medical Research Institute (IMIM), Barcelona, Spain

*These authors contributed equally.

#These authors equally supervised this work.

Address corresponding to Bru Cormand, Ph.D., and Noèlia Fernàndez-Castillo, Ph.D., Departament

de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de Barcelona, Avinguda

Diagonal 643, edifici Prevosti, 3ª planta, 08028 Barcelona, Catalonia, Spain; E-mail:

bcormand@ub.edu, noefernandez@ub.edu

WORD COUNT: 3425 Introduction, Results and Discussion + 2409 Materials and methods.

ABSTRACT

Background and Purpose: Cocaine addiction causes serious health problems and no effective

treatment is available yet. We previously identified a genetic risk variant for cocaine addiction in

the *PLCB1* gene and found this gene upregulated in postmortem brains of cocaine abusers and in

human dopaminergic neuron-like cells after an acute cocaine exposure. Here, we functionally

tested the contribution of *PLCB1* gene to cocaine addictive properties in mice.

Experimental approach: We used heterozygous *Plcb1* knockout mice (*Plcb1+/-*) and characterized

their behavioral phenotype. Subsequently, mice were trained for operant conditioning and self-

administered cocaine for 10 days. *Plcb1+/-* mice were assessed for cocaine motivation, followed

by 26 days of extinction and finally evaluated for cue-induced reinstatement of cocaine seeking.

Gene expression alterations after reinstatement were assessed in medial prefrontal cortex (mPFC)

and hippocampus (HPC) by RNAseq.

Key Results: *Plcb1+/-* mice showed normal behavior, although they had increased anxiety and

impaired short-term memory. Importantly, after cocaine self-administration and extinction, we

found a reduction in the cue-induced reinstatement of cocaine-seeking behavior in *Plcb1+/-* mice.

After reinstatement, we identified transcriptomic alterations in the medial prefrontal cortex of

Plcb1+/- mice, mostly related to pathways relevant to addiction like the dopaminergic synapse and

long-term potentiation.

Conclusions and Implications: To conclude, we found that heterozygous deletion of the *Plcb1*

gene decreases cue-induced reinstatement of cocaine seeking, pointing at PLCB1 as a possible

therapeutic target for preventing relapse and treating cocaine addiction.

KEYWORDS: addiction, relapse, cocaine self-administration, RNAsequencing, transcriptomic, resilience, medial prefrontal cortex (mPFC)

INTRODUCTION

Cocaine is the most used psychostimulant illicit drug worldwide (UNODC, 2019), causing severe health problems that include the development of cocaine addiction in around 15-16% of cocaine users (O'Brien and Anthony, 2005). Cocaine addiction is a complex psychiatric disorder that results from the interaction of genetic, epigenetic, and environmental risk factors (Pierce et al., 2018). The heritability of cocaine addiction is one of the highest among psychiatric disorders, estimated around 65% for women (Kendler et al., 2000) and 79% for men (Kendler and Prescott, 1998). However, the genetic factors and mechanisms that underlie the transition from drug use to addiction and its establishment remain unknown.

In a previous study, we identified a single nucleotide polymorphism in the *PLCB1* gene associated with drug dependence (rs1047383), and especially with a subgroup of cocaine-addicted patients, that was replicated in an independent clinical sample (Cabana-Domínguez et al., 2017). Also, genetic variants in this gene were found nominally associated with an illegal substance and cocaine addiction in two GWAS performed in European-American samples (Drgon et al., 2012). On the other hand, we found that cocaine increased the expression of *PLCB1* both in human dopaminergic neuron-like cells (differentiated SH-SY5Y cells) after acute cocaine exposure and in postmortem samples of the nucleus accumbens (NAc) of cocaine abusers (Cabana-Domínguez et al., 2017). Interestingly, this gene was also found over-expressed in the same brain region in mice

after cocaine administration for 7 days and also during withdrawal (Eipper-Mains et al., 2013). All

this evidence suggest that *PLCB1* may play a role in cocaine addiction.

The *PLCB1* gene encodes phospholipase C beta 1, and it is highly expressed in the brain, mainly in

the frontal cortex, basal ganglia (caudate, putamen, and NAc) and hippocampus. These brain

regions are crucial for drug reward and the formation of drug-context associations, both

contributing to the development and maintenance of addiction (Bisagno et al., 2016; Castilla-

Ortega et al., 2016, 2017; Kutlu and Gould, 2016; Pitts et al., 2016; Cooper et al., 2017). Several

neurotransmitters activate this protein, including dopamine through DRD1 and DRD2 (Lee et al.,

2004; Rashid et al., 2007), serotonin by 5-HT2A and 2C receptors (Hagberg et al., 1998; Chang et

al., 2000) and glutamate by mGluR1 (Conn and Pin, 1997; Hannan et al., 2001). Thus, PLCB1 might

be a point of convergence of neurotransmitter systems that play an essential role in the

development of addiction (recently reviewed (Howell and Negus, 2014; Moretti et al., 2020)). The

activation of PLCB1 produces the cleavage of phosphatidylinositol 4,5-bisphosphate (PIP2) into the

second messengers diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3), responsible for

intracellular signal transduction. Alterations in PLCB1-mediated signaling in the brain have been

associated with other neuropsychiatric disorders such as epilepsy, schizophrenia, and bipolar

disorder (Yang et al., 2016).

Here we studied the contribution of the *PLCB1* gene to cocaine addiction and dissected its

participation in the different aspects of the addictive process. We used heterozygous knockout

(KO) mice (*Plcb1+/-*), as the homozygous KO (*Plcb1-/-*) showed seizure attacks and low viability

after birth (Kim et al., 1997). In addition, the use of a constitutive heterozygous KO mouse model

allowed us to assess *Plcb1* haploinsufficiency during neurodevelopment in mice similarly to

humans, where inherited genetic risk or protective variants can modulate the susceptibility to

addiction (Cabana-Domínguez et al., 2017). First, we performed a general phenotypic

characterization using a battery of behavioral tests, including memory, anxiety, locomotor activity,

coordination, food and water intake, and sucrose preference. Then, we evaluated cocaine operant

self-administration, extinction, and cue-induced reinstatement. Finally, we studied transcriptional

alterations to further understand the molecular mechanisms involved.

RESULTS

In the present work, we aimed to assess the contribution of the *PLCB1* gene in the different stages

of cocaine addiction using heterozygous knockout mice (*Plcb1+/-*). To do so, we performed a

general phenotypic characterization and, then we evaluated cocaine operant self-administration,

extinction, and cue-induced reinstatement in those animals. Also, we studied transcriptional

changes in mPFC and HPC to understand the mechanisms involved.

Behavioral tests for phenotype characterization

We first evaluated the effects of the heterozygous deletion of *Plcb1* on general behavioral

responses, including locomotor, cognitive and emotional responses, food, and water intake.

Plcb1+/- mice showed a lower discrimination index in the novel object recognition compared to

WT (*t*-test=3.21, $P<0.01$, Figure 1B), suggesting an impairment in short-term memory. This

difference was not influenced by exploration time, as this variable was equal between genotypes

(Figure 1C). Furthermore, no differences in locomotor activity were reported between genotypes,

discarding an involvement of the *Plcb1* heterozygous deletion in locomotion (Figure 1D). Mutant

mice showed increased exploratory activity with a higher number of rearings than the WT mice (*U*-

Mann Whitney=67.5, $P<0.01$, Figure 1E). An anxiogenic profile was revealed in the elevated plus

maze in mutants, as shown by the reduced time spent in the open arms (t-test=3.02, $P<0.001$,

Figure 1F) and the percentage of time (t-test=3.16, $P<0.001$, Figure 1G). Finally, the rota rod test

revealed that the heterozygous deletion of *Plcb1* did not affect motor coordination (Figure 1H-I).

Several consummatory and locomotor parameters were long-term monitored in the PheComp

boxes (Supplementary Figure 1A-H, n=6 per genotype). No differences between genotypes were

revealed during the whole experimental period in body weight, food, and water intake, levels of

sucrose preference, stereotyped movements, horizontal and vertical locomotor activity, indicating

no altered behavior in the *Plcb1+/-* mice.

Operant conditioning maintained by cocaine

Then, we investigated the effects of the heterozygous deletion of *Plcb1* in cocaine behavioral

responses related to its addictive properties. For this purpose, *Plcb1+/-* mice and their WT

littermates were trained for cocaine operant self-administration (0.5mg/kg/infusion) during FR1

and FR3, progressive ratio, extinction and cue-induced reinstatement (Figure 2A). Control mice

trained with saline were included for both genotypes. Results showed that the percentage of mice

reaching the criteria of operant conditioning was 100% for both genotypes trained with cocaine

and 33% for mice trained with saline: [*Chi-square* test=9.50; $P<0.01$, *Plcb1+/-* cocaine vs. *Plcb1+/-*

saline] and [*Chi-square* test=14.00; $P<0.001$, WT cocaine vs. WT saline], as expected.

The primary reinforcing effects and the motivation for cocaine were similar in both genotypes

(Figure 2B-C). During FR1, both genotypes similarly increased the number of cocaine infusions

across sessions, whereas mice trained with saline remained steady (repeated measures ANOVA,

interaction between genotype x drug x sessions, $P<0.05$, Figure 2B and Supplementary Table S1).

Thus, the evolution of operant responding was different in mice trained with cocaine and saline,

with an increased responding over nearly all sessions in cocaine-trained mice . This enhancement

in cocaine responding was more pronounced in mutants than WT mice. Indeed, WT mice showed a

decrease in cocaine intake in the second session. The maximum number of infusions was reached

on session 5 in both genotypes and was slightly superior in *Plcb1+/-* mutants (36.35±1.47) than in

WT (29.61±2.23) but post hoc analyses demonstrated that this difference in cocaine intake was

not significant (Figure 2B). Similar findings were observed for the total number of nose-pokes in

which operant responding increased in both genotypes with cocaine but remained stable with

saline (Supplementary Figure S2A). When the effort to obtain one dose of cocaine increased to

FR3, the number of infusions was stable across sessions in WT and *Plcb1+/-* mice. Similarly,

operant responding was higher for all groups than in FR1 with stable higher levels of responding

(Supplementary Figure S2A) and a higher number of infusions (repeated measures ANOVA, the

main effect of the drug, *P*<0.001, Figure 2B) in mice trained with cocaine than with saline,

independently of the genotype.

Motivation for cocaine was evaluated in a progressive ratio schedule, and no significant

differences were obtained between genotypes (Figure 2C). The levels of extinction of the operant

behavior were similar between genotypes and decreased progressively across sessions (repeated

measures ANOVA, interaction genotype x sessions, *P*<0.001, Figure 2D). The percentage of mice

reaching cocaine-seeking extinction criteria was similar in *Plcb1+/-* (73%) and WT (78%) mice.

Importantly, *Plcb1+/-* mice showed significantly reduced cue-induced reinstatement of cocaine-

seeking compared to WT mice (*U* Mann-Whitney, *P*<0.05, Figure 2E), with 27.59% less active nose-

pokes compared to WT mice trained with cocaine. Furthermore, mice trained with saline from

both genotypes exhibited 55.84% reduction of active nose-pokes than WT mice trained with

cocaine and 39.20% less than mutants trained with cocaine. No significant differences were

obtained between genotypes in inactive nose-pokes during operant conditioning maintained by

cocaine nor during extinction (Supplementary Figure S2B-C). Both genotypes trained with cocaine

acquired the reinstatement criterion (double nose pokes in the active hole than the number of

nose pokes during the 3 consecutive days when the mice acquired the extinction criteria) showing

their capability to maintain this conditioning learning task. These data showed that *Plcb1+/-*

resulted in a phenotype of resistance to cue-induced reinstatement with reduced cocaine-seeking

(Figure 2E).

Brain transcriptomic analysis after the reinstatement of cocaine-seeking behavior

To further understand the role of PLCB1 in the molecular mechanisms involved in this cocaine

relapse-related phenotype, we analyzed the transcriptomic profiles of mPFC and HPC immediately

after the reinstatement of cocaine-seeking behavior in WT and *Plcb1+/-* mice trained with cocaine.

We identified 2,115 protein-coding genes differentially expressed (DEGs) in mPFC (1,231

downregulated and 943 upregulated) and only 12 in HPC, when comparing *Plcb1+/-* and WT

animals (Supplementary Tables S2-3). In accordance, PCA analysis and heatmap plots revealed that

individuals with the same genotype plotted together only in mPFC, but not in HPC (Supplementary

Figure S3). This suggested a more prominent role of mPFC, so further studies were carried out only

with DEGs of this brain area. Analysis of functional group over-representation identified several

processes previously related to cocaine addiction, including dopaminergic synapse, learning, long-

term potentiation (LTP), neurotransmitter secretion, and axon guidance, as well as other relevant

signaling pathways such as MAPK, mTOR, and neurotrophin (Figure 3A-B and Supplementary

Tables S3-5). We focused on the dopaminergic synapse pathway (enriched in the mPFCs DEGs;

P_{Adj}<0.05), in which phospholipase c (such as Plcb1) is directly participating in signal transmission

(Figure 4). Interestingly, many genes coding for proteins in this pathway are differentially

expressed in *Plcb1+/-* mice after cue-induced reinstatement (Figure 4 and Supplementary Table

S6). Furthermore, we found that several TFBS were over-represented in the DEGs of mPFC,

including YY1, MYOD, NRF1, ERR1, FREAC2, NFY and E4F1 (complete list of TFBS in Supplementary

Table S7). Then, we filtered the DEGs on mPFC based on fold-change (FC>|1.2|) and obtained a list

of 238 genes that we used for gene network construction. This analysis showed a highly scored

network (score=62, Figure 3C) that includes 31 DEGs involved in "cellular development, cellular

growth and proliferation, nervous system development and function".

DISCUSSION

Here we investigated, for the first time, the contribution of the *PLCB1* gene to cocaine addictive

properties using *Plcb1+/-* mice. We found that the heterozygous *Plcb1* genotype resulted in a

phenotype of resistance to cue-induced reinstatement of cocaine-seeking behavior. Furthermore,

we found relevant transcriptomic differences in *Plcb1+/-* mice compared to WT after cue-induced

reinstatement of cocaine seeking in mPFC, a brain area essential in relapse (Jasinska et al., 2015;

Moorman et al., 2015). This study supports a role for *PLCB1* in cocaine addiction, confirming

previous findings in humans (Cabana-Domínguez et al., 2017), and suggests it may be relevant in

relapse to cocaine addiction.

The phenotype of *Plcb1+/-* mice was characterized at different levels to evaluate consummatory

and general behavior. In general, mutant mice presented normal levels of body weight, food and

water intake, locomotor activity, and motor coordination, demonstrating that this animal model is

valid to study the effects of the heterozygous deletion of the *Plcb1* gene in other behavioral

responses. *Plcb1+/-* showed increased anxiety in the elevated plus maze, reduced short-term

memory in the novel object recognition paradigm, without any sign of depressive-like behavior in

the anhedonia sucrose preference test. Thus, the single targeting of the *Plcb1* gene has an impact

on selective emotional and cognitive responses.

The short-term memory impairment observed in the mutants did not affect the acquisition of

operant associative conditioning nor the instrumental learning that drives the goal-directed action,

as shown by similar levels of operant cocaine self-administration and extinction learning than WT.

In agreement, both genotypes trained with cocaine accomplish cue-induced reinstatement

criterion with high enhancement of responding during this test compared to extinction. Thus,

different brain circuits are involved in each kind of learning, with perirhinal-hippocampal

structures (Winters et al., 2004) participating in short-term memory in the novel object

recognition paradigm and mPFC-dorsal striatum pathway in cue-associated seeking. Also

differences in each paradigm such as the acute retrieval or repetitive exposure involved

respectively in each task may have a crucial influence in cognition. Hence, cocaine seeking can be

multidimensional, involving different types of associative learning that together lead to an

extensive repertoire of conditioned and instrumental responding.

The anxiogenic profile of *Plcb1+/-* is in accordance with the results previously reported in the

Plcb4-/- mice, which were associated with alterations in the cholinergic activity of the medial

septum (Shin et al., 2009). However, the selective knock-down of *Plcb1* in the mPFC did not

replicate this phenotype in a previous study (Kim et al., 2015), suggesting that other areas may be

involved. The anxiogenic profile of *Plcb1+/-* mice had no effect on cocaine self-administration since

the acquisition and extinction of this operant behavior was not modified in the mutants. Besides,

mutants showed protection against cue-induced reinstatement of cocaine seeking, instead of the

expected cocaine seeking promoted by an anxiogenic phenotype (Mantsch et al., 2016).

Therefore, the association of an anxiogenic profile with a phenotype resilient to cocaine seeking in

the *Plcb1+/-* mice suggests modifications in specific brain areas involved in cocaine relapse, such

as the mPFC. Recently, a phenotype resilient to develop food addiction has been associated with

increased strength of pyramidal glutamatergic synaptic transmission in the mPFC related to

decreased compulsivity in the face of negative consequences (Domingo-Rodriguez et al., 2020) in

mice with an anxiogenic profile (Lutz et al., 2015). Concerning cue-induced cocaine seeking, the

mPFC has a crucial role and the network of glutamatergic projections from the prelimbic mPFC to

the dorsal striatum participates in this phenotype (Lüscher et al., 2020). In our model, *Plcb1*

haploinsufficiency is linked to reduced cue-induced cocaine seeking possibly associated with

modifications in this top-down corticolimbic brain network. Furthermore, these glutamatergic

projections from mPFC to dorsal striatum receive mesocortical dopaminergic inputs from the VTA

that could be crucially involved in the protective effects of *Plcb1,* since this gene plays an essential

role in the dopaminergic signal transmission in the mPFC and many of the genes encoding for

proteins in this pathway are differentially expressed in *Plcb1+/-* mice after cue-induced

reinstatement (Figure 4).

Transcriptomic analyses performed after the cue-induced reinstatement of cocaine-seeking

behavior revealed some of the molecular underpinnings that underlie the cocaine relapse-related

phenotype observed in the mutant mice. Interestingly, differences in gene expression between

Plcb1+/- and WT were predominantly found in the mPFC, whereas almost no differences were

observed in the HPC. This evidence highlights mPFC as a key region to explain these differences in

cocaine-seeking reinstatement observed in *Plcb1+/-* mice. Consistently, in DEGs in mPFC, we found

enrichment on pathways that are essential for the development of addiction, including the

dopaminergic neurotransmission (Figures 3-4) (Wise and Robble, 2020). *PLCB1* is part of the

dopamine-DARPP-32 signaling pathway (Figure 4), which plays a key role in cocaine reward

(Borgkvist and Fisone, 2007; Nishi and Shuto, 2017). In a previous study, we also found this

pathway enriched in DEGs in the frontal cortex and ventral striatum of mice that showed

frustrated expected reward, produced by the cue in the absence of expected reward after a high

level of effort in a progressive ratio schedule of reinforcement with palatable food (Martín-García

et al., 2015b). Notably, *Plcb1* was upregulated in the frontal cortex of those frustrated mice with

increased responses to obtain the reward (palatable food). These data are in line with our findings

in which decreased expression of *Plcb1* (*Plcb1*+/- mice) results in decreased cue-induced responses

to obtain cocaine after extinction. All these results support the participation of *Plcb1* in cue-

induced cocaine seeking revealed in the present study and the involvement of mPFC .

Our transcriptomic analyses also found enrichment in genes related to learning and memory

processes, such as LTP in the mPFC. The transition from drug use to drug addiction is a

maladaptive process that directly affects learning and memory (Torregrossa et al., 2011; Goodman

and Packard, 2016). LTP produces long-lasting activity-dependent synaptic modifications that

underlie memory, and drugs of abuse alter this mechanism in several brain regions such as mPFC

(Huang et al., 2007; Lu et al., 2010), mesocorticolimbic system (Thomas et al., 2008), VTA (Placzek

et al., 2016) and hippocampus (Keralapurath et al., 2017; Preston et al., 2019), among others.

Importantly, genes related to the glutamatergic system such as *Gria2-4* (glutamate ionotropic

receptor AMPA2-4 (alpha 2-4)) and *Grik2* (ionotropic glutamate receptor kainate 2 (beta 2)) were

found to be upregulated in *Plcb1*+/- mice. Meanwhile, *Grin1, 2c, and 2d* (glutamate ionotropic

receptor NMDA1 (zeta 1, epsilon3 and epsilon4)) were downregulated, suggesting an increase in

AMPAR/NMDAR ratio in cocaine-experienced and abstinent *Plcb1 +/-* mice as compared to WT,

one of the key indicators of LTP induction and increased synaptic strength (Lüscher and Malenka,

2011). Furthermore, we found enrichment in other neuroplasticity (synapse organization, neuron

projection development, and axon guidance) (Bahi and Dreyer, 2005; O'Brien, 2009; Cooper et al.,

2017; Dong et al., 2017) and signaling pathways (MAPK, mTOR, and neurotrophin) (Li and Wolf,

2015; García-Pardo et al., 2016) related to addiction. The assessment of gene expression in the

mPFC of the *Plcb1+/-* mice highlighted alterations in relevant pathways for the addictive process,

which could contribute to the results observed in the cue-induced reinstatement.

Nowadays, there are few effective pharmacological treatments available for cocaine use disorders,

and frequently, psychosocial interventions in combination with pharmacotherapy are needed

(Kampman, 2019). Studies performed in mice have pinpointed multiple potential therapeutic

approaches for the reinstatement of cocaine-seeking behavior, targeting the reward circuit

(Buchta et al., 2020). In humans, treatment with bupropion, topiramate, or disulfiram has been

widely used. Bupropion, a non-tricyclic antidepressant that inhibits dopamine and norepinephrine

reuptake, is effective in reducing craving (Frishman, 2007). Topiramate, a GABA/glutamatergic

medication, has also been used to treat cocaine use disorder as it reduces the activity of the

mesocorticolimbic dopaminergic system (Kampman, 2019). A completely different strategy is the

use of disulfiram which potentially inhibits the oxidoreductase dopamine β-hydroxylase (DβH,

encoded by the *DBH* gene), which converts dopamine to norepinephrine (Kampangkaew et al.,

2019). Significantly, a genetic variation in the *SLC6A3* gene (encoding DAT) has been associated

with disulfiram treatment for cocaine addiction, with patients with higher DAT levels having better

treatment outcomes than those with lower DAT levels (Kampangkaew et al., 2019). Thus, further

studies and new therapeutic targets are needed to obtain effective treatment for cocaine

addiction. The results obtained in the present study underscore the relevance of the *Plcb1* gene in

the cue-induced reinstatement of cocaine seeking after extinction. Together with previous findings

in humans (Drgon et al., 2012; Cabana-Domínguez et al., 2017) and mice (Eipper-Mains et al.,

2013), *PLCB1* merits to be further evaluated as a promising novel therapeutic target for preventing

relapse and treating cocaine addiction .

The experimental approach used in our study, the *Plcb1+/-* mouse model, allowed us to reproduce

better the molecular context observed in humans in comparison to the use of a complete KO

mouse (Hall et al., 2013), as these animals preserved, at least, half of the expression of *Plcb1*.

However, the haploinsufficiency of *Plcb1* during neurodevelopment in these animals could

produce alterations that may contribute to the phenotype observed in the present study.

Nevertheless, this may also be similar in humans with genetic risk variants that decrease *PLCB1*

expression. Therefore, this approach is appropriate to study a specific genetic alteration that

confers susceptibility to drug addiction and to delineate the precise contribution of *PLCB1*.

To sum up, we studied, for the first time, the contribution of the *PLCB1* gene to cocaine addictive

properties using *Plcb1+/-* mice. Previous studies have revealed an up-regulation of *Plcb1* in brain

areas related to the reward circuit after cocaine exposure in animals (Eipper-Mains et al., 2013)

and in human cocaine abusers (Cabana-Domínguez et al., 2017). These changes, together with our

results, suggest that cocaine increases the expression of *Plcb1,* and this mechanism plays an

essential role in cocaine addiction, as revealed now by the resistance to cue-induced

reinstatement of cocaine-seeking behavior exhibited by *Plcb1+/-* mutant mice. These results

highlight the importance of the *Plcb1* gene in the development of cocaine addiction and relapse

and pinpoint PLCB1 as a promising therapeutic target for cocaine addiction and perhaps other

types of addiction.

METHODS

Animals

Male mice, 8 weeks old, were housed individually at temperature- and humidity-controlled

laboratory conditions (21±1°C, 55±10%) maintained with food and water ad libitum. Mice were

tested during the dark phase of a reverse light cycle (lights off at 8.00h and on at 20.00h). *Plcb1+/-*

mice in a C57BL/6J background and their wild-type (WT) littermates were used (Kim et al., 1997).

All experimental protocols were performed in accordance with the guidelines of the European

Communities Council Directive 2010/63/EU and approved by the local ethics committee (Comitè

Ètic d'Experimentació Animal-Parc de Recerca Biomèdica de Barcelona, CEEA-PRBB, agreement

N°9213). In agreement, maximal efforts were made to reduce the suffering and the number of

mice used.

Genotyping of transgenic mice

Plcb1+/- mice in a C57BL/6J background were kindly provided by H.-S. Shin (Kim et al., 1997). Very

briefly, the knockout animals were generated by the replacement of exons that encode aminoacid

residues 50–82 by a Neomycin cassette that was used both for gene disruption and for positive

selection.

To genotype the mice, genomic DNA from ear punches or tail tips was extracted as previously

described (Gómez-Grau et al., 2017). Genotyping was carried out by PCR on extracted genomic

DNA using three specific PCR primers: a sense primer, 5⬚-GTTAAGTCCTCAGGCAAACACC-3', and

two antisense primers, 5⬚-ACCTTGGGAGCTTTGGCGTG3' and 5⬚-CTGACTAGGGGAGGAGTAGAAG-

3', that allowed us to amplify a 180bp *Plcb1+/+* band and a 290bp *Plcb1-/-* band (Filis et al., 2009).

Fragments were separated by 2.5% agarose gel electrophoresis. Known WT and *Plcb1+/-* controls

were always used to validate genotyping results. The genotype of all animals used was confirmed

after the cue-induced reinstatement.

Behavioral tests for phenotype characterization

A group of 25 WT and 12 *Plcb1+/-* mice underwent different behavioral tests for phenotype

characterization for 10 days, as described in Figure 1:

Short-term memory. To measure short-term memory, the novel object recognition (NOR) test was

used. This test was performed in a V-maze apparatus (40 cm per side, Panlab, Spain) as previously

described by us and others (Bura et al., 2010; Planaguma et al., 2015). On day 1 mice were

habituated for 9 minutes in the V-maze. On day 2, mice were put back into the V-maze for 9

minutes; two identical objects were presented, and the time the mice spent exploring each object

was recorded. After a retention phase of 3 hours, the mice were placed for another 9 minutes into

the V-maze; in each paradigm, one of the familiar objects was replaced with a novel object and the

total time spent exploring each object (novel and familiar) was registered. Object exploration was

defined as the orientation of the nose to the object at a distance of less than 2 cm. A

discrimination index was calculated as the difference in the time spent exploring the novel and the

time spent exploring the familiar object divided by the total time exploring both objects. A higher

discrimination index is considered to reflect more excellent memory retention for the familiar

object.

Locomotor activity. Locomotor activity was assessed in locomotor activity boxes (9×20×11 cm;

Imetronic, Passac, France), equipped with 2 rows of photocell detectors, and placed in a low-

luminosity environment (20–25 lux), as previously described (Martin-Garcia et al., 2013). The

locomotor activity was recorded for 2 hours, and 2 variables were measured: horizontal activity as

horizontal displacement and vertical activity as vertical exploration/elevation.

Elevated plus maze test. This test was performed as previously reported (Planaguma et al., 2015).

The test measures the conflict between the natural tendency of mice to avoid an illuminated and

elevated surface, and the natural tendency to explore new environments. It consisted of a black

plastic cross with arms 40 cm long and 6 cm wide placed 50 cm above the floor. Two opposite

arms were surrounded by walls (15 cm high, closed arms, 10 lux), while the two other arms were

devoid of such walls (open arms, 200 lux). A central platform connected the four arms. At the start

of the session, the mouse was placed at the end of a closed arm facing the wall. During the 5-

minutes trial, the number of entries and the time spent in each arm were recorded. Anxiety was

assessed as both the time spent avoiding the open arms and the number of entries into them.

Motor coordination. Rota rod apparatus was used to assess motor coordination. Precisely, this test

measures the ability of the mouse to remain on a rotating rod in which the speed of rotation is

gradually increased and evaluate general motor coordination. The device is a round drum (5 cm

diameter) suspended between 2 plexiglass walls. The drum is suspended at 24 cm from a soft mat

covered tabletop. The speed of rotation is controlled by an electric engine with a digital revolution

per min (rpm) display. The test consisted in five trial sessions. Each trial started with the mouse

being placed on the rotating rod at 4.0 rpm, and then every 3 s the speed increased by 1.0 rpm

until 20.0 rpm. The trial terminated when the mouse fell from the rod or after 90 s, whichever

occurred first. There was a 10 s interval between trials. The average maximum rota rod speed and

time to fall was calculated for each mouse (Martin-Garcia et al., 2013).

Experimental procedure for food and drink monitoring.

Habituation. Six animals per genotype were taken from their home cages and placed in

experimental chambers named PHECOMP boxes (Panlab and Harvard Apparatus) equipped with a

food, drink and motor activity monitoring system

(https://www.panlab.com/en/products/phecomp-system-panlab) for two weeks of habituation.

Animals were monitored in these boxes 24 h a day during the whole experimental period.

Sucrose preference test. After a period of habituation, a sucrose preference test was performed on

the third week of PheComp cage housing during one week, as previously described (Bura et al.,

2010; Planaguma et al., 2015). Two bottles of water, one with 2% sucrose and the other without,

were placed in the cage. The position of bottles were exchanged everyday, and the consumption

was measured after a 24h interval. The preference for sucrose was calculated as the relative

amount of water with sucrose versus total liquid (water with and without sucrose) consumed by

the mice (Bura et al., 2010).

Operant conditioning maintained by cocaine

Operant self-administration apparatus. The self-administration experiments were conducted in

mouse operant chambers (Model ENV-307A-CT; Med Associates Inc., Georgia, VT, USA) equipped

with two holes, one randomly selected as the active hole and the other as the inactive hole as

previously described (Gutierrez-Cuesta et al., 2014).

Operant conditioning maintained by cocaine. Cocaine self-administration experiments were

performed as previously described (Gutierrez-Cuesta et al., 2014; Martín-García et al., 2015a).

Mice were deeply anesthetized by intraperitoneal injection of a mixture of ketamine (75mg/kg;

Imalgène; Merial Laboratorios S.A., Barcelona, Spain) and medetomidine (1mg/kg; Domtor;

Esteve, Barcelona, Spain). After surgery, anesthesia was reversed by subcutaneous injection of the

synthetic α2 adrenergic receptor antagonist, atipamezole (2.5mg/kg; Revertor; Virbac, Barcelona,

Spain) indicated for the reversal of the sedative and analgesic effects of medetomidine (α2

adrenergic receptor agonist). In addition, mice received an intraperitoneal injection of gentamicin

(1 mg/kg; Genta-Gobens; Laboratorios Normon, S.A., Madrid, Spain) along with subcutaneous

administration of the analgesic meloxicam (2 mg/kg; Metacam; Boehringer Ingelheim, Rhein,

Germany). Each 2h daily self-administration session started with a priming injection of the drug.

Cocaine (obtained from The Spanish Agency of Drugs, Madrid, Spain) was intravenously infused in

23.5µl over 2s (0.5 mg/kg per injection). Cue light, located above the active hole, was paired with

the delivery of the reinforcer. Mice (WT n=36; *Plcb1+/-* n=26) were trained under a fixed ratio 1

schedule of reinforcement (FR1; one nose-poke lead to the delivery of one dose of cocaine) over 5

consecutive daily sessions and under a fixed ratio 3 (FR3) over 5 consecutive daily sessions. Control

mice trained with saline were included for both genotypes (WT n=6; *Plcb1+/-* n=6). The timeout

period after infusion delivery was 10 s. Responses on the inactive hole and all responses elicited

during the 10-s timeout period were also recorded. Responses during the 10-s timeout period

were considered a measure of impulsivity reflecting the inability to stop motor behavior once it is

initiated. The criteria for self-administration behavior were achieved when all of the following

conditions were met: (1) mice maintained stable, responding with 20% deviation from the mean of

the total number of reinforcers earned in three consecutive sessions (80% of stability); (2) at least

75% of mice responding on the active hole; and (3) a minimum of 10 reinforcers per session.

After the 10 FR sessions, animals were tested in a progressive ratio schedule of 4h where the

response requirement to earn the cocaine escalated according to the following series: 1–2–3–5–

12–18–27–40–60–90–135–200–300–450–675–1000. Then, we proceed with the extinction phase,

2h daily sessions following the same experimental conditions than the cocaine self-administration

sessions except that cocaine were not available, and cue-light was not presented. Mice were given

2-h daily sessions until they achieved the extinction criterion with a maximum of 26 sessions. The

criterion for extinction was achieved when, during 3 consecutive sessions, mice completed a mean

number of nose pokes in the active hole consisting of 30% of the mean responses obtained during

the 3 days to achieve the acquisition criteria for cocaine self-administration training. On day 27,

only mice that accomplished extinction criterion were tested in the cue-induced reinstatement

during a 2h session, to evaluate the reinstatement of cocaine-seeking behavior. The test for cue-

induced reinstatement was conducted under the same conditions used in the training phase

except that cocaine was not available. The reinstatement criterion was achieved when nose pokes

in the active hole were double the number of nose pokes in the active hole during the 3

consecutive days when the mice acquired the extinction criteria.

The catheter was flushed daily with heparinized saline (30 USP units/ml). The patency of

intravenous catheters was evaluated after the last cocaine self-administration session and

whenever the behavior appeared to deviate dramatically from that observed previously by

infusion of thiopental through the catheter. If prominent signs of anesthesia were not apparent

within 3s of the infusion, the mouse was removed from the experiment. The success rate for

maintaining patency of the catheter (mean duration of 11 days) until the end of the cocaine self-

administration training was 90%.

After the cue-induced reinstatement, animals were euthanized by decapitation, brains were

quickly removed, and the medial prefrontal cortex (mPFC) and hippocampus (HPC) were dissected.

Brain tissues were then frozen by immersion in 2-methylbutane surrounded by dry ice and stored

at −80°C for later RNA isolation and transcriptomic analyses.

RNA extraction and RNA sequencing

Total RNA of 12 WT and 11 *Plcb1+/-* mice from mPFC and HPC were isolated using the RNeasy

Lipid Tissue Mini Kit (Qiagen Düsseldorf, Germany) according to the manufacturer's protocol. RNA

concentration was determined using the NanoDrop ND-1000 spectrophotometer (NanoDrop

Technologies, Wilmington, DE, USA), and integrity was evaluated using the Bioanalyzer2100

platform (Agilent Technologies, Santa Clara, CA, USA). RNA samples were grouped in 4 pools

consisting of 3 mice per pool for each experimental group, except for 1 group in the *Plcb1+/-* mice

were only 2 animals were pooled. The pools were organized to homogenize the average number

of nose pokes in the different pools. The pooled individuals were the same for both mPFC and

HPC.**RNA sequencing**

RNA sequencing (RNAseq) was performed by the Centre de Regulació Genòmica (CRG, Barcelona,

Spain). Libraries were prepared using the TruSeq Stranded mRNA Sample Prep Kit_v2 (Illumina,

San Diego, CA, USA) according to the manufacturer's protocol and sequenced 2x75 on Illumina's

HiSeq3000 system for both mPFC and HPC. The Bioinformatics service of CRG carried out the

analysis of RNAseq. Briefly, FastQC v0.11.5(Andrews S., 2010) was used to inspect the reads

quality and CutAdapt 1.7.1 (Martin, 2011) to clean the data of adapters and low-quality reads.

Then, reads were mapped to the *Mus musculus* genome of reference (GRCm38/mm10) with STAR

2.5.3a (Dobin et al., 2013), and the differential expression analysis was done by DESeq2 (Love et

al., 2014a) to compare WT and *Plcb1+/-*. Corrections for multiple testing were applied by adjusting

the p-values with a 5% False Discovery Rate.

RNAseq data of mPFC and HPC were explored on a principal component analysis (PCA) plot using

the "plotPCA" method from the DESeq2 package (Love et al., 2014b) and log2 gene expression

data. The PCAs were performed with the 500 genes showing the highest variance among the

samples to calculate the distance among them. The heatmaps were performed using the

"heatmap" function on R, and the hierarchical clustering considered the euclidean distance

between the samples considering all the genes or only those with corrected p-value < 1e-05.

Functional annotation of RNAseq results

We performed a functional group enrichment of differentially expressed genes (DEGs) in mPFC

using the DAVID Annotation Tool (http://david.abcc.ncifcrf.gov)(Jiao et al., 2012) considering GO

(Gene Ontology) biological processes and KEGG pathways (Kyoto Encyclopedia of Genes and

Genomes). Then, we searched for over-represented transcription factor-binding sites (TFBS) using

the information of MsigDB (https://www.gsea-msigdb.org/gsea/msigdb) integrated on

WebGestalt2019 (http://www.webgestalt.org/)(Liao et al., 2019), and the default parameters,

applying the weighted set cover method to reduce redundancy. In both analyses, the Benjamin–

Hochberg procedure was performed for multiple testing. Finally, we investigated the existence of

gene networks with Ingenuity Pathway Analysis 8.8 software (IPA,

http://www.ingenuity.com/products/ipa; Ingenuity Systems, Redwood City, CA, USA) (Krämer et

al., 2014) after selecting genes with fold-change >|1.2|.

Statistical analysis

Three-way ANOVA with repeated measures was used to test the evolution over sessions or days.

Sessions or days were used as within-subject factors and genotype (*Plcb1+/-* or WT) and drug

(cocaine or saline) were used as between-subjects factors. Post-hoc analyses (Newman-Keuls)

were performed when required. Comparisons between two groups were analyzed by Student *t*-

test or *U*-Mann-Whitney depending on the distribution defined by the Kolmogorov-Smirnov

normality test and the sample size. The chi-square analyses were performed to compare the

percentage of mice that acquired operant learning criteria in the different experimental groups.

Results are expressed as mean ± SEM or individual values with the median and the interquartile

range specified in the figure legend. Differences were considered significant at $P<0.05$. The sample

size was calculated based on power analysis. The statistical analyses were performed using the

Statistical Package for Social Science tool SPSS®25.0 (SPSS Inc, Chicago, USA).

See supplementary information for more details of genotyping of transgenic mice, drugs,

behavioral tests of phenotype characterization, operant conditioning maintained by cocaine, RNA

extraction and sequencing.

500   ACKNOWLEDGEMENTS AND FUNDING

We are grateful to H.-S. Shin for providing breeder mice to establish the colony. We thank M.

Linares, R. Martín, D. Real, F. Porrón for their technical support, and the Genomics Unit at the CRG

for assistance with the mRNAseq.

This work was supported by the Spanish 'Ministerio de Economía y Competitividad-MINECO'

(#SAF2017-84060-R-AEI/FEDER-UE), the Spanish 'Instituto de Salud Carlos III, RETICS-RTA'

(#RD12/0028/0023), the 'Generalitat de Catalunya, AGAUR' (#2017 SGR-669), 'ICREA-Acadèmia'

(#2015) and the Spanish 'Ministerio de Sanidad, Servicios Sociales e Igualdad, ´Plan Nacional Sobre

Drogas of the Spanish Ministry of Health' (#PNSD-2017I068) to RM, 'Fundació La Marató-TV3'

(#2016/20-30) and 'Plan Nacional Sobre Drogas of the Spanish Ministry of Health' (#PNSD-

2019I006) to E.M-G., Spanish 'Ministerio de Ciencia, Innovación y Universidades' (#RTI2018-

100968-B-100), Spanish 'Ministerio de Economía y Competitividad' (#SAF2015-68341-R), 'AGAUR-

Generalitat de Catalunya' (#2017-SGR-738) and Spanish 'Ministerio de Sanidad, Servicios Sociales e

Igualdad, ´Plan Nacional Sobre Drogas of the Spanish Ministry of Health' (#PNSD-2017I050) to BC.

The research leading to these results has also received funding from the European Union H2020

Program [H2020/2014-2020] under grant agreements n° 667302 (CoCA), 643051 (MiND) and

728018 (Eat2beNICE) to BC. JC-D was supported by the H2020 CoCA and Eat2beNICE projects and

NF-C by 'Centro de Investigación Biomédica en Red de Enfermedades Raras' (CIBERER).

J.C-D, B.C. and N.F.-C. obtained and maintained the mutant mice (*Plcb1+/-*); E.M.-G. and R.M.

conceived and designed the behavioral studies with input from J.C.-D, B.C. and N. F.-C; E.M.-G. and

J.C.-D. performed the behavioral phenotype characterization of mutant mice. E.M.-G. performed

the surgery for i.v. catheterization and the operant conditioning maintained by cocaine. A.G.-R.

collaborated in extinction and cue-induced reinstatement. E.M.-G. and J.C.-D. performed statistical

analyses and graphs with the supervision of R.M., B.C. and N.F.-C.; A.G.R. collaborated with the

extraction of the samples; J.C.-D. performed the RNA extractions, RNA sequencing, and the

bioinformatic analyses supervised by B.C. and N.F.-C.; E.M.-G. and J.C.-D. wrote the manuscript and

N.F.C., R.M. and B.C. provided a critical review of the manuscript with inputs from all the other

authors.

We have uploaded it to the preprint server bioRxiv (https://doi.org/10.1101/2020.06.18.158964).

DISCLOURES

The authors have no conflicts of interest.

REFERENCES

Andrews S. (2010). FastQC: A quality control tool for high throughput sequence data.

Bahi, A., and Dreyer, J.L. (2005). Cocaine-induced expression changes of axon guidance molecules
in the adult rat brain. Mol. Cell. Neurosci. *28*: 275–291.

Bisagno, V., González, B., and Urbano, F.J. (2016). Cognitive enhancers versus addictive
psychostimulants: The good and bad side of dopamine on prefrontal cortical circuits. Pharmacol.
Res. *109*: 108–118.

Borgkvist, A., and Fisone, G. (2007). Psychoactive drugs and regulation of the cAMP/PKA/DARPP-
32 cascade in striatal medium spiny neurons. Neurosci. Biobehav. Rev. *31*: 79–88.

Buchta, W.C., Moutal, A., Hines, B., Garcia-Keller, C., Smith, A.C.W., Kalivas, P., et al. (2020).
Dynamic CRMP2 Regulation of CaV2.2 in the Prefrontal Cortex Contributes to the Reinstatement
of Cocaine Seeking. Mol. Neurobiol. *57*: 346–357.

Bura, S.A., Burokas, A., Martin-Garcia, E., and Maldonado, R. (2010). Effects of chronic nicotine on
food intake and anxiety-like behaviour in CB(1) knockout mice. Eur Neuropsychopharmacol. *20*:
369–378.

Cabana-Domínguez, J., Roncero, C., Pineda-Cirera, L., Palma-Álvarez, R.F., Ros-Cucurull, E., Grau-
López, L., et al. (2017). Association of the PLCB1 gene with drug dependence. Sci. Rep. *7*: 1–8.

Castilla-Ortega, E., Ladrón de Guevara-Miranda, D., Serrano, A., Pavón, F.J., Suárez, J., Rodríguez
de Fonseca, F., et al. (2017). The impact of cocaine on adult hippocampal neurogenesis: Potential
neurobiological mechanisms and contributions to maladaptive cognition in cocaine addiction
disorder. Biochem. Pharmacol. *141*: 100–117.

Castilla-Ortega, E., Serrano, A., Blanco, E., Araos, P., Suárez, J., Pavón, F.J., et al. (2016). A place for
the hippocampus in the cocaine addiction circuit: Potential roles for adult hippocampal
neurogenesis. Neurosci. Biobehav. Rev. *66*: 15–32.

Chang, M., Zhang, L., Tam, J.P., and Sanders-Bush, E. (2000). Dissecting G protein-coupled receptor
signaling pathways with membrane- permeable blocking peptides. Endogenous 5-HT(2C) receptors
in choroid plexus epithelial cells. J. Biol. Chem. *275*: 7021–7029.

Conn, P.J., and Pin, J.P. (1997). Pharmacology and functions of metabotropic glutamate receptors.
Annu. Rev. Pharmacol. Toxicol. *37*: 205–237.

Cooper, S., Robison, A.J., and Mazei-Robison, M.S. (2017). Reward Circuitry in Addiction.
Neurotherapeutics *14*: 687–697.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., et al. (2013). STAR: Ultrafast
universal RNA-seq aligner. Bioinformatics *29*: 15–21.

Domingo-Rodriguez, L., Ruiz de Azua, I., Dominguez, E., Senabre, E., Serra, I., Kummer, S., et al.
(2020). A specific prelimbic-nucleus accumbens pathway controls resilience versus vulnerability to
food addiction. Nat. Commun. *11*:.

Dong, Y., Taylor, J.R., Wolf, M.E., and Shaham, Y. (2017). Circuit and synaptic plasticity mechanisms
of drug relapse. J. Neurosci. *37*: 10867–10876.

Drgon, T., Johnson, C., Nino, M., Drgonova, J., Walther, D., and Uhl, G.R. (2012). 'Replicated'
genome wide association for dependence on illegal substances: genomic regions identified by
overlapping clusters of nominally positive SNPs. *156(2)*: 125–138.

Eipper-Mains, J.E., Kiraly, D.D., Duff, M.O., Horowitz, M.J., Mcmanus, C.J., Eipper, B.A., et al.

(2013). Effects of cocaine and withdrawal on the mouse nucleus accumbens transcriptome. Genes,

Brain Behav. *12*: 21–33.

Filis, P., Lannagan, T., Thomson, A., Murray, A.A., Kind, P.C., and Spears, N. (2009). Phospholipase

C-β1 signaling affects reproductive behavior, ovulation, and implantation. Endocrinology *150*:

3259–3266.

Frishman, W.H. (2007). Smoking Cessation Pharmacotherapy — Nicotine and Non-Nicotine

Preparations. 10–22.

García-Pardo, M.P., Roger-Sanchez, C., Rodríguez-Arias, M., Miñarro, J., and Aguilar, M.A. (2016).

Pharmacological modulation of protein kinases as a new approach to treat addiction to cocaine

and opiates. Eur. J. Pharmacol. *781*: 10–24.

Gómez-Grau, M., Albaigès, J., Casas, J., Auladell, C., Dierssen, M., Vilageliu, L., et al. (2017). New

murine Niemann-Pick type C models bearing a pseudoexon-generating mutation recapitulate the

main neurobehavioural and molecular features of the disease. Sci. Rep. *7*:.

Goodman, J., and Packard, M.G. (2016). Memory systems and the addicted brain. Front. Psychiatry

7: 24.

Gutierrez-Cuesta, J., Burokas, A., Mancino, S., Kummer, S., Martin-Garcia, E., and Maldonado, R.

(2014). Effects of genetic deletion of endogenous opioid system components on the reinstatement

of cocaine-seeking behavior in mice. Neuropsychopharmacology. *39*: 2974–2988.

Hagberg, G.-B., Blomstrand, F., Nilsson, M., Tamir, H., and Hansson, E. (1998). Stimulation of 5-

HT2A receptors on astrocytes in primary culture opens voltage-independent Ca channels.

Neurochem. Int. *32*: 153–162.

Hall, F.S., Drgonova, J., Jain, S., and Uhl, G.R. (2013). Implications of genome wide association

studies for addiction: Are our a priori assumptions all wrong? Pharmacol. Ther. *140*: 267–279.

Hannan, A.J., Blakemore, C., Katsnelson, A., Vitalis, T., Huber, K.M., Bear, M., et al. (2001). PLC-β1,

activated via mGluRs, mediates activity-dependent differentiation in cerebral cortex. Nat.

Neurosci. *4*: 282–288.

Howell, L.L., and Negus, S.S. (2014). Monoamine transporter inhibitors and substrates as

treatments for stimulant abuse. In Advances in Pharmacology, (NIH Public Access), pp 129–176.

Huang, C.C., Lin, H.J., and Hsu, K. Sen (2007). Repeated cocaine administration promotes long-term

potentiation induction in rat medial prefrontal cortex. Cereb. Cortex *17*: 1877–1888.

Jasinska, A.J., Chen, B.T., Bonci, A., and Stein, E.A. (2015). Dorsal MPFC circuitry in rodent models

of cocaine use: Implications for drug-addiction therapies. Addict. Biol. *20*: 215.

Jiao, X., Sherman, B.T., Huang, D.W., Stephens, R., Baseler, M.W., Lane, H.C., et al. (2012). DAVID-

WS: A stateful web service to facilitate gene/protein list analysis. Bioinformatics *28*: 1805–1806.

Kampangkaew, J.P., Spellicy, C.J., Nielsen, E.M., Harding, M.J., Ye, A., Hamon, S.C., et al. (2019).

Pharmacogenetic role of dopamine transporter (SLC6A3) variation on response to disulfiram

treatment for cocaine addiction. Am. J. Addict. *28*: 311–317.

Kampman, K.M. (2019). The treatment of cocaine use disorder. 1–8.

Kendler, K.S., Karkowski, L.M., Neale, M.C., and Prescott, C.A. (2000). Illicit Psychoactive Substance

Use, Heavy Use, Abuse, and Dependence in a US Population-Based Sample of Male Twins. Arch.

Gen. Psychiatry *57*: 261–269.

Kendler, K.S., and Prescott, C.A. (1998). Cocaine use, abuse and dependence in a population-based

sample of female twins. Br. J. Psychiatry *173*: 345–350.

Keralapurath, M.M., Briggs, S.B., and Wagner, J.J. (2017). Cocaine self-administration induces

changes in synaptic transmission and plasticity in ventral hippocampus. Addict. Biol. *22*: 446–456.

Kim, D., Jun, K.S., Lee, S.B., Kang, N.G., Min, D.S., Kim, Y.H., et al. (1997). Phospholipase C isozymes

selectively couple to specific neurotransmitter receptors. Nature *389*: 290–293.

Kim, S.-W., Seo, M., Kim, D.-S., Kang, M., Kim, Y.-S., Koh, H.-Y., et al. (2015). Knockdown of

phospholipase C-β1 in the medial prefrontal cortex of male mice impairs working memory among

multiple schizophrenia endophenotypes. J. Psychiatry Neurosci. *40*: 78.

Krämer, A., Green, J., Pollard, J., and Tugendreich, S. (2014). Causal analysis approaches in

ingenuity pathway analysis. Bioinformatics *30*: 523–530.

Kutlu, M.G., and Gould, T.J. (2016). Effects of drugs of abuse on hippocampal plasticity and

hippocampus-dependent learning and memory: contributions to development and maintenance

of addiction. Learn. Mem. *23*: 515–533.

Lee, S.P., So, C.H., Rashid, A.J., Varghese, G., Cheng, R., Lança, A.J., et al. (2004). Dopamine D1 and

D2 receptor co-activation generates a novel phospholipase C-mediated calcium signal. J. Biol.

Chem. *279*: 35671–35678.

Li, X., and Wolf, M.E. (2015). Multiple faces of BDNF in cocaine addiction. Behav. Brain Res. *279*:

240–254.

Liao, Y., Wang, J., Jaehnig, E.J., Shi, Z., and Zhang, B. (2019). WebGestalt 2019: gene set analysis

toolkit with revamped UIs and APIs. Nucleic Acids Res. *47*: W199–W205.

Love, M.I., Huber, W., and Anders, S. (2014a). Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol. *15*: 550.

Love, M.I., Huber, W., and Anders, S. (2014b). Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol. *15*: 550.

Lu, H., Cheng, P., Lim, B.K., Khoshnevisrad, N., and Poo, M. (2010). Elevated BDNF after Cocaine

Withdrawal Facilitates LTP in Medial Prefrontal Cortex by Suppressing GABA Inhibition. Neuron *67*:

821–833.

Lüscher, C., and Malenka, R.C. (2011). Drug-Evoked Synaptic Plasticity in Addiction: From

Molecular Changes to Circuit Remodeling. Neuron *69*: 650–663.

Lüscher, C., Robbins, T.W., and Everitt, B.J. (2020). The transition to compulsion in addiction. Nat.

Rev. Neurosci. *21*: 247–263.

Lutz, B., Marsicano, G., Maldonado, R., and Hillard, C.J. (2015). The endocannabinoid system in

guarding against fear, anxiety and stress. Nat. Rev. Neurosci. *16*: 705–718.

Mantsch, J.R., Baker, D.A., Funk, D., Lê, A.D., and Shaham, Y. (2016). Stress-induced reinstatement

of drug seeking: 20 years of progress. Neuropsychopharmacology *41*: 335–356.

Martín-García, E., Bourgoin, L., Cathala, A., Kasanetz, F., Mondesir, M., Gutiérrez-Rodriguez, A., et

al. (2015a). Differential Control of Cocaine Self-Administration by GABAergic and Glutamatergic

CB1 Cannabinoid Receptors. Neuropsychopharmacology 1–14.

Martín-García, E., Fernández-Castillo, N., Burokas, A., Gutiérrez-Cuesta, J., Sánchez-Mora, C.,

Casas, M., et al. (2015b). Frustrated expected reward induces differential transcriptional changes

in the mouse brain. Addict. Biol. *20*: 22–37.

Martin-Garcia, E., Mannara, F., Gutierrez-Cuesta, J., Sabater, L., Dalmau, J., Maldonado, R., et al.

(2013). Intrathecal injection of P/Q type voltage-gated calcium channel antibodies from

paraneoplastic cerebellar degeneration cause ataxia in mice. J.Neuroimmunol. *261*: 53–59.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.Journal *17*: 10.

Moorman, D.E., James, M.H., McGlinchey, E.M., and Aston-Jones, G. (2015). Differential roles of

medial prefrontal subregions in the regulation of drug seeking. Brain Res. *1628*: 130–146.

Moretti, J., Poh, E.Z., and Rodger, J. (2020). rTMS-Induced Changes in Glutamatergic and

Dopaminergic Systems: Relevance to Cocaine and Methamphetamine Use Disorders. Front.

Neurosci. *14*: 137.

Nishi, A., and Shuto, T. (2017). Potential for targeting dopamine/DARPP-32 signaling in

neuropsychiatric and neurodegenerative disorders. Expert Opin. Ther. Targets *21*: 259–272.

O'Brien, C.P. (2009). Neuroplasticity in addictive disorders. Dialogues Clin. Neurosci. *11*: 350–353.

O'Brien, M.S., and Anthony, J.C. (2005). Risk of Becoming Cocaine Dependent: Epidemiological

Estimates for the United States, 2000–2001. Neuropsychopharmacology *30*: 1006–1018.

Pierce, R.C., Fant, B., Swinford-Jackson, S.E., Heller, E.A., Berrettini, W.H., and Wimmer, M.E.

(2018). Environmental, genetic and epigenetic contributions to cocaine addiction.

Neuropsychopharmacology *43*: 1471–1480.

Pitts, E.G., Taylor, J.R., and Gourley, S.L. (2016). Prefrontal cortical BDNF: A regulatory key in

cocaine- and food-reinforced behaviors. Neurobiol. Dis. *91*: 326–335.

Placzek, A.N., Prisco, G.V. Di, Khatiwada, S., Sgritta, M., Huang, W., Krnjević, K., et al. (2016). EIF2α-

mediated translational control regulates the persistence of cocaine-induced LTP in midbrain
dopamine neurons. Elife *5*:.

Planaguma, J., Leypoldt, F., Mannara, F., Gutierrez-Cuesta, J., Martin-Garcia, E., Aguilar, E., et al.
(2015). Human N-methyl D-aspartate receptor antibodies alter memory and behaviour in mice.
Brain. *138*: 94–109.

Preston, C.J., Brown, K.A., and Wagner, J.J. (2019). Cocaine conditioning induces persisting changes
in ventral hippocampus synaptic transmission, long-term potentiation, and radial arm maze
performance in the mouse. Neuropharmacology *150*: 27–37.

Rashid, A.J., So, C.H., Kong, M.M.C., Furtak, T., El-Ghundi, M., Cheng, R., et al. (2007). D1-D2
dopamine receptor heterooligomers with unique pharmacology are coupled to rapid activation of
Gq/11 in the striatum. Proc. Natl. Acad. Sci. U. S. A. *104*: 654–659.

Shin, J., Gireesh, G., Kim, S.W., Kim, D.S., Lee, S., Kim, Y.S., et al. (2009). Phospholipase C β4 in the
medial septum controls cholinergic theta oscillations and anxiety behaviors. J. Neurosci. *29*:
15375–15385.

Thomas, M.J., Kalivas, P.W., and Shaham, Y. (2008). Neuroplasticity in the mesolimbic dopamine
system and cocaine addiction. Br. J. Pharmacol. *154*: 327–342.

Torregrossa, M.M., Corlett, P.R., and Taylor, J.R. (2011). Aberrant learning and memory in
addiction. Neurobiol. Learn. Mem. *96*: 609–623.

UNODC (2019). World Drug Report 2019.

Winters, B.D., Forwood, S.E., Cowell, R.A., Saksida, L.M., and Bussey, T.J. (2004). Double
dissociation between the effects of peri-postrhinal cortex and hippocampal lesions on tests of
object recognition and spatial memory: Heterogeneity of function within the temporal lobe. J.
Neurosci. *24*: 5901–5908.

Wise, R.A., and Robble, M.A. (2020). Dopamine and addiction. Annu. Rev. Psychol. *71*: 79–106.

Yang, Y.R., Kang, D.S., Lee, C., Seok, H., Follo, M.Y., Cocco, L., et al. (2016). Primary phospholipase C
and brain disorders. Adv. Biol. Regul. *61*: 80–85.

A

Behavioral tests for phenotype characterization

Days 1 - 2 - 3 - 4 - 5 - 6 - 7 - 8 - 9 - 10

Short-term memory

Locomotor activity

Anxiety-like behavior

Motor coordination

V-maze

Actimetry box

Elevated plus maze

Rota rod

B

Short term memory
Novel object recognition (V-Maze)

Discrimination index

WT   Plcb1+/-

**

C

Short term memory
Novel object recognition (V-Maze)

Exploration time (s)

WT   Plcb1+/-

D

Locomotor activity

Horizontal counts in 2h

WT   Plcb1+/-

E

Rearing activity

Vertical counts in 2h

WT   Plcb1+/-

**

F

Elevated plus maze

Time spent in the open arms (s)

WT   Plcb1+/-

**

G

Elevated plus maze

Percentage of time in the open arms

WT   Plcb1+/-

**

H

Rota rod

Number of r.p.m.

WT   Plcb1+/-

I

Rota rod

Time in the rota rod (s)

WT   Plcb1+/-

Figure 1. Behavioral tests for phenotype characterization. (A) Timeline of the experimental sequence. (B,C) Short-term memory measured in the V-maze for novel object recognition task, (B) discrimination index and (C) exploration. (D) Locomotor and (E) rearing activity measured in actimetry boxes. (F) Anxiety-like behavior measured by the time spent in the open arms and (G) the percentage of time in the open arms in the elevated plus maze. Motor coordination measured in the rota rod test by the (H) number of r.p.m. that the mice performed in the average trials or the (I) mean time they remain in the apparatus. All data are expressed in median and interquartile range and individual data are shown (WT n=25; Plcb1+/- n=12).

A

B

Infusions
(cocaine, 0.5 mg/kg/inf)

☆   ***

C

Motivation for cocaine

D

Extinction of cocaine self-administration

&&&

E

Cue-induced
reinstatement of cocaine seeking

Figure 2. Operant conditioning maintained by cocaine in Plcb1+/-, wild-type (WT) and corresponding control saline mice. (A) Timeline of the experimental sequence. (B) The number of cocaine infusions (0.5 mg/kg/infusion) in both genotypes increased progressively across sessions during FR1 (repeated measures ANOVA, interaction between genotype x drug x sessions, $P<0.05$) and remained stable during FR3 and higher in mice trained with cocaine (repeated measures ANOVA, main effect of drug, ***$P<0.001$). The maximum number of infusions was reached on session 5 in both genotypes and was slightly superior in Plcb1+/- (36.35±1.47) than in WT (29.61±2.23) but post hoc analyses demonstrated that this difference was not significant. (C) The motivation for cocaine measured by the breaking point achieved in the progressive ratio schedule of reinforcement (4h) was equivalent in both genotypes trained with cocaine but higher than those trained with saline independently of the genotype (individual data with interquartile range, U Mann-Whitney, ##$P<0.01$ vs WT cocaine, \$\$$P<0.01$ vs Plcb1+/- cocaine). (D) Both genotypes also showed similar levels of extinction that decreased during sessions, but the curve was more pronounced in WT mice than in mutants (repeated measures ANOVA, interaction genotype x sessions, &&&$P<0.001$). (E) Decreased cue-induced reinstatement of cocaine seeking in Plcb1+/- mice was obtained compared to WT (individual data with interquartile range, U Mann-Whitney, #$P<0.05$, ##$P<0.01$ vs WT cocaine). All data are expressed in mean ± SEM when sessions are represented or median and interquartile range when individual data are shown (WT cocaine n=28-36; Plcb1+/- cocaine n=19-26; saline n=6 per genotype). Statistical details are included in Supplementary Table S1.

Figure 3. Gene expression changes in mPFC after cue-induced reinstatement of cocaine seeking comparing Plcb1+/- versus wild-type (WT) mice. (A) Selection of over-represented KEGG pathways (Kyoto Encyclopedia of Genes and Genomes) and (B) GO (Gene Ontology) identified by DAVID software among the differentially expressed genes. The number of genes with altered expression included in each category is indicated on the right side of the bar. In red, relevant pathways for the addictive process. (C) Gene network involved in cellular development, cellular growth and proliferation, nervous system development and function (score = 62). The green nodes in the pathway indicate genes with downregulated expression in Plcb1+/- identified in RNAseq.

Figure 4. Alterations in expression in the dopaminergic synapse in mPFC of Plcb1+/- mice after cue-induced reinstatement of cocaine seeking. Adapted from KEGG (Kyoto Encyclopedia of Genes and Genomes) pathways (mmu04728). Enriched pathway in DEGs in mPFC, 25 out of 131 genes in the pathway were differently expressed (Praw=2e-03; Pajd=0.02). In red, upregulated genes and in green, downregulated genes in mPFC. *Protein complexes including upregulated and downregulated genes. Correspondence between genes and proteins can be found in Supplementary Table S6.