Cognitive fluctuations surrounding sleep in daily living

Reto Huber[1] and Arko Ghosh[2]*

[1]Child Development Center, University Children's Hospital Zurich, Switzerland &
Department of Child and Adolescent Psychiatry and Psychotherapy, Psychiatric Hospital
University of Zurich, Switzerland

[2] Institute of Psychology, Cognitive Psychology Unit, Leiden University, The Netherlands

Correspondence:
8
Arko Ghosh
Institute of Psychology, Leiden University
Wassenarseweg 52, Leiden 2333 AK
The Netherlands
a.ghosh@fsw.leidenuniv.nl

Author contributions: AG conceived the study with the help of RH. AG gathered and analysed
the data. AG drafted the manuscript with the help of RH.

Acknowledgements: We thank the students at the Applied Cognitive Psychology master's
program at Leiden University for aiding in the data collection. We thank Andrew Westbrook
for sharing the smartphone data collected at The Donders Institute. We thank Leonardo Cohen
for his advice during the preparation of this manuscript and suggesting the title.

Funding: Investigator AG was supported by intramural funding from Leiden University and a
research grant from VELUX Stiftung during this study (no. 1283). Investigator RH was
supported by the University Medicine Zurich flagship grant "SleepLoop" and research grants
from the Swiss National Science Foundation (320030_153387 & 320030_179443).

Conflict of interest: Arko Ghosh is a co-founder of QuantActions Ltd, Lausanne, Switzerland.
This company focuses on converting smartphone taps to mental health indicators. Software
and data collection services from QuantActions were used to monitor smartphone activity.

Abstract

Cognitive output fluctuates surrounding sleep. These fluctuations are rarely investigated in subjects' daily living, partially due to the inaccessibility of sleep lab technology. Here we leverage the continuous and long-term smartphone touchscreen interaction logs to study the patterns of cognitive output surrounding sleep and contrasted these to the patterns underlying physical activity captured from wrist-worn actigraphy. The wearable also captured the ambient luminescence levels. According to spectral density analysis, both cognitive and physical activity was dominated by 24 h rhythm. However, the power of these rhythms differed in a domain-specific manner. The proxy measures of cognitive performance – tapping speed [TS], unlocking speed [US] and app locating speed [ALS] – contained lower-powered rhythm than physical activity. Still, this did not mean cognitive processes, in general, are low powered, as the amount of smartphone usage showed the strongest 24 h rhythm. Fixed 24 h (cosine) curve fits, revealed that the US peaked later in the day in comparison to physical activity and inter-individual differences in smartphone measures were not related to the experienced luminescence. Interestingly, subjects interacted with their smartphones during the actigraphy labelled sleep period. The cognitive measures in this period of fragmented sleep were worse in comparison to the hour before or after sleep. We conclude that the diurnal rhythms underlying cognitive activity in the real world are distinct from physical activity. Smartphones can seamlessly capture the dynamic fluctuations in cognitive output including when in bed and apparently asleep.

Introduction

We all experience that performance in daily tasks fluctuates strongly according to rhythms of different length. The best-studied rhythms in a laboratory setting are diurnal (1, 2). Essentially, cognitive functioning across the 24 h depends on both circadian rhythms and sleep-wake duration-dependent fluctuations of sleep pressure. Specifically, the endogen circadian rhythm significantly impacts various performance measures in a ~ 24 h rhythm, with performance being high during the day and deteriorating during the night-time (3, 4). On the other hand, performance suffers also under high sleep pressure condition as a function of the duration of prior wakefulness (3, 5). Additionally, sleep inertia, i.e. morning grogginess, reduces performance in the first 30 to 90 minutes after wake up (6). Ultimately the actual output of the system depends on the interplay between these different factors (circadian, sleep pressure, sleep inertia). We also know that misalignment of these internal processes which typically come along with extreme conditions (like shift work, jet lag or sleep restriction) can have dramatic consequences e.g. in terms of accidents and work-related errors (7). Though we have a substantial understanding of the fluctuations of cognitive functions based on the described factors, it is difficult to capture and study the consequences of these fluctuations in real life. Moreover, rhythms with a period longer than 24 hours are understudied because of the difficulty in capturing them in a laboratory setting. One promising way to overcome these empirical challenges is to leverage the daily digital interactions which may yield proxy measures of cognitive functions (8–10). According to one recent report leveraging the keypresses while on the web search engine, the speed of the keypresses fluctuates according to the time-of-the day akin to what has been found based on reaction time like tasks in the laboratory (11).

The unobtrusive sensing of cognitive performance in daily living conditions may provide some much-needed insights into the nature of the fluctuations under regular sleeping

conditions i.e., without instructed sleep deprivation. Our understanding of these regular

fluctuations is largely driven by the baseline measures obtained before sleep deprivation and

other protocols in controlled laboratory settings(12). For instance, sensorimotor cognition

captured by using a psychomotor vigilance performance (PVP) test remains stable through the

waking hours before the sleep deprivation but memory performance captured using a different

test fluctuates through the day in the same period (4, 13). The lack of data on cognitive

fluctuations also percolates to the periods in daily living where sleep is fractured i.e., during

the brief periods of interruptions in sleep due to either endogenous or exogenous reasons. This

gap perhaps persists simply due to the unpredictability of these events – in particular when the

interruptions are endogenously triggered - and thus out of reach of conventional performance

assessments. Regardless, if people are woken up from their sleep to perform tasks akin to

emergency response by a medical worker, executive functions, in particular, appear

vulnerable (14).

In addition to the unavailability of the measures from particular times of the day, the

generally sparse nature makes it difficult to estimate the precise nature of the rhythms that

govern cognition. Towards this, the dominating periods must be estimated without imposing a

priori waveform with a 24 h period. For instance, continuously measured physical activity

(actigraphy) can be processed using spectral density analysis to establish the inter-individual

differences in the near 24 h period, and these variations may be markers of clinical conditions

(15–17). For cognitive measures, the fundamental issue of which period and the

corresponding inter-individual differences remains unclear.

The ubiquitous smartphone interaction logs termed as tappigraphy can be converted

into proxy measures of cognitive performance (9) and may provide the dense data needed to

capture the fluctuations surrounding sleep. For instance, the tapping speed can be used as a

proxy for cognitive processing speed that is conventionally based on reaction time tasks

conducted in the laboratory or laboratory-like settings (8). Moreover, the near-continuous

nature of tappigraphy will help detect a range of rhythms as done for actigraphy. In a recent

study, we found that users typically engage on the smartphone through all the waking hours,

and also into sleep to an extent that the longest smartphone usage gaps at night provide a

reliable proxy for actigraphy estimated sleep (18). Interestingly, virtually all the users

engaged on the smartphone – even if briefly – during the actigraphy estimated sleep (18). This

phenomenon of sleep fracture, in particular, opens the opportunity to measure the cognitive

status during the period of sleep fragmentation by using tappigraphy.

In this study we defined three different tappigraphy parameters to ubiquitously capture

the cognitive processes akin to that measured in reaction experiments in the laboratory: (a) the

time that is taken to go from one touch to another, (b) the time that is taken to unlock the

phone and (c) the time that is taken to locate app icons on the home screen before launch. This

yielded a time series of measurements enabling spectral density analysis of the cognitive

fluctuations to identify the oscillations that dominate the cognitive outputs. We contrasted

these measures to actigraphy (including ambient luminescence) captured using a wrist-worn

wearable. Finally, through the combination of actigraphy and tappigraphy, we estimated the

cognitive status during the actigraphy labeled periods of pre-bed, 'sleep', and rise times.

Results

Periodicity in luminescence, physical activity, and tappigraphy

The wearable and tappigraphy signals fluctuated through the recording period (Fig. 1).

To quantify the periodicity of these fluctuations, we used the Lomb-Scargle method. The

population average traces were used to quantify the consistency of the behavioural patterns in

the sampled population. The periodograms revealed a consistent ~24 h periodicity across all

of the signals and a less prominent ~7 d periodicity for the ambient luminescence, physical

activity, smartphone usage (no. of touches), and the two proxy measures of cognitive

processing speed: tapping speed (TS) and unlocking speed (US) (Fig. 2).

By using the periodogram peaks we next estimated which period consistently

dominates the signals. First, we contrasted the location of the periodogram peaks with 24 h

periods using one sample t-tests to establish deviations from this anticipate period (α_{FWER} =

0.0083). The mean peak periodicity of ambient light fluctuations was 23.96 ($p = 0.002$, $t(69)$

= -3.27), for movements it was 24.19 ($p = 0.22$, $t(70) = 1.23$), for smartphone usage it was

23.97 ($p = 4.24$ x 10^{-6}, $t(183) = -4.74$), for TS 23.38 ($p = 0.01$, $t(184) = -2.57$), for US it

was 23.82 ($p = 0.43$, $t(184) = -0.78$) and for ALS it was 23.00 ($p = 0.01$, $t(156) = - 2.65$).

Next, we compared the peak locations across the signals to find that the periods were domain

dependent (Fig. 2, $p = 0.01$, $f(5,846) = 3.01$, ANOVA).

There were pronounced inter-parameter differences between the power estimates (i.e.,

in the normalised peak amplitudes of the periodogram, $p = 7.22$ x 10^{-74} , $f(5,848) = 87.02$,

ANOVA, Fig. 2c). The power of smartphone usage showed the highest ~ 24 h peaks relative

to any of the signals and the ALS showed the weakest peaks. Notably, the proxy measures of

cognitive processing (TS, US, and ALS) speed show a lower amplitude than the wearable

measures of luminescence or physical activity (Fig. 2).

Time-of-the-day effects in physical and cognitive signals

Upon establishing the dominance of $\sim$ 24 h rhythms in the physical and cognitive

fluctuations we next focused on how the gathered measures relate to the time-of-the-day. As

anticipated from the periodograms, the central tendencies revealed substantial time-of-the-day

fluctuations across the parameters (Fig. 3). To systematically address at which hour the

performance consistently peaked we relied upon cosinor analysis with a fixed 24 h waveform

(with individual fits corresponding to $p < 0.05$). The hour at which performance peaked (the

cosinor acrophase) depended on the parameter ($p = 0$, $f(5,673) = 25.64$, Watson-Williams

multi-sample test). Follow-up tests revealed that the peak for ambient light exposure

preceded all other measures, whereas the peak for unlocking speed lagged all other measures

except for ALS (Fig. 3). Although the hour of peak performance for the proxy measures of

cognitive processing occurred between 16 to 17 h, the differences between the peak vs. off-

peak performance was the most pronounced for TS ($p = 3.23 \times 10^{-29}$, $t(125) = 14.78$, paired t-

test), followed by US ($p = 3.48 \times 10^{-12}$, $t(117) = 7.76$) and the ALS ($p = 0.0062$, $t(39) = 2.89$).

Next, we addressed whether the inter-individual differences in the acrophase were

correlated across parameters (Fig. 3). In particular, we were interested in the putative

determinants of the cognitive processing speed. We used paired circular correlations to

address these relationships. Interestingly, the subtle variations in the ambient light acrophase

were not correlated to any of the measures. However, physical activity was correlated to only

one of the cognitive measures: US. Furthermore, smartphone usage was related to all

measures of the cognitive processing speed.

Similarly, we also followed-up on the $\sim$7-day rhythm identified using the periodogram

based on time-of-the week analysis to find systematic variance according to the day-of-the-

week, for all measures except the US and ALS (Supplementary Fig.). The day on which the

signals peaked varied according to the measured parameter ($p = 0$, $f(5,838) = 52.77$, Watson-

Williams multi-sample test). Furthermore, while physical activity and luminescence peaked

around the weekend, smartphone usage and tapping speed peaked around the weekday.

Poor cognitive performance during actigraphy labelled 'sleep'

The results from the time-of-the-day effects revealed that there were smartphone

interactions captured at all hours of the day in the sampled population. Struck by this

phenomenon we set out to explore how the cognitive performance fluctuates surrounding

sleep and, more uniquely, while in actigraphy defined sleep (i.e., while lying still in bed and

using the smartphone and yet classified as sleep by the Cole–Kripke algorithm on actigraphy).

From each individual, we pooled all the instances of usage at three different periods: *pre-bed*

defined as 1 h immediately preceding actigraphy defined sleep time, *bed* defined as during the

putative sleep (actigraphy defined) time and *rise* defined as 1 h immediately following the

sleep period (Fig. 4). We focused on the subset of the population where the parameter

estimation requirements were satisfied to yield a measure in each of these periods. TS

fluctuated only marginally across the three periods ($p = 0.07$, $f(2,65) = 2.73$, ANOVA).

Follow-up paired t-tests revealed marginal slowing in *bed* vs. *pre-bed* (p = 0.05, t(65) = -2.0),

and the *bed* vs. *rise* ($p = 0.05$, $t(65) = 2.04$). Furthermore, there was no difference between the

pre-bed vs. *rise* ($p = 0.59$, $t(65) = -0.54$). US fluctuated substantially through these periods (p

$= 5.62$ x 10-9, $f(2,62) = 22.23$), and the follow-up t-tests revealed a similar pattern as for TS

albeit more exaggerated. The *bed* period compared to *pre-bed* was substantially slower ($p =$

5.01 x 10^{-7}, $t(62) = -5.6$), and the *bed* period was also slower vs. *rise* period ($p = 6.12$ x 10^{-5},

$t(62) = 4.3$). There was a marginal difference between the *pre-bed* and *rise* period, with the

rise being slower ($p = 0.03$, $t(62) = -2.20$). ALS too fluctuated through these periods ($p = 5.50$

x 10^{-30}, $f(2,47) = 150.08$). Interestingly, the *bed* period was no different vs. the *pre-bed* period

($p = 0.52$, $t(47) = -0.64$). However, the *rise* period was faster than both *pre-bed* ($p = 0$, $t(47) =$

12.98) and *bed* periods ($p = 0$, $t(47) = 15.53$).

Discussion

The proxy measures of cognitive processes captured using tappigraphy revealed a range of systematic fluctuations and here we contrasted these to ambient luminescence and physical activity. The tappigraphy measures – like luminescence and physical activity – was dominated by ~ 24 h rhythm. But there were visible differences between the cognitive and the non-cognitive rhythms in terms of the exact period, the power and even the time-of-the day when they peak. Some of these differences also extended to the ~ 7 d rhythm. Intriguingly, the tappigraphy measures also allowed us to assess performance at odd hours including when in bed revealing a distinctly slow cognitive output.

As continuous measures related to cognitive output are mostly unexplored the general patterns of the signals observed in tappigraphy are of interest. The periodograms of the three proxy measures of cognitive processing speed (TS, US, ALS) revealed dominant powers at frequencies with periods of ~ 24 h and ~ 7 d. While all of the measures were dominated by 24 h cycles, there were substantial inter-individual differences and the exact period varied according to the parameter. This is in line with previous observations demonstrating deviations from the 24 h period in cognitively rooted performance parameters such as handgrip strength (19). We also observed ~ 7 d in the cognitive parameters of smartphone usage and TS. This supports the idea that the time information – both in terms of time-of-the-day and time-of-the-week – is encoded in cognitive output (20).

Luminescence is a primary *zeitgeber* for circadian physiological rhythms according to observations mostly in the sleep laboratory (21–23). Still, cognitive processes may not be faithfully tethered to this in the real world. First, the cognitive processing measures were less dominated by ~24 h rhythms in comparison to the luminescence captured at the wrist. Secondly, according to the fixed (24 h), cosinor analysis all of the proxy measures of

cognitive processing peaked (became faster) later in the day in comparison to the experienced

luminescence. Finally, population-level variance in tappigraphy time to peak was unrelated to

the variance in the experienced luminescence. We speculate that cognitive processes follow

diurnal rhythms that are partially independent of the experienced luminescence.

The processes that drive the well-documented rhythmicity in overall physical activity

may not entirely overlap with the oscillators underlying cognitive activity in the real world.

Towards this, cosinor analysis revealed some important separations between cognitive and

physical activity. The US peaked later than physical activity and the variations in time to peak

in physical activity were only related to one of the tappigraphy measures – US. The

differences between physical and cognitive activity were further widened in the ~ 7 d cycles

and while tappigraphy measures (smartphone usage & TS) peaked during weekdays, physical

activity (and experienced luminescence) peaked during weekends.

Interestingly, smartphone behaviour showed the strongest ~ 24 h power (signal

normalised) in comparison to the other measures considered here. This underscores the

habitual nature of smartphone behaviour where it may be more driven by daily cycles than

overall physical activity or the amount of light exposure. Nevertheless, the daily cycles were

less powerful for the proxy measures of cognitive output suggesting discord between

engaging in smartphone behaviour and the cognitive processing speed proxied on the

smartphone. This discord was vivid in the time-of-the-day analysis and by using the fixed 24-

hour cosinor analysis we found that UT peaked (was the fastest) later in the day in

comparison to smartphone usage. This raises the possibility that the need or desire to engage

on the smartphone and cognitive processes are out of sync – i.e., phase shifted.

By leveraging smartphone touchscreen behaviour, we could sample cognitive

fluctuations at the grey zone between sleep and wakefulness. People spontaneously interacted

with their smartphones in the actigraphy labelled sleep periods and we leveraged these

interactions to address the cognitive status in this 'sleep' period in comparison to the

performance 1 h before and after this period. Now, admittedly actigraphy can overestimate

sleep and people may engage on their smartphones while at rest in bed (18). Still, this

provided us with an opportunity to assess cognition in this period of sleep fracture. Across

the different proxy measures of cognitive processing, the performance was poor in this

obscure period. The mechanism underlying this as in sleep inertia vs. pressure could not be

clarified without polysomnography; it is possible the participants intermittently woke-up from

sleep in the bed (inducing inertia) and it is equally possible they may have remained still

without sleep (building sleep pressure). Still the current pattern suggests a dual contribution.

In the hour after sleep, inertia can be considered to be maximal and yet the performance at

sleep fracture was worse than this period. This suggests that an additional factor – such as

sleep pressure – is compounded with sleep inertia to additionally degrade cognitive output in

the obscure sleep period.   Conversely, in the hour before sleep, sleep pressure can be

considered to be maximal and yet TS and US degraded further at sleep fracture. Interestingly,

ALS did not degrade further and perhaps the underlying processes are particularly sensitive to

pressure rather than inertia. Such specific variations or lack there off are in keeping with the

general notion that sleep impacts cognitive processes in a domain-specific manner (24, 25).

Our approach of assessing cognitive fluctuations surrounding sleep in daily living

conditions raises some important questions. First, we observed different rhythms in cognitive

vs. the physical activity measures including the presumable *zeitgeber* of ambient light. It was

not clear if these asynchronies were introduced by smartphone behaviour or if they are

intrinsic properties captured on the smartphone. On a related note, the consequences of the

differences between smartphone usage vs. the proxy measures of cognitive processing too

need further exploration. These asynchronies may have important consequences for mental

and physical well-being (26). Second, there is much to be addressed on why and how people

behave at physical rest while in bed (i.e., actigraphy defined sleep). What are the cognitive

and behavioural processes underlying these behaviours that spontaneously occur so close to

sleep? Finally, unlike traditional cognitive testing, the parameters extracted from spontaneous

smartphone behaviour do not allow us to specify the cognitive processes with precession even

if they are correlated to conventional reaction time or even if they intuitively correspond to

cognitive processing speed. Nevertheless, meaningful cognitive processes are inherently

complicated and the approach of tappigraphy and the findings presented here is a key step to

help unravel that complexity.

Methods

Participants

A total of 189 individuals were sufficiently sampled (with 235 addressing the recruitment call, full data is made available and see below). To be included in the study the subjects self-reported that they were healthy and without any ongoing neurological disease or medication. The data collection and analysis were approved by the ethical committees of Leiden University (Psychology Research Ethics Committee) and the medical ethics committee of Arnhem-Nijmegen. All subjects provided informed consent for the study. The age (reported by 165 participants) was a median of 23 years (min, 16 and max, 45).

Actigraphy measurement

Actigraphy measures were obtained from a subset of participants (n = 79) and reported in a previous study (18). Participants wore GENEACTIV watches (Activinsights, Cambridgeshire, UK) on both the wrists, but only the measures from the left wrist were used here. The watches measured the 3-axis accelerometry along with the ambient luminesce and near body temperature, but only the former two measures were used here. The participants were instructed to wear the watches for 3 weeks continuously and this yielded measure lasting for a median of 21 days (min, 7 and max, 32). The 3 axis accelerometry was reduced by using $M = \sqrt{(x^2 + y^2 + z^2)}$, where M is the value used here and x, y and z correspond to the accelerations on the distinct axis. The Cole–Kripke algorithm was used to label sleep periods based on these measures as described in detail elsewhere along with the corresponding MATLAB codes (18, 27).

Smartphone measurements

The timestamp of touchscreen interactions and the corresponding app labels (as in

Facebook, Launcher screen, Weather) were recorded using an app running in the background

of the user's device (TapCounter, QuantActions, Lausanne, Switzerland). Based on this

labelled time-series of events the following parameters were estimated in hourly bins: (a)

Smartphone usage, in the form of a number of touchscreen interactions in each bin while the

phone was in an unlocked state, (b) tapping speed, in the form of the 25th percentile inter-

touch interval accumulated from all of the screen ON sessions in each bin, (c) unlocking

speed, in the form of 25th percentile inter-event interval between the two intervals, one, the

touchscreen turning ON and two, the touch on the unlocked screen and, (d) app locating

speed, as in the inter-touch interval between two consecutive touches on the home/launch

screen (identified using the corresponding app label) before the launching of any app. As with

the previous measures the 25^{th} percentile of the intervals in each hour bin was recorded. All

of the smartphone parameters were transformed by using log_{10}.

Estimating the periodogram and the corresponding metrics

Lomb-Scargle periodograms were estimated (MATLAB, Mathworks, Natick, USA)

and the power was scaled by the input variance. The periodogram was estimated between 0.05

and 12 cycles per day with a step of 0.001 cycles. The statistical significance ($\alpha = 0.001$) of

the power fluctuations were estimated against 0 using t-tests (LIMO EEG(28)) and multiple

comparisons corrected using the false discovery rate (FDR, also on LIMO EEG). Inputs

spanning longer than 10 days were used for this analysis. To compare the periodogram peaks

at ~1 cycle per day across the different smartphone and wearable parameters, the peak was

determined within the range of 0.7 and 1.6 cycles per day. First, the peaks from the different

measures were compared using one-way ANOVA (MATLAB, MathWorks, Natick, USA).

These were followed-up with t-tests comparing all possible pairs of measures. The tests were

corrected using Bonferroni correction of Family-Wise Error Rate (FWER, $\alpha = 0.05$, Victor

Martinez's Multiple Testing Toolbox as implemented MATLAB)(29). The 95% confidence

intervals were estimated using the inverse of Student's T cumulative distribution function

(MATLAB). Follow up t-tests after ANOVA to compare the periodogram peaks (location and

amplitude) across the different parameters were also corrected using FWER. Inputs spanning

longer than 7 days were used for this analysis block focused on ~1 cycle per day rhythm.

Finding peak-of-performance in time-of-the-day using cosinor analysis

357   The acrophase of the sine wave fits obtained using Cosinor.m (implemented in

MATLAB by Casey Cox)(30). Inputs spanning longer than 7 days were used for this analysis.

The time-of-the day fluctuations were compared across the different parameters using the

Parametric Watson-Williams multi-sample test (Circular Statistics Toolbox for MATLAB)

and as a follow-up, the same test was used in pairs. The inter-individual differences in the

acrophase across the different parameters were tested for correlation using circular correlation

(Circular Statistics Toolbox for MATLAB) (31). The statistical output was corrected for

multiple comparisons using the Bonferroni correction of Family-Wise Error Rate (FWER, $\alpha = $

0.05). The 95.0% confidence intervals were estimated using the same toolbox.

Finding peak-of performance in day-of-the-week

368   The hourly smartphone and wearable parameters as described above were sorted

according to the day of the week. Inputs spanning longer than 10 days were used for this

analysis. The mean value from each day of the week was used to derive the location of the

peak. These locations were converted into radians towards circular mean and confidence

intervals (95%). The measures were compared for day-of-the-week differences across the

different parameters as stated above for time-of-the day analysis, that is by using the

Parametric Watson-Williams multi-sample test and follow-up paired tests were corrected for

FWER.

Estimating performance surrounding sleep

The hourly smartphone parameters were time-locked to the sleep times estimated

using the Cole-Kripke algorithm on the actigraphy measures from the left wrist (27). The

median values in the hour bin preceding, during and after the sleep period was estimated from

each individual. To address if there were differences between these three measures were

contrasted using two-way ANOVA (MATLAB, MathWorks, Natick, and $\alpha = 0.05$).

Code availability

All of the codes that were programmed for this study are made available at the

repository link. Pre-existing codes are provided as links in the shared scripts. (Codes are

attached during peer-review)

Data availability

A reduced and de-identified data at 1 h bins are made available at the following

repository link. (This data is attached during peer-review)

Figure 1. The range of measures captured in daily living conditions in this study. Wearable

(a-b) and smartphone data from one exemplary subject monitored over 21 days. (a) The

amount of ambient light was captured on the wrist along with (b) the acceleration used here as

a proxy for physical activity. We quantified the smartphone behaviour in hour-long bins in

terms of (c) the number of touches, (d) the speed of the interactions measured as the interval

between subsequent touches (fastest 25^{th} percentile at each bin, tapping speed, TS), (e) the

time taken to unlock the screen (unlocking speed, US) and (f) the time to select an app icon

while on the home screen (app locating speed, ALS).

Figure 1

Figure 2. Lomb-Scargle periodogram reveals the periods that dominate the wearable

(luminescence and physical activity) and smartphone parameters. (a) Mean periodogram and

their corresponding confidence intervals (95%), with significant differences from zero

amplitude signal marked in red. (b) Mean peak periods derived from the periodograms and

their corresponding confidence intervals. (c) Mean peak normalised powers and their

corresponding confidence intervals. The size of the shapes in b & c correspond to the sample

size (see main text for the corresponding degrees of freedom).

Figure 2

Figure 3. Time of the day reflects on physical activity and processing speeds captured on the

smartphone. (a) Time of the day fluctuations in mean values and the corresponding 95%

confidence intervals. (b) Cosinor fits revealed the peak performance times on the 24 h clock,

the confidence intervals marked with triangles. (c) The comparison of processing at the

acrophase vs. off phase in the sample, with each individual represented with a connecting line.

(d) The inter-individual differences in the time of peak performance (cosinor, acrophase) are

related to each other. The circular correlation coefficient is shown for the significant

relationships.

Figure 3

Figure 4. Cognitive processing speed captured on the smartphone during actigraphy labelled

sleep. (a) The median processing speeds – in terms of tapping speed, unlocking time and

locating the app icon - captured for each individual accumulated over the observation period.

(b) The differences in mean processing speed captured during actigraphy estimated sleep in

contrast to the values accumulated in the hour before sleep and after sleep (95% confidence

interval). The size of the shapes in b correspond to the sample size (see the main text for the

corresponding degrees of freedom).

Figure 4

436   **Supplementary Figure.** Day of the week reflects on physical activity and processing speeds

437   captured on the smartphone. (a) Mean values and the corresponding confidence intervals

438   (95%). (b) The peak performance in terms of the best (mean) performing day of the week and

439   corresponding confidence intervals.

a

Luminiscance

Physical activity

Smartphone usage

Tapping speed

Unlocking speed

Finding app icon

Day of the week

b

Peak Performance time
(max. day of the weak)

Suppl. Fig.

References

1. D.-J. Dijk, J. F. Duffy, C. A. Czeisler, Circadian and sleep/wake dependent aspects of subjective alertness and cognitive performance. *J. Sleep Res.* **1**, 112–117 (1992).

2. K. P. Wright, C. A. Lowry, M. K. LeBourgeois, Circadian and wakefulness-sleep modulation of cognition in humans. *Front. Mol. Neurosci.* **5** (2012).

3. A. A. Borbély, S. Daan, A. Wirz-Justice, T. Deboer, The two-process model of sleep regulation: a reappraisal. *J. Sleep Res.* **25**, 131–143 (2016).

4. C. Schmidt, F. Collette, C. Cajochen, P. Peigneux, A time to think: Circadian rhythms in human cognition. *Cogn. Neuropsychol.* **24**, 755–789 (2007).

5. J. C. Lo, J. L. Ong, R. L. F. Leong, J. J. Gooley, M. W. L. Chee, Cognitive Performance, Sleepiness, and Mood in Partially Sleep Deprived Adolescents: The Need for Sleep Study. *Sleep* **39**, 687–698 (2016).

6. A. T. Wertz, J. M. Ronda, C. A. Czeisler, K. P. Wright, Effects of Sleep Inertia on Cognition. *JAMA* **295**, 159–164 (2006).

7. M. E. Jewett, *et al.*, Time course of sleep inertia dissipation in human performance and alertness. *J. Sleep Res.* **8**, 1–8 (1999).

8. M. Balerna, A. Ghosh, The details of past actions on a smartphone touchscreen are reflected by intrinsic sensorimotor dynamics. *Npj Digit. Med.* **1**, 4 (2018).

9. Insel TR, Digital phenotyping: Technology for a new science of behavior. *JAMA* **318**, 1215–1216 (2017).

10. R. W. White, E. Horvitz, Population-scale hand tremor analysis via anonymized mouse cursor signals. *Npj Digit. Med.* **2**, 1–7 (2019).

11. T. Althoff, E. Horvitz, R. W. White, J. Zeitzer, Harnessing the Web for Population-Scale Physiological Sensing: A Case Study of Sleep and Performance in *Proceedings of the 26th International Conference on World Wide Web*, WWW '17., (International World Wide Web Conferences Steering Committee, 2017), pp. 113–122.

12. K. Blatter, C. Cajochen, Circadian rhythms in cognitive performance: Methodological constraints, protocols, theoretical underpinnings. *Physiol. Behav.* **90**, 196–208 (2007).

13. E. C.-P. Chua, *et al.*, Sustained Attention Performance during Sleep Deprivation Associates with Instability in Behavior and Physiologic Measures at Baseline. *Sleep* **37**, 27–39 (2014).

14. J. Horne, R. Moseley, Sudden early-morning awakening impairs immediate tactical planning in a changing 'emergency' scenario. *J. Sleep Res.* **20**, 275–278 (2011).

15. A. Germain, D. J. Kupfer, Circadian rhythm disturbances in depression. *Hum. Psychopharmacol. Clin. Exp.* **23**, 571–585 (2008).

477   16.   R. T. Krafty, *et al.*, Measuring Variability in Rest-Activity Rhythms from Actigraphy
478   with Application to Characterizing Symptoms of Depression. *Stat. Biosci.* **11**, 314–333
479   (2019).

480   17.   Y. Leng, *et al.*, Association of Circadian Abnormalities in Older Adults With an
481   Increased Risk of Developing Parkinson Disease. *JAMA Neurol.* (2020)
482   https:/doi.org/10.1001/jamaneurol.2020.1623 (June 17, 2020).

483   18.   J. N. Borger, R. Huber, A. Ghosh, Capturing sleep–wake cycles by using day-to-day
484   smartphone touchscreen interactions. *Npj Digit. Med.* **2**, 1–8 (2019).

485   19.   I. E. Ashkenazi, A. Reinberg, A. Bicakova-Rocher, A. Ticher, The genetic background
486   of individual variations of circadian-rhythm periods in healthy human adults. *Am. J.*
487   *Hum. Genet.* **52**, 1250–1259 (1993).

488   20.   J. Huttenlocher, L. V. Hedges, V. Prohaska, Memory for day of the week: A 5 + 2 day
489   cycle. *J. Exp. Psychol. Gen.* **121**, 313–325 (1992).

490   21.   S. L. Chellappa, M. C. M. Gordijn, C. Cajochen, "Chapter 7 - Can light make us bright?
491   Effects of light on cognition and sleep" in *Progress in Brain Research*, Human Sleep
492   and Cognition Part II., H. P. A. Van Dongen, G. A. Kerkhof, Eds. (Elsevier, 2011), pp.
493   119–133.

494   22.   R. A. Wever, Light Effects on Human Circadian Rhythms: A Review of Recent Andechs
495   Experiments. *J. Biol. Rhythms* **4**, 49–73 (1989).

496   23.   T. A. Bedrosian, R. J. Nelson, Timing of light exposure affects mood and brain circuits.
497   *Transl. Psychiatry* **7**, e1017–e1017 (2017).

498   24.   T. M. Burke, F. A. J. L. Scheer, J. M. Ronda, C. A. Czeisler, K. P. Wright, Sleep inertia,
499   sleep homeostatic and circadian influences on higher-order cognitive functions. *J. Sleep*
500   *Res.* **24**, 364–371 (2015).

501   25.   M. Ferrara, L. De Gennaro, M. Casagrande, M. Bertini, Selective slow-wave sleep
502   deprivation and time-of-night effects on cognitive performance upon awakening.
503   *Psychophysiology* **37**, 440–446 (2000).

504   26.   E. J. W. Van Someren, R. F. Riemersma-Van Der Lek, Live to the rhythm, slave to the
505   rhythm. *Sleep Med. Rev.* **11**, 465–484 (2007).

506   27.   R. J. Cole, D. F. Kripke, W. Gruen, D. J. Mullaney, J. C. Gillin, Automatic sleep/wake
507   identification from wrist activity. *Sleep* **15**, 461–469 (1992).

508   28.   C. R. Pernet, N. Chauveau, C. Gaspar, G. A. Rousselet, LIMO EEG: A Toolbox for
509   Hierarchical LInear MOdeling of ElectroEncephaloGraphic Data. *Comput. Intell.*
510   *Neurosci.* (2011) https:/doi.org/10.1155/2011/831409 (January 25, 2019).

511   29.   Y. Benjamini, D. Yekutieli, The Control of the False Discovery Rate in Multiple Testing
512   under Dependency. *Ann. Stat.* **29**, 1165–1188 (2001).

513   30.   W. Nelson, Y. L. Tong, J. K. Lee, F. Halberg, Methods for cosinor-rhythmometry.
514   *Chronobiologia* **6**, 305–323 (1979).

515   31. P. Berens, CircStat: A MATLAB Toolbox for Circular Statistics. *J. Stat. Softw. Vol 1*
516   *Issue 10 2009* (2009).

517