Upregulation of gluconeogenesis leads to increased tolerance to antibiotics

targeting the MreB elongosome in *E. coli*

Brody Barton, Addison Grinnell, Randy M. Morgenstein[#]

Department of Microbiology and Molecular Genetics, Oklahoma State University,

Stillwater, OK

Running title: Gluconeogenesis and A22 resistance in *E. coli*

[#] Address correspondence to Randy Morgenstein, randy.morgentein@okstate.edu

Abstract word count: 146

Manuscript word count: 4352

Abstract

Antibiotic resistant bacteria are a global threat to human health. One way to combat the rise of antibiotic resistance is to make new antibiotics that target previously ignored proteins. The bacterial actin homolog, MreB, is highly conserved among rod-shaped bacteria and essential for growth, making MreB a good focus for antibiotic targeting. Therefore, it is imperative to understand mechanisms that can give rise to resistance to MreB targeting drugs. Using the MreB targeting drug, A22, we show that changes to central metabolism through deletion of TCA cycle genes, leads to the upregulation of gluconeogenesis resulting in cells with an increased minimal inhibitory concentration to A22. This phenotype can be recapitulated through the addition of glucose to the media. Finally, we show that this increase in minimal inhibitory concentration is not specific to A22 but can be seen in other cell wall targeting antibiotics, such as mecillinam.

Importance

The spread of antibiotic resistance has made bacterial infections harder to treat. Finding new targets for antibiotic development is critical to overcoming the variety of resistance mechanism that are already crippling our ability to treat infections with current antibiotics. The bacterial actin homolog MreB is a good target for new antibiotic development because it is essential for growth and highly conserved among rod-shaped pathogens. The significance of this research is in understanding the mechanisms cells can develop toward the inhibition of MreB to better understand how to make MreB targeting antibiotics in the future.

Introduction

The spread of antibiotic resistance among bacteria is a global human health concern (1). In May 2015 the World Health Organization (WHO) outlined a global action plan on antimicrobial resistance. Due to the spread of antibiotic resistance among pathogens against currently used therapeutics, development of new antibiotics is required (2-4). One strategy for the development of new antimicrobials is the modifications of current drugs, such as has been done with the penicillin family of antibiotics; however resistance to these new drugs requires the simple modification of beta-lactamase enzymes (5). For example, the second-generation penicillin, methicillin, was developed in 1959 with resistant strains isolated only two years later (6). Another strategy is to develop new therapeutics against novel bacterial targets, for which resistance has not yet been selected, which should increase the longevity of these new antibiotics.

The bacterial actin homolog, MreB, is a highly conserved and essential protein necessary for shape and growth in many rod-shaped human pathogens, including *Escherichia coli, Salmonella enterica, Pseudomonas aeruginosa,* and *Vibrio cholerae* (7, 8). Therefore, MreB is a good target candidate for new antibiotic development. While there are no current FDA-approved MreB-targeting antibiotics, the compound A22 is used in laboratory settings to study the role of MreB in the cell. A22 binds adjacent to the ATP binding pocket of MreB and alters its polymerization dynamics (9-13). MreB directs the localization of cell wall synthesis, and disruption of MreB, either through the use of A22 or genetic modifications, causes cells to become misshapen and lyse. One

issue with developing A22-like compounds for human use is the high rate of suppressor

mutations leading to A22 resistance, which almost always map to MreB (12, 14-16).

Therefore, it is imperative that we understand mechanisms of resistance against MreB-

targeting antibiotics, including mutations outside of MreB, before exploring the

expensive process of MreB-targeted drug development.

Mutations in genes encoding enzymes of the tricarboxylic acid cyle (TCA) cycle

have been shown to provide antibiotic tolerance. It was shown that a mutation in *icd*,

which encodes isocitrate dehydrogenase, results in toxic intermediates which activate

efflux pumps. *icd* mutants have increased resistance to nalidixic acid (NA), a DNA-

targeting drug, because the intermediates that accumulate due to this mutation activate

the AcrAB-TolC efflux pump, which can pump NA out of the cell (17). More recently, it

was shown that antibiotic tolerance was induced by inhibiting ATP production (18, 19).

One way this could be achieved is by blocking enzymes in the TCA cycle such as

aconitase (*acnB*) or succinate dehydrogenase (*sdhA*), which lowers the reducing power

available for oxidative respiration (20). In addition to the connection between the TCA

cycle and antibiotic resistance, a connection between the TCA cycle and cell size has

been shown in *Caulobacter crescentus*. In *C. crescentus*, mutations that result in the

accumulation of α-ketoglutarate cause cell shape defects by reducing the synthesis of

cell wall precursors (21).

Here, we performed a screen of *E. coli* deletion mutants to find mutations that

caused an increase in the minimal inhibitory concentration of A22 (MIC_{A22}) in *E. coli*.

This screen uncovered many possible pathways that could lead to a higher MIC_{A22}.

Because of the well-known connection between cell shape and size and metabolism

and the connections between the TCA cycle and antibiotic tolerance, we focused our

efforts on understanding how disruption of the TCA cycle protein, Mdh, leads to a higher

MIC_{A22}. We show that deletion of *mdh* results in a higher MIC_{A22} through a novel

mechanism that activates gluconeogenesis, causing an increase in the levels of cell wall

precursors. Additionally, we found that this higher MIC is phenocopied by the addition of

glucose to the growth medium and holds true when cells are treated with mecillinam, an

inhibitor of the cell wall synthesis enzyme penicillin binding protein (PBP) 2.

Results

Metabolic gene deletions cause an increase in MIC_{A22}. We wanted to determine if there

were gene deletions that result in a cell's ability to live without the essential bacterial

actin homolog, MreB. It is known that mutations that cause an increase in expression of

the cell division genes *ftsZAQ* suppresses the growth defects of deletions of many cell

shape determinants, including MreB (22-24). Because suppressor mutations arise

quickly when *mreB* is deleted, we used the MreB depolymerizing drug A22 to mimic the

loss of MreB in order to determine if cells can bypass the need for MreB (25). A random

subset of the Keio deletion library was screened for gene deletions that cause an

increase in the concentration of A22 that results in less than half the growth ($O.D._{600}$)

compared to the LB only control (henceforth referred to as, MIC_{A22}). (26). Of the 276

gene deletions from the Keio collection screened, we found 86 gene deletions (31%)

that showed a higher MIC value than wild-type (WT) cells (Table S1). Gene ontology

analysis revealed that the largest class of mutants with increased MIC_{A22} were in genes

involved in cellular metabolism (Fig. 1A), and, of these, two were involved in the TCA

cycle.

Our initial subset of the Keio collection used in the screen contained three strains

with deletions in TCA cycle genes: malate dehydrogenase (*mdh*), isocitrate

dehydrogenase (*icd*), and fumarase (*fumE*) (Fig. 1BC). Of these, only *mdh* and *icd*

showed a higher MIC_{A22} than WT cells from our initial screen (Table S1). We confirmed

the presence of the deletion in each of these genes by PCR and retested the MIC_{A22}.

We found that the *mdh* deletion resulted, but not the *icd* deletion resulted in an increase

in the MIC_{A22} (Fig. 1B). This suggests that some of the 86 genes in table S1 are false

positives. However, *mdh* was found both in the initial screen and after PCR confirmation

to have a large increase in the MIC_{A22}.

FumE is not the major fumarase in *E. coli* and its function was only discovered

through a large-scale metabolomics screen, before which it was predicted to be a

transcription factor (27, 28). Because the *fumE* deletion did not show an increased

MIC_{A22}, we tested if deletion of the major aerobic fumarase, FumA, or other enzymes

involved in the TCA cycle increase the MIC_{A22} by going back to the Keio collection and

performing MIC assays on citrate synthase (*gltA*), aconitate hydratase (*acnB*), 2-

oxoglutarate dehydrogenase (*sucA*), succinyl-CoA synthetase (*sucC*),

succinate:quinone oxidoreductase (*sdhA*), and the major anaerobic fumarase (*fumA*)

after PCR confirmation of the deletions (29). Of these six tested mutants, only two

showed a higher MIC_{A22}: *acnB* and *sucC* (Fig 1C).

Previous studies have shown changes to antibiotic sensitivity when mutations are

made in the genes involved in the TCA cycle (17, 20, 21). *acnB* mutations are thought

to alter antibiotic susceptibility through changes in cellular ATP levels (19), a

mechanism that might explain our observations. To our knowledge, neither mutations in

mdh nor *sucC* have been reported to cause a decrease in antibiotic susceptibly. In this

paper, we focus our efforts on understanding the effects of the *mdh* mutation on A22

tolerance, as we do not expect the *mdh* mutation to function in ATP synthesis because

the action of Mdh occurs after the steps in the TCA cycle where reducing power is

created.

Slow growth is known to help cells grow without MreB (22). Growth curves of WT

and *mdh* cells show that in our test conditions (LB medium) there is no appreciable

change in growth rates, indicating that slow growth is not contributing to the increase in

MIC_{A22} (Fig. S1) and that ATP production is not severely affected in this strain. Because

the growth rate was the same between WT and *mdh* cells and increased tolerance to

A22 is not a universal feature of TCA cycle gene deletions (Fig. 1BC), we hypothesize

that the Mdh enzyme or changes in malate levels (the substrate of Mdh) is the most

likely cause of the increased MIC_{A22} seen in the *mdh* mutant.

A22 disrupts MreB in mdh cells. Cells could show an increased MIC_{A22} for three

reasons: 1) A22 cannot accumulate in the cell to appropriate concentrations, 2) A22

cannot enter the cell due to membrane changes, or 3) MreB polymers are protected

from A22. With respect to the first option, it is possible that an intermediate accumulates

in the *mdh* mutant that activates an efflux pump that removes A22 from the cells or that

A22 is degraded faster in *mdh* cells due to changes in the metabolic activity of different

enzymes. To determine if any of the above mechanisms are the cause for the increased

MIC_{A22} in *mdh* cells, we imaged WT cells, an MreB point mutant resistant to A22, and

mdh cells after growth in LB alone, LB with a sublethal concentration of A22 (1 µg/ml)

for 4 hours, or LB spiked with 10 µg/ml of A22 for 2 hours after 2 hours of growth. As

expected, both A22 treatments cause WT cells to become round, whereas the MreB

point mutant remains as a rod regardless of A22 treatment (16). However, *mdh* cells

became round, like WT cells, confirming that A22 is able to accumulate in the mutant

cells at a high enough concentration to inhibit MreB polymers (Fig. 2S). These data

further suggest that either Mdh or malate levels are responsible for the increased

MIC_{A22}.

mdh mutants up-regulate gluconeogenesis leading to an increase in MIC_{A22}. There is

precedent for metabolic proteins having additional cellular roles separate from their

known enzymatic activities (moonlighting). For example, aconitase, an enzyme in the

TCA cycle, that forms D-*threo*-isocitrate from citrate with aconitate as an intermediate,

also plays a role in iron sensing regulation by binding to specific mRNAs for post-

transcriptional regulation (30, 31). Additional moonlighting enzymes have been shown

to help regulate cell shape/size based on changes in nutrient availability (recently

reviewed (32, 33)). An example in *E. coli* is OpgH, a metabolic enzyme which can

directly interact with the main division protein, FtsZ, to help modulate cell length in

different nutrient conditions (34).

As a deletion of *mdh* would inhibit both its normal enzymatic function and any

moonlighting activity, we performed metabolomic analysis to identify changes in

metabolites in *mdh* cells to determine if these changes affect the MIC_{A22}. Malate levels

act as a positive control in these experiments, as malate should be increased in cells

with a defective malate dehydrogenase. As expected, malate levels were higher in the

mdh mutant than WT cells (Fig. 2A). In the *mdh* mutant, the increased levels of malate

should increase flux into pyruvate through malate dehydrogenase (*maeAB*), which is

observed by elevated pyruvate levels in the *mdh* mutant (Fig. 2B). This increase in

pyruvate could either flow back into the TCA cycle or into gluconeogenesis to produce

sugar (35). We saw an increase in glucose-6-phosphate (G6P), the end product of

gluconeogenesis, in the *mdh* mutant, which could only come from increased

gluconeogenesis because the cells were grown in LB medium without any sugar (Fig.

2C). G6P can be converted into fructose-6-phosphate which can be used to form N-

acetylglucosamine (NAG), while pyruvate can be converted to phosphoenolpyruvate,

another metabolite used in the synthesis of cell wall precursors (36).

Thus, the increased levels of both G6P and pyruvate led us to hypothesize that

the *mdh* deletion causes an increase in gluconeogenesis resulting in increased levels of

cell wall precursor which may cause an increase in cell wall synthesis (37). To this end,

we looked for changes in the level of cell wall precursors in the metabolomics data and

saw an increase in UDP-NAG, confirming our hypothesis (Fig. 2D). These results

suggest that increasing the amount of cell wall precursors can suppress the lethal

effects of MreB disruption and that there is not a moonlighting function of Mdh.

The addition of glucose mimics deletion of mdh. To confirm our hypothesis that the

increase of cell wall precursors caused by an increase in gluconeogenesis is the reason

for a higher MIC_{A22}, we sought to increase cell wall precursor synthesis through another

mechanism. Because G6P levels were increased in *mdh* cells, we reasoned that the

addition of glucose to the medium would have a similar effect on the levels of cell wall

precursors as deleting *mdh*. Additionally, glucose has been shown to compensate for

mutations in aconitase to restore antibiotic susceptibility (20).

MIC assays were performed on WT cells grown on LB medium with increasing

amounts of glucose or α-methyl-D-glucoside (αMG), a nonhydrolyzable form of glucose

(38, 39). While very low levels (<0.125%) of glucose had no effect on the MIC_{A22} in WT

cells, even glucose levels as low as 0.125% resulted in a statistically significant increase

in the fold change of the MIC_{A22} in cells grown with glucose vs LB alone compared with

cells grown with αMG vs LB (Fig. 3). An increase in the glucose levels resulted in even

higher MIC_{A22} fold changes. The significant increase in MIC_{A22} during glucose versus

αMG treatment suggests that the changes are not due to osmotic protection caused by

the sugar. These results further support our hypothesis that increasing the levels of cell

wall precursors leads to an increase in MIC_{A22}.

Deletion of mdh leads to an increased MIC to mecillinam. The major role of MreB is to

direct the location of the cell wall synthesis enzymes. It has been suggested that MreB

forms a complex with the SEDS family of cell wall synthesis enzymes (40-42).

Specifically, recent work has suggested that MreB interacts with RodA and PBP2 to

regulate cell elongation, while PBP1A/1B work independently of MreB (40). Because the

mdh deletion results in cells with a higher MIC_{A22}, it stands to reason that cells might

also have a higher MIC against mecillinam, a PBP2-targeting antibiotic (43). To that end,

we performed MIC assays on mecillinam and other antibiotics that target cell wall

synthesis proteins to determine how general the effects of the *mdh* deletion are on the

MICs of cell wall-targeting drugs.

We performed MIC assays with a variety of cell wall-targeting antibiotics that

inhibit either cell elongation (ampicillin, mecillinam, cefsulodin) or cell division

(cephalexin), by blocking the activity of different PBPs involved in cell wall synthesis. An

MIC fold change >1 indicates that the *mdh* deletion causes cells to be more resistant

and a fold change <1 indicates that WT cells are more sensitive to the specified

antibiotic. The only other cell wall-targeting antibiotic to which the *mdh* deletion shows

an increased MIC is mecillinam, which targets the MreB complex partner PBP2 (Fig. 4).

The fact that the *mdh* deletion strain does not have a higher MIC_{cef} adds more evidence

that PBP1A/B are not part of the MreB complex because cefsulodin specifically targets

PBP1A/1B. Cephalexin targets PBP3 (FtsI) which is part of the division machinery, a

different complex than the MreB-elongosome. We did not see an increase in the MIC_{ceph}

in the *mdh* deletion, suggesting that an increase in gluconeogenesis cannot overcome

cell division inhibition. Ampicillin targets multiple PBPs, with the highest affinity for

PBP4, PBP3, and PBP2 in that order (44, 45). Because ampicillin binds to PBP3 tightly

and there was no increase in the MIC_{Ceph}, it leads to reason that deletion of *mdh* would

not have an effect on the MIC_{Amp}. These results suggest that the up-regulation of

gluconeogenesis caused by the *mdh* deletion specifically causes an increase to the MIC

of drugs that target the MreB elongation synthesis complex (A22 and mecillinam), but

not to other cell wall-targeting antibiotics.

Inactivation of aconitase (*acnB*) and succinyl-CoA synthetase (*sucC*), leads to an

increase in the MIC_{A22} (Fig. 1D); therefore, we tested whether deletions of these genes

also leads to an increase in the MIC_{mec}. The *acnB* mutation displays an increased

MIC_{mec}, while deletion of *sucC* does not (Fig. 3S). This suggests that the role of SucC in

A22 resistance is different than that of Mdh. However, it is unclear if AcnB acts through

a general antibiotic resistance mechanism via ATP depletion or through a specific

MreB-elongosome mechanism.

Up-regulation of gluconeogenesis leads to an increased MIC to mecillinam. The addition

of glucose to the media was able to phenocopy the effects of deleting *mdh* on the

MIC_{A22} (Fig. 3). Because we saw a similar increase in the MIC_{Mec} as we saw with the

MIC_{A22} in the *mdh* deletion, we wanted to confirm that the addition of glucose would also

lead to an increased MIC_{Mec} in WT cells. The addition of glucose leads to a statistically

significant increase in the MIC_{Mec} versus αMG (Fig. 3). Taken together, these data

indicate that an increase in cell wall precursors, either through an increase in

gluconeogenesis or the addition of glucose to the medium, leads to an increased MIC of

antibiotics that specifically target the MreB elongosome, but not other cell wall synthesis

enzymes.

Discussion

Our findings demonstrate that changes in cellular metabolism can have drastic

effects on antibiotic resistance. An increase in gluconeogenesis, through mutations of

metabolic enzymes or the addition of glucose to the growth medium, leads to increased

MICs of both the MreB targeting drug, A22, and the PBP2-targeting drug, mecillinam.

This effect appears to be specific for the MreB cell wall synthesis complex as we do not

see an increase in MIC of antibiotics that target other enzymes in the cell wall synthesis

pathway.

Cellular metabolism is a complicated and interconnected network. It is difficult to

predict how mutations in one pathway may lead to changes in other pathways in a cell.

While trying to understand novel mechanisms for A22 resistance, we found that loss of

MDH, an enzyme in the TCA cycle, leads to an increase in the MIC_{A22} by inducing

gluconeogenesis (Fig. 1). The induction of gluconeogenesis leads to the overproduction

of cell wall precursors (Fig. 3). MreB is thought to control where cell wall synthesis

occurs, with the localization of MreB, enriched in the middle of the cell, correlating with

the sites of new cell wall growth in *E. coli* (46, 47). When MreB is disrupted, the location

of cell wall synthesis becomes random, causing misshapen cells and eventually cell

death due to lysis (46, 48, 49). We hypothesize that the increase in cell wall precursors

allows for more cell wall synthesis and cell survival by prohibiting cell lysis. The fact that

both inhibition of MreB with PBP2 is suppressed through this mechanism, suggests that

the increase in cell wall precursors allows for a secondary synthesis system to dominate

(see below). Below, we discuss the implications for these results on understanding

bacterial cell wall synthesis and the development of drug treatment plans.

MreB is complexed with PBP2 but not class A PBPs.

The bacterial cell wall is a macromolecule made up of sugars crosslinked by

proteins to provide shape and support to the cell. In rod-shaped bacteria, such as *E.*

coli, B. subtilis, and C. cresentus, MreB is thought to form the main protein of the

elongosome complex consisting of multiple PBPs, MreBCD, RodA, and RodZ. Work in

the above species led to the hypothesis that MreB interacts with the cytoplasmic cell

wall synthesis components and helps to direct the localization of the periplasmic-acting

enzymes (16, 23, 46, 50-54). MreB is a cytoplasmic protein that is associated with the

inner face of the cytoplasmic membrane (9). How does this cytoplasmic protein

communicate with the extracellular cell wall? Our recent work showed that the integral

membrane protein RodZ acts as a linker between MreB and the cell wall synthesis

machinery, enabling MreB dynamics and assembly (16, 47). When *rodZ* is deleted, cells

grow as irregular round cells because the localization of MreB and the number of MreB

polymers are disrupted.

Recent work has questioned the model that the MreB elongosome is a large

complex composed of both the bifunctional class A PBPs (aPBPs), and monofunctional

class B PBPs (bPBPs). The aPBPs contain both transglycosylation and transpeptidation

activities, and thus should be functional alone, while the bPBPs, such as PBP2,

possess only transpeptidation activity. However, PBP2 is essential in *E. coli*, and in both

B. subtilis and *E. coli*, cell elongation has been shown to proceed without the aPBPs

(40, 42). This suggests that the MreB complex is not as large as previously thought. If

bPBPs can cause cell elongation without aPBPs, then there must be another enzyme

capable of transglycosylation reactions. RodA has been suggested to be the

transglycosylase that works with PBP2 (bPBP) (40, 42). These authors also showed

that the RodA-PBP2 complex works with the MreB cytoskeleton while the aPBPs act

separately from this complex. Our previous work studying MreB dynamics suggested

that RodZ helps modulate MreB motion through interactions with RodA/PBP2, further

confirming the idea that they are in a complex together (16). Here we show that

resistance to antibiotics that target MreB (A22) or PBP2 (mecillinam), but not aPBPs

(cefsulodin) are less effective when gluconeogenesis is upregulated (Fig. 3-4).

The results together reinforce the model that MreB forms a cell wall synthesis

complex with PBP2 but not PBP1A/B, and suggest that cell wall synthesis deriving from

the activity of aPBPs may be upregulated when there is an abundance of precursor

molecules in the cell. Upregulation of an aPBP-specific synthesis mechanism would

explain how cells can grow when MreB or PBP2 are inhibited. How aPBP activity is

modulated when other cell wall synthesis mechanisms are inhibited is currently not

known but would be an interesting topic for further study.

Drug efficacy is affected by the metabolic state of cells.

Because antibiotic resistance is more prevalent now than ever before, it is

essential that we understand how and when to use the antibiotics to which bacteria are

susceptible. Antibiotic resistance normally comes from genetic changes in a cell. This

can be through the acquisition of degrading enzymes, such as beta-lactamases for

resistance to penicillins; the modification of a drug target, such as mutations of the *rpsL*

gene for resistance to streptomycin; or a cell's ability to limit the intracellular

concentration of an antibiotic, such as through upregulation of efflux pumps (55-57).

The efficacy of antibiotics is higher against growing cells, as slow growing cells show an

increase tolerance to many antibiotics (58, 59). When only a subfraction of the

population is tolerant to antibiotic exposure, the surviving cells are termed 'persisters'

(60).

Here we show that mutations in metabolic genes such as *mdh,* seemingly

unrelated to the target of both the antibiotics, A22 and mecillinam, leads to increases in

antibiotic susceptibility (Fig.1 and 4). Additionally, the addition of glucose to the growth

media phenocopies the *mdh* mutation and leads to an increase in the MIC of both

drugs. Previous research has shown that while persister cells are more tolerant of

aminoglycoside antibiotics, the addition of different sugars specifically potentiated the

effects of gentamycin and other aminoglycosides, resulting in a reduction of persister

cell survival after treatment (61). The work presented here suggests that the addition of

sugar has the opposite effect on the efficacy of mecillinam and decreases its

effectiveness. Additionally, inhibiting ATP production can lead to increases in antibiotic

tolerance which can be bypassed through the addition of glucose (18-20).Therefore, it

will be imperative when designing an antibiotic treatment protocol that one remembers

that there is not a single solution. While the addition of sugars may help to treat cells

when using aminoglycosides by killing both actively growing cells and persister cells, we

show the opposite is true when using mecillinam, as the addition of sugar actually

increases the required dose (MIC).

Furthermore, gluconeogenic metabolism has been shown to promote infection.

Colonization by *E. coli* of both cows and mice has been shown to increase during

gluconeogenic growth, especially when undergoing competition in the gut (62, 63).

Moreover, a gluconeogenic environment leads to an increase in virulence factor

expression of enterohemorrhagic *E. coli* (64). Therefore, the use of metabolites to

potentially help eliminate a subset of cells, such as persister cells, may have the

unintended consequences of both increasing colonization and virulence of other cells,

and diminishing the efficacy of other antibiotics.

The properties that make MreB an attractive drug target — conservation across

many pathogens and essentiality — are true for other members of the MreB complex,

including RodA and RodZ. Our results suggest that drugs that target these proteins

would be less effective if gluconeogenesis is activated. As new antibiotics are

developed against novel targets, these results show it is important to think about the

broad function of the target protein as the efficacy of a drug may be affected in

seemingly unknown ways as we have shown that antibiotics tolerance to MreB can be

increased through a mutation in the TCA cycle gene *mdh*.

Material and Methods

Bacterial Growth. Bacteria were grown using standard laboratory conditions. Cultures

were grown overnight in LB medium (10g/L NaCl, 10g/L tryptone, 5g/L yeast extract)

and subcultured in the morning 1:1000 and grown to exponential phase (O.D.$_{600}$ 0.3-0.6)

at 37°C. Keio collection mutants were moved in MG1655 using P1 transduction.

Transductants were selected for on Kanamycin (30µg/mL) and confirmed by PCR.

MIC assay.

Optical densities of all cultures used in the minimum inhibitory concentration assays

(MIC) were checked at 600 nm (OD$_{600}$) with a Thermo Scientific Biomate-3S. Overnight

cultures were grown in LB and normalized to the same starting OD$_{600}$. 1:100 dilutions

were made into 96-well plates filled with 100ul of LB medium plus indicated antibiotics

and/or specified sugars and tricarboxylic acids. Two-fold dilutions of each antibiotic was

made.

Microscopy.

For all imaging, cells were grown at 37°C in LB medium. Imaging was done on 1% LB-

agarose pads at room temperature. Phase contrast images were collected on a Nikon

Ni-E epifluorescent microscope equipped with a 100X/1.45 NA objective (Nikon), Zyla

4.2 plus cooled sCMOS camera (Andor), and NIS Elements software (Nikon).

Metabolomics.

Collection. Cells were grown overnight in LB medium and then subcultured 1:1000 in

fresh LB for ~4 hours until in exponential phase. Cells were passed through a filter and

quenched with cold acetonitrile:methanol:water (40:40:20) + 0.5% formic acid. The

solution was neutralized with 1.9M ammonium bicarbonate before being centrifuged for

5 minutes. Supernatants were collected for mass spectrometry

Analysis. Supernatants were analyzed via the methods of Su et al. (65). Specificity was

achieved through a combination of chromatographic separations followed by high-

resolution MS. This method allows for the identification of ~300 water soluble

metabolites. Covariant ion analysis (COVINA) was used to identify peaks.

Acknowledgements

Research reported in this publication was supported by NIGMS of the National Institutes

of Health under award number R15GM129636-01A1. 100% of this project was financed

with Federal money. The content is solely the responsibility of the authors and does not

necessarily represent the official view of the NIH.

The authors would like to thank Dr. Elizabeth Ohneck and Dr. Tyrrell Conway for editing

of the manuscript.

1.   W. H. O. (WHO) (Antimicrobial Resistance: Global Report on Surveillance. ed WHO

(Geneva, Switzerland).

2.   S. A. McEwen, P. J. Collignon, Antimicrobial Resistance: a One Health Perspective.

Microbiology Spectrum **6** (2018).

3.   A. Pragasam, B. Veeraraghavan, E. Nalini, S. Anandan, K. Kaye, An Update on

Antimicrobial Resistance and the Role of Newer Antimicrobial Agents for *Pseudomonas*

aeruginosa. *Indian Journal of Medical Microbiology* **36**, 303-316 (2018).

4.   S. Lakhundi, K. Zhang, Methicillin-Resistant *Staphylococcus aureus*: Molecular

Characterization, Evolution, and Epidemiology. *Clinical Microbiology Reviews* **31**,

e00020-00018 (2018).

5.   S. S. Castle, "Penicillins" in xPharm: The Comprehensive Pharmacology Reference*,* S. J.

Enna, D. B. Bylund, Eds. (Elsevier, New York, 2007), pp. 1-3.

6.   M. Lobanovska, G. Pilla, Penicillin's Discovery and Antibiotic Resistance: Lessons for the

Future? *The Yale journal of biology and medicine* **90**, 135-145 (2017).

7.   F. van den Ent, L. A. Amos, J. Lowe, Prokaryotic Origin of the Actin Cytoskeleton. *Nature*

413, 39-44 (2001).

8.   S. A. Alyahya *et al.*, RodZ, a Component of the Bacterial Core Morphogenic Apparatus.

Proceedings of the National Academy of Sciences **106**, 1239-1244 (2009).

436 9. F. van den Ent, T. Izoré, T. B. Bharat, C. M. Johnson, J. Lowe, Bacterial Actin MreB Forms

437  Antiparallel Double Filaments. *eLife* **3** (2014).

438 10. E. Awuni, Y. Mu, Effect of A22 on the Conformation of Bacterial Actin MreB.

439  *International journal of molecular sciences* **20**, 1304 (2019).

440 11. Y. Awuni, S. Jiang, R. C. Robinson, Y. Mu, Exploring the A22-Bacterial Actin MreB

441  Interaction through Molecular Dynamics Simulations. *The Journal of Physical Chemistry*

442  *B* **120**, 9867-9874 (2016).

443 12. Z. Gitai, N. A. Dye, A. Reisenauer, M. Wachi, L. Shapiro, MreB Actin-Mediated

444  Segregation of a Specific Region of a Bacterial Chromosome. *Cell* **120**, 329-341 (2005).

445 13. G. J. Bean *et al.*, A22 Disrupts the Bacterial Actin Cytoskeleton by Directly Binding and

446  Inducing a Low-Affinity State in MreB. *Biochemistry* **48**, 4852-4857 (2009).

447 14. N. Ouzounov *et al.*, MreB Orientation Correlates with Cell Diameter in *Escherichia coli*.

448  *Biophysical Journal* **111**, 1035-1043 (2016).

449 15. N. A. Dye, Z. Pincus, I. C. Fisher, L. Shapiro, J. A. Theriot, Mutations in the Nucleotide

450  Binding Pocket of MreB can Alter Cell Curvature and Polar Morphology in *Caulobacter*.

451  *Molecular Microbiology* **81**, 368-394 (2011).

452 16. R. M. Morgenstein *et al.*, RodZ links MreB to Cell Wall Synthesis to Mediate MreB

453  Rotation and Robust Morphogenesis. *Proceedings of the National Academy of Sciences*,

454  12510-12515 (2015).

455 17. R. B. Helling *et al.*, Toxic Waste Disposal in *Escherichia coli*. *Journal of bacteriology* **184**,

456  3699-3703 (2002).

457   18.   Y. Shan *et al.*, ATP-Dependent Persister Formation in *Escherichia coli*. *mBio* **8**, e02267-

458   02216 (2017).

459   19.   B. P. Conlon *et al.*, Persister Formation in *Staphylococcus aureus* is Associated with ATP

460   Depletion. *Nature Microbiology* **1**, 16051 (2016).

461   20.   S. E. Rowe *et al.*, Reactive Oxygen Species Induce Antibiotic Tolerance During Systemic

462   *Staphylococcus aureus* Infection. *Nature Microbiology* **5**, 282-290 (2020).

463   21.   I. Irnov *et al.*, Crosstalk Between the Tricarboxylic Acid Cycle and Peptidoglycan

464   Synthesis in *Caulobacter crescentus* through the Homeostatic Control of α-

465   ketoglutarate. *PLoS genetics* **13**, e1006978-e1006978 (2017).

466   22.   F. O. Bendezú, P. A. J. de Boer, Conditional Lethality, Division Defects, Membrane

467   Involution, and Endocytosis in *mre* and *mrd* Shape Mutants of *Escherichia coli*. *Journal of*

468   *Bacteriology* **190**, 1792-1811 (2008).

469   23.   T. Kruse, J. Bork-Jensen, K. Gerdes, The Morphogenetic MreBCD Proteins of *Escherichia*

470   *coli* form an Essential Membrane-Bound Complex. *Molecular Microbiology* **55**, 78-89

471   (2005).

472   24.   D. Vinella, D. Joseleau-Petit, D. Thévenet, P. Bouloc, R. D'Ari, Penicillin-binding Protein 2

473   Inactivation in *Escherichia coli* Results in Cell Division Inhibition, Which is Relieved by

474   FtsZ Overexpression. *Journal of bacteriology* **175**, 6704-6710 (1993).

475   25.   N. Iwai, K. Nagai, M. Wachi, Novel S-Benzylisothiourea Compound That Induces

476   Spherical Cells in Escherichia coli Probably by Acting on a Rod-shape-determining

477   Protein(s) Other Than Penicillin-binding Protein 2. *Bioscience, Biotechnology, and*

478   *Biochemistry* **66**, 2658-2662 (2002).

26.   T. Baba *et al.*, Construction of *Escherichia coli* K-12 in-frame, Single-gene Knockout

Mutants: the Keio Collection. *Molecular Systems Biology* **2**, 2006.0008-2006.0008

(2006).

27.   D. C. Sévin, T. Fuhrer, N. Zamboni, U. Sauer, Nontargeted *in vitro* Metabolomics for

High-throughput Identification of Novel Enzymes in *Escherichia coli*. *Nature Methods* **14**,

187-194 (2017).

28.   E. Flores-Bautista, C. L. Cronick, A. R. Fersaca, M. A. Martinez-Nuñez, E. Perez-Rueda,

Functional Prediction of Hypothetical Transcription Factors of *Escherichia coli* K-12

Based on Expression Data. *Computational and Structural Biotechnology Journal* **16**, 157-

166 (2018).

29.   C.-P. Tseng, Regulation of Fumarase (f*umB*) Gene Expression in *Escherichia coli* in

Response to Oxygen, Iron and Heme Availability: Role of the *arcA*, *fur*, and *hemA* Gene

Products. *FEMS Microbiology Letters* **157**, 67-72 (1997).

30.   J.-A. M. Benjamin, E. Massé, The iron-sensing aconitase B binds its own mRNA to

prevent sRNA-induced mRNA cleavage. *Nucleic acids research* **42**, 10023-10036 (2014).

31.   Y. Tang, J. R. Guest, Direct Evidence for mRNA Binding and Post-transcriptional

Regulation by *Escherichia coli* Aconitases. *Microbiology* **145**, 3069-3079 (1999).

32.   C. S. Westfall, P. A. Levin, Bacterial Cell Size: Multifactorial and Multifaceted. *Annual*

review of microbiology **71**, 499-517 (2017).

33.   A. M. Sperber, J. K. Herman, Metabolism Shapes the Cell. *Journal of Bacteriology* **199**,

e00039-00017 (2017).

34.   N. S. Hill, P. J. Buske, Y. Shi, P. A. Levin, A Moonlighting Enzyme Links Escherichia coli Cell

501   Size with Central Metabolism. *PLOS Genetics* **9**, e1003663 (2013).

502   35.   U. Sauer, B. J. Eikmanns, The PEP–pyruvate–oxaloacetate Node as the Switch Point for

503   Carbon Flux Distribution in Bacteria. *FEMS Microbiology Reviews* **29**, 765-794 (2005).

504   36.   H. Barreteau *et al.*, Cytoplasmic Steps of Peptidoglycan Biosynthesis. *FEMS Microbiology*

505   *Reviews* **32**, 168-207 (2008).

506   37.   D. Mengin-Lecreulx, B. Flouret, J. van Heijenoort, Pool Levels of UDP N-

507   acetylglucosamine and UDP N-acetylglucosamine-enolpyruvate in *Escherichia coli* and

508   Correlation with Peptidoglycan Synthesis. *Journal of Bacteriology* **154**, 1284-1290

509   (1983).

510   38.   D. Rogers, S.-H. Yu, Substrate Specificity of a Glucose Permease of *Escherichia coli*.

511   *Journal of Bacteriology* **84**, 877-881 (1962).

512   39.   M. Hernandez-Asensio, J. M. Ramirez, F. F. Del Campo, The Control by Respiration of the

513   Uptake of α-methyl-glucoside in *Escherichia coli* K12. *Archives of Microbiology* **103**, 155-

514   162 (1975).

515   40.   H. Cho *et al.*, Bacterial Cell Wall Biogenesis is Mediated by SEDS and PBP Polymerase

516   Families Functioning Semi-Autonomously. *Nature Microbiology* **1**, 16172 (2016).

517   41.   S. Leclercq *et al.*, Interplay Between Penicillin-binding Proteins and SEDS Proteins

518   Promotes Bacterial Cell Wall Synthesis. *Scientific Reports* **7**, 43306 (2017).

519   42.   A. J. Meeske *et al.*, SEDS Proteins are a Widespread Family of Bacterial Cell Wall

520   Polymerases. *Nature* **537**, 634-638 (2016).

43.   B. G. Spratt, The Mechanism of Action of Mecillinam. *Journal of Antimicrobial*

Chemotherapy **3**, 13-19 (1977).

44.   D. A. Preston, C. Y. Wu, L. C. Blaszczak, D. E. Seitz, N. G. Halligan, Biological

Characterization of a New Radioactive Labeling Reagent for Bacterial Penicillin-binding

Proteins. *Antimicrobial agents and chemotherapy* **34**, 718-721 (1990).

45.   N. A. Curtis, D. Orr, G. W. Ross, M. G. Boulton, Affinities of Penicillins and

Cephalosporins for the Penicillin-binding Proteins of *Escherichia coli* K-12 and their

Antibacterial Activity. *Antimicrobial agents and chemotherapy* **16**, 533-539 (1979).

46.   T. S. Ursell *et al.*, Rod-Like Bacterial Shape is Maintained by Feedback Between Cell

Curvature and Cytoskeletal Localization. *Proceedings of the National Academy of*

Sciences **111**, E1025-E1034 (2014).

47.   B. P. Bratton, J. W. Shaevitz, Z. Gitai, R. M. Morgenstein, MreB Polymers and Curvature

Localization are Enhanced by RodZ and Predict *E. coli's* Cylindrical Uniformity. *Nature*

Communications **9**, 2797 (2018).

48.   S. Wang, Ned S. Wingreen, Cell Shape Can Mediate the Spatial Organization of the

Bacterial Cytoskeleton. *Biophysical Journal* **104**, 541-552 (2013).

49.   L. Furchtgott, N. S. Wingreen, K. C. Huang, Mechanisms for Maintaining Cell Shape in

Rod-Shaped Gram-negatve Bacteria. *Molecular Microbiology* **81**, 340-353 (2011).

50.   C. L. White, A. Kitich, J. W. Gober, Positioning Cell Wall Synthetic Complexes by the

Bacterial Morphogenetic Proteins MreB and MreD. *Molecular Microbiology* **76**, 616-633

(2010).

51.   A. V. Divakaruni, C. Baida, C. L. White, J. W. Gober, The Cell Shape Proteins MreB and

MreC Control Cell Morphogenesis by Positioning Cell Wall Synthetic Complexes.

Molecular Microbiology **66**, 174-188 (2007).

52.   A. V. Divakaruni, R. R. O. Loo, Y. Xie, J. A. Loo, J. W. Gober, The Cell-shape Protein MreC

Interacts with Extracytoplasmic Proteins Including Cell Wall Assembly Complexes in

Caulobacter crescentus. *Proceedings of the National Academy of Sciences of the United*

States of America **102**, 18602-18607 (2005).

53.   R. M. Figge, A. V. Divakaruni, J. W. Gober, MreB, the Cell Shape-Determining Bacterial

Actin Homologue, Co-Ordinates Cell Wall Morphogenesis in *Caulobacter crescentus*.

Molecular Microbiology **51**, 1321-1332 (2004).

54.   Y. Kawai, R. A. Daniel, J. Errington, Regulation of Cell Wall Morphogenesis in *Bacillus*

subtilis by Recruitment of PBP1 to the MreB Helix. *Molecular Microbiology* **71**, 1131-

1144 (2009).

55.   N. Honoré, S. T. Cole, Streptomycin Resistance in Mycobacteria. *Antimicrobial Agents*

and Chemotherapy **38**, 238-242 (1994).

56.   R. Cantón *et al.*, Prevalence and Spread of Extended-spectrum β-lactamase-producing

Enterobacteriaceae in Europe. *Clinical Microbiology and Infection* **14**, 144-153 (2008).

57.   M. A. Webber, L. J. V. Piddock, The Importance of Efflux Pumps in Bacterial Antibiotic

Resistance. *Journal of Antimicrobial Chemotherapy* **51**, 9-11 (2003).

58.   E. Tuomanen, R. Cozens, W. Tosch, O. Zak, A. Tomasz, The Rate of Killing of *Escherichia*

coli by β-Lactam Antibiotics Is Strictly Proportional to the Rate of Bacterial Growth Free.

Journal of general microbiology **132** (1986).

59.   M. H. Pontes, E. A. Groisman, Slow Growth Determines Nonheritable Antibiotic

Resistance in *Salmonella enterica*. *Science Signaling* **12**, eaax3938 (2019).

60.   A. Brauner, O. Fridman, O. Gefen, N. Q. Balaban, Distinguishing Between Resistance,

Tolerance and Persistence to Antibiotic Treatment. *Nature Reviews Microbiology* **14**,

320-330 (2016).

61.   K. R. Allison, M. P. Brynildsen, J. J. Collins, Metabolite-enabled Rradication of Bacterial

Persisters by Aminoglycosides. *Nature* **473**, 216-220 (2011).

62.   R. L. Miranda *et al.*, Glycolytic and Gluconeogenic Growth of *Escherichia coli* O157:H7

(EDL933) and *E. coli* K-12 (MG1655) in the Mouse Intestine. *Infection and Immunity* **72**,

1666-1676 (2004).

63.   Y. Bertin *et al.*, The Gluconeogenesis Pathway is Involved in Maintenance of

Enterohaemorrhagic *Escherichia coli* O157:H7 in Bovine Intestinal Content. *PloS one* **9**,

e98367-e98367 (2014).

64.   J. W. Njoroge, Y. Nguyen, M. M. Curtis, C. G. Moreira, V. Sperandio, Virulence Meets

Metabolism: Cra and KdpE Gene Regulation in Enterohemorrhagic *Escherichia coli*. *mBio*

3, e00280-00212 (2012).

65.   X. Su *et al.*, In-Source CID Ramping and Covariant Ion Analysis of Hydrophilic Interaction

Chromatography Metabolomics. *Anal Chem* **92**, 4829-4837 (2020).

Figure 1. Mutations in the genes of the TCA cycle increase the MIC of A22. A)

Screening of a subset of the Keio deletion library for mutants with increased MIC_{A22}

yielded 86 mutations in different gene ontology groups. B) Model of the TCA cycle.

Indicated proteins have higher MIC_{A22} when deleted. C) MIC_{A22} of mutants from the

initial screen. Only *mdh* mutants show an increased MIC_{A22}. D) MIC_{A22} of mutants that

break the other steps in the TCA cycle. *acnB* and *sucC* deletions also show a higher

MIC$_{A22}$. MIC fold change is represented as mutant/wild-type cells. A fold change of 1

(dotted line) indicates no change. * p<.05, *** p>.001 comparing each strain to 1.

Figure 2. Metabolomics analysis reveals upregulation of gluconeogenesis in *mdh* cells.

Cells were grown in LB media without the addition of sugar. The only way to produce

pyruvate and glucose is through gluconeogenesis. A) Ion counts of malate levels in WT

and *mdh* cells. B) Ion counts of pyruvate levels in WT and *mdh* cells. C) Ion counts of

glucose-6P levels in WT and *mdh* cells. D) Ion counts of UD-NAG levels in WT and *mdh*

cells. Average of 3 independent experiments with standard deviation shown.

610

Figure 3. The addition of glucose to the media phenocopies the MIC phenotypes of the

mdh deletion. WT cells were grown in LB medium supplemented with the indicated

amount of either glucose or a non-hydrolizable glucose mimic, α-methyl D-glucoside (α

md). Cells were treated with either A22 (black bars) or mecillinam (blue bars). Error

shown in standard deviation from 3 independent experiments. * p<.05, ***p<.001

comparing glucose to α md for each condition.

Figure 4. *mdh* mutation provides increased MIC specifically of A22 and mecillinam. MIC

assays were performed for the indicated antibiotics. A fold change of 1 (dotted line)

indicates no change. *** p>.001 comparing each strain to 1.

Category	Count
Metabolism	24
Genes of Unknown Function	14
Protein Activity	13
DNA Maintenance	12
Envelope	12
Transport	7
Gene Expression	4
Total	86

629 Table S1 Genes with high A_{22} MICs are enriched in metabolism. Gene ontology (GO) analysis of

630 the 86 strains displaying increased A_{22} MICs revealed a high concentration of genes focused in

631 metabolism.

Metabolism	Fold Change	Standard Deviation
fucO	12	4.62
gatZ	12	4.618802154
mdh	12	4.618802154
rsxA	12	4.618802154
bcsA	10.125	7.44283772
lipB	7	2
bioB	6.5	7.141428429
caiD	3.0125	3.44972825
hemE	2.75	3.5
purM	2.75	3.570714214
uxuB	2.3875	3.761731649
proB	2.375	3.772156766
purK	2.2625	3.830877837
hyfC	2.25	1.258305739
eutJ	2	0
garK	2	1.632993162
acs	1.7625	1.690845449
icd	1.5	0.577350269
rbsR	1.5	0.577350269
yodB	1.5	0.577350269
allB	1.3875	1.784364966
moeB	1.375	1.796988221

	Fold Change	Standard Deviation
prpE	1.375	0.75
yjcS	1.375	0.75
Gene of Unknown Function	**Fold Change**	**Standard Deviation**
yedS	12	4.618802154
yeeN	10	7.659416862
yeeP	10	4
yjgZ	10	7.659416862
yrhA	4.375	4.190763654
ybaV	4.25	4.330127019
yobF	2.875	3.473110997
yfjR	2	0
yifN	1.75	1.658312395
glvG	1.625	1.600781059
yhcC	1.625	0.75
yhfK	1.625	0.75
yoeF	1.625	0.75
yjgX	1.375	1.75
Protein Activity	**Fold Change**	**Standard Deviation**
rpsT	10	4
rsuA	8	0
prfC	6.125	7.531876703
yihP	3.5	3

ykgM	3.5	3
tfaE	3.125	1.75
allR	2.625	3.591076904
uspG	2.5	3.674234614
ytfE	2.5	3.674234614
mtfA	2.375	3.75
fabR	2	1.632993162
napD	2	0
ibpB	1.5	1.683250823
DNA Maintenance	**Fold Change**	**Standard Deviation**
dnaT	12	4.618802154
hsdS	4	0
yjjB	3.5	1
caiT	2.5	3.674234614
dinF	2.5	3.674234614
rnhB	2.3875	3.761731649
aroP	2.375	3.75
priC	2.375	3.772156766
srmB	1.5	0.577350269
ttk (slmA)	1.5	0.577350269
torI	1.375	0.75
yrdD	1.375	0.75
Envelope	**Fold Change**	**Standard Deviation**

gmhB	10	4
clpP	9	5.033222957
wcaC	8.125	6.329494451
mltG	6.625	7.04006392
wcaK	6	4
bamB	3.125	3.326033674
ldtB	2.75	3.570714214
dtpA	2.25	3.840572874
oppC	1.75	1.658312395
mpl	1.625	1.600781059
ybbY	1.625	0.75
yeaY	1.5	1
Transport	**Fold Change**	**Standard Deviation**
wzzE	12	4.618802154
cls (clsA)	4.625	7.586995453
kdpF	2.75	3.5
kefG	2.375	3.772156766
mdtB	1.625	0.75
yhdY	1.625	0.75
ssuC	1.375	0.75
Gene Expression	**Fold Change**	**Standard Deviation**
narQ	12	4.618802154
yafT	8	0

yaeH	8	0
creC	1.5	1

Table S2. List of genes with screened for increased MIC_{A22}. Genes are sorted by ontology groups

and then listed in order of decreasing MIC_{A22}.

Figure 1S. *mdh* mutant grows the same as WT. WT and *mdh* cells were grown

overnight in LB. Cells were equalized to the same O.D.$_{600}$ and subcultured into fresh LB

medium in a 96 well plate at a 1:100 dilution. The plate was grown at 37°C with shaking

in a Biotek Synergy plate reader. Error bars are standard deviation across wells in a

plate. Data is a representative experiment.

Figure 2S. *mdh* cells become round during A22 treatment. Cells were grown overnight

and subcultured 1:1000 into fresh LB medium. Cells were grown for 4 hours in the 0 and

1 µg/ml A22. Cells were grown in LB for 2 hours before the addition of 10 µg/ml A22 for

2 hours. Images are representative of an experiment done in triplicate. Scale bar = 5 µ

m.

Figure 3S. *sucC* does not cause a change in mecillinam resistance. MIC_{mec} of *acnB* and

sucC mutants compared to WT type cells.

655