Balanced JAK/STAT signaling is critical to maintain the functional

and structural integrity of the *Drosophila* respiratory epithelium

Xiao Niu[1], Christine Fink[1], Kimberley Kallsen[2,4], Viktoria Mincheva[1], Sören Franzenburg[3],

Ruben Prange[1], Judith Bossen[1], Holger Heine[2, 5], Thomas Roeder[1, 5]

[1]Kiel University, Zoology, Dept. of Molecular Physiology, Kiel, Germany

[2]Research Center Borstel, Dept. of Innate Immunity, Borstel, Germany

[3]Kiel University, IKMB, Kiel, Germany

[4]Current address: Boehringer Ingelheim, Ingelheim, Germany

[5]German Center for Lung Research, Airway Research Center North, Kiel, Germany

Summary

Signals mediated by the Janus kinase (JAK)/signal transducer and activator of transcription

(STAT) pathway play a central role in maintaining homeostasis in a multitude of tissues. A

large number of studies have shown that this role is particularly prominent in the lungs.

Deregulation of the JAK/STAT signaling pathway is causally linked with various, mostly

chronic, lung diseases, including lung cancer, asthma, and chronic obstructive pulmonary

disease. To elucidate the molecular framework that explains how deregulated JAK/STAT

signaling gives rise to pathogenic states, we used the fruit fly *Drosophila* as a model. While

the JAK/STAT pathway is characterized by high structural diversity and complexity in

vertebrates, it is relatively simple in *Drosophila*. The JAK/STAT pathway was active in

almost all respiratory epithelial cells of larvae and adult flies. Stressful stimuli, such as

cigarette smoke, evoked strong and regionalized activation of the JAK/STAT pathway,

which was most likely driven by the concurrently induced ligand Unpaired 2. Inhibition of

JAK/STAT signaling induced apoptotic processes in epithelial cells. The aforementioned

chronic lung diseases are associated with increased activity of the JAK/STAT pathway and

are treated with specific JAK inhibitors. Therefore, we investigated the effects of increased

JAK/STAT signaling in the respiratory epithelium of *Drosophila*. Ectopic activation of the

JAK/STAT pathway led to premature death at the larval or pupal stage. Furthermore, it

induced major structural changes in epithelial cells, which almost completely lost their

typical characteristics. These structural changes led to considerable thickening of the

epithelium, substantial narrowing of the air-conducting space, and disruption of the

tracheal epicuticular structure. Pharmacological interference of JAK/STAT signaling

reverted this phenotype. Activation of the JAK/STAT pathway also affected vesicle-

mediated transport, which led to erroneous trafficking of typical junction proteins. In

summary, these results demonstrate that balanced JAK/STAT signaling is essential for the

normal functionality of the respiratory epithelium and thus the entire organ. A basal level

of JAK/STAT signaling is required for cellular processes such as growth and division.

However, chronic overactivation of this signaling leads to massive structural changes that

are closely related to pathologies typically seen in chronic inflammatory lung diseases.

Introduction

The Janus kinase (JAK)/signal transducer and activator of transcription (STAT) signaling

system is of central importance for several critical physiological processes including

development, tissue homeostasis, and immune responses [1]. Various cytokines, growth

factors, and related signaling compounds signal via this evolutionarily conserved system

[2]. The JAK/STAT signaling pathway effectively transduces external signals into desired

cellular responses despite having relatively few essential components. Signaling via this

pathway is therefore straightforward and allows environmental factors to directly

influence transcriptional activity in cells, thereby linking important biological processes

with environmental cues [3]. Moreover, JAK/STAT signaling is tightly associated with a

great variety of immune responses. JAK/STAT signaling acts downstream of a plethora of

cytokines that transmit immune-related information and is, therefore, a central hub in

various immunocompetent cells [4-7]. Deregulation of this signaling pathway is directly

linked with numerous human diseases, including cancer and inflammatory diseases [5, 8,

9].

Lung diseases are often associated with deregulated JAK/STAT signaling. Such

deregulation can arise due to mutations and polymorphisms in genes associated with

JAK/STAT signaling. In addition, this signaling is activated by stressors, infections, and

injuries. This might cause sustained chronic inflammation of the airways and/or alveoli,

which is closely associated with the onset and chronification of various lung diseases.

Chronic obstructive pulmonary disease (COPD), asthma, idiopathic pulmonary fibrosis,

and lung cancer are causally associated with deregulated JAK/STAT signaling [10]. This

signaling is not only of great importance during organ development, but also plays a

central role in maintaining tissue and immune homeostasis, especially in response to

stressors, infection, and damage [11, 12]. Functional JAK/STAT signaling is required to

cope with stressful stimuli such as hyperoxia in the airway epithelium [10]. Deregulation

of this signaling in the airways is associated with pathological states. Specifically,

chronically reduced JAK/STAT signaling is associated with impaired repair capacities [11],

while increased JAK/STAT signaling leads to cell proliferation that can cause cancer [13].

Despite its simple general organization, the JAK/STAT signaling pathway is characterized

by multiple redundancies in vertebrates. A multitude of elements function at each level

of the JAK/STAT signaling pathway; for example, more than 50 cytokines can activate this

pathway [6]. Moreover, deregulation of JAK/STAT signaling in different motile and

resident cell types found in the lungs is associated with chronic lung diseases. Therefore,

models with a much simpler JAK/STAT pathway and a less complicated cellular

composition in the airways should help to elucidate the effects of dysfunctional JAK/STAT

signaling, especially in airway epithelial cells.

Simple models such as the fruit fly *Drosophila melanogaster* are thus a reasonable

alternative to analyze the relevance of JAK/STAT signaling in the airways. While this

signaling pathway exhibits broad diversification at all levels in mammals, it involves a

single receptor (Domeless), one JAK (Hopscotch), and one STAT transcription factor

(Stat92E) in *Drosophila* [2, 14]. This low level of redundancy offers a unique advantage to

investigate the JAK/STAT pathway in tissues of interest. JAK/STAT signaling is highly

important for many processes, including development, in flies. The tracheal system does

not form when JAK/STAT signaling is abolished in tracheal primordia [15, 16]. Unpaired

(Upd), a major ligand of this pathway, is expressed in tracheal pits and initiates JAK/STAT

signaling in this organ primordium [17]. A lack of JAK/STAT signaling during very early

phases of tracheal development leads to a lack of expression of developmentally relevant

genes such as trachealess (*trh*), which is essential to instruct important phases of tracheal

development including cell movement and elongation, as well as invagination processes

that lead to tube formation [15, 18]. This is consistent with the important role of JAK/STAT

signaling during lung development in vertebrates [19, 20].

The current study aimed to determine the physiological significance of JAK/STAT signaling

in a simple, but fully functional, airway system. Therefore, we studied the larval airway

system of the fruit fly *D. melanogaster* and evaluated if the JAK/STAT signaling pathway is

active in functional airway epithelial cells and if stress signals modify the activity of this

pathway. Furthermore, we investigated if deregulation of this pathway leads to disease-

associated phenotypes in this simple airway system.

Results

The JAK/STAT pathway is strongly activated in the trachea during embryogenesis and is

important for organ development [21]. However, the exact pattern of its activation is

unclear. In this study, we used the *Stat92E-GFP* reporter [21] to monitor activation of the

JAK/STAT pathway concurrently with *btl>LacZ.nls* [22], which specifically labels the trachea.

Figure 1A schematically depicts the JAK/STAT pathway in *Drosophila*. Activation of the

Stat92E was enhanced in a central region in the trachea, mainly in the transverse

connective area (Fig. 1B–D). To confirm the importance of the JAK/STAT pathway in the

development of the *Drosophila* respiratory system, we manipulated its activity in the

trachea by driving expression of a dominant-negative isoform of Domeless (*Dome.DN*)

using *btl-Gal4*. Although some animals grew to the larval stage, all died before reaching

the pupal stage (Fig. 1E). Microscopic analysis of surviving larvae showed that the dorsal

trunk (DT) was absent (Fig. 1E''). Time-lapse imaging of trachea-specific *Dome.DN*

expression in embryos revealed that all tracheal segments were fused at stage 15;

however, the DT subsequently separated (Fig. 1F and H). Next, we tested the effects of

ectopic activation of the JAK/STAT pathway induced by expression of the ligand (Upd3) or

the constitutively active JAK-allele (Hop.CA) on tracheal development. These animals died

prematurely at the embryo or larval stage (Fig. 1E). Microscopic analysis of larvae and

embryos showed that the tracheal segment separated at the larval stage (Fig. 1E'''–E'''');

however, the disconnection of segments was due to failure of their fusion at the beginning

of tracheal development (Fig. 1G and H). In contrast with animals that expressed *upd3*,

most *Hop.CA*-expressing animals exhibited a full tracheal tree. However, they all died at

the first larval stage, which indicates that JAK/STAT signaling influences tracheal

development after the first tracheal structures have formed.

To determine if the JAK/STAT pathway also operates in functional airway epithelium,

namely, the larval tracheal system, we used the STAT reporter line mentioned above. STAT

signaling was active in all regions of the tracheal system in third instar larvae (Fig. 2A).

JAK/STAT activity was reduced in the posterior zone of the trachea, tenth tracheal

metamere (Tr10), where cells progressively undergo apoptosis in response to trachea

metamorphosis (Fig. 2B). However, JAK/STAT activity was much stronger in regions such

as Tr2, spiracular branch (SB) and dorsal branch (DB), where cells re-enter the cell cycle at

this developmental stage than in nearby areas containing quiescent cells (Fig. 2C and D).

We next evaluated if JAK/STAT activity is observed in airway epithelial cells of adults. The

same approach detected substantial reporter activity in airway epithelial cells of fully

developed adults (Fig. 2E and E'). Inhibition of JAK/STAT signaling changed the fate and

morphology of larval airway epithelial cells (Fig. 2F). This inhibition was achieved by

ectopically expressing *Dome.DN*. These cells gradually lost their typical shape, and

apoptotic processes were initiated 2 days after induction of ectopic expression. This

induction of apoptosis was confirmed by detection of Dcp1 in cells with typical apoptotic

features (Fig. 2G and H). Apoptosis was induced to an even greater extent (and faster) in

stem cells of the larval tracheal system, where Dcp1-positive cells were observed on 1 day

after induction of ectopic *Dome.DN* expression (Fig. 2I–K). The JAK/STAT pathway was

activated throughout the entire tracheal system. Therefore, we evaluated expression of

the three Upd ligands in this system. Experiments using the corresponding enhancer-*Gal4*

lines demonstrated that *upd2* was highly expressed in the tracheal system (Fig. S1).

To elucidate the effects of JAK/STAT deregulation in the tracheal system initiated at

different time points during larval development, we employed the temperature-inducible

Gal4/Gal80[ts] system (Fig. 3A). We first tested the effects of inhibition of the tracheal

JAK/STAT pathway induced by ectopic expression of *Dome.DN* on survival and larval

structure (Fig. 3B). Ectopic expression was initiated at three time points during larval

development (Fig. 3B). Larval mortality was higher, the earlier expression was initiated.

No larvae survived to the pupal stage (Fig. 3C). On the other hand, activation of JAK/STAT

signaling induced by ectopic expression of *upd3* or *Hop.CA* led to lower levels of larval

death. In both cases, no pupae developed into adults. Furthermore, we analyzed the

effects of this intervention on tracheal structure. Whereas *Dome.DN* overexpression

elicited only minor effects on tracheal structure, *upd3* or *Hop.CA* overexpression caused

substantial epithelial thickening (Fig. 3D–H). Quantitative evaluation revealed that ectopic

overexpression of *Hop.CA* greatly increased epithelial thickness (Fig. 3H). Experiments

involving mosaic expression of *Hop.CA* driven by *vvl-FLP, CoinFLP-Gal4, UAS-EGFP (vvl-coin)*

demonstrated that this effect was cell-autonomous. Cells that expressed *Hop.CA* were

much thicker than neighboring cells that did not (Fig. 3I). Treatment with various JAK

inhibitors [23] reversed epithelial thickening induced by ectopic overexpression of *Hop.CA*

in the trachea (Fig. 3J–L). Treatment with Baricitinib, Oclacitinib, or Filgotinib reduced

epithelial thickening by more than 60% (Fig. 3L).

To elucidate the mechanisms underlying this structural response, we analyzed the

phenotype in more detail. The length of a defined area of the tracheal system (between

Tr8 and Tr9), the length-thickness product, and the cell number were quantified (Fig. 4A

and B). Length was unaffected by manipulation of JAK/STAT signaling in the trachea, with

the exception that it was reduced after 4 days of ectopic *Dome.DN* expression (Fig. 4B).

Consequently, the length-thickness product mirrored the findings made when assessing

thickness. However, the number of cells in the Tr8-Tr9 region of the trachea with

manipulated JAK/STAT signaling was similar to that in matching controls (Fig. 4B''). Hence,

the thickening was due to an increase in cell volume, not an increase in cell number. To

determine if this thickening is an artifact caused by the isolation method, we directly fixed

the trachea in situ or isolated it in cell culture medium with the same osmolarity as

hemolymph. The same thickening was observed in both cases (Supplemental Fig. 2). To

quantify the time course of this thickening, we subjected the trachea to different

treatment regimens prior to analysis. At least 2 days of ectopic expression were necessary

to induce this phenotype (Fig. 4C–E).

After demonstrating that JAK/STAT signaling operates in a fully functional tracheal system

and that deregulation of its activation substantially impacts the structure of the airway

system, we aimed to identify conditions under which JAK/STAT signaling activity is

increased. To this end, we subjected flies to two stressful conditions that are related to

the airway system, namely, cigarette smoke exposure and hypoxia. Whereas hypoxia did

not affect the activities of the JAK/STAT pathway reporter or the corresponding ligands

(Upd2 (Fig. 5D–F) and Upd3 (Fig. 5J–L)), smoke exposure induced expression of both *upd2*

and *upd3* as well as activation of the JAK/STAT pathway reporter (Fig. 5A–C, F–I, and L).

Induced expression was mostly confined to a highly sensitive region close to the spiracles

(Tr9 and Tr10) (Fig. 5B, F, H, and L).

To investigate the molecular mechanisms underlying the effects of chronic JAK/STAT

activation on the airways, we performed RNA-sequencing analysis of tracheal cells

expressing *Hop.CA* in third instar larvae. The validity of this experimental procedure was

supported by the induced expression of *hopscotch* (including *Hop.CA*) and *Socs36E* (one

of the best-characterized Stat92E target genes) [21], which were upregulated 35-fold and

2.7-fold, respectively ($p < 10^{-12}$). Principal component analysis of biological replicates

separated control and overexpression samples into different groups (Fig. 6A, middle).

Finally, the expression of 2004 genes was statistically significantly regulated, with 1128

downregulated genes and 876 upregulated genes ($p < 0.05$). A circular heatmap of 707

differentially expressed genes ($p < 0.01$ and fold change > 2) is presented (Fig. 6A).

Moreover, we analyzed changes in expression upon ectopic silencing of the JAK/STAT

pathway using *Dome.DN*. A Venn diagram revealed that there was a statistically significant

overlap in the cohorts of genes regulated by both interventions (Fisher's exact test, *p* <

0.0001; Fig. 6B).

To further elucidate the molecular mechanisms functioning in the thickened epithelium,

we performed promotor scan studies and Gene Ontology (GO) analyses. Genes with the

highest rates of induction supported by the lowest p-values were subjected to promoter

scanning (Pscan) analysis. Putative genes directly regulated by the JAK/STAT pathway were

identified (Tables 1 and 2). Among these highly regulated genes, the largest group

contained genes that encoded products involved in innate immunity, including seven

antimicrobial peptide genes (*IM1, IM2, IM4, IM14, CG5791, CG5778,* and *Drs*). All these

antimicrobial peptides, except for Drs, contain one or two CXXC regions. These CXXC-

containing peptides belong to the family of Bomanins, whose expression is highly induced

following bacterial or fungal infection, and high levels of the corresponding mature

peptides are found in the hemolymph of infected flies [24].

Next, we analyzed all significantly regulated genes based on KEGG and GO annotations.

Six terms in KEGG pathway analysis and nine categories in GO analysis were enriched

among these genes according to their predicted functions (Table S1 and Fig. 7).

One striking feature of the epithelium with chronically activated JAK/STAT signaling was

induced expression of genes involved in vesicle-mediated transport (Fig. 7). GO analysis

demonstrated that this term was enriched among 32 regulated genes that had been

assigned to vesicle-mediated transport. All these genes were upregulated, except for one

(CG5946). However, according to a Pscan analysis, these genes did not seem to be direct

JAK/STAT targets since a Stat92E promoter binding was not enriched (table S2 and S3). The

transcript levels of genes related to epicuticle development were reduced. Fifty-five genes

relevant to cuticle development were regulated, of which 40 were downregulated. In

addition, other biological processes such as muscle cell differentiation, cellular protein-

containing complex assembly, RNA processing, translation, and detection of chemical

stimuli were significantly enriched. In the latter three categories, the number of

upregulated genes was similar to the number of downregulated genes. A schematic

representation of the different biological processes superimposed on a scheme of an

epithelial cell is shown (Fig. 7B). One interesting result was that endocytosis and vesicle-

mediated transport were impaired. To analyze this further, we focused on two peripheral

membrane-binding proteins, namely, Coracle (Cora), a component of septate junctions

and Armadillo (Arm), a component of adherens junctions. Both proteins depend on vesicle

transport to reach their destinations [25, 26] (Fig. 8A). Immunofluorescence analysis of

Hop.CA-overexpressing animals revealed that Arm and Cora accumulated in the cytoplasm,

which is indicative of dysfunctional endosomal transport (Fig. 8B–E). We also observed a

stenosis phenotype with tube narrowing. This arose when epithelial cells expressed

Hop.CA at very high levels, as indicated by strong green fluorescence (Fig. 8F and G and

Fig. S2). The lumen narrowing observed here is reminiscent of the phenotype caused by

mutations of the coatomer protein complex (COP)I or COPII complexes, which are

required for the efficient secretion of proteins that drive tube expansion [27, 28] (Fig. 8H).

Cuticle development was identified as another regulated biological process based on

transcriptome data. This process is dependent on secretion in the larval trachea, and

chitin-rich structures, called taenidia, are believed to make tubes flexible as well as

sufficiently strong in order to avoid collapse [29]. The chitin structure in the trachea was

unsorted in *Hop.CA*-expressing animals compared with control tubes (Fig. 9A and B).

Chitin staining revealed that the highly organized structure of the chitinous intima was

almost completely lost in *Hop.CA*-expressing animals (Fig. 9C). Mosaic analysis

demonstrated that the trachea was structurally disorganized where *Hop.CA* was expressed

(Fig. 9D–F).

The effects of activating the JAK/STAT pathway on the epithelium were time-dependent.

To determine if a weaker expression of *Hop.CA* induces similar phenotypes, we used

another trachea-specific Gal4-driver, namely, PPK4 (also called *nach-Gal4*). In the *nach-*

Gal4 line, *Gal4* is expressed from the late embryo stage, is strongly expressed in the early

L1 and late L2 stages, at which point expression is stronger than that driven by *btl-Gal4*,

but is weakly expressed at other larval stages [30, 31]. Microscopy analysis of the trachea

showed that weak expression of *Hop.CA* in the trachea promoted epithelial thickening (Fig.

10A and B). However, epithelial thickening after 4 days of *Hop.CA* induction was slightly

lower in *nach>Hop.CA* than in *btl>Hop.CA* (Fig. 10D). On the other hand, weakly driven

upd3 expression in the trachea resulted in identical changes as those observed using *btl-*

Gal4 (Fig. 10C and E). Finally, we used the same driver to inhibit JAK/STAT signaling in the

trachea by expressing *Dome.DN*. Animals died prematurely at the larval stage and their

tracheas were filled with liquid, demonstrating that functionality was impaired (Fig. 10F).

Discussion

The present study aimed to elucidate the significance of the JAK/STAT signaling pathway

in the respiratory epithelium. We used the fruit fly *D. melanogaster* owing to the

experimental advantages of this system and the unique simplicity of the JAK/STAT

signaling pathway. JAK/STAT signaling exhibited tonic activity in almost all respiratory

epithelial cells and its activity was substantially enhanced by exposure to cigarette smoke.

Stress-induced induction of the JAK/STAT signaling pathway and the corresponding

cytokines is also observed in primary human respiratory epithelial cells, suggesting that

the activation and mechanism-of-action of JAK/STAT signaling are similar in human and

Drosophila respiratory epithelial cells [32]. Our finding that the JAK/STAT signaling

pathway operated in airway epithelial cells of embryos, larvae, and adults implies that it

plays a central role in these cells at all developmental stages. The JAK/STAT pathway was

previously reported to be important for embryonic development of the tracheal system

[33]. We clarified the sequence of developmental steps in which this signaling pathway

plays a central role. The JAK/STAT signaling pathway is particularly important for

development and maintenance of the DT. Tracheal JAK/STAT signaling is required during

development of the adult tracheal system at the pupal stage [34]. This is consistent with

our finding that the various interventions induced lethality in the pupal stage at the latest.

It is a matter of debate whether JAK/STAT signaling is essential for mammalian lung

development. However, the prevailing view that JAK/STAT signaling is not essential for

embryonic lung development [11] has been challenged by recent studies [20]. Although

JAK/STAT signaling is essential for different aspects of tracheal development, the major

focus of the current study was on the role of this signaling in maintaining homeostasis in

the fully functional airway epithelium. Therefore, we chose *Drosophila* larvae as a model

because they possess fully functional airways that must perform gas exchange for survival

and growth of the animal. Respiratory epithelial cells in larvae are terminally

differentiated and cannot divide, but increase in size as larvae grow. We demonstrated

that the JAK/STAT signaling pathway was essential in the larval airway epithelium and that

its blockade induced apoptosis. This implies that the JAK/STAT pathway plays a central role

in homeostatic processes in this organ, which also appears to be the case in mammalian

airway epithelia. Perturbation of JAK/STAT signaling induces apoptosis in cell lines derived

from airway epithelia [35], and this seems to be particularly relevant for growing tissues

and cells that can still divide or grow [36]. This ability to grow might be one reason for the

high susceptibility of *Drosophila* airway epithelial cells to blockade of JAK/STAT signaling.

Thus, cytokines that activate the JAK/STAT signaling pathway act as survival factors at least

in part, especially where repair mechanisms are operative. Functional JAK/STAT signaling

is also required for regenerative processes after epithelial damage [11, 12], the response

to infections [37], and the reaction to hyperoxia [38]. This indicates that JAK/STAT signaling

in mammalian airway epithelial cells and the *Drosophila* trachea is particularly relevant

for a protective reaction to stressful stimuli. Consistently, we demonstrated that JAK/STAT

signaling was normally active in airway epithelial cells of *Drosophila* larvae. A basal level

of JAK/STAT signaling was observed in these cells and this signaling was further activated

in response to very strong stressors such as chronic exposure to smoke particles. There

appears to be an organ-autonomous JAK/STAT signaling system, with expression and

release of the cytokines Upd2 and Upd3, and activation of a JAK/STAT-dependent response

induced in the same regions of the tracheal system. A comparable, stress-induced system

has been identified in the intestinal epithelium of flies, where highly stressful insults

targeting absorptive enterocytes induce production and release of the cytokine Upd3 [39,

40]. In contrast with the intestines, where cytokines produced by stressed enterocytes

induce proliferation of stem cells to replenish the enterocyte pool, the typical structure of

epithelial cells in the tracheal system is altered without activation of stem cells.

Although a threshold level of JAK/STAT signaling is required for the functionality and

survival of airway epithelial cells, excessive activation of this signaling is associated with

several lung diseases including lung cancer, acute lung injury, asthma, pulmonary fibrosis,

and COPD [13, 41-45]. This implies that different facets of this highly complex signaling

pathway are critically involved in the development of the majority of chronic lung diseases.

Thus, maintaining homeostasis in the airway epithelium, especially with respect to its

potential damage and repair, is essential for a healthy life. In this context, the JAK/STAT

pathway seems to play a decisive role [11]. To study the imbalance of JAK/STAT signaling

that is causally associated with the majority of chronic lung diseases and is responsible for

the structural changes typically seen in these diseases, we ectopically activated JAK/STAT

signaling specifically in the airway epithelium of *Drosophila* larvae. This activation induced

major structural changes in the airway epithelium, mainly to the architecture of tracheal

cells. Mosaic analysis demonstrated that this effect was cell-autonomous. These major

structural changes of the trachea are reminiscent of those observed in asthma, COPD,

acute lung injury, and lung cancer. Structural changes that permit to fulfill the original

function of epithelial cells are often observed during the epithelial-to-mesenchymal

transition. JAK/STAT signaling is essential for the epithelial-to-mesenchymal transition

intrinsic to lung cancer [46].

Ectopic JAK/STAT pathway activation induced complex alterations to the transcriptome in

airway epithelial cells. These changes affected immune-relevant genes such as those

encoding antimicrobial peptides, mainly from the Bomanin family. JAK/STAT signaling

typically regulates antimicrobial responses in the human airway epithelium [41]. However,

chronically activated JAK/STAT signaling perturbed secretory processes and the formation

of extracellular structures. The extracellular matrix defects caused by *Hop.CA*

overexpression are reminiscent of the pathophysiology of the aforementioned chronic

lung diseases that involve responses to acute or chronic injury [47]. Therefore, attempts

to enhance the repair capacities of epithelial cells by inducing structural changes via

excessive JAK/STAT signaling would also perturb normal cellular functions, such as the

transport of junction proteins to the membrane. This would lead to reduced barrier

function of the epithelium, which is a hallmark of chronic lung diseases such as asthma

and COPD [48-50]. We also observed epicuticular changes, which considerably influenced

the structure of the whole organ. However, it is difficult to identify an equivalent change

in vertebrates.

The JAK/STAT pathway is an excellent target for therapeutic intervention in many lung

diseases, including asthma, COPD, acute lung injury, idiopathic pulmonary fibrosis, and

lung cancer [43, 51-55]. We demonstrated that pharmacological interference of the

JAK/STAT signaling pathway reverted the structural phenotype observed upon ectopic

activation of this signaling and consequently flies survived. This shows that the *Drosophila*

model is not only suitable to study the structural effects of excessive JAK/STAT signaling

and the underlying molecular mechanisms, but also to screen compounds and thereby

identify novel therapeutic strategies.

It must be remembered that the vertebrate lung and insect trachea are not homologous.

Nevertheless, they share a high degree of similarity in terms of their development,

physiology, innate immunity, and operative signaling systems [56, 57]. Therefore, the fruit

fly is a very valuable tool to study genes associated with a great variety of chronic lung

diseases including asthma, COPD, and lung cancer [58-63]. This simple model can be used

as part of an experimental toolbox to elucidate the role of JAK/STAT signaling in the

airways and the effects of chronic deregulation of this signaling. In addition, it provides a

readily accessible experimental system that is amenable to pharmacologic interventions,

and allows hypotheses and intervention strategies to be easily tested.

Material and Methods

***Drosophila* strains and husbandry**

Stat92E-GFP was used to monitor the activation of the pathway [21]; the Gal4-UAS system

[64] was used to target ectopic expression to the tracheal system. Gal4 drivers used were:

btl-Gal4, UAS-GFP on the 3rd chromosome, and *btl-Gal4, UAS-GFP* on the 2nd

chromosome (obtained from the Leptin group, Heidelberg, Germany); *upd2-Gal4*; *upd3-*

Gal4 [60]; *nach-Gal4* [30]. The UAS responders included: *UAS-LacZ.nls* (BDSC 3956), *UAS-*

domeΔcyt2.1 (*UAS-Dome.DN*) and *UAS-Hop.CA* (*UAS-hopTumL*) were obtained from N.

Perrimon [15, 65]. The *UAS-upd3* was constructed in our lab. *TubP-Gal80[ts]* (BDSC 7018)

were obtained from the Bloomington stock center. Unless otherwise stated, the flies were

raised on standard medium at 25 °C with 50–60% relative humidity under a 12:12 h

light/dark cycle as described earlier [66].

Coin-FLP expression system

Vvl-FLP/CyO; btl-moe.mRFP (BDSC 64233), *tubP-Gal80[ts]* and *CoinFLP-Gal4, UAS-2xEGFP*

(BDSC 58751) were used to construct animals for the tracheal mosaic analysis. Ventral

veins lacking (*vvl*) was expressed in larval tracheal clones that covered approximately 30

to 80% of the trachea [67, 68]. The genotype of the flies was *vvl-FLP, CoinFLP-Gal4, UAS-*

2xEGFP/CyO (*vvl-coin*) and *vvl-FLP, CoinFLP-Gal4, UAS-2xEGFP/CyO; tub-Gal80[ts]* (*vvl-*

coin.ts).

Developmental Viability

For developmental viability of eggs, the eggs were collected overnight and were not

physically handled in any way. The number of the hatched eggs and the pupae were

counted starting from two days after the collection. Each group has 4 replicates that

included more than 20 eggs. For developmental viability of larvae, *tub-Gal80[ts]* was used

to limit UAS responder expression at the larvae stage. Animals were raised at 18 °C to keep

the UAS responder gene silent. Larvae at different instar stages were transferred to new

medium at 29 °C. In this study, usually 4 replicates were performed with 30 larvae. The

stage of larvae was determined via the appearance of anterior spiracles.

Determination of epithelial thickness

The trachea of L2 and L3 Larvae were carefully dissected out from the posterior side of

the body in PBS. The isolated trachea was immersed in 50% glycerol and digital images

captured within 15 min. L2 Larvae were distinguished from L3 larvae by the appearance

of anterior spiracles. The relative ages of L3 larvae were inferred from the size of the

animal. 30 larvae were used in each group and specimens were analysed by Image J.

Drug application and determination of epithelial thickness

JAK inhibitors (Baricitinib #16707 , Oclacitinib #18722, Filgotinib #17669 - Cayman

Chemicals, Michigan, USA) were diluted in DMSO [100 mM]. For later application the

inhibitors were diluted 1:10 in 100% EtOH. We used 20 µl of each diluted inhibitor for 2

ml of concentrated medium (5% yeast extract, 5% corn flour, 5% sucrose, 1% low-melt

agarose, 1 ml of 10% propionic acid and 3 ml of 10% Nipagin). The eggs of each crossing

have been applicated on the modified medium and kept at 20 °C until the larvae reach

the L2 stage. Afterwards they were incubated at 30 °C for 2 days to induce the *btl-Gal4,*

UAS-GFP; tub-Gal80[ts] (btl.ts) –driver. The trachea of L3 Larvae were carefully dissected

out from the posterior side of the body in the PBS. Isolated trachea was immersed in 50%

glycerol and digital images captured in 15 min. 10 larvae were used for each group.

Time-lapse microscopy

All images were acquired using a ZEISS Axio Image Z1 fluorescent microscope. Embryos

were dechlorinated in 3% sodium hypochlorite and immersed in Halocarbon oil 700

(Sigma Aldrich, 9002-83-9). Then the embryos were imaged after stage 15 when the

tracheal tree formed at 3 hours intervals.

Cigarette smoke and hypoxic exposure

All cigarette smoke exposure experiments were carried out in a smoking chamber,

attached to a diaphragm pump. Common research 3R4F cigarettes (CTRP, Kentucky

University, Lexington, USA) were used for all experiments. The vials containing animals

were capped with a monitoring grid to allow the cigarette smoke to diffuse into the vial.

For long-time smoke experiments, L2 larvae were exposed to smoke three times a day for

30 min each, on two consecutive days. For heavy smoke experiments, L3 larvae were

exposed to 2 cigarettes smoke one time for 45 min, which led to about 35-50% mortality.

To study the effects of hypoxia on the activity of JAK/STAT signalling pathway, larvae were

exposed to long-term hypoxia and short-term hypoxia separately. For long-term hypoxia

experiments, L2 larvae were exposed to 5% oxygen three times a day for 30 min each, on

two consecutive days. For short-term hypoxia experiments, L3 larvae were exposed to 1%

oxygen once for 5 hours.

Antibody and tracheal stains

Larvae were dissected by ventral filleting and fixed in 4% paraformaldehyde for 30 min.

Embryos were staged according to Campos-Ortega and Hartenstein [69] and fixed in 4%

formaldehyde for 30 min. Immunostaining followed standard protocols as described

earlier [70, 71]. GFP signals were amplified by immunostaining with polyclonal rabbit anti-

GFP (used at 1:500, Sigma-Aldrich, Merck KGaA, Darmstadt, Germany, SAB4301138). 40-

1a (used at 1:50, DSHB, Iowa City, USA) was used to detect Beta-galactosidase. Coracle

protein was detected with a monoclonal mouse anti-coracle antibody (used at 1:200,

DSHB, Iowa City, USA, C566.9). Armadillo protein was detected with a monoclonal mouse

anti-armadillo antibody (DSHB, US, N2 7A1, used at 1:500). A monoclonal rabbit Cleaved

Drosophila Dcp1 (used at 1:200, Cell Signaling, Frankfurt/M, Germany, #9578) was used

to detect apoptotic cells. Secondary antibodies used were: Cy3-conjugated goat-anti-

mouse, Cy3-conjugated goat-anti-rabbit, Alexa488-conjugated goat-anti-mouse (Jackson

Immunoresearch, Dianova, Hamburg, Germany), Alexa488-conjugated goat-anti-rabbit

(used at 1:500, Cell signaling, Frankfurt/M, Germany, #9578). Tracheal chitin was stained

with 505 star conjugated chitin-binding probe (NEB, Germany, used at 1:300). Nuclei were

stained with 4',6-Diamidino-2-Phenylindole, Dihydrochloride (DAPI) (Roth, Karlsruhe,

Germany, 6843). Specimens were analyzed and digital images captured either with a

confocal (CLSM Leica TCS SP1, Leica, Wetzlar, Germany) or a conventional fluorescence

microscope (ZEISS Axio Imager Z1, Zeiss, Oberkochen, Germany), respectively.

RNA isolation and RNA sequencing

For the gene expression analysis of 3rd instar larvae trachea, animals were dissected in

cold PBS and isolated trachea transferred to RNA Magic (BioBudget, Krefeld, Germany)

and processed essentially as described earlier [60] with slight modifications. The tissue

was homogenized in a Bead Ruptor 24 (BioLab products, Bebensee, Germany) and the

RNA was extracted by using the PureLink RNA Mini Kit (Thermo Fisher, Waltham, MA, USA)

for phase separation with the RNA Magic reagent. An additional DNase treatment was

performed following the on-column PureLink DNase treatment protocol (Thermo Fisher,

Waltham, MA, USA).

Sequencing libraries were constructed using the TruSeq stranded mRNA kit (Illumina, San

Diego, USA) and 50 bp single-read sequencing was performed on an Illumina HiSeq 4000

with 16 samples per lane. Resulting sequencing reads were trimmed for low-quality bases

and   adapters   using   the   fastq   Illumina   filter

(http://cancan.cshl.edu/labmembers/gordon/fastq_illumina_filter/)   and   cutadapt

(version 1.8.1) [72]. Transcriptomics analysis including gene expression values and

differential expression analysis was done using CLC Genomics Workbench. The detailed

protocols can be obtained from the CLC Web site (http://www. clcbio.com/products/clc-

genomics-workbench). *Drosophila melanogaster* reference genome (Release 6) [73] was

used for mapping in this research.

The transcription factor binding site enrichment and the Gene Ontology enrichment

analyses of the differentially expressed genes were carried out using Pscan and Panther,

respectively. Date was visualized through the circos software.

References

481
1.   Rawlings JS, Rosler KM, Harrison DA: **The JAK/STAT signaling pathway**. *J Cell Sci* 2004, **117**(Pt
8):1281-1283.
2.   Zeidler MP, Bausek N: **The Drosophila JAK-STAT pathway**. *JAKSTAT* 2013, **2**(3):e25353.
3.   Yan R, Small S, Desplan C, Dearolf CR, Darnell JE, Jr.: **Identification of a Stat gene that functions in**
Drosophila development. *Cell* 1996, **84**(3):421-430.

4.   Shuai K, Liu B: **Regulation of JAK-STAT signalling in the immune system**. *Nat Rev Immunol* 2003,
3(11):900-911.
5.   Villarino AV, Kanno Y, Ferdinand JR, O'Shea JJ: **Mechanisms of Jak/STAT signaling in immunity**
and disease. *J Immunol* 2015, **194**(1):21-27.
6.   Villarino AV, Kanno Y, O'Shea JJ: **Mechanisms and consequences of Jak-STAT signaling in the**
immune system. *Nat Immunol* 2017, **18**(4):374-384.
7.   Agaisse H, Perrimon N: **The roles of JAK/STAT signaling in Drosophila immune responses**.
Immunol Rev 2004, **198**:72-82.
8.   O'Shea JJ, Schwartz DM, Villarino AV, Gadina M, McInnes IB, Laurence A: **The JAK-STAT pathway:**
impact on human disease and therapeutic intervention. *Annu Rev Med* 2015, **66**:311-328.
9.   Constantinescu SN, Girardot M, Pecquet C: **Mining for JAK-STAT mutations in cancer**. *Trends*
Biochem Sci 2008, **33**(3):122-131.
10.   Pernis AB, Rothman PB: **JAK-STAT signaling in asthma**. *J Clin Invest* 2002, **109**(10):1279-1283.
11.   Tadokoro T, Wang Y, Barak LS, Bai Y, Randell SH, Hogan BL: **IL-6/STAT3 promotes regeneration of**
airway ciliated cells from basal stem cells. *Proc Natl Acad Sci U S A* 2014, **111**(35):E3641-3649.
12.   Kida H, Mucenski ML, Thitoff AR, Le Cras TD, Park KS, Ikegami M, Muller W, Whitsett JA: **GP130-**
STAT3 regulates epithelial cell migration and is required for repair of the bronchiolar
epithelium. *Am J Pathol* 2008, **172**(6):1542-1554.
13.   Dutta P, Sabri N, Li J, Li WX: **Role of STAT3 in lung cancer**. *JAKSTAT* 2014, **3**(4):e999503.
14.   Arbouzova NI, Zeidler MP: **JAK/STAT signalling in Drosophila: insights into conserved regulatory**
and cellular functions. *Development* 2006, **133**(14):2605-2616.
15.   Brown S, Hu N, Hombria JC: **Identification of the first invertebrate interleukin JAK/STAT receptor,**
the Drosophila gene domeless. *Curr Biol* 2001, **11**(21):1700-1705.
16.   Perrimon N, Mahowald AP: **l(1)hopscotch, A larval-pupal zygotic lethal with a specific maternal**
effect on segmentation in Drosophila. *Dev Biol* 1986, **118**(1):28-41.
17.   Harrison DA, McCoon PE, Binari R, Gilman M, Perrimon N: **Drosophila unpaired encodes a**
secreted protein that activates the JAK signaling pathway. *Genes Dev* 1998, **12**(20):3252-3263.
18.   Isaac DD, Andrew DJ: **Tubulogenesis in Drosophila: a requirement for the trachealess gene**
product. *Genes Dev* 1996, **10**(1):103-117.
19.   Nogueira-Silva C, Santos M, Baptista MJ, Moura RS, Correia-Pinto J: **IL-6 is constitutively**
expressed during lung morphogenesis and enhances fetal lung explant branching. *Pediatr Res*
2006, **60**(5):530-536.
20.   Piairo P, Moura RS, Baptista MJ, Correia-Pinto J, Nogueira-Silva C: **STATs in Lung Development:**
Distinct Early and Late Expression, Growth Modulation and Signaling Dysregulation in
Congenital Diaphragmatic Hernia. *Cell Physiol Biochem* 2018, **45**(1):1-14.
21.   Bach EA, Ekas LA, Ayala-Camargo A, Flaherty MS, Lee H, Perrimon N, Baeg GH: **GFP reporters**
detect the activation of the Drosophila JAK/STAT pathway in vivo. *Gene Expr Patterns* 2007,
7(3):323-331.
22.   Shiga Y, M. TM, Hayashi S: **A nuclear GFP/ β‐galactosidase fusion protein as a marker for**
morphogenesis in living Drosophila. *Development, Growth & Differentiation* 1996, **38**:99-106.

23.   Roskoski R, Jr.: **Janus kinase (JAK) inhibitors in the treatment of inflammatory and neoplastic**
diseases. *Pharmacol Res* 2016, **111**:784-803.

24.   Lindsay SA, Lin SJH, Wasserman SA: **Short-Form Bomanins Mediate Humoral Immunity in**
Drosophila. *J Innate Immun* 2018, **10**(4):306-314.

25.   Lock JG, Stow JL: **Rab11 in recycling endosomes regulates the sorting and basolateral transport**
of E-cadherin. *Mol Biol Cell* 2005, **16**(4):1744-1755.

26.   Oshima K, Fehon RG: **Analysis of protein dynamics within the septate junction reveals a highly**
stable core protein complex that does not include the basolateral polarity protein Discs large. *J*
Cell Sci 2011, **124**(Pt 16):2861-2871.

27.   Jayaram SA, Senti KA, Tiklova K, Tsarouhas V, Hemphala J, Samakovlis C: **COPI vesicle transport is**
a common requirement for tube expansion in Drosophila. *PLoS One* 2008, **3**(4):e1964.

28.   Tsarouhas V, Senti KA, Jayaram SA, Tiklova K, Hemphala J, Adler J, Samakovlis C: **Sequential pulses**
of apical epithelial secretion and endocytosis drive airway maturation in Drosophila. *Dev Cell*
2007, **13**(2):214-225.

29.   Glasheen BM, Robbins RM, Piette C, Beitel GJ, Page-McCaw A: **A matrix metalloproteinase**
mediates airway remodeling in Drosophila. *Dev Biol* 2010, **344**(2):772-783.

30.   Liu L, Johnson WA, Welsh MJ: **Drosophila DEG/ENaC pickpocket genes are expressed in the**
tracheal system, where they may be involved in liquid clearance. *Proc Natl Acad Sci U S A* 2003,
100(4):2128-2133.

31.   Wagner C, Isermann K, Roeder T: **Infection induces a survival program and local remodeling in**
the airway epithelium of the fly. *Faseb J* 2009, **23**(7):2045-2054.

32.   Kode A, Yang SR, Rahman I: **Differential effects of cigarette smoke on oxidative stress and**
proinflammatory cytokine release in primary human airway epithelial cells and in a variety of
transformed alveolar epithelial cells. *Respir Res* 2006, **7**:132.

33.   Hombria JC, Sotillos S: **JAK-STAT pathway in Drosophila morphogenesis: From organ selector to**
cell behavior regulator. *JAKSTAT* 2013, **2**(3):e26089.

34.   Powers N, Srivastava A: **JAK/STAT signaling is involved in air sac primordium development of**
Drosophila melanogaster. *FEBS Lett* 2019, **593**(7):658-669.

35.   Zheng XJ, Yang ZX, Dong YJ, Zhang GY, Sun MF, An XK, Pan LH, Zhang SL: **Downregulation of leptin**
inhibits growth and induces apoptosis of lung cancer cells via the Notch and JAK/STAT3
signaling pathways. *Biol Open* 2016, **5**(6):794-800.

36.   Quinton LJ, Mizgerd JP: **NF-kappaB and STAT3 signaling hubs for lung innate immunity**. *Cell*
Tissue Res 2011, **343**(1):153-165.

37.   Matsuzaki Y, Xu Y, Ikegami M, Besnard V, Park KS, Hull WM, Wert SE, Whitsett JA: **Stat3 is required**
for cytoprotection of the respiratory epithelium during adenoviral infection. *J Immunol* 2006,
177(1):527-537.

38.   Hokuto I, Ikegami M, Yoshida M, Takeda K, Akira S, Perl AK, Hull WM, Wert SE, Whitsett JA: **Stat-3**
is required for pulmonary homeostasis during hyperoxia. *J Clin Invest* 2004, **113**(1):28-37.

39.   Jiang H, Patel PH, Kohlmaier A, Grenley MO, McEwen DG, Edgar BA: **Cytokine/Jak/Stat signaling**
mediates regeneration and homeostasis in the Drosophila midgut. *Cell* 2009, **137**(7):1343-1355.

40.   Miguel-Aliaga I, Jasper H, Lemaitre B: **Anatomy and Physiology of the Digestive Tract of**
568   **Drosophila melanogaster**. *Genetics* 2018, **210**(2):357-396.
569   41.   Simeone-Penney MC, Severgnini M, Tu P, Homer RJ, Mariani TJ, Cohn L, Simon AR: **Airway**
570   **epithelial STAT3 is required for allergic inflammation in a murine model of asthma**. *J Immunol*
571   2007, **178**(10):6191-6199.
572   42.   Zhang X, Yue P, Page BD, Li T, Zhao W, Namanja AT, Paladino D, Zhao J, Chen Y, Gunning PT *et al*:
573   **Orally bioavailable small-molecule inhibitor of transcription factor Stat3 regresses human**
574   **breast and lung cancer xenografts**. *Proc Natl Acad Sci U S A* 2012, **109**(24):9623-9628.
575   43.   Yew-Booth L, Birrell MA, Lau MS, Baker K, Jones V, Kilty I, Belvisi MG: **JAK-STAT pathway**
576   **activation in COPD**. *Eur Respir J* 2015, **46**(3):843-845.
577   44.   Prele CM, Yao E, O'Donoghue RJ, Mutsaers SE, Knight DA: **STAT3: a central mediator of**
578   **pulmonary fibrosis?** *Proc Am Thorac Soc* 2012, **9**(3):177-182.
579   45.   Gao H, Guo RF, Speyer CL, Reuben J, Neff TA, Hoesel LM, Riedemann NC, McClintock SD, Sarma JV,
580   Van Rooijen N *et al*: **Stat3 activation in acute lung injury**. *J Immunol* 2004, **172**(12):7703-7712.
581   46.   Liu RY, Zeng Y, Lei Z, Wang L, Yang H, Liu Z, Zhao J, Zhang HT: **JAK/STAT3 signaling is required for**
582   **TGF-beta-induced epithelial-mesenchymal transition in lung cancer cells**. *Int J Oncol* 2014,
583   **44**(5):1643-1651.
584   47.   Fahy JV, Dickey BF: **Airway mucus function and dysfunction**. *N Engl J Med* 2010, **363**(23):2233-
585   2247.
586   48.   Georas SN, Rezaee F: **Epithelial barrier function: at the front line of asthma immunology and**
587   **allergic airway inflammation**. *J Allergy Clin Immunol* 2014, **134**(3):509-520.
588   49.   Gon Y, Hashimoto S: **Role of airway epithelial barrier dysfunction in pathogenesis of asthma**.
589   *Allergol Int* 2018, **67**(1):12-17.
590   50.   Heijink IH, Brandenburg SM, Postma DS, van Oosterhout AJ: **Cigarette smoke impairs airway**
591   **epithelial barrier function and cell-cell contact recovery**. *Eur Respir J* 2012, **39**(2):419-428.
592   51.   Milara J, Ballester B, Morell A, Ortiz JL, Escriva J, Fernandez E, Perez-Vizcaino F, Cogolludo A,
593   Pastor E, Artigues E *et al*: **JAK2 mediates lung fibrosis, pulmonary vascular remodelling and**
594   **hypertension in idiopathic pulmonary fibrosis: an experimental study**. *Thorax* 2018, **73**(6):519-
595   529.
596   52.   Athari SS: **Targeting cell signaling in allergic asthma**. *Signal Transduct Target Ther* 2019, **4**:45.
597   53.   Loh CY, Arya A, Naema AF, Wong WF, Sethi G, Looi CY: **Signal Transducer and Activator of**
598   **Transcription (STATs) Proteins in Cancer and Inflammation: Functions and Therapeutic**
599   **Implication**. *Front Oncol* 2019, **9**:48.
600   54.   Severgnini M, Takahashi S, Rozo LM, Homer RJ, Kuhn C, Jhung JW, Perides G, Steer M, Hassoun
601   PM, Fanburg BL *et al*: **Activation of the STAT pathway in acute lung injury**. *Am J Physiol Lung Cell*
602   *Mol Physiol* 2004, **286**(6):L1282-1292.
603   55.   Song L, Rawal B, Nemeth JA, Haura EB: **JAK1 activates STAT3 activity in non-small-cell lung**
604   **cancer cells and IL-6 neutralizing antibodies can suppress JAK1-STAT3 signaling**. *Mol Cancer Ther*
605   2011, **10**(3):481-494.
606   56.   Andrew DJ, Ewald AJ: **Morphogenesis of epithelial tubes: Insights into tube formation,**
607   **elongation, and elaboration**. *Dev Biol* 2010, **341**(1):34-55.

608   57.   Bergman P, Seyedoleslami Esfahani S, Engstrom Y: **Drosophila as a Model for Human Diseases-**
609   **Focus on Innate Immunity in Barrier Epithelia**. *Curr Top Dev Biol* 2017, **121**:29-81.
610   58.   Bossen J, Uliczka K, Steen L, Pfefferkorn R, Mai MM, Burkhardt L, Spohn M, Bruchhaus I, Fink C,
611   Heine H *et al*: **An EGFR-Induced Drosophila Lung Tumor Model Identifies Alternative**
612   **Combination Treatments**. *Mol Cancer Ther* 2019, **18**(9):1659-1668.
613   59.   Levine BD, Cagan RL: **Drosophila Lung Cancer Models Identify Trametinib plus Statin as**
614   **Candidate Therapeutic**. *Cell Rep* 2016, **14**(6):1477-1487.
615   60.   Prange R, Thiedmann M, Bhandari A, Mishra N, Sinha A, Hasler R, Rosenstiel P, Uliczka K, Wagner
616   C, Yildirim AO *et al*: **A Drosophila model of cigarette smoke induced COPD identifies Nrf2**
617   **signaling as an expedient target for intervention**. *Aging (Albany NY)* 2018, **10**(8):2122-2135.
618   61.   Kallsen K, Zehethofer N, Abdelsadik A, Lindner B, Kabesch M, Heine H, Roeder T: **ORMDL**
619   **deregulation increases stress responses and modulates repair pathways in Drosophila airways**.
620   *J Allergy Clin Immunol* 2015, **136**(4):1105-1108.
621   62.   Roeder T, Isermann K, Kabesch M: **Drosophila in asthma research**. *Am J Respir Crit Care Med*
622   2009, **179**(11):979-983.
623   63.   Roeder T, Isermann K, Kallsen K, Uliczka K, Wagner C: **A Drosophila asthma model - what the fly**
624   **tells us about inflammatory diseases of the lung**. *Adv Exp Med Biol* 2012, **710**:37-47.
625   64.   Brand AH, Perrimon N: **Targeted gene expression as a means of altering cell fates and generating**
626   **dominant phenotypes**. *Development* 1993, **118**(2):401-415.
627   65.   Harrison DA, Binari R, Nahreini TS, Gilman M, Perrimon N: **Activation of a Drosophila Janus**
628   **kinase (JAK) causes hematopoietic neoplasia and developmental defects**. *EMBO J* 1995,
629   **14**(12):2857-2865.
630   66.   Fink C, Hoffmann J, Knop M, Li Y, Isermann K, Roeder T: **Intestinal FoxO signaling is required to**
631   **survive oral infection in Drosophila**. *Mucosal Immunol* 2016, **9**(4):927-936.
632   67.   Bosch JA, Tran NH, Hariharan IK: **CoinFLP: a system for efficient mosaic screening and for**
633   **visualizing clonal boundaries in Drosophila**. *Development* 2015, **142**(3):597-606.
634   68.   Chen F, Krasnow MA: **Progenitor outgrowth from the niche in Drosophila trachea is guided by**
635   **FGF from decaying branches**. *Science* 2014, **343**(6167):186-189.
636   69.   Campos-Ortega JA, Hartenstein V: **The Embryonic Development of Drosophila melanogaster**:
637   Springer-Verlag Berlin Heidelberg; 1997.
638   70.   Jeon M, Nguyen H, Bahri S, Zinn K: **Redundancy and compensation in axon guidance: genetic**
639   **analysis of the Drosophila Ptp10D/Ptp4E receptor tyrosine phosphatase subfamily**. *Neural Dev*
640   2008, **3**:3.
641   71.   Levi BP, Ghabrial AS, Krasnow MA: **Drosophila talin and integrin genes are required for**
642   **maintenance of tracheal terminal branches and luminal organization**. *Development* 2006,
643   **133**(12):2383-2393.
644   72.   Martin M: **Cutadapt Removes Adapter Sequences From High-Throughput Sequencing Reads**.
645   *EMBnetjournal* 2011, **17**:10-12.
646   73.   Hoskins RA, Carlson JW, Wan KH, Park S, Mendez I, Galle SE, Booth BW, Pfeiffer BD, George RA,
647   Svirskas R *et al*: **The Release 6 reference sequence of the Drosophila melanogaster genome**.
648   *Genome Res* 2015, **25**(3):445-458.

650　Table 1 Genes that were most strongly upregulated in response to Hop.CA

651　overexpression.

Name	Maximum group mean	Fold change	FDR p-value
CG16772	11.85472435	135.4356918	0
hop	286.0640796	35.42957686	0
CG14570	42.20996514	35.3698493	0
CG14147	55.30940549	32.87679686	0
Hsp70Bb	161.511126	29.98530152	0
Ubx	200.4588187	23.51089416	0
pre-mod(mdg4)-I	1.124681971	18.66759173	2.22648E-06
IM14	11.45782606	15.13546529	3.80132E-07
IM4	26.42664387	14.97315552	1.74645E-12
CG14569	1576.880077	14.72575164	0
IM2	7.261319979	14.4714161	2.01792E-06
CG5791	1.786460103	13.80267454	2.90502E-05
CG9121	3.412245019	11.53962271	0
IM1	7.159415315	10.97306621	7.52122E-05
CG11413	35.69960698	10.63636792	3.44242E-07
CG16710	2.132413445	10.32026533	1.72352E-07
CG31960	267.2645814	10.04792875	0
CG16789	1.452007625	9.460811492	5.58576E-09
CG6034	6.915088882	8.468387575	4.7943E-11
Ace	2.056513226	8.187696842	0
CG13059	9.898481009	7.578675956	1.33893E-06
CG31809	8.642528741	7.388263435	0
CG7203	9.406607055	7.213889526	8.66989E-09
Vha14-2	2.510881225	7.050881817	2.31024E-07
wb	7.35456153	6.689915209	0
Drs	8166.473129	6.635790672	0
CG5778	3.202417196	6.607038471	0.000294835
CG3457	2.693960701	6.337465056	2.38755E-06
Lcp65Ad	9.932204556	5.792567795	2.58249E-08
lectin-22C	29.06148461	5.674130819	6.88538E-06
fuss	3.973734058	5.673577323	0

CG43333	2.390923691	5.632143859	1.48392E-09
CG18336	2.963052606	5.485479621	5.52167E-05
CG32563	3.062298877	5.352654098	0.000696584
FASN2	1.651227873	5.340581227	2.95545E-10
Cyp4d21	45.69613978	5.308508635	0
Obp57d	1.797549513	5.276761495	0.000102926
CG17560	2.313559171	5.218521032	0.000766701
CG5888	68.16302548	5.201444372	0

652 $p < 1 \times 10^{-4}$, fold change > 5, maximum group mean > 1. The list is restricted to 39 genes.

654 Table 2 Transcription factor-binding site motifs enriched in 41 highly upregulated genes.

Matrix ID	Matrix name	P-value
MA0023.1	dl (var.2)	0.000443261
MA0022.1	dl	0.00407113
MA0532.1	Stat92E	0.00794732
MA0450.1	hkb	0.00811969
MA0197.2	nub	0.0154201
MA0242.1	Bgb::run	0.0371696
MA0204.1	Six4	0.0443521
MA0444.1	CG34031	0.0488169

655 A total of 133 transcription factor profiles were used.

Figure 1: Effects of JAK/STAT signaling in the airway development in *Drosophila* embryos. (A) shows a
model of JAK/STAT signaling pathway in *Drosophila* comprising all major components. Three ligands
(Upd, Upd2 and Upd3) bind to a single receptor, Domeless (Dome), which transmits this information
via a single JAK (Hopscotch (Hop)) to a single STAT transcription factor (Stat92E). (B-D) Fluorescence
micrographs of *btl-Gal4*; *Stat92E-GFP*; *UAS-LacZ.nls* embryos stained for GFP (green, JAK/STAT pathway
activated zones), Beta-galactosidase (red, tracheal metamere), respectively. The JAK/STAT signaling
pathway is activated in the whole trachea metamere (Tr). The activation of JAK/STAT pathway in the

terminal regions is weaker than that in the central regions before (B) and after the fuse of tracheal
invaginations of all segments (C). (D) shows the pattern of activation of JAK/STAT signaling in the
trachea metamere of embryos. DB, dorsal branch; DTa, dorsal trunk anterior; DTp, dorsal trunk
posterior; VB, visceral branch; LTa, lateral trunk anterior; LTp, lateral trunk posterior; and GB, ganglionic
branch; SB, spiracular branch; transverse connective, TC. The stronger signal in the central regions
mainly belongs to TC. Scale bar: 20 µm. (E) Ectopic blocking or activation of JAK/STAT signaling in the
trachea was done by expressing of *Dome.DN* or *Hop.CA* (or *upd3*), respectively under the control of
btl-Gal4. The percentage of embryos that hatched and the percentage of larva that developed into
pupa is shown at the left side. Right (E'-E''''): Micrographs surviving larvae. *Btl* driving the expression
of *upd3* and *Dome.DN* inhibited the formation of an intact tracheal tree. *Btl* driving the expression of
Hop.CA in the trachea showed more intact trachea trees compared to expressing *upd3* and *Dome.DN*
in the trachea. X/XX: broken trachea/intact trachea; 50 animals were observed in each group. (F) Time-
lapse imaging the trachea of *btl>Dome.DN* embryo continuously for 9 hours after the trachea formed
(9hr ATF). 40 specimens were investigated. (G-H) Activation of the JAK/STAT pathway in the trachea led
to incompleted tracheal development during tracheal morphogenesis. Fluorescence micrographs of
btl>Hop.CA (or *>upd3*) embryos (G), and numbers of the trachea which had no fused Tr. (H)
Quantification of the numbers of defect tracheal branches in response to the different treatments.
Each group contains 3 replicates and each replicate contains 30 embryos in G and H. Arrows mark the
broken sites in the trachea. ns means not significant, * $p < 0.05$, ** $p < 0.01$ by Student's t-test. Scale
bar: 20 µm in B-D; 50 µm in E-G.

685
686

Figure 2: Inhibition of JAK/STAT signaling pathway induced cell apoptosis. (A-D) Fluorescence
micrographs of *btl-Gal4; Stat92E-GFP*; *UAS-LacZ.nls* larvae. The activation of JAK/STAT signaling was
detected by using the *Stat92E-GFP* reporter. In general, JAK/STAT signaling was induced in the whole
trachea of the larvae. The region in the posterior region of the trachea (orange dash line in B) that is
going to degenerate will lose JAK/STAT signaling. Conversely, some regions like Tr2, DB (C), and SB (D)

where the cell re-enter into cell cycle (white dash line) exhibited stronger JAK/STAT signaling compared

to the cell quiescent areas close to them. (E) Fluorescence micrographs of *btl-Gal4; Stat92E-GFP*; *UAS-*

LacZ.nls adult. JAK/STAT signaling was induced in the trachea of the adult. (F) Fluorescence micrographs

of the mutant clones in the trachea of *vvl-FLP*, *CoinFLP-Gal4, UAS-EGFP; tub-Gal80[ts]* (*vvl-*

coin.ts)>*UAS-Dome.DN* larvae. Negative regulation of JAK/STAT pathway by expressing *Dome.DN* in the

trachea led to the loss of cell shape and disintegration afterward (green arrows). (G-K) Fluorescence

micrographs of the trachea of *vvl-coin.ts* larvae (G), and the trachea of *vvl-coin.ts>UAS-Dome.DN* larvae

(H) stained for cleaved Dcp1 (red). Modified stem cells (J and K) could be stained by cleaved Dcp1

earlier than the mutant somatic cells. Inhibition of JAK/STAT signaling induced blockade of cell division

(I and K). 50 specimens were used in each assay.

Figure 3: Ectopic activation of epithelial JAK/STAT signaling induces cell thickening. (A) *btl-Gal4, UAS-*
GFP; tub-Gal80[ts] (*btl.ts*) larvae were reared at 18 ℃ (nonpermissive, A) and 29 ℃ (permissive, A´),
respectively. Using the Gal4/UAS system, comprising the temperature-sensitive repressor Gal80[ts],
were used to time ectopic gene expression. (B) Developmental viability of larvae with different
genotypes (including *Dome.DN*, *Hop.CA*, and *upd3* expression in the trachea driven by *btl.ts*) and
different start points of expression (indicated by black arrows). Statistical analysis of survival times is
included. In this experiment, 4 replicates were performed with 30 larvae in B. (C) In general, none of
these manipulations allowed survival up to adults. (D-G) Micrographs of Tr8-Tr9 of *btl.ts* L3 larvae (D,
Control); *btl.ts>Dome.DN* L3 larvae (E); *btl.ts>Hop.CA* L3 larvae (F) and *btl.ts>upd3* L3 larvae (G) that
were raised at 29 ℃ for 2 and 4 days, respectively. (H) Statistical analysis of the epithelial thicknesses
in Tr8-Tr9 of the corresponding larvae. The thickness of 30 trachea per group were measured in H. (I)
Micrographs of Tr8 of *vvl-FLP*, *CoinFLP-Gal4, UAS-EGFP* (*vvl-coin*)*>UAS-Hop.CA* larvae. The thickening
of the tracheal epithelium is observed in those cells that express *Hop.CA*. Changes in epithelial
thickness could be rescued by application of specific JAK inhibitors (J-L). (J-K) Microscopy of the tracheal
epithelium (L3 larvae). Control crossings *btl.ts>w[1118]* (J) compared to JAK/STAT activated crossings
btl.ts>UAS-Hop.CA (K). (L) Illustration and quantification of epithelial thickness (highlighted in red) of
the JAK inhibitors (Baricitinib, Oclacitinib, Filgotinib) compared to DMSO control. The thickness of 10
trachea per group were measured in J,K. ns means no significant, * $p < 0.05$, ** $p < 0.01$ by Student's
t-test. Scale bar: 500 µm in A; 20 µm in D-G; 50 µm J-K.

Figure 4: Changes in epithelial thickness could be attributed to the increase in cell volume and is time-dependent. (A-B) Illustration and quantification of the length, length X thickness, and the cell number of the Tr8-Tr9 region of larvae with different types of ectopic manipulation (including *Dome.DN*, *Hop.CA* and *upd3* expression in the trachea driven by *btl.ts*) for 2 or 4 days, respectively. 30 trachea were analysed in each group. (C-E) Illustration of the time point and period of activation of *Hop.CA* expression by using the *btl.ts* (C) and quantification of the thickness of the Tr8-Tr9 region of control larvae (*btl.ts* line) and those experiencing ectopic manipulation (*btl.ts>Hop.CA* line). Quantitative analyses of the thicknesses (D). 1-day expression of *Hop.CA* starting at L2 larvae (E), but that started at the early L3 stage promotes epithelial thickening (E′). Each group contains 50 replicates in D-E. ns means not significant, * $p < 0.05$, ** $p < 0.01$ by Student's t-test.

Figure 5: JAK/STAT signaling in airway epithelial cells is activated by long term cigarette smoke exposure.

Fluorescence micrographs of the trachea of *upd2* (A-E) or *upd3* (G-K)-*Gal4*; *Stat92E-GFP*; *UAS-LacZ.nls*

larvae that were exposed for 2 days smoke (B and H), heavy smoke (C and I), strong hypoxia (D and J),

and 2 days of hypoxia (E and K). Trachea were stained for GFP (red, JAK/STAT pathway activated zones),

Beta-galactosidase (green, cells that expressed upd2 and upd3), respectively. (F and L) The numbers

of cells that expressed *upd2* and *upd3* were counted under these different conditions. 40 trachea in

each group were assayed. ns means not significant, * $p < 0.05$, ** $p < 0.01$ by Student's t-test. Scale bar:

200 μm.

Figure 6: Changes at the transcript level in the airway epithelium of JAK/STAT mutants. Altered gene

expression levels in the trachea caused by 16 hours persistent expression of *Hop.CA* driven by *btl.ts*. In

total, 2004 genes were regulated significantly ($p < 0.05$), in which 707 genes of (fold change > 2, $p <$

0.01) showing even higher levels of significance. (A) These 707 genes were used to visualize the

differences of the transcript levels between *Hop.CA* expressed trachea and control trachea. From the

outside to the inside, a histogram for expression mean values (red, control; blue, *Hop.CA* group), a

heatmap for controls, a heatmap for *Hop.CA* overexpression, gene names, and PCA analysis, are shown.

According to transcriptome analysis of the tracheal epithelium 1128 genes ($p < 0.05$) were down-

regulated, 876 genes ($p < 0.05$) were up-regulated. The changes in the gene expression total value were

not significant in the ectopic expression trachea compared to the controls. (B) The inhibition of the

JAK/STAT signaling by ectopic expressing *Dome.DN* for 18 hours affects the transcription of 814 genes,

however, which only 102 genes were regulated in both types of airways; with ectopically activated and

inhibited JAK/STAT signaling.

Figure 7: Gene Ontology (GO) enrichment analysis of the 2004 annotated differentially expressed genes in *Hop.CA* overexpressing airways vs controls. (A) Gene ontology analysis showed 10 biological processes that were enriched in *Hop.CA* overexpressing airways compared with matching controls. Most regulated genes in the GO1, GO2, and GO3 groups were down-regulated. Most regulated genes that involved in the COPI-and COPII-mediated vasicular transport between endoplasmic reticulum and Golgi (GO9 and GO10) were up-regulated. GO1, cuticle development; GO2, muscle system process; GO3, muscle cell differentiation; GO4, glutathione metabolic process; GO5, detection of chemical

stimulus; GO6, cellular protein-containing complex assembly; GO7, RNA processing; GO8, translation;

GO9, endoplasmic reticulum to Golgi vesicle-mediated transport; GO10, retrograde vesicle-mediated

transport Golgi to the endoplasmic reticulum. (B) Enriched biological processes were superimposed on

a sketch depicting a tracheal epithelial cell, with the corresponding *p*-value added. Blue indicates

processes where genes were mainly down-regulated; red indicates processes where genes were mainly

up-regulated; green indicates processes where genes were both down-regulated and up-regulated at

similar extents. See also table S10 for annotation of genes involved in each process. All processes

shown displayed $p < 0.05$ (Fisher's exact test).

Figure 8: Activation of airway epithelial JAK/STAT signaling affects vesicle-mediated transport of proteins. (A-E) Coracle (Cora) and Armadillo (Arm) localization in airway epithelial cells. (A) shows the schematic orientation of Arm and Cora only at the contact zone between two airway epithelial cells.

Adherens junction (AJ), Septate junction (SJ). Trachea of *vvl-coin* larvae (control; B and D) and *vvl-*
coin>Hop.CA larvae (treatment; C and E) stained for GFP (green, Gal4 positive cells), Arm or Cora (red),
respectively. 30 specimens were investigated in each group. (F-G) Micrographs of the trachea of *vvl-*
coin larvae (control; F) and *vvl-coin>Hop.CA* larvae (treatment; G). White are places where tracheal
stenosis took place in the regions with cells expressing *Hop.CA* to a high degree. Scale bar: 20 µm in A-
E; 50 µm in F-G. (H) Schematic illustration of the thickening of the epithelium and narrowing of the
tube. In the control epithelium, a secretory burst of luminal proteins drives the diametric expansion of
the tubes and this process depends on vesicle-mediated transport. COPI-and COPII-mediated vesicular
transport plays a central role in this process, mutation members in the process underly the tube size
defect in previous reports [27, 28]. However, long-term induced JAK/STAT activation impedes vesicle-
mediated transport that can be reflected by the deregulation of massive genes involved in COPI-and
COPII-mediated vesicular transport. This dysfunction of transport led to the appearance of tube size
defect such as the increase in the cell volume and narrowing the tube size.

Figure 9: Activation of JAK/STAT signaling in airway epithelial cells affects the development of the epicuticle. (A-C) Activation of JAK/STAT signaling in airway epithelia observed in whole larvae by expressing *UAS-Hop.CA* under the control of the *btl.ts*. Controls (A-C) and induction of expression by shifting the temperature to 29 °C causes the defects in the tracheal epicuticle (A'-C'). 50 control larvae

and *btl.ts>Hop.CA* larvae were investigated in A-C. (A, A') Overview of the whole larvae showing the

whole tracheal system. In (B, B') the region containing Tr7-Tr9 is shown. (C, C') Chitin staining of isolated

airways (at the Tr8 region). (D-F) Micrographs of the trachea of *vvl-coin* larvae under different

conditions. In D, D' the control is shown, in E, E', the undisturbed larval of animals experiencing *Hop.CA*

overexpression are displayed. F, F' similar animals as in E, E', but the animals covered with a slide,

inducing a certain degree of compression. The arrows show the collapsed sites in the tracheal tube.

30 larvae in each group were investigated in D-F. Scale bar: 50 µm in B; 20 µm in C; 200 µm in D-F.

Figure 10: Mild activation of epithelial JAK/STAT signaling mitigated the thickening phenotype. (A-C)

Micrographs of the Tr8-Tr9 regions of *nach-Gal4* larvae (control, A), *nach>Hop.CA* (treatment, B), and

nach>upd3 larvae (treatment, C). (D-E) Quantification of the epithelial thickness of the Tr8-Tr9 region

of those larvae whose JAK/STAT signaling were activated by expressing *Hop.CA* or *upd3* under the

control of *nach-Gal4* and *btl-Gal4*. (F) Micrographs of *nach>Dome.DN* larvae experiencing suppression

of the JAK/STAT pathway make trachea translucent and filled with fluids (arrows); triangle indicates the

tracheal position. 30 larvae in each group were used to quatify the thickness. ns means not significant,

* *p* < 0.05, ** *p* < 0.01 by Student's t-test. Scale bar: 20 µm in A-C; 200 µm in F.