Title: Short sequence motif dynamics in the SARS-CoV-2 genome suggest a role for

cytosine deamination in CpG reduction.

Authors: Mukhtar Sadykov[1*], Tobias Mourier[1*], Qingtian Guan[1], Arnab Pain[1,2,3**]

Affiliation:

[1]King Abdullah University of Science and Technology (KAUST), Pathogen Genomics

Laboratory, Biological and Environmental Science and Engineering (BESE), Thuwal-

Jeddah, 23955-6900, Saudi Arabia;

[2]Research Center for Zoonosis Control, Global Institution for Collaborative Research

and Education (GI-CoRE); Hokkaido University, N20 W10 Kita-ku, Sapporo, 001-0020

Japan;

[3]Nuffield Division of Clinical Laboratory Sciences (NDCLS), University of Oxford,

Headington, Oxford, OX3 9DU, United Kingdom

*These authors contributed equally

**Correspondence: Arnab Pain Email: arnab.pain@kaust.edu.sa. King Abdullah

University of Science and Technology, Jeddah, Saudi Arabia. Phone: (+966) 54 470

0687

Abbreviations: C>U stands for cytosine to uracil substitution, the same applies to other

nucleotide substitutions; APOBEC - Apolipoprotein B Editing Complex; ZAP – zinc-

finger antiviral protein.

Keywords: virus evolution, genome evolution, genome biology, virus-host interaction.

ABSTRACT

RNA viruses use CpG reduction to evade the host cell defense, but the driving

mechanisms are still largely unknown. In an attempt to address this we used a rapidly

growing genomic dataset of SARS-CoV-2 with relevant metadata information.

Remarkably, by simply ordering SARS-CoV-2 genomes by their date of collection, we

find a progressive increase of C-to-U substitutions resulting in 5'-UCG-3' motif reduction

that in turn have reduced the CpG frequency over just a few months of observation.

This is consistent with APOBEC-mediated RNA editing resulting in CpG reduction, thus

allowing the virus to escape ZAP-mediated RNA degradation. Our results thus link the

dynamics of target sequences in the viral genome for two known host molecular

defense mechanisms, mediated by the APOBEC and ZAP proteins.

Introduction

Viruses utilize numerous mechanisms to avoid the host cell defense. One such

mechanism is the CpG dinucleotide reduction observed in many single-stranded RNA

viruses (Atkinson et al. 2014). Low CpG abundance is typically a consequence of two

factors: constraints of amino-acid conservation and selection pressure from diverse host

cell defense mechanisms targeting CpG (Gu et al. 2019).

Prior to the involvement of the host immune system, the infected cells have to rely on

their own antiviral restriction factors. These factors are mostly proteins induced by

interferon signaling cascade (Chemudupati et al. 2019). The antiviral proteins directly

involved in the restriction of viral replication and transcription are zinc-finger antiviral

protein (ZAP) and Apolipoprotein B Editing Complex (APOBEC) (Kluge et al. 2015). The

defensive role of ZAP on HIV-1 manifestation in the host cell has been extensively

studied (Ficarelli et al. 2019). ZAP selectively binds to viral CpG regions and recruits

RNA exosome complex resulting in viral RNA degradation (Harris and Dudley 2015;

Takata et al. 2017). Recently, ZAP was shown to restrict SARS-CoV-2 production upon

the induction by interferons (Nchioua et al. 2020).

The APOBEC protein family is another example of host antiviral enzymes. Seven

human APOBEC proteins are known for catalyzing cytosine to uracil deamination in

foreign single-stranded DNA (ssDNA) and RNA (ssRNA) (Sharma et al. 2015; Salter

and Smith 2018; Sharma et al. 2019; Giorgio et al. 2020). APOBEC proteins induce

extensive nucleotide changes, which is restrictive to viral replication, but they also

contribute to the genetic diversity and adaptation of a virus (Kim et al. 2014; Harris and

Dudley 2015). Enzymatic target motifs for most of the APOBEC enzymes have been

experimentally identified, among which the most common were 5'-[T/U]C-3' and 5'-CC-

3' for DNA/RNA substrates (Thielen et al. 2010; Sharma et al. 2015; Salter and Smith

2018; Sharma et al. 2019). Some members of APOBEC proteins inhibit replication of

human coronavirus, HCoV-NL63, but their role in hypermutation of the viral genome is

not clear (Milewska et al. 2018). However, it was recently suggested that the SARS-

CoV-2 undergoes genome editing by host-dependent RNA-editing proteins such as

APOBEC (C>U) and ADAR (A>G) (Giorgio et al. 2020; Simmonds 2020). mRNA

transcriptional response data from COVID-19 patients demonstrated that APOBEC3A

(A3A) belongs to one of the most expressed genes in response to SARS-CoV-2, and

the highest among APOBEC family (Blanco-Melo et al. 2020).

SARS-CoV-2 is an enveloped positive-sense ssRNA virus, belonging

to *Coronaviridae* family, which is characterized by large genomes (~30K nucleotide

long), high recombination frequencies and genomic proofreading mechanism (Fehr and

Perlman 2015). The COVID-19 pandemic is bringing intensive scientific attention to the

virus, and thousands of viral genomic sequences with metadata are publicly available

(Elbe and Buckland-Merrett 2017; Hadfield et al. 2018). Given the large amount of

available data and the relatively low mutation rate of the virus (Salemi et al. 2004), we

aimed to monitor its genomic evolution on a very brief time scale during the COVID-19

pandemic. We show progressive C>U substitutions in SARS-CoV-2 genome within the

timeframe of five months. We highlight the role of C>U substitutions in the reduction of

5'-UCG-3' motifs and hypothesize that this progressive decrease is driven by host

APOBEC and ZAP enzymes.

Results and Discussion

Figure 1. (A) SNV events observed between individual SARS-CoV-2 sample

sequences (n=22,164) and the reference genome. (B) The number of C>U substitutions

across sample dates. The average number of substitutions for each sampling day is

plotted (blue line, left y-axis) with plus/minus one standard deviations as error bars. The

number of samples for each day is shown as red bars (right y-axis). (C) Folding

potential of positions with C>U changes (see Materials and Methods). P-values from

Fisher's exact test are shown above bars. (D) The fraction of [A/C/G/U]CG triplets that

are changed to [A/C/G/U]UG over time. The average fractions, relative to the reference

genome, are shown as circles for each sampling day (x-axis). Error bars denote

plus/minus one standard deviation. (E) A model for the consequences of host-driven

evolution by APOBEC enzymes on viral CpG dinucleotide composition.

GISAID data of viral sequences including their date of collection allowed us to analyze

viral nucleotide change dynamics over time. We have analyzed 22,164 SARS-CoV-2

genomes from GISAID deposited from December 24[th], 2019 to May 23[rd], 2020 (Elbe

and Buckland-Merrett 2017) (Figure S1 & S2; Table S1). By aligning these genomes to

the SARS-CoV-2 reference (GenBank accession: NC_045512), we observed a total of

9,210 single nucleotide variations (SNVs) with C>U being the most abundant

substitution and representing almost one third of all substitution events (Figure 1A). As

nearly half of the C positions in the reference are substituted to U in at least one

sample, we expect a high degree of saturation. Downsizing the number of samples

resulted in a consistent increase in C>U frequencies, suggesting that a considerable

fraction of the observed C>U changes represent multiple, independent events (Figure

S3). The reported C>U frequency is therefore most likely an underestimate. Over this

period of five months, we find a steady increase in C>U substitutions (Figure 1B). One

potential driver behind this increase could be the recently proposed APOBEC-mediated

viral RNA editing (Giorgio et al. 2020; Simmonds 2020). Since APOBEC3 family

members display a preference for RNA in open conformation as opposed to forming

secondary structures (McDaniel et al. 2020), we calculated the folding potential of all

genomic sites that include C>U substitutions (Figure 1C). Positions with C>U changes

are more often located in regions with low potential for forming secondary RNA

structures. These observations are in agreement with the notion that members of the

APOBEC family are the main drivers of cytosine deamination in SARS-CoV-2, thus

contributing to the extensive viral genome editing (Giorgio et al. 2020; Simmonds 2020).

The majority of APOBEC enzymes preferentially acts on a 5'-UC-3' (5'-TC-3' for DNA

substrate) motif (Thielen et al. 2010; Sharma et al. 2015; Salter and Smith 2018;

Sharma et al. 2019). We searched for possible APOBEC genetic footprints (5'-UC-3' >

5'-UU-3') in viral dinucleotide frequencies for the 5-month period (Figure S4). Among all

dinucleotides, UpC showed the highest degree of decrease, while UpU exerted the

highest rates of increase. CpG, CpA, CpC, CpU and GpA also showed a negative net

gain but not as prominent as UpC, CpA or ApG. We find that the majority of dinucleotide

losses were due to C>U changes, in agreement with a recent study by Rice and

colleagues (Rice et al. 2020). The remaining dinucleotides showed relative stability

across the time period. Interestingly, recent reports showed the reduction of CpG

dinucleotides in 5'-UTR and ORF1ab regions (Wang et al. 2020; Wei et al. 2020; Zhang

et al. 2020). When analyzing the context of genomic sites undergoing C>U changes we

noticed an enrichment for 5'-UCG-3' motifs (Table S2). To assess the contribution of

C>U changes in CpG loss, we examined the dynamics of [A/C/G/U]CG trinucleotides

over time (Figure 1D). The progressive change (~1% over a 5-month period) of 5'-UCG-

3' to 5'-UUG-3' is most striking when supported by a larger number of genomes (days

70 to 115), whereas no such pattern is observed for the other trinucleotides (Figure 1D).

The association between cytosine deamination and CpG loss is further underlined by

the rapid, progressive increase in 5'-UCG-3' > 5'-UUG-3' changes compared to other 5'-

UC[A/C/U]-3' motifs (Figure S5).

Our CpG net gain analysis showed a decline, indicating that the number of total CpG

losses overrides their gain (Figure S4), in agreement with recently published

observations (Wang et al. 2020; Wei et al. 2020; Zhang et al. 2020). Similar to other

coronavirus genomes, SARS-CoV-2 has a low CpG frequency around 1.5%, while the

expected frequency is 3.6% (Wang et al. 2020). The observed CpG decline could

indicate the ongoing adaptation of the virus to the new host or tissues where antiviral

proteins targeting foreign CpG regions are differently expressed (Xia 2020). It was

recently shown that synonymous substitutions disrupting CpG dinucleotides appear to

contribute to the increased virulence of attenuated poliovirus (Stern et al. 2017). We

observe that more than half of the C>U substitutions disrupting CpG dinucleotides were

synonymous, consistent with recent studies (Di Gioacchino et al. 2020; Rice et al.

2020) (Figure S6). In the future, the SARS-CoV-2 virus may level up its CpG frequency

with its host, achieving so-called host-cell mimicry (Greenbaum et al. 2008), but the

observed CpG loss suggests that equilibrium between gains and losses have not yet

been reached.

Previously, from a comparative analysis of different coronaviruses, Woo and colleagues

concluded that CpG reduction and cytosine deamination are two independent selective

forces that shape coronavirus codon bias (Woo et al. 2007). Although selective forces

may act differently upon motifs resulting from C>U changes, our results suggest a direct

link between cytosine deamination and CpG reduction in SARS-CoV-2.

Previous studies reported that a reduced number of CpG motifs in HIV and other

viruses played an important role in the viral replication inside the host cell, thus

escaping ZAP protein activity (Takata et al. 2017). Given the high expression levels of

APOBEC and ZAP genes in COVID-19 patients (Blanco-Melo et al. 2020), the

involvement of ZAP-mediated restriction of SARS-CoV-2 (Nchioua et al. 2020), and our

observations, we hypothesize that as a consequence of APOBEC-mediated RNA

editing, SARS-CoV-2 genome may escape host cell ZAP activity. Both APOBEC and

ZAP are interferon-induced genes that act preferentially on ssRNA in open conformation

(Chemudupati et al. 2019; Luo et al. 2020; McDaniel et al. 2020). Initially, APOBEC and

ZAP enzymes may have overlapping preferred target motifs for their enzymatic

functions (Figure 1E). The catalytic activity of APOBEC on 5'-UC-3' leads to cytosine

deamination, which destroys ZAP's extremely specific acting site (CpG) (Meagher et al.

2019). The conversion of C>U allows viral RNA to escape from ZAP-mediated RNA

destruction. A recent study hypothesized that both ZAP and APOBEC provide selective

pressure that drives the adaptation of SARS-CoV-2 to its host (Wei et al. 2020). Here

we provided one of the potential mechanisms that contribute to CpG reduction in SARS-

CoV-2. Our analysis together with other recent studies strongly supports the proposed

model, but it merits future experimental validation. Since APOBEC enzymes can act on

both RNA and DNA sequences, it is tempting to speculate that some other viruses may

undergo CpG reduction similarly to SARS-CoV-2.

In summary, the rapidly growing genomic data from SARS-CoV-2 (that importantly

includes relevant metadata) and the moderate mutation rate of the virus has allowed us

to observe a progressive increase of C-to-U substitutions resulting in CpG loss over a

brief period of time. Notably, this was achieved using a phylogeny-free approach,

exclusively considering the temporal aspect of the samples. Given that APOBEC

proteins induce extensive mutations in a viral genome which are mostly detrimental,

here we provide an example of a rather unexpected role of these proteins in facilitating

the adaptation of a virus to the host. To our knowledge, this is the first study linking the

dynamics of viral genome mutation to two known host molecular defense mechanisms,

involving the APOBEC and ZAP proteins.

Materials and Methods

Samples and recording of substitutions

Samples were downloaded from GISAID (https://www.gisaid.org/) on May 23rd, 2020

(Elbe and Buckland-Merrett 2017). Only samples with available collection dates and

with no more than 500 N's in the assembled sequence were used. December 24th,

2019 was set to day1. Assembled virus genomes were separately aligned to the

reference genome (GenBank accession: NC_045512) using MAFFT (Katoh and

Standley 2013), and mismatches were collected. Alignments with more than 500 gaps

in the sample (due to incomplete assembly) and any alignments with gaps in the

reference sequence were discarded. This resulted in 22,164 alignments. Only non-

ambiguous bases and only mismatches without gaps in 3 positions up- and downstream

were kept. To exclude alignment artifacts, mismatches in the first and last 100 positions

in the reference genome were discarded.

The total number of substitutions is recorded from the combined set of pair-wise

alignments, and a substitution at a given position in the reference is counted once

regardless of the number of samples in which it is found. The timeline analysis of

dinucleotide gains and losses were inferred from each sample's individual alignment to

the reference.

Nucleotide motifs

For all di-, tri-, and tetra-nucleotide motifs containing C in the reference genome, the

number of motifs with a C-to-U substitution, and the total number of the motifs were

noted. The ratio between these two measures was compared to the expected ratio,

defined as the number of C's with a C-to-U substitution divided by the total number of

C's in the genome. The probability was calculated using a binomial distribution. All

statistical tests were performed using Rstudio v. 1.1.414 (Booth et al. 2018).

<u>Folding potential</u>

The reference sequence was divided into overlapping 500-nucleotides windows, each

shifted by 10 nucleotides. These sequences were folded using RNAfold (Lorenz et al.

2011). For each position in the reference genome, the number of windows in which the

position was predicted to form a base-pair in a secondary structure was recorded. The

folding potential of a position was defined as the fraction of windows with predicted

base-pairing of that position. Hence, a position predicted to participate in a base-pair in

all 500-nt windows has a folding potential of 1.

Supplementary Material

Figure S1. Number of samples from different time points. December 24th 2019 is set to

day 1. Samples are colored according to their geographic region.

Figure S2. Substitution occurrences in samples. Most substitutions (15,750) were

observed in just a single sample, whereas a single substitution was found in 4,683

samples.

Figure S3. The inverse relationship of C>U frequency and the number of samples

included. Down-sizing the number of samples leads to higher C>U frequency (0.1 on

the x-axis means 10% of the total samples), suggesting that as more and more samples

are included the prominent C>U changes happen as independent events at identical

positions (hence only counted once in this analysis). As this saturation is expected to be

lower for other, less abundant changes this will result in an apparent drop in the fraction

of changes constituted by C>U. The blue dots are the individual simulated data sets,

and the boxes denote their distribution.

Figure S4. Number of dinucleotide gains and losses over time.

Figure S4 (cont.). Number of dinucleotide gains and losses over time.

Figure S4 (cont.). Number of dinucleotide gains and losses over time.

251
252 **Figure S4 (cont.).** Number of dinucleotide gains and losses over time.

253
254 **Figure S5.** The fraction of UC[A/C/G/U] to UU[A/C/G/U] changes over time. The fraction

255 of trinucleotide changes relative to the reference genome are shown as circles for each

256 sampling day (x-axis). Error bars denote plus/minus one standard deviation.

Figure S6. Plots showing the number C positions (top) and CpG positions (bottom) in

the reference genome (black bars) and the number of these that have C>U changes

(grey bars). Numbers are shown for different codon positions as indicated below the

plots. CpG positions are defined as the position of the C in the motif, and positions with

no changes are defined as either synonymous or non-synonymous based on the effect

264   of a potential C>U change. The percentage of bases and motifs that have C>U changes

265   are shown as open circles (right y-axis).

266 **Table S2.** The most enriched motifs undergoing C>U changes.

	C>U		
motif	motifs with C-to-U change/total number of motifs	odds-ratio	uncorrected p-value
UCG	72/113	1.30	0.0012387
UUCG	27/39	1.41	0.0088244
CG	257/439	1.19	4.56E−05
CGG	55/76	1.47	3.19E−05
CGGA	13/16	1.65	0.0088056
CGGU	22/29	1.54	0.0030566
GUGC	75/113	1.35	0.0001577
UGCU	148/255	1.18	0.0026032
CUA	307/561	1.11	0.0044054
UUAC	137/241	1.16	0.0096264
UGAC	87/137	1.29	0.0004834
CAAC	112/194	1.18	0.0098096
UAAC	88/135	1.33	0.00012

Acknowledgments

We thank all laboratories which have contributed sequences to the GISAID database

and Zhadyra Yerkesh for giving her comments and helpful discussions.

This work was supported by funding from King Abdullah University of Science and

Technology (KAUST), Office of Sponsored Research (OSR). Work in AP's laboratory is

supported by the KAUST faculty baseline fund (BAS/1/1020-01-01).

Author Contributions

A.P. supervised the project. M.S. and T.M. designed experiments. T.M. and QG

performed bioinformatic analysis. M.S. wrote the draft of the manuscript. All authors

discussed, edited, read, and agreed to the final version of the manuscript.

Availability of Data

The data underlying this article are available in GISAID, at https://gisaid.org. The ID

numbers of genomes used are provided in Table S1.

References

Atkinson NJ, Witteveldt J, Evans DJ, Simmonds P. 2014. The influence of CpG and UpA dinucleotide frequencies on RNA virus replication and characterization of the innate cellular pathways underlying virus attenuation and enhanced replication. Nucleic Acids Res. doi:10.1093/nar/gku075.

Blanco-Melo D, Nilsson-Payant BE, Liu WC, Uhl S, Hoagland D, Møller R, Jordan TX, Oishi K, Panis M, Sachs D, et al. 2020. Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19. Cell. doi:10.1016/j.cell.2020.04.026.

Booth DS, Szmidt-Middleton H, King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, Fairclough SR, Hellsten U, et al. 2018. RStudio: Integrated Development for R. Nature. doi:10.1108/eb003648.

Chemudupati M, Kenney AD, Bonifati S, Zani A, McMichael TM, Wu L, Yount JS. 2019. From APOBEC to ZAP: Diverse mechanisms used by cellular restriction factors to inhibit virus infections. Biochim Biophys Acta - Mol Cell Res. doi:10.1016/j.bbamcr.2018.09.012.

Elbe S, Buckland-Merrett G. 2017. Data, disease and diplomacy: GISAID's innovative contribution to global health. Glob Challenges. doi:10.1002/gch2.1018.

Fehr AR, Perlman S. 2015. Coronaviruses: An overview of their replication and pathogenesis. In: Coronaviruses: Methods and Protocols.

Ficarelli M, Antzin-Anduetza I, Hugh-White R, Firth AE, Sertkaya H, Wilson H, Neil SJD, Schulz R, Swanson CM. 2019. CpG Dinucleotides Inhibit HIV-1 Replication through Zinc Finger Antiviral Protein (ZAP)-Dependent and -Independent

Mechanisms. J Virol. doi:10.1128/jvi.01337-19.

Di Gioacchino A, Sulc P, Komarova A V., Greenbaum BD, Monasson R, Cocco S. 2020.

The Heterogeneous Landscape and Early Evolution of Pathogen-Associated

CpG Dinucleotides in SARS-CoV-2. SSRN Electron J.

doi:10.2139/ssrn.3611280.

Giorgio S Di, Martignano F, Torcia MG, Mattiuz G, Conticello SG. 2020. Evidence for

host-dependent RNA editing in the transcriptome of SARS-CoV-2. Sci Adv.

doi:10.1126/sciadv.abb5813.

Greenbaum BD, Levine AJ, Bhanot G, Rabadan R. 2008. Patterns of evolution and host

gene mimicry in influenza and other RNA viruses. PLoS Pathog.

doi:10.1371/journal.ppat.1000079.

Gu H, Fan RLY, Wang D, Poon LLM. 2019. Dinucleotide evolutionary dynamics in

influenza A virus. Virus Evol. doi:10.1093/ve/vez038.

Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, Sagulenko P,

Bedford T, Neher RA. 2018. NextStrain: Real-time tracking of pathogen

evolution. Bioinformatics. doi:10.1093/bioinformatics/bty407.

Harris RS, Dudley JP. 2015. APOBECs and virus restriction. Virology.

doi:10.1016/j.virol.2015.03.012.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability. Mol Biol Evol.

doi:10.1093/molbev/mst010.

Kim EY, Lorenzo-Redondo R, Little SJ, Chung YS, Phalora PK, Maljkovic Berry I,

Archer J, Penugonda S, Fischer W, Richman DD, et al. 2014. Human APOBEC3

Induced Mutation of Human Immunodeficiency Virus Type-1 Contributes to

Adaptation and Evolution in Natural Infection. PLoS Pathog.

doi:10.1371/journal.ppat.1004281.

Kluge SF, Sauter D, Kirchhoff F. 2015. SnapShot: Antiviral Restriction Factors. Cell.

doi:10.1016/j.cell.2015.10.019.

Lorenz R, Bernhart SH, Höner zu Siederdissen C, Tafer H, Flamm C, Stadler PF,

Hofacker IL. 2011. ViennaRNA Package 2.0. Algorithms Mol Biol.

doi:10.1186/1748-7188-6-26.

Luo X, Wang X, Gao Y, Zhu J, Liu S, Gao G, Gao P. 2020. Molecular Mechanism of

RNA Recognition by Zinc-Finger Antiviral Protein. Cell Rep.

doi:10.1016/j.celrep.2019.11.116.

McDaniel YZ, Wang D, Love RP, Adolph MB, Mohammadzadeh N, Chelico L, Mansky

LM. 2020. Deamination hotspots among APOBEC3 family members are defined

by both target site sequence context and ssDNA secondary structure. Nucleic

Acids Res. doi:10.1093/nar/gkz1164.

Meagher JL, Takata M, Gonçalves-Carneiro D, Keane SC, Rebendenne A, Ong H, Orr

VK, MacDonald MR, Stuckey JA, Bieniasz PD, et al. 2019. Structure of the zinc-

finger antiviral protein in complex with RNA reveals a mechanism for selective

targeting of CG-rich viral sequences. Proc Natl Acad Sci U S A.

doi:10.1073/pnas.1913232116.

Milewska A, Kindler E, Vkovski P, Zeglen S, Ochman M, Thiel V, Rajfur Z, Pyrc K. 2018.

351  APOBEC3-mediated restriction of RNA virus replication. Sci Rep.

352   doi:10.1038/s41598-018-24448-2.

353 Nchioua R, Kmiec D, Mueller J, Conzelmann C, Gross R, Swanson CM, Neil S, Stenger

354   S, Sauter D, Muench J, et al. 2020. The Zinc Finger Antiviral Protein restricts

355   SARS-CoV-2. bioRxiv. doi:10.1101/2020.06.04.134379.

356 Rice AM, Morales AC, Ho AT, Mordstein C, Mühlhausen S, Watson S, Cano L, Young

357   B, Kudla G, Hurst LD. 2020. Evidence for strong mutation bias towards, and

358   selection against, U content in SARS-CoV-2: implications for vaccine design. Mol

359   Biol Evol. doi:10.1093/molbev/msaa188.

360 Salemi M, Fitch WM, Ciccozzi M, Ruiz-Alvarez MJ, Rezza G, Lewis MJ. 2004. Severe

361   Acute Respiratory Syndrome Coronavirus Sequence Characteristics and

362   Evolutionary Rate Estimate from Maximum Likelihood Analysis. J Virol.

363   doi:10.1128/jvi.78.3.1602-1603.2004.

364 Salter JD, Smith HC. 2018. Modeling the Embrace of a Mutator: APOBEC Selection of

365   Nucleic Acid Ligands. Trends Biochem Sci. doi:10.1016/j.tibs.2018.04.013.

366 Sharma S, Patnaik SK, Thomas Taggart R, Kannisto ED, Enriquez SM, Gollnick P,

367   Baysal BE. 2015. APOBEC3A cytidine deaminase induces RNA editing in

368   monocytes and macrophages. Nat Commun. doi:10.1038/ncomms7881.

369 Sharma S, Wang J, Alqassim E, Portwood S, Cortes Gomez E, Maguire O, Basse PH,

370   Wang ES, Segal BH, Baysal BE. 2019. Mitochondrial hypoxic stress induces

371   widespread RNA editing by APOBEC3G in natural killer cells. Genome Biol.

372   doi:10.1186/s13059-019-1651-1.

Simmonds P. 2020. Rampant C->U hypermutation in the genomes of SARS-CoV-2 and

other coronaviruses - causes and consequences for their short and long

evolutionary trajectories. biorxiv. doi:10.1101/2020.05.01.072330.

Stern A, Yeh M Te, Zinger T, Smith M, Wright C, Ling G, Nielsen R, Macadam A,

Andino R. 2017. The Evolutionary Pathway to Virulence of an RNA Virus. Cell.

doi:10.1016/j.cell.2017.03.013.

Takata MA, Gonçalves-Carneiro D, Zang TM, Soll SJ, York A, Blanco-Melo D, Bieniasz

PD. 2017. CG dinucleotide suppression enables antiviral defence targeting non-

self RNA. Nature. doi:10.1038/nature24039.

Thielen BK, McNevin JP, McElrath MJ, Hunt BVS, Klein KC, Lingappa JR. 2010. Innate

immune signaling induces high levels of TC-specific deaminase activity in

primary monocyte-derived cells through expression of APOBEC3A isoforms. J

Biol Chem. doi:10.1074/jbc.M110.102822.

Wang Y, Mao J-M, Wang G-D, Luo Z-P, Yang L, Yao Q, Chen K-P. 2020. Human

SARS-CoV-2 has evolved to reduce CG dinucleotide in its open reading frames.

Sci Rep. 10(1):12331. doi:10.1038/s41598-020-69342-y.

https://doi.org/10.1038/s41598-020-69342-y.

Wei Y, Silke JR, Aris P, Xia X. 2020. Coronavirus genomes carry the signatures of their

habitats. bioRxiv. doi:10.1101/2020.06.13.149591.

Woo PCY, Wong BHL, Huang Y, Lau SKP, Yuen KY. 2007. Cytosine deamination and

selection of CpG suppressed clones are the two major independent biological

forces that shape codon usage bias in coronaviruses. Virology.

doi:10.1016/j.virol.2007.08.010.

Xia X. 2020. Extreme genomic CpG deficiency in SARS-CoV-2 and evasion of host

antiviral defense. Mol Biol Evol. doi:10.1093/molbev/msaa094.

Zhang J, Kang J, Liu M, Han B, Li L, He Y, Yi Z, Chen L. 2020 Jan 1. Multi-site co-

mutations and 5'UTR CpG immunity escape drive the evolution of SARS-CoV-2.

bioRxiv.:2020.07.21.213405. doi:10.1101/2020.07.21.213405.

http://biorxiv.org/content/early/2020/07/27/2020.07.21.213405.abstract.

402