Exploring the complexity of soybean (*Glycine max*) transcriptional regulation using global gene co-expression networks

Fabricio Almeida-Silva[1], Kanhu C. Moharana[1], Fabricio B. Machado[1], Thiago M. Venancio[1]*

[1]Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e Biotecnologia, Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos Goytacazes, RJ, Brazil.

*TMV: Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e Biotecnologia, Universidade Estadual do Norte Fluminense Darcy Ribeiro. Av. Alberto Lamego 2000, P5, sala 217, Campos dos Goytacazes, RJ, Brazil. Email: thiago.venancio@gmail.com

ABSTRACT

Soybean (*Glycine max* (L.) Merr.) is one of the most important crops worldwide, constituting a major source of protein and edible oil. Gene co-expression networks (GCN) have been extensively used to study transcriptional regulation and evolution of genes and genomes. Here, we report a soybean GCN using 1,284 publicly available RNA-Seq samples from 15 distinct tissues. We found modules that are differentially regulated in specific tissues, comprising processes such as photosynthesis, gluconeogenesis, lignin metabolism, and response to biotic stress. We identified transcription factors among intramodular hubs, which probably integrate different pathways and shape the transcriptional landscape in different conditions. The top hubs for each module tend to encode proteins with critical roles, such as succinate dehydrogenase and RNA polymerase subunits. Importantly, gene essentiality was strongly correlated with degree centrality and essential hubs enriched in genes involved in nucleic acids metabolism and regulation of cell replication. By using a using a guilt-by-association approach, we predicted functions for 93 of 106 hubs without functional description in soybean. Most of the duplicated genes had different transcriptional profiles, supporting their functional divergence, although paralogs originating from whole-genome duplications (WGD) are more often preserved in the same module than those from other mechanisms. Together, our results highlight the importance of GCN analysis in unraveling key functional aspects of the soybean genome, in particular those associated with hub genes and WGD events.

Keywords: transcription factors, gene duplication, gene function, genome evolution, systems biology.

INTRODUCTION

Soybean (*Glycine max* (L.) Merr.) is one of the most important legume crops worldwide, being important for human nutrition, animal feed, and biotechnological applications. It has a paleopolyploid genome that resulted from two whole-genome duplication (WGD) events that happened 58 and 13 million years ago [1]. Because of these WGD events, ~75% of the soybean genes are present in multiple copies [2]. Over the last decade, several transcriptomic studies have investigated soybean transcriptional programs in different tissues and conditions [3-6]. However, most works explore only a few conditions and developmental stages, typically using pathway/function enrichment methods. Such approaches can overlook a myriad of interesting transcriptional patterns, which could otherwise be unraveled by integrative studies using hundreds of samples submitted to systems-level analyses.

A powerful approach to further explore transcriptomic data is the use of co-expression networks. Co-expression networks are undirected graphs comprising nodes representing genes that are connected by edges whenever significantly co-expressed. Co-expression networks have been used to understand transcriptional regulation in many plant species, such as maize (*Zea mays* L.), rice (*Oryza sativa* L.), tomato (*Solanum lycopersicum* L.), thale cress (*Arabidopsis thaliana* (L.) Heynh), and soybean [7-10]. However, most co-expression network analyses focus on case-control experiments, limiting their potential to uncover broader biological patterns [11-13]. A recent global co-expression network analysis of 1,072 soybean microarray samples revealed a gene module that is likely involved in the evolution of nodulation in plants [14]. In spite of these interesting results, hybridization-based transcriptional quantification often suffer from poor reproducibility [15]. Over the past few years, data from hundreds of soybean RNA-seq samples have accumulated and were recently integrated by our group in a large and publicly available soybean transcriptome atlas [16]. Nevertheless, a network analysis using this collection remains to be performed.

Co-expression analyses have recently been used to study the evolution of gene function in plants [17, 18]. After a duplication event, selective constraints are relaxed and gene copies typically undergo two possible fates: i. gene fractionation, when one copy accumulate loss-of-function mutations and eventually erode from the genome and; ii. gene retention via subfunctionalization or neofunctionalization [19]. Although pseudogenization is the most common fate of duplicated genes [20], genes encoding regulatory proteins (e.g. transcription factors – TFs – and signal transduction proteins) or subunits of protein complexes (e. g. proteasome and ribosome) are retained more often than expected [21, 22], mostly because of the maintenance of their stoichiometry [21]. However, in case of local duplications (e.g. tandem duplications), TFs and subunits of protein complexes tend to be fractionated because of dosage sensitivity [23].

Gene retention is thought to be more likely if the newly duplicated genes remain in the genome for an extended window of time, allowing mutation and

selection pressures to rewire gene functions [24]. The vast majority of retained genes
undergo subfunctionalization and, more rarely, neofunctionalization [25]. At the
transcriptional level, subfunctionalization of a gene pair typically happens through
mutations at cis-regulatory regions, resulting in two possible scenarios: i. both genes
reduce their transcription, making them both necessary to perform the original
function or; ii. genes from the pair become expressed in different tissues or conditions
(transcriptional divergence) [26]. Thus, a network-level gene expression analysis
could shed light on the possible fates of duplicated genes.
Here, we report a GCN built with data from 1,284 RNA-seq samples to
investigate the global transcriptional landscape of soybean in different tissues and
conditions. We identified biological processes and metabolic pathways that are
enhanced or repressed in particular tissues and pointed TF families that probably
control their expression, including candidates for downstream experiments (e.g.
ectopic expression). We predicted the function of 93 unannotated hubs using a guilt-
by-association approach. Further, we used this large GCN to gain insight into the fate
of duplicated genes. We demonstrate that most soybean duplicated genes diverge in
their expression profiles. However, duplicate pairs originating from WGDs are more
often preserved in the same module than those arising by other mechanisms,
suggesting an increased conservation of gene function for WGD-derived copies. We
also demonstrate that duplicate pairs in the same module are likely under stronger
purifying selection, as revealed by their lower Ka/Ks ratios. Collectively, our findings
reveal potentially novel regulators and enlighten our understanding of the fate of
duplicated genes during the evolution of the soybean genome.
89
MATERIAL AND METHODS
Data collection and processing
The normalized expression data (TPM, Transcripts Per Million) for 1,298 samples were
downloaded from the recently published Soybean Expression Atlas [16]. Samples
were filtered by using the standardized connectivity (Z.k) method in the R package
WGCNA [27]. For that, we constructed a sample network based on squared Euclidean
distances and assessed the overall scaled connectivity (Z.k) of the network. This
method is efficient in filtering outlying samples and has been largely used in gene co-
expression analyses [28-30]. Samples with Z.k < -2.5 were considered outliers and
were filtered out. The resulting 1,284 samples represent 15 different tissues
(Supplementary Figure S1). Further, to reduce noise, we excluded genes with TPM <
5 in more than 20% of the samples. Finally, the 30,563 x 1,284 matrix was $\log_2$
transformed by using log (TPM+1).
103
Network reconstruction and module detection
The signed network was inferred by using the R package *WGCNA* [27]. We assessed
the scale-free topology fit to choose the most suitable β power. The lowest β power
for which the network reached a scale-free topology (R^2 = 0.8) was 17

(Supplementary Figure S2). Pairwise correlations between genes were calculated by
using the Pearson Correlation Coefficient (PCC). The PCC matrix was then
transformed to an adjacency matrix, which was used to construct a topological
overlap matrix (TOM). The TOM considers both PCC and shared neighbors to
determine similarity, which amplifies disparity between gene pairs, making it
effective for module detection. Genes were clustered into modules with hierarchical
clustering based on dissimilarity (1 – TOM). Finally, we calculated pairwise similarity
between modules and merged the ones that had PCC ≥ 0.8. Minimum module size
was set to 30. The R package *igraph* (http://igraph.org) [31] was used for extraction
and visualization of subgraphs.
118
Hub gene inference and functional enrichment
Hub genes are those with the highest number of connections (i. e. degree) in a
network. Intramodular connectivity (kWithin) was used as degree to detect
intramodular hubs. The choice of thresholds for hub identification is somewhat
arbitrary. While some groups consider the top 10% genes with the highest degree as
hubs [32, 33], others use module membership (MM) to define such genes [34]. MM
is a measure of how close a given gene is to its module, and it can be calculated by
correlating a gene to the module eigengene. Here, we combined both parameters
and considered hubs the top 10% genes with the highest intramodular connectivity
and with MM ≥ 0.8. Enrichment analysis for GO terms, pathways and conserved
protein domains was performed in PhytoMine
(https://phytozome.jgi.doe.gov/phytomine/begin.do), and REVIGO was used to
remove redundancy [35]. Soybean TFs were downloaded from PlantTFDB [36], and
TF enrichment was determined by performing an one-sided Fisher's exact test with
the R package *bc3net* [37]
134
Analysis of upstream cis-regulatory elements
We searched for regulatory motifs in the regions between -1000bp and +200bp
relative to the transcription start site, as done in *A. thaliana* [38]. These regions were
extracted with *bedtools* [39]. A *de novo* motif discovery was performed by using
MEME [40]. As background, we used an order 0 Hidden Markov Model (HMM) with all
expressed genes, using the script *fasta-get-markov* available in MEME suite.
141
Simulated networks, paralogous genes and selection analysis
We simulated 10,000 degree-preserving networks in R through successive
permutations of node labels to verify the probability of finding our results by chance.
Paralogous gene pairs derived from dispersed, tandem, proximal, transposed, and
whole-genome duplications were download from the Plant Duplicate Gene Database
[41]. We used Ks thresholds to subdivide WGD-derived pairs into the WGDs that
happened 13 mya (Ks ≤ 0.4) and 58 mya (Ks > 0.4). For simplicity, a smaller number
of pairs potentially arising from older WGD events were combined in this latter

category. Ka, Ks and Ka/Ks ratios were computed using the Perl script
calculate_Ka_Ks_pipe.pl, available in [41]. Paralog pairs in the same module were
interpreted as having similar transcriptional profiles. Pairs were interpreted as having
different transcriptional profiles if: i. genes were distributed in different modules; ii.
only one gene from the pair was present in the network or; iii. both genes were in
the *grey* module, which stores genes that were not assigned to modules.
156
RESULTS AND DISCUSSION
Degree centrality analysis reveals that hubs are enriched in essential genes
After hierarchically clustering the genes based on TOM dissimilarity and merging the
modules with 80% or more similarity, we generated a GCN containing 29 modules
(Figure 1; Supplementary Table S1). The largest and smallest modules were
chocolate3 (n = 6,250) and *steelblue3* (n = 30), respectively. The *grey* module, which
stores unassigned genes, had 4,766 genes. We selected the top hub from each
module and analyzed their functions in PhytoMine. Top hub genes tend to encode
proteins of paramount importance, such as succinate dehydrogenase
(Glyma.19G08600), an RNA polymerase subunit (Glyma.06G217400), DNA
topoisomerase (Glyma.14G137100), and cellulose synthase (Glyma.06G324300)
(Supplementary Table S2).
Jeong *et al* found that hubs tend to be essential in yeast (*S. cerevisiae*) protein-
protein interaction networks [42]. We hypothesized that degree centrality in a GCN
is also associated with gene essentiality in soybean. Essential genes in plants are
often referred to as embryonic lethal genes (i. e. genes that compromise embryo
viability when not expressed). *A. thaliana* essential genes have been diligently
catalogued over the years by Meinke's group, who recently published a curated list
of 510 *EMBRYO-DEFECTIVE (EMB)* genes [43], out of which 481 have soybean
orthologs (1,010 soybean genes). Importantly, the soybean orthologs of the essential
genes (Supplementary Table S3) are significantly overrepresented among GCN hubs
(Fisher's exact test; p = 5.8e-05). Further, essential genes that are hubs are enriched
in functions related to nucleic acids metabolism, translation and regulation of cell
division, such as replication factors, ribosomal proteins, and elongation factors.
Next, we investigated if any of our modules was individually enriched for
essential genes and essential hub genes. The modules *darkmagenta, yellow2,*
lightskyblue1, lightblue1, chocolate3, darkgoldenrod3, and *darkviolet* were
significantly enriched in essential genes (Fisher's exact test; p = 7.8e-05, 2.3e-19,
4.5e-02, 5.5e-09, 7.8e-06, 5.8e-09, and 6.0e-03, respectively). A functional
enrichment analysis revealed that, respectively, these modules are related to
photosynthesis and gluconeogenesis; translation and ribosome biogenesis; RNA
synthesis and processing; RNA metabolism; DNA replication; protein ubiquitination
and regulation of transcription and; *darkviolet* had the enriched protein domain
acetyl-CoA carboxylase alpha subunit. The latter three modules were also enriched
with essential hubs (Fisher's exact test; p = 9.6e-08, 1.3e-04, 4.8e-02, respectively).

These findings show that these gene clusters are essential to embryo viability and
could be experimentally validated in the future. Overall, our findings corroborate the
well-established role of hub genes as essential players of the cell physiology [44, 45].
195
Module hubs uncover biological processes associated with specific tissues
We selected the hub genes from each module and performed an enrichment analysis
for GO terms, pathways and protein domains (see methods for details). Information
on the number of hubs per module are in Supplementary Table S4. Only 14 of 29
modules (48.3%) had enriched GO terms, pathways or protein domains, and we
identified the function of nine of the 14 modules based on enriched biological
processes (GO-BP) and pathways (Supplementary Table S5). Further, we identified
module-tissue relationships by correlating each module eigengene to each of the 15
tissues available in our dataset (Figure 2; Supplementary Table S6), some of which
are discussed below.
Genes in the module *aliceblue* are related to lignin and phenylpropanoid
metabolism, such as laccases (Glyma.01G112600, Glyma.03G077900,
Glyma.08G359100), peroxidases (Glyma.01G163100, Glyma.11G049600,
Glyma.11G078400, Glyma.12G195500), xylem cysteine proteinase
(Glyma.04G014700, Glyma.06G014800), and cytochrome b561
(Glyma.02G306800). Module-tissue correlation showed that they are up-regulated in
leaf, root, and hypocotyl (r = 0.33, 0.31, 0.22; p = 1e-34, 3e-29, 1e-15, respectively)
and down-regulated in cotyledons, embryo and seeds (r = -0.21, -0.29, -0.33; p =
1e-14, 8e-26, 2e-34, respectively). Lignins are polymers made up of monolignols,
which are polymerized into lignin by laccases and peroxidases [46]. Because of their
strong covalent bonds to cellulose in secondary cell walls, lignins provide mechanical
support to plant organs such as stems and leaves, which explains the up-regulation
of *aliceblue* genes in this tissue [47]. In addition, lignins are difficult for herbivores
to digest, constituting a defense mechanism. Lignin is also well known for its role in
the stability of vascular tissues such as xylem [48]. The xylem cysteine proteinase
encoded by Glyma.04G014700, also in *aliceblue*, is likely involved in the autolysis of
tracheary elements during xylogenesis in roots [49]. Lignin is primarily synthesized
in hypocotyls during seedling establishment, as shown in growing hemp hypocotyls
[50]. Since lignin biosynthesis begins in the hypocotyl at the seedling stage, such
genes are also expected to be down-regulated in embryo, seeds and cotyledons,
which are mostly represented by developing seeds in our dataset. Thus, our results
recapitulate some of the physiological and anatomical roles of lignin.
Hub genes in the module *antiquewhite4* encode proteins related to the
tricarboxylic acid (TCA) cycle and aerobic respiration, such as isocitrate
dehydrogenase (Glyma.13G144900, Glyma.10G058100, Glyma.16G118900) and
ureide permeases (Glyma.01G058500 and Glyma.02G116300). These genes are up-
regulated in leaf (r = 0.53; p = 5e-93) and down-regulated in seed, embryo and
cotyledon (r = -0.43, -0.43, -0.2; p = 7e-60, 6e-60, 2e-13, respectively). As most

seed-related tissues in our dataset represent developing seeds, these genes are
expected to be down-regulated, since seed reserves are accumulated during
development. Storage proteins and other nutrients are then mobilized during
germination to promote early seedling development [51]. The up-regulation of TCA
cycle-related genes in leaf is likely because the TCA cycle is one of the major sources
of redox equivalents for oxidative phosphorylation, which provides ATP for sucrose
synthesis [52].

In *blue4,* we found genes related to transcriptional regulation and signaling,
such as ABC transporters (Glyma.13G361900, Glyma.07G015800), ferrochelatase
(Glyma.04G050400), phosphatases (Glyma.06G290200, Glyma.08G071900),
kinases (Glyma.04G088800, Glyma.06G090800, Glyma.13G073900,
Glyma.14G001300, Glyma.02G311200), TFs (Glyma.03G127600,
Glyma.02G232600, Glyma.07G023300, Glyma.14G200200, Glyma.09G005700,
Glyma.12G212300, Glyma.13G289400), and adenylate kinase (Glyma.10G048200).
Blue4 genes were slightly up-regulated in leaf and root (r = 0.21 and 0.24; p = 6e-
14, 5e-18, respectively) and down-regulated in embryo (r = -0.34; p = 6e-36). We
analyzed the annotation of *blue4* orthologs in *A. thaliana* and found that this module
is likely related to defense against pathogens, especially fungi. Among these
orthologs, we found the ABC transporters PEN3 and PIS1 (AT1G59870 and
AT3G53480), which participate in salicylic acid-dependent resistance against direct
pathogen penetration and in secretion of phenolic compounds involved in Fe uptake
[53]. Absorbed Fe ions can then be used by the ferrochelatase FC1 (AT5G26030) to
supply heme for defensive hemoproteins [54]. Similarly, the TFs WRKY33 and TGA6
from this module are involved in the activation of salicylic acid-responsive genes and
pathways mediating response to fungal pathogens, respectively [54, 55]. Innate
immune responses rely on the detection of microbe-associated molecular patterns
(MAMPs) by pattern recognition receptors (PRRs). Most *A. thaliana* PRRs are leucine-
rich repeat (LRR)-receptor kinases [56, 57]. *Blue4* also harbor orthologs of *A. thaliana*
innate immunity genes, such as an LRR-receptor kinase (AT5G01950) and an
interleukin-1 receptor-associated kinase (AT1G14370). We also found an ortholog of
the receptor kinase CERK1 (AT3G21630), which is involved in response to
chitooligosaccharides, a very common plant MAMP [56, 58]. Hence, the up-regulation
of these genes in leaf and root is coherent, whereas their down-regulation in embryo
is likely because of its physical protection by other organs and tissues [59].

Hubs from *darkmagenta* encode proteins related to photosynthesis and
gluconeogenesis, such as phosphoribulokinase (Glyma.01G010200), glycine
transaminase (Glyma.01G026700), photosystem II enhancer protein 1
(Glyma.02G061100), NADPH-dependent thioredoxin reductase 3
(Glyma.02G185300), fructose-bisphosphate aldolase (Glyma.04G008300),
glyceraldehyde-3-phosphate dehydrogenase (Glyma.04G015900), and fructose-1,6-
biphosphatase (Glyma.07G142700). These genes are remarkably up-regulated in leaf
(r = 0.8; p = 7e-292) and down-regulated in embryo, root, seed, and seed coat (r =

-0.28, -0.29, -0.34, and -0.28; p = 4e-25, 2e-26, 4e-36, 2e-25, respectively). This
is expected because in leaves, photosynthesis is responsible for CO_2 fixation in triose
phosphate, which feeds sucrose synthesis [60]. Hence, our findings support a strong
transcriptional co-regulation of sucrose synthesis and photosynthesis genes. In root
and seed-related parts, however, the down-regulation of these genes is due to their
very low photosynthetic potential as compared with other plant parts, making them
predominantly heterotrophic [61-64].

Finally, hubs from *orange2* are enriched in genes involved in cytochrome C
biogenesis and assembly (e.g. Glyma.06G254100, Glyma.12G147300). This module
is up-regulated in leaf (r = 0.59; p = 2e-119) and down-regulated in seed-related
parts (embryo and seed coat, r = -0.35, -0.32; p = 1e-37, 6e-32, respectively). The
Arabidopsis ortholog (AT5G54290) of these genes encodes CcdA, a thylakoid
membrane cytochrome C biogenesis protein that transfers reducing equivalents from
stromal thioredoxins to the thylakoid lumen [65]. Many proteins located in the
thylakoid lumen depend on redox regulation, and it has been shown that the
Arabidopsis ccda mutant is incapable of assembling the cytochrome b_6f in thylakoids
[66].

Functional prediction of unannotated hubs
Guilt-by-association (GBA) approaches have been widely used for function inference
[67-70]. In the context of GCN, it assumes that genes with similar expression
patterns are co-regulated and more likely to share the same pathway or biological
process. We explored the GCN and found 106 unannotated (i.e. genes without brief
description in PhytoMine) hubs, of which 93 were distributed in the nine modules for
which we could assign some biological function, as described above. By using a GBA
approach, we predicted functions for all these 93 hubs (Supplementary Table S7),
which might unveil interesting players in soybean transcriptional regulation.

We also found that 28 out of 29 modules had annotated top hubs. The top hub
in the module *antiquewhite4* (Glyma.19G074200) encoded a protein of unknown
function. We performed a functional enrichment analysis of its direct neighbors (r ≥
0.7) and found that they are involved in response to auxin (p = 1.19e-3), which is
likely the function of this unannotated gene. These genes encode auxin-responsive
GH3 family proteins (Glyma.03G256200 and Glyma.19G258800), and SAUR-like
auxin-responsive   proteins   (Glyma.04G231100,   Glyma.06G134000,
Glyma.08G004100, Glyma.19G254000).

The soybean genome remains far from being completely annotated. While
80.71% of soybean genes have some, although often minimal, functional annotation
in PhytoMine's brief description, the remaining 19.29% encode proteins of unknown
function. Network-based approaches like the one reported here might help us
understand the functions of many interesting genes, in particular those potentially
associated with agronomic traits. As hub genes are typically major players of the cell
physiology, our results can be relevant for the soybean genomics community.

Enrichment of transcription factor families uncovers major regulators of important biological processes

319
320
We applied a Fisher's exact test and found that 10 of the 29 modules are enriched in
TFs (Supplementary Table S8), suggesting that the functions involved in these
modules require a more complex transcriptional regulatory architecture. We also
found 6 modules enriched in specific TF families (FDR ≤ 0.05; Table 1). We
hypothesize that the overrepresented TF families in each module have key regulators
of the biological processes enriched in the module (Table 1). We also performed a *de*
novo motif discovery in promoter regions of genes in each module with enriched TF
families (Supplementary Figure S3), although none of the identified motifs matched
those from the same family in JASPAR and other Arabidopsis databases available in
Tomtom (MEME suite) [71]. Because of their regulatory roles, TFs are often reported
as important targets for developing transgenic lines with increased efficiency in
several biological processes and metabolic pathways [72-74]. Hence, the genes
reported in Table 1 could be interesting targets for plant transformation studies.

Network topology and the possible fate of soybean duplicated genes

335
We sought to explore the constructed GCN to explore the fates of duplicate genes.
All soybean duplicated gene pairs were downloaded from Plant Duplicate Gene
Database [41]. Duplicated gene pairs were classified in tandem, proximal,
transposed, dispersed and whole-genome duplication (TD, PD, TRD, DD and WGD,
respectively). Based on the Ks distribution, WGD-derived duplicates were subdivided
into WGD 58 mya and WGD 13 mya, representing WGD events that happened 58 and
13 million years ago, respectively (Supplementary Figure S4). The co-occurrence of
duplicates in modules can help understand their fates after duplication, as each
module has a unique expression profile (Supplementary Figure S5). In 27.2% of the
gene pairs (22,003/80,946), both genes were excluded during data filtering (see
methods for details) and are thus not present in the network, while in 27.6%
(22,355/80,946) only one gene of the pair was present in the network. Finally, both
genes were present in the network for 45.2% of the pairs (36,589/80,946).
Expectedly, these three categories have distinct gene expression levels
(Supplementary Figure S6).
We analyzed the co-occurrence of duplicated genes and the prevalence of their
modes of duplication in modules. Duplicate pairs in which both genes were absent in
the network were not considered. The co-occurrence of duplicate pairs in the same
module was interpreted as an indicative of functional conservation, as they reflect
similar expression profiles after duplication. On the other hand, when duplicate pairs
were in different modules or when only one gene from the pair was present, we
interpreted it as an indicative of functional divergence. In order to assess the
significance of these results, we created 10,000 degree-preserving simulated
networks (Figure 3; see methods for details). We observed that most pairs displayed

divergent expression profiles or signs of fractionation (Figure 3). Strikingly, WGD
pairs, particularly those from the most recent WGD, showed greater levels of
retention when compared to other modes of gene duplication, both in the same and
in different modules. These observations are possibly correlated with previous reports
on the increased retention of WGD duplicates involved in intricate systems (e.g.
regulatory genes) and protein complexes [23].
As previously shown, subfunctionalization is by far the most common fate of
retained duplicated genes [75], although it is not totally clear how this process
happens at the transcriptional level. Our findings suggest that the most common fate
of duplicated genes is functional divergence, as 59,4% to 91.7% of the duplicates do
not belong to the same module. Nevertheless, the frequency of genes in the same
module is higher than the expected by chance, supporting that the transcriptional
similarity of part of the duplicates is under selective pressure. This finding provides
new insights into how retained copies are regulated throughout the soybean genome.
Finally, we hypothesized that paralog pairs with similar expression profiles (i.e
same module) are under stronger purifying selection as compared to paralog pairs
with different expression profiles (i.e. members of the pair in different modules, both
in *grey* module or only one gene from the pair present in the network). We observed
that, except for proximal and tandem duplication-derived pairs, paralog pairs with
similar expression profiles have significantly lower Ka/Ks ratios as compared to those
with different expression profiles (Wilcoxon test; $p < 0.05$; Figure 3C). This result
suggests that duplicates that are retained in the same module are under stronger
negative selection, both at transcriptional and sequence levels. These pairs could be
involved in systems that rely on tightly regulated stoichiometric balance of its
components.
385
CONCLUSION
Here, we employed a thorough GCN analysis to explore the systems-wide
transcriptional landscape of soybean tissues. We explored the functions of several
modules and hubs using enrichment and GBA methods. Several of the identified
modules are strongly associated with specific tissues, giving important clues about
their physiological roles. We also found modules enriched in TFs, which probably
correspond to systems dependent on more intricate transcriptional regulation
architectures. Finally, we used our GCN to explore evolution of duplicate genes in
soybean and found that most of the retained duplicated genes tend to diverge in
expression, although genes from WGD are more often found in the same modules
than those duplicated by other mechanisms. The GCN reported here provides insights
into the roles of several soybean genes and co-expression modules, which might be
important not only in the elucidation of gene functions, but also in exploring genes
involved in relevant agronomic traits.

AUTHOR CONTRIBUTIONS

400 Conceived the study: FA-S, KCM and TMV. Data analysis: FA-S, KCM and FBM.
401 Funding, project coordination and infrastructure: TMV. Manuscript writing: FA-S and
402 TMV.

ACKNOWLEDGEMENTS

403 This work was supported by funding from Coordenação de Aperfeiçoamento de
404 Pessoal de Nível Superior (CAPES, Finance code 001), Conselho Nacional de
405 Desenvolvimento Científico e Tecnológico (CNPq), and Fundação Carlos Chagas Filho
406 de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ). The authors declare no
407 conflict of interest.

REFERENCES

1. Gill, N., et al., *Molecular and chromosomal evidence for allopolyploidy in soybean.*
Plant Physiol, 2009. **151**(3): p. 1167-74.
2. Schmutz, J., et al., *Genome sequence of the palaeopolyploid soybean.* Nature,
2010. **463**(7278): p. 178-83.
3. Bellieny-Rabelo, D., et al., *Transcriptome analysis uncovers key regulatory and*
metabolic aspects of soybean embryonic axes during germination. Sci Rep, 2016.
6: p. 36009.
4. Collakova, E., et al., *Metabolic and Transcriptional Reprogramming in Developing*
Soybean (Glycine max) Embryos. Metabolites, 2013. **3**(2): p. 347-72.
5. Gazara, R.K., et al., *Transcriptional landscape of soybean (Glycine max)*
embryonic axes during germination in the presence of paclobutrazol, a gibberellin
biosynthesis inhibitor. Sci Rep, 2019. **9**(1): p. 9601.
6. Severin, A.J., et al., *RNA-Seq Atlas of Glycine max: a guide to the soybean*
transcriptome. BMC Plant Biol, 2010. **10**: p. 160.
7. Huang, J., et al., *Construction and Optimization of a Large Gene Coexpression*
Network in Maize Using RNA-Seq Data. Plant Physiol, 2017. **175**(1): p. 568-583.
8. Jiang, Z., et al., *Differential Coexpression Analysis Reveals Extensive Rewiring of*
Arabidopsis Gene Coexpression in Response to Pseudomonas syringae Infection.
Sci Rep, 2016. **6**: p. 35064.
9. Ozaki, S., et al., *Coexpression analysis of tomato genes and experimental*
verification of coordinated expression of genes found in a functionally enriched
coexpression module. DNA Res, 2010. **17**(2): p. 105-16.
10. Pereira Lima, J.J., et al., *Molecular characterization of the acquisition of longevity*
during seed maturation in soybean. PLoS One, 2017. **12**(7): p. e0180282.

11. Gamboa-Tuz, S.D., et al., *Transcriptomics and co-expression networks reveal tissue-specific responses and regulatory hubs under mild and severe drought in papaya (Carica papaya L.).* Sci Rep, 2018. **8**(1): p. 14539.

12. Tan, M., et al., *Co-expression network analysis of the transcriptomes of rice roots exposed to various cadmium stresses reveals universal cadmium-responsive genes.* BMC Plant Biol, 2017. **17**(1): p. 194.

13. You, Q., et al., *Co-expression network analyses identify functional modules associated with development and stress response in Gossypium arboreum.* Sci Rep, 2016. **6**: p. 38436.

14. Wu, Z., et al., *A global coexpression network of soybean genes gives insights into the evolution of nodulation in nonlegumes and legumes.* New Phytol, 2019. **223**(4): p. 2104-2119.

15. Jaksik, R., et al., *Microarray experiments and factors which affect their reliability.* Biol Direct, 2015. **10**: p. 46.

16. Brum Machado, F., et al., *Systematic analysis of 1,298 RNA-Seq samples and construction of a comprehensive soybean (Glycine max) expression atlas.* Plant J, 2020.

17. Hansen, B.O., et al., *Elucidating gene function and function evolution through comparison of co-expression networks of plants.* Front Plant Sci, 2014. **5**: p. 394.

18. Romero-Campero, F.J., et al., *A contribution to the study of plant development evolution based on gene co-expression networks.* Front Plant Sci, 2013. **4**: p. 291.

19. Roulin, A., et al., *The fate of duplicated genes in a polyploid plant genome.* Plant J, 2013. **73**(1): p. 143-53.

20. Rensing, S.A., *Gene duplication as a driver of plant morphogenetic evolution.* Curr Opin Plant Biol, 2014. **17**: p. 43-8.

21. Birchler, J.A., et al., *Dosage balance in gene regulation: biological implications.* Trends Genet, 2005. **21**(4): p. 219-26.

22. Papp, B., C. Pal, and L.D. Hurst, *Dosage sensitivity and the evolution of gene families in yeast.* Nature, 2003. **424**(6945): p. 194-7.

23. Moharana, K.C. and T.M. Venancio, *Polyploidization events shaped the transcription factor repertoires in legumes (Fabaceae).* Plant J, 2020.

24. Teufel, A.I., et al., *The Many Nuanced Evolutionary Consequences of Duplicated Genes.* Mol Biol Evol, 2019. **36**(2): p. 304-314.

25. Panchy, N., M. Lehti-Shiu, and S.H. Shiu, *Evolution of Gene Duplication in Plants.* Plant Physiol, 2016. **171**(4): p. 2294-316.

26. Braasch, I., et al., *Reply to: 'Subfunctionalization versus neofunctionalization after whole-genome duplication'.* Nat Genet, 2018. **50**(7): p. 910-911.

27. Langfelder, P. and S. Horvath, *WGCNA: an R package for weighted correlation network analysis.* BMC Bioinformatics, 2008. **9**: p. 559.

28. Koh, F.M., et al., *Emergence of hematopoietic stem and progenitor cells involves a Chd1-dependent increase in total nascent transcription.* Proc Natl Acad Sci U S A, 2015. **112**(14): p. E1734-43.

29. Levine, M.E., P. Langfelder, and S. Horvath, *A Weighted SNP Correlation Network Method for Estimating Polygenic Risk Scores.* Methods Mol Biol, 2017. **1613**: p. 277-290.

30. Patel, H., R.J.B. Dobson, and S.J. Newhouse, *A Meta-Analysis of Alzheimer's*
Disease Brain Transcriptomic Data. J Alzheimers Dis, 2019. **68**(4): p. 1635-1656.
31. Csardi, G. and T. Nepusz, *The igraph software package for complex network*
research. InterJournal, complex systems, 2006. **1695**(5): p. 1-9.
32. Bi, D., et al., *Gene expression patterns combined with network analysis identify*
hub genes associated with bladder cancer. Comput Biol Chem, 2015. **56**: p. 71-
83.
33. Borin, G.P., et al., *Gene Co-expression Network Reveals Potential New Genes*
Related to Sugarcane Bagasse Degradation in Trichoderma reesei RUT-30. Front
Bioeng Biotechnol, 2018. **6**: p. 151.
34. Mack, K.L., et al., *Gene Expression Networks Across Multiple Tissues Are*
Associated with Rates of Molecular Evolution in Wild House Mice. Genes (Basel),
2019. **10**(3).
35. Supek, F., et al., *REVIGO summarizes and visualizes long lists of gene ontology*
terms. PLoS One, 2011. **6**(7): p. e21800.
36. Jin, J., et al., *PlantTFDB 4.0: toward a central hub for transcription factors and*
regulatory interactions in plants. Nucleic Acids Res, 2017. **45**(D1): p. D1040-
D1045.
37. de Matos Simoes, R. and F. Emmert-Streib, *Bagging statistical network inference*
from large-scale gene expression data. PLoS One, 2012. **7**(3): p. e33624.
38. Yu, C.P., J.J. Lin, and W.H. Li, *Positional distribution of transcription factor binding*
sites in Arabidopsis thaliana. Sci Rep, 2016. **6**: p. 25164.
39. Quinlan, A.R. and I.M. Hall, *BEDTools: a flexible suite of utilities for comparing*
genomic features. Bioinformatics, 2010. **26**(6): p. 841-2.
40. Bailey, T.L., et al., *MEME SUITE: tools for motif discovery and searching.* Nucleic
Acids Res, 2009. **37**(Web Server issue): p. W202-8.
41. Qiao, X., et al., *Gene duplication and evolution in recurring polyploidization-*
diploidization cycles in plants. Genome Biology, 2019. **20**.
42. Jeong, H., et al., *Lethality and centrality in protein networks.* Nature, 2001.
411(6833): p. 41-2.
43. Meinke, D.W., *Genome-wide identification of EMBRYO-DEFECTIVE (EMB) genes*
required for growth and development in Arabidopsis. New Phytol, 2019.
44. Han, J.D., et al., *Evidence for dynamically organized modularity in the yeast*
protein-protein interaction network. Nature, 2004. **430**(6995): p. 88-93.
45. Luscombe, N.M., et al., *Genomic analysis of regulatory network dynamics reveals*
large topological changes. Nature, 2004. **431**(7006): p. 308-12.
46. Heldt, H.-W., *Plant Biochemistry.* 2011.
47. Liu, Q., L. Luo, and L. Zheng, *Lignins: Biosynthesis and Biological Functions in*
Plants. Int J Mol Sci, 2018. **19**(2).
48. Bhuiyan, N.H., et al., *Gene expression profiling and silencing reveal that*
monolignol biosynthesis plays a critical role in penetration defence in wheat
against powdery mildew invasion. J Exp Bot, 2009. **60**(2): p. 509-21.
49. Zhao, C., et al., *XND1, a member of the NAC domain family in Arabidopsis*
thaliana, negatively regulates lignocellulose synthesis and programmed cell death
in xylem. Plant J, 2008. **53**(3): p. 425-36.

50. Behr, M., et al., *Insights into the molecular regulation of monolignol-derived product biosynthesis in the growing hemp hypocotyl.* BMC Plant Biol, 2018. **18**(1): p. 1.

51. Rajjou, L., et al., *Seed germination and vigor.* Annu Rev Plant Biol, 2012. **63**: p. 507-33.

52. Kromer, S., *Respiration during Photosynthesis.* Annual Review of Plant Physiology and Plant Molecular Biology, 1995. **46**: p. 45-70.

53. Marmagne, A., et al., *A high content in lipid-modified peripheral proteins and integral receptor kinases features in the Arabidopsis plasma membrane proteome.* Molecular & Cellular Proteomics, 2007. **6**(11): p. 1980-1996.

54. Nagai, S., et al., *Induction of isoforms of tetrapyrrole biosynthetic enzymes, AtHEMA2 and AtFC1, under stress conditions and their physiological functions in arabidopsis.* Plant Physiology, 2007. **144**(2): p. 1039-1051.

55. Libault, M., et al., *Identification of 118 Arabidopsis transcription factor and 30 ubiquitin-ligase genes responding to chitin, a plant-defense elicitor.* Molecular Plant-Microbe Interactions, 2007. **20**(8): p. 900-911.

56. Belkhadir, Y., et al., *Brassinosteroids modulate the efficiency of plant immune responses to microbe-associated molecular patterns.* Proceedings of the National Academy of Sciences of the United States of America, 2012. **109**(1): p. 297-302.

57. Ronald, P.C. and B. Beutler, *Plant and Animal Sensors of Conserved Microbial Signatures.* Science, 2010. **330**(6007): p. 1061-1064.

58. Osman, G.H., et al., *Development of insect resistant maize plants expressing a chitinase gene from the cotton leaf worm, Spodoptera littoralis.* Scientific Reports, 2015. **5**.

59. Raviv, B., et al., *The Dead Can Nurture: Novel Insights into the Function of Dead Organs Enclosing Embryos.* International Journal of Molecular Sciences, 2018. **19**(8).

60. Ward, J.M., et al., *Sucrose transport in higher plants.* International Review of Cytology - a Survey of Cell Biology, Vol 178, 1998. **178**: p. 41-71.

61. Adams, C.A. and R.W. Rinne, *Carbon-Dioxide Fixation in Developing Soybean Seeds.* Plant Physiology, 1979. **63**(5): p. 17-17.

62. Asokanthan, P.S., et al., *The photosynthetic potential of canola embryos.* Physiologia Plantarum, 1997. **101**(2): p. 353-360.

63. Harvey, D.M., C.L. Hedley, and R. Keely, *Photosynthetic and Respiratory Studies during Pod and Seed Development in Pisum-Sativum-L.* Annals of Botany, 1976. **40**(169): p. 993-1001.

64. Rolletschek, H., H. Weber, and L. Borisjuk, *Energy status and its control on embryogenesis of legumes. Embryo photosynthesis contributes to oxygen supply and is coupled to biosynthetic fluxes.* Plant Physiology, 2003. **132**(3): p. 1196-1206.

65. Motohashi, K. and T. Hisabori, *CcdA Is a Thylakoid Membrane Protein Required for the Transfer of Reducing Equivalents from Stroma to Thylakoid Lumen in the Higher Plant Chloroplast.* Antioxidants & Redox Signaling, 2010. **13**(8): p. 1169-1176.

566   66.   Page, M.L.D., et al., *A homolog of prokaryotic thiol disulfide transporter CcdA is*
567   *required for the assembly of the cytochrome b(6)f complex in Arabidopsis*
568   *chloroplasts.* Journal of Biological Chemistry, 2004. **279**(31): p. 32474-32482.
569   67.   Usadel, B., et al., *Co-expression tools for plant biology: opportunities for*
570   *hypothesis generation and caveats.* Plant Cell Environ, 2009. **32**(12): p. 1633-51.
571   68.   Shahan, R., et al., *Consensus Coexpression Network Analysis Identifies Key*
572   *Regulators of Flower and Fruit Development in Wild Strawberry.* Plant Physiol,
573   2018. **178**(1): p. 202-216.
574   69.   Petereit, J., et al., *petal: Co-expression network modelling in R.* BMC Syst Biol,
575   2016. **10 Suppl 2**: p. 51.
576   70.   De Smet, R., et al., *Coordinated Functional Divergence of Genes after Genome*
577   *Duplication in Arabidopsis thaliana.* Plant Cell, 2017. **29**(11): p. 2786-2800.
578   71.   Sandelin, A., et al., *JASPAR: an open-access database for eukaryotic transcription*
579   *factor binding profiles.* Nucleic Acids Res, 2004. **32**(Database issue): p. D91-4.
580   72.   Rabara, R.C., P. Tripathi, and P.J. Rushton, *The Potential of Transcription Factor-*
581   *Based Genetic Engineering in Improving Crop Tolerance to Drought.* Omics-a
582   Journal of Integrative Biology, 2014. **18**(10): p. 601-614.
583   73.   Sharma, V., et al., *An apple transcription factor, MdDREB76, confers salt and*
584   *drought tolerance in transgenic tobacco by activating the expression of stress-*
585   *responsive genes.* Plant Cell Reports, 2019. **38**(2): p. 221-241.
586   74.   Wang, N.N., et al., *The cotton WRKY transcription factor (GhWRKY33) reduces*
587   *transgenic Arabidopsis resistance to drought stress.* Scientific Reports, 2019. **9**.
588   75.   Freeling, M., *The evolutionary position of subfunctionalization, downgraded.*
589   Genome Dyn, 2008. **4**: p. 25-40.

Figures

Figure 1. Dendrogram and module colors before and after merging similar modules. In the WGCNA standard pipeline, each module is given a color. Modules were merged based on similarity of their eigengenes. The module eigengene is the first principal component of a principal component analysis, which summarizes the expression of the module. Modules with eigengenes more than 80% similar (r = 0.8) were merged.

Figure 2. Hubs from modules with enriched biological processes (GO) and pathways. For each subgraph, green squares represent hubs that are TFs. Below each subgraph, we illustrate tissues where the genes tend to be down-regulated (blue) and up-regulated (red). Up- and down-regulation in each tissue were obtained by calculating Pearson correlation coefficients between module eigengenes and a binary matrix m_{ij} containing 1 if the sample i corresponds to the tissue j, and 0 otherwise. P-values were derived by using the function corPvalueStudent() in WGCNA. Given the heterogeneity of the data, correlations were considered significant if $r \geq 0.2$ and $p < 0.05$. Ct, cotyledon; Em, embryo; S, seed; Hc, hypocotyl; Lf, leaf; Rt, root; SC, seed coat; En, endosperm.

Figure 3. Distribution and Ka/Ks analysis of duplicate pairs in the network. A. We observed that the minority of the duplicate pairs derived from all modes of duplication were in the same module in the GCN, supporting functional divergence as the most common fate of duplicated genes. B. We created 10,000 degree-preserving simulated networks by permuting the node labels of the GCN. Next, we compared the distribution of frequencies in which duplicate pairs were in the same module (histograms) with that observed in the real GCN (dashed lines). When both genes from a pair fell into the *grey* module, we treated them as being in different modules, as *grey* stores unassigned genes. C. Distribution of Ka/Ks ratios per mode of duplication and presence in the same module. Ka/Ks values were log transformed using log(Ka/Ks + 0.1) for better visualization. Except for proximal and tandem duplication-derived gene pairs, Ka/Ks ratios are significantly lower for paralog pairs within the same module, supporting that they are under stronger purifying selection at both transcriptional and sequence levels. DD, dispersed duplications; PD, proximal duplications; TD, tandem duplications; TRD, transposed duplications; WGD, whole-genome duplications; ns, no significant difference; ****, p<0.0001.

Tables

Table 1. Overrepresented TF families in soybean co-expression modules.
One-sided Fisher's exact test with Benjamini-Hochberg correction was used. TF
families were considered significantly overrepresented if FDR < 0.05.

Module name	TF family	FDR	Module function	Gene ID
aliceblue	WOX	8.83e-4	Lignin metabolism	Glyma.04G040900, Glyma.14G084600
antiquewhite4	TALE	1.27e-3	TCA cycle and aerobic respiration	Glyma.09G095700, Glyma.10G081700, Glyma.12G188800, Glyma.13G312900, Glyma.17G104800
blue4	WRKY	2.98e-6	Response to biotic stress	Glyma.02G232600, Glyma.07G023300, Glyma.08G218600, Glyma.09G005700, Glyma.12G212300, Glyma.13G289400, Glyma.14G200200, Glyma.15G003300, Glyma.15G110300
darkmagenta	CO-like	7.28e-3	Photosynthesis and gluconeogenesis	Glyma.05G233700, Glyma.07G091400
lightskyblue1	MYB-related	5.41e-4	mRNA synthesis and processing	Glyma.09G257700, Glyma.13G128500, Glyma.15G236400, Glyma.18G234900
yellow2	C2H2	1.04e-2	Translation and ribosome biogenesis	Glyma.11G189500, Glyma.12G084700, Glyma.12G181400