The 20-hydroxyecdysone agonist, halofenozide, primes anti-*Plasmodium* immunity in *Anopheles gambiae* via the ecdysone receptor

Rebekah A. Reynolds and Ryan C. Smith

Department of Entomology, Iowa State University, Ames, Iowa, USA

*Corresponding author: smithr@iastate.edu

Abstract

Mosquito physiology and immunity are integral determinants of malaria vector

competence. This includes the principal role of hormonal signaling in *Anopheles gambiae*

initiated shortly after blood-feeding, which stimulates immune priming and promotes

vitellogenesis through the function of 20-hydroxyecdysone (20E). Previous studies

demonstrated that manipulating 20E signaling through the direct injection of 20E or the

application of a 20E agonist can significantly impact *Plasmodium* infection outcomes,

reducing oocyst numbers and the potential for malaria transmission. In support of these

findings, we demonstrate that a 20E agonist, halofenozide, is able to prime anti-

Plasmodium immune responses that limit *Plasmodium* ookinete survival. We

demonstrate that halofenozide requires the function of ultraspiracle (USP), a component

of the canonical heterodimeric ecdysone receptor, to prime malaria parasite killing

responses. Additional experiments suggest that the effects of priming are temporal, such

that the application of halofenozide post-infection has no effect on malaria parasite

infection. Unlike 20E, halofenozide does not influence cellular immune function, AMP

production, or influence bacteria survival as in previous experiments. Together, our

results further demonstrate the potential of targeting 20E signaling pathways to reduce

malaria parasite infection in the mosquito vector and provide new insight into the

mechanisms of halofenozide-mediated immune priming that differ from 20E.

Introduction

Malaria kills over 400,000 people each year, with the majority of deaths in children under the age of five (World Health Organization, 2019). Transmission requires an *Anopheles* mosquito to acquire and transmit *Plasmodium* parasites through the act of blood-feeding, a behavior that evolved in mosquitoes to acquire nutrient-rich blood for egg development (Clements, 1992). Strategies to interrupt mosquito blood-feeding, and subsequent parasite transmission, are an essential step in reducing disease transmission. The use of long-lasting insecticide treated bed nets (LLITNs) and indoor residual spraying (IRS) were integral in reducing malaria deaths by approximately 50 percent over the last 20 years (Fernandes *et al*, 2018; Slutsker & Kachur, 2013; World Health Organization, 2019). These preventative measures have proven effective in Africa, where over 90 percent of the malaria cases occur globally, yet progress in the remaining regions of the world have plateaued in recent years (World Health Organization, 2019). Moreover, the increasing prevalence of insecticide and drug resistance now threaten a global resurgence in malaria cases (World Health Organization, 2019), therefore requiring new approaches to combat malaria transmission.

In order to be transmitted, malaria parasites must survive several bottlenecks in their mosquito host, where parasite numbers at the oocyst stage are arguably the lowest in the malaria life cycle (Smith *et al*, 2014). These bottlenecks are mediated in part by blood-meal derived factors, the mosquito microbiota, and components of the innate immune system that ultimately determine vector competence and vectorial capacity (Smith *et al*, 2014). Therefore, targeting components of mosquito physiology and/or immunity can serve as excellent targets for malaria control strategies to interrupt *Plasmodium* development in the mosquito host.

Previous studies have demonstrated that the steroid hormone 20-hydroxyecdysone (20E) is essential for multiple aspects of mosquito physiology; such as development, mating, reproduction, and immunity (Pondeville *et al*, 2008; Deitsch *et al*, 1995; Reynolds *et al*, 2020; Baldini *et al*, 2013). This includes the demonstration that 20E injection primes *An. gambiae* cellular immunity and reduces both *E. coli* and *P. berghei* survival in the mosquito host (Reynolds *et al*, 2020). 20E agonists, which were initially developed as

insecticides to reduce larval populations, have shown promise for use on adult *An.*
gambiae mosquitoes (Childs *et al*, 2016). Methoxyfenozide (DBH) reduced mosquito
lifespan, decreased fecundity, and significantly lowered the prevalence of *P. falciparum*
parasite infection in the mosquito host (Childs *et al*, 2016). However, the mechanisms by
which 20E agonists prime anti-*Plasmodium* immunity remain unexplored.

Here, we examine the 20E agonist, halofenozide, to better understand the mechanisms
of anti-*Plasmodium* immune priming, which are stimulated by the direct application to the
mosquito cuticle. In agreement with previous work on another 20E agonist (Childs *et al*,
2016), we demonstrate that halofenozide significantly reduces *P. berghei* parasite
numbers when applied to mosquitoes prior to infection. Furthermore, we demonstrate that
halofenozide priming influences the success of *Plasmodium* ookinete invasion and
requires the function of *ultraspiracle* (USP) as part of the heterodimeric ecdysone receptor
(USP/EcR). When compared to earlier studies of 20E immune priming (Reynolds *et al*,
2020), we find that halofenozide and 20E differentially influence mosquito cellular
immunity, suggesting that halofenozide acts as a nonsteroidal agonist to promote distinct
differences in cellular function (Okamoto *et al*, 2018). In summary, our study provides
additional detail into the mechanisms by which 20E agonists prime mosquito immunity
and limit *Plasmodium* parasite survival.

Results

Halofenozide application reduces *P. berghei* survival and infection prevalence

Previous studies have demonstrated the ability of 20E and the 20E agonist DBH to prime
mosquito responses that limit malaria parasite infection in *An. gambiae* (Reynolds *et al*,
2020; Childs *et al*, 2016). While recent evidence provides insight into the priming effects
of 20E (Reynolds *et al*, 2020), the mechanisms of priming by 20E agonists have not been
examined. To explore this question, we performed experiments with halofenozide, a
similar 20E agonist, and examined its influence on malaria parasite infection.
Halofenozide was topically applied to adult female mosquitoes without significant effect
on mosquito survival (Figure S1) and challenged with *P. berghei* ~24hrs post-application.
Compared to control (acetone-treated) mosquitoes, halofenozide treatment with either
0.25 µg/µL or 0.5 µg/µL significantly reduced the intensity of *P. berghei* oocysts eight days

post infection (Figure 1A). Moreover, halofenozide reduced the prevalence of infection

from 85% in control mosquitoes to <50% in halofenozide-treated mosquitoes (Figure 1B).

Given the low mortality rate and the strong anti-*Plasmodium* effect, the remainder of the

halofenozide experiments were conducted using the 0.5 µg/µL concentration.

Halofenozide priming promotes ookinete killing

To better determine how and when halofenozide limits parasite numbers, we examined

the effects of halofenozide priming on distinct stages of malaria parasite infection. When

early oocyst numbers were examined two days post-*P. berghei* infection, halofenozide-

treated mosquitoes displayed a significant reduction in early oocyst survival (Figure 2A),

arguing that halofenozide priming promotes ookinete killing. To further validate this point

and determine if halofenozide application also influenced other stages of parasite

development, halofenozide applications were performed approximately 24 hours post-

infection. Halofenozide treatment did not influence parasite survival when applied after

an established *P. berghei* infection (Figure 2B), suggesting that the effects of

halofenozide treatment are only effective against invading *Plasmodium* ookinetes when

applied prior to infection.

Halofenozide requires USP to stimulate immune priming

To confirm the known function of halofenozide acting through the canonical 20E receptor

(Retnakaran *et al*, 2003), we examined the expression of *vitellogenin* and *cathepsin B,*

genes responsive to 20E signaling (Upton *et al*, 2015; Roy *et al*, 2015). Halofenozide

application significantly increased the expression of *vitellogenin* (Figure 3A; Figure S2)

and *cathepsin B* (Figure 3B; Figure S2) to comparable levels as the injection of 20E

(Figures 3A and 3B), supporting that halofenozide activates canonical 20E signaling. To

determine if the heterodimeric ecdysone receptor comprised of the ecdysone receptor

(EcR) and ultraspiracle (USP) is critical for halofenozide priming of anti-*Plasmodium*

immunity, we used RNAi to examine the role of the respective components in

halofenozide priming (Figure S3). In RNAi experiments, the injection of dsRNA targeting

the heterodimeric receptor had no effect on *EcR*, yet significantly depleted levels of *USP*

(Figure S3). As a result, efforts to ascertain the function of the ecdysone receptor were

evaluated with *USP*-silencing. In control, *dsGFP*-silenced mosquitoes, halofenozide

application prior to *P. berghei* challenge resulted in a significant reduction in parasite
numbers (Figure 3C), similar to previous results (Figures 1 and 2). However, the topical
application of halofenozide in the *dsUSP*-silenced background (Figure S3) did not
influence *P. berghei* survival (Figure 3C). Together, this suggests that halofenozide
priming of anti-*Plasmodium* immunity requires the function of canonical 20E signaling
through the heterodimeric ecdysone receptor (EcR/USP).

Mechanisms of halofenozide priming are distinct from 20E priming
We previously examined the influence of 20E on cellular immunity and the effects of 20E
priming that limit bacteria and malaria parasite infection (Reynolds *et al*, 2020). Similarly,
we wanted to determine if halofenozide application promotes similar changes to 20E in
gene expression, cellular immunity, and anti-microbial defense. To compare gene
expression, we examined the expression of *cecropin 1* (*CEC 1*) and *cecropin 3* (*CEC 3*),
which previously were up-regulated in response to 20E injection in naïve mosquitoes
(Reynolds *et al*, 2020). However, halofenozide did not stimulate the expression of *CEC1*
or *CEC3* (Figure 4A and 4B), suggesting that halofenozide initiates a different
transcriptional repertoire than that of 20E (Reynolds *et al*, 2020). This led us to question
whether halofenozide application primed cellular immunity similar to 20E in naïve
mosquitoes (Reynolds *et al*, 2020). Contrary to previous work with 20E, halofenozide
treatment did not influence the phagocytic activity of mosquito immune cells (Figure 4C).
Moreover, halofenozide application did not reduce bacterial survival in *E. coli* challenge
experiments (Figure 4D). Together, these results suggest that the mechanisms of
halofenozide priming are distinct from that of 20E in limiting malaria parasites.

Discussion
To date, the most effective malaria control strategies target the mosquito host by reducing
mosquito habitats, preventing transmission through the use of bed nets, or by killing adult
mosquitoes through the use of insecticides (Bhatt *et al*, 2015). However, due to increasing
insecticide resistance in many malaria-endemic regions of the world, it is critical to
develop new methods or improve on existing tools to interrupt malaria transmission. This
includes the use of commonly used insecticides to manipulate mosquito host physiology,
influencing fitness or rendering them less likely to acquire and transmit mosquito-borne

pathogens. Childs *et al.* demonstrated that the application of a 20E agonist, DBH,
negatively influenced mosquito survival, reproduction, and reduced *P. falciparum* survival
in the mosquito host (Childs *et al*, 2016), supporting that 20E agonists have the potential
to improve existing malaria control strategies (Childs *et al*, 2016; Brown *et al*, 2020). This
is further supported by recent studies demonstrating that 20E confers anti-*Plasmodium*
immune priming (Reynolds *et al*, 2020), yet prior to our presented work herein, the manner
in which 20E agonists influence *Plasmodium* survival have remained unknown.

In agreement with previous work with DBH (Childs *et al*, 2016), we found the topical
application of halofenozide, a similar 20E agonist, significantly impaired *P. berghei* oocyst
survival and reduced infection prevalence. Additionally, we provide new details into the
temporal effects of halofenozide priming, demonstrating that the anti-*Plasmodium*
properties of halofenozide treatment only function to reduce *Plasmodium* numbers when
applied prior to malaria parasite challenge. Therefore, if 20E agonists were integrated into
bed nets or employed by indoor residual spraying (IRS) as previously proposed (Childs
et al, 2016; Brown *et al*, 2020), mosquitoes must to come into contact with 20E agonists
prior to taking an infectious blood meal to influence parasite survival. However, further
studies are required to determine if other 20E agonists, such as DBH, share similar
mechanisms of limiting malaria parasites.

The influence of halofenozide priming on early oocyst numbers suggests that
halofenozide acts on "early-phase" components of anti-*Plasmodium* immunity that
influence the success of ookinete invasion (Smith *et al*, 2014, 2015; Smith & Barillas-
Mury, 2016). This is further supported by our results demonstrating that halofenozide
application ~24h post-infection, and after ookinete invasion, did not influence malaria
parasite survival. As a result, halofenozide (and potentially other 20E agonists) may
influence the mosquito microbiota (Dong *et al*, 2009), complement function (Fraiture *et al*,
2009; Povelones *et al*, 2009; Oliveira *et al*, 2012), or a yet undescribed mechanism to
limit *Plasmodium* ookinete survival. However, further experiments are required to define
the direct mechanisms by which halofenozide promotes malaria parasite killing.

The integral role of USP in conferring the effects of halofenozide priming provide strong

support that halofenozide activates ecdysone signaling, which is in agreement with
previous evidence that 20E agonists competitively bind to the EcR/USP heterodimer
(Dhadialla *et al*, 1998). Although we were unable to knock-down *EcR*, EcR alone is
incapable of high affinity binding to 20E to promote the activation of downstream 20E-
regulated genes (Yao *et al*, 1993). This is further supported by the activation of *Vg* and
cathepsin B, two highly responsive 20E-regulated genes (Dhadialla & Raikhel, 1990; Roy
et al, 2015)*,* following halofenozide application at comparable levels to the injection of
20E. Therefore, our data support that halofenozide priming occurs through the
heterodimeric ecdysone receptor.

Previous work has identified the downstream targets of 20E signaling in *An. gambiae*, as
well as demonstrated the influence of 20E on cellular immunity and anti-pathogen
defense responses that limit bacterial and parasite survival (Reynolds *et al*, 2020). Since
halofenozide functions through the canonical 20E receptor, we originally hypothesized
that halofenozide and 20E would similarly influence mosquito physiology and immunity
(Reynolds *et al*, 2020). While both 20E and halofenozide stimulate *Vg* and *cathepsin B*,
halofenozide application does not stimulate immune gene expression, cellular immunity,
or anti-bacterial responses similar to 20E (Reynolds *et al*, 2020). Together, these results
suggest that halofenozide and 20E work through similar, yet distinct mechanisms to
influence mosquito physiology and prime anti-*Plasmodium* immunity. This may potentially
be explained by differences between steroid hormones such as 20E and their non-
steroidal agonist to enter target cells and stimulate cellular function (Nakagawa, 2005).
In support of this hypothesis, previous work in *Drosophila* demonstrates that 20E and
chromafenozide, a nonsteroidal 20E agonist, differentially enter cells to stimulate cellular
function (Okamoto *et al*, 2018). 20E requires the ecdysone importer (*EcI*) to gain cell
entry, while chromafenozide can stimulate cellular function independent of EcI (Okamoto
et al, 2018). Based on these notable differences between steroid hormones and non-
steroidal agonists, additional studies are required to further delineate the mechanisms of
20E and halofenozide priming on anti-*Plasmodium* immunity.

In summary, we demonstrate that the 20E agonist, halofenozide, is an effective tool to
prime physiological responses that reduce the prevalence and intensity of malaria

parasite infection, similar to previous studies with the 20E agonist DBH (Childs *et al*,

2016). Moreover, we define the temporal requirements of halofenozide application, where

anti-*Plasmodium* priming only occurs in mosquitoes treated pre-infection, consistent with

a role in halofenozide mediating physiological responses that limit the success of ookinete

invasion. We demonstrate that halofenozide priming requires the function of USP,

implicating ecdysone signaling in mediating the physiological responses that limit

Plasmodium infection. However, through comparative analysis of gene expression and

cellular immune function, we establish that halofenozide application produces

physiological responses not directly comparable to the effects of 20E (Reynolds *et al*,

2020). From these data, we provide new fundamental insight into the mechanisms of

halofenozide priming to better understand 20E agonist function. Together, the evidence

presented here and in previous studies (Childs *et al*, 2016; Brown *et al*, 2020) suggest

that 20E agonists could be an effective tool to help reduce the transmission of malaria.

Methods

Ethics statement

222   The protocols and procedures used in this study were approved by the Animal Care and

223   Use Committee at Iowa State University (IACUC-18-228).

Mosquito rearing and *Plasmodium* infections

An. gambiae (G3 strain) were maintained at 27°C and 80% relative humidity, with a 14/10-

hour light/dark cycle. Larvae were maintained on a ground fish food diet (Tetramin).

Pupae were isolated using a pupal separator (John W. Hock Company) and placed in

containers of ~50 mosquitoes where they were allowed to eclose in mixed populations of

male and female mosquitoes. Adult mosquitoes were maintained on a 10% sucrose

solution.

For mosquito infections with *P. berghei*, a mCherry *P. berghei* strain (Graewe *et al*, 2009)

was passaged into female Swiss Webster mice as previously described (Kwon *et al*,

2017; Smith *et al*, 2015; Kwon & Smith, 2019; Reynolds *et al*, 2020). After the confirmation

of an active infection by the presence of exflagellation, mice were anesthetized and

placed on mosquito cages for feeding. Infected mosquitoes were sorted and then

maintained at 19°C. Mosquito midguts were dissected eight days post-infection in 1x
PBS, and oocyst numbers were measured by fluorescence using a Nikon Eclipse 50i
microscope.

Halofenozide topical application on adult mosquitoes
For topical applications with halofenozide (Chem Services Inc.), a stock solution of
40µg/µL was prepared using 0.04 grams of halofenozide in 1 mL of 100% acetone. Prior
to application, working concentration were prepared by diluting the stock solution 1:79 or
1:159 in 100% acetone to achieve respective working concentrations of 0.25 µg/µL and
0.5µg/µL. Adult female *An. gambiae* (3-5 days post eclosion) were topically applied with
0.2µL of either 100% acetone (control) or halofenozide (0.25 µg/µL and 0.5µg/µL) in
100% acetone. Applications were performed on individual mosquitoes using a repeating
syringe dispenser (PB600-1 Hamilton syringe with a 10-microliter syringe). Surviving
mosquitoes were challenged (*E. coli* or *P. berghei*) 24 hours post-topical application or
were used to collect samples for RNA isolation to determine the effects of halofenozide
treatment on gene expression.

Gene expression analysis
Total RNA was isolated from whole mosquitoes (~10 mosquitoes) using TRIzol (Thermo
Fisher Scientific) according to the manufacturer's protocol. RNA samples were quantified
using a Nanodrop spectrophotometer (Thermo Fisher Scientific) and ~2ug of total RNA
was used as a template for cDNA synthesis using the RevertAid First Strand cDNA
Synthesis kit (Thermo Fisher). Gene expression was measured by qRT-PCR using gene-
specific primers (Table S1) and PowerSYBR Green (Invitrogen). Samples were run in
triplicate and normalized to an S7 reference gene and quantified using the $2^{-\Delta\Delta CT}$ method
as previously (Livak & Schmittgen, 2001).

dsRNA synthesis and gene-silencing
T7 primers for GFP and *ultraspiricle* (USP) were used to amplify cDNA prepared from
whole *An. gambiae* mosquitoes and cloned into a pJET1.2 vector using a CloneJET PCR
Cloning Kit (Thermo Fisher). The resulting plasmids were used as a template for
amplification using the corresponding T7 primers (Table S2). PCR products were purified

using the DNA Clean and Concentrator kit (Zymo Research) and used as a template for

dsRNA synthesis as previously described (Reynolds *et al*, 2020; Kwon *et al*, 2017). The

resulting dsRNA was resuspended in RNase-free water to a concentration of 3 µg/uL. For

gene-silencing experiments, 69nL of dsRNA was injected per mosquito and evaluated by

qRT-PCR to establish gene knockdowns at 1-5 days post-injection. Time points with the

highest efficiency of gene-silencing were chosen for downstream experiments.

Phagocytosis Assays Phagocytosis assays were performed *in vivo* as previously

described (Kwon & Smith, 2019; Reynolds *et al*, 2020) using red fluorescent FluoSpheres

(1 µm Molecular Probes) at a 1:10 dilution in 1x PBS. Following injection, mosquitoes

were allowed to recover for two hours at 19°C, then perfused onto a multi-test slide. To

visualize hemocytes, cells were stained using a 1:500 dilution of FITC-labeled wheat

germ agglutinin (WGA; Sigma) in 1x PBS, while nuclei were stained with ProLong Gold

anti-fade reagent with DAPI (Invitrogen). Hemocytes were identified by the presence of

WGA and DAPI signals, with the number of cells containing beads by the total number of

cells to calculate the percent phagocytosis. The phagocytic index was calculated by

counting the total number of beads per cell (this is summed for all of the cells) dividing by

the number of phagocytic cells. Approximately 200 cells were counted per mosquito

sample.

E. coli **challenge**

To evaluate the effect of halofenozide priming on bacterial infection, challenge

experiments were performed as previously (Benoit *et al*, 2017; Reynolds *et al*, 2020) with

control- or halofenozide-treated mosquitoes by injecting 69nL of *E. coli* into the thorax of

individual mosquitoes from each experimental condition. Following challenge,

mosquitoes were kept at 19°C for 18 to 24 hours before perfusion. Pooled hemolymph

from five mosquitoes (~50µL) was diluted in 450µL of room temperature LB broth, then

100µL was plated in triplicate on LB-kanamycin plates and incubated overnight at 37°C.

The number of *E. coli* colonies per plate were recorded, with the average used to compare

between control and halofenozide-treated samples (number of *E. coli* colonies/average

of control). This standardization was used to normalize for variation in *E. coli* numbers

between experiments.

Acknowledgments

We would like to acknowledge Hyeogsun Kwon for his helpful advice and laboratory assistance throughout this work, as well as David Hall for mosquito rearing. This work was supported by a National Science Foundation Graduate Research Fellowship Program under Grant No. 1744592 to R.A.R and by support from the Agricultural Experiment Station at Iowa State University to R.C.S.

References

Baldini F, Gabrieli P, South A, Valim C, Mancini F & Catteruccia F (2013) The interaction between a sexually transferred steroid hormone and a female protein regulates oogenesis in the malaria mosquito *Anopheles gambiae*. *PLoS Biol.* **11:** e1001695

Benoit JB, Vigneron A, Broderick NA, Wu Y, Sun JS, Carlson JR, Aksoy S & Weiss BL (2017) Symbiont-induced odorant binding proteins mediate insect host hematopoiesis. *Elife* **6:** 709–719

Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, Battle KE, Moyes CL, Henry A, Eckhoff PA, Wenger EA, Briët O, Penny MA, Smith TA, Bennett A, Yukich J, Eisele TP, Griffin JT, Fergus CA, Lynch M, et al (2015) The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature* **526:** 207–211

Brown F, Paton DG, Catteruccia F, Ranson H & Ingham VA (2020) A steroid hormone agonist reduces female fitness in insecticide-resistant *Anopheles* populations. *Insect Biochem. Mol. Biol.* **121:** 1033722

Childs LM, Cai FY, Kakani EG, Mitchell SN, Paton D, Gabrieli P, Buckee CO & Catteruccia F (2016) Disrupting mosquito reproduction and parasite development for malaria control. *PLoS Pathog.* **12:** e1006060

Clements A (1992) The biology of mosquitoes: Development, Nutrition and Reproduction.

Deitsch KW, Chen JS & Raikhel AS (1995) Indirect control of yolk protein genes by 20-hydroxyecdysone in the fat body of the mosquito, *Aedes aegypti*. *Insect Biochem. Mol. Biol.* **25:** 449–454

Dhadialla TS, Carlson GR & Le DP (1998) New Insecticides with ecdysteroidal and juvenile hormone activity. *Annu. Rev. Entomol.* **43:** 545–569

Dhadialla TS & Raikhel AS (1990) Biosynthesis of mosquito vitellogenin. *J. Biol. Chem.* **265:** 9924–9933

Dong Y, Manfredini F & Dimopoulos G (2009) Implication of the mosquito midgut microbiota in the defense against malaria parasites. *PLoS Pathog.* **5:** e1000423.

Fernandes JN, Moise IK, Maranto GL & Beier JC (2018) Revamping mosquito-borne disease control to tackle future threats. *Trends Parasitol.* **34:** 359–368

Fraiture M, Baxter RHG, Steinert S, Chelliah Y, Frolet C, Quispe-Tintaya W, Hoffmann J

a., Blandin SA & Levashina EA (2009) Two mosquito LRR proteins function as
complement control factors in the TEP1-mediated killing of *Plasmodium*. *Cell Host*
Microbe **5:** 273–284

Graewe S, Retzlaff S, Struck N, Janse CJ & Heussler VT (2009) Going live: A comparative
analysis of the suitability of the RFP derivatives RedStar, mCherry and tdTomato for
intravital and in vitro live imaging of *Plasmodium* parasites. *Biotechnol. J.* **4:** 895–
902

Kwon H, Arends BR & Smith RC (2017) Late-phase immune responses limiting oocyst
survival are independent of TEP1 function yet display strain specific differences in
Anopheles gambiae. *Parasit. Vectors* **10:** 369

Kwon H & Smith RC (2019) Chemical depletion of phagocytic immune cells in *Anopheles*
gambiae reveals dual roles of mosquito hemocytes in anti-*Plasmodium* immunity.
Proc. Natl. Acad. Sci. **116:** 14119–14128

Livak KJ & Schmittgen TD (2001) Analysis of relative gene expression data using real-
time quantitative PCR. *Methods* **25:** 402–408

Nakagawa Y (2005) Nonsteroidal ecdysone agonists. *Vitam. Horm.* **73:** 131–173

Okamoto N, Viswanatha R, Bittar R, Li Z, Haga-Yamanaka S, Perrimon N & Yamanaka
N (2018) A membrane transporter is required for steroid hormone uptake in
Drosophila. *Dev. Cell***:** 1–12

Oliveira GDA, Lieberman J & Barillas-Mury C (2012) Epithelial nitration by a
peroxidase/NOX5 system mediates mosquito antiplasmodial immunity. *Science.*
335: 856–859

Paton DG, Childs LM, Itoe MA, Holmdahl IE, Buckee CO & Catteruccia F (2019) Exposing
Anopheles mosquitoes to antimalarials blocks *Plasmodium* parasite transmission.
Nature 567:239-243

Pondeville E, Maria A, Jacques J-C, Bourgouin C & Dauphin-Villemant C (2008)
Anopheles gambiae males produce and transfer the vitellogenic steroid hormone 20-
hydroxyecdysone to females during mating. *Proc. Natl. Acad. Sci.* **105:** 19631–19636

Povelones M, Waterhouse RM, Kafatos FC & Christophides GK (2009) Leucine-rich
repeat protein complex activates mosquito complement in defense against
Plasmodium parasites. *Science* **324:** 258–261

Retnakaran A, Krell P, Feng Q & Arif B (2003) Ecdysone agonists: Mechanism and
importance in controlling insect pests of agriculture and forestry. *Arch. Insect*
Biochem. Physiol. **54:** 187–199

Reynolds RA, Kwon H & Smith RC (2020) 20-hydroxyecdysone primes innate immune
responses that limit bacterial and malarial parasite survival in *Anopheles gambiae*.
mSphere **5:** e00983-19

Roy S, Saha TT, Johnson L, Zhao B, Ha J, White KP, Girke T, Zou Z & Raikhel AS (2015)
Regulation of Gene Expression Patterns in Mosquito Reproduction. *PLoS Genet.* **11:**
e1005450

Slutsker L & Kachur SP (2013) It is time to rethink tactics in the fight against malaria.
Malar. J. **12:** 1–2

Smith RC & Barillas-Mury C (2016) *Plasmodium* oocysts: overlooked targets of mosquito
immunity. *Trends Parasitol.* **32**: 979-990.

Smith RC, Barillas-Mury C & Jacobs-Lorena M (2015) Hemocyte differentiation mediates
the mosquito late-phase immune response against *Plasmodium* in *Anopheles*
gambiae. Proc. Natl. Acad. Sci. **112:** E3412-20

Smith RC, Vega-Rodríguez J & Jacobs-Lorena M (2014) The *Plasmodium* bottleneck:
malaria parasite losses in the mosquito vector. *Mem. Inst. Oswaldo Cruz* **109:** 644–
661.

Upton LM, Povelones M & Christophides GK (2015) *Anopheles gambiae* blood feeding
initiates an anticipatory defense response to *Plasmodium berghei. J. Innate Immun.*
7: 74–86.

World Health Organization (2019) World Malaria Report 2019. Geneva.

Yao TP, Forman BM, Jiang Z, Cherbas L, Chen JD, McKeown M, Cherbas P & Evans
RM (1993) Functional ecdysone receptor is the product of EcR and Ultraspiracle
genes. *Nature* **366:** 476–479

Figures

Figure 1. Halofenozide priming reduces *P. berghei* survival and infection prevalence. (A) 100% acetone (control) or halofenozide dissolved in 100% acetone (0.25 µg/µL and 0.5 µg/µL) were applied to *Anopheles gambiae* adult females 24 hours prior to challenge with a *P. berghei*-infected blood meal. Eight days post-infection, oocyst numbers were examined from dissected midguts. The red bar delineates the median number of oocysts from pooled data from three independent experiments. Data were analyzed by Kruskal-Wallis with a Dunn's post-test using GraphPad Prism 6.0. n= the number of mosquitoes examined for each condition. **(B)** The prevalence of infection (% infected/total) is depicted for mosquitoes under each treatment and examined by X^2 analysis to determine significance.

Figure 2. Temporal effects of halofenozide application on *Plasmodium* infection. The effects of halofenozide application were more closely examined to determine the influence of pre- **(A)** and post-infection **(B)** application on malaria parasite infection. **(A)** After priming with halofenozide (0.5 µg/µL), *P. berghei* oocyst numbers were examined two days post-infection. The significant influence of halofenozide on early oocyst numbers suggests that priming limits the success of ookinete invasion. This is supported by experiments where halofenozide (0.5 µg/µL) was applied ~24 hours post-*P. berghei* infection **(B)** to assess whether halofenozide impacts parasite survival if applied at a later time point. No differences in oocyst numbers were detected when oocyst number were evaluated eight days post-infection. The red bar delineates the median number of oocysts from pooled data from three independent experiments. Data were analyzed by Mann-Whitney analysis using GraphPad Prism 6.0. n= the number of mosquitoes examined for each condition.

Figure 3. Halofenozide stimulates ecdysone signaling and requires ultraspiracle

(USP) to confer anti-*Plasmodium* immune priming. The effects of halofenozide (0.5

µg/µL), application on mosquito gene expression were evaluated on two downstream

components of ecdysone signaling, *vitellogenin* **(A)** and *cathepsin B* **(B)**, and compared

to 20E as a positive control. Statistical significance was determined by Mann-Whitney

analysis (**, $P < 0.01$; ns, not significant) from three or more independent biological

samples. **(C)** To determine the involvement of the heterodimeric EcR/USP receptor in

mediating the effects of halofenozide priming, we examined *USP* function by RNAi in *GFP*

(control)- or *USP*-silenced mosquitoes. Two days post dsRNA injection, we topically

applied acetone (-, control) or halofenozide (+, 0.5 µg/µL) and challenged mosquitoes

with *P. berghei* ~24 hours post-halofenozide topical application. Oocyst numbers were

evaluated eight days for parasite development. The red bar delineates the median

number of oocysts from pooled data from three independent experiments. Data were

analyzed by Mann-Whitney analysis using GraphPad Prism 6.0. n= the number of

mosquitoes examined for each condition.

Figure 4. Halofenozide does not influence AMP production or cellular immunity.
The effects of halofenozide topical application and 20E injection were compared by
examining their respective influence on gene expression using the 20E-responsive anti-
microbial peptides (AMPs), cecropin 1 (CEC1) **(A)** and cecropin 3 (CEC3) **(B)**. Statistical
significance was determined by Mann-Whitney analysis (*, $P<0.05$; **, $P< 0.01$) from
three or more independent biological samples. The influence of halofenozide was also
examined on the phagocytic activity of mosquito immune cells, evaluating the percentage
of phagocytic cells and the phagocytic index (number of beads per cell) **(C)**. Bacterial
challenge experiments were performed with *E. coli* to measure the potential immune
effects of halofenozide priming. For each experiment, halofenozide-treated mosquitoes
were challenged by intrathoracic infection ~24h after halofenozide application. Bacterial
titers were examined by hemolymph perfusion at either 18- or 24-hours post-*E. coli*
challenge **(D)**. Pooled hemolymph from five mosquitoes was plated in duplicate on LB-
kanamycin plates, and the bacterial colonies were counted to determine the titers. Data
were normalized from three or more independent experiments and visualized using box
plots (whiskers show minimum/maximum values, lines denotes median values). Data was

449   analyzed by Mann-Whitney analysis using GraphPad Prism 6.0 to determine significance.

450   ns, not significant.