Stress enhances hippocampal neuronal synchrony and prolongs sharp-wave ripples

Anupratap Tomar[1,2]*, Denis Polygalov [1], Sumantra Chattarji[3,4], Thomas McHugh[1]*.

[1] Laboratory for Circuit & Behavioral Physiology, RIKEN Center for Brain Science, 2-1 Hirosawa, Wako-shi, Saitama, Japan

[2] Current address: Center for Synaptic Plasticity, School of Physiology, Pharmacology and Neuroscience, University of Bristol, University Walk, Bristol BS8 1TD, UK

[3] National Centre for Biological Sciences, Bellary Road, Bangalore, 560065, India

[4] Centre for Discovery Brain Sciences, Deanery of Biomedical Sciences, University of Edinburgh, Hugh Robson Building, 15 George Square, Edinburgh, EH89XD, UK

* Correspondence: anupratap.tomar@bristol.ac.uk and thomas.mchugh@riken.jp

Keywords: stress, hippocampus, CA1, SPW-Rs, acute-stress, chronic-stress, pyramidal cells, ripples

Abstract

Chronic stress affects hippocampal function at multiple levels of neural organization. However,

much of this understanding is derived from postmortem analyses of molecular, morphological,

physiological and behavioral changes at fixed time points. Neural signatures of an ongoing

stressful experience in the intact brain of awake animals and their links to later hippocampal

dysfunction remain poorly understood. Here we used *in vivo* tetrode recordings to analyze the

dynamic impact of 10 days of immobilization stress on neuronal activity in area CA1 of mice.

Unexpectedly, there was a net decrease in pyramidal cell activity in stressed animals. Although

these results, suggest a lack of stress-induced hyperexcitability, more detailed analysis revealed

that a greater fraction of spikes occurred specifically during sharp-wave ripples, resulting in an

increase in neuronal synchrony. After repeated stress some of these alterations were visible

during rest even in the absence of stress. These findings offer new insights into stress-induced

alterations in ripple-spike interactions and mechanisms through which chronic stress may

interfere with subsequent information processing.

Introduction

The hippocampus is a medial temporal lobe structure that is crucial for encoding, updating and retrieving episodic memories (Eichenbaum, 2017; Tulving, 1985). Unfortunately, the same plasticity mechanisms that enable the hippocampus to perform these important functions, also make it vulnerable to damage caused by severe and repeated stress (Chattarji et al., 2015; McEwen et al., 2015). Stress-induced changes in the rodent hippocampus include shrinkage and debranching of pyramidal cell dendrites, (McEwen et al., 2015; Sousa et al., 2000), loss of dendritic spines (Magariños et al., 1997; Sandi et al., 2003) and alterations in synaptic plasticity mechanisms, including long-term potentiation (Alfarez et al., 2003; Shors et al., 1989). Together, these detrimental effects of stress at the cellular and synaptic levels are thought to contribute to impairments in hippocampal learning and memory. However, our current understanding is based primarily on postmortem analyses of stressed versus unstressed animals at fixed time points after the end of stress. The gradual and cumulative impact of stress on hippocampal function in the same animal over the course of repeated stress has not been explored in detail. Further, relatively little is known about how the intact, drug-free hippocampus is involved in the quick appraisal of an ongoing stressful situation (Cadle and Zoladz, 2015; Joëls, 2009).

These gaps in knowledge were addressed in several rodent studies that examined the effects of stress on theta-associated foraging behaviour (Kim et al., 2007; Park et al., 2015; Passecker et al., 2011; Tomar et al., 2015) during which hippocampal pyramidal 'place' cells (O'Keefe and Dostrovsky, 1971) form a cognitive map of the animals' surroundings (O'Keefe and Nadel, 1978). Surprisingly, earlier reports found only modest changes in place cell properties between stressed and control subjects. However, in addition to theta-associated exploratory states, pyramidal cells also exhibit highly coordinated activity during off-line behavioural states such as rest and sleep (Buzsáki, 1989). These offline states are dominated by high-frequency (100-200 Hz) transients termed sharp-wave ripples (SPW-Rs) which have been shown to play key roles in learning and memory (Ego-Stengel and Wilson, 2010; Girardeau et al., 2009; Jadhav et al., 2012) and the planning of future actions (Joo and Frank, 2018; Pfeiffer and Foster, 2013; Schmidt and Redish, 2013). However, no information is available on the effects of either acute or chronic stress on SPW-R properties or their modulation of pyramidal cell activity. Further, if and how SPW-Rs are affected during stress, and if such effects differ between acute versus chronic stress remains to characterized. Hence, the current study aimed to address these outstanding issues. To this end, we examined hippocampal neural dynamics in mice on the first

(acute) and last (chronic) day of a chronic immobilization stress (CIS) protocol. To characterize

a neural signature of stress in CA1 we recorded local field potentials (LFPs) and neuronal

activity during stress and compared it with activity recorded in an adjacent quiescence/rest

state.

Results

We examined the dynamics of CA1 pyramidal cell activity and ongoing LFPs on the first

(acute) and last day (chronic) of 10-day chronic immobilization stress (CIS) (Fig. 1a),

replicating a protocol that has been previously used to examine the effects of chronic stress on

hippocampal dendrites (Vyas et al., 2002), spatial coding (Tomar et al., 2015), and behaviour

(Ramirez et al., 2015; Suvrathan et al., 2010). Specifically, we examined the effects of stress

at two time-points during CIS. First, we compared activity during the first episode of 2-hour

stress (stress-state) to the preceding stress-free period (rest-state) on Day 1; this is referred to

as the "acute" condition as this involves only a single exposure to stress. In other words, the

effects of acute stress on the first day of CIS were analysed by comparing two conditions: i)

acute-rest, and ii) acute-stress. Second, we carried out the same analyses on the last day of CIS,

when the same animal has already experienced 9 exposures of the same stressor. Thus, we

again compared activity during the 10^{th} episode of 2-hour stress (stress-state) to an adjacent

stress-free period (rest-state). This is referred to as the "chronic" condition as it involves

quantifying the cumulative effects of repeated stress over 10 days. Here again we compared

two more conditions: iii) chronic-rest, and iv) chronic-stress, and together data from these four

conditions will be presented in the following sections. The CIS protocol led to a gradual

decrease in body weight (acute, 28.82 ± 0.97 vs chronic, 26.65 ± 0.69, N = 4 mice, paired t-

test: t = 3.8411, p = 0.031) (supplementary Fig. 1a), confirming the efficacy of this chronic

stress paradigm as it is consistent with previous reports (Vyas et al., 2002). Lesions in the

stratum pyramidale confirmed that recordings were made from area CA1 (Fig. 1b).

CA1 pyramidal cell spiking differs between stress-state and rest-state

A long-held view on the detrimental effects of stress centers on the idea that severe and repeated

stress leads to hippocampal hyperactivity, which in extreme cases may cause excitotoxic

damage. For instance, both the glucocorticoid cascade (Sapolsky, 1996) and synaptic saturation

hypotheses of stress (Cadle and Zoladz, 2015; Diamond et al., 2004) suggest that enhanced

calcium and glutamate release during stress alter subsequent synaptic plasticity and mnemonic

114 processes. The implicit assumption underlying both these hypotheses is that hippocampal

115 neuronal networks undergo hyperexcitability during stress though no study has directly tested

116 this possibility in the intact brain *in vivo*. Surprisingly, we found hippocampal multiunit activity

117 to be suppressed during stress (Fig. 1c and supplementary Fig. 1b, c). Comparison of the

118 average firing rate of CA1 pyramidal cells between stress and rest across days revealed a

119 significant decrease during the stress-state (Fig 1d) with the difference being most pronounced

120 during acute stress on the first day (acute-rest, 0.80 ± 0.06 Hz vs acute-stress, 0.38 ± 0.03 Hz,

121 p = 1.32×10^{-8}, Tukey's HSD). Following repeated stress, the mean firing rate continued to be

122 low not only during the last session of chronic stress, but also during the adjacent rest period,

123 resulting in a significant reduction of firing rates was both the rest and stress periods on Day

124 10, indicative of carryover of the stress phenotype into non-stress states as well (acute-rest,

125 0.80 ± 0.06 Hz, vs chronic-rest, 0.59 ± 0.04 Hz, p = 0.017; chronic-rest, 0.59 ± 0.04, vs chronic-

126 stress, 0.48 ± 0.05, p = 0.425, Tukey's HSD). Similarly, we observed a significant lowering of

127 peak firing rates (Hz) (Fig. 1e**)** on both the first day (acute-rest, 7.77 ± 0.59 Hz, vs acute-stress,

128 3.37 ± 0.49 Hz, p = 7.41×10^{-7}, Tukey's HSD) and the last day (chronic-rest, 8.80 ± 0.69 Hz vs

129 chronic-stress, 4.31 ± 0.62 Hz, p = 2.53×10^{-6}, Tukey's HSD) of CIS. While stress did not alter

130 the length (ms) of bursting activity, the stress-state was associated with longer inter-burst-

131 intervals (Fig. 1f), indicating that time gaps (s) between burst activity were lengthened,

132 consistent with the overall decrease in activity (acute-rest, 33.29 ± 3.33 vs acute-stress, 78.69

133 ± 6.42, p = 2.2×10^{-9}; chronic-rest, 40.41 ± 3.68 vs chronic-stress, 65.33 ± 6.82, p = 5.35×10^{-3},

134 Tukey's HSD). Taken together, these data demonstrate that stress decreases the activity of CA1

135 pyramidal cells.

137 **LFP profile and SPW-R properties differ between the stress-state and rest-state**

138 Considering stress did not cause a straightforward increase in CA1 firing rates, we next focused

139 our attention on other measures of hippocampal neural dynamics. Hippocampal behavioral

140 states display characteristic patterns of LFP activity (Buzsáki, 1989) wherein rest and

141 immobility states are defined by a low ratio of LFP power in the theta (6-12 Hz) and delta (1-

142 4 Hz) bands, along with periodic high-frequency sharp-wave ripples (SPW-Rs, 150-200 Hz)

143 (Grosmark et al., 2012). Consistent with earlier reports, we observed a negative correlation

144 between the theta/delta ratio and SPW-R probability during the unstressed rest-state. But, this

145 relationship was disrupted during both acute (Fig. 1g, left; acute-rest, R = -0.56, p = 0.02, vs

acute-stress, R = -0.19, p = 0.47) and chronic stress (Fig. 1a, right; chronic-rest, R = -0.54, p =

0.029 vs chronic-stress, R = -0.17, p = 0.52), raising the possibility that CIS induces a unique

physiological state.

The above findings underscore the need to examine in greater detail the effects of stress on

SPW-Rs. Moreover, there is growing evidence for SPW-Rs playing an important role in

memory consolidation, which is affected by stress (Fernández-Ruiz et al., 2019; Nakashiba et

al., 2009). Hence, we next analyzed SPW-R properties in both the early and late stages of

chronic stress. While no differences were observed in the intrinsic oscillation frequency

(supplementary Fig. 2a), the average SPW-R duration was significantly longer during stress at

both acute and chronic time points (Fig. 2b; acute-rest, 104.94 ± 3.29 ms vs acute-stress, 137.05

± 4.65 ms, p= 3.86×10^{-5}; chronic-rest, 107.54 ± 2.44 ms vs chronic-stress, 130.35 ± 6.996 ms,

p = 0.0055, Tukey's HSD). The population distribution of SPW-R duration (Fig. 2c) clearly

showed a greater fraction of long-duration ripples during both acute and chronic stress (acute-

rest (median = 90.93) vs acute-stress (median = 122.27), $p < 2.22 \times 10^{-16}$; chronic-rest (median

= 92.16) vs chronic-stress (median = 105.68), $p < 2.22 \times 10^{-16}$, Kolmogorov-Smirnov test).

Further, the increase in SPW-Rs duration was characterized by a rapid onset and sustained

increase following stress initiation (Fig. 2d).

Recent work has suggested that longer SPW-Rs could result from the merging or concatenation

of single ripple events, which is reflected as an increase in power in the underlying gamma

oscillation (Oliva et al., 2018). To examine this possibility we next measured the low-gamma

power (17-40 Hz) during ripple events, but found no differences between stress and rest during

both the acute (Day 1) and chronic (Day 10) time points (Fig. 2e and f), suggesting absence of

excessive merging of SPW-Rs. We also examined if stress leads to increased ripple bursts, a

phenomenon ascribed to input from the entorhinal cortex (Yamamoto and Tonegawa, 2017).

The proportion of ripples occurring as singlets (Fig. 2g) was similar between rest and stress at

both the acute and chronic time points (singlets, acute-rest, 0.81 ± 0.02 vs acute-stress, $0.80 \pm$

0.02, p= 0.999; chronic-rest, 0.78 ± 0.02 vs chronic-stress, 0.78 ± 0.03, p= 1.0, Tukey's HSD)

again suggesting that the observed SPW-Rs were longer single events.

We next compared the normalized amplitudes (supplementary Fig. 2b) of SPW-Rs and found

them to be significantly larger during both acute and chronic stress sessions (acute-rest, $4.98 \pm$

0.21 vs acute-stress, 6.22 ± 0.30, p = 0.013; chronic-rest, 5.14 ± 0.20 vs chronic-stress, $6.50 \pm$

0.37, p = 0.006, Tukey's HSD). Once again, the increase in SPW-R amplitude showed a rapid

onset and sustained increase following the beginning of the 2-hour stress on both the first and
last days (supplementary Fig. 2c). These data demonstrated that stress induction quickly alters
basic SPW-R properties in hippocampal area CA1.

Stress alters excitability and participation of CA1 pyramidal cells during SPW-Rs

In addition to its hypothesized role as a biomarker of cognition, SPW-Rs have also been
described as a "built in" physiological sensor of hippocampal excitability because of network
wide activation of hippocampal neurons which accompanies their occurrence (Buzsáki, 2015).
Thus, next we searched for evidence of hyperexcitability during both acute and chronic stress
by comparing the pyramidal cell firing patterns during SPW-R events between rest and stress
states. In agreement with suppressed average firing rates reported above, multiunit activity was
muted during SPW-Rs across both acute and chronic stress sessions, though the magnitude of
difference was larger during chronic stress (Fig. 3a). Neither acute nor chronic stress altered
the relationship between CA1 pyramidal cell firing in SPW-Rs and their overall mean firing
rate (supplementary Fig. 3a: acute-rest, $R^2 = 0.56$, $p < 2.22 \times 10^{-16}$; acute-stress, $R^2 = 0.47$, $p <$
2.22×10^{-16}, likelihood ratio test, $p = 0.80$; chronic-rest, $R^2 = 0.44$, $p < 2.22 \times 10^{-16}$; chronic-
stress, $R^2 = 0.27$, $p < 2.22 \times 10^{-16}$, likelihood ratio test, $p = 0.105$). Further, similar to overall
activity, we observed a decrease in pyramidal cell firing rate (Hz) inside SPW-Rs during acute
stress (Fig 3b acute-rest, 2.20 ± 0.19 Hz vs acute-stress, 1.38 ± 0.12 Hz, $p = 1.78 \times 10^{-4}$, Tukey's
HSD, while following chronic stress, mean firing remained low both during rest and stress on
the last day (acute-rest, 2.20 ± 0.19 Hz vs chronic-rest, 1.64 ± 0.11 Hz, $p = 0.024$, chronic-rest,
1.64 ± 0.11 Hz vs chronic-stress, 1.43 ± 0.11 Hz, $p = 0.749$, Tukey's HSD). Further, the impact
of chronic stress on firing in SPW-Rs during the rest-state was also evident in the cumulative
distribution plots (supplementary Fig 3b), as the rest-state on the last day showed a leftward
shift (acute-rest vs chronic-rest, $p = 0.013$, Tukey's HSD). Firing outside of SPW-Rs changed
in a similar fashion and decreased during acute-stress and chronic-rest states (supplementary
Fig. 3c &d). Interestingly, despite this overall drop in activity, during stress-states pyramidal
cells discharged a much larger percentage of their total spikes inside of SPW-Rs (Fig. 3c). The
fraction of spikes inside SPW-Rs during stress almost doubled on the first day (acute-rest,
$14.49 \pm 0.85\%$ vs acute-stress, $26.19 \pm 1.21\%$ $p = 1.37 \times 10^{-12}$, Tukey's HSD), and remained
elevated on the last day (chronic-rest, $15.93 \pm 1.05\%$ vs chronic-stress, $25.0. \pm 1.45\%$, $p =$
3.62×10^{-7}, Tukey's HSD) of CIS.

We next assessed the effects of stress on the participation of pyramidal cells in SPW-Rs events.

In agreement with a previous reports (Fernández-Ruiz et al., 2019; O'Keefe and Nadel, 1978),

the baseline firing rate of pyramidal cells displayed a positive relationship with the cell's

participation probability in SPW-Rs during both acute and chronic stress, though slight but

significant differences were observed during acute stress (Fig. 3d: acute rest, $R^2 = 0.55$, p <

$2.22\text{x}10^{-16}$; acute-stress, $R^2 = 0.40$, $p < 2.22\text{x}10^{-16}$; likelihood ratio test, $p = 0.038$; chronic-rest,

$R^2 = 0.42$, $p < 2.22\text{x}10^{-16}$; chronic-stress, $R^2 = 0.21$, $p = 2.97\text{x}10^{-13}$, likelihood ratio test, $p =$

0.94). Further, the extent of pyramidal cell participation in SPW-Rs varied widely (Fig.

3e;(Grosmark et al., 2012; Ylinen et al., 1995). However, a comparison across sessions and

days revealed a significant effect of day (Fig. 3e); while pyramidal cell participation was

similar on the first day (acute-rest, $13.39 \pm 0.84\%$ vs acute-stress, $11.41 \pm 0.73\%$, $p = 0.204$,

Tukey's HSD), the chronic rest-state showed a significantly lower participation than that of

the acute rest-state (acute-rest, $13.39 \pm 0.84\%$ vs chronic-rest, $10.57 \pm 0.58\%$, $p = 0.027$,

Tukey's HSD), suggesting that chronic stress suppresses pyramidal cell participation in SPW-

Rs. No further decrease was observed between behaviour states on the last day (chronic-rest,

$10.57 \pm 0.60\%$ vs chronic-stress, $10.77 \pm 0.73\%$, $p = 0.997$, Tukey's HSD) of CIS. Cumulative

distribution plots further confirmed the impact of chronic stress on SPW-R participation during

the rest state (Fig. 3f: acute-rest vs chronic-rest, $p = 0.006$, Kolmogorov-Smirnov test). Overall,

these data demonstrated that despite suppressed firing rates, SPW-R-specific activation of CA1

pyramidal cells was enhanced during both acute and chronic stress. However, with progression

of the CIS protocol, pyramidal cells participated in fewer SPW-Rs during the rest period

adjacent to the stress exposure.

Co-firing of CA1 pyramidal cells is altered by stress

Hippocampal neuronal synchrony peaks during SPW-Rs (Buzsáki et al., 1992; Csicsvari et al.,

1999a; Wilson and McNaughton, 1994) and this has been suggested as a key mechanism in

hippocampal mnemonic function (Buzsáki and Mizuseki, 2014; Cheng and Frank, 2008). Our

finding that a greater fraction of pyramidal spiking occurs inside SPW-Rs, raised the possibility

of enhanced synchrony of pyramidal cell firing during stress. However, the overall decrease in

firing rates, and reduced SPW-R participation by pyramidal cells with repeated stress,

suggested the contrary. To examine this, we quantified the co-activation of pairs of pyramidal

cells exhibiting a positive correlation in their firing patterns in a given session and observed a

significant elevation during stress periods (Fig. 4a). Co-activity Z-scores were significantly

higher during acute (acute-rest, 1.228 ± 0.027 vs acute-stress, 1.495 ± 0.031, $p = 1.2\text{x}10^{-8}$,

Tukey's HSD) as well as chronic stress (chronic-rest, 1.388 ± 0.03, vs chronic-stress, $1.622 \pm$

0.043, $p = 3.8 \times 10^{-6}$, Tukey's HSD). Further, the rest-state on the last day showed significantly

larger co-activation values as compared to the first day (acute rest vs chronic rest, $p = 0.0016$,

Tukey's HSD). Finally, to compare the coordinated activity more directly, we calculated co-

activity scores of cell-pairs with positive values simultaneously recorded both rest and stress

states (Fig. 4b). This analysis revealed a positive relationship between cell pairs during both

rest and stress states on the first day (acute: $R^2 = 0.022$, $p = 0.033$), which grew stronger with

further episodes of stress over time (chronic: $R^2 = 0.019$, $p = 0.076$) and significantly differed

between days ($p = 1.28 \times 10^{-7}$, likelihood ratio test) suggesting that chronic stress led to

reorganization of network activity even during non-stress periods.

Discussion

Accumulating evidence has identified stress-induced changes in the hippocampus across

biological scales – from behavioral deficits to their neuronal, synaptic and molecular correlates.

However, how the intact hippocampal circuitry in an awake behaving animal responds to, and

encodes information during, a stressful situation is poorly understood (McEwen, 2007). Here,

we report a reduction in spiking in CA1 pyramidal neurons during the first (acute) exposure to

2-hour immobilization stress (Fig. 1d and e). However, multiple exposures to stress led to

suppressed spiking even during rest periods outside of the stress itself. During both acute and

chronic stress, theta/delta ratio remained low (Fig. 1g) and SPW-R events were of longer

duration compared to rest (Fig. 2b and c). Further, stress altered the firing of CA1 pyramidal

cells during SPW-Rs, leading to a greater fraction of spikes occurring inside SPW-Rs during

both acute and chronic stress (Fig. 3c). Finally, pyramidal cells displayed enhanced co-activity

which was also carried over to the rest-state on the final day of the 10-day chronic stress

paradigm (Fig. 4a and b).

Considering that stress leads to elevated levels of glutamate and calcium, a long-held view in

research on stress-induced plasticity is that stress leads to enhanced excitability in hippocampal

circuits (Gunn and Baram, 2017; Sapolsky et al., 1986). However, our data from *in vivo*

recordings revealed a robust decrease in firing rates of CA1 pyramidal cells during acute stress

(Fig 1d). This result is consistent with a previous study that found suppression of pyramidal

'place cell' activity in an immobilized rat even when it was moved through its place field on a

track (Foster et al., 1989) as well as other studies which reported suppressed place cell firing

rates after the termination of stress (Park et al., 2015; Passecker et al., 2011). We also explored
other measures of hippocampal activity by assessing pyramidal cell activity during SPW-Rs,
which are high-frequency transient oscillations, referred to as "built in" sensors of hippocampal
excitability (Buzsáki, 2015). Strikingly, this analysis revealed evidence in support of stress-
induced enhanced excitability as pyramidal cells discharged a greater fraction of spikes inside
SPW-Rs during stress states (Fig 3c). Together these results offer a new framework for
analyzing stress-induced hyperexcitability at the network level, manifested as SPW-R events
during stress.

Our analyses also revealed stress-induced enhancement in CA1 neuronal synchrony as
evidenced by increased pyramidal cell co-activation during SPW-Rs (Fig. 4). Such co-activity
during repeated exposures to stress is likely to cause CA1 pyramidal neurons to fire together,
thereby creating conditions at the synaptic level that are consistent with the synaptic saturation
hypothesis of stress (Diamond et al., 2004). This gives rise to questions that will require further
investigation, including if such synchronous activity would saturate synaptic mechanisms that
occlude further plasticity necessary for subsequent encoding? If so, this may offer insights into
earlier findings that chronic stress causes rigidity in hippocampal networks (Tomar et al.,
2015), thus leading to context generalization (Krugers et al., 1997) and spatial learning deficits
(Kim et al., 2007; Park et al., 2015). Interestingly, a recent rodent study demonstrated a causal
relationship between long duration SPW-Rs and learning (Fernández-Ruiz et al., 2019). In light
of this study, our findings of increased pyramidal cell synchrony, along with long-duration
SPW-Rs suggests that during stress, the hippocampus may engage in a learning state involved
in encoding aversive memories related to the stressful experience (Cadle and Zoladz, 2015;
Diamond et al., 2004, 2007; Schwabe et al., 2012).

The observation that altered SPW-R properties were present from the onset of immobilization
stress (Fig. 2d and supplementary Fig. 2c) also raises interesting questions about underlying
mechanisms. For instance, this timeline suggest that both altered SPW-R characteristics and
suppressed hippocampal spiking during the acute stress are mediated by sympatho-adrenal
medullary (SAM) system-activated neurochemicals, (Cadle and Zoladz, 2015; Gunn and
Baram, 2017). Further, reports that norepinephrine application in hippocampal slices alters
SPW-R properties (Haq et al., 2012) and suppresses CA1 pyramidal spiking (Bergles et al.,
1996; Pang and Rose, 1987), suggests norepinephrine release as a likely mechanism
contributing to the stress phenotypes observed in this study. Finally, our findings on suppressed
CA1 spiking during the chronic-rest state is in agreement with the role of glucocorticoid

receptor (GR) mediated suppression of hippocampal neuronal activity during stress (Joëls,

2018).

Following repeated stress, the enhanced pyramidal cell co-activation during SPW-Rs was also

accompanied by participation of CA1 pyramidal cells in fewer SPW-Rs than during non-stress

states (Fig. 3e and f). Whether suppressed participation of pyramidal cells is an epiphenomenon

or is to counter enhanced network synchrony as stress becomes chronic, is not clear. However,

these results show that repeated stress alters hippocampal ripple-spike interactions, a

phenomenon previously linked to altered cognition in rodent models of disease (Middleton et

al., 2018; Raveau et al., 2018; Suh et al., 2013; Witton et al., 2016). Two major afferents to

area CA1 that influence information processing and SPW-R properties are the Schaffer

collaterals from area CA3 (Davoudi and Foster, 2019; Nakashiba et al., 2009) and

temporoammonic inputs from the entorhinal cortex (Oliva et al., 2018; Yamamoto and

Tonegawa, 2017). We found no change in either SPW-R-associated low-gamma power (Fig.

2f) or ripple-burst patterns (Fig. 2g), suggesting that the stress phenotypes we observed are

likely caused by changes occurring within areas CA1 and CA3 and are minimally influenced

by entorhinal inputs. In addition, extrahippocampal regions, including the amygdala, are

known to influence CA1 neural dynamics (Ghosh et al., 2013; Kim et al., 2012, 2015) , LTP

(Kim et al., 2005; Vouimba and Richter-Levin, 2005) and the release of stress hormones,

including catecholamines (Roozendaal et al., 2009; Schwabe et al., 2012). Thus, future studies

will be needed to characterize the relative contributions of CA3 and amygdalar inputs to SPW-

R duration, pyramidal cell synchrony, and altered participation of pyramidal cells in SPW-Rs,

reported in the current study.

In conclusion, a large body of earlier work has characterized the morphological, molecular,

physiological and behavioral changes in the hippocampus following either acute or repeated

stress. Our study adds a new layer of understanding by providing a window into the dynamics

of hippocampal network activity during episodes of stress and identifies altered ripple-spike

interactions as a potential biomarker of stress.

Acknowledgements

We thank all members of the CBP laboratory for their support, the Advanced Manufacturing Support Team, RIKEN Center for Advanced Photonics for their assistance in microdrive production and Lalitha Krishnan for assistance with figure generation. We also thank Dr. Kazumasa Z. Tanaka for inputs and support. This works was supported by Grant-in-Aid for Scientific Research from MEXT (19H05646; T.J.M), Grant-in-Aid for Scientific Research on Innovative Areas from MEXT (19H05233; T.J.M) and RIKEN BSI and CBS (T.J.M).

Author contributions

A.T. and T.J.M conceived the study. A.T. performed all experiments. A.T. D.P. and T.J.M. analysed the data. A.T., T.J.M. and S.C. wrote the manuscript with input from D.P. Funding provided by T.J.M.

Declaration of interests

The authors declare no competing financial interests

Correspondence and request for material

Any request for data or code should be addressed to A.T. or T.J.M.

Methods

Animals. All experiments were performed using male mice. In total 21 mice were used; of these, 4 were used for measuring the stress effects on bodyweight while the remaining 17 were used for in vivo electrophysiology. All mice were aged between 3 and 6 months at the start of experiments and were maintained on a 12-h light-dark cycle with *ad libitum* access to food and water. All procedures were approved by the RIKEN Institutional Animal Care and Use Committee and complied with all relevant ethical regulations.

Experimental design and stress protocol. Mice underwent the same chronic immobilization stress (CIS) protocol as described previously (Tomar et al., 2015). Briefly, mice experienced complete immobilization (2h/d, 10 consecutive days: Fig. 1a) in rodent immobilization bags, without access to food or water. All mice underwent the same experimental protocol as described previously, except 5 mice that experienced a familiar track on the first and 10th day of experiment followed by stress exposure. Here we examined the first 30 minutes of data recorded during CIS (stress-state) and the preceding quiescence period (rest-state). Recordings on the first day of CIS were termed 'acute' while those on the last day of experiment were termed 'chronic', providing four time points: i) acute-rest, ii) acute-stress, iii) chronic-rest and iv) chronic-stress. Rest-state data was recorded for ~ 15-30 min and hence for temporal distributions, theta/delta ratio, correlation between theta/delta and SPW-R occurrence, the first 15 minutes of data was used.

Surgery, recording and histology. Mice were anaesthetized using Avertin (2, 2, 2-tribromoethanol; Sigma-Aldrich, 476 mg/kg, i.p.) and were surgically implanted with a microdrive (manufactured with the assistance of the Advanced Manufacturing Support Team, RIKEN Center for Advanced Photonics, Japan). The microdrive housed eight independently movable tetrodes (14 µm diameter, nichrome) and was placed above right dorsal hippocampus (coordinates from bregma: AP -1.8mm; ML + (1.2mm). Prior to surgery, tetrodes were gold plated to lower impedance down to a range of 100–250 kΩ. Tetrodes were gradually lowered over the course of several days, such that by the start of the experiment they reached the CA1 stratum pyramidale. Data were acquired using a 32-channel Digital Lynx 4SX acquisition system (Neuralynx, Bozeman, MT). Signals were sampled at 32,556 Hz and spike waveforms were filtered between 600 Hz and 6 kHz. Skull screws located above the cerebellum served as a ground, and a tetrode that was seated in the superficial layers of the neocortex, and devoid of spiking activity, was used for referencing. 3-4 weeks after surgery, when all tetrodes reached the CA1 stratum pyramidale, evident by multiple large amplitude spikes and SPW-Rs, the experiment was initiated. During both rest-state and stress-state recordings the mice were located in a small circular sleep box (15 cm diameter). At the conclusion of the experiment mice underwent terminal anaesthesia (Avertin), and electric current (30 µA, for 8 s) was administered through each electrode to mark their locations. Transcardial perfusion was carried by using saline followed by 4% paraformaldehyde (PFA) followed by a further 24 h fixation in 4% PFA. Brains were sliced using a vibratome (Leica) to prepare coronal slices (50 µm thick), which were subsequently stained with Eosin Y, and inspected by standard light microscopy to confirm electrode placement.

Unit isolation and spike analysis. Spike sorting was performed by automatic spike sorting program (KlustaKwik; (Harris et al., 2000)), followed by manual adjustments of the cluster

boundaries using SpikeSort3D software (Neuralynx). Candidate clusters with < 0.5% of spikes
displaying an inter-spike-interval shorter than 2 ms, a total number of spikes exceeding 50,
having a cluster isolation distance value (Schmitzer-Torbert et al., 2005) >15, spike width
(peak-to-trough) >170 µs and complex spike index (CSI) (McHugh et al., 1996) >5 were
considered as pyramidal cells and were used for further analysis.

Local field potential analysis. The raw LFP data were first downsampled using a custom
software written in C to 1627.8 Hz (a factor of 20), followed by a quality control measure and
channel selection via visual inspection and largest power in SPW-R frequency band (80-250
Hz). A low-pass filter with a cut-off frequency equal to half the target sampling frequency was
applied to the LFP prior to downsampling to prevent any signal distortion. Power spectral
density (PSD) was calculated by using Welch's averaged modified periodogram method
(pwelch function in Matlab) with a 2048 sample window size (1.26 s), 50% overlap and 4096
FFT points (2.52 s) resulting in a time-varying spectrogram. To account for power fluctuations
caused by difference in position/impedance of the electrodes and to make PSD values
comparable across mice, each PSD curve was normalized by its own mean power within the 0
– 3 Hz frequency band. Temporal dynamics between behaviour states were assessed by binning
LFP data in 1 min time-bins and by comparing this data between stress-states and rest-states.

SPW-R detection. SPW-R events were detected using modifications to the method described
in (Csicsvari et al., 1999b). Previously selected LFP channels were first band-pass filtered (80-
250 Hz) using a 69 order Kaiser-window FIR zero-phase shift filter. Subsequently, the absolute
value of Hilbert transform (instantaneous ripple power) was smoothed with 50 ms Gaussian
window and candidate SWR-events were detected as periods where magnitude exceeded 3
standard deviation (SD) above its mean for longer than 30 ms. Further, the initiation and
termination points of candidate SWR events were defined as points when the magnitude
returned to the mean. Summed multi-unit activity (MUA) across all pyramidal cells fired
during any given recorded session was converted to instantaneous firing rate using time bin
size equal to the underlying LFP trace sampling rate and smoothed, to allow detection of firing
bursts using the same thresholds as described for the candidate SPW-R events detection.
Candidate SPW-R events not coincident with MUA bursts were excluded from subsequent
analysis. For precise peak PSD frequency detection of the SPW-R events, a multitaper method
was employed on the product of each filtered SPW-R waveform and Hanning window of the
same length.

Single Unit properties. Mean firing rate of individual single units was calculated as total
number of spikes emitted by the unit during sleep/stress trial divided by the trial's duration.
Peak firing rate of each single unit was calculated by smoothing ISIs of the unit with a 5SD
Gaussian kernel and taking maximum value of the resulting firing rate curve (firing rate over
time). Spike bursting analysis was performed as described previously (Bakkum et al., 2014) by
defining two or more spikes occurring within 10 ms time bin as a burst.

SPW-R triggered spectrograms. The spectrograms were calculated by using complex
wavelet transform (CWT) (Morlet type wavelet, parameter = 7) method applied to a segment
of the wide band LFP in 400 ms window centred on each SPW-R. Resulting spectrograms were
averaged across individual SPW-R events in each recording session/trial. To compensate for
1/f power loss each power value within each spectrogram was multiplied with the frequency
correspondent to that power value. To make power values comparable across subjects, each
spectrogram was normalized by its own mean power within (0 – 5 Hz) frequency band.

SPW-R bursting analysis. The concentration of SPW-R events in time (bursting) was estimated by first taking middle time stamp of each SPW-R event and calculating the difference between time stamps of the previous and next event. If adjacent events occurred further than 200ms apart the current event marked as 'singlet'. Number of timestamps in remaining events was counted, and each event marked as member of 'doublet', 'triplet' or 'other' (for more than 3 events in a SPW-R burst).

Participation of single units in SPW-R events. Each ripple event's start and end timestamps were used to quantify single unit activity corresponding to the co-occurred SPW-R events. Within SPW-R firing rate of any given single unit within each trial was calculated as total number of spikes generated by the single unit divided by combined duration of all SPW-R events of the trial. Between SPW-R firing rate for any given trial was calculated by first, removing all spikes fired within all SPW-R events occurred during the trial and then calculating mean firing rate of resulting spike train. Participation of a single unit in SPW-R events was calculated as % of SPW-R events the unit fired at least single spike in any given trial.

Coactivity Z-score. A likelihood of any given pair of single units (limited to pyramidal cells) firing together during SPW-R events or "coactivity Z-score" was calculated as described previously (Singer and Frank, 2009). Briefly, for any given trial a set of start and stop timestamps of SPW-R events and a set of spike train timestamps, fired by pyramidal cells were prepared. Then a 'coactivity matrix' of size (Ncells x Nspw-r events) was calculated. Each element of the coactivity matrix is set to 1 if a given cell was active (e.g. fired at least one spike) during given SPW-R event or 0 otherwise. Next, for every possible pair of cells a raw coactivity score was calculated as number of SPW-R events during which both cells of the pair were active. Finally, a z-cored coactivity value was calculated by normalisation of difference between raw coactivity score and variance by standard deviation as described in (Singer and Frank, 2009).

Statistical analysis. All statistical analyses were performed in R software (3.3.2). All boxplots were analysed using two-way ANOVAs (*aov* function, stats package) followed by Tukey's honestly significant difference (HSD) test. (*TukeyHSD* function, stats package). Distributions of various SPW-R properties as a function of SPW-R durations/amplitude were assessed using linear mixed effects models (LMMs) where mouse identity was specified as a random factor and behaviour states and categorical variables were specified as fixed factors (*lmer* function, lme4 package). The output of the *lmer* function was summarized as an ANOVA table (*anova* function, stats package). Post hoc pair-wise comparisons were made using the least-squares-means (LSM) approach (*lsmeans* function, lsmeans package). Correlation between parameters was calculated using Pearson's correlation coefficient analysis (base package). Dependence of a parameter on another was calculated by employing standardized major axis (SMA) regression (*sma* function, smatr package). For the log-transformed scatter plots, the slope and the R^2 values were calculated using log-transformed data and accordingly reported. Comparisons between regression lines were made by likelihood ratio tests (*sma* function, smatr package). For cumulative distribution analysis, the Kolmogorov-Smirnov test was employed (*ks.test,* stats package). Boxplots represent Interquartile Range (IQR, 25th-75th percentiles), median is the thick line housed in the box and whiskers extend to 1.5 times the IQR. No data point was removed as outlier either for making boxplots or for statistical analysis. Unless noted, the level of statistical significance was set to 0.05 and p values are shown as follows: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Data and code availability. The data and MATLAB codes that support the findings of this study are available from corresponding authors upon request.

References

Alfarez, D.N., Joëls, M., and Krugers, H.J. (2003). Chronic unpredictable stress impairs long-term potentiation in rat hippocampal CA1 area and dentate gyrus in vitro. Eur. J. Neurosci. *17*, 1928–1934.

Bakkum, D.J., Radivojevic, M., Frey, U., Franke, F., Hierlemann, A., and Takahashi, H. (2014). Parameters for burst detection. Front. Comput. Neurosci. *7*.

Bergles, D.E., Doze, V.A., Madison, D.V., and Smith, S.J. (1996). Excitatory actions of norepinephrine on multiple classes of hippocampal CA1 interneurons. J. Neurosci. Off. J. Soc. Neurosci. *16*, 572–585.

Buzsáki, G. (1989). Two-stage model of memory trace formation: a role for "noisy" brain states. Neuroscience *31*, 551–570.

Buzsáki, G. (2015). Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. Hippocampus *25*, 1073–1188.

Buzsáki, G., and Mizuseki, K. (2014). The log-dynamic brain: how skewed distributions affect network operations. Nat. Rev. Neurosci. *15*, 264–278.

Buzsáki, G., Horváth, Z., Urioste, R., Hetke, J., and Wise, K. (1992). High-frequency network oscillation in the hippocampus. Science *256*, 1025–1027.

Cadle, C.E., and Zoladz, P.R. (2015). Stress time-dependently influences the acquisition and retrieval of unrelated information by producing a memory of its own. Front. Psychol. *6*.

Chattarji, S., Tomar, A., Suvrathan, A., Ghosh, S., and Rahman, M.M. (2015). Neighborhood matters: divergent patterns of stress-induced plasticity across the brain. Nat. Neurosci. *18*, 1364–1375.

Cheng, S., and Frank, L.M. (2008). New Experiences Enhance Coordinated Neural Activity in the Hippocampus. Neuron *57*, 303–313.

Csicsvari, J., Hirase, H., Czurkó, A., Mamiya, A., and Buzsáki, G. (1999a). Oscillatory coupling of hippocampal pyramidal cells and interneurons in the behaving Rat. J. Neurosci. Off. J. Soc. Neurosci. *19*, 274–287.

Csicsvari, J., Hirase, H., Czurkó, A., Mamiya, A., and Buzsáki, G. (1999b). Fast network oscillations in the hippocampal CA1 region of the behaving rat. J. Neurosci. Off. J. Soc. Neurosci. *19*, RC20.

Davoudi, H., and Foster, D.J. (2019). Acute silencing of hippocampal CA3 reveals a dominant role in place field responses. Nat. Neurosci. *22*, 337–342.

Diamond, D.M., Park, C.R., and Woodson, J.C. (2004). Stress generates emotional memories and retrograde amnesia by inducing an endogenous form of hippocampal LTP. Hippocampus *14*, 281–291.

Diamond, D.M., Campbell, A.M., Park, C.R., Halonen, J., and Zoladz, P.R. (2007). The Temporal Dynamics Model of Emotional Memory Processing: A Synthesis on the Neurobiological Basis of

Stress-Induced Amnesia, Flashbulb and Traumatic Memories, and the Yerkes-Dodson Law. Neural
Plast. *2007*.

Ego-Stengel, V., and Wilson, M.A. (2010). Disruption of ripple-associated hippocampal activity during
rest impairs spatial learning in the rat. Hippocampus *20*, 1–10.

Eichenbaum, H. (2017). On the Integration of Space, Time, and Memory. Neuron *95*, 1007–1018.

Fernández-Ruiz, A., Oliva, A., Oliveira, E.F. de, Rocha-Almeida, F., Tingley, D., and Buzsáki, G. (2019).
Long-duration hippocampal sharp wave ripples improve memory. Science *364*, 1082–1086.

Foster, T.C., Castro, C.A., and McNaughton, B.L. (1989). Spatial selectivity of rat hippocampal
neurons: dependence on preparedness for movement. Science *244*, 1580–1582.

Ghosh, S., Laxmi, T.R., and Chattarji, S. (2013). Functional connectivity from the amygdala to the
hippocampus grows stronger after stress. J. Neurosci. Off. J. Soc. Neurosci. *33*, 7234–7244.

Girardeau, G., Benchenane, K., Wiener, S.I., Buzsáki, G., and Zugaro, M.B. (2009). Selective
suppression of hippocampal ripples impairs spatial memory. Nat. Neurosci. *12*, 1222–1223.

Grosmark, A.D., Mizuseki, K., Pastalkova, E., Diba, K., and Buzsáki, G. (2012). REM Sleep Reorganizes
Hippocampal Excitability. Neuron *75*, 1001–1007.

Gunn, B.G., and Baram, T.Z. (2017). Stress and Seizures: Space, Time and Hippocampal Circuits.
Trends Neurosci. *40*, 667–679.

Haq, R.U., Liotta, A., Kovacs, R., Rösler, A., Jarosch, M.J., Heinemann, U., and Behrens, C.J. (2012).
Adrenergic modulation of sharp wave-ripple activity in rat hippocampal slices. Hippocampus *22*,
516–533.

Harris, K.D., Henze, D.A., Csicsvari, J., Hirase, H., and Buzsáki, G. (2000). Accuracy of Tetrode Spike
Separation as Determined by Simultaneous Intracellular and Extracellular Measurements. J.
Neurophysiol. *84*, 401–414.

Jadhav, S.P., Kemere, C., German, P.W., and Frank, L.M. (2012). Awake Hippocampal Sharp-Wave
Ripples Support Spatial Memory. Science *336*, 1454–1458.

Joëls, M. (2009). Stress, the hippocampus, and epilepsy. Epilepsia *50*, 586–597.

Joëls, M. (2018). Corticosteroids and the brain. J. Endocrinol. *238*, R121–R130.

Joo, H.R., and Frank, L.M. (2018). The hippocampal sharp wave-ripple in memory retrieval for
immediate use and consolidation. Nat. Rev. Neurosci. *19*, 744–757.

Kim, E.J., Kim, E.S., Park, M., Cho, J., and Kim, J.J. (2012). Amygdalar Stimulation Produces Alterations
on Firing Properties of Hippocampal Place Cells. J. Neurosci. *32*, 11424–11434.

Kim, E.J., Pellman, B., and Kim, J.J. (2015). Stress effects on the hippocampus: a critical review. Learn.
Mem. *22*, 411–416.

Kim, J.J., Koo, J.W., Lee, H.J., and Han, J.-S. (2005). Amygdalar Inactivation Blocks Stress-Induced
Impairments in Hippocampal Long-Term Potentiation and Spatial Memory. J. Neurosci. *25*, 1532–
1539.

Kim, J.J., Lee, H.J., Welday, A.C., Song, E., Cho, J., Sharp, P.E., Jung, M.W., and Blair, H.T. (2007). Stress-induced alterations in hippocampal plasticity, place cells, and spatial memory. Proc. Natl. Acad. Sci. U. S. A. *104*, 18297–18302.

Krugers, H.J., Douma, B.R., Andringa, G., Bohus, B., Korf, J., and Luiten, P.G. (1997). Exposure to chronic psychosocial stress and corticosterone in the rat: effects on spatial discrimination learning and hippocampal protein kinase Cgamma immunoreactivity. Hippocampus *7*, 427–436.

Magariños, A.M., Verdugo, J.M.G., and McEwen, B.S. (1997). Chronic stress alters synaptic terminal structure in hippocampus. Proc. Natl. Acad. Sci. *94*, 14002–14008.

McEwen, B.S. (2007). Physiology and Neurobiology of Stress and Adaptation: Central Role of the Brain. Physiol. Rev. *87*, 873–904.

McEwen, B.S., Bowles, N.P., Gray, J.D., Hill, M.N., Hunter, R.G., Karatsoreos, I.N., and Nasca, C. (2015). Mechanisms of stress in the brain. Nat. Neurosci. *18*, 1353–1363.

McHugh, T.J., Blum, K.I., Tsien, J.Z., Tonegawa, S., and Wilson, M.A. (1996). Impaired hippocampal representation of space in CA1-specific NMDAR1 knockout mice. Cell *87*, 1339–1349.

Middleton, S.J., Kneller, E.M., Chen, S., Ogiwara, I., Montal, M., Yamakawa, K., and McHugh, T.J. (2018). Altered hippocampal replay is associated with memory impairment in mice heterozygous for the Scn2a gene. Nat. Neurosci. *21*, 996–1003.

Nakashiba, T., Buhl, D.L., McHugh, T.J., and Tonegawa, S. (2009). Hippocampal CA3 output is crucial for ripple-associated reactivation and consolidation of memory. Neuron *62*, 781–787.

O'Keefe, J., and Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. Brain Res. *34*, 171–175.

O'Keefe, J., and Nadel, L. (1978). The Hippocampus as a Cognitive Map (Oxford University Press, USA).

Oliva, A., Fernández-Ruiz, A., Fermino de Oliveira, E., and Buzsáki, G. (2018). Origin of Gamma Frequency Power during Hippocampal Sharp-Wave Ripples. Cell Rep. *25*, 1693-1700.e4.

Pang, K., and Rose, G.M. (1987). Differential effects of norepinephrine on hippocampal complex-spike and θ-neurons. Brain Res. *425*, 146–158.

Park, M., Kim, C.-H., Jo, S., Kim, E.J., Rhim, H., Lee, C.J., Kim, J.J., and Cho, J. (2015). Chronic Stress Alters Spatial Representation and Bursting Patterns of Place Cells in Behaving Mice. Sci. Rep. *5*, 16235.

Passecker, J., Hok, V., Della-Chiesa, A., Chah, E., and O'Mara, S.M. (2011). Dissociation of dorsal hippocampal regional activation under the influence of stress in freely behaving rats. Front. Behav. Neurosci. *5*, 66.

Pfeiffer, B.E., and Foster, D.J. (2013). Hippocampal place-cell sequences depict future paths to remembered goals. Nature *497*, 74–79.

Ramirez, S., Liu, X., MacDonald, C.J., Moffa, A., Zhou, J., Redondo, R.L., and Tonegawa, S. (2015). Activating positive memory engrams suppresses depression-like behaviour. Nature *522*, 335–339.

Raveau, M., Polygalov, D., Boehringer, R., Amano, K., Yamakawa, K., and McHugh, T.J. (2018).
Alterations of in vivo CA1 network activity in Dp(16)1Yey Down syndrome model mice.

Roozendaal, B., McEwen, B.S., and Chattarji, S. (2009). Stress, memory and the amygdala. Nat. Rev.
Neurosci. *10*, 423–433.

Sandi, C., Davies, H.A., Cordero, M.I., Rodriguez, J.J., Popov, V.I., and Stewart, M.G. (2003). Rapid
reversal of stress induced loss of synapses in CA3 of rat hippocampus following water maze training.
Eur. J. Neurosci. *17*, 2447–2456.

Sapolsky, R.M. (1996). Stress, Glucocorticoids, and Damage to the Nervous System: The Current
State of Confusion. Stress *1*, 1–19.

Sapolsky, R.M., Krey, L.C., and McEwen, B.S. (1986). The Neuroendocrinology of Stress and Aging:
The Glucocorticoid Cascade Hypothesis. Endocr. Rev. *7*, 284–301.

Schmidt, B., and Redish, A.D. (2013). Navigation with a cognitive map. Nature *497*, 42–43.

Schmitzer-Torbert, N., Jackson, J., Henze, D., Harris, K., and Redish, A.D. (2005). Quantitative
measures of cluster quality for use in extracellular recordings. Neuroscience *131*, 1–11.

Schwabe, L., Joëls, M., Roozendaal, B., Wolf, O.T., and Oitzl, M.S. (2012). Stress effects on memory:
An update and integration. Neurosci. Biobehav. Rev. *36*, 1740–1749.

Shors, T.J., Seib, T.B., Levine, S., and Thompson, R.F. (1989). Inescapable versus escapable shock
modulates long-term potentiation in the rat hippocampus. Science *244*, 224–226.

Singer, A.C., and Frank, L.M. (2009). Rewarded outcomes enhance reactivation of experience in the
hippocampus. Neuron *64*, 910–921.

Sousa, N., Lukoyanov, N.V., Madeira, M.D., Almeida, O.F., and Paula-Barbosa, M.M. (2000).
Reorganization of the morphology of hippocampal neurites and synapses after stress-induced
damage correlates with behavioral improvement. Neuroscience *97*, 253–266.

Suh, J., Foster, D.J., Davoudi, H., Wilson, M.A., and Tonegawa, S. (2013). Impaired Hippocampal
Ripple-Associated Replay in a Mouse Model of Schizophrenia. Neuron *80*, 484–493.

Suvrathan, A., Tomar, A., and Chattarji, S. (2010). Effects of chronic and acute stress on rat behaviour
in the forced-swim test. Stress Amst. Neth. *13*, 533–540.

Tomar, A., Polygalov, D., Chattarji, S., and McHugh, T.J. (2015). The dynamic impact of repeated
stress on the hippocampal spatial map. Hippocampus *25*, 38–50.

Tulving, E. (1985). Memory and consciousness. Can. Psychol. Can. *26*, 1–12.

Vouimba, R.-M., and Richter-Levin, G. (2005). Physiological Dissociation in Hippocampal Subregions
in Response to Amygdala Stimulation. Cereb. Cortex *15*, 1815–1821.

Vyas, A., Mitra, R., Shankaranarayana Rao, B.S., and Chattarji, S. (2002). Chronic stress induces
contrasting patterns of dendritic remodeling in hippocampal and amygdaloid neurons. J. Neurosci.
22, 6810.

Wilson, M.A., and McNaughton, B.L. (1994). Reactivation of hippocampal ensemble memories during
sleep. Science *265*, 676–679.

Witton, J., Staniaszek, L.E., Bartsch, U., Randall, A.D., Jones, M.W., and Brown, J.T. (2016). Disrupted
hippocampal sharp-wave ripple-associated spike dynamics in a transgenic mouse model of
dementia. J. Physiol. *594*, 4615–4630.

Yamamoto, J., and Tonegawa, S. (2017). Direct Medial Entorhinal Cortex Input to Hippocampal CA1
Is Crucial for Extended Quiet Awake Replay. Neuron *96*, 217-227.e4.

Ylinen, A., Bragin, A., Nadasdy, Z., Jando, G., Szabo, I., Sik, A., and Buzsaki, G. (1995). Sharp wave-
associated high-frequency oscillation (200 Hz) in the intact hippocampus: network and intracellular
mechanisms. J. Neurosci. *15*, 30–46.

Fig. 1. CA1 pyramidal cell activity is altered during stress. a Schematic representation of the chronic immobilization stress (CIS) protocol that mice received every day for 10 consecutive days. **b** Coronal section of the hippocampus showing the tetrode locations (black arrows) at the CA1 pyramidal layer. **c** Time matched (30 min) representative examples of unsorted spikes recorded during rest-state (left) and stress-state (right). Numbers next to each cluster depict total number of spikes recorded during that state. **d** Mean firing rate between rest-state and stress-state (two-way ANOVA: behaviour-state, $F(1, 1074) = 29.611$, p = 6.54×10^{-8}; day, $F(1, 1074) = 0.908$, p = 0.341; behaviour-state x day interaction, $F(1,1074) = 9.118$, p = 0.0026; Tukey's HSD: acute-rest vs acute-stress, p = 1.32×10^{-8}; acute-rest vs chronic-rest, p=0.017; chronic-rest vs chronic-stress, p = 0.42). **e** Peak firing rate between rest-state and stress-state (two-way ANOVA: behaviour-state, $F(1, 1074) = 54.234$, p = 3.54×10^{-13}; day, $F(1, 1074) = 4.014$, p = 0.0454; behaviour-state x day interaction, $F(1, 1074) = 0.005$, p

$= 0.945$; Tukey's HSD: acute-rest vs acute-stress, $p = 7.0 \times 10^{-7}$; chronic-rest, vs chronic-stress,
$p = 2.5 \times 10^{-6}$). **f** Inter-burst-interval between rest-state and stress-state (two-way ANOVA:
behaviour-state, $F(1, 1066) = 47.294$, $p = 1.04 \times 10^{-11}$; day, $F(1, 1066) = 0.749$, $p = 0.387$;
behaviour-state x day interaction, $F(1, 1066) = 3.685$, $p = 0.055$; Tukey's HSD: acute-rest vs
acute-stress, 2.2×10^{-9}; chronic-rest vs chronic-stress, $p = 0.005$). **g** Correlation between
theta/delta ratio and SPW-Rs differs between behaviour-states on the first day (left: acute-rest,
slope $= -0.044$, $R = -0.56$, $p = 0.02$; acute-stress, slope $= 0.00$, $R = -0.19$, $p = 0.47$) and last day
(right: chronic-rest, slope $= -0.04$, $R = -0.54$, $p=0.029$; chronic-stress, slope $= -0.01$, $R = -0.17$,
$p = 0.52$) of CIS. All box plots represent interquartile range (IQR, 25^{th}-75^{th} percentiles),
median is the thick line in the box and whiskers extend to 1.5 times the IQR. Circles depict
rest-state (black) stress-state (red). * $p<0.05$, ** $p < 0.01$, *** $p < 0.001$. acute-rest: n = 288
cells, N = 17 mice, acute-stress: n = 282 cells, N = 16 mice, chronic rest: n = 282 cells, N = 16
mice, chronic stress: n = 226 cells, N = 16 mice).

Fig. 2. CA1 SPW-R duration differs between rest and stress states. a Representative examples of local field potential (LFP) for non-filtered (top) and filtered (bottom) SPW-R events recorded during rest-state (left) and stress-state (right) from CA1 *stratum pyramidale*. **b** SPW-R duration differs between rest-state and stress-state; (two-way ANOVA: behaviour-state, $F_{(1, 61)} = 35.233$, $p = 1.5 \times 10^{-7}$; day, $F_{(1, 61)} = 0.114$, $p = 0.737$; behaviour-state x day, $F_{(1, 61)} = 1.002$, $p = 0.321$; Tukey's HSD: acute-rest vs acute-stress, $p = 3.86 \times 10^{-5}$, chronic-rest vs chronic-stress, $p = 0.0055$). **c** Distributions of the duration of SPW-Rs differs between

behaviour states on first day (left: acute-rest vs acute-stress, Kolmogorov-Smirnov test, $D = 0.236$, $p < 2.22 \times 10^{-16}$) and last day (right: chronic-rest vs chronic-stress, Kolmogorov-Smirnov test, $D = 0.113$, $p < 2.22 \times 10^{-16}$) showed a rightward shift. **d** Temporal dynamics of averaged SPW-R duration (1-min bins) differ between behaviour states on the first day (left: LMMs; behaviour-state, $F_{(1, 447)} = 249.26$, $p < 2.22 \times 10^{-16}$; time, $F_{(14, 447)} = 0.703$, $p = 0.772$; behaviour-state x time, $F_{(14, 447)} = 0.456$, $p = 0.955$). Similarly, on the last day, SPW-R duration increased during stress (right: LMMs; behaviour-state, $F_{(1, 412)} = 68.75$, $p = 1.74 \times 10^{-15}$; time: $F_{(14, 412)} = 1.05$, $p = 0.404$; behaviour-state x time, $F_{(14, 412)} = 1.226$, $p = 0.253$). **e** Averaged peri-SPW-R wavelet spectrograms during acute-rest (top left), acute-stress (bottom left), chronic-rest (top right) and chronic-stress (bottom right). **f** Low gamma (17-40Hz) power during SPW-R events does not differ between behaviour-states (two-way ANOVA: behaviour-state, $F_{(1, 61)} = 1.31$, $p = 0.26$; day, $F_{(1, 61)} = 0.55$, $p = 0.46$; behaviour-state x day, $F_{(1, 61)} = 0.07$, $p = 0.79$). **g** Fraction of different types of ripple bursts during acute stress (two-way ANOVA: behaviour-state, $F_{(1, 124)} = .00$, $p = 1$; category, $F_{(3, 124)} = 1521.05$, $p < 2.22 \times 10^{-16}$; behaviour-state x category, $F_{(3, 124)} = 0.22$, $p = 0.88$; Tukey's HSD: acute-rest vs acute-stress: singlets, $p = 0.999$; doublets, $p = 0.999$; triplets, $p = 0.999$; others, $p = 0.999$) and chronic stress (two-way ANOVA: behaviour-state, $F_{(1, 120)} = .00$, $p = 1$; category, $F_{(3, 120)} = 981.21$, $p < 2.22 \times 10^{-16}$; behaviour-state x category, $F_{(3, 124)} = 0.169$, $p = 0.917$; Tukey's HSD: chronic-rest vs chronic-stress: singlets, $p = 1.0$; doublets, $p = 0.999$; triplets, $p = 0.999$; others, $p = 0.999$). All box plots represent median and 25th-75th percentiles, with whiskers extending to the extreme data points. Circles depict rest-state (black) stress-state (red). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Acute-rest: n = 7220 SPW-Rs, N = 17 mice; acute-stress: n = 9851 SPW-Rs, N = 16 mice; chronic rest: n = 7041 SPW-Rs, N = 16 mice; chronic stress: n = 8978 SPW-Rs, N = 16 mice).

Fig. 3. Altered CA1 pyramidal activation in SPW-Rs during stress-state.

a The relationship between SPW-R duration (20-ms bins) and multiunit activity during SPW-Rs for first (left) and last (right) day of CIS (LMMs: behaviour-state, $F(1, 16896) = 0.2325$, p

= 0.63; duration, $F_{(16, 16896)} = 715.93$, $p < 2.22\text{x}10^{-16}$; behaviour-state x duration, $F_{(16, 16896)} = 7.615$, $p < 2.22\text{x}10^{-16}$) and the last day (LMMs: behaviour-state, $F_{(1, 15877)} = 1.877$, $p = 0.17$; duration, $F_{(16, 15877)} = 525.5$, $p < 2.22\text{x}10^{-16}$; behaviour-state x day, $F_{(16, 15877)} = 36.435$, $p < 2.22\text{x}10^{-16}$) of CIS). Green dots on top of graph indicate significant differences between behaviour-states. **b** Within SPW-R average firing rate differs between behaviour-states (two-way ANOVA: behaviour-state, $F_{(1, 1074)} = 13.981$, $p = 1.9\text{x}10^{-4}$; day, $F_{(1, 1074)} = 2.892$, $p = 0.089$; behaviour-state x day, $F_{(1, 1074)} = 4.564$, $p = 0.033$; acute-rest vs acute-stress, $p = 1.78\text{x}10^{-4}$; chronic-rest vs chronic-stress, $p = 0.749$; acute-rest vs chronic-rest, $p = 0.024$). **c** Percentage of spikes discharged by CA1 pyramidal cells in SPW-R differs between behaviour-states (two-way ANOVA: behaviour-state, $F_{(1, 1074)} = 84.83$, $p < 2.22\text{x}10^{-16}$; day $F_{(1, 1074)} = 0.079$, $p = 0.778$; behaviour-state x day, $F_{(1, 1074)} = 1.35$, $p = 0.25$; Tukey's HSD: acute-rest vs acute-stress, $p = 1.4\text{x}10^{-12}$; chronic-rest vs chronic-stress, $p = 3.62\text{x}10^{-7}$). **d** Dependency of pyramidal cell participation in SPW-Rs on their mean firing rate on the first day (acute-rest: slope = 0.848, $R^2 = 0.55$, $p < 2.22\text{x}10^{-16}$; acute-stress: slope = 0.96, $R^2 = 0.40$, $p < 2.22\text{x}10^{-16}$; likelihood ratio test (df = 1) = 4.318, $p = 0.038$) and the last day (chronic-rest: slope = 0.903, $R^2 = 0.42$, $p < 2.22\text{x}10^{-16}$; chronic-stress: slope=0.898, R^2=0.21, p=$2.97\text{x}10^{-13}$; likelihood ratio test (df = 1) = 0.006, p = 0.94) of CIS. **e** Pyramidal cell participation in SPW-Rs differs with the progression of stress protocol (two-way ANOVA: behaviour-state, $F_{(1, 1074)} = 1.686$, $p = 0.194$; day, $F_{(1, 1074)} = 5.695$, $p = 0.017$; behaviour-state x day, $F_{(1, 1074)} = 2.19$, $p = 0.139$; Tukey's HSD: acute-rest vs acute-stress, $p = 0.204$; chronic-rest vs chronic-stress, $p = 0.998$; acute-rest vs chronic rest, $p = 0.027$). **f** Cumulative distribution of pyramidal cell participation in SPW-Rs differs between rest-states on the first day (black) and last day (blue) of CIS (Kolmogorov-Smirnov test: D = 0.142, p = 0.006). All box plots, represent interquartile range (IQR, 25th-75th percentiles), median is the thick line in the box and whiskers extend to 1.5 times the IQR. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Acute-rest: n = 288 cells, N = 17 mice; acute-stress: n = 282 cells, N = 16 mice; chronic rest: n = 282 cells, N = 16 mice; chronic stress: n = 226 cells, N = 16 mice).

Fig. 4. Altered CA1 pyramidal cell coactivation in SPW-Rs during stress-state.

a Co-activity Z-scores of pyramidal cells during SPW-Rs differ between rest-state (black) and stress-state (red) (two-way ANOVA: behaviour-state, $F(1, 6195) = 60.67$, $p = 7.87\times10^{-15}$; day, $F(1, 6195) = 15.39$, $p = 8.84\times10^{-5}$; behaviour-state x day, $F(1, 6195) = 0.26$, $p = 0.61$; Tukey's HSD: acute-rest, $n = 1625$ cell pairs vs acute-stress, $n = 1610$ cell pairs, $p = 1.2\times10^{-8}$; chronic-rest, 1701 cell pairs vs chronic-stress, $n = 1263$ cell pairs, $p = 3.8\times10^{-6}$; acute-rest vs chronic-rest, $p = 0.0016$). **b** Relationship between cell pairs of positive co-activity Z-scores during both rest-state and stress-state significantly differs between days (acute: slope = 0.99, $R^2 = 0.022$, $p = 0.033$ n = 208 cell pairs; chronic: slope = 1.74, $R^2 = 0.019$, $p = 0.076$, n = 165 cell pairs; likelihood ratio test, $p = 1.28\times10^{-7}$) of CIS. Boxplots represent interquartile range (IQR, 25th-75th percentiles), median is the thick line in the box and whiskers extend to 1.5 times the IQR. $* p < 0.01$, $** p < 0.005$, $*** p < 0.001$. Acute-rest: N = 17 mice; acute-stress, N = 16 mice; chronic-rest, N = 16 mice; chronic-stress, N = 16 mice)