Title Page

Title:

Simultaneously monitoring aquatic and terrestrial biodiversity using riverine water eDNA: seasonal

variation of monitoring effectiveness

Running title:

Monitoring aquatic and terrestrial biodiversity

Authors:

Haile Yang [1], Hao Du [1,*], Hongfang Qi [2], Luxian Yu [2], Xindong Hou [3], Hui Zhang [1], Junyi Li [1], Jinming

Wu [1], Chengyou Wang [1], Qiong Zhou [1], Qiwei Wei [1,*]

[1] Key Laboratory of Freshwater Biodiversity Conservation, Ministry of Agriculture and Rural Affairs of China,

Yangtze River Fisheries Research Institute, Chinese Academy of Fishery Sciences, Wuhan 430223, China

[2] Qinghai Key Laboratory of Qinghai-Lake Naked Carps Breeding and Conservation, The Rescue and

Rehabilitation Center of Naked Carps of Qinghai Lake, Xining 810016, China

[3] State Key Laboratory of Biogeology and Environmental Geology, China University of Geoscience, Wuhan

430074, China

Corresponding authors:

Hao Du, duhao@yfi.ac.cn

Qiwei Wei, weiqw@yfi.ac.cn

Abstract

Both aquatic and terrestrial biodiversity information can be detected in riverine water environmental DNA (eDNA). However, the monitoring effectiveness (i.e., the proportion of aquatic and terrestrial biodiversity information detected in riverine water eDNA samples) is unknown. To investigate the monitoring effectiveness, we introduced the concept of watershed biological information flow (WBIF) and proposed that the monitoring effectiveness depended on the transportation effectiveness of the WBIF. Then, the monitoring effectiveness could be assessed in the WBIF framework. Here, we conducted a monitoring effectiveness assessment case study in a watershed on the Qinghai-Tibet Plateau according to analysis of the bacterial operational taxonomic unit (OTU) assemblages detected in riverine water eDNA samples and riparian soil eDNA samples during three seasons. The results showed that (1) the downstream-to-upstream monitoring effectiveness: only 76% of the bacterial OTUs could be detected 1 km downstream in spring and more than 97% and 96% could be detected in summer and autumn, respectively. (2) The river-to-land monitoring effectiveness: more than 62% of the bacterial OTUs in riparian soil eDNA samples could be detected in adjacent riverine water eDNA samples on rainy summer days and 16% and 48% could be detected on cloudy spring and autumn days, respectively. These results suggested that riverine water eDNA was viable for simultaneously monitoring aquatic and terrestrial biodiversity and that rainy days in summer or autumn were suitable sampling times on the Qinghai-Tibet Plateau. More studies on monitoring effectiveness in other taxonomies and in other watersheds with different climatic conditions are needed to support simultaneous aquatic and terrestrial biodiversity assessments.

Keywords: environmental DNA, biodiversity monitoring, monitoring effectiveness, watershed biological

44   information flow, transportation effectiveness, Qinghai-Tibet Plateau

Introduction

Biodiversity monitoring is the basal content for ecological research, ecosystem management and

conservation action (Cardinale, et al.,2012; Hooper, et al.,2012; Carrizo, et al.,2017; Mallick &

Chakraborty,2018). While abiotic environmental conditions and ecosystem property information are now

available at highly resolved spatial and temporal scales (Vina, et al.,2013; Jetz, et al.,2016), biodiversity is

still often studied from a local and accumulative perspective and is generally not available at a wide

taxonomic breadth, high-resolution temporal scale and spatial coverage (Anderson,2018; Chase, et al.,2018;

Mcglinn, et al.,2019; Altermatt, et al.,2020). This limitation slows the development of ecological research,

ecosystem management and conservation action. Now, meta-barcoding and high-throughput sequencing of

environmental DNA (eDNA, i.e., DNA extracted from environmental samples such as water, soil, and air)

provide novel opportunities to monitor biodiversity (Deiner, et al.,2016; Cristescu & Hebert,2018;

Altermatt, et al.,2020), and this approach is nonlethal for most classically sampled taxonomic groups,

minimizes habitat disruption and can assess diversity across the tree of life with a single-field sampling

protocol, making it extremely cost effective (Deiner, et al.,2016; Valentini, et al.,2016; Stat, et al.,2017).

As an efficient and easy-to-standardize monitoring approach (Thomsen & Willerslev,2015; Valentini, et

al.,2016; Lugg, et al.,2018; Seymour,2019), and with the continuous advancements in DNA sequencing

technology, using eDNA metabarcoding to monitor biodiversity would be an appropriate method to

revolutionize biodiversity monitoring by enabling the census of wide taxonomic species on a highly

resolved spatial and temporal scale in near real time (Thomsen & Willerslev,2015; Cristescu &

Hebert,2018; Altermatt, et al.,2020).

Streams and rivers connect upstream regions with downstream regions, connect land with waterbodies, and

transport natural and anthropogenic materials and information through extensive and heterogeneous

network systems (Luo, et al.,2011; Deiner, et al.,2016; Wang, et al.,2016; Matsuoka, et al.,2019). Riverine

water eDNA incorporates biodiversity information across terrestrial and aquatic biomes (Deiner, et

al.,2016; Matsuoka, et al.,2019; Yang, et al.,2019). By analyzing riverine water eDNA, information on

terrestrial and aquatic biological compositions and even their spatial structures can be obtained (Deiner, et

al.,2016; Matsuoka, et al.,2019). Therefore, the eDNA that is transported in river networks offers a novel

and spatially integrated component that can be used to simultaneously monitor aquatic and terrestrial

biodiversity and will transform biodiversity data acquisition in research, management and conservation

(Deiner, et al.,2016; Yang, et al.,2019). However, to achieve this objective, the monitoring effectiveness

(i.e., the proportion of aquatic and terrestrial biodiversity information that could be detected using limited

riverine water eDNA samples) needs to be assessed first. Until now, there has been no systemic research on

monitoring effectiveness.

The concept of watershed biological information flow (WBIF) (Yang, et al.,2019) is a good conceptual

framework to assist with the assessment of monitoring effectiveness. In the WBIF framework, the

monitoring effectiveness depends on the transportation effectiveness of the WBIF from the upstream to

downstream regions and from the land to river (Deiner & Altermatt,2014; Sansom & Sassoubre,2017; Pont,

et al.,2018; Seymour,2019; Yang, et al.,2019). The WBIF framework integrates the processes of origin,

state, transport, and fate of eDNA (Barnes & Turner,2016; Shogren, et al.,2017). Many studies have shown

that eDNA degrades over time in a logistic manner (Barnes, et al.,2014; Nukazawa, et al.,2018;

Seymour,2019; M, et al.,2020), and the detection distance varies from less than 1 km in a small stream to

more than 100 km in a large river (Deiner & Altermatt,2014; Stoeckle, et al.,2016; Pont, et al.,2018;

Seymour,2019). Deiner and colleagues (2016) indicated that much terrestrial biological information has

been input to rivers and that a large number of eukaryotic phyla from terrestrial taxa can be detected from

riverine water eDNA (Deiner, et al.,2016). Our previous WBIF study conducted on the Qinghai-Tibet

Plateau in summer showed that the transportation effectiveness from adjacent riparian sites to river sites

was 62.76% on rainy days and 44.16% on sunny days; additionally, the transportation effectiveness from

adjacent upstream sites to downstream sites (distance varied from 7 km to 23.5 km) was approximately

80.30% (Yang, et al.,2019). Moreover, it was shown that the transportation effectiveness of WBIF relied

on transport capacity, degradation rate and environmental filtration (Yang, et al.,2019). The transport

capacity mainly depended on erosion and runoff, and degradation rate mainly depended on environmental

features, and environmental filtration mainly depended on environmental change, all of which were related

to season and weather conditions (Yang, et al.,2019). Because the monitoring effectiveness depends on the

transportation effectiveness of WBIF and WBIF is related to season and weather conditions, we propose

that the monitoring effectiveness varies with season and weather conditions.

The aim of this study was to identify the effectiveness of monitoring the biodiversity information of

upstream and riparian zones using riverine water eDNA in different seasons and weather conditions. In this

study, based on the WBIF framework, we conducted a case study in the Shaliu River basin, which is a

typical watershed on the Qinghai-Tibet Plateau; specifically, we analyzed the transportation effectiveness

of two types of WBIFs, i.e., the WBIF from upstream to downstream regions and the WBIF from riparian

zones to rivers, in different seasons and weather conditions, as indicated by environmental microbes. Then,

we used the transportation effectiveness of the WBIF to identify the corresponding monitoring

effectiveness. Our objectives were twofold: first, we sought to establish whether riverine water eDNA was

viable for monitoring the biodiversity information of upstream and riparian zones. Second, we sought to

identify which season and weather conditions were optimal for monitoring the biodiversity information of

the upstream and riparian zones.

Materials and Methods

Study Area

The Shaliu River basin (37°10′-37°52′ N, 100°17′-99°32′ E), as a sub-basin of the Qinghai Lake basin, is

located 3196 m above sea level on the Qinghai-Tibet Plateau (Fig. 1). The mean annual precipitation is

423.4 mm, and the average annual evaporation is 1674.7 mm. The annual mean temperature is −0.6 ℃, the

monthly mean temperature of January is -17.5 ℃ and that of July is 11.0 ℃. The Shaliu River freezes in

October and unfreezes in the following April. The Shaliu River is 106 km long, with a catchment area of

1320 km^2. Grassland is the main land cover type, accounting for more than 90% of the watershed area.

Less than 5% of the watershed area was seriously changed by human activity, such as transformation into

cultivated land and building land[1]. Due to its simple ecosystem assemblages and weak disturbance by

human activity, the Shaliu River basin is a natural simplified model for investigating the effectiveness of

---

[1] http://www.gangcha.gov.cn/html/2125/item.html

monitoring aquatic and terrestrial biodiversity information using riverine water eDNA.

Field Sampling

We collected eDNA samples three times, including 27 soil eDNA samples and 27 water eDNA samples,

from 9 transects (Fig. 1) in the Shaliu River on April 8 and 9, June 25 and 26 and September 19 and 20,

2019. A 1.5 L surface water sample was collected from the river site of each transect and transported at 0 ℃

to the laboratory of the Rescue and Rehabilitation Center of Naked Carps of Qinghai Lake. Then, water

samples were filtered using 0.2-μm membrane filters to obtain the eDNA sample in the laboratory, and the

water eDNA samples were frozen, transported at -20 ℃ and stored at -80 ℃ until DNA extraction. A 5 mL

soil eDNA sample was collected from the riparian zone site (5 m distance from the river) of each transect

and transported to the laboratory at 0 ℃. Then, the soil eDNA samples were frozen, transported at -20 ℃

and stored at -80 ℃ until DNA extraction.

In the first sampling period (spring group), during April 8 and 9, the air temperature was -6~8 ℃, the water

temperature was -0.5~0.7 ℃, the frozen river was starting to thaw, the runoff volume was 1.8~3.9 m³/s, the

soil was still frozen, and both days were cloudy with freezing, heavy winds. On the frozen days of April 8

and 9, the river was clear; on these days, we sampled 18 samples (9 water samples and 9 soil samples) at 9

transects from the downstream to upstream regions.

In the second sampling period (summer group), during June 25 and 26, the air temperature was 7~17 ℃,

the water temperature was 4.3~16.4 ℃, and the runoff volume was 29.9~45.5 m³/s. On the sunny day of

June 25, the river was clear; on this day, we collected 4 samples (2 water samples and 2 soil samples) from

the transects of SL1 and SL2. It began to rain on the night of June 25, and on the rainy day of June 26, the

river was turbid; we collected samples (7 water samples and 7 soil samples) from the last 7 transects from

the downstream to upstream regions.

In the third sampling period (autumn group) on September 19 and 20, the air temperature was 0~10 ℃, the

water temperature was 0.2~8.8 ℃, part of the transects started to freeze, and the runoff volume was

5.7~12.8 m³/s. On the rainy day of September 19, the river was turbid; we collected 4 samples (2 water

samples and 2 soil samples) from the SL1 and SL2 transects. On the cloudy day of September 20, the river

was clear, and we collected 14 samples (7 water samples and 7 soil samples) from the last 7 transects from

the downstream to upstream regions.

DNA Extraction and Sequence Analysis

Microbial DNA was extracted from eDNA samples using an E.Z.N.A.® Stool DNA Kit (Omega BioTek,

Norcross, GA, USA) according to the manufacturer's protocols. Then, the final DNA concentration and

purity were determined by a NanoDrop 2000 UV-vis spectrophotometer (Thermo Scientific, Wilmington,

USA), and DNA quality was checked by 1% agarose gel electrophoresis. The V3-V4 hypervariable regions

of the bacterial 16S rRNA gene were amplified with the primers 338F (5'-

ACTCCTACGGGAGGCAGCAG-3') and 806R (5'-GGACTACHVGGGTWTCTAAT-3') by using a PCR

thermocycler system (GeneAmp 9700, ABI, USA). The PCRs were conducted using the following

program: 3 min of denaturation at 95 ℃; 29 cycles of 30 s at 95 ℃, 30 s for annealing at 55 ℃, and 45 s for

elongation at 72 ℃; and a final extension at 72 ℃ for 10 min. The PCRs were performed in triplicate 20-μL

mixtures containing 4 μL of 5× FastPfu Buffer, 2 μL of 2.5 mM dNTPs, 0.8 μL of each primer (5 μM), 0.4

μL of FastPfu Polymerase, 0.2 μL of BSA and 10 ng of template DNA. The resulting PCR products were

extracted from a 2% agarose gel, further purified using an AxyPrep DNA Gel Extraction Kit (Axygen

Biosciences, Union City, CA, USA) and quantified using QuantiFluor™-ST (Promega, USA) according to

the manufacturer's protocol.

Purified amplicons were pooled in equimolar amounts and subjected to paired-end sequencing on an

Illumina MiSeq platform (Illumina, San Diego, USA) according to standard protocols by Majorbio

Bio-Pharm Technology Co., Ltd. (Shanghai, China). Raw fastq files were demultiplexed, quality-filtered

by Trimmomatic and merged by FLASH. Operational taxonomic units (OTUs) were clustered with a 97%

similarity cutoff using UPARSE, and chimeric sequences were identified and removed using UCHIME.

The taxonomy of each 16S rRNA gene sequence was analyzed by the RDP classifier Bayesian algorithm

against the Silva132/16S_Bacteria database using a confidence threshold of 70%. The data were analyzed

on the Majorbio Cloud Platform (www.majorbio.com). The raw data have been deposited in the CNSA

(https://db.cngb.org/cnsa/) of CNGBdb with accession number CNP0001046.

Statistical Analysis

The OTU numbers, types and taxonomic features of the samples of the three groups, i.e., those sampled on

April 8 and 9, June 25 and 26 and September 19 and 20, 2019, were analyzed. Community richness (Chao

richness index) was examined to reveal the variation among the three groups. The WBIF (including the

WBIF from the upstream to downstream regions and the WBIF from land to river) of each group was

assessed to reveal the effectiveness of monitoring the biodiversity information of the upstream and riparian

zones using riverine water eDNA. The analysis of the WBIF follows the processing approach proposed by

Yang et al. (2019). In the analysis of the WBIF, all statistics used the types of OTUs of each sample rather

than the numbers of OTUs of each sample. The processing approach can be described simply as follows.

As the WBIF driven by watershed ecosystem processes was transported from land to river and from

upstream to downstream regions, the transportation effectiveness of the WBIF could be estimated by

comparing the OTU assemblages between adjacent soil eDNA samples and water eDNA samples and by

comparing the OTU assemblages between two adjacent water eDNA samples. The transportation

effectiveness of the WBIF was indicated by the proportion of input OTU types (i.e., the common types

between the source site sample and the pool site sample) to output OTU types (the total types of source site

sample) (Eq. 1).

$$e = \mathrm{Num}(S_{OTU} \cap P_{OTU})/\mathrm{Num}(S_{OTU}) \qquad \text{(Eq. 1)}$$

where e denotes the transportation effectiveness of the WBIF; S_{OTU} denotes the OTU assemblage of the

source site sample (i.e., the adjacent soil eDNA sample in the land-river WBIF or the adjacent upstream

water eDNA sample in the upstream-downstream WBIF); and P_{OTU} denotes the OTU assemblage of the

pool site sample (i.e., the adjacent water eDNA sample in the land-river WBIF or the adjacent downstream

water eDNA sample in the upstream-downstream WBIF).

As the transportation effectiveness of the WBIF relied on transport capacity, degradation rate and

environmental filtration and the distance of the land-to-river WBIF was less than 5 m, the transportation

effectiveness of the land-to-river WBIF was assumed to be constructed by transport capacity and

environmental filtration. The transportation effectiveness of the land-to-river WBIF could be indicated by

the proportion of the common types shared between adjacent soil eDNA samples and water eDNA samples

to the total types of soil eDNA samples (Eq. 1). The transport capacity of the land-to-river WBIF could be

indicated by the proportion of the common types shared between adjacent soil eDNA samples and water

eDNA samples to the common types shared between the soil eDNA sample and all water eDNA samples in

the corresponding group (Eq. 2). The environmental filtration of the land-to-river WBIF could be indicated

by the proportion of the types included in the soil eDNA sample but not in any water eDNA sample to the

total types in the soil eDNA sample (Eq. 3).

$$t = \mathrm{Num}(S_{OTU} \cap P_{OTU})/\mathrm{Num}(S_{OTU} \cap W_{OTU}) \quad \text{(Eq. 2)}$$

$$f = 1 - \mathrm{Num}(S_{OTU} \cap W_{OTU})/\mathrm{Num}(S_{OTU}) \quad \text{(Eq. 3)}$$

where t denotes the transport capacity; f denotes the environmental filtration; S_{OTU} denotes the OTU

assemblage of the source site sample (i.e., the soil eDNA sample); and W_{OTU} denotes the OTU assemblage

of all water eDNA samples.

The upstream-to-downstream WBIF that is indicated by environmental microbes includes the effective

WBIF (i.e., the flow of living organisms) and the noneffective WBIF (i.e., the flow of dead organisms).

The effective WBIF was impacted by transport capacity and environmental filtration. The noneffective

WBIF was impacted by transport capacity and degradation rate. We established the following assumptions:

the transport capacity was consistent in a defined runoff condition; the proportion of noneffective WBIF at

each site was consistent; the noneffective WBIF degraded over time (i.e., distance) in a logistic manner;

and the environmental filtration was consistent in a definite environmental change. The transportation

effectiveness of the upstream-to-downstream WBIF could be constructed by the transport capacity of the

WBIF, the environmental filtration of the effective WBIF and the degradation rate of the noneffective

WBIF (Eq. 4). In practice, as watershed ecosystem processes are impacted by varied influencing factors at

any site and time, the analytical solution of Eq. 4 is impossible. Therefore, Eq. 4 will be

programming-solved according to the evolutionary algorithm.

$$e = (t\char`^d)[(1-k)(1-f) + k(1/2)\char`^(d/D)] \qquad \text{(Eq. 4)}$$

where e denotes the transportation effectiveness of the WBIF; t denotes the transport capacity; d denotes

the distance of the WBIF; k denotes the proportion of the noneffective WBIF; f denotes the environmental

filtration; and D denotes the half-life distance.

Results

Biological Information Features of the Samples of the Three Groups

A total of 1,030,826, 968,122 and 842,317 clean sequences were obtained, respectively, from the 18

samples of the spring group, summer group and autumn group (more details in Fig. 2a), and the average

lengths of these sequences were 447.54 bp, 416.06 bp, and 416.32 bp, respectively. A total of 10,602,

13,766 and 16,500 types of bacterial OTUs were detected (UPARSE, 97% cutoff), respectively, from the

18 samples of the three groups (more details in Fig. 2b and Fig. 3), which belonged to 58 phyla, 141

classes, 424 orders, 782 families, 1,895 genera and 4,537 species. The OTU counts and compositions of

each group were highly variable (Fig. 2b and Fig. 2c). The rarefaction curves of each sample showed that

the number of clean sequences of each sample in five groups (except for the spring water eDNA samples)

was lower than it should be, especially for the water eDNA samples in summer and autumn (Fig. 3).

The total types of OTUs detected in the soil eDNA samples estimated using the species accumulation curve

were similar among the spring, summer and autumn groups (Fig. 2d), although there were some spatially

and temporally heterogeneous OTU compositions (Fig. 2c and Fig. 4). The total types of OTUs detected in

the water eDNA samples estimated using the species accumulation curve showed that the richest OTU

types occurred in autumn (Fig. 2d). Moreover, the temporal heterogeneity of OTU composition among the

three groups was obvious (Fig. 4). The common OTUs shared between the soil and water eDNA samples

totaled 2,834, 5,651 and 5,279 in the spring group, summer group and autumn group, respectively, which

accounted for 36.30%, 71.98% and 67.58% of the total types of OTUs detected in the soil eDNA samples

of each group, respectively.

Land-river WBIF Features of the Three Groups

The transportation effectiveness of the WBIF from the riparian zone to the river was 16.62%

(0.1662 ± 0.1254, 95%) in the frozen spring, 62.76% (0.6276 ± 0.0873, 95%) in the rainy summer, and 48.09%

(0.4809 ± 0.0522, 95%) in the cloudy autumn. The transport capacity of the WBIF from the riparian zone to

the river was 26.88% (0.2688 ± 0.2024, 95%) in the frozen spring, 68.49% (0.6849 ± 0.0913, 95%) in the

rainy summer, and 57.36% (0.573579 ± 0.052897, 95%) in the cloudy autumn. The environmental filtration

of the WBIF from the riparian zone to the river was 38.54% (0.3854 ± 0.0293, 95%) in the frozen spring,

8.38% (0.0838 ± 0.0206, 95%) in the rainy summer, and 16.18% (0.1618 ± 0.0451, 95%) in the cloudy

autumn. More details are shown in Table 1.

During the spring sampling period, the ice water in the estuary (transect SL1) covered the riparian

grassland. Because the water sample was likely to be the eluate from the soil at SL1, the transportation

effectiveness and transport capacity of the WBIF from the riparian zone to the river were high. As Qinghai

Lake is saline, the environmental filtration of the WBIF from the riparian zone to the river in the estuary

(SL1) was high. During the summer sampling period, because the samples from SL1 and SL2 were

sampled on sunny days and the other samples were sampled on rainy days, the transport capacity of the

WBIF from the riparian zone was low, the environmental filtration of the WBIF from the riparian zone was

high, and the transportation effectiveness of the WBIF from the riparian zone was low at SL1 and SL2.

During the autumn sampling period, because the samples from SL1 and SL2 were sampled on rainy days

and the other samples were sampled on cloudy days, the transport capacity of the WBIF from the riparian

zone was high at SL2. Because of the environmental stress caused by estuary saline water, the

environmental filtration of the WBIF from the riparian zone was high at SL1.

Upstream-downstream WBIF Features of the Three Groups

Along with the flow from upstream to downstream, there were 4 chains that connected the sampling

transects. Their cumulative transportation effectiveness of the upstream-to-downstream WBIF varied with

cumulative distance (Table 2). The transportation effectiveness of the upstream-to-downstream WBIF

relied on transport capacity, degradation rate and environmental filtration. In spring, the transport capacity

of the upstream-to-downstream WBIF was 99.97% (0.9997 ± 0.0003, 95%) per km, of which there was

66.85% (0.6685 ± 0.0034, 95%) noneffective WBIF, the half-life distance of the noneffective WBIF was

1.55 (1.5490 ± 0.1269, 95%) km, and the environmental filtration from SL2 to SL1 (from the freshwater

ecosystem to saline water ecosystem) was 16.04% (0.1604 ± 0.0082, 95%). In summer, the transport

capacity of the upstream-to-downstream WBIF was 99.42% (0.9942 ± 0.0009, 95%) per km, of which there

was 43.46% (0.4346 ± 0.0417, 95%) noneffective WBIF, the half-life distance of the noneffective WBIF

was 14.52 (14.5234 ± 1.4405, 95%) km, the environmental filtration from SL3 to SL2 (from rainy sampling

conditions to sunny sampling conditions) was 0.57% (0.0057 ± 0.0055, 95%) and the environmental

filtration from SL2 to SL1 (from the freshwater ecosystem to saline water ecosystem) was 54.42%

(0.5442 ± 0.0100, 95%). In autumn, the transport capacity of the upstream-to-downstream WBIF was 99.23%

(0.9923 ± 0.0016, 95%) per km, of which there was 49.35% (0.4935 ± 0.0410, 95%) noneffective WBIF, the

half-life distance of the noneffective WBIF was 10.40 (10.3981 ± 0.7111, 95%) km, and the environmental

filtration from SL2 to SL1 (from the freshwater ecosystem to saline water ecosystem) was 12.87%

(0.1287 ± 0.0171, 95%). The transportation effectiveness of the upstream-to-downstream WBIF was

75.86%, 97.41% and 96.07% in spring, summer and autumn, respectively, without regard to environmental

filtration in defined areas and times, with environmental change.

Discussion

Effectiveness of Monitoring Riparian Biodiversity Information Using Riverine Water eDNA

Because riverine water eDNA incorporates terrestrial biodiversity information (Deiner, et al.,2016; Yang,

et al.,2019), by analyzing riverine water eDNA, information can be obtained on terrestrial biological

compositions (Deiner, et al.,2016; Matsuoka, et al.,2019). The effectiveness of monitoring riparian

biodiversity information depends on the transportation effectiveness of the WBIF from the riparian zone to

the river. Following the WBIF framework (Yang, et al.,2019), the transportation effectiveness of the WBIF

from the riparian zone to the river is mainly constructed by the capacity of transporting eDNA in the

riparian zone into the river (driven by runoff, animals, wind, etc.) and environmental filtration (driven by

soil/water environmental shifts, soil aggregates, etc.). Therefore, the effectiveness of monitoring riparian

biodiversity information is also impacted by these factors.

In the frozen spring period, the maximum effectiveness of monitoring riparian biodiversity information

was 16.62%, as the transportation effectiveness of the WBIF from the riparian zone to the river was 16.62%

(estimated based on the OTU compositions detected from the soil eDNA sample and the water eDNA

sample at the same sampling transect). In other words, a water eDNA sample could include information on

as much as 16.62% of the biodiversity information from the adjacent riparian zone. Moreover, nearly 36.30%

of the types of OTUs detected from the soil eDNA samples could be detected in water eDNA samples. This

result suggested that as much as 36.30% of the riparian zone biodiversity information could be monitored

using water eDNA samples. This effectiveness was mainly controlled by the low transport capacity of the

WBIF (26.88%) and by the high environmental filtration (38.54%).

In the rainy summer period, the maximum effectiveness of monitoring the riparian biodiversity information

was 62.76%, as the transportation effectiveness of the WBIF from the riparian zone to the river was 62.76%

(estimated based on the OTU compositions detected from the soil eDNA sample and the water eDNA

sample at the same sampling transect). In other words, a water eDNA sample could monitor as much as

62.76% of biodiversity information from the adjacent riparian zone. Moreover, nearly 71.98% of the types

of OTUs detected from the soil eDNA samples could be detected in the water eDNA samples. This result

suggested that as much as 71.98% of the biodiversity information in the riparian zone could be monitored

using water eDNA samples. This effectiveness was mainly controlled by the transport capacity of the

WBIF (68.49%).

In the cloudy autumn period, the maximum effectiveness of monitoring riparian biodiversity information

was 48.09%, as the transportation effectiveness of the WBIF from the riparian zone to the river was 48.09%

(estimated based on the OTU compositions detected from the soil eDNA sample and the water eDNA

sample at the same sampling transect). In other words, a water eDNA sample could monitor as much as

48.09% of the biodiversity information from the adjacent riparian zone. Moreover, nearly 67.58% of the

types of OTUs detected from the soil eDNA samples could be detected in the water eDNA samples. This

result suggested that as much as 67.58% of the biodiversity information in the riparian zone could be

monitored using water eDNA samples. This effectiveness was mainly controlled by the transport capacity

of the WBIF (57.36%) and by environmental filtration (16.18%).

The transport capacity of the WBIF from the riparian zone to the river was driven by animal movement,

wind erosion and transport, rainfall and surface runoff erosion, and surface runoff transport (Yang, et

al.,2019). In the frozen spring period, there was little animal movement, as most animals were hibernating

during Qinghai-Tibet Plateau's long winter. Although there was strong wind, the frozen soil limited the

amount of wind erosion and transport. Moreover, the precipitation form was snow, and there was little

surface runoff, which provided suboptimal conditions for significant erosion and transport. In summer and

autumn, animal movement was frequent, wind erosion and transport were permitted, and rainfall and

surface runoff erosion and surface runoff transport were significant. Therefore, the transport capacity of the

WBIF from the riparian zone to the river in spring was obviously lower than that in summer and autumn

and that on sunny or cloudy days was obviously lower than that on rainy days.

The environmental filtration of the WBIF was impacted by soil-to-water environmental shifts and soil

aggregates (Wilpiszeski, et al.,2019; Yang, et al.,2019). In the frozen spring period, the microbial activities

were low in both soil and water, and most microbes (input to rivers from soil) could not be effectively

preserved in water. Because there was no significant surface runoff, there was no soil aggregate to

conserve the microbial input to the river. In summer and autumn, the microbial activities were high in both

soil and water, and most microbes (input to rivers from soil) could be effectively preserved in water.

Rainfall and surface runoff promoted the input of soil aggregates into rivers and promoted the conservation

of the microbes living in them. Therefore, the environmental filtration of the WBIF in spring was

obviously higher than that in summer and autumn and that on sunny or cloudy days was obviously higher

than that on rainy days.

Sales et al. (2020) indicated that although the detection probability of riverine water eDNA was 40%~67, it

provided comparable results to conventional survey methods per unit of survey effort for three species

(water vole, field vole and red deer); in other words, the results from 3~6 water replicates would be

equivalent to the results from 3~5 latrine surveys and 5~30 weeks of single camera deployment (Sales, et

al.,2020). Considering that the number of clean sequences of each summer water eDNA sample and

autumn water eDNA sample is obviously lower than it should be (Fig. 3), it is probable that the water

eDNA samples contain more types of OTUs than the current results indicate. In other words, the

transportation effectiveness of the WBIF from the riparian zone to the river in summer and autumn was

obviously underestimated; therefore, the effectiveness of monitoring riparian biodiversity information in

summer and autumn was obviously underestimated. However, the monitoring effectiveness was still

62%~72%. This result suggests that riverine water eDNA is viable for monitoring riparian biodiversity

information on rainy days in summer or autumn.

Effectiveness of Monitoring Upstream Biodiversity Information Using Downstream Water eDNA

The fact that species information at some distance upstream could be detected is the key for eDNA

application in lotic systems (Stoeckle, et al.,2016; Carraro, et al.,2018; Pont, et al.,2018). The effectiveness

of monitoring upstream biodiversity information depends on the transportation effectiveness of the

upstream-to-downstream WBIF (Deiner & Altermatt,2014; Sansom & Sassoubre,2017; Pont, et al.,2018;

Seymour,2019; Yang, et al.,2019). eDNA could originate from living and dead organisms and could be

detected at distances downstream, which determined the eDNA transport and degradation processes

(Stoeckle, et al.,2016; Nukazawa, et al.,2018; Tillotson, et al.,2018; Yang, et al.,2019). Following the

framework of the WBIF (Yang, et al.,2019), the transportation effectiveness of the

upstream-to-downstream WBIF was mainly constructed by the transport capacity of the

upstream-to-downstream WBIF (impacted by eDNA evenness dispersed in water) and the degradation rate

(constructed by the proportion and the half-life distance of noneffective WBIF). Therefore, the

effectiveness of monitoring upstream biodiversity information was also impacted by these factors.

In spring, the maximum effectiveness of biodiversity information monitoring 1 km upstream was 75.86%,

as the transportation effectiveness of the upstream-to-downstream WBIF was 75.86% per km. This result

suggested that a water eDNA sample could monitor as much as 75.86% of the biodiversity information 1

km upstream. This effectiveness was mainly controlled by the high proportion of the noneffective WBIF

(66.85%) and the small half-life distance of the noneffective WBIF (1.55 km).

In summer, the maximum effectiveness of biodiversity information monitoring 1 km upstream is 97.41%,

as the transportation effectiveness of the upstream-to-downstream WBIF is 97.41% per km. This result

suggested that a water eDNA sample could monitor as much as 97.41% of the biodiversity information 1

km upstream. This effectiveness was mainly controlled by the proportion of the noneffective WBIF

(43.46%) and its half-life distance (14.52 km).

In autumn, the maximum effectiveness of biodiversity information monitoring 1 km upstream was 96.07%,

as the transportation effectiveness of the upstream-to-downstream WBIF was 96.07% per km. This result

suggested that a water eDNA sample could monitor as much as 96.07% of the biodiversity information 1

km upstream. This effectiveness was mainly controlled by the proportion of the noneffective WBIF

(49.35%) and its half-life distance (10.40 km).

The transport capacity of the upstream-to-downstream WBIF was impacted by the even dispersal of eDNA

in water. When the community richness is too high to spread each type of OTU in every liter of water, the

transport capacity of the upstream-to-downstream WBIF declines. Therefore, along with the increase in the

types of bacterial OTUs from spring to summer to autumn (Fig. 2d), the transport capacity declined from

spring to summer to autumn. Fortunately, in the Shaliu River basin, the transport capacity was high (more

than 99%) in the three seasons.

The degradation rate in the present study was constructed by the proportion and the half-life distance of the

noneffective WBIF. The proportion of noneffective WBIF relies on microbial activity, which is impacted

by temperature (Lin, et al.,2016). Therefore, the highest proportion of noneffective WBIF was detected in

the frozen spring period, and the lowest proportion as detected in summer. The half-life distance of

noneffective WBIF was mainly impacted by the flow rate of river runoff and the half-life period of eDNA

degradation (Yang, et al.,2019). The half-life period of eDNA degradation is impacted by many

environmental conditions, such as biochemical oxygen demand, temperature, pH, and organic matter

(Barnes, et al.,2014; Eichmiller, et al.,2016; Nukazawa, et al.,2018; Seymour, et al.,2018; van Bochove, et

al.,2020). However, the highest flow rate and the largest half-life distance of noneffective WBIF are found

in summer. In other words, the half-life distance of the noneffective WBIF was mainly controlled by the

flow rate in the Shaliu River.

The number of clean sequences of each water eDNA sample in summer and autumn was obviously lower

than it should be (Fig. 3), and this result indicated that the water eDNA samples contained more types of

OTUs than the current results presented; in other words, the transportation effectiveness of the

upstream-to-downstream WBIFs in summer and autumn were obviously underestimated. Therefore, the

effectiveness of monitoring upstream biodiversity information in summer and autumn was also obviously

underestimated. However, the monitoring effectiveness was greater than 96% 1 km downstream in summer

and autumn. This result suggested that riverine water eDNA was viable for monitoring upstream

biodiversity information in summer and autumn.

Cost-effective Proposal for Monitoring the Biodiversity Information of Upstream and Riparian

Zones Using Water eDNA

Because eDNA can be used to assess diversity across the tree of life with a single-field sampling protocol

(Deiner, et al.,2016), eDNA sampling is more cost-effective (including cost and time) than conventional

biodiversity monitoring methods (Mächler, et al.,2014; Valentini, et al.,2016; Seymour,2019). Moreover,

simultaneously monitoring aquatic and terrestrial biodiversity using riverine water eDNA represents one

step toward   obtaining a more cost-effective biodiversity monitoring method.

As the highest effectiveness of monitoring riparian biodiversity information (62.76% for adjacent sites, or

71.98% for overall sites) appeared on rainy days in summer, the cost-effective sampling time for

monitoring riparian biodiversity information using water eDNA is a rainy day in summer. Considering that

the effectiveness was mainly impacted by rainfall and surface runoff, a rainy day in autumn would also be

a cost-effective sampling time. Considering the temporal heterogeneity of biodiversity information,

although there was variation in the OTU composition (Fig. 4), we do not suggest monitoring riparian

biodiversity information using water eDNA during the frozen spring period because the monitoring

effectiveness would be too low to detect the different biodiversity information in spring from that in

summer and autumn. If sampling is required in the frozen spring period, it would be better to monitor

riparian biodiversity information using soil eDNA directly.

As the effectiveness of biodiversity information monitoring 1 km upstream was 97.41% in summer, 96.07%

in autumn, and higher than 75.86% in spring, the cost-effective sampling time for monitoring riparian

biodiversity information using water eDNA is summer or autumn. In term of the temporal heterogeneity of

biodiversity information, the water samples in autumn had the highest community richness, followed by

those collected in summer and spring (Figs. 2, 3). Considering the degradation of noneffective WBIF, the

flow rate mainly controlled the half-life distance of noneffective WBIF. We suggest that, to monitor

microbial biodiversity information, autumn is the first choice for sampling, followed by summer because

of the relatively high flow rate. Because there were low monitoring effectiveness and high OTU

composition differences with other seasons in spring, the microbial biodiversity in spring should be

monitored with a high sampling site density, if spring sampling is required.

After examining the rarefaction curves of each sample, it was obvious that the number of clean sequences

of each sample in the five groups (except for the spring water eDNA samples) was lower than it should be

(Fig. 3). Perhaps more than 60,000 clean sequences of each sample or more than one duplicate sample is

needed, especially for the water eDNA samples in summer and autumn. Therefore, to monitor the

biodiversity information of upstream and riparian zones using riverine water eDNA, a rainy day in autumn

or summer would be the most cost-effective sampling time, and more than one duplicate sample would be

a better sampling choice for water eDNA samples in summer and autumn. If sampling is needed in spring,

riparian biodiversity information should be monitored by soil eDNA rather than by water eDNA, and

aquatic biodiversity information should be monitored using a high-density sampling method. All

biodiversity assessments based on the monitoring results using riverine water eDNA should be revalued

based on the monitoring effectiveness.

Moreover, the monitoring effectiveness in this study is indicated by environmental microbes, and whether

the effectiveness of monitoring other taxonomies using riverine water eDNA is higher or lower is still

unknown, and more research is required. Although the seasonal variation in monitoring effectiveness was

delineated based on the results indicated by environmental microbes, it would be effective for other

taxonomic groups. In other words, a rainy day in autumn or summer is the most cost-effective sampling

time for monitoring the biodiversity information of upstream and riparian zones using riverine water eDNA

for other taxonomic groups. Therefore, a study on monitoring the effectiveness assessment of other

taxonomic groups would be useful for implementation on rainy days in autumn or summer to assess a

cost-effective monitoring proposal. If the monitoring effectiveness in all taxonomic groups was verified to

be higher than that obtained using conventional methods, the claim of revolutionizing biodiversity science

(Thomsen & Willerslev,2015; Cristescu & Hebert,2018; Altermatt, et al.,2020) would come true.

The results showed that the effectiveness of monitoring the biodiversity information of the upstream and

riparian zones was 96%~97% (1 km upstream) and 62%~72% (5 m distance from the river) on rainy days

in summer or autumn, respectively, which was investigated in the Shaliu River basin (a typical watershed

on the Qinghai-Tibet Plateau), and this information is likely a useful reference for other watersheds on the

Qinghai-Tibet Plateau. Moreover, the following points may be applicable to watersheds in other regions: (1)

rainfall, surface runoff and animal movements increased the effectiveness of monitoring riparian

biodiversity information using riverine water eDNA; (2) frozen soil limited the effectiveness of monitoring

riparian biodiversity information using riverine water eDNA; and (3) a high flow rate increased the

effectiveness of monitoring upstream biodiversity information using downstream water eDNA. We believe

that our framework for assessing monitoring effectiveness is reasonable and general. We encourage more

research on monitoring effectiveness in other watersheds with different climatic conditions to support

simultaneous aquatic and terrestrial biodiversity assessments.

Acknowledgements

493   This work was supported by the Central Public-Interest Scientific Institution Basal Research Fund (Grant

494   No. 2019HY-XKQ02, 2020TD08), the Basic Research Program of Science and Technology Department

495   Qinghai Provence (Grant No. 2018-ZJ-703), the Provincial Natural Science Foundation of Qinghai (Grant

496   No. 2018-ZJ-908).

References

Altermatt F, Little CJ, Mächler E, *et al.* (2020) Uncovering the complete biodiversity structure in spatial
networks: the example of riverine systems. *Oikos*.
Anderson CB (2018) Biodiversity monitoring, earth observations and the ecology of scale. *Ecology Letters*
21, 1572-1585.
Barnes MA, Turner CR (2016) The ecology of environmental DNA and implications for conservation
genetics. *Conservation Genetics* **17**, 1-17.
Barnes MA, Turner CR, Jerde CL*, et al.* (2014) Environmental Conditions Influence eDNA Persistence in
Aquatic Systems. *Environmental Science & Technology* **48**, 1819-1827.
Cardinale BJ, Duffy JE, Gonzalez A*, et al.* (2012) Biodiversity loss and its impact on humanity. *Nature* **486**,
59-67.
Carraro L, Hartikainen H, Jokela J, Bertuzzo E, Rinaldo A (2018) Estimating species distribution and
abundance in river networks using environmental DNA. *Proceedings of the National Academy of*
Sciences of the United States of America **115**, 11724-11729.
Carrizo SF, Lengyel S, Kapusi F*, et al.* (2017) Critical catchments for freshwater biodiversity conservation

in Europe: identification, prioritisation and gap analysis. *Journal of Applied Ecology* **54**, 1209-1218.

Chase JM, McGill BJ, McGlinn DJ*, et al.* (2018) Embracing scale-dependence to achieve a deeper understanding of biodiversity and its change across communities. *Ecology Letters* **21**, 1737-1751.

Cristescu ME, Hebert PDN (2018) Uses and Misuses of Environmental DNA in Biodiversity Science and Conservation. *Annual Review of Ecology, Evolution, and Systematics* **49**, 209-230.

Deiner K, Altermatt F (2014) Transport Distance of Invertebrate Environmental DNA in a Natural River. *PLoS ONE* **9**, e88786.

Deiner K, Fronhofer EA, Mächler E, Walser J, Altermatt F (2016) Environmental DNA reveals that rivers are conveyer belts of biodiversity information. *Nature Communications* **7**, 12544.

Eichmiller JJ, Best SE, Sorensen PW (2016) Effects of Temperature and Trophic State on Degradation of Environmental DNA in Lake Water. *Environmental Science & Technology* **50**, 1859-1867.

Hooper DU, Adair EC, Cardinale BJ*, et al.* (2012) A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* **486**, 105-108.

Jetz W, Cavender-Bares J, Pavlick R*, et al.* (2016) Monitoring plant functional diversity from space. *Nature Plants* **2**, 16024.

Lin Q, De Vrieze J, Li J, Li X (2016) Temperature affects microbial abundance, activity and interactions in anaerobic digestion. *Bioresource Technology* **209**, 228-236.

Lugg WH, Griffiths J, van Rooyen AR, Weeks AR, Tingley R (2018) Optimal survey designs for environmental DNA sampling. *Methods in Ecology and Evolution* **9**, 1049-1059.

Luo Y, Xu L, Rysz M*, et al.* (2011) Occurrence and Transport of Tetracycline, Sulfonamide, Quinolone, and Macrolide Antibiotics in the Haihe River Basin, China. *Environmental Science & Technology* **45**, 1827-1833.

M L, XJ S, WJ W*, et al.* (2020) Studying the retention time of Fenneropenaeus chinensis eDNA in water. *Progress in Fishery Sciences* **41**, 51-57.

Mächler E, Deiner K, Steinmann P, Altermatt F (2014) Utility of environmental DNA for monitoring rare and indicator macroinvertebrate species. *Freshwater Science* **33**, 1174-1183.

Mallick PH, Chakraborty SK (2018) Forest, wetland and biodiversity: Revealing multi-faceted ecological services from ecorestoration of a degraded tropical landscape. *Ecohydrology & Hydrobiology* **18**, 278-296.

Matsuoka S, Sugiyama Y, Sato H*, et al.* (2019) Spatial structures of fungal DNA assemblages revealed with eDNA metabarcoding in a forest river network in western Japan. *bioRxiv*, 637686.

McGlinn DJ, Xiao X, May F*, et al.* (2019) Measurement of Biodiversity (MoB): A method to separate the scale-dependent effects of species abundance distribution, density, and aggregation on diversity change. *Methods in Ecology and Evolution* **10**, 258-269.

Nukazawa K, Hamasuna Y, Suzuki Y (2018) Simulating the Advection and Degradation of the Environmental DNA of Common Carp along a River. *Environmental Science & Technology* **52**, 10562-10570.

Pont D, Rocle M, Valentini A*, et al.* (2018) Environmental DNA reveals quantitative patterns of fish biodiversity in large rivers despite its downstream transportation. *Scientific Reports* **8**, 10313-10361.

Sales NG, McKenzie MB, Drake J*, et al.* (2020) Fishing for mammals: Landscape-level monitoring of terrestrial and semi-aquatic communities using eDNA from riverine systems. *Journal of Applied Ecology* **57**, 707-716.

Sansom BJ, Sassoubre LM (2017) Environmental DNA (eDNA) Shedding and Decay Rates to Model

Freshwater Mussel eDNA Transport in a River. *Environmental Science & Technology* **51**, 14244-14253.

Seymour M (2019) Rapid progression and future of environmental DNA research. *Communications*
Biology **2**, 80.

Seymour M, Durance I, Cosby BJ*, et al.* (2018) Acidity promotes degradation of multi-species
environmental DNA in lotic mesocosms. *Communications Biology* **1**, 4.

Shogren AJ, Tank JL, Andruszkiewicz E*, et al.* (2017) Controls on eDNA movement in streams: Transport,
Retention, and Resuspension. *Scientific Reports* **7**, 5065.

Stat M, Huggett MJ, Bernasconi R*, et al.* (2017) Ecosystem biomonitoring with eDNA: metabarcoding
across the tree of life in a tropical marine environment. *Scientific Reports* **7**, 12240.

Stoeckle BC, Kuehn R, Geist J (2016) Environmental DNA as a monitoring tool for the endangered
freshwater pearl mussel (Margaritifera margaritifera L.): a substitute for classical monitoring approaches?
Aquatic Conservation: Marine and Freshwater Ecosystems **26**, 1120-1129.

Thomsen PF, Willerslev E (2015) Environmental DNA – An emerging tool in conservation for monitoring
past and present biodiversity. *Biological Conservation* **183**, 4-18.

Tillotson MD, Kelly RP, Duda JJ*, et al.* (2018) Concentrations of environmental DNA (eDNA) reflect
spawning salmon abundance at fine spatial and temporal scales. *Biological Conservation* **220**, 1-11.

Valentini A, Taberlet P, Miaud C*, et al.* (2016) Next‐generation monitoring of aquatic biodiversity using
environmental DNA metabarcoding. *Molecular Ecology* **25**, 929-942.

van Bochove K, Bakker FT, Beentjes KK*, et al.* (2020) Organic matter reduces the amount of detectable
environmental DNA in freshwater. *Ecology and Evolution* **10**, 3647-3654.

Vina A, Chen X, Yang W*, et al.* (2013) Improving the efficiency of conservation policies with the use of
surrogates derived from remotely sensed and ancillary data. *Ecological Indicators* **26**, 103-111.

Wang S, Fu B, Piao S*, et al.* (2016) Reduced sediment transport in the Yellow River due to anthropogenic
changes. *Nature Geoscience* **9**, 38-41.

Wilpiszeski RL, Aufrecht JA, Retterer ST*, et al.* (2019) Soil aggregate microbial communities: towards
understanding microbiome interactions at biologically relevant scales. *Applied and Environmental*
Microbiology **85**, e319-e324.

Yang H, Du H, Qi H, Yu L, Wei Q (2019) Quantifying the watershed biological information flow driven by
runoff based on the indicator of environmental microbes. *ChinaXiv* **201911**, 75.

Data Accessibility

588 The datasets generated for this study can be found in the CNSA (https://db.cngb.org/cnsa/) of CNGBdb

589 with accession number CNP0001046.

Author Contributions

HY acquired fund, designed and performed research, analyzed data, wrote and edited the paper.

HD acquired fund, designed and supervised research, reviewed and edited the paper.

HQ acquired fund, designed research, reviewed and edited the paper.

LY performed field sampling.

XH performed laboratory experiment.

HZ, JL, JW and CW reviewed and edited the paper.

QZ administrated project.

QW supervised and validated research.

Tables and Figures (with captions)

Fig. 1 Sampling transects.

SL1 denotes the first sampling transect on the Shaliu River. The distances labeled in parentheses under the tags of sampling transects denote the distances from the estuary to the sampling transects, such as SL1 (1.8 km), which means the distance from the estuary to SL1 is 1.8 km.

Fig. 2 Biological information features of the samples: numbers of clean sequences in each sample (a), types of OTUs in each sample (b), community richness of each sample at the OTU level (c) and species accumulation curves at the OTU level (d).

201904S denotes the soil eDNA samples that were sampled during April 2019; 201909W denotes the water eDNA samples that were sampled during September 2019.

Fig. 3 Rarefaction curves of each sample.

201904S denotes the soil eDNA samples that were sampled during April 2019; 201909W denotes the water
eDNA samples that were sampled during September 2019.

Fig. 4 The OTU types in riparian soil samples and riverine water samples shared by three groups.
201904S denotes the soil eDNA samples that were sampled during April 2019; 201909W denotes the water
eDNA samples that were sampled during September 2019.

Table 1 Transport capacity, environmental filtration and transportation effectiveness of watershed biological information flow (WBIF) from the riparian zone to the river in each sampling transect

	201904			201906			201909		
	Transport capacity	**Environmental filtration**	**Transportation effectiveness**	**Transport capacity**	**Environmental filtration**	**Transportation effectiveness**	**Transport capacity**	**Environmental filtration**	**Transportation effectiveness**
S-W_SL6b	0.099499	0.414501	0.058256	0.504529	0.064307	0.472084	0.566649	0.138540	0.488145
S-W_SL6	0.859828	0.383092	0.530435	0.756467	0.048747	0.719591	0.655770	0.153814	0.554904
S-W_SL5m	0.221932	0.311564	0.152786	0.776533	0.080882	0.713725	0.475058	0.199631	0.380221
S-W_SL5	0.226740	0.360500	0.145000	0.655574	0.109190	0.583992	0.597921	0.254486	0.445759
S-W_SL4b	0.175403	0.401328	0.105009	0.667238	0.107198	0.595712	0.539223	0.121664	0.473619
S-W_SL4	0.184110	0.388603	0.112564	0.784735	0.080409	0.721635	0.570324	0.120697	0.501487
S-W_SL3	0.205840	0.409836	0.121480	0.649059	0.095981	0.586761	0.610110	0.143771	0.522394
S-W_SL2	0.176972	0.414116	0.103685	0.568211	0.222821	0.441602	0.758528	0.269508	0.554098
S-W_SL1	0.428305	0.503238	0.212766	0.257329	0.216037	0.201736	0.579158	0.470276	0.306794

S-W denotes the WBIF from the soil eDNA sample (riparian zone) to the water eDNA sample (river). SL1, SL2, SL3, SL4, SL4b, SL5m, SL5, SL6 and SL6b denote the sampling transections. S-W_SL1 denotes the WBIF from the riparian zone to the river at SL1. 201904, 201906, and 201909 indicate the groups that were sampled in April, June and September 2019, respectively. Transport capacity indicates the proportion of soil microbes that were transported into the river. Environmental filtration indicates that the proportion of soil microbes that could not be kept in rivers. Transportation effectiveness indicates the proportion of soil microbes that could be detected in rivers.

Table 2 The accumulative runoff distance and accumulative transportation effectiveness from the
first sampling transection in each chain of the watershed biological information flow (WBIF)

Transport Chain	Cumulative Distance/km	Cumulative Effectiveness		
		201904	201906	201909
Chain A				
SL6b-SL5	9	0.521352	0.798283	0.598885
##-SL4	24.5	0.412811	0.534830	0.471711
##-SL3	48	0.368327	0.486959	0.415639
##-SL2	63	0.330961	0.383295	0.308563
##-SL1	70	0.119217	0.197425	0.257389
Chain B				
SL6-SL5	8	0.217131	0.762412	0.605484
##-SL4	23.5	0.111554	0.590463	0.469171
##-SL3	47	0.093625	0.517251	0.408623
##-SL2	62	0.067231	0.414025	0.299731
##-SL1	69	0.032371	0.192146	0.248692
Chain C				
SL5m-SL4	8.5	0.430074	0.765880	0.667929
##-SL3	32	0.334385	0.579820	0.540136
##-SL2	47	0.228181	0.448004	0.361037
##-SL1	54	0.072555	0.196740	0.298750
Chain D				
SL4b-SL3	23	0.521333	0.738029	0.719376
##-SL2	38	0.352000	0.555864	0.437211
##-SL1	45	0.108000	0.237335	0.335324

Chains A, B, C and D denote the 4 chains of the upstream-to-downstream WBIF. In Chain A, SL6b-SL5
denotes the WBIF from sampling transects SL6b to SL5; ##-SL4 denotes the WBIF from sampling
transects SL6b to SL5 and then to SL4; ##-SL3 denotes the WBIF from sampling transects SL6b to SL5,
SL4 and then to SL3; ##-SL2 denotes the WBIF from sampling transects SL6b to SL5, SL4, SL3 and then
to SL2; ##-SL1 denotes the WBIF from sampling transects SL6b to SL5, SL4, SL3, SL2 and then to SL1
(estuary). 201904, 201906, and 201909 indicate the groups that were sampled in April, June and September
2019, respectively.