1

Show me your Dufour's gland and I'll tell you who you are: caste and physiology

specific signals in *Bombus impatien*s

Nathan T. Derstine[1*], Gabriel Villar[1,] Abraham Hefetz[2,3], Jocelyn Millar[4], Etya Amsalem[1]

[1]Department of Entomology, Center for Chemical Ecology, Center for Pollinator Research, Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, PA 16802 U.S.A.

[2]School of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Israel

[3]School of Marine Sciences, Ruppin Academic Center, Israel

[4]Departments of Entomology and Chemistry, University of California Riverside, CA 92521

[*] Corresponding author

Abstract

Reproductive division of labor in insect societies is regulated through multiple concurrent

mechanisms, primarily chemical and behavioral. Here, we examined if the Dufour's gland

secretion in the primitively eusocial bumble bee *Bombus impatiens* signals information about

caste, social condition, and reproductive status. We chemically analyzed Dufour's gland contents

across castes, age groups, social and reproductive conditions, and examined worker behavioral

and antennal responses to gland extracts.

We found that workers and queens each possess caste-specific compounds in their Dufour's

glands. Queens and gynes differed from workers based on the presence of diterpene compounds

which were absent in workers, whereas four esters were exclusive to workers. These esters, as

well as the total amounts of hydrocarbons in the gland, were produced in larger quantities by

queenless, fertile workers at specific age groups as compared to queenright, sterile workers.

Olfactometer bioassays demonstrated attraction of workers to Dufour's gland extracts that did

not represent a reproductive conflict, while electroantennogram recordings showed higher

overall antennal sensitivity in queenless workers. Our results demonstrate that compounds in

the Dufour's gland act as caste- and physiology-specific signals and are used by workers to

discriminate between workers of different social and reproductive status.

Introduction

Introduction

Reproductive division of labor in insect societies is regulated through multiple concurrent

mechanisms. These often involve a combination of interactions with brood[1,2], agonistic and

policing behaviors among nestmates[3–5], and chemical signaling among individuals in the colony[6–

8]. In the latter, chemical signals from the queen can affect worker physiology and behavior by

reducing their ovarian activation[9] and by marking eggs to indicate their maternity, which

prevents their consumption by "policing" workers[10–12]. In workers, chemical signals can indicate

fertility status[13–15], which increases aggressive behavior towards fertile workers[16] or indicate

sterility, which appease aggressive individuals[8,17]. Across social insects, these signals are

chemically diverse and have multiple glandular origins[18,19].

In female Hymenoptera, one such glandular origin is the Dufour's gland (DG), which is associated

with the sting apparatus in females and produces a remarkable diversity of compounds with

myriad functions[20,21]. In solitary bees, the DG secretion functions primarily in waterproofing

brood cells[22,23], although there is also evidence that it is the source of a sex pheromone in two

parasitoids[24], and may serve as a nest-marking pheromone in certain solitary bee species[25,26]. In

social bees, analyses of the DG secretion have suggested additional functions in social

communication[27], such as signaling of reproductive status and the possible regulation of worker

aggressive and foraging behaviors in honey bees and bumble bees[8,17,28,29].

Bumble bees provide an excellent opportunity to study the role of chemical signals in the

regulation of reproduction because they exhibit characteristics of both simple and highly eusocial

insect societies. In the first part of the colony cycle, reproduction is dominated by a single queen,

with workers being functionally sterile. This changes in the latter half of the colony cycle, the

competition phase, where workers initiate egg laying and produce males in competition with the

queen[30]. In *Bombus terrestris,* the DG secretion varies with caste, reproductive state, age, and

task[8,29]. In queens, the DG contains aliphatic saturated and unsaturated hydrocarbons, whereas

in workers these are accompanied by long chain esters. The reduction of these esters to non-

detectable amounts in queenless workers with activated ovaries suggests that they may signal

sterility[8]. There is also a negative correlation between the amount of esters a worker possesses

and the aggression directed toward her during worker-worker competition over reproduction,

suggesting an appeasement effect of these esters, in line with their hypothesized function as

sterility signals[17]. It is unknown if these signals are widespread in *Bombus*, or represents a unique

chemical signaling mechanism in *B. terrestris*. In the present study, we leverage our

understanding of *B. terrestris* chemical ecology in order to glean new insights into potential

shared or elaborated mechanisms mediating social organization across bumble bee species*.

To examine the role of the gland from a signal production standpoint and discover if *B. impatiens*

females produce a social signal in the DG linked to their fertility status, we examined the chemical

composition of the secretion as a function of caste, age, social, and reproductive status. To

examine the signaling role of the gland from a sensory standpoint, we tested the behavioral and

antennal responses of workers, either queenless or queenright, to the DG secretion of either

queenless or queenright workers. The DG may carry multiple social signals which are caste or

social status dependent. Depending on what these roles are, DG compounds may induce

attraction or avoidance to facilitate cooperation or competition. Therefore, we hypothesize that

if workers produce a cooperative signal (e.g., sterile workers producing esters), a DG extract of

these workers will elicit higher attraction by cooperative workers in comparison with a control

extract, and if workers produce a fertility signals, a DG extract of these workers will elicit

avoidance in competing workers as compared to control. Additionally, we hypothesize that the

secretion from workers carrying a cooperative or competitive signal will elicit stronger antennal

responses in other workers, reflecting increased sensitivity towards compounds that may impact

their fitness.

Materials and Methods

Bee colonies and sample preparation for chemical analysis

Worker samples were collected from *B. impatiens* colonies (n = 8) that were obtained from

Koppert Biological Systems (Howell, Michigan, USA) or Biobest Canada Ltd. (Leamington, ONT).

Upon arrival to the laboratory, colonies were approximately 2-3 weeks old (based on first worker

emergence), with less than 30 workers each, a queen, and all stages of brood. Colonies were
maintained in nest-boxes at a constant temperature of 28–30°C and 60% relative humidity,
constant darkness and supplied *ad libitum* with a 60% sugar solution and honeybee-collected
pollen. These colonies were used as sources of newly-emerged workers that were sampled prior
to sexual production phase (ie, gynes and males). Newly-emerged workers were assigned to one
of three treatments: queenless (QL, n=70), queenless and brood-less (QLBL, n=70), and
queenright (QR, n=70) workers. QR workers were individually labeled and remained in their natal
colony in the presence of the queen until the day of collection while QR and QLBL workers were
housed in a plastic cage (11 cm diameter × 7 cm height) in groups of 3-6 workers without a queen.
Queenless groups of workers typically lay eggs within 6-8 days[31], and the presence of brood may
also affect worker reproduction. Therefore, in the QL groups, eggs laid by workers remained in
the cages, whereas in the QLBL groups, eggs laid by workers were removed daily. QR workers
were collected from young queenright colonies showing no signs of competition over
reproduction between the queen and the workers. Workers ranging in age from one to fourteen
days old (5/age group) were collected from all treatments. All workers were collected and stored
at −80° C until dissection.
Egg-laying queens (n=20) were obtained from full-sized colonies with >100 workers. All egg-laying
queens were actively producing female workers prior to sampling. Unmated queens (gynes) were
collected from 3 gyne-producing colonies. They were separated from their natal colonies upon
emergence to prevent mating, housed in small cages in groups of 3-5 gynes and sampled at 4 age
points: 3, 6, 10, and 14 days after emergence (5/age group, 20 queens in total).
Olfactometer and electrophysiology bioassays were conducted using 7-day old QR and QLBL
workers exposed to DG extracts of either 7-day old QR or QLBL workers, based on differences
observed in the chemical analysis and ovarian activation in these groups.
115

<u>Dissection of bees</u>
Ovaries were dissected under a stereomicroscope in distilled water, and the largest three
terminal oocytes across both ovaries (at least one from each ovary) were measured with an
eyepiece micrometer. The mean of these oocytes for each bee was used as a measure of ovary

activation. Mean terminal oocytes were used either as a continuous variable or to categorize

workers into three groups: inactive (<0.5mm), intermediate (0.5 - 2mm), and active (>2mm).

DGs from each individual were dissected and immediately immersed in hexane. Glands were

extracted for 48 hours after which each sample was evaporated and reconstituted in 100 μL

hexane spiked with an internal standard of 1 μg eicosane (C_{20}).

<u>Chemical analysis</u>

To identify compounds in the DG, extracts were analyzed on two separate mass spectrometers.

At UC Riverside, samples were analyzed on a Hewlett-Packard (H-P, now Agilent Technologies,

Santa Clara CA, USA) 6890 gas chromatograph (GC) equipped with a DB-17 mid-polarity column

(0.25 mm id x 30 m x 0.25 μm film thickness; J&W Scientific, Folsom CA, USA) interfaced to an H-

P 5973 mass spectrometer. The temperature program was as follows: 100˚C (0 min. hold) to

160˚C at 20˚C/min, then 4˚C/min to 280˚C (hold for 25 min). Injections were made in split-less

mode, with injector and transfer line temperatures of 280˚C. At Penn State, samples were

analyzed on an Agilent 7890A GC equipped with a HP-5ms column (0.25 mm id x 30 m x 0.25 μm

film thickness, Agilent) and interfaced to an Agilent 5975C mass spectrometer. Compounds were

tentatively identified based on diagnostic ions in the resulting spectra and retention indices, and

where possible, identifications were confirmed by matching retention times and mass spectra

with those of authentic standards.

Compounds in DG extracts were quantified on a Trace 1310 GC (Thermo Fisher, Waltham, MA

USA) equipped with a flame-ionization detector (FID) and a TG-5MS column (0.25 mm id x 30 m

x 0.25 μm film thickness; Thermo Fisher). The temperature program was as follows: 60˚C to 120˚C

at 15˚C/min followed by 120˚C to 300˚C (5m hold) at 4˚C/min. The injector port and FID were

held constant at 250˚C and 320˚C, respectively.

<u>Olfactometer Bioassays</u>

To examine if *B. impatiens* workers perceived the DG secretion by olfaction, and if their response

depended on reproductive or social state, bioassays were conducted using two-choice

olfactometers as previously described[32]. Olfactometers (see Fig. S1 for illustration) were

fashioned from plastic petri dishes (150 × 15 mm) with two equidistant holes (2 cm diameter) in

the bottom providing access to small plastic cups that held treatment or control stimuli (QR or

QLBL DG extract or control). Bees were prevented from returning to the main arena after a choice

was made via a 3 cm section of plastic straw glued to each hole. After lifting the petri dish lid

enough to allow entry, workers were placed in the center of the olfactometer and given 60 min

to make a choice. One gland equivalent of extract (100 µl total volume) was pipetted onto glass

coverslips and the solvent was allowed to evaporate before being introduced into the cups, which

were subsequently affixed to the olfactometer. Bioassays were conducted under a red light and

olfactometers were washed with soap and water and allowed to air dry before being reused.

Cups containing test stimuli, extracts and bees were only used once. All behavioral assays were

conducted in the rearing chamber between noon to 4 PM.

7-day old QR (n=39) or QLBL workers (n=48) were tested for their preference between 2 choices

in three bioassays: QR DG extract vs. callow DG extract, QLBL DG extract vs. callow DG extract,

and QR DG extract vs. QLBL DG extract. Overall, we conducted 122 bioassays (16-25 replicates

per treatment group and social condition) which resulted in 71.3% response rate. Bioassays

where bees did not respond within the time frame were not included in the analysis. Non-

responder bees were not associated with any specific treatment (proportion test, χ^2 = 0.16, p =

0.69, 25.9% and 30.9% of QL and QR non-responders, respectively). Callow DG extract was used

as control as it is undifferentiated into a specific chemical profile, being mostly devoid of potential

signal molecules, but went through the same extraction process as 7-day old worker DG,

capturing methodological noise that a solvent control would not. In addition, the preference to

a solvent control and callow extract were not significantly different when compared directly

(binomial exact test, p = 0.77, n = 12, 20% non-responders). In all bioassays, extracts were

presented to workers as a dose of 1-bee equivalent. Extracts were prepared as described above

for the chemical analyses, pooled, and the concentration was adjusted to 1 gland equivalent per

100 µL hexane.

<u>Electroantennogram recordings</u>

Odor cartridges were created using 7-day old worker DG extracts, either QR or QLBL, as follows.

1 DG equivalent per 100 µL of hexane (DG extracts were prepared as above) was administered

onto strips of Whatman filter paper. After the solvent had evaporated, the filter paper strips were

inserted into glass pipettes and sealed with aluminum foil until use. A "blank" control consisting

of 100 µL of solvent was allowed to evaporate from the filter paper before being placed and

sealed in the cartridge. For the electroantennograms (QR n=13, QLBL n=16), the right antenna

was excised and the tip snipped with a fine scissors prior to mounting on a quadroprobe

electroantennogram system with Spectra 360 electroconductive gel[33]. A constant airflow of

charcoal-purified humidified air was passed across the antenna through a 10 mm glass tube for

the duration of each experiment. Each antenna was allowed to reach a steady baseline before

starting the experiment, at which point the test odorants were delivered into a steady airstream

using the odor cartridges in conjunction with a stimulus flow controller (Syntech, Hilversum,

Netherlands). A 40 ml/s air pulse was pushed through the cartridge for 0.05 seconds, effectively

delivering 2 ml of volatiles into the air stream and across the antenna. Responses to each stimulus

were amplified using a quadroprobe, 10x pre-amplifier (Syntech, Netherlands) and further

amplified 10x before being recorded and analyzed on a desktop computer using custom software

(as in[34]). Odorants from control, QLBL, or QR extracts were presented to each antenna 30 seconds

apart, after the antennal response had returned to baseline activity. Responses to a final blank

puff were subtracted from the responses to the test odorants and the corrected values were used

in analyses. Each worker antenna received all treatments, and each odor cartridge was used

twice.

<u>Statistical analysis</u>

Statistical analyses were performed using R (version 3.6.1) in RStudio (version 1.2.1335). To

examine differences in the DG secretion between females of differing caste (queen, workers) and

social conditions (QR, QL, QLBL worker groups), linear discriminant analyses (LDA) were

performed using Z-transformed relative peak areas as predictor variables, and social condition,

caste, or age as grouping variables. Only peaks > 0.5% of the total peak area were used. These

selected peaks were then re-standardized to 100% and transformed using Aitchinson's

formula $[Z_{ij} = \ln[Y_{ij}/g(Y_j)]]$[35], to reduce collinearity prior to multivariate analysis[36,37]. For peaks that

were not present in all groups, a small nominal value was added as the transformation cannot be

calculated with zero values. In some cases, poorly separated clusters of peaks prevented accurate

integration, and after mass spectral confirmation that these were of the same compound class

(e.g. esters), they were integrated together as a "complex" and used as a single variable.

Amounts of DG compounds and oocyte size were not normally distributed (Shapiro-Wilks test p

< 0.05) and were therefore analyzed with a generalized linear model (GLM) using the R package

lme4. To examine the effect of social condition and age on worker oocyte size, social condition

and age were used as fixed effects and colony as a random effect, with a gaussian distribution

and log as link function. When two numeric independent variables were included in the model,

these were centered and scaled to allow comparison of their coefficients. To examine the effect

of social condition, age, and oocyte size on the total amount of esters in worker DGs, those terms

were used as fixed effects, and colony as a random effect. Wilcoxon tests were used to compare

terminal oocyte size, and ester and hydrocarbon amounts between QLBL and QR 7-day workers.

Choice data from olfactometer bioassays were analyzed using exact binomial tests, calculating

two-tailed probabilities for a difference in the proportion of choices against the null hypothesis

of 50%, or no preference. The average strength of antennal responses from electroantennogram

recordings were compared using t-tests when comparing responses based on antennal social

condition, and paired t-tests when comparing antennal responses within social condition and

between extract type. Significance was evaluated at the α = 0.05 level.

Results

Chemistry of *Bombus impatiens* Dufour's gland

DG extracts and the average oocyte size were analyzed in 210 workers (5/age group, ages 1-14

days for 3 treatment groups), 20 unmated gynes (5/age group, ages 3, 6, 10, 14 days), and 20

active queens (4-15 weeks from the emergence of the first worker). During analysis, we

discovered 11 outlier worker samples, which were excluded from further analyses. One QR

worker had fully activated ovaries at 8 days of age, which was a clear outlier, and 10 QL workers

on days 9 and 10 sharing the same 2 cages and from the same colony had unusually undeveloped

ovaries which likely resulted from a shared effect related to rearing conditions (e.g.,

underfeeding) or maternal colony. Thus, 199 workers were included in subsequent analyses.

Overall, GC-FID analysis of *B. impatiens* DG extracts in females revealed 98 compounds comprised

of aliphatic alkanes (C_{21}-C_{31}), alkenes, wax esters, and diterpenes (Fig. S2, Table S1).

Examining caste specificity in Dufour's gland secretion

The DGs of different castes of *B. impatiens* contain compounds that are both queen-specific and

worker-specific, in addition to linear alkanes and alkenes found in both castes (Fig S2). Gynes and

queens differed from workers based on the presence of the diterpenes ß-springene and isomers,

which were absent in workers. Conversely, three dodecyl esters (dodecyl octanoate, dodecyl

decanoate, and dodecyl dodecanoate), and hexadecyl decanoate were specific to workers (Fig

S2). In addition, workers produced large quantities of a fifth ester, dodecyl oleate, which was

present only in trace amounts in gynes. In gynes and queens, β-springene accounted for 3.2% ±

0.19 and 1.3% ± 0.33 (mean ± SE, n=20), on average, of the total secretion, respectively. The

worker-specific esters comprised, on average, 1.0% ± 0.07, 2.6% ± 0.08, 1.8% ± 0.08, 1.5% ± 0.13,

and 5.1% ± 0.28 (mean ± SE, n=199) of the total secretion across workers of all ages and

treatments, respectively. Long chain esters (tentatively identified as terpene esters 6 and 7)

comprised, on average, 1.2% ± 0.18 and 2.2% ± 0.26 (mean ± SE, n=20) of the gyne secretion,

while only being detected in trace amounts in workers, and at <1% in DG extracts of queens.

Discriminant analyses using either all compounds or specific chemical classes (esters, alkanes,

and alkenes) as predictor variables showed that esters enabled differentiation between active

queens, gynes, and workers with similar explanatory power as the analyses using all compounds

(Fig 1). Figure 1 shows plots of the data according to the first two linear discriminant functions

based on the compound class used as predictor variables. Of the worker esters, dodecyl

decanoate provided the greatest separation between queen and worker castes along LD1

(coefficient of LD1 1.71, % variation explained = 93%), while of the gyne esters, terpene esters 6

and 7 accounted for the most variability along LD2 (coefficients of LD2 0.94 and -1.22

respectively, % variation explained = 7%).

Examining the effect of age, social, and reproductive state on worker Dufour's gland secretion

The discriminant analysis data indicate clear separation between queens, gynes, and workers, but produced little to no separation between the worker groups (Fig. 1). However, the large differences between the castes may have masked the more subtle differences between workers. Therefore, we analyzed the three worker groups separately and tested the effects of worker age, social conditions, and reproductive status) on their DG composition. GLM models were fit using social condition (categorical variable with 3 levels: QR, QL, QLBL), age (as a scaled continuous variable, days 1-14 after eclosion), and ovarian activation (as a scaled continuous variable, 0-4 mm as measured by the average terminal oocyte size), and interactions between these variables when significant.

Reproductive Status. Ovarian activation increased with age in all treatment groups. However, workers in the QR social condition progressed at a slower pace compared to the QL and QLBL groups, which was especially evident during days 7-12 (Fig. 2A). There was a significant positive effect of ovarian activation on the amount of the total DG secretion (GLM estimate = 0.35, SE = 0.09, t-value = 4.02, $p < 0.0001$), and total hydrocarbons (GLM estimate = 0.22, SE = 0.07, t-value = 3.01, $p = 0.003$), but not total esters ($p = 0.94$) (Figs 2B & 2C. The total secretion is the sum of hydrocarbons and esters and therefore not shown separately).

Social condition. There was no significant effect of social condition on the amount of total DG secretion (QLBL GLM estimate = -0.09, SE = 0.10, t-value = -0.89, $p = 0.37$; QR GLM estimate = -0.11, SE = 0.09, t-value = -1.22, $p = 0.22$), or amount of total hydrocarbons (QLBL GLM estimate = -0.07, SE = 0.13, t-value = -0.70, $p = 0.48$; QR GLM estimate = -0.16, SE = 0.10, t-value = -1.68, $p = 0.09$). There was, however, a significant positive effect of social condition on the amount of esters in the QR (GLM estimate = 3.99, SE = 1.22, t-value = 3.25, $p = 0.01$), and QLBL groups (GLM estimate = 3.55, SE = 1.2, t-value = 2.94, $p = 0.003$) with higher amounts of esters produced by QR and QLBL workers vs. QL workers.

Age. There was a significant positive effect of age on the total DG secretion (GLM estimate = 0.47,

SE = 0.08, t-value = 6.18, p < 0.0001), the total amount of hydrocarbons (GLM estimate = 0.4, SE

= 0.07, t-value = 5.66, p < 0.0001), and the total amount of esters (GLM estimate = 0.37, SE =

0.09, t-value = 4.28, p < 0.001). There were positive effects of age (GLM estimate = 0.116, SE =

0.009, t-value = 12.27, p < 0.0001) and the QLBL social condition (GLM estimate = 0.429, SE =

0.075, t-value = 5.71, p < 0.0001) on worker ovarian activation (Fig. 2A). All variables tested in

this study were affected by age, demonstrating the need to control for age in order to understand

the signaling role of the glandular secretion in females.

Overall, the reproductive state of workers was positively correlated with the total secretion,

hydrocarbons and ester amounts, while the social condition was associated mostly by the ester

amounts. To further examine the link between DG chemistry and the reproductive state of

workers, we performed a discriminant analysis with three categorial classifications of the ovaries

(see Methods for how categories were determined; Fig. 3). This analysis demonstrates that either

esters or the entire secretion provide more group differentiation than alkenes or alkanes when

workers are grouped according to ovarian activation.

Examining the behavioral responses of *B. impatiens* workers to Dufour's gland extracts

The largest differences in ovary activation, amounts of esters and hydrocarbons between the two

most distinct social conditions (QR and QLBD) were at the age 6-7 days (ovaries: Wilcoxon test,

W = 4, p < 0.001; QR – 0.7 ± 0.04, QLBL – 1.9 ± 0.31; mean mm ± SE, n = 10), esters (W = 0.8, p <

0.001; QR – 4.8 ± 1.1, QLBL – 0.9 ± 0.1; mean µg ± SE, n = 10), and hydrocarbons (W = 86, p =

0.005; QR – 14.4 ± 2.6, QLBL – 5.6 ± 1.1; mean µg ± SE, n = 10; Fig. 4). Therefore, this time point

was chosen for the behavioral and antennal assays. In olfactometer bioassays, workers from both

social conditions (QR, QLBL), given a choice between a callow worker DG extract (which served

as a control) and QLBL DG extract, avoided QLBL DG extract (QLBL responders, n = 14, p = 0.01;

QR responders n = 16, p = 0.004, binomial exact test; Fig. 5). When provided with a choice

between callow extract vs. QR DG extract, workers of neither social condition showed a

significant preference (QLBL responder, n = 16, p = 1; QR responder, n = 16, p = 0.21). Similarly,

when provided with a choice between QLBL and QR DG extracts, neither showed a significant
preference (QLBL responder, n = 15, p = 1). Both 7-day old QR and QLBL workers preferred
extracts that represented lack of competition (i.e. callow or queenright workers). Seven day old
QLBL workers that had, on average, lesser amounts of esters and higher ovary activation (Fig. 4),
preferred the control over QLBL extract and QLBL over QR extract, while 7-day old QR that had,
on average, more esters and hydrocarbons and lower ovary activation (Fig. 4), preferred the
control over either QLBL or QR extracts.

Examining the antennal responses of *B. impatiens* workers to Dufour's gland extracts
Electroantennogram recordings from antennae of 7-day-old workers of different social
conditions responding to QLBL or QR DG extracts (also from 7-day-old workers) revealed that
antennae of QLBL workers produced stronger responses, on average, than those of QR workers
(p = 0.02, Fig. 6). Within social condition, QR workers produced stronger responses to DG extract
from their own social condition than the QLBL extract (p < 0.0001, Fig. 6), while QLBL workers
showed no difference in response based on the extract type (p = 0.64). Overall, QLBL that had,
on average, greater ovary activation and lesser amounts of esters (Fig. 4) exhibited stronger
responses than QR, and QR that had, on average, lower ovary activation and more esters were
more responsive to QR extracts.

Discussion
The results of this study reveal the presence of caste-specific and physiology-specific compounds
in the DG of *B. impatiens* females. Our results further demonstrate that the DG secretion not only
contains important information about caste, age, social, and reproductive condition, but also that
the signaling properties of the gland contents are perceived and used by workers to discriminate
among workers. This discrimination suggests that workers are attracted to DG extracts of workers
that represent a lack of competition, and that QLBL workers, as opposed to QR workers, have
higher sensitivity to DG extracts, representing an increased investment in sensory capability.

The discriminant analysis of DG secretion showed clear differentiation of worker and queen
groups, highlighting the caste-specific chemistry of the gland. This separation was primarily
explained by worker-specific esters, gyne-specific esters, and queen-specific diterpene
compounds, although alkanes and, to some extent, alkenes contributed to some of the variability
(Fig. 1). The long chain esters that separate gynes from workers are also found in much lower
amounts in active queens, pointing towards a signal that functions either in caste recognition by
workers, or sexual attraction of males. While the latter was not investigated in this study, caste
recognition by workers is supported by characteristic behaviors of workers towards gynes. For
example, although workers can be quite aggressive towards other workers or active queens from
a different colony, they remain docile in the presence of gynes (Pers. obs., ND and EA).

The diterpene compounds that characterize the queen caste DG are primarily known from the
labial glands of bumble bee males, where they are hypothesized to serve as sex-pheromone
components[38]. Female sex pheromones were identified from head and body surface extracts of
gynes[39], but DG components have not been investigated as sex pheromones. Previous research
comparing the DG secretion in unmated gynes of 5 species of European bumble bee found large
variation in long-chain wax esters, with *B. pascuorum* and *B. lucorum* possessing up to 5% of
hexadecenyl hexadecenoate or octadecyl octadec-9-enoate, while *B. hypnorum, B. lapidarius,*
and *B. terrestris* had none or less 1%[40]. Interestingly, diterpenes were not found in the DG of
these species, indicating that they may be species and caste specific. The DG's diterpene
compounds in *B. impatiens* could also function in caste recognition, a role that is often ascribed
to cuticular hydrocarbons[41,42]. The DG of the honey bee contains ~12 times more secretion in
queens than workers and contains caste specific compounds, leading previous researchers to
hypothesize that the gland deposits a signal of egg-maternity that distinguishes queen- from
worker-laid eggs[43,44]. Caste-specific compounds, likely from the DG, have also been identified
from the eggs of the bumble bee *B. terrestris*[45]. In *B. impatiens*, it is unclear why both the queen
and worker castes would each need a unique set of compounds to differentiate egg-maternity,
or why workers would advertise egg maternity, because this is likely to result in oophagy[3],
suggesting that these compounds might have another as yet unknown signaling function.

Workers produced a series of dodecyl esters that were not detected in extracts from gynes or active queens (Fig S2, Table S1). A signal is defined as a trait, structure, or behavior that evolved to convey specific information to, and elicit a response from, a receiver, increasing both the emitter's and receiver's overall fitness[46,47]. Because worker signals are specific to workers and are detected by other workers, they must benefit workers. We did not focus on the ability of queens, gynes, or males to differentiate workers based on their DG signals, but the secretion may contain multiple signals targeting different individuals, as occurs with CHCs which signal both fertility and nestmate identity in many social insects[42,48], or with the honey bee queen mandibular pheromone, which functions as a primer pheromone to both workers and males, and a sex pheromone to males[49,50]. The worker-specific signal could play a role in the interactions among workers, as in *Bombus terrestris*[8,17,29,51], or have an additional unknown signaling role. The benefit of the signal to the receiver in the interactions between workers also may change according to the social condition of the workers. QR workers may gain from both producing a signal that conveys harmony and respond to such a signal in order to maintain the social phase they are in (i.e., protect the queen and ensure the production of future gynes), while QL workers, who enter a reproductive mode, may gain from being tuned to signals that convey potential competition that may threaten the fate of their eggs. By manipulating social condition to produce either QLBL reproductive workers or QR non-reproductive workers, and then testing their perception and behavioral response to DG signals from those same groups, we show that workers preferentially avoid bees of a non-harmonious reproductive state (Fig. 5), with both QR and QL workers choosing callow over QL extract, and being indifferent between control and QR extracts.

In electroantennography experiments, antennae from QLBL workers produced stronger responses than QR antennae (Fig. 6), suggesting that QLBL antennae are more sensitive to perceiving DG signals. Queenlessness is known to affect olfactory development and performance in the honey bee, where the worker olfactory system develops substantially during the first 6 days of life, and is strongly affected by the queen's presence[52,53]. Bumble bees and honey bees have similar life span and 6-7 days is the same time period that workers in this study were kept

queenless. Additionally, in honey bee workers, queenlessness is associated with a reduction in

glomerular volume in the antennal lobe, and decreased learning of conditioned stimuli[54]. It is not

known if similar processes govern bumble bee worker development. Reproductively active 7-day-

old workers are likely capable of laying eggs and would benefit from detecting and avoiding the

queen, who would aggress the workers that challenge her reproductive monopoly. Additionally,

QR workers showed a reduced antennal response to QLBL extract in comparison to QLBL workers

in electroantennography experiments (Fig 6). This could indicate that QR bees are operating in

"harmony mode" within the colony with a higher threshold for detecting and aggressing other

individuals. It is not known at what level this differential response is regulated. QR and QLBL bees

could have differing numbers of olfactory receptors tuned to the DG compounds, or these

chemical signals could be interpreted at higher levels of signal integration and processing in the

brain.

Within workers, the composition of the DG secretion varied between social conditions, and the

amount of the secretion increased with age (Fig. 2), with differences in the amounts of esters and

hydrocarbons at certain age groups (eg, in 6-7-day-old QR and QLBL workers; Fig. 4). While the

broad qualitative composition of DG extracts in *B. impatiens* workers is similar to related species

of *Bombus*[8,45,55,56], the influence of social condition and reproductive state on gland chemistry is

different than the most closely studied species, *B. terrestris*. In *B. terrestris* workers, the

percentages of DG esters are negatively correlated with ovarian activation[8,29], and the size of the

gland is positively correlated with ovarian activation[57]. Here, we demonstrate that in *B.*

impatiens, the amounts of compounds of all chemical classes, including esters, are positively

correlated with age and ovarian activation, regardless of social condition, while the percent

composition is stable over time (Fig. S3). This indicates that differences in the amount of

compounds observed at specific ages result from upregulation of the gland as a whole, instead

of specific components changing independently. In some systems, when a worker transitions into

fertility, their pheromonal secretion becomes more queenlike[28]. In *B. impatiens*, fertile worker

secretions do not mimic queen secretions, instead both maintain the production of compounds

unique to their own caste. Thus, the quantitative differences in DG components appears to

represent a more subtle fertility signal than in *B. terrestris,* where the wax esters are only present

workers with inactivated ovaries[8]. This may require a more finely tuned threshold for aggression

and help explain the relatively docile conflict between queen and workers over reproduction of

B. impatiens compared with *B. terrestris*[58,59]. Furthermore, the DG esters also vary between the

two species, with dodecyl esters predominating in *B. impatiens* and octyl esters predominating

in *B. terrestris*[8]. Interestingly, dodecyl esters are also found in the labial glands of *B. terrestris*

females[60] whereas predominately octyl esters are found in the labial glands of *B. impatiens*

(Orlova M, Villar G, Hefetz, A, Millar J, Amsalem E, in preparation), presenting an interesting case

of inversion and suggesting a regulatory system that is shared but not identical in different

bumble bee species.

Thus, our data suggest that the DG contains multiple concurrent signals that contain information

about caste, age, social and reproductive condition. The presence of diterpene and ester

compounds, which are much less common than aliphatic alkanes and alkenes in insects, may

serve as additional channels with which to communicate this information. It is also possible that

the signal carries different information to different perceiving individuals. For instance, diterpene

compounds may signal "gyne" to conspecific workers and "mate" to conspecific males. Future

research should investigate where compounds from the DG can be found in a colony, whether

they are deposited onto eggs and/or the cuticular surface of the bee, and their role in worker-

worker, worker-queen and queen-male interactions.

Acknowledgments

The authors are grateful to Dr. Tom Baker for use of his electrophysiology equipment, and to

Freddy Purnell for assistance in bumble bee colony maintenance and rearing.

Funding

This work was funded by NSF-CAREER IOS-1942127 to EA and JM

References

1. Orlova, M., Starkey, J. & Amsalem, E. A small family business: synergistic and additive

effects of the queen and the brood on worker reproduction in a primitively eusocial bee.

J. Exp. Biol. **223**, (2020).

2. Starkey, J., Brown, A. & Amsalem, E. The road to sociality: brood regulation of worker

reproduction in the simple eusocial bee *Bombus impatiens*. *Anim. Behav.* **154**, 57–65

(2019).

3. Zanette, L. R. S. *et al.* Reproductive conflict in bumblebees and the evolution of worker

policing. *Evolution (N. Y).* **66**, 3765–3777 (2012).

4. Wenseleers, T. *et al.* A test of worker policing theory in an advanced eusocial wasp,

Vespula rufa. *Evolution* **59**, 1306–14 (2005).

5. Duchateau, M. J. & Velthuis, H. H. W. Ovarian development and egg-laying in workers of

Bombus-terrestris. *Entomol. Exp. Appl.* **51**, 199–213 (1989).

6. Hoover, S. E. R., Keeling, C. I., Winston, M. L. & Slessor, K. N. The effect of queen

pheromones on worker honey bee ovary development. *Naturwissenschaften* **90**, 477–80

(2003).

7. Vargo, E. L. & Laurel, M. Studies on the mode of action of a queen primer pheromone of

the fire ant *Solenopsis invicta*. *J. Insect Physiol.* **40**, 601–610 (1994).

8. Amsalem, E., Twele, R., Francke, W. & Hefetz, A. Reproductive competition in the

bumble-bee *Bombus terrestris*: do workers advertise sterility? *Proc. R. Soc. B* **276**, 1295–

304 (2009).

9. Le Conte, Y. & Hefetz, A. Primer pheromones in social Hymenoptera. *Annu. Rev. Entomol.*

53, 523–542 (2008).

10. Oi, C. A. *et al.* Dual effect of wasp queen pheromone in regulating insect sociality. *Curr.*

Biol. **25**, 1638–1640 (2015).

11. Smith, A. a., Hölldobler, B. & Liebig, J. Hydrocarbon signals explain the pattern of worker

and egg policing in the ant *Aphaenogaster cockerelli*. *J. Chem. Ecol.* **34**, 1275–1282

(2008).

12. Endler, A. *et al.* Surface hydrocarbons of queen eggs regulate worker reproduction in a

social insect. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 2945–50 (2004).

13.   Dietemann, V., Peeters, C., Liebig, J., Thivet, V. & Hölldobler, B. Cuticular hydrocarbons

mediate discrimination of reproductives and nonreproductives in the ant *Myrmecia*

gulosa. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 10341–6 (2003).

14.   Cuvillier-Hot, V., Cobb, M., Malosse, C. & Peeters, C. Sex, age and ovarian activity affect

cuticular hydrocarbons in *Diacamma ceylonense*, a queenless ant. *J. Insect Physiol.* **47**,

485–493 (2001).

15.   Visscher, P. K. & Dukas, R. Honey bees recognize development of nestmates' ovaries.

Anim. Behav. **49**, 542–544 (1995).

16.   Smith, A., Hölldober, B., Liebig, J., Holldobler, B. & Liebig, J. Cuticular hydrocarbons

reliably identify cheaters and allow enforcment of altruism in a social insect. *Curr. Biol.*

19, 78–81 (2009).

17.   Amsalem, E. & Hefetz, A. The appeasement effect of sterility signaling in dominance

contests among *Bombus terrestris* workers. *Behav. Ecol. Sociobiol.* **64**, 1685–1694 (2010).

18.   Fletcher, D. J. C. & Ross, K. G. Regulation of reproduction in eusocial Hymenoptera. *Annu.*

Rev. Entomol. **30**, 319–343 (1985).

19.   Keeling, C. I., Plettner, E. & Slessor, K. N. Hymenopteran semiochemicals. *Top. Curr.*

Chem. **239**, 133–77 (2004).

20.   Mitra, A. Function of the Dufour's gland in solitary and social Hymenoptera. *J. Hymenopt.*

Res. **35**, 33–58 (2013).

21.   Abdalla, F. C. & Cruz-Landim, C. Dufour glands in the hymenopterans (Apidae,

Formicidae, Vespidae): a review. *Brazilian J. Biol.* **61**, 95–106 (2001).

22.   Hefetz, A., M. Fales, H. & Batra, S. Natural polyesters: Dufour's gland macrocyclic

lactones form brood cell laminesters in *Colletes* bees. *Science (80-.).* **204**, 415–417

(1979).

23.   Cane, J. H. Chemical evolution and chemosystematics of the dufour's gland secretions of

the lactone-producing bees (Hymenoptera: Colletidae, Halictidae, and Oxaeidae).

Evolution (N. Y). **37**, 657–674 (1983).

24.   Weseloh, R. M. Dufour's gland: source of sex pheromone in a hymenopterous parasitoid.

Science (80-.). **193**, 695–697 (1976).

25.   Pitts-Singer, T. L. *et al.* Comparison of the chemical compositions of the cuticle and

Dufour's gland of two solitary bee species from laboratory and field conditions. *J. Chem.*

Ecol. **43**, 451–468 (2017).

26.   Shimron, O., Hefetz, A. & Tengo, J. Structural and communicative functions of Dufour's

gland secretion in *Eucera palestinae* (Hymenoptera; Anthophoridae). *Insect Biochem.* **15**,

635–638 (1985).

27.   Hefetz, A. Exocrine glands and their products in non-Apis bees: chemical, functional, and

evolutionary perspectives. in *Pheromone Communication in Social Insects* (eds. Vander

Meer, R. K., Breed, M. D., Espelie, K. E. & Winston, M. L.) (Westview Press, 1998).

28.   Dor, R., Katzav-Gozansky, T. & Hefetz, A. Dufour's gland pheromone as a reliable fertility

signal among honeybee (*Apis mellifera*) workers. *Behav. Ecol. Sociobiol.* **58**, 270–276

(2005).

29.   Amsalem, E., Shpigler, H., Bloch, G. & Hefetz, A. Dufour's gland secretion, sterility and

foraging behavior: Correlated behavior traits in bumblebee workers. *J. Insect Physiol.* **59**,

1250–1255 (2013).

30.   Duchateau, M. J. & Velthuis, H. H. W. Development and reproductive strategies in

Bombus terrestris colonies. *Behavior* **107**, 186–207 (1988).

31.   Amsalem, E., Grozinger, C. M., Padilla, M. & Hefetz, A. *The Physiological and Genomic*

Bases of Bumble Bee Social Behaviour. *Advances in Insect Physiology* **48**, (Elsevier Ltd.,

2015).

32.   Starkey, J., Derstine, N. & Amsalem, E. Do bumble bees produce brood pheromones? *J.*

Chem. Ecol. **45**, 725–734 (2019).

33.   Villar, G., Baker, T. C., Patch, H. M. & Grozinger, C. M. Neurophysiological mechanisms

underlying sex- and maturation-related variation in pheromone responses in honey bees

(*Apis mellifera*). *J. Comp. Physiol. A* **201**, 731–739 (2015).

34.   Park, K. C., Ochieng, S. a, Zhu, J. & Baker, T. C. Odor discrimination using insect

electroantennogram responses from an insect antennal array. *Chem. Senses* **27**, 343–52

(2002).

35.   Aitchison, J. *The Statistical Analysis of Compositional Data*. (Chapman and Hall, 1986).

36.   Martin, S. J., Takahashi, J.-I. & Katada, S.-I. Queen condition, mating frequency, queen

553   loss, and levels of worker reproduction in the hornets Vespa affinis and V. simillima. *Ecol.*

554   *Entomol.* **34**, 43–49 (2009).

555   37.   Derstine, N. T., Gries, R., Zhai, H., Jimenez, S. I. & Gries, G. Cuticular hydrocarbons

556   determine sex, caste, and nest membership in each of four species of yellowjackets

557   (Hymenoptera: Vespidae). *Insectes Soc.* **65**, 581–591 (2018).

558   38.   Valterová, I., Martinet, B., Michez, D., Rasmont, P. & Brasero, N. Sexual attraction: A

559   review of bumblebee male pheromones. *Zeitschrift fur Naturforsch. - Sect. C J. Biosci.* **74**,

560   233–250 (2019).

561   39.   Krieger, G. M. *et al.* Identification of queen sex pheromone components of the

562   bumblebee *Bombus terrestris*. *J. Chem. Ecol.* **32**, 453–71 (2006).

563   40.   Chalikova, L., Hovorka, O., Ptacek, V. & Valterova, I. Exocrine gland secretions of virgin

564   queens of five bumblebee species (Hymenoptera: Apidae, Bombini). *Naturforsch. B* **59c**,

565   582–589 (2004).

566   41.   Ren, W. *et al.* Maculatic Acids—Sex Attractant Pheromone Components of Bald-Faced

567   Hornets. *Angew. Chemie - Int. Ed.* **57**, 11618–11622 (2018).

568   42.   van Zweden, J. S. & Ettorre, P. Nestmate recognition in social insects and the role of

569   hydrocarbons. in *Insect Hydrocarbons: Biology, Biochemistry, and Chemical Ecology* (eds.

570   Blomquist, G. J. & Bagnéres, A.-G.) 222–243 (Cambridge University Press, 2010).

571   43.   Katzav-Gozansky, T., Soroker, V. & Hefetz, A. Plasticity of caste-specific Dufour's gland

572   secretion in the honey bee (*Apis mellifera* L.). *Naturwissenschaften* **84**, 238–241 (1997).

573   44.   Ratnieks, F. L. W. Egg-laying, egg-removal, and ovary development by workers in

574   queenright honey bee colonies. *Behav. Ecol. Sociobiol.* **32**, 191–198 (1993).

575   45.   Ayasse, M. *et al.* Caste- and colony-specific chemical signals on eggs of the bumble bee,

576   *Bombus terrestris* L. (Hymenoptera: Apidae). *Chemoecology* **9**, 112–126 (1999).

577   46.   Wyatt, T. D. *Pheromones and animal behavior: Chemical signals and signatures*.

578   *Cambridge* (Cambridge Univ. Press, 2014).

47.   Smith, M. J. & Harper, D. G. C. Animal signals: models and terminology. *J. Theor. Biol.*

177, 305–311 (1995).

48.   Cournault, L. & de Biseau, J. C. Hierarchical perception of fertility signals and nestmate

recognition cues in two dolichoderine ants. *Behav. Ecol. Sociobiol.* **63**, 1635–1641 (2009).

49.   Slessor, K. N., Winston, M. L. & Le Conte, Y. Pheromone communication in the honeybee

(*Apis mellifera* L.). *J. Chem. Ecol.* **31**, 2731–45 (2005).

50.   Villar, G. & Grozinger, C. M. Primer effects of the honeybee, *Apis mellifera*, queen

pheromone 9-ODA on drones. *Anim. Behav.* **127**, 271–279 (2017).

51.   Amsalem, E., Shamia, D. & Hefetz, A. Aggression or ovarian development as

determinants of reproductive dominance in *Bombus terrestris*: interpretation using a

simulation model. *Insectes Soc.* **60**, 213–222 (2013).

52.   Masson, C. & Arnold, G. Ontogeny, maturation and plasticity of the olfactory system in

the workerbee. *J. Insect Physiol.* **30**, 7–14 (1984).

53.   Allan, S. A., Slessor, K. N., Winston, M. L. & King, G. G. S. The influence of age and task

specialization on the production and perception of honey bee pheromones. *J. Insect*

Physiol. **33**, 917–922 (1987).

54.   Morgan, S. M., Butz Huryn, V. M., Downes, S. R. & Mercer, A. R. The effects of

queenlessness on the maturation of the honey bee olfactory system. *Behav. Brain Res.*

91, 115–126 (1998).

55.   Tengö, J. A. N. *et al.* Species specificity and complexity of Dufour's gland secretion of

bumble bees. *Comp. Biochem. Physiol.* **99B**, 641–646 (1991).

56.   Hefetz, A., Taghizadeh, T. & Francke, W. The exocrinology of the queen bumble bee

Bombus terrestris (Hymenoptera: Apidae, Bombini). *Zeitschrift fur Naturforsch. Sect. C - J.*

Biosci. **51**, 409–422 (1996).

57.   Abdalla, F. C., Velthuis, H., Duchateau, M. J. & Cruz-Landim, C. D. Secretory cycle of the

Dufour's gland in workers of the bumble bee *Bombus terrestris* L. (Hymenoptera: Apidae,

Bombini). *Netherlands J. Zool.* **49**, 135–156 (1999).

58.   Cnaani, J., Schmid-Hempel, R. & Schmidt, J. O. Colony development, larval development

and worker reproduction in *Bombus impatiens* Cresson. *Insectes Soc.* **49**, 164–170 (2002).

59.   Padilla, M., Amsalem, E., Altman, N., Hefetz, A. & Grozinger, C. M. Chemical

communication is not sufficient to explain reproductive inhibition in the bumblebee

Bombus impatiens. *R. Soc. Open Sci.* **3**, 160576 (2016).

60.   Amsalem, E., Kiefer, J., Schulz, S. & Hefetz, A. The effect of caste and reproductive state

on the chemistry of the cephalic labial glands secretion of *Bombus terrestris*. *J. Chem.*

Ecol. **40**, 900–912 (2014).

Author Contributions

EA, AH, GV designed the experiments. GV and ND performed dissections, ovary measurements, and gland extracts. ND performed data and statistical analyses. GV performed olfactometer and GC-EAD bioassays. ND and JM performed GC-MS analysis. ND, and EA wrote the manuscript. All authors provided feedback and edits.

Additional Information

Competing interests

The authors declare no competing interests.

Figure 1 - Linear discriminant analyses on Z-transformed relative peak areas of Dufour's gland
extracts of workers of three social condition treatment groups (queenright – QR, queenless -
QL, and queenless, brood-less - QLBL), unmated gynes, and active queens, with
caste/treatment as the grouping variable. Each point is an individual bee. Panels **a-d** shows the
clustering generated by plotting the first two linear discriminant functions when using only the
esters, alkanes, alkenes, or all compounds.

630
631

ur's gland chemistry as a function of
enright (QR), queenless (QL), and
oocyte length (mm), (B) mean μg of
SE. Data are grouped into two-day
group.

638 Figure 3 - Linear discriminant analyses on Z-transformed relative peak areas of Dufour's gland
639 extracts of workers with three ovarian activation categories as the grouping variable. Each point
640 is the value from an individual bee. Panels **a-d** show the clustering generated by plotting the
641 first two linear discriminant functions when using only the esters, alkanes, alkenes, or all
642 compounds.
643

644
645

646 Figure 4 - Change in 6- and 7-day-old worker reproductive state and Dufour's gland chemistry
647 as a function of social condition in queenless (QR) and queenless-broodless (QLBL) workers. (A)
648 mean terminal oocyte length (mm), (B) mean µg of total esters and hydrocarbons ± SE (n = 10).
649 Each data point is the value from an individual worker.

Figure 5 – The proportion of queenless brood-less (QLBL) and queenright workers (QR) responding to 3 different combinations of stimuli, designated on the left and right of the plot: control vs. QLBL DG extract, control vs. QR DG extract, and QLBL vs. QR DG extract. Control stimuli consisted of callow worker DG extract. All stimuli were presented in the amount of 1 gland equivalent. The number of workers responding to each treatment is given in parenthesis on each bar. Blue bars represent QLBL response bees, and orange bars represent QR response bees. Asterisks indicate significance (p <0.05) using exact binomial tests.

660   Figure 6 – Electroantennogram responses (μV) of queenless broodless and queenright worker
661   antennae to Dufour's gland extracts (1 worker equivalent) from queenless broodless and
662   queenright. Blue and orange boxplots represent responses to queenless broodless and
663   queenright Dufour's gland extracts, respectively. Asterisks indicate significance (* p < 0.05, ***
664   p < 0.0001).