Patients with ventromedial prefrontal lesions show an implicit approach bias to angry faces

Macià Buades-Rotger*[1,2,3], Anne-Kristin Solbakk[4,5,6,7], Matthias Liebrand[1], Tor Endestad[4,5,6,7], Ingrid Funderud[4,5,7], Paul Siegwardt[1], Dorien Enter[3,8], Karin Roelofs[3,8], Ulrike M. Krämer[1,2]

[1] Department of Neurology, University of Lübeck, Lübeck, Germany

[2] Department of Psychology, University of Lübeck, Lübeck, Germany

[3] Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands

[4] Department of Psychology, University of Oslo, Oslo, Norway

[5] RITMO Centre for Interdisciplinary Studies in Rhythm, Time and Motion, University of Oslo, Norway

[6] Oslo University Hospital, Rikshospitalet, Oslo, Norway

[7] Department of Neuropsychology, Helgeland Hospital, Mosjøen, Norway

[8] Behavioural Science Institute, Radboud University, Nijmegen, The Netherlands

* Corresponding author:

Macià Buades-Rotger, PhD

Radboud University

Montessorilaan 3

6525 HR Nijmegen

The Netherlands

+24 3655935

m.buadesrotger@donders.ru.nl

ORCID: http://orcid.org/0000-0003-1219-9216

1   **Abstract**

Damage to the ventromedial prefrontal cortex (VMPFC) can cause maladaptive social behavior, but the

cognitive processes underlying these behavioral changes are still uncertain. Here, we tested whether

patients with acquired VMPFC lesions show altered approach-avoidance tendencies to emotional facial

expressions. Thirteen patients with focal VMPFC lesions and 31 age- and gender-matched healthy

controls performed an implicit approach-avoidance task in which they either pushed or pulled a joystick

depending on stimulus color. While controls avoided angry faces, VMPFC patients displayed an

incongruent response pattern characterized by both increased approach and reduced avoidance of

angry facial expressions. The approach bias was stronger in patients with higher self-reported

impulsivity and disinhibition, and in those with larger lesions. We further used linear ballistic accumulator

modelling to investigate latent parameters underlying approach-avoidance decisions. Controls

displayed negative drift rates when approaching angry faces, whereas VMPFC lesions abolished this

pattern. In addition, VMPFC patients had weaker response drifts than controls during avoidance.

Finally, patients showed reduced drift rate variability and shorter non-decision times, indicating

impulsive and rigid decision-making. Our findings thus suggest that VMPFC damage alters the pace of

evidence accumulation in response to social signals, eliminating a default, protective avoidant bias and

facilitating dysfunctional approach behavior.

Keywords: approach, avoidance, anger, threat, ventromedial, lesion

Introduction

Patients with damage to the ventromedial prefrontal cortex (VMPFC) often show disruptive

social behavior (Anderson et al., 2006; Barrash et al., 2000; Beer et al., 2006; Blair, 2004). Ventromedial

lesions typically impact adjacent white matter, thereby hindering VMPFC-amygdala cross-talk (Folloni

et al., 2019) and rendering individuals more emotionally reactive (Jenkins et al., 2018; Motzkin et al.,

2015). Consequently, antisocial behavior related to VMPFC dysfunction has been classically attributed

to deficits in emotion regulation (Davidson et al., 2000). However, this view has proven difficult to

reconcile with the many other functions ascribed to the VMPFC, such as subjective value computation

(Clithero & Rangel, 2014). This apparent discrepancy has been mended by recent theories that

conceptualize emotion regulation as a special case of value-based decision-making, wherein the brain

must choose among mutually contradicting affective states and behaviors (Dixon et al., 2017; Gross,

2015; Koch et al., 2018). Recent investigations adhere to this model in ascribing an *evaluative* and

generative role to the VMPFC (Hiser & Koenigs, 2018). According to this view, the VMPFC codes for

the potential hedonic or threatening value of a given stimulus in order to steer the organism towards or

away from it (Rudebeck & Rich, 2018). In this framework the VMPFC is assumed to generate cognitive

maps of current internal states and external sensory information, enabling the selection of the most

appropriate course of action (Stalnaker et al., 2015; R. C. Wilson et al., 2014). Such a process has

been termed model-based or goal-directed behavior because it operates on the basis of internal

representations of oneself and the environment rather than by force of habit (Lucantonio et al., 2012).

From this rationale, it follows that antisocial behavior after VMPFC damage could arise from

inaccurate assessment and selection processes. More specifically, VMPFC lesions might impair the

ability to correctly predict the consequences of one's own actions in response to social signals

(Rudebeck & Murray, 2014), e.g., wrongly expecting rewards from approaching potential punishment

cues. Nevertheless, evidence to support this tenet is scarce in humans with VMPFC lesions. One report

suggests that VMPFC-damaged patients display an altered sense of personal distance, e.g., they get

closer to strangers (Perry et al., 2016). Comparably, a study showed that persons with VMPFC lesions

judge negative facial expressions (i.e., angry, disgusted, fearful and sad) as *more* approachable (Willis

et al., 2010). It remains to be tested, however, whether these tendencies can be attributed to implicit

biases during action selection, and whether these putative alterations are linked with actual impairments

in daily functioning. Moreover, it is unclear which precise cognitive mechanisms underlie such abnormal

behavioral dispositions. These are important steps in understanding how VMPFC-dependent

disturbances in social behavior play out in everyday life.

In order to clarify these issues, we investigated whether VMPFC lesions lead to implicit

response biases towards or away from negative, positive, or neutral facial expressions. We used a

version of the approach-avoidance task (AAT) wherein subjects have to either push or pull a joystick

depending on the color (e.g. red or green) of a human face (Roelofs et al., 2010). Faces are

programmed to grow or shrink in size accordingly, giving the impression that they loom closer or recede

upon pulling and pushing, respectively. Hence, the AAT allows measuring implicit response tendencies

to task-irrelevant features of the faces such as their emotional expression. A study with this task

suggested that psychopaths lack automatic avoidance of directly-gazing angry faces, and that this effect

was correlated with aggressiveness (von Borries et al., 2012). Following a similar approach, we tested

whether task scores correlated with patients' daily emotional behavior as measured with validated

clinical scales in order to assess the practical relevance of possible approach-avoidance biases.

In addition, we scrutinized the putative cognitive mechanisms underlying altered task

performance in VMPFC patients using Linear Ballistic Accumulator (LBA) modelling on response times

(Brown & Heathcote, 2008). LBA modelling assumes that decisions arise from a sequential evidence

accumulation process, the speed of which is determined by multiple latent variables (e.g., pre-existing

response tendencies or shorter decision latencies) that can be quantified and compared between

experimental conditions and/or groups. Previous modelling studies on an explicit version of the AAT

reported relatively faster evidence accumulation in healthy subjects when threatening stimuli are to be

avoided (Krypotos et al., 2015; Tipples, 2019). LBA modelling might hence offer insights not captured

by standard methods of reaction time analysis.

Materials and methods

Participants and lesion localization

The clinical sample consisted of 13 patients with chronic (> 6 months post-injury or surgery),

focal damage to the ventral prefrontal cortex (mean age=50.8 [27-62], 7 women, 12 right-handed).

Lesions were mainly ventromedial (Fig. 1A; Table 1), namely Brodmann Areas (BA) 10, 11, 24, 25, and

32. There was also substantial rostromedial damage (BA 9 and anterior BA 10) and, to a lesser extent,

in the ventrolateral prefrontal cortex (BA 45, 46, and 47). Harm to posterior dorsomedial areas (BA 6

and 8) was minimal, with a small number of lesions extending to the anterior insula (BA 13). Lesions

were predominantly right-sided in ten patients, one had an exclusively left-sided lesion, and two had

comparable damage in both hemispheres. All lesioned tissue was restricted to the frontal lobe. Etiology

of the lesions was either meningioma (n=9), traumatic brain injury (n=2), oligodendroglioma (n=1), or

astrocytoma (n=1). The control sample was composed of 31 age- and gender-matched neurologically

healthy individuals (mean age=50.1 [43-54], 19 women, all right-handed). As previously reported,

patients had normal or corrected to normal vision, showed no deficits in standard neuropsychological

testing, and had no motor dysfunction of the hands. However, they reported greater difficulties in

executive function, metacognition, and behavioral regulation as compared to a separate control sample

(see Løvstad et al., 2012 for a complete report). All patients were recruited and measured at Oslo

University Hospital and the University of Oslo, whereas the behavioral control sample was recruited

and measured at the University of Lübeck. All participants provided informed consent and the study

procedures adhered to the Declaration of Helsinki. The study was approved by the ethics committee of

the University of Lübeck and the Regional Committee for Medical Research Ethics - South East Norway.

Clinical scales

Patients filled out the self-report form of the Behavior Rating Inventory of Executive Function –

Adult version (BRIEF-A; Roth et al., 2005) and the Urgency, Premeditation, Perseverance, Sensation

Seeking (UPPS) Impulsive Behavior Scale (Whiteside & Lynam, 2001), both ad-hoc translated into

Norwegian. The BRIEF-A is a standardized rating scale consisting of 75 items that tap into everyday

executive functioning within the past 6 months. Internal consistency and test-retest reliability of the

BRIEF-A are reportedly high and construct validity has been established in healthy and clinical

populations (Waid-Ebbs et al., 2012). For the purposes of this study we discarded all BRIEF-A scales

not directly related to the control of automatic emotional tendencies ("Working Memory",

"Plan/Organize", "Organization of Materials", "Shift" and "Initiate") and thus only considered the scales

"Inhibit", "Emotional Control", and "Self-Monitor". The Inhibit scale measures deficits in inhibitory control

and impulsivity; the Emotional Control scale assesses a person's inability to regulate emotional

responses; and the Self-Monitor scale evaluates difficulties in social or interpersonal awareness. The

UPPS Impulsive Behavior Scale (Whiteside & Lynam, 2001) is a 45-item self-report, assessing different

facets of impulsivity on four subscales. The UPPS has been shown to display good internal consistency

and construct validity (Whiteside et al., 2005). We used the total UPPS score for correlational analyses

because we deemed all subscales ("Negative urgency", "Lack of Premeditation", "Lack of

Perseverance", "Sensation Seeking") to be theoretically associated with emotional action control.

Implicit Approach-Avoidance Task (AAT)

Subjects performed the implicit approach-avoidance task (AAT; Fig. 1B) as previously

described (Roelofs et al., 2010; von Borries et al., 2012). Stimuli were photographs (Ekman & Friesen,

1976; Lundqvist et al., 1998) showing the face of one out of eight actors (four male and four female)

displaying angry, happy or neutral expressions with either direct (straight) or averted (sideways) gaze.

Photographs were cut out ovally and tinted red or green, amounting to a total number of 384 trials.

Participants performed 18 practice trials comprising only straight-gazing neutral faces, followed by the

experimental trials. After half of the trials, subjects had a break, performed two additional practice trials

with directly gazing neutral expressions to recall task demands and completed the second half. Stimuli

were presented randomly, with no more than three of the same emotion-response combinations in

succession. The neutral faces from the practice trials were also presented in the task proper.

Pictures were presented at a 1024 x 768 pixels resolution on a computer screen. We placed

the joystick (Logitech Attack 3) between subject and screen to allow for comfortable pull and push

movements. Participants started each trial by pressing the fire button with the index finger of the

dominant hand. A face stimulus appeared in the center of the screen. Participants were instructed to

ignore the facial expression and only respond to the color of the face. Half the participants had to push

the joystick in response to red and pull in response to green stimuli, the other half had the opposite

instruction. To visually emphasize that pull movements meant approach, and push movements meant

avoidance, pictures grew or shrank in size following pull or push movements, respectively. Stimuli had

a starting size of 9.5º by 13º and could shrink to a minimum of 3.5º by 4.5º when pushing or grow to a

maximum of 15.5º by 20° when pulling. In practice trials, pictures remained visible after erroneous

responses to allow for response correction, whereas in the task proper stimuli disappeared after they

had reached minimal or maximal size. Participants were instructed to respond as quickly and accurately

as possible. Importantly, trials could only be initiated once the joystick was placed back in its original

centered position.

Behavioral data analysis

Reaction times (RT) were recorded as time from stimulus onset until the first joystick movement.

We excluded incorrect trials as well as those with RT shorter than 150ms or longer than 1000ms, and

extracted mean log-transformed RT per cell (as in Bertsch et al., 2018). We then ran an analysis of

variance (ANOVA) on the resulting values with within-subject factors emotion (happy, neutral, angry),

actor gender (male, female), gaze (left, right, and direct), movement (pull or push), and the between-

subject factor group (VMPFC vs healthy controls) using the *ez* package (version 4.4-0). We modelled

all relevant task factors as in previous studies with the implicit AAT (Roelofs et al., 2010; von Borries et

al., 2012). In order to control for multiple testing, we applied a False Discovery Rate (FDR) correction

as recommended for exploratory ANOVAs (Cramer et al., 2016). Color and condition were

counterbalanced across participants (green=pull for one half, green=push for the other half) and are

thus controlled for by design, though this randomization was not stratified by gender or other participant

characteristics. We inspected significant effects with post-hoc t-tests.

Error rates in the implicit AAT are often low, due to which between-condition differences in error

rates are not analyzed (Roelofs et al., 2010; von Borries et al., 2012). Here, too, errors were few and

unevenly distributed across conditions. Therefore, as in previous studies (Roelofs et al., 2010; von

Borries et al., 2012), we simply compared the mean error rate between groups. We used Welch's t-test,

which is robust to unequal variances and uneven sample sizes (Ruxton, 2006). Subsequently, we

computed Pearson correlation coefficients between AAT scores (between-condition differences in RT

and overall error rates) and each of the four clinical scales. We assessed the robustness of significant

correlations with bootstrap resampling to obtain 95% bias-corrected accelerated confidence intervals

(BCa CI) with 10000 iterations using the *bootstrap* package (version 2019.5). We performed all

analyses described in this section in R (version 3.6.1) running on R Studio (version 1.1.423).

Linear ballistic accumulator (LBA) modelling of reaction times

We subsequently implemented Linear Ballistic Accumulator (LBA) modelling on reaction time

data (Brown & Heathcote, 2008). LBA models assume that decisions stem from a sequential evidence

accumulation process (Fig. 3A). Evidence for each response option is gathered linearly by a separate

accumulator, which races against the other/s until one of them reaches a decision threshold. Evidence

accumulation starts after a variable period of non-decision time and its speed is given by the drift rate,

which is sampled from a normal distribution. The standard deviation of this distribution constitutes what

we here label drift noise, i.e., variability in the pace of evidence accumulation. In addition, the

accumulators might begin each trial from a different starting point, which is drawn from a uniform

distribution. Therefore, a response option will be taken more quickly if starting point and decision

threshold are nearer, if the drift rate is higher and less variable, and if the non-decision time is shorter.

LBA models are akin to the now-popular drift diffusion models (DDM), but are simpler and more

tractable computationally and thus well-suited for the relatively low amount of trials available in the

present dataset (see Heathcote and Hayes, 2012, for a detailed empirical comparison between LBA

and DDM).

Here, we fitted a series of LBA models with two accumulators (approach and avoidance) and

four parameters: decision threshold, starting point, drift rate, and drift noise. We tested a total of 16

models in which a given combination of these parameters was allowed to vary between the six

experimental conditions of interest: pull angry, pull happy, pull neutral, push angry, push happy, and

push neutral. We could not test for a modulation of experimental condition on non-decision time

because models including this effect failed to converge in most subjects. See Table 2 for a summary of

all models. We fitted each model on the reaction time data of each individual participant using full

information maximum likelihood estimation as implemented in the *glba* package version 0.2

(https://CRAN.R-project.org/package=glba). We used raw RT excluding errors and responses quicker

than 150ms or slower than 1s. For model comparison we inspected which model yielded the lowest

Bayesian Information Criterion (BIC) values across participants. BIC is a standard fit measure that

penalizes model complexity (Burnham & Anderson, 2004; Raftery, 1995). Our model fitting and

comparison approach is highly comparable to that of a recent DDM study on social approach-avoidance

decisions (Mennella et al., 2020). Afterwards, we simulated data per group using the *rlba()* function and

the average parameter estimates from the winning model. We next ran an ANOVA on the parameters

of the winning model with factors emotion, movement, and group in order to test which model

parameters varied as a function of group (Mennella et al., 2020). Finally, we compared parameters

between groups with independent-samples Welch t-tests. We used R (version 3.6.1) running on R

Studio (version 1.1.423) for all analyses in this section.

Neuroimaging data acquisition and analysis

Structural brain volumes were recorded at the Intervention center at Oslo University hospital

(Norway) on a Philips Ingenia 3-T scanner. We acquired structural images with a T1-weighted 3D turbo

gradient-echo sequence with the following settings: repetition time (TR)=1.900ms, echo time

(TE)=2.23ms, flip angle=8°, voxel size=1mm^3, field-of-view (FOV)=256x256mm. Members of the team

at the University of Oslo, trained in lesion reconstruction, manually delineated lesion masks on each

patient's anatomical images. We normalized these masks as recommended for lesioned brains

(Ripollés et al., 2012) and created lesion overlap maps using MRIcron (Rorden & Brett, 2000). We

further inquired which Brodmann Areas (BA) were damaged, following previous work (Jenkins et al.,

2014). We extracted all BA masks from the Wake Forest University PickAtlas (Maldjian et al., 2003)

and co-registered them with the Montreal Neurological Institute-152 template to ensure that lesions and

masks were in the same space (Jenkins et al., 2014). We then computed the number of overlapping

voxels between each lesion and BA mask. For each non-intact BA, we report the number of patients

with damage in that region along with the mean percentage of damaged tissue (Table 1; see also

Participants and lesion localization above).

Figure 1. A) Lesion overlap. Warmer colors depict more overlap between patients. Peak overlap was located in x=4, y=58, z=-14 (Montreal Neurological Institute space). **B)** Schematic depiction of the implicit Approach Avoidance Task (AAT). Subjects had to either push or pull a joystick in response to the color of the presented face while ignoring its facial expression (angry, happy, or neutral), gaze (direct or averted), gender (male or female), and identity (eight actors). Pushing made faces shrink in size, whereas pulling made them grow larger. The 384 trials were self-paced.

Table 1: Lesioned Brodmann areas

Brodmann area	N	% damage (M $\pm$ SE)
6	3	0.99 ± 0.89 %
8	3	4.87 ± 4.30 %
9	12	5.07 ± 1.93 %
10	12	18.44 ± 3.35 %
11	10	7.62 ± 2.16 %
13	4	0.55 ± 0.46 %
24	9	3.72 ± 0.90 %
25	8	18.73 ± 3.54 %
32	12	20.97 ± 4.48 %
45	4	0.85 ± 0.44 %
46	6	2.97 ± 1.79 %
47	7	1.95 ± 0.95 %

N patients: number of patients (out of 13) with damage in a given region; % damage: average percentage of lesioned tissue across patients; M: mean; SE: standard error.

We also inspected whether lesion size was linked with reaction times and error rates in the

task. We correlated lesion size with behavioral parameters showing a group difference in the AAT and

obtained the 95% bootstrapped CIs with 10000 iterations using the *bootstrap* R package to assess

these effects' robustness. We proceeded identically with all lesioned Brodmann areas, and compared

the lesion-behavior correlations with each other using the William's test as implemented in the *r.test()*

function from the *psych* R package.

Results

Approach-Avoidance Task (AAT) results

In our primary analysis of reaction times we observed main effects of group ($F_{1,42}$=11.92,

p=.001, pFDR=.010) and emotion ($F_{2,84}$=6.89, p=.001, pFDR=.010) which were qualified by an emotion

x movement interaction that did not survive multiple comparison correction ($F_{2,84}$=4.36, p=.015,

pFDR=.083), and, crucially, by a group x emotion x movement interaction ($F_{2,84}$=12.64, p<.001,

pFDR<.001). In order to dissect the latter three-way interaction, we computed the difference between

push and pull (i.e., avoidance minus approach) for each emotion and inspected for differences between

emotion categories in each group, following previous work (Roelofs et al., 2010; von Borries et al.,

2012). As shown in Fig. 2A, VMPFC patients showed a stronger approach bias toward angry relative

to both happy (t_{12}=3.17, p=.008) and neutral faces (t_{12}=4.32, p<.001), with no difference between happy

and neutral faces (p=.416). In comparison (Fig. 2B), controls showed a trend-level avoidant bias for

angry relative to neutral faces (t_{30}=1.75, p=.089), with no further differences between categories (all

p>.272). Thus, VMPFC patients were generally slower when pushing angry faces away relative to

pulling them close.

In order to ascertain whether these effects were predominantly driven by approach or

avoidance, we computed the difference in reaction times between emotions separately for push and

pull movements in each group. Regarding approach movements, VMPFC patients were faster to pull

angry relative to neutral (t_{12}=3.66, p=.003) but not happy faces (p=.107). Controls showed no between-

emotion differences in pull movements (all p>.278). For avoidance movements, VMPFC patients were

slower to push angry relative to happy faces (t_{12}=2.88, p=.013) but comparably fast when pushing angry

and neutral ones (p=.284). Controls were quicker to avoid angry as compared to neutral faces (t_{30}=2.27,

p=.030) but not happy ones (p=.605). Therefore, controls specifically showed avoidance of angry in

comparison with neutral expressions. In contrast, VMPFC patients showed increased approach of angry

relative to neutral faces, and reduced avoidance of angry as compared to happy ones. We used these

significant between-emotion differences for later correlation analyses, as they index the increased

threat approach (pull angry minus pull neutral) and reduced threat avoidance (push angry minus push

happy) demonstrated by VMPFC patients.

Additionally, there was an emotion x gaze interaction across the whole sample ($F_{4,168}$=4.63,

p=.001, pFDR=.011). We computed the difference in reaction times between direct and averted gaze

and compared between emotions over all participants to further investigate this effect. The interaction

was driven by slower reactions to directly-gazing neutral faces relative to happy (t_{43}=3.09, p=.003) and,

at trend level, angry ones (t_{43}=1.81, p=.077).

We subsequently compared error rates between groups. Although both groups performed the

task well, VMPFC patients committed about twice as many errors (6.87±1.13%) than healthy controls

(3.47±0.46%), $t_{16.14}$=2.77, p=.013 (Fig. 1C).

Figure 2. A) Patients with ventromedial prefrontal cortex (VMPFC) lesions showed an approach bias (reaction times [RT] for push minus pull) towards angry relative to happy and neutral faces. **B)** Healthy controls (HC) showed no bias in either direction, with a trend towards avoidance of angry relative to neutral faces. **C)** VMPFC patients made more errors than HC. **D)** Shorter RT for pull angry minus pull neutral trials were linked with greater self-rated impulsivity in VMPFC patients. **D)** Shorter RT for pull angry minus pull neutral trials were correlated with greater disinhibition in VMPFC patients. **F)** Error rates were correlated with all clinical self-reports in VMPFC patients, including greater self-rated disinhibition. ~p<.1, *p<.05, **p<.01, ***p<.001. BCa CI: Bias-Corrected accelerated Confidence Intervals obtained with bootstrapping.

Correlations between task scores and clinical scales

We then inspected for associations between clinical scales and task-derived scores, with the

aim of testing the clinical relevance of approach-avoidance biases as measured with the AAT. The

approach bias for angry minus neutral faces was linked with increased self-reported impulsivity (Fig.

1D; r=-.63, p=.020, 95% BCa CI=[-.85, -.28]), and greater disinhibition (Fig. 1E; r=-.57, p=.041, 95%

BCa CI=[-.85, -.15]), but there were no correlations with either of the other two clinical scales, or

between the angry push minus happy push difference and any of the scales (all p>.160). Error rates

were associated with impairment in all scales, namely impulsivity (r=.57, p=.040, 95% BCa CI=[.05,

.85]), disinhibition (Fig. 1F; r=.84, p<.001, 95% BCa CI=[.58, .95]), difficulties in emotional control (r=.59,

p=.033, 95% BCa CI=[.23, .78]), and worse self-monitoring (r=.70, p=.007, 95% BCa CI=[.25, .90]).

Correlations between lesion anatomy and AAT scores

Subsequently, we tested whether task-derived response biases were linked with lesion size.

Patients with larger lesions were quicker to approach angry relative to neutral faces (r=-.77, p=.001,

95% BCa CI=[-.92, -.55]). Correlations between this bias and the percentage of lesioned tissue were

negative and sizeable for each individual Brodmann area (between r=-.25 and r=-.70) and did not

significantly differ from each other (all p>.099). This indicates that the tendency to approach angry as

compared to neutral faces cannot be specifically attributed to any of these subregions. Lesion size was

not correlated with the push angry minus push happy difference (p=.369) or with error rates (p=.469).

Lesion extension was thus exclusively associated with threat approach, but not with the reduced threat

avoidance and increased error rates displayed by VMPFC patients.

Linear Ballistic Accumulator (LBA) modelling results

Next, we turned to Linear Ballistic Accumulator (LBA) modelling in order to uncover which latent

decision parameters might account for VMPFC patients' response patterns. We provide the complete

list of models in Table 2. The winning model assumed that emotional expression and movement

modulated drift rates exclusively. This model had the lowest Bayesian Information Criterion (BIC) across

subjects (median BIC=-646.62, k=10 free parameters) and was the best-fitting model in all 13 VMPFC

patients as well as in 90% (28/31) of control participants. According to model-comparison guidelines

(Burnham & Anderson, 2004; Raftery, 1995), the evidence for this model can be considered substantial

relative to the two next best-fitting (and slightly more complex) models, one assuming an effect of

emotional expression on drift rate and drift noise (median BIC=-629.92, k=15 free parameters), and one

in which emotional expression impacted drift rate and decision threshold (median BIC=-629.57, k=15

free parameters). The winning model could reproduce reaction times in pull angry trials with a precision

of around ~30-50ms across successive simulations for both VMPFC patients (example mean simulated

data=495ms; mean real data=544ms) and control participants (example mean simulated data=593ms;

mean real data=624ms).

An ANOVA on parameters of the winning model revealed main effects of group ($F_{1,42}$=15.40,

p<.001, pFDR<.001), emotion ($F_{2,84}$=653.88, p<.001, pFDR<.001), and movement ($F_{1,42}$=129.80,

p<.001, pFDR<.001) as well as significant pairwise interactions between all factors (group x movement:

$F_{2,84}$=15.76, p<.001, pFDR<.001; group x emotion: $F_{2,84}$=4.10, p=.019, pFDR=.019; emotion x

movement: $F_{2,84}$=134.10, p<.001, pFDR<.001). These effects were nonetheless qualified by a

significant emotion x movement x group interaction ($F_{2,84}$=11.31, p<.001, pFDR<.001), paralleling

reaction times.

Table 2: Summary of Linear Ballistic Accumulator models tested

Modulated parameters in model	K	Median BIC
Threshold, starting point, drift rate, drift noise	25	-546.16
Threshold, starting point, drift rate	20	-575.52
Threshold, starting point, drift noise	20	-75.84
Threshold, drift rate, drift noise	20	-606.32
Starting point, drift rate, drift noise	20	-606.47
Threshold, starting point	15	-117.53
Threshold, drift rate	15	-629.57
Threshold, drift noise	15	-119.90
Starting point, drift rate	15	-624.78
Starting point, drift noise	15	-157.82
Drift rate, drift noise	15	-629.92
Threshold	10	-161.38
Starting point	10	-107.24
Drift rate	**10**	**-646.62**
Drift noise	10	-172.87
Null model	5	-169.83

K: number of free parameters; BIC: Bayesian Information Criterion. The model marked in **bold** had the best fit to the data across participants.

We subsequently tested which parameters differed between groups (Table 3). Controls had

negative response drifts when pulling angry faces close, whereas the mean value for this parameter

was centered around zero in VMPFC patients (Fig. 3B, left; $t_{17.46}$=3.51, p=.002). VMPFC patients also

showed lower drift rates than control participants when pushing angry faces away (Fig. 3B, right;

$t_{15.56}$=2.92, p=.010). Therefore, response drifts in VMPFC patients were weaker when avoiding angry

faces and relatively less negative (i.e. centered around null) when approaching them. VMPFC patients

also had reduced response drifts when pushing happy faces away (Fig. 2C, right; $t_{41.99}$=2.29, p=.026),

but not when pulling them close (Fig. 2C, left; p=.258). This pattern was also present at trend level for

neutral expressions (Fig. 2D; avoid: $t_{39.10}$=1.85, p=.070; approach: p=.723). Thus, VMPFC patients had

generally lower drift rates than controls during avoidance movements, especially for angry faces.

Regarding the remaining parameters, the patient group displayed reduced drift noise (Fig. 2E;

$t_{41.11}$= 3.42, p=.001; HC: 0.18± 0.02, VMPFC: 0.08±0.01), and non-decision times (Fig. 2F; $t_{13.58}$=2.54,

p=.023; HC: -.15±.10, VMPFC: -1.21±.40). There were no group differences in decision threshold

(p=.126) or starting point (p=.364). Hence, evidence accumulation began earlier and was less variable

across conditions in VMPFC patients.

Figure 3. A) Schematic depiction of a Linear Ballistic Accumulator (LBA) model, which operationalizes decisions as the result of a sequential evidence accumulation process. The model assumes separate, competing accumulators for each response option, with faster decisions when the response threshold is lower, starting point is higher, non-decision time is shorter, and the drift towards a given option is stronger and less variable (i.e. higher drift rate and lower drift noise). We estimated the parameters from each participant's reaction time distribution with a maximum likelihood algorithm. **B)** Ventromedial prefrontal cortex (VMPFC) patients showed less negative (i.e. around zero) drift rates than healthy controls (HC) when pulling angry faces close (left), and lower drift rates when pushing angry faces away (right). **C)** VMPFC patients had lower drift rates than HC when pushing happy faces away. **D)** VMPFC patients displayed trend-level lower drift rates than HC when avoiding neutral faces. **E)** VMPFC patients had lower drift noise. **F)** VMPFC patients showed shorter non-decision times. A.u.: arbitrary units. ~p<.1, *p<.05, **p<.01.

Table 3: Group-wise means and standard errors of free parameters from the winning model

Parameter	HC	VMPFC
Drift rate pull angry**	-.24±.03	.04±.07
Drift rate pull happy	-.01±.009	-.001±.005
Drift rate pull neutral	-.002±.008	-.005±.005
Drift rate push angry*	1.46±.06	.94±.16
Drift rate push happy*	.04±.02	-.01±.01
Drift rate push neutral	.03±.02	-.01±.01
Drift noise**	.18±.02	.08±.01
Starting point	.13±.02	.08±.03
Threshold	.87±.14	1.56±.39
Non-decision time*	-.15±.10	-1.21±.40

HC: healthy controls; VMPFC: orbitofrontal cortex patients. Asterisks denote significant between-group differences in parameter estimates at *p<.05 or **p<.01.

In a final exploratory analysis, we tested for linear associations between LBA parameters

altered in VMPFC patients and clinical impairment. For drift rates, we limited these analyses to threat

approach (pull angry minus pull neutral) and threat avoidance (push angry minus push happy), as these

were the same contrasts that we computed for correlations with reaction times. We also correlated drift

noise and non-decision times with clinical scores. There were no associations between either score

and any of the clinical scales (all p>.234).

Discussion

Maladaptive social behavior is common after ventromedial prefrontal cortex (VMPFC) damage

(Anderson et al., 2006; Blair, 2004), but the neurocognitive processes underlying these symptoms

remain elusive. Here, we tested whether patients with acquired VMPFC lesions show altered automatic

responses to emotional facial expressions. VMPFC patients displayed both reduced avoidance of, and

increased approach to angry faces. Modelling of reaction times indicated that these biases are due to

differences in stimulus processing rather than to pre-existing preferences for either type of stimuli.

Between-group comparisons revealed relatively slower evidence accumulation when avoiding angry

faces in VMPFC patients relative to controls. Moreover, patients lacked the negative response drifts

that controls showed during approach of angry expressions. VMPFC patients further evidenced less

variable and earlier-starting evidence accumulation. The approach bias in VMPFC patients was

associated with self-reported clinical measures of impulsive and disinhibited behavior. Patients also

committed more errors, which was in turn correlated with greater self-reported impulsivity, disinhibition,

problems in emotional control, and worse self-monitoring. Finally, larger lesions were linked with a

relatively more pronounced approach bias to angry faces, but not with error rates or avoidance biases.

All in all, these findings suggest that VMPFC damage can precipitate maladaptive behavior by altering

the implicit processing of threatening social information during action selection.

VMPFC lesions increase approach and reduce avoidance of threatening stimuli

Our findings expand on a previous report indicating that VMPFC-damaged individuals report

negative facial expressions to be more approachable (Willis et al., 2010). Here, we showed that this

translates into observable, automatic motor behavior, such that VMPFC patients were quicker to

actively approach angry faces (i.e., pull them towards themselves), but slower to avoid them (i.e., push

them away). Reduced implicit avoidance of angry faces has been reported in psychopathic offenders

(von Borries et al., 2012), who also display dampened physiological reactivity to threatening distractors

(Newman et al., 2010) and deficits in VMPFC-dependent tasks (Blair, 2010). Therefore, VMPFC

dysfunction seems to confer both lower threat aversion and enhanced threat approach, features that

may facilitate antisocial behavior in some individuals.

The present results broadly converge with clinical (Blair, 2004), volumetric (Chester et al.,

2017), and functional (Beyer et al., 2015; Gilam et al., 2015) studies asserting that the VMPFC is

essential for the regulation of aggressive urges. Our data further indicate that the VMPFC does not

merely suppress automatic impulses but rather directs the course of approach-avoidance reactions, in

line with recent proposals (Hiser & Koenigs, 2018; Rudebeck & Rich, 2018), and with the well-known

association between damage to this region and disadvantageous decision-making (Koenigs & Tranel,

2007). Given that the VMPFC is involved in the anticipation and evaluation of actions related to certain

stimuli (R. C. Wilson et al., 2014), we suggest that VMPFC dysfunction gives rise to an altered

processing of threat signals. Specifically, it might be that VMPFC damage compromises the prediction

of behavioral outcomes associated with potentially punishing stimuli, i.e., tagging angry faces as neutral

or even potentially rewarding (Rudebeck & Murray, 2014). These abnormal value forecasts can in turn

enable the impulsive, rule-breaking behavior that characterizes the sequelae of some VMPFC lesions.

In line with the latter statement, approach towards angry relative to neutral faces was linked

with greater self-reported disinhibition and impulsive behavior. Paralleling our results, it has been

reported that patients with borderline personality disorder, who regularly engage in antagonistic and

aggressive behavior, also show an approach bias to angry faces (Bertsch et al., 2018) and comparable

levels of impulsivity and self-reported anger as those of VMPFC patients (Berlin et al., 2005). Similarly,

healthy individuals with high trait anger are quicker to approach angry relative to happy faces (Veenstra

et al., 2017). The current results thus provide further evidence that threat signals might act as appetitive

stimuli for individuals with externalizing symptomatology (Chester, 2017), and further add that VMPFC

lesions might precipitate such dysfunctional evaluation processes. Although VMPFC patients'

externalizing behavior is typically less severe than that of clinically antisocial individuals (Berlin et al.,

2005), lesions to the VMPFC are the most frequently associated with subsequent criminality (Darby et

al., 2018). Therefore, VMPFC lesions are likely to confer at least some risk for antisocial behavior

through other mediating features such as problems in value representation (Blair, 2010), as the present

data support.

Of note, the response tendencies observed in VMPFC patients were independent of gaze

direction. This pattern deviates from previous studies reporting group-specific approach-avoidance

biases exclusively for directly-gazing angry faces (Roelofs et al., 2010; von Borries et al., 2012). Hence,

the present findings tentatively suggest that VMPFC lesions might be associated with reduced

sensitivity to gaze direction. We did find, however, that straight-looking neutral faces were linked with

slower reaction times across the whole sample irrespective of movement type. The latter observation

insinuates that neutral expressions, due to their inherent ambiguity (Blasi et al., 2009), are more

thoroughly evaluated when directed to oneself.

Importantly, VMPFC patients performed generally worse than controls in the implicit approach-

avoidance task (AAT), which is suggestive of difficulties in ignoring task-irrelevant stimulus features.

This observation concurs with other studies in showing that VMPFC patients are more susceptible to

distraction by to-be-ignored stimulus characteristics (Kuusinen et al., 2018; Mäki-Marttunen et al.,

2017), and agrees with the general idea that VMPFC damage hinders the implementation of goal-

directed behavior (Rudebeck & Rich, 2018). Moreover, error rates were associated with greater self-

reported impulsivity and disinhibition in VMPFC patients, as well as with worse emotional control and

self-monitoring. Such findings speak for the predictive validity of the AAT and support its potential

usefulness for assessing emotional dysfunction in neurological patients (Fricke & Vogel, 2020).

Neuroanatomical analyses showed that lesion size strongly predicted threat approach, and that

no specific prefrontal regions accounted for this effect. These analyses confirmed nevertheless that

lesions were mostly localized in the VMPFC, with additional extensive damage to the frontal pole. Thus,

the current data suggest that ventromedial and anterior prefrontal cortex are most strongly linked with

the observed threat approach bias. It has been suggested that all prefrontal subregions carry out

evaluative functions, with a relative local specialization for certain types of information (Dixon et al.,

2017). In the context of emotional control, the frontal pole has been postulated to monitor current and

alternative strategies in order to facilitate response switching as dictated by current task demands, e.g.

from approach to avoidance (Bramson et al., 2020; Koch et al., 2018). Therefore, we speculate that the

implementation of adaptive approach-avoidance behavior might rely on both VMPFC-dependent value

representations as well as emotional action monitoring in frontopolar areas.

VMPFC lesions affect latent decision parameters

We used Linear Ballistic Accumulator (LBA) modelling to delve deeper into the decision

processes underlying approach-avoidance responses in VMPFC patients. These analyses indicated

that emotional facial expressions modulated drift rates (i.e., the speed of evidence accumulation after

a stimulus appears) but no other parameters. These findings extend previous drift diffusion modelling

work using an explicit version of the AAT in which emotional expressions impacted not only drift rates

but also response thresholds and non-decision times (Tipples, 2019). Hence, the influence of emotional

expressions on latent decision variables may be less pronounced when facial expressions are to be

ignored. The present data do however fully dovetail previous modelling studies in that response drifts

were maximal when threatening stimuli were to be avoided (Krypotos et al., 2015; Mennella et al., 2020;

Tipples, 2019). Our results complement these findings by showing that angry faces automatically bias

evidence accumulation towards avoidance even in the absence of explicit response contingencies.

Between-group comparisons of model parameters revealed profound differences between

VMPFC patients and control participants. VMPFC patients showed near-zero drift rates when

approaching (i.e., pulling) angry faces, whereas healthy controls showed negative values in this

parameter. VMPFC lesions might thus eliminate a default bias against threat approach. In addition, we

observed weaker response drifts during avoidance responses (i.e., push movements) in patients

relative to controls. The group difference in this parameter was strongest for angry facial expressions

but also present in happy and, at trend level, neutral trials. Evidence accumulation leading to avoidance

decisions is hence more sluggish in VMPFC patients, and especially so in the presence of angry facial

expressions. Therefore, the incongruent approach behavior often observed in VMPFC patients (Perry

et al., 2016; Willis et al., 2010) might be partly attributable to an altered evidence accumulation process

in response to social signals. Specifically, evidence accumulation in VMPFC patients seems to lack a

bias against threat approach and is generally slower during avoidance. In control participants, in

contrast, the positive drift rates when pushing angry faces away might have outweighed the negative

drifts when pulling them close, resulting in threat avoidance. These observations agree with the idea

that the VMPFC encodes the currently relevant state-space (Stalnaker et al., 2015; R. C. Wilson et al.,

2014). Angry facial expressions should, on the basis of previous experience, evoke a representation of

possible negative outcomes and thereby facilitate avoidance, as seen in the reaction times of control

participants. This negative outcome representation is abolished after VMPFC lesions, presumably

producing the observed alterations in evidence accumulation and the resulting abnormal approach-

avoidance responses.

In addition, VMPFC patients displayed relatively shorter non-decision times and lower drift rate

variability irrespective of experimental condition. This implies that approach-avoidance decision

processes start earlier and are more rigid in VMPFC patients as compared to control participants. The

lower non-decision times are in consonance with the generally speeded responding and higher error

rates incurred by VMPFC patients, as well as with the enhanced impulsivity often observed in VMPFC-

damaged individuals (Berlin et al., 2004, 2005). On the other hand, the reduced drift rate variability

observed in patients parallels the deficits in goal-directed behavior subsequent to VMPFC damage, i.e.,

a failure to update stimulus value resulting in perseverative responses (Rudebeck et al., 2013;

Rudebeck & Murray, 2014).

Surprisingly, clinical impairment could be predicted by reaction times but not by any single LBA

parameter. Thus, VMPFC patients' emotional dysregulation appears to be more strongly influenced by

the combination of multiple latent decision processes (which putatively produce the observed reaction

times) rather than by any single one of them in isolation. The absence of group differences in starting

point or decision threshold is also noteworthy, as it indicates that the approach bias observed in VMPFC

patients is likely due to post-stimulus processing rather than to pre-existing response tendencies. This

finding agrees with a recent modeling study on approach-avoidance decisions in which the emotional

valence of the stimuli modulated drift rates but no other parameters in the model (Mennella et al., 2020).

Taken together, LBA results suggest that damage to the VMPFC might lead to rapid and invariant

evidence accumulation, which is in turn slower when avoiding threatening stimuli but relatively faster

when approaching these signals.

Limitations

The cross-sectional nature of the design, along with the reduced sample size common in

studies with focal lesion patients (Motzkin et al., 2015; Pujara et al., 2016), constrain the generalizability

of the present results. Special caution should be exercised regarding the correlations: even though we

used bootstrapping to assess their robustness, the ability of the implicit AAT to track interindividual

differences is uncertain due to the lack of data on this instrument's reliability (Hedge et al., 2018). In

general, effect sizes from discovery studies such as the present one should be assumed to be inflated

until replication or follow up studies permit a more precise estimation of the true effect (B. M. Wilson et

al., 2020). It should also be noted that there was no lesion control group, which renders the regional

specificity of the results uncertain. Nonetheless, anatomical analyses revealed that the strongest lesion

overlap was located in ventromedial and rostral-anterior aspects. Finally, due to time constraints, we

were not able to measure patients' explicit emotion recognition abilities, which are sometimes

(Heberlein et al., 2008) but not always (Willis et al., 2010) impaired in VMPFC patients. Control tasks

involving emotion recognition and distraction susceptibility as well as additional measures such as eye-

tracking (Goursaud & Bachevalier, 2020) would permit a more precise dissection of the mechanisms

underlying altered approach-avoidance behavior after VMPFC damage. This limitation is minimized by

the fact that the task did not require emotion recognition to be performed.

Conclusion

The present study provides insight on how VMPFC dysfunction impacts the processing of

threatening information during approach-avoidance decisions. This was manifested in altered evidence

accumulation in response to threatening stimuli in combination with markers of premature and inflexible

decision-making. Intervention programs to improve social functioning in VMPFC patients might

therefore benefit from a focus on correctly interpreting and reacting to emotional information as well as

on ameliorating impulsivity (Levine et al., 2008). In sum, our study demonstrates that VMPFC damage

can steer individuals towards maladaptive approach behavior by biasing the automatic evaluation of

threat signals.

Acknowledgements and funding

This study was funded by the German Science Foundation (grant number KR3691/5-1). MBR

was supported by a German Science Foundation stipend (grant number BU 3756/1-1). AKS, TE, and

IF were supported by the Research Council of Norway through a grant (project number 240389) and

through the Centres of Excellence scheme (project number 262762 RITMO). KR was supported by a

VICI grant (#453-12-001) from the Netherlands Organization for Scientific Research (NWO) and a

consolidator grant from the European Research Council (ERC_CoG-2017_772337). We thank Dr.

Torstein R. Meling for help with patient recruitment and Dr. Per Kristian Hol for clinical evaluation of the

patients' MRI scans. Dr. Buades-Rotger, Dr. Solbakk, Dr. Liebrand, Dr. Endestad, Dr. Funderud, Mr.

Siegwardt, Ms. Enter, Dr. Roelofs, and Dr. Krämer report no biomedical financial interests or potential

conflicts of interest.

References

Anderson, S. W., Barrash, J., Bechara, A., & Tranel, D. (2006). Impairments of emotion and real-world complex behavior following childhood- or adult-onset damage to ventromedial prefrontal cortex. *Journal of the International Neuropsychological Society*, *12*(2), 224–235. https://doi.org/10.1017/S1355617706060346

Barrash, J., Tranel, D., & Anderson, S. W. (2000). Acquired Personality Disturbances Associated With Bilateral Damage to the Ventromedial Prefrontal Region. *Developmental Neuropsychology*, *18*(3), 355–381. https://doi.org/10.1207/S1532694205Barrash

Beer, J. S., John, O. P., Scabini, D., & Knight, R. T. (2006). Orbitofrontal cortex and social behavior: Integrating self-monitoring and emotion-cognition interactions. *Journal of Cognitive Neuroscience*, *18*(6), 871–879. PubMed. https://doi.org/10.1162/jocn.2006.18.6.871

Berlin, H. A., Rolls, E. T., & Iversen, S. D. (2005). Borderline Personality Disorder, Impulsivity, and the Orbitofrontal Cortex. *American Journal of Psychiatry*, *162*(12), 2360–2373. https://doi.org/10.1176/appi.ajp.162.12.2360

Berlin, H. A., Rolls, E. T., & Kischka, U. (2004). Impulsivity, time perception, emotion and reinforcement sensitivity in patients with orbitofrontal cortex lesions. *Brain*, *127*(5), 1108–1126. https://doi.org/10.1093/brain/awh135

Bertsch, K., Roelofs, K., Roch, P. J., Ma, B., Hensel, S., Herpertz, S. C., & Volman, I. (2018). Neural correlates of emotional action control in anger-prone women with borderline personality disorder. *Journal of Psychiatry & Neuroscience : JPN*, *43*(3), 161–170. Pmc. https://doi.org/10.1503/jpn.170102

Beyer, F., Münte, T. F., Göttlich, M., & Krämer, U. M. (2015). Orbitofrontal Cortex Reactivity to Angry Facial Expression in a Social Interaction Correlates with Aggressive Behavior. *Cerebral Cortex*, *25*(9), 3057–3063. https://doi.org/10.1093/cercor/bhu101

Blair, R. J. R. (2004). The roles of orbital frontal cortex in the modulation of antisocial behavior. *Brain and Cognition*, *55*(1), 198–208. https://doi.org/10.1016/S0278-2626(03)00276-8

Blair, R. J. R. (2010). Psychopathy, frustration, and reactive aggression: The role of ventromedial prefrontal cortex. *British Journal of Psychology (London, England: 1953)*, *101*(Pt 3), 383–399. https://doi.org/10.1348/000712609X418480

Blasi, G., Hariri, A. R., Alce, G., Taurisano, P., Sambataro, F., Das, S., Bertolino, A., Weinberger, D. R., & Mattay, V. S. (2009). Preferential amygdala reactivity to the negative assessment of neutral faces. *Biological Psychiatry*, *66*(9), 847–853. PubMed. https://doi.org/10.1016/j.biopsych.2009.06.017

Bramson, B., Folloni, D., Verhagen, L., Hartogsveld, B., Mars, R. B., Toni, I., & Roelofs, K. (2020). Human Lateral Frontal Pole Contributes to Control over Emotional Approach–Avoidance Actions. *Journal of Neuroscience*, *40*(14), 2925–2934. https://doi.org/10.1523/JNEUROSCI.2048-19.2020

Brown, S. D., & Heathcote, A. (2008). The simplest complete model of choice response time: Linear ballistic accumulation. *Cognitive Psychology*, *57*(3), 153–178. PubMed. https://doi.org/10.1016/j.cogpsych.2007.12.002

Burnham, K. P., & Anderson, D. R. (2004). Multimodel Inference: Understanding AIC and BIC in Model Selection. *Sociological Methods & Research*, *33*(2), 261–304. https://doi.org/10.1177/0049124104268644

Chester, D. S. (2017). The Role of Positive Affect in Aggression. *Current Directions in Psychological Science*, *26*(4), 366–370. https://doi.org/10.1177/0963721417700457

Chester, D. S., Lynam, D. R., Milich, R., & DeWall, C. N. (2017). Physical aggressiveness and gray matter deficits in ventromedial prefrontal cortex. *Cortex*, *97*, 17–22. https://doi.org/10.1016/j.cortex.2017.09.024

Clithero, J. A., & Rangel, A. (2014). Informatic parcellation of the network involved in the computation of subjective value. *Social Cognitive and Affective Neuroscience*, *9*(9), 1289–1302. https://doi.org/10.1093/scan/nst106

Cramer, A. O. J., van Ravenzwaaij, D., Matzke, D., Steingroever, H., Wetzels, R., Grasman, R. P. P. P., Waldorp, L. J., & Wagenmakers, E.-J. (2016). Hidden multiplicity in exploratory multiway ANOVA: Prevalence and remedies. *Psychonomic Bulletin & Review*, *23*(2), 640–647. https://doi.org/10.3758/s13423-015-0913-5

Darby, R. R., Horn, A., Cushman, F., & Fox, M. D. (2018). Lesion network localization of criminal behavior. *Proceedings of the National Academy of Sciences*, *115*(3), 601. https://doi.org/10.1073/pnas.1706587115

Davidson, R. J., Putnam, K. M., & Larson, C. L. (2000). Dysfunction in the Neural Circuitry of Emotion Regulation—A Possible Prelude to Violence. *Science*, *289*(5479), 591–594. https://doi.org/10.1126/science.289.5479.591

Dixon, M. L., Thiruchselvam, R., Todd, R., & Christoff, K. (2017). Emotion and the prefrontal cortex: An integrative review. *Psychological Bulletin*, *143*(10), 1033–1081. https://doi.org/10.1037/bul0000096

Ekman, P., & Friesen, W. V. (1976). Pictures of Facial Affect. *Consulting Psychologists Press*. https://ci.nii.ac.jp/naid/10011335061/

Folloni, D., Sallet, J., Khrapitchev, A. A., Sibson, N., Verhagen, L., & Mars, R. B. (2019). Dichotomous organization of amygdala/temporal-prefrontal bundles in both humans and monkeys. *ELife*, *8*, e47175. https://doi.org/10.7554/eLife.47175

Fricke, K., & Vogel, S. (2020). How interindividual differences shape approach-avoidance behavior: Relating self-report and diagnostic measures of interindividual differences to behavioral

measurements of approach and avoidance. *Neuroscience & Biobehavioral Reviews*, *111*, 30–56. https://doi.org/10.1016/j.neubiorev.2020.01.008

Gilam, G., Lin, T., Raz, G., Azrielant, S., Fruchter, E., Ariely, D., & Hendler, T. (2015). Neural substrates underlying the tendency to accept anger-infused ultimatum offers during dynamic social interactions. *NeuroImage*, *120*(Supplement C), 400–411. https://doi.org/10.1016/j.neuroimage.2015.07.003

Goursaud, A.-P. S., & Bachevalier, J. (2020). Altered face scanning and arousal after orbitofrontal cortex lesion in adult rhesus monkeys. *Behavioral Neuroscience*, *134*(1), 45–58. https://doi.org/10.1037/bne0000342

Gross, J. J. (2015). Emotion Regulation: Current Status and Future Prospects. *Psychological Inquiry*, *26*(1), 1–26. https://doi.org/10.1080/1047840X.2014.940781

Heathcote, A., & Hayes, B. (2012). Diffusion versus linear ballistic accumulation: Different models for response time with different conclusions about psychological mechanisms? *Canadian Journal of Experimental Psychology = Revue Canadienne de Psychologie Experimentale*, *66*(2), 125–136. PubMed. https://doi.org/10.1037/a0028189

Heberlein, A. S., Padon, A. A., Gillihan, S. J., Farah, M. J., & Fellows, L. K. (2008). Ventromedial Frontal Lobe Plays a Critical Role in Facial Emotion Recognition. *Journal of Cognitive Neuroscience*, *20*(4), 721–733. https://doi.org/10.1162/jocn.2008.20049

Hedge, C., Powell, G., & Sumner, P. (2018). The reliability paradox: Why robust cognitive tasks do not produce reliable individual differences. *Behavior Research Methods*, *50*(3), 1166–1186. https://doi.org/10.3758/s13428-017-0935-1

Hiser, J., & Koenigs, M. (2018). The Multifaceted Role of the Ventromedial Prefrontal Cortex in Emotion, Decision Making, Social Cognition, and Psychopathology. *Biological Psychiatry*, *83*(8), 638–647. https://doi.org/10.1016/j.biopsych.2017.10.030

Jenkins, L. M., Andrewes, D. G., Nicholas, C. L., Drummond, K. J., Moffat, B. A., Phal, P. M., & Desmond, P. (2018). Emotional reactivity following surgery to the prefrontal cortex. *Journal of Neuropsychology*, *12*(1), 120–141. https://doi.org/10.1111/jnp.12110

Jenkins, L. M., Andrewes, D. G., Nicholas, C. L., Drummond, K. J., Moffat, B. A., Phal, P. M., Desmond, P., & Kessels, R. P. C. (2014). Social cognition in patients following surgery to the prefrontal cortex. *Psychiatry Research: Neuroimaging*, *224*(3), 192–203. https://doi.org/10.1016/j.pscychresns.2014.08.007

Koch, S. B. J., Mars, R. B., Toni, I., & Roelofs, K. (2018). Emotional control, reappraised. *Neuroscience & Biobehavioral Reviews*, *95*, 528–534. https://doi.org/10.1016/j.neubiorev.2018.11.003

Koenigs, M., & Tranel, D. (2007). Irrational economic decision-making after ventromedial prefrontal damage: Evidence from the Ultimatum Game. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *27*, 951–956. https://doi.org/10.1523/JNEUROSCI.4606-06.2007

Krypotos, A.-M., Beckers, T., Kindt, M., & Wagenmakers, E.-J. (2015). A Bayesian hierarchical diffusion model decomposition of performance in Approach–Avoidance Tasks. *Cognition and Emotion*, *29*(8), 1424–1444. https://doi.org/10.1080/02699931.2014.985635

Kuusinen, V., Cesnaite, E., Peräkylä, J., Ogawa, K. H., & Hartikainen, K. M. (2018). Orbitofrontal Lesion Alters Brain Dynamics of Emotion-Attention and Emotion-Cognitive Control Interaction in Humans. *Frontiers in Human Neuroscience*, *12*. https://doi.org/10.3389/fnhum.2018.00437

Levine, B., Turner, G. R., & Stuss, D. T. (2008). Rehabilitation of frontal lobe functions. In *Cognitive neurorehabilitation: Evidence and application, 2nd ed* (pp. 464–486). Cambridge University Press. https://doi.org/10.1017/CBO9781316529898.033

Løvstad, M., Funderud, I., Endestad, T., Due-Tønnessen, P., Meling, T. R., Lindgren, M., Knight, R. T., & Solbakk, A. K. (2012). Executive functions after orbital or lateral prefrontal lesions: Neuropsychological profiles and self-reported executive functions in everyday living. *Brain Injury*, *26*(13–14), 1586–1598. PubMed. https://doi.org/10.3109/02699052.2012.698787

Lucantonio, F., Stalnaker, T. A., Shaham, Y., Niv, Y., & Schoenbaum, G. (2012). The impact of orbitofrontal dysfunction on cocaine addiction. *Nature Neuroscience*, *15*(3), 358–366. https://doi.org/10.1038/nn.3014

Lundqvist, D., Flykt, A., & Öhman, A. (1998). The Karolinska directed emotional faces (KDEF). *CD ROM from Department of Clinical Neuroscience, Psychology Section, Karolinska Institutet*, *91*, 630.

Mäki-Marttunen, V., Kuusinen, V., Peräkylä, J., Ogawa, K. H., Brause, M., Brander, A., & Hartikainen, K. M. (2017). Greater Attention to Task-Relevant Threat Due to Orbitofrontal Lesion. *Journal of Neurotrauma*, *34*(2), 400–413. https://doi.org/10.1089/neu.2015.4390

Maldjian, J. A., Laurienti, P. J., Kraft, R. A., & Burdette, J. H. (2003). An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *NeuroImage*, *19*(3), 1233–1239. https://doi.org/10.1016/S1053-8119(03)00169-1

Mennella, R., Vilarem, E., & Grèzes, J. (2020). Rapid approach-avoidance responses to emotional displays reflect value-based decisions: Neural evidence from an EEG study. *NeuroImage*, *222*, 117253. https://doi.org/10.1016/j.neuroimage.2020.117253

Motzkin, J. C., Philippi, C. L., Wolf, R. C., Baskaya, M. K., & Koenigs, M. (2015). Ventromedial Prefrontal Cortex Is Critical for the Regulation of Amygdala Activity in Humans. *Biological Psychiatry*, *77*(3), 276–284. https://doi.org/10.1016/j.biopsych.2014.02.014

Newman, J. P., Curtin, J. J., Bertsch, J. D., & Baskin-Sommers, A. R. (2010). Attention moderates the fearlessness of psychopathic offenders. *Biological Psychiatry*, *67*(1), 66–70. Pmc. https://doi.org/10.1016/j.biopsych.2009.07.035

Perry, A., Lwi, S. J., Verstaen, A., Dewar, C., Levenson, R. W., & Knight, R. T. (2016). The role of the orbitofrontal cortex in regulation of interpersonal space: Evidence from frontal lesion and frontotemporal dementia patients. *Social Cognitive and Affective Neuroscience*, *11*(12), 1894–1901. https://doi.org/10.1093/scan/nsw109

Pujara, M. S., Philippi, C. L., Motzkin, J. C., Baskaya, M. K., & Koenigs, M. (2016). Ventromedial Prefrontal Cortex Damage Is Associated with Decreased Ventral Striatum Volume and Response to Reward. *The Journal of Neuroscience*, *36*(18), 5047. https://doi.org/10.1523/jneurosci.4236-15.2016

Raftery, A. E. (1995). Bayesian Model Selection in Social Research. *Sociological Methodology*, *25*, 111–163. JSTOR. https://doi.org/10.2307/271063

Ripollés, P., Marco-Pallarés, J., de Diego-Balaguer, R., Miró, J., Falip, M., Juncadella, M., Rubio, F., & Rodriguez-Fornells, A. (2012). Analysis of automated methods for spatial normalization of lesioned brains. *NeuroImage*, *60*(2), 1296–1306. https://doi.org/10.1016/j.neuroimage.2012.01.094

Roelofs, K., Putman, P., Schouten, S., Lange, W.-G., Volman, I., & Rinck, M. (2010). Gaze direction differentially affects avoidance tendencies to happy and angry faces in socially anxious individuals. *Behaviour Research and Therapy*, *48*(4), 290–294. https://doi.org/10.1016/j.brat.2009.11.008

Rorden, C., & Brett, M. (2000). Stereotaxic Display of Brain Lesions. *Behavioural Neurology*, *12*(4). https://doi.org/10.1155/2000/421719

Roth, R., Isquith, P., & Gioia, G. (2005). *Behavior Rating Inventory of Executive Function—Adult Version (BRIEF-A)* (Vol. 20). Psychological Assessment Resources.

Rudebeck, P. H., & Murray, E. A. (2014). The Orbitofrontal Oracle: Cortical Mechanisms for the Prediction and Evaluation of Specific Behavioral Outcomes. *Neuron*, *84*(6), 1143–1156. https://doi.org/10.1016/j.neuron.2014.10.049

Rudebeck, P. H., & Rich, E. L. (2018). Orbitofrontal cortex. *Current Biology*, *28*(18), R1083–R1088. https://doi.org/10.1016/j.cub.2018.07.018

Rudebeck, P. H., Saunders, R. C., Prescott, A. T., Chau, L. S., & Murray, E. A. (2013). Prefrontal mechanisms of behavioral flexibility, emotion regulation and value updating. *Nature Neuroscience*, *16*(8), 1140–1145. https://doi.org/10.1038/nn.3440

Stalnaker, T. A., Cooch, N. K., & Schoenbaum, G. (2015). What the orbitofrontal cortex does not do. *Nat Neurosci*, *18*(5), 620–627. https://doi.org/10.1038/nn.3982

Tipples, J. (2019). Recognising and reacting to angry and happy facial expressions: A diffusion model analysis. *Psychological Research*, *83*(1), 37–47. https://doi.org/10.1007/s00426-018-1092-6

Veenstra, L., Schneider, I. K., Bushman, B. J., & Koole, S. L. (2017). Drawn to danger: Trait anger predicts automatic approach behaviour to angry faces. *Cognition and Emotion*, *31*(4), 765–771. https://doi.org/10.1080/02699931.2016.1150256

von Borries, A. K. L., Volman, I., de Bruijn, E. R. A., Bulten, B. H., Verkes, R. J., & Roelofs, K. (2012). Psychopaths lack the automatic avoidance of social threat: Relation to instrumental aggression. *Psychiatry Research*, *200*(2), 761–766. https://doi.org/10.1016/j.psychres.2012.06.026

Waid-Ebbs, J. K., Wen, P.-S., Heaton, S. C., Donovan, N. J., & Velozo, C. (2012). The item level psychometrics of the behaviour rating inventory of executive function-adult (BRIEF-A) in a TBI sample. *Brain Injury*, *26*(13–14), 1646–1657. https://doi.org/10.3109/02699052.2012.700087

Whiteside, S. P., & Lynam, D. R. (2001). The Five Factor Model and impulsivity: Using a structural model of personality to understand impulsivity. *Personality and Individual Differences*, *30*(4), 669–689. https://doi.org/10.1016/S0191-8869(00)00064-7

Whiteside, S. P., Lynam, D. R., Miller, J. D., & Reynolds, S. K. (2005). Validation of the UPPS impulsive behaviour scale: A four-factor model of impulsivity. *European Journal of Personality*, *19*(7), 559–574. https://doi.org/10.1002/per.556

Willis, M. L., Palermo, R., Burke, D., McGrillen, K., & Miller, L. (2010). Orbitofrontal cortex lesions result in abnormal social judgements to emotional faces. *Neuropsychologia*, *48*(7), 2182–2187. https://doi.org/10.1016/j.neuropsychologia.2010.04.010

Wilson, B. M., Harris, C. R., & Wixted, J. T. (2020). Science is not a signal detection problem. *Proceedings of the National Academy of Sciences*, *117*(11), 5559–5567. https://doi.org/10.1073/pnas.1914237117

Wilson, R. C., Takahashi, Y. K., Schoenbaum, G., & Niv, Y. (2014). Orbitofrontal cortex as a cognitive map of task space. *Neuron*, *81*(2), 267–279. https://doi.org/10.1016/j.neuron.2013.11.005

A

B

Push

Time

Pull

n

1   13

A Linear Ballistic Accumulator Model

B Drift rate angry

C Drift rate happy

D Drift rate neutral

E Drift noise

F Non-decision time