Title: Vegetative phase change in *Populus tremula x alba*.

Authors: Erica H. Lawrence[1*], Aaron R. Leichty[1,2], Cathleen Ma[3], Steven H. Strauss[3] and R.

Scott Poethig[1]

[1]Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania, 19104 USA

[2]Department of Plant Biology, University of California, Davis, California, 95616 USA

[3]Department of Forest Ecosystems and Society, Oregon State University, Corvallis, Oregon

97331

*Corresponding author: lawrence.erica.h@gmail.com

Summary

• Plants transition through juvenile and adult phases of vegetative development in a process

known as vegetative phase change (VPC). In poplars (genus *Populus*) the differences

between these stages are subtle, making it difficult to determine when this transition

occurs. Previous studies of VPC in poplars have relied on plants propagated *in vitro,* so

the normal nature of this process is still unknown.

• We examined developmental morphology of seed-grown and *in vitro* derived *Populus*

tremula x alba (clone 717-1B4), and compared the phenotype of these, to plants over-

and under-expressing miR156, the master regulator of VPC.

• In seed-grown plants, most traits changed from node-to-node during the first 3 months of

development but remained relatively constant starting around node 25. Most of these

traits remained unchanged in clones over-expressing miR156, demonstrating they are

regulated by the miR156/SPL pathway. Leaf fluttering--a characteristic of *Populus*

tremula ("quaking aspen")--is one of these miR156-regulated traits.

•   Vegetative development in plants grown from culture mirrored that of seed-grown plants,

32   allowing direct comparison between plants often used in research and those found in

33   nature. These results provide a foundation for further research on the role of VPC in the

34   ecology and evolution of this economically important genus.

Introduction

During their ontogeny, plants transition through distinct phases, which is reflected in

morphological variation in the organs produced at different times in development. These

developmental transitions include the visually obvious transition between vegetative and

reproductive growth as well as the often more subtle change between juvenile and adult

vegetative stages, known as vegetative phase change (Ashby, 1948; Brink, 1962; Doorenbos,

1965; Poethig, 2013). In species with abrupt changes in vegetative morphology—termed

heteroblastic—the distinction between the juvenile and adult phases is unambiguous (Zotz *et al.*,

2011). For example, many species of *Acacia* transition from producing bipinnate compound

leaves to producing a simple undivided leaf termed a phyllode (Walters & Bartholomew, 1984).

Other examples include *Eucalyptus globulus*, which transitions from producing ovate, wax-

covered leaves to producing lanceolate glaucous leaves (James & Bell, 2001) and *Ipomoea*

caerulea, which transitions from producing monolobed to trilobed leaves (Ashby, 1948). The

juvenile-to-adult transition is more difficult to identify in species that experience less drastic

changes in leaf and shoot morphology, with the result that we know very little about the duration

of these phases in different taxa, or the relationship between this developmental transition and

other aspects of plant life history, such as the response of plants to various biotic and abiotic

conditions. However, even in these so-called *homoblastic* species (Goebel, 1900), organs

produced early in shoot development differ from organs produced later in shoot development in a

variety of morphological, anatomical, and physiological traits (Feng *et al.*, 2016). Identifying

these traits and determining which are components of vegetative phase change and which are

regulated by other factors, such as plant size or chronological age, is an important problem in

plant development.

In *Arabidopsis* and most other plants, the transition from the juvenile to the adult phase of

vegetative development is regulated by the closely related microRNAs, miR156 and miR157

(Poethig, 2013; He *et al.*, 2018). These miRNAs directly repress members of the Squamosa

Promoter Binding Protein-Like (SPL) family of transcription factors (Rhoades *et al.*, 2002;

Schwab *et al.*, 2005) which promote adult traits (Xu *et al.*, 2016). SPL expression is inhibited

during the juvenile phase by high levels of miR156/miR157. As plants age, miR156/miR157

expression declines, allowing for the increased expression of their SPL targets. This coordinated

change in miR156 and SPL gene expression leads to the onset of adult traits, with different traits

typically appearing at slightly different times in shoot development, depending on their

sensitivity to the level of SPL proteins (He *et al.*, 2018).

Vegetative phase change is a particularly important in long-lived perennial species, such

as trees, because the ability of these species to survive long enough to become reproductively

competent depends on their ability to respond to changing conditions over a long period of time.

Poplars are of considerable economic and ecological importance and have become popular

model systems for tree biology because they can be readily propagated in culture and are

susceptible to *Agrobacterium*-mediated transformation (Taylor, 2002). Temporal variation in leaf

and shoot morphology, and the phenotype of plants over-expressing miR156, have been

described in clones of *Populus x canendensis* (Wang *et al.*, 2011) and *P. tremula x alba*

(Rubinelli *et al.*, 2013)*.* However, the significance of this information for vegetative phase

change in *Populus* is still unknown because the developmental morphology of seed-grown plants

has never been described. Here, we compare the vegetative phenotype of seed-grown *Populus*

tremula x alba with plants originating from explants in tissue culture and identify phase-specific

traits by characterizing the phenotype of plants over- and under-expressing miR156. We found

that shoot development in plants propagated *in vitro* closely resembles the development of seed-

grown plants, and that vegetative phase change occurs very early in both types of shoots. Most of

the traits we examined changed significantly as the shoot developed, and most of these traits

were sensitive to the level of miR156. One of these was leaf fluttering, a trait that has a

significant effect on photosynthetic efficiency (Roden & Pearcy, 1993b). Our results provide

basic information about the nature of vegetative phase change in *Populus tremula x alba and*

demonstrate that shoots regenerated *in vitro* can be used as models of primary shoot development

in this species.

Materials and Methods

Plant transformation

Assembly of Ubi10:AthMIR156a and Ubi10:MIM156 constructs have previously been

described (Feng *et al.*, 2016). Additionally, a second MIM156 construct was designed using

Golden Gate Cloning (Engler *et al.*, 2014). Further details of construct assembly can be found in

Table S1.

Poplar line 717-1B4 was transformed using *Agrobacterium tumefaciences (strain AGL1)*

following the protocol described in Filichkin *et al.* (2006). Briefly, leaf and stem explants were

selected on callus-induction, shoot induction, shoot elongation and rooting media using BASTA.

A total of 30 Ubi:AthMIR156a events and 25 Ubi:MIM156 events were isolated. Of these, two

lines for each construct type were selected based on levels of miR156 in 5 cm leaf primordia

(data not shown). For the Ubi:AthMIR156a and Ubi:MIM156 constructs made with the pCambia

3300 backbone, the Ubi:AthMIR156a construct had a higher rate of producing transgenic shoots

(percent of explants forming shoots was 52% versus 33.9%). We did not track efficiency for the

other MIM156 construct.

Plant material and growth conditions

Populus tremula x alba line 717-1B4, two independent miR156 over-expressor lines, 40

and 78, and two independent MIM156 lines, 22 and 84, were obtained by *in vitro* propagation

and hardening on propagation media as described in Meilan and Ma (2006). Plants were then

transplanted to Fafard-2 growing mix (Sungro Horticulture, Massachusetts, USA) in 0.3-L pots

in the greenhouse at the University of Pennsylvania (39.9493°N, 75.1995°W, 22.38 m a.s.l.) and

kept in plastic bags for increased humidity for 2 weeks. Plants were transferred to 4.2-L pots

with Fafard-52 growing mix 3 weeks later and fertilized with Osmocote 14-14-14 (The Scotts

Company, Marysville, OH, USA), supplemented by weekly applications of Peters 20-10-20 (ICL

Fertilizers, Dublin, OH, USA). Plants were grown at a temperature ranging from 22 and 27°C on

a 16 hr photoperiod, with illumination provided by a combination of natural light and 400-W

metal halide lamps (P.L. Light Systems, Ontario, Canada) with daily irradiances between 300

and 1,500 µmol m^{-2} s^{-1}. Planting of individuals was staggered across three months with the

position of plants in the greenhouse randomized and individuals rotated frequently.

Populus tremula x alba seeds were purchased from Sheffield's Seed Company (Locke,

NY), and were germinated on a layer of vermiculite on top of Fafard-2 growing mix in 0.64-L

pots in the greenhouse under conditions described above. Seedlings were transplanted one month

after germination into Fafard-52 growing mix in 1.76-L pots, fertilized with Osmocote 14-14-14.

They were transplanted to 4.2-L pots 3 months later. Planting of individuals was staggered across

five months and individuals were rotated within the greenhouse frequently.

Mir156/7 expression

The abundance of miR156 and miR157 RNA was measured in <1 cm leaf primordia.

Total RNA was isolated using the Spectrum™ Plant Total RNA Kit (Sigma-Aldrich), according

to the manufacturer's instructions. RNA levels were measured by RT-qPCR, using primers

specific for miR156, miR157 and miR159 in combination with the stem-loop RT primers

described in Varkonyi-Gasic et al. (2007) (Table S2). miR159 was used as an endogenous

control (Leichty & Poethig, 2019) and validated across all time points in seed-grown plants.

cDNA was synthesized using Invitrogen SuperScript III following the methods of Varkonyi-

Gasic et al. (2007). Platinum Taq (Invitrogen) was used with the Roche universal hydrolysis

probe #21, and a three-step amplification protocol. Relative measures of abundance were

calculated using the $2^{-\Delta\Delta Ct}$ method (Livak & Schmittgen, 2001).

Stomatal impressions, epidermal traits, petiole morphology and abaxial trichome density.

Stomatal density and epidermal cell size were measured using clear nail polish

impressions of adaxial and abaxial surfaces. Trichomes were removed from the abaxial side of

leaves with tape when necessary, and three sections per leaf were imaged and photographed

using an Olympus BX51 light microscope. Hand sections of leaf petioles within 1 cm of leaf

attachment site were imaged as described above. Trichome density on the abaxial side of the leaf

was measured using an Olympus MVX10 dissecting scope. Measurements were made along the

mid vein, where the density was low enough to permit accurate counts. All measurements and

counts were made using Fiji software.

Leaf vein densities

Sections from fully expanded leaves were cleared using NaOH and bleach, and then

stained with Safranin following the protocol described in Scoffoni and Sack (2013). Vein lengths

of all visible vein orders were measured from microscope images using Fiji software, and vein

densities were calculated following Scoffoni and Sack (2013).

Leaf shape and size

Fully expanded leaves were attached to paper using double-sided tape and scanned. The

scanned images were measured using the Morphological Analysis of Size and Shape (MASS)

software (Chuanromanee *et al.*, 2019).

Leaf initiation rate, internode distance and branching

The leaf initiation rate was determined by counting the node of the uppermost visible leaf

at least twice a week. The number of branches at least 1 cm in length between nodes 1 and 25

was determined after 230 days of growth.

Fluttering

To measure leaf fluttering, plants were placed 100 cm away from a fan, with the leaf to

be measured facing perpendicular to the wind. Wind speeds at the sample leaf position were 1.5

m/s as measured with an anemometer. Leaf movement was filmed at 240 fps for 30 seconds

(representative example in supplemental file "Poplar Fluttering Video"). Leaf movement was

analyzed by following the movement of a mark placed at the tip of the leaf using Tracker video

analysis and modeling tool v5.0.7. The height and width of the petiole were measured using a

digital caliper.

Statistical Analysis

All statistical analyses were performed in JMP $^{®}$ Pro v. 14.0.0 (SAS Institute Inc., Cary,

NC). Traits in seed grown poplar across leaf positions were measured from a minimum of 42

individuals and fit with a linear or second-degree polynomial as appropriate with *p*-values and R^2

values reported. Traits that were examined at multiple leaf positions in 717 wildtype, MIM156

lines 84 and 22 and miR156 over-expression lines 40 and 78 were measured from a minimum of

6 individuals and compared by two-way *ANOVA* and a Student's *t* test ($\alpha = 0.05$) where leaf

position and genotype were the main effects. Whole-plant measures of branching and internode

length in these plants was compared by one-way *ANOVA* where genotype was the main effect

and a Student's *t* test ($\alpha = 0.05$), and leaf emergence was compared by two-way *ANOVA* with

days after transplant and genotype as the main effects.

Results

miR156/7 expression in seed grown plants and 717-1B4 lines

The miR156/7 expression pattern in the leaf primordia of seed-grown plants matched that previously reported for other species, such as *A. thaliana* and *Vachellia* sp. (He *et al.*, 2018; Leichty & Poethig, 2019). miR156/7 abundance dropped 7-fold between nodes 1 and 10, followed by a slower decline in successive nodes (Fig. **1A**). We first determined that the control small RNA, miR159, was expressed at the same level in leaves 15 and 25 of seed-grown plants and the clone 717-1B4, and then measured the level of miR156/7 in wild-type and transgenic genotypes of 717-1B4. The abundance of miR156/7 in the primordia of leaf 15 and 25 in wildtype 717-1B4 was similar to the amount of miR156/7 in these same leaves in seed-grown poplar (Fig. **1A, B**), suggesting that miR156/7 are expressed at comparable levels at the same node in plants propagated by these methods. Leaf 15 and 25 of the miR156 over-expressor lines had, respectively 4.9 and 5.6 times the amount of miR156a as wild-type leaves, a level comparable to that of leaf 3 and 5 of seed-grown plants (Fig. **1A, B**). Although the amount of miR156/7 in leaves 15 and 25 of the MIM156 lines was similar to that in 717-1B4 and seed-grown poplar (Fig. **1A, B**), line 84 had approximately 1/3 the amount of miR156 at an earlier stage of development, before it was transferred to soil (data not shown). Target mimics do not necessarily reduce the level of their target miRNAs (Franco-Zorrilla *et al.*, 2007), so the significance of these MIM156 data is difficult to interpret. In any case, the similarity between the amount of miR156/7 in MIM156 and wild-type lines shows that the transformation protocol and vector components have no independent effect on miR156/7 abundance.

Changes in miR156 expression alter plant growth rate and form

Plants grown from seed produced leaves at a relatively slow rate for the first few weeks after planting, and transitioned to a faster, linear growth rate after producing 15-20 leaves (Fig. **3A**). Leaf initiation in wild-type and transgenic clones of 717-1B4 was delayed for 35-40 days after transplanting to soil, and then occurred at increasing rate over the next 2-3 weeks of measurement. During this latter period, lines transformed with Ubi10:AthMIR156a had higher rates of leaf initiation than either wild-type or MIM156 lines (Fig. **3B**). Ubi10:AthMIR156a lines also had shorter internodes than either the wild-type and MIM156 lines (Fig. **3C**) and produced

significantly more axillary branches than these genotypes (Fig. **3D**). The MIM156 lines were not

significantly different from 717-1B4 for any of these traits, although these lines did have slightly

longer internodes than 717-1B4.

Juvenile and adult leaves differ in leaf shape and size

The leaves of seed-grown plants became gradually rounder (decreased L:W ratio) (Fig.

4A) and larger (Fig. **4C**) from node 1 to node 25 and did not change significantly in shape or size

after this point (Fig. **1C**). 717-1B4 displayed a similar trend, although the leaves of this clone

reached their climax shape and size a few nodes earlier than seed-grown plants (Fig. **2** and **4B,**

D). MIM156 lines did not differ significantly in size from 717-1B4, but the leaves of line 84

were rounder than 717-1B4 beginning at node 15. In contrast, the leaves of lines over-expressing

miR156 were significantly more elongated and smaller than wild-type leaves starting from node

15 (Student's *t* test *p* <0.05). These results suggest that changes in leaf shape and size during

shoot development in poplar are partly regulated by a decrease in level of miR156.

To obtain a comprehensive picture of the effect of miR156 on leaf morphology in

Populus tremula x alba, we also measured the abaxial trichome density, stomatal density,

epidermal cell area, specific leaf area, vein density, and petiole morphology of successive leaves.

We examined every 5th leaf of seed-grown plants from germination to node 45-50, and every 5th

leaf of 717-1B4 and transgenic lines from node 10 to 25.

The leaves of *Populus alba* appear white because of a dense layer of trichomes on the

abaxial surface of the leaf blade, and this trait is also characteristic of *P. tremula x alba.* All the

leaves of *P. tremula x P. alba* produced abaxial trichomes, but trichome density increased

dramatically during shoot growth (Fig. **4G, H**). In seed-grown plants, abaxial trichome density

increased more than 10-fold between leaf 5 and leaf 40 and increased by about 6-fold between

leaf 10 and leaf 25 in 717-1B4. In contrast, abaxial trichome density remained at the same

relatively low level from node 10 to node 25 in the miR156 over-expressors, and it increased

significantly more than 717-1B4 in both MIM156 lines (Student's *t* test; *p* <0.01 and *p* <0.0001

for line 22 and 84 respectively). This result suggests that abaxial trichome production is a phase-

specific trait in *P. tremula x alba* and is hypersensitive to the level of miR156.

One factor that could contribute to this increase in trichome density is a decrease in

epidermal cell size, which would increase the number of epidermal cells per unit area. We

measured this trait by taking impressions of the adaxial epidermis, because the high density of

trichomes made it impractical to measure epidermal cell area on the abaxial surface. In seed-

grown plants, cell area decreased approximately 4-fold over the first 30 leaves and remained

relatively constant after this point (Fig. **4M**). In 717-1B4, cell area decreased about 2-fold from

leaf 10 to leaf 20, and decreased much less dramatically, if at all, between leaf 20 and leaf 25.

The MIM156 lines were not significantly different from 717-1B4. In contrast, cell area in the

miR156 over-expressors did not change significantly between leaf 10 and leaf 25. These results

suggest that epidermal cell area is a phase-specific trait, and also suggest that the increase in

abaxial trichome density is partly attributable to an increase in the cell density in the epidermis.

This is not the only explanation for the increase in abaxial trichome density, however, because

the magnitude of the change in trichome density and cell area is different, and because the

MIM156 had a significant effect on trichome density, but not cell area.

Stomatal density on both the adaxial and abaxial side of the leaf changed across leaf

positions, but in opposite directions (Fig. **4I, K**). In seed-grown plants, adaxial stomatal density

decreased linearly from leaf 5 to leaf 40, whereas abaxial stomatal density increased linearly

over this interval; at every node, the density of stomata on the adaxial surface of the leaf was

significantly less than that on the abaxial surface. Both 717-1B4 and the MIM156 lines had a

very low density of adaxial stomata on leaf 10, and this number was reduced to nearly zero on

leaf 25. In contrast, the density of abaxial trichomes in these genotypes increased significantly

between leaf 10 and leaf 25. In the miR156 over-expressing lines, the density of stomata on both

the adaxial and abaxial surfaces of leaves 10 and 25 was essentially identical to that of the basal-

most leaves of seed-grown plants. Thus, miR156 regulates stomatal density on both sides of the

leaf. However, other factors also contribute to the developmental changes in stomatal density

because this parameter changes linearly during shoot development, rather than in the curvilinear

pattern characteristic of the change in miR156 levels (Fig. **4J, L**).

Specific leaf area (SLA), is the area of the leaf blade per unit mass, and is determined by

the thickness of the leaf blade and cell density. SLA decreased in a curvilinear fashion from leaf

5 to leaf 40 in seed-grown plants and decreased linearly from leaf 10 to leaf 25 in 717-1B4 and

the 4 transgenic lines (Figure **4E, F**). However, SLA decreased nearly linearly between leaf 10

and leaf 25 in seed-grown plants, so these data do not reveal if there is a significant difference in

the pattern in seed-grown and clonally propagated plants. On the other hand, we found that—at

nodes 15-25—lines over-expressing miR156 had a significantly higher SLA, and one MIM156

line (84) had a significantly lower SLA, than 717-1B4 at node 10 (Student's t test, p <0.05 and p

<0.0001 for miR156 over-expressors at nodes 15 and 20, and node 25 respectively and p =0.019

for line 84 node 10). These results suggest that SLA is a phase-specific trait and is regulated by

miR156.

Leaf vein density increases as plants transition from juvenile to adult

Leaf vein density increased 1.8-fold with increasing leaf position in 717-1B4 and in the

MIM156 lines, although MIM156 line 84 had a significantly higher vein density than 717-1B4 in

leaf 10 (Student's t test, p <0.01). Similar results were obtained for the density of vein endings.

In the miR156 over-expressors, vein density and the density of vein endings did not differ much

between leaf 10 and leaf 25, and were identical to, or slightly lower than, the values observed for

leaf 10 in 717-1B4 (Fig. **5**).

Petiole morphology changes during development leading to altered leaf fluttering behavior

Many species of poplar, including *Populus tremula x alba,* exhibit leaf fluttering, a trait

that has been attributed to the vertical elongation of the petiole (Niklas, 1991). In a previous

study of *Populus x canadensis*, we found that the petioles of *in vitro* propagated plants became

increasing elongated as the shoot developed, and that over-expression of miR156 suppressed this

process (Wang et al. 2011). To determine if this phenomenon occurs naturally, we examined the

L:W ratio of a cross-section of the petiole in seed-grown *Populus tremula x alba* (Fig. **6A**). We

found that the petiole of leaves at node 10 was slightly broader than tall (L:W < 1), whereas the

petioles of leaves at higher nodes became vertically elongated (L:W >1), reaching their climax

shape around node 40. In wild-type 717-1B4, and in the MIM156 and miR156 over-expressors,

the petiole also became progressively elongated (Fig. **6B**). However, the miR156 over-expressors

had significantly less elongated petioles than the 717-1B4 and the MIM156 lines, and reached a

maximum L:W ratio at an earlier node. We also examined the effect of these transgenes on

petiole anatomy. In 717-1B4, the petioles of leaves at nodes 10 and 15 had an adaxial

indentation, a large central vascular bundle, and two smaller vascular bundles located in a

horizontal plane above the central bundle. The petioles of leaves 20 and 25 lacked an adaxial

indentation and had three vertically oriented vascular bundles that decreased in size from the

abaxial to the adaxial side of the petiole. In the MIM156 lines, the vascular anatomy of leaf 15

was intermediate between that of leaf 10 and leaf 20 in 717-1B4, indicating that the transition to

the adult shape is accelerated in these lines, despite the lack of a significant change in the L:W

ratio of the petiole (Fig. **6C**). Consistent with their L:W ratio, lines over-expressing miR156

retained a juvenile petiole morphology up to at least node 25. Petiole anatomy is therefore an

obvious qualitative indicator of the phase identity of a leaf in *Populus tremula x alba*.

316   To determine if this change in petiole morphology has an effect on the behavior of the

leaf, we looked at two measures of leaf fluttering—the total distance moved by the leaf, and the

maximum horizontal span of the leaf tip—when leaves were exposed to a constant wind velocity

(Fig. **7A**). There was a statistically significant (p <0.0001) positive linear relationship between

petiole L:W ratio and these fluttering measures in 717-1B4, the MIM156 and miR156 over-

expressing lines. Leaves at basal node 10 in the 717-1B4 and in the MIM156 lines fluttered less

than leaves at more apical node 25, whereas juvenile leaves at node 25 of the miR156 over-

expressing lines fluttered less than leaves at node 25 of the 717-1B4 and in the MIM156 lines

(Fig.**7B, C**). Thus, the petiole morphology of juvenile and adult leaves of *Populus tremula x alba*

has a significant effect on their response to wind.

Discussion

328   Many plants do not undergo visually abrupt changes in form during shoot development,

making the onset of vegetative phase change difficult to observe. As a result, the existence of this

phenomenon and its significance for plant ecology and evolution are often under appreciated. In

this study, we characterized the growth rate, developmental morphology, and temporal pattern of

miR156 expression in *Populus tremula x alba,* an ecologically and economically important tree.

We found that many leaf traits change during shoot development in plants grown from seeds, as

well as in shoots arising *de novo* in culture, and that most of these changes are correlated with

and sensitive to changes in the level of miR156. These data confirm and extend previous studies

of the effect of miR156 on *Populus* development (Wang *et al.*, 2011; Rubinelli *et al.*, 2013)*,* and

provide a foundation for future research on the function of vegetative phase change in the life

history of this woody species.

339   In nature, poplars propagate primarily from adventitious shoots, whereas clones used for

research or commercial purposes are propagated from axillary buds or by inducing shoot

regeneration *in vitro* (Taylor, 2002). The extent to which these plants recapitulate the

development of the primary shoot is still unknown. We found by at least the 10th node, the

development of a regenerated shoot is remarkably similar to that of a seed-grown plant. This

result suggests that the phase-identity of poplar cells is either re-set during the de-differentiation

of the primary tissue sample, or during shoot regeneration. This observation also raises the long-

standing question of whether differentiated traits can be maintained in tissue culture.

347   Although vegetative phase change sometimes involves traits unique to a particular

species (Wang *et al.*, 2011; Leichty & Poethig, 2019), analyses of plants with altered levels of

miR156 have revealed a number of phase-specific traits that are common to phylogenetically

diverse species. In *Zea mays*, *Arabidopsis thaliana*, *Nicotiana tabacum,* and now *Populus*

tremula x alba, these traits include patterns of trichome production, leaf shape and size,

epidermal and stomatal cell traits, vein density, leaf initiation rate, and branching (Bongard-

Pierce *et al.*, 1996; Telfer *et al.*, 1997; Feng *et al.*, 2016; He *et al.*, 2018). At least some of these

traits are also regulated by miR156 in *Populus x canadensis* (Wang *et al.*, 2011), rice (Xie *et al.*,

2006), alfalfa (Aung *et al.*, 2015), switchgrass (Fu *et al.*, 2012), *Lotus japonicus* (Wang, 2014),

and tomato (Zhang *et al.*, 2011). This observation indicates that vegetative phase change is an

ancient and fundamental aspect of shoot development, and also suggests that the differential

expression of various traits during shoot development contributes to plant fitness and is therefore

under natural selection (Leichty & Poethig, 2019).

360   Determining how phase-specific traits contribute to plant fitness is challenging, however.

A number of studies have shown that plants over-expressing miR156 are more or less sensitive

to various environmental stresses (Stief *et al.*, 2014; Cui *et al.*, 2014; Arshad *et al.*, 2017; Kang

et al., 2020), but there is still little evidence that juvenile and adult phases of wild type plants

display these same differences, or how they might contribute to plant fitness. There are relatively

few examples of phase-specific vegetative traits that are either known or expected to be

selectively advantageous (Williams *et al.*, 1998; Holeski *et al.*, 2009; Leichty & Poethig, 2019).

Our observation that adult leaves flutter more than juvenile leaves is another such example. In

poplar, leaf fluttering is beneficial under high light and high heat environments, but is

disadvantageous under low light conditions because it reduces photosynthesis (Roden & Pearcy,

1993a,b). This suggests that reduced leaf fluttering is advantageous early in development, when

plants are likely to be shaded by other plants, whereas increased fluttering is advantageous once

plants are no longer in the understory.

During shoot development, different morphological and physiological traits change in a

coordinated fashion, but not necessarily at the same node or at the same rate (Hackett & Murray,

1997; Leichty & Poethig, 2019). Furthermore, under certain conditions, traits can become

dissociated, resulting in leaves or shoots that have unusual combinations of juvenile and adult

traits (Robbins, 1960; Doorenbos, 1965). In our experiments, this was apparent from the

phenotype of transgenic lines expressing a miR156/miR157 target site mimic (MIM156). The

stronger of these transgenes had a significant effect on leaf L:W ratio, SLA, epidermal cell area,

petiole morphology and leaf vein density, but did not affect leaf initiation rate, stomatal density

and petiole L:W ratio. These observations are consistent with studies in in *Arabidopsis*, which

show that different miR156-regulated SPL genes have different transcription patterns and

sensitivity to miR156 abundance (Xu *et al.*, 2016; He *et al.*, 2018). Along with the evidence that

SPL genes also have different loss-of-function phenotypes (Xu *et al.*, 2016), these results suggest

that variability in the expression of different phase-specific traits may arise from the differential

response of functionally distinct SPL genes to changes in the level of miR156.

miR156 and its SPL targets control many economically important traits, including

flowering time, shoot architecture, biomass and lignin production, pest and disease resistance

and abiotic stress responses (Wang & Wang, 2015). In the biofuel crop *P. virgatum*, for example,

over-expressing miR156 leads to increases in biomass, digestibility, and lignin production, a

delay in flowering time, and altered susceptibility to the pathogen *Puccinia emaculata* (Chuck *et*

al., 2011; Fu *et al.*, 2012; Baxter *et al.*, 2018*)*. Similar effects of miR156 over-expression have

been observed in the forage legumes *Trifolium pretense* (Zheng *et al.*, 2016) and *Medicago*

sativa (Aung *et al.*, 2015). Poplars are fast-growing trees useful for a wide variety of industrial

purposes, as well for the re-forestation of degraded land (Taylor, 2002). The ability to engineer

their chemistry and vegetative development through the manipulation of miR156 offers new

opportunities for the improvement of this important group of plants.

Acknowledgements

We thank Samara Gray and Joshua Darfler for their assistance in caring for the plants used in

this study. This research was funded by the NSF Graduate Research Fellowship (DGE-1321851),

Author Contributions

406 E.H.L., A.R.L., and R.S.P. planned and designed the research. E.H.L. and A.R.L. performed the

407 experiments. C.M., and S.H.S. helped create the transgenics used in this study. E.H.L performed

408 statistical analyses and wrote the first draft of the manuscript. E.H.L., A.R.L., C.M., S.H.S., and

409 R.S.P revised and provided comments on the manuscript.

References

Arshad M, Feyissa BA, Amyot L, Aung B, Hannoufa A. **2017**. MicroRNA156 improves

drought stress tolerance in alfalfa (Medicago sativa) by silencing SPL13. *Plant Science* **258**:

122–136.

Ashby E. **1948**. Studies in the morphogenesis of leaves. II. The area, cell size and cell number of

leaves of Ipomoea in relation to their position on the shoot. *New Phytologist* **47**: 177–195.

Aung B, Gruber MY, Amyot L, Omari K, Bertrand A, Hannoufa A. **2015**. Ectopic

expression of LjmiR156 delays flowering, enhances shoot branching, and improves forage

quality in alfalfa. *Plant Biotechnology Reports* **9**: 379–393.

Baxter HL, Mazarei M, Dumitrache A, Natzke JM, Rodriguez M, Gou J, Fu C, Sykes RW,

Turner GB, Davis MF, *et al.* **2018**. Transgenic miR156 switchgrass in the field: growth,

recalcitrance and rust susceptibility. *Plant Biotechnology Journal* **16**: 39–49.

Bongard-Pierce DK, Evans MMS, Poethig RS. **1996**. Heteroblastic features of leaf anatomy in

maize and their genetic regulation. *International Journal of Plant Sciences* **157**: 331.

Brink RA. **1962**. Phase change in higher plants and somatic cell heredity. *The Quarterly Review*

of Biology **37**: 1–22.

Chuanromanee TS, Cohen JI, Ryan GL. **2019**. Morphological Analysis of Size and Shape

(MASS): An integrative software program for morphometric analyses of leaves. *Applications in*

Plant Sciences **7**: e11288.

Chuck GS, Tobias C, Sun L, Kraemer F, Li C, Dibble D, Arora R, Bragg JN, Vogel JP,

Singh S, *et al.* **2011**. Overexpression of the maize Corngrass1 microRNA prevents flowering,

improves digestibility, and increases starch content of switchgrass. *Proceedings of the National*

Academy of Sciences **109**: 995–995.

Cui L-G, Shan J-X, Shi M, Gao J-P, Lin H-X. **2014**. The miR156-SPL9-DFR pathway

coordinates the relationship between development and abiotic stress tolerance in plants. *The*

Plant Journal **80**: 1108–1117.

Doorenbos J. **1965**. Juvenile and adult phases in woody plants. *Encyl. Plant Physiol.* **15**: 1222–

35.

Engler C, Youles M, Gruetzner R, Ehnert T-M, Werner S, Jones JDG, Patron NJ,

Marillonnet S. **2014**. A Golden Gate modular cloning toolbox for plants. *ACS Synthetic Biology*

3: 839–843.

Feng S, Xu Y, Guo C, Zheng J, Zhou B, Zhang Y, Ding Y, Zhang L, Zhu Z, Wang H, *et al.*

2016. Modulation of miR156 to identify traits associated with vegetative phase change in

tobacco (Nicotiana tabacum). *Journal of Experimental Botany* **67**: 1493–1504.

Filichkin SA, Meilan R, Busov VB, Ma C, Brunner AM, Strauss SH. **2006**. Alcohol-

inducible gene expression in transgenic Populus. *Plant Cell Reports* **25**: 660–667.

Franco-Zorrilla JM, Valli A, Todesco M, Mateos I, Puga MI, Rubio-Somoza I, Leyva A,

Weigel D, García JA, Paz-Ares J, *et al.* **2007**. Target mimicry provides a new mechanism for

regulation of microRNA activity. *Nature Genetics* **39**: 1033–1037.

Fu C, Sunkar R, Zhou C, Shen H, Zhang J-YY, Matts J, Wolf J, Mann DGJ, Stewart CN,

Tang Y, *et al.* **2012**. Overexpression of miR156 in switchgrass (Panicum virgatum L.) results in

various morphological alterations and leads to improved biomass production. *Plant*

Biotechnology Journal **10**: 443–452.

Goebel K. **1900**. *Organography of Plants Part I. General Organography. (English Translation*

by I. B. Balfour). Oxford: Clarendon Press.

Hackett WP, Murray JR. **1997**. Approaches to understanding maturation or phase change. In:

Geneve RL, Preece JE, Merkle SA, eds. Biotechnology of Ornamental Plants. New York: CAB

International, 73–86.

He J, Xu M, Willmann MR, McCormick K, Hu T, Yang L, Starker CG, Voytas DF, Meyers

BC, Poethig RS. **2018**. Threshold-dependent repression of SPL gene expression by

miR156/miR157 controls vegetative phase change in Arabidopsis thaliana. *PLOS Genetics* **14**:

e1007337.

Holeski LM, Kearsley MJC, Whitham TG. **2009**. Separating ontogenetic and environmental

determination of resistance to herbivory in cottonwood. *Ecology* **90**: 2969–2973.

James SA, Bell DT. **2001**. Leaf morphological and anatomical characteristics of heteroblastic

Eucalyptus globulus ssp. Globulus (Myrtaceae). *Australian Journal of Botany* **49**: 259–269.

Kang T, Yu CY, Liu Y, Song WM, Bao Y, Guo XT, Li B, Zhang HX. **2020**. Subtly

manipulated expression of ZmmiR156 in tobacco improves drought and salt tolerance without

changing the architecture of transgenic plants. *Frontiers in Plant Science* **10**: 1664.

Leichty AR, Poethig RS. **2019**. Development and evolution of age-dependent defenses in ant-

acacias. *Proceedings of the National Academy of Sciences* **116**: 15596–15601.

Livak KJ, Schmittgen TD. **2001**. Analysis of relative gene expression data using real-time

quantitative PCR and the 2-$\Delta\Delta$CT method. *Methods* **25**: 402–408.

Meilan R, Ma C. **2006**. Poplar (Populus spp.). In: Wang K, ed. Methods in Molecular Biology:

Agrobacterium Protocols. Totowa, NJ: Humana Press Inc., 143–151.

Niklas KJ. **1991**. The elastic moduli and mechanics of Populus tremuloides (Salicaceae) petioles

in bending. *American Journal of Botany* **78**: 989–996.

Poethig RS. **2013**. Vegetative phase change and shoot maturation in plants. *Current Topics in*

Developmental Biology **105**: 125–152.

Rhoades MW, Reinhart BJ, Lim LP, Burge CB, Bartel B, Bartel DP. **2002**. Prediction of

plant microRNA targets. *Cell* **110**: 513–520.

Robbins WJ. **1960**. Further observations on juvenile and adult Hedera. *American Journal of*

Botany **47**: 485–491.

Roden JS, Pearcy RW. **1993a**. Effect of leaf flutter on the light environment of poplars.

Oecologia **93**: 201–207.

Roden JS, Pearcy RW. **1993b**. The effect of flutter on the temperature of poplar leaves and its

implications for carbon gain. *Plant, Cell and Environment* **16**: 571–577.

Rubinelli PM, Chuck G, Li X, Meilan R. **2013**. Constitutive expression of the Corngrass1

microRNA in poplar affects plant architecture and stem lignin content and composition. *Biomass*

and Bioenergy **54**: 312–321.

Schwab R, Palatnik JF, Riester M, Schommer C, Schmid M, Weigel D. **2005**. Specific

effects of microRNAs on the plant transcriptome. *Developmental Cell* **8**: 517–527.

Scoffoni C, Sack L. **2013**. Quantifying leaf vein traits. *Prometheus Wiki*.

Stief A, Altmann S, Hoffmann K, Pant BD, Scheible W-R, Bäurle I. **2014**. Arabidopsis

miR156 regulates tolerance to recurring environmental stress through SPL transcription factors. *The Plant Cell* **26**: 1792–1807.

Taylor G. **2002**. Populus: Arabidopsis for forestry. Do we need a model tree? *Annals of Botany* **90**: 681–689.

Telfer A, Bollman KM, Poethig RS. **1997**. Phase change and the regulation of trichome distribution in Arabidopsis thaliana. *Development* **124**: 645–654.

Varkonyi-Gasic E, Wu R, Wood M, Walton EF, Hellens RP. **2007**. Protocol: A highly sensitive RT-PCR method for detection and quantification of microRNAs. *Plant Methods* **3**: 12.

Walters GA, Bartholomew DP. **1984**. Acacia Koa leaves and phyllodes: Gas exchange, morphological, anatomical, and biochemical characteristics. *Botanical Gazette* **145**: 351–357.

Wang J-W. **2014**. Regulation of flowering time by the miR156-mediated age pathway. *Journal of Experimental Botany* **65**: 4723–4730.

Wang JW, Park MY, Wang LJ, Koo Y, Chen XY, Weigel D, Poethig RS. **2011**. MiRNA control of vegetative phase change in trees. *PLoS Genetics* **7**: e1002012.

Wang H, Wang H. **2015**. The miR156/SPL module, a regulatory hub and versatile toolbox, gears up crops for enhanced agronomic traits. *Molecular Plant* **8**: 677–688.

Williams WP, Davis FM, Buckley PM, Hedin PA, Baker GT, Luthe DS. **1998**. Factors associated with resistance to fall Armyworm (Lepidoptera: Noctuidae) and southwestern Corn Borer (Lepidoptera: Crambidae) in corn at different vegetative stages. *Journal of Economic Entomology* **91**: 1471–1480.

Xie K, Wu C, Xiong L. **2006**. Genomic organization, differential expression, and interaction of SQUAMOSA Promoter-Binding-Like Transcription Factors and microRNA156 in Rice. *Plant Physiology* **142**: 280–293.

Xu M, Hu T, Zhao J, Park M-Y, Earley KW, Wu G, Yang L, Poethig RS. **2016**. Developmental functions of miR156-regulated SQUAMOSA PROMOTER BINDING PROTEIN-LIKE (SPL) genes in arabidopsis thaliana. *PLOS Genetics* **12**: e1006263.

Zhang X, Zou Z, Zhang J, Zhang Y, Han Q, Hu T, Xu X, Liu H, Li H, Ye Z. **2011**. Over-expression of sly-miR156a in tomato results in multiple vegetative and reproductive trait alterations and partial phenocopy of the *sft* mutant. *FEBS Letters* **585**: 435–439.

Zheng Q-L, Liu J, Goff BM, Dinkins RD, Zhu H. **2016**. Genetic manipulation of miR156 for improvement of biomass production and forage quality in Red Clover. *Crop Science* **56**.

526　**Zotz G, Wilhelm K, Becker A**. **2011**. Heteroblasty-a review. *Botanical Review* **77**: 109–151.

Figure 1. miR156/7 relative abundance in *Populus tremula x alba* leaf primordia grown from seed (A) or in transgenic and wildtype plants of the 717-1B4 background grown from tissue culture (B). miR156/7 abundance relative to the average of 4 replicates from seed grown leaves 1 & 2. Data presented as individual biological replicates depicted by grey open circles and a mean best fit line in black (A) or as means ± s.e.m (n = 3) with 717 wildtype data depicted by purple circles, miR156a over-expressor lines depicted by blue squares and MIM156 lines depicted by red triangles. Representative leaf shape for leaves 1-17, 20 and 30 from plants grown from seed (C).

Figure 2. Representative plants from the wild-type 717-1B4, miR156a over-expressor line 40 and MIM156 line 84 two months following transplant from tissue culture. Representative leaf shape for leaves 10, 15, 20 and 25 in the 717-1B4 and transgenic poplar plants from tissue culture.

Figure 3. Number of leaves initiated by *P. tremula x alba* grown from seed (A) and the 717-1B4 wildtype and four transgenic lines from tissue culture (n = 3-6) (B). Internode distance between nodes 10-20 (n = 6) (C), and number of branches of > 1cm in length between nodes 1 and 25 after 230 days of growth (n = 6) (D) in the 717-1B4 and four transgenic lines. Symbols and colors are the same as figure 1 with seed grown data presented as individual biological replicates and 717-1B4 lines as means ± s.e.m. Data in panel A show a second-degree polynomial relationship with $p < 0.0001$ and $R^2 = 0.92$ and a two-way *ANOVA* for panel B found $p < 0.0001$ for genotype and days after transplant but no significance for an interaction between effects. One-way *ANOVA* for panels C and D found $p < 0.0001$ with the results of a *Student's T* between WT and mutants represented by * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

Figure 4. Leaf length:width (L:W) ratio, leaf area, specific leaf area (SLA), abaxial trichome density, adaxial stomatal density, abaxial stomatal density, and adaxial epidermal cell area of seed grown poplar (A, C, E, G, I, K, M) and the 717-1B4 and transgenic lines (B, D, F, H, J, L, N). Symbols and colors are the same as figure 1 with seed grown data presented as individual biological replicates and 717-1B4 lines as means ± s.e.m (n = 5-11). Data in panels A, C, E, G and M show a second-degree polynomial relationship with $p < 0.0001$ and R^2 values of 0.17, 0.6,

0.32, 0.56 and 0.85 respectively. Data in panels I and K show a linear relationship with p

<0.0001 and R^2 values of 0.58 and 0.46 respectively. Two-way *ANOVA*s with leaf position and

genotype as the effects for data in panels B, D, F, H, L and N found p <0.001 for both leaf

position and genotype. In panel J, p <0.001 for genotype but N.S. for leaf position. For the

interaction between genotype and leaf position, p <0.05 for data in panels B, F, H, and L and

N.S. for panels D, J, and N.

Figure 5. Total leaf vein density (A) and the number of vein endings (B) in the 717-1B4 and

transgenic plants. Symbols and colors are the same as figure 1 with data presented as means ±

s.e.m (n = 5-6). A two-way *ANOVA* with leaf position and genotype as the effects found p

<0.0001 for both effects and the interaction for data in panels A and B. Representative samples

of leaves cleared and stained with safranin-O to visualized vein density and architecture in leaves

10 and 25 of 717-1B4 and transgenic poplar lines (C).

Figure 6. Petiole length:width ratio from cross sections made within 1 cm of leaf attachment in

seed grown (A) and 717-1B4 and transgenic poplar (B). Symbols and colors are the same as

figure 1 with seed grown data presented as individual biological replicates and 717-1B4 lines as

means ± s.e.m (n = 7-12). Data from panel A show a second-degree polynomial relationship with

p <0.0001 and $R^2 = 0.62$. For panel B, a two-way *ANOVA* with leaf position and genotype as the

effects found p< 0.0001 for leaf position and genotype and $p=0.01$ for the interaction.

Representative samples of petiole cross sections stained with safranin-O from leaves 10, 15, 20

and 25 of the 717-1B4 and four transgenic lines (C).

Figure 7. Diagram showing the path length and horizontal span measurements of leaf fluttering

of a WT 717-1B4 adult leaf at node 25, juvenile leaf at node 10, and a miR156 over-expressor

line 40 juvenile leaf at node 25 (A). Quantification of leaf fluttering through total path length

over a 1 second period (B) and maximum horizontal span of the tip of the leaf (C) and their

relationship with petiole length:width ratio. WT 717-1B4 data is depicted by circles,

miR156aOX line 40 by squares and MIM156 line 84 by open triangles. Leaf developmental

stage is noted by color with adult leaves (node 25) in red, juvenile (node 10) in dark blue and

juvenilized (node 25) from miR156 overexpression in light blue. Linear fit for panel A) path

length = -0.34288 + 0.544(petiole L:W ratio) and B) horizontal span = -0.0852 + 0.0939(petiole

L:W ratio).

WT 717-1B4

10   15   20   25

miR156a ox
Line 40

miR156a ox
Line 78

MIM156
Line 22

MIM156
Line 84

3 cm

A

path length

horizontal span

Adult: WT node 25

Juvenile: WT node 10

Juvenilized: miR156ox node 25

B

R^2= 0.431
P < 0.001

Total Path Length (m)

0.8

0.6

0.4

0.2

1.0 1.2 1.4 1.6 1.8 2.0

Petiole L:W Ratio

C

R^2= 0.413
P < 0.001

Horizontal Span (m)

0.12

0.10

0.08

0.06

0.04

0.02

0.00

1.0 1.2 1.4 1.6 1.8 2.0

Petiole L:W Ratio

● Adult: WT node 25   ● Juvenile: WT node 10   ■ Juvenilized: miR156OX node 25
△ Adult: MIM156 node 25   △ Juvenilie: MIM156 node 10