microRNA-19b regulates proliferation & patterning in the avian forebrain

Archita Mishra[±1], Suvimal Kumar Sindhu[±2*], Niveda Udaykumar[3], Jonaki Sen[1*]

[1]Indian Institute of Technology Kanpur, Kanpur-208016, UP, India
[2]Yale School of Medicine, New Haven, CT, 06511, USA
[3]St. Jude Children's Hospital, Memphis, TN, 38105, USA

[±] These authors contributed equally.

*For Correspondence: suvimal.sindhu@yale.edu (SKS)**;** jonaki.sen@iitk.ac.in (JS)

Competing interest: The authors declare that no competing interest exist.

Abstract

Unlike the six-layered organization of neurons in the mammalian neocortex, the avian pallium features spatially-segregated neuronal clusters with partially conserved gene expression. While efforts to uncover similarities between the avian pallium and mammalian neocortex have persisted, the mechanisms of pallial development in birds have not been comprehensively explored. Here, we have established the role of a highly conserved microRNA, miR-19b, in controlling proliferation in the embryonic avian forebrain, by regulating the expression of E2f8, a cell-cycle inhibitor and NeuroD1, a neuronal differentiation factor. Further, investigation revealed that there exists a proliferation-dependent specification of pallial neurons in birds, similar to that in mammals. In summary, these findings suggest a profound role of miR-19b in shaping the avian pallium and indicate a closer resemblance of developmental mechanisms between the avian pallium and the mammalian neocortex.

Keywords: miR-19b, Pallium, Chick, Neocortex, Fezf2, Mef2c, Senescence

Introduction

Elucidating the mechanism(s) driving the evolution of the mammalian laminar neocortex and its relationship to the non-laminar pallium found in birds and reptiles has been a long-standing interest in neuroscience (Butler et al., 2011). The mammalian neocortex features a six-layered laminar pallium, with neurons being born from a common pool of neural stem cells (NSCs)/radial glial cells (RGCs) located in the ventricular zone (VZ). Intermediate basal progenitors (IPCs) produced from the RGCs, reside in the subventricular zone (SVZ), where they undergo a few rounds of cell division to produce more neural progenitor cells (NPCs). These NPCs then differentiate and migrate radially from the dorsal region and tangentially from the ventral region to form neocortical neurons (Haubensak et al., 2004; McConnell & Kaznowski, 1991; Miyata et al., 2004; Molyneaux et al., 2007; Noctor et al., 2004). In contrast, the differentiating immature

neurons produced from NPCs in birds, undergo radial migration without the generation of IPCs from RGCs (Cardenas et al., 2018; Nomura et al., 2016). The avian pallium is segregated into clusters of neurons known as nuclei (Ball & Balthazart, 2021; Karten, 1997). Interestingly, in birds laminar organization is observed in limited regions such as the hyperpallium and the auditory cortex (Stacho et al., 2020; Wang et al., 2010). Furthermore, neurons in the developing avian pallium exhibit neocortex-like subtypes, as revealed by marker and projection analyses (Abellan et al., 2014; Atoji & Wild, 2012; Dugas-Ford et al., 2012; Gupta et al., 2012; Puelles et al., 2000; Suzuki & Hirata, 2014; Suzuki et al., 2012). However, the mechanisms governing the production and patterning of these homologous neurons in the avian brain remain poorly understood.

In mammals, several signaling pathways (Ohtaka-Maruyama & Okado, 2015) and microRNAs (miRNAs), which cause post-transcriptional silencing of gene expression of multiple target mRNAs (Agarwal et al., 2015; Bartel, 2004), regulate the production and specification of neocortical neurons (Diaz et al., 2020; Shu et al., 2019; Todorov et al., 2024). Loss of miRNA activity through Dicer ablation reduces the size of the neocortex and negatively impacts NSC survival and differentiation (Andersson et al., 2010; De Pietri Tonelli et al., 2008; Kawase-Koga et al., 2010; Nowakowski et al., 2011). A highly conserved microRNA cluster, miR-17~92, has been shown to be crucial for survival of mouse embryos (Ventura et al., 2008) and plays a role in expanding NSCs/RGCs in the mammalian brain (Bian et al., 2013). Furthermore, a member of this cluster, miR-19a, regulates the number of NSCs in mammals by controlling the expression of Pten, a tumor suppressor that regulates NSC renewal (Bian et al., 2013; Zheng et al., 2008).

Another member of this cluster miR-19b, an ortholog of miR-19a, has been identified as a prime oncogenic miRNA (Fan et al., 2014; Mendell, 2008; Olive et al., 2009). Our previous study demonstrated that miR-19b is expressed in the VZ of the developing chicken forebrain (Sindhu et al., 2019). These findings motivated us to further investigate the function of miR-19b in regulating neurogenesis. We first conducted a comprehensive expression analysis of miR-19b across different developmental stages in chicken. Interestingly, this analysis revealed an expression pattern that is spatially and temporally synchronous with the neurogenic wave in the developing chicken pallium. Further analysis showed that miR-19b regulates the number of NSCs and controls the expression of E2f8, a crucial regulator of the cell-cycle (Christensen et al., 2005; Maiti et al., 2005), as well as NeuroD1, a key factor in neuronal differentiation (Lee et al., 1995; Pataskar et al., 2016). In fact, we found that altering proliferation and neurogenic differentiation through the functional manipulation of miR-19b affected neuronal patterning in the developing avian forebrain. In summary, this study reveals some conserved mechanisms of neurogenesis and patterning between the avian pallium and the mammalian neocortex during development, reinforcing the notion of their shared evolutionary origin.

Results
Mutually exclusive expression of NeuroD1 and miR-19b
We have previously reported that miR-19b is expressed in the VZ of the chick forebrain at Hamilton-Hamburger stage 24 (HH24) (Hamburger & Hamilton, 1951; Sindhu et al., 2019), further analysis revealed that its expression persists in the VZ across various developmental stages e.g. at HH29 and, HH35 (Fig. 1 A-C). Interestingly, NeuroD1 - a transcription factor reported to be necessary and sufficient for neural specification and

differentiation, (Lee et al., 1995; Pataskar et al., 2016) is expressed in the subventricular
zone (SVZ) in the developing chicken forebrain, in a mutually exclusive expression
domain to that of miR-19b, particularly at HH24 (Fig. 1A). Further, at HH29 these two
factors are expressed differentially in complementary regions, with miR-19b being
expressed at higher levels in the dorsal ventricular ridge (DVR) region (yellow-dashed
box) as compared to the wulst region (magenta-dashed box) (Fig. 1B). Interestingly, at
HH35, miR-19b expression is relatively low in the wulst region, where NeuroD1
expression is higher and distributed throughout (white arrows in magenta-dashed boxed
region) (Fig. 1C). In contrast, NeuroD1 is only present in the SVZ in the DVR region,
while miR-19b is expressed at higher levels throughout the DVR region (Fig. 1C). This
suggests that miR-19b and NeuroD1 are expressed in a spatio-temporally dynamic
manner in mutually exclusive domains in the wulst and DVR region.
99
Spatio-temporal dynamics of neurogenesis
Neurogenesis in the avian pallium follows a distinct spatio-temporally dynamic pattern,
where it begins in the presumptive wulst region followed by the DVR region as is evident
from differential labelling with EdU and immunohistochemistry for phospho-Histone 3
(PH3) (Abdel-Mannan et al., 2008; Cheung et al., 2007; Gupta et al., 2012; Suzuki et al.,
2012). To correlate this dynamic spatio-temporal pattern of neurogenesis with the
complementary expression patterns of miR-19b and NeuroD1, we compared the
expression pattern of NeuroD1 with PH3 immunostaining and found that indeed PH3
and NeuroD1 are present in complementary regions across developmental stages. At
HH24, PH3+ and NeuroD1+ cells are homogeneously distributed in the VZ and SVZ,
marking the presence of proliferating NSCs and post-mitotic immature neurons,
respectively (Fig. 1D). However, at HH29 proliferation ceases in the wulst region as
demonstrated by low PH3 immunoreactivity and there is a higher expression of NeuroD1
in the SVZ relative to the DVR (Fig. 1D). Further, at HH35 there are negligible number of
PH3+ cells in both the regions, whereas DVR shows enriched expression of NeuroD1 in
the SVZ. This indicates that while NSC proliferation ceases at both locations, the DVR is
still producing post-mitotic immature neurons (Fig. 1D). These findings suggest that the
VZ of the presumptive wulst region ceases proliferation earlier than the DVR and
NeuroD1 is primarily expressed in SVZ (D'Amico et al., 2013; Lee et al., 2000) of the
regions where proliferation have ceased and neuronal differentiation is initiated (FIG.
1E).
121
Taken together, it appears that the VZ, a proliferative zone, expresses miR-19b whereas
the SVZ, a post-mitotic zone, expresses NeuroD1, hence they are in mutually exclusive
domains. Hence, we hypothesized that miR-19b might serve as a key regulator of cell
proliferation by enhancing proliferation through its expression in the VZ and/or by
inhibiting neuronal differentiation through its regulation of NeuroD1, as observed in the
mouse pancreas.
128
miR-19b as a proliferation regulator
To explore the function(s) of miR-19b, we identified its potential target mRNAs using
tools like - miRDB (Chen & Wang, 2020) and TargetScan (Agarwal et al., 2015) and
selected 423 candidates from their intersection (Fig. 2A). Gene ontology analysis
indicated that miR-19b targets are mostly associated with cellular senescence (Fig. 2B).
Enriched expression of miR-19b in the VZ which contains the NSCs, prompted us to
explore the possible function of miR-19b in regulation of cell proliferation. Furthermore,

among the candidate regulators of cellular-senescence, we chose a cell-cycle inhibitor,
E2f8 (Christensen et al., 2005; Maiti et al., 2005), due to its strong expression in the
developing chick forebrain at HH30 (Fig. 2E) and previous reports of its regulation by
miR-19b in CD4+ T-cells (Lim et al., 2024). TargetScan analysis showed that chicken
E2f8 has an evolutionarily conserved target region (Fig. 2 D) (Agarwal et al., 2015). We
then performed a sensor assay (Sindhu et al. 2019) to determine if miR-19b regulates
the E2f8 transcript via its target regions by co-expressing miR-19b with transcripts of
E2f8 fused with an mCherry reporter in HEK293 T cells (Fig. 2C). This resulted in
decreased mCherry expression in the presence of miR-19b as compared to the control,
suggesting that miR-19b can regulate these transcripts through their target regions *in*
vitro (Fig. 2C). Further, *in ovo* gain-of-function (GoF) and loss-of-function (LoF)
experiments were performed with overexpression of miR-19b and Sponge-19b,
respectively (Sindhu et al., 2019). A Sponge is an artificial transcript that comprises of
multiple copies of the miRNA target sequence, which sequester the endogenously
expressed functional miRNAs, thus it competitively prevents miRNAs from interacting
with its cognate target mRNAs (Ebert & Sharp, 2010). Expression of Sponge-19b that,
sequesters miR-19b leading to a decrease in availability of miR-19b in the cell (Sindhu et
al., 2019), showed increase in E2f8 transcripts in the developing chick forebrain at HH25
(Fig. 2F). Such manipulations in the levels of miR-19b demonstrated that miR-19b is
both necessary as well as sufficient to regulate the expression of E2f8 (Fig. 2E, F). To
further establish the role of miR-19b in regulation of the cell-cycle, we employed flow-
cytometry to sort chicken fibroblast cells (DF1) and observed that miR-19b over-
expression increased the proportion of cells in the S-phase and M-phase, while
decreasing the proportion in G0/G1. On the other hand, depletion of functional miR-19b
through expression of Sponge-19b had the opposite effect (Fig. 2G). We also performed
EdU-labeling of the chick forebrain neurons under conditions of miR-19b GoF and LoF,
adding EdU 2 hours before harvesting at HH25. Quantification of GFP+;EdU+ cells
showed that EdU incorporation was promoted by miR-19b overexpression, while it was
reduced by expression of Sponge-19b (Fig. 2H). In addition, immunohistochemistry for
PH3 conducted at HH34 after functional manipulation of the miR-19b, yielded similar
results with over-expression of miR-19b increasing GFP+;PH3+ cells, and expression of
Sponge-19b leading to a reduction of the same (Fig. 2I). These findings strongly indicate
that miR-19b regulates cell proliferation in the developing chick pallium.
169
Perturbation in proliferation affects pallial patterning
Since birds show differential proliferation in two distinct regions of the avian pallium i.e.
the wulst and the DVR regions (Abdel-Mannan et al., 2008; Cheung et al., 2007; Gupta
et al., 2012; Suzuki et al., 2012) that express separate neocortical layer markers (Dugas-
Ford et al., 2012; Puelles et al., 2000; Suzuki & Hirata, 2014; Suzuki et al., 2012),
therefore, we wanted to study the relationship between proliferation and specification of
distinct regions in the avian pallium. Prior to manipulating proliferation to examine its
effect on regional specification, we confirmed the expression of patterning markers -
Fezf2 and Mef2c, expressed in the wulst and DVR regions respectively in the avian
pallium, which correspond to the deep-layer and upper-layer neurons in the mammalian
neocortex (Chen et al., 2008; Molyneaux et al., 2007; Suzuki et al., 2012). Upon
analysis, we found that at HH28, Fezf2 was expressed in the ventricular zone (VZ) of the
dorsomedial region and the subventricular zone (SVZ) of the dorsal pallium (Fig.
3A). Immunostaining for NeuroD1 - a postmitotic immature neuron marker, confirmed
Fezf2+ cells in the SVZ as postmitotic immature neurons, due to their co-expression of

NeuroD1 (Fig. 3A, Inset). Additionally, Mef2c expression was observed in cells atop the
Fezf2+ cells in the SVZ (Fig. 3A). Thus, Fezf2 and Mef2c expression at HH28 appeared
in a bi-laminar form, resembling the neocortical laminar order present in mammals
(Molyneaux et al., 2007), indicating the possibility of primitive pallial lamina formation in
the common ancestors of birds and mammals. However, at advanced stages such as
HH32, HH36 and post-hatching day 0, the expression domains of these markers
became spatially segregated, with Fezf2 in the hyperpallium and parahippocampal
region (APH) in the wulst and Mef2c in the mesopallium (Msp) in the DVR region (Fig.
3B) (Dugas-Ford et al., 2012; Suzuki et al., 2012). Therefore, as the development of the
avian pallium progresses, Fezf2+ and Mef2c+ neurons segregate spatially, departing
from their earlier bi-layered arrangement and residing in distinct regions such as
presumptive wulst and DVR regions.
197
To test the role of proliferation in patterning, we inhibited the cell-proliferation by
overexpressing CDKN1A gene that encodes P21 protein, a cell-cycle inhibitor (Deng et
al., 1995; Harper et al., 1993; Zhang et al., 1999), and found ectopic expression of Fezf2
and NeuroD1, indicating that inhibiting the cell-cycle alone is sufficient to generate these
neuronal subtypes in the developing chick pallium (Fig. 3C). Interestingly, chicken NPCs
that exit the cell-cycle earlier than other NPCs *in vitro* have been shown to preferentially
produce more Fezf2+ neurons at the expense of Mef2c+ neurons, much like in
mammals (Shen et al., 2006; Suzuki et al., 2012). Thus, *in vitro* findings support the *in*
vivo evidence that the early cessation of proliferation leads to the production of Fezf2+
neurons. Further analysis of the CDKN1A overexpressing regions revealed a decrease
in the Mef2c+ neuronal population (Fig. 3D). These results indicate that precocious
cessation of proliferation leads to preferential production of deep-layer like Fezf2+
neurons and suppresses the alternative upper-layer-like Mef2c+ fate (Chen et al., 2008;
McConnell & Kaznowski, 1991).
212
Our observation of cell-cycle inhibition resulting in ectopic Fezf2 expression (Fig. 3C),
along with the demonstration of miR-19b as a cell proliferation regulator, prompted us to
investigate if miR-19b might also regulate the expression of deep-layer neuronal
markers. We conducted a LoF experiment for miR-19b by expressing Sponge-19b in the
VZ and interestingly this resulted in ectopic Fezf2 expression in comparison to the
control (Fig. 3E). This suggests that miR-19b plays a crucial role in specification of deep-
layer-like neurons in the chick pallium. Such mechanism of proliferation-dependent
specification of pallial neurons resembles what is observed in the mammalian neocortex
(Shen et al., 2006) suggesting conservation in the mechanisms for cortical neuronal
specification between birds and mammals.
223
miR-19b regulates NeuroD1 which in turn regulates pallial patterning
Interestingly, miR-19b has been shown to also regulate the NeuroD1 transcript, albeit in
the mouse pancreas (Zhang et al., 2011). Based on this and the mutually exclusive
expression pattern of miR-19b and NeuroD1 (Fig. 1A-E, 4A), along with the fact that
chicken NeuroD1 mRNAs have evolutionarily conserved miR-19b binding sites (Fig. 4B),
we hypothesized that miR-19b regulates NeuroD1 in the developing chicken forebrain. A
sensor assay for NeuroD1 was performed, similar to the one described for E2f8 (Fig. 2C)
and we found that indeed miR-19b can regulate the NeuroD1-target region in a
sequence-dependent manner *in vitro* (Fig. 4 C). However, experiments carried out *in ovo*
suggested that while miR-19b is sufficient for regulating the expression of NeuroD1, it is

not necessary for the same (Fig. 4D-E). This is because miR-19b over-expression could decrease NeuroD1+ cells whereas Sponge-19b expression did not lead to an increase in NeuroD1+ cells relative to the control (Fig. 4D-E). This may be explained by the presence of other inhibitors of NeuroD1 *in vivo*.

Since NeuroD1 has also been implicated in causing cell-cycle exit and promoting neural specification and differentiation (Lee et al., 1995; Pataskar et al., 2016) and cell-cycle perturbation affects avian pallial patterning (Fig. 3C-D), therefore we wanted to find out if manipulation of NeuroD1 expression can also affect patterning in the avian pallium. Prior to this, we established the relationship of expression domains of lateral pallial marker i.e. Mef2c with NeuroD1 to better score the changes in regional expression of Mef2c upon perturbation of NeuroD1. The expression pattern comparison of NeuroD1 and Mef2c in the forebrain at HH29 showed that NeuroD1 is expressed in the SVZ region, while Mef2c is expressed in the mesopallium region (Fig. 4F). This suggests that NeuroD1 is transiently expressed in immature neurons, followed by Mef2c as they mature, similar to the upper layer of the mouse neocortex (Lee et al., 1995; Molyneaux et al., 2007; Pataskar et al., 2016). Subsequently, to test the role of NeuroD1 in patterning, we manipulated the function of NeuroD1 by expressing a dominant-negative version, NeuroD1-EnR, where DNA-binding domain of chicken NeuroD1 was fused to the Engrailed-Repressor domain (Yan & Wang, 2004) and found that this indeed inhibits the expression of Mef2c mRNA as well as protein (Fig. 4G, H). On the other hand, NeuroD1 overexpression induced the expression of Mef2c mRNA and Protein (Fig. 4I). These data indicate that NeuroD1 is both necessary and sufficient to regulate the expression of Mef2c, which is crucial for pallial patterning.

Since miR-19b is sufficient to regulate NeuroD1 expression (Fig. 4D), which in turn regulates Mef2c expression (Fig. 4G-J), we directly tested the effect of manipulating miR-19b on Mef2c expression. We observed that upon miR-19b GoF there was a down-regulation of Mef2c expression while miR-19b LoF, mediated by expression of Sponge-19b, led to a decrease in Mef2c expression (Fig. 4K). The ability of Sponge-19b to decrease Mef2C was surprising as we had shown earlier that it does not downregulate NeuroD1 (Fig 4E). The possible explanation for this could be as follows: Sponge-19b can inhibit proliferation (Fig. 2H-I) and proliferation inhibition leads to induction of Fezf2, while inhibiting Mef2c expression (Fig. 3C-D). Thus, it is likely that the decrease in Mef2c expression with Sponge-19b could result from inhibition of cell proliferation rather than through inhibition of NeuroD1 expression.

Discussion

The avian pallial organization and the mammalian neocortical arrangement are quite different, despite sharing homologous neurons. Therefore, in this study, we examined the chicken pallium to explore key mechanisms of neurogenesis and patterning that may be evolutionarily conserved and hence influence mammalian neocortical development. Here, we have uncovered the following striking similarities between the two species, in terms of mechanisms of production of neurons: a) miR-19a/b-mediated control of neurogenic proliferation, b) miR-19b-mediated repression of NeuroD1 expression and c) proliferation-dependent specification of neurons leading to regulation of pallial patterning. In addition, we have also found a transient pallial bi-layer comprising of Fezf2+ and Mef2c+ neurons in the early embryonic chick forebrain. Overall, this study

282 provides strong evidence that miR-19b could be a major player in shaping avian pallial
283 development using developmental mechanisms partially conserved in mammals.
284
285 **miR-19b regulates proliferation of neural progenitor cells**
286 This study has revealed that miR-19b is expressed in the VZ and has been shown to
287 regulate the NSCs number. Further, it also targets E2f8, an inhibitor of cell-proliferation
288 and NeuroD1, a neuronal differentiation factor. Thus, miR-19b is very likely to be a
289 crucial regulator of proliferation in the developing chicken forebrain.
290
The E2f family of transcription factors regulates the cell-cycle, with some acting as
activators and others as repressors (Maiti et al., 2005). Atypical member of E2f family,
E2f8, along with E2f7, supports cell-survival in developing mouse embryos (Li et al.,
2008) and inhibits proliferation of skin cancer cells (Thurlings et al., 2017). E2f8 has
been shown to act as inhibitor of transcription as well (Park et al., 2022). Additionally,
miR-19b has been also shown to repress expression of E2f8 in CD4+ T-cell (Lim et al.,
2024).Therefore, E2f8 is likely to play a similar role in the chick forebrain, in regulating
cell proliferation by promoting cell-cycle exit, and its control by miR-19b may significantly
impact neuron production in the avian embryonic pallium.
300
301 NeuroD1 has been shown to initiate the neuronal differentiation program (Lee et al.,
302 1995; Pataskar et al., 2016; Schwab et al., 2000) and promote cell-cycle exit, when
303 misexpressed in the mitotic retinal cells (Ochocinska & Hitchcock, 2009). Therefore,
304 suppressing NeuroD1 in the VZ may be essential for maintaining the proliferative state of
305 NSCs, and the role of miR-19b could be important in this context.
306
Based on our observation that miR-19b inhibits E2f8 expression as well as NeuroD1
expression, we speculate that miR-19b is required for maintaining the proliferative state
of NPCs by blocking their cell-cycle exit through suppression of E2f8 and preventing
premature neurogenesis through inhibition of NeuroD1 expression in the VZ. All the
above support our hypothesis that miR-19b regulates cell proliferation in the VZ in the
developing chick forebrain. Moreover, since the ortholog of miR-19b i.e. miR-19a has
also been shown to regulate neurogenesis (Bian et al., 2013), our study suggests that
miR19-mediated mechanism of regulation of neurogenesis is conserved between birds
and mammals.
316
317 **Progression of the cell-cycle and specification of neocortical neurons**
318 In the mouse there is an intrinsic program that regulates sequential formation of deep-
319 layer and upper-layer neurons in a temporally regulated manner (Shen et al., 2006). A
320 similar process has been reported for *in vitro* neurogenesis from chicken pallial neural
321 stem cell (Suzuki et al., 2012).
322
Overexpression of CyclinD1-Cdk4 in the mouse embryonic pallium increases Tbr2+
basal progenitors (IPCs) in the SVZ, initially inhibiting neuronal differentiation (Lange et
al., 2009). Over time, CyclinD1-Cdk4 effects weaken, leading to more IPCs and mature
upper-layer neurons in the cortical plate. Ectopic expression of CyclinD1-Cdk4 in chick
embryos at HH24 also produces postmitotic cells that are Tbr2+ immature neurons,
although their subtypes were not further identified to explore the link between cell-
progression and neuronal specification (Nomura et al., 2016).
330

In our study, we found that in the early chick embryo suppressing the progression of cell-
cycle led to the production of postmitotic neurons, as shown by NeuroD1 expression
along with Fezf2 in these cells. This is similar to what has been observed when chick
NSCs at an early stage produce deep-layer-like Fezf2+ neurons under *in vitro*
conditions (Suzuki et al., 2012). Interestingly, in mice, the gene CDKN1B, which is
related to CDKN1A, produces the P27 protein, a cell-cycle inhibitor. P27 is important for
the formation of upper-layer neurons because it slows down the cell-cycle and helps
activate Ngn2, which controls the expression of NeuroD1 (Nguyen et al., 2006; Roybon
et al., 2009). Thus, there is a striking similarity between the chick and the mouse with
respect to the link between the time of exit from cell-cycle and the specification of
neuronal subtype.
342
miR19b as a regulator of NeuroD1 expression
NeuroD1 has been previously shown to be both necessary and sufficient for neural
specification and differentiation (Lee et al., 1995; Pataskar et al., 2016). Recently, it has
been demonstrated that transient expression of NeuroD1 is sufficient to activate the
neurogenic program in progenitor cells (Pataskar et al., 2016). Therefore, it is likely that
one or more factors in the neural progenitor cells in the VZ maybe necessary to inhibit
any precocious expression of NeuroD1.
350
Our study suggests that miR-19b is sufficient but not necessary to regulate NeuroD1
expression. This could possibly be due to the presence of other inhibitory factors such
as other microRNAs and transcriptional repressors, in the VZ. In mammals, additional
microRNAs, such as miR30a-5p (Kim et al., 2013), miR-190 (Zheng et al., 2010) and
miR-124 (Liu et al., 2011), have been reported to regulate the expression of NeuroD1 in
various tissue contexts. Hence, it is possible that one or more of these microRNAs are
present in the VZ of the developing chick forebrain and contribute to the inhibition of
NeuroD1 expression, both under natural conditions and in experimental settings
involving the suppression of miR-19b.
360
NeuroD1 and production of neocortical neurons
NeuroD1 was discovered as a gene that is both necessary and sufficient for the
differentiation of neurons in several regions of the nervous system, including the
forebrain (Boutin et al., 2010; Gao et al., 2009; Lee et al., 1995; Schwab et al., 2000).
Interestingly, NeuroD1 GoF experiments in the mouse brain showed increased cell-cycle
exit and more postmitotic neurons in the intermediate zones compared to controls,
although layer markers were not analyzed separately (Pataskar et al., 2016). However,
NeuroD1 overexpression in the muller cells in retina showed its re-specification into
different neuronal subtypes depending on its expression level (Xu et al., 2023). In our
study, we found that NeuroD1 is both necessary and sufficient for the expression of
Mef2c+ upper-layer-like neocortical neurons in the chick pallium. Hence, NeuroD1
appears to play a key role in regulating production and specification of neurons during
pallial development.
374
375 In conclusion, our study establishes miR-19b as a significant contributor to the regulation
376 of NSC proliferation, differentiation, and pallial patterning in birds in an evolutionarily
377 conserved manner. Thus, our findings provide novel insight into some aspects of shared
378 evolutionary and developmental mechanisms that regulate pallial development.
379

Materials and methods

Key resource table

Reagent type or resource	Source	Identifier
Antibodies		
Rabbit Polyclonal anti-Mef2c	Millipore-Sigma	Cat# HPA005533; RRID:AB_1079352
Mouse Monoclonal anti-NeuroD1	Abcam	Cat# ab60704; RRID:AB_943491
Rabbit Polyclonal anti-PH3	Millipore-Sigma	Cat# H0412; RRID:AB_477043
Alexa Fluor 594 anti-Mouse IgG	Jackson ImmunoResearch Labs	Cat# 115-585-003; RRID:AB_2338871
Alexa Fluor 594 anti-Rabbit IgG	Jackson ImmunoResearch Labs	Cat# 111-585-045; RRID:AB_2338062
Chemicals, Peptides, and Recombinant Proteins		
Poly-Freeze	Sigma	Cat# P0091
Turbofect	ThermoFisher	Cat# R0533
Fast Green Dye	Sigma-Aldrich	Cat# 44715
To Pro Far Red	Invitrogen	Cat# T3605
Critical Commercial Assays		
EdU Kit	Thermo Fisher Scientific	Cat# C10338
Deposited Data Resources		
miRDB	http://mirdb.org/miRDB/	RRID:SCR_010848
TargetScan	http://targetscan.org/	RRID:SCR_010845
Experimental Models: Cell Lines		
HEK293T	ATCC	Cat# CRL-3216; RRID:CVCL_0063
DF1	ATCC	Cat# CRL-3586
Experimental Models: Organisms/Strains		
Chick Embryo (White leg horn)	Ganesh Enterprises, Nankari, Kanpur, India	N/A
Oligonucleotides		
NeuroD1-Target-Region	Macrogen	5'TACTACTTGTTACCTTTCCCGCGCCGAATTATTTTGTTGTGGTTTTGTACAG AATTTTTAATGACTTTTTATAACGTG3'
E2f8-Target-Region	Macrogen	5'AAGGTACAGTCTGTCAATCAGTGCATATTTTGCACAAAGGAAGTACTGTGAGCATGTACA3'.
Sponge-19b	Macrogen	5'TGCTGCCGtcagttttgcaccttttgcacaCCGGtcagttttgcaccttttgcacaCCGGtcagtt ttgcaccttttgcacaCCGG3'
Recombinant DNA		
pRmiR-Empty	Smith et al. 2009	N/A

pRmiR-19b	Sindhu et al. 2019	N/A
pSponge-19b	Sindhu et al. 2019	N/A
Sensor-NeuroD1-Target-Region	This Paper	N/A
Sensor-E2f8-Target-Region	This Paper	N/A
pCAG-GFP	Cepko Lab, Harvard University	N/A
pCMV-SPORT6-CDKN1A	MGC Clones	N/A
Software and Algorithms		
Venn Diagram Tool	https://bioinformatics.psb.ugent.be/webtools/Venn/	N/A
Enrichr Tool	https://maayanlab.cloud/Enrichr/#	RRID:SCR_001575
BD FACSDiva software	Becton Dickinson, USA	RRID:SCR_001456
Zen 2011	Zeiss	RRID:SCR_0013672
ImageJ	https://imagej.net/ij/	RRID:SCR_003070
Prism 10	Graph Pad	RRID:SCR_002798

Animals

383
384 Fertilized chicken eggs from the White Leghorn variety were purchased from Ganesh
385 Enterprises, Nankari, Indian Institute of Technology Kanpur, Uttar Pradesh, India. These
386 eggs were incubated in a humidified incubator at 38 °C until desired Hamilton
387 Hamburger (HH) stages (Hamburger and Hamilton 1992).

Cell-culture

389
390 Human embryonic kidney (HEK-293T) (ATCC; CRL-3216) and chicken embryonic
391 fibroblast, DF-1 (ATCC; CRL-3586) cells were cultured with 10% Fetal Bovine Serum in
392 Dulbecco's Modified Eagle's Medium (DMEM) media.

Tissue preparation

394
The chick embryo at the desired HH stages were taken and brains were dissected out.
The fixation was performed at 4°C overnight in 4% Paraformaldehyde (PFA). These
brains were washed with phosphate buffer saline (PBS), and after dehydration was
performed in 30% sucrose, they were embedded in Poly-freeze (Sigma, P0091). Finally,
a cryostat (Leica CM 1850) was used to obtain 12-16 µm thick sections of the chick
pallium.

RNA *in situ* hybridization

402
For microRNA *in situ* hybridization, a probe was designed where the 3'end of the
antisense sequence of miR-19b was labeled with digoxigenin (Eurofins MWG Operon).
This probe was used to perform section RNA *in situ* hybridization using a published
protocol (Thompson, Deo, and Turner 2007). Finally, the detection was performed using
another published method (Trimarchi et al. 2007).

The mRNA *in situ* hybridization was performed on the tissue sections using a published
method (Trimarchi et al. 2007). The following cDNA clones were used to prepare probes:

(1) Fezf2 (pTA2-cFezf2) was a gift from Prof. T. Hirata, National Insitutute of Genetics,
Japan, (2) E2f8 (ChEST 638e24) was procured from Chicken EST collection from
Source Biosciences (previously MRC Geneservice, UK).
414

In silico analysis of miR-19b target mRNAs

416 The list of target mRNAs was obtained using Targetscan & miRDB. Using venn diagram
417 tool (https://bioinformatics.psb.ugent.be/webtools/Venn/) intersection revealed 423 target
418 mRNAs. After that, plot & gene ontology analysis were performed using online tool
419 Enrichr (https://maayanlab.cloud/Enrichr/#).
420

Plasmid construction

1. pRmiR-19b: pRmiR-Empty construct (Smith et al. 2009) after BsaI digestion, was
ligated with the synthesized primary sequence of miR-19b (Macrogen Inc.) to
generate pRmiR-19b.
2. Sensor-ND1-Target-Region: The chicken NeuroD1 3'UTR having miR-19b target
region with flanking NotI and HindIII sites were synthesized (Macrogen Inc.) and
ligated in pCAG-mCherry construct (a gift from Prof. Constance Cepko, Harvard
Medical School, USA), downstream of the mCherry. Chicken NeuroD1 target
region: (NM_204920.1 from 1380[th] nucleotide position to 1458[th] nucleotide
position)
5'TACTACTTGTTACCTTTCCCGCGCCGAATTATTTTGTTGTGGTTTTGTACAG
AATTTTTAATGACTTTTTATAACGTG3'.
3. Sensor-E2f8-Target-Region: The chicken E2f8 3'UTR having miR-19b target
region with flanking NotI and HindIII sites were synthesized (Macrogen Inc.) and
ligated in pCAG-mCherry construct (a gift from Prof. Constance Cepko, Harvard
Medical School, USA), downstream of the mCherry. Chicken E2f8 target region:
XM_040672655.2 from 3108[th] nucleotide position to 3167[th] nucleotide position)
5'AAGGTACAGTCTGTCAATCAGTGCATATTTTGCACAAAGGAAGTACTGTGA
GCATGTACA3'.
4. pSponge-19b: For this first three copies of miR-19b target sequence (written in
small letters) was ligated in the pRmiR vector (Smith et al. 2009) to obtain
pSponge-19b-3x construct
5'TGCTGCCGtcagttttgcaccttttgcacaCCGGtcagttttgcaccttttgcacaCCGGtcagtt
ttgcaccttttgcacaCCGG3'. After that, using SalI, BglII, and BamHI Sponge-19b-3X
sequence was chained in tandem to obtain twelve copies of sponge-19b and this
construct was named pSponge-19b (Ebert and Sharp 2010) results in loss-of-
function of miR-19b.
5. pCAG-GFP was a gift from Prof. C. Cepko, Harvard Medical School, USA.
6. pCMV-SPORT6-CDKN1A from MGC clones.
450

Sensor assay

452 HEK-293T cells were transfected using Turbofect (Thermo Fisher Scientific Inc., R0533)
453 as per manufacturer's protocol, with pRmiR vector(s) and respective Sensor vector(s)
454 following which GFP and mCherry fluorescence were observed after 72 hours.

In ovo electroporation

Fertilized chicken eggs were incubated overnight and lowered by taking out 2-3ml of
albumin. Then, a window cut in the shell on the top and into one of the forebrain vesicles
the DNA construct (2µg/µl) with 0.1% of fast green (Sigma-Aldrich, 44715) was injected.

Platinum hockey stick electrodes (Nepagene, Japan) were used to deliver an electric
current of 13V in five pulses for 50ms duration with a gap of 950ms, to the HH22 chick
embryo head, using an electroporator (ECM830 Harvard instruments, USA). Before
sealing with the tape, Penicillin, and Streptomycin with sterile PBS, were added over the
embryos. Finally, harvesting was done at the desired HH stage.
464
Immunofluorescence
Briefly, after 4% PFA fixation for 5 minutes, three PBT washes were performed. Blocking
was performed with 5% of heat-inactivated goat serum (HINGS) for 1 hour. Primary
antibody was added and incubated overnight at 4°C. The primary antibodies were used
at the following dilution: anti-Mef2c, 1:200 (Millipore-Sigma, HPA005533); anti-NeuroD1,
1:1500 (Abcam, ab60704), anti-PH3, 1:300 (Sigma, H0412). After washing three times in
PBT for five minutes each, the appropriate secondary antibody was added. Secondary
antibodies used were as follows: Alexa Flour 594-conjugated Anti-Mouse IgG, 1:250
(Jackson ImmunoResearch Laboratories, INC., 115-585-003) and Alexa Flour 594-
conjugated Anti-Rabbit IgG, 1:250 (Jackson ImmunoResearch Laboratories, INC., 111-
585-045).
476
Cell-Cycle analysis
Chicken embryonic fibroblast, DF1 cells were transfected with respective DNA
constructs using Turbofect (Thermo Fisher Scientific Inc., R0533), as per manufacturer's
protocol. Cells were harvested after Trypsin treatment for 2-4 minutes then after cells
were washed with DMEM media. After that cells were passed through strainer to obtain
single-cell suspension, followed by treatment with 70% ice-cold ethanol for 30 minutes at
4dC. After that, two washes of PBT (PBS with 0.02% Tween-20) were done. After that
cells were treated with RNase A (10mg/ml) for 30 minutes at room temperature. For
labelling 10μl of 10/ng/μl   To-Pro-Far Red (Invitrogen, T3605) was added to the cells
for 15 minutes at room temperature after that washed with PBS. Fluorescence of
individual cells were quantified using flow cytometer (BD FACS Calibur, Becton
Dickinson, USA) using the facility at Central Drug Research Institute, Lucknow, India.
Data were analyzed using BD FACSDiva software (Becton Dickinson, USA).
490
EdU labeling and detection
After electroporation of desired DNA constructs at HH23, and then 2 hours before
harvesting 25μg of EdU (ThermoFisher Scientific, C10338) was overlaid in 200μl of 1X
Phosphate Buffer Saline (PBS). After harvesting at E4.5 (HH25), brains were dissected
out in PBS and emersion fixed in 4% paraformaldehyde and after sequential dehydration
with up to 30% sucrose, brains were embedded in the Poly-freeze media (Millipore-
Sigma, SHH0026) and 14 micrometer sections were generated and mounted on glass
slide. EdU detection was performed on these sections using CLICK-IT EdU-labeling kit
(ThermoFisher Scientific, C10338) as per manufacturer's protocol.
500
Image acquisition and statistical analysis
To obtain the images of RNA *in situ* hybridization as well as immunofluorescence a
Leica stereomicroscope (DM500B) equipped with a DFC500 camera was used. Other
fluorescent images were obtained using the multi-photon laser scanning confocal
microscope LSM780 (Carl Zeiss Inc.) and ZEN 2011 software.

Each set of experiments was repeated at least three times; for animals, n=1 means on
one animal experiment was performed; for cell culture, n=1 means in one dish
experiment was performed. Only samples with GFP and/or mCherry expression after
transfection or electroporation were included in the analysis. ImageJ software was used
to measure the fluorescence intensity. Graph Pad Prism 10 were used to plot the graph
and to perform unpaired student's t-test and calculate the significance.

Acknowledgements
We acknowledge Prof. Amitabha Bandhopadhyay for suggestions on the manuscript and Ms.
Neetu Dey (IIT Kanpur, India) for microscopy. Mr. D.N. Vishwakarma (CDRI, Lucknow, India)
for flow-cytometry. Funded by SERB-DST, Govt. of India. (EMR/2016/000886) to J.S. CSIR,
Govt. of India to S.K.S. for PhD (09/092(0824)/2011-EMR-I). MHRD, Govt. of India. to A.M.
and N.U. for PhD.

____

____

Additional information

Funding

Funder	Grant reference number	Author
Science and Engineering Research Board - Department of Science & Technology, India	EMR/2016/000886	Jonaki Sen
Council of Scientific and Industrial Engineering-Government of India	09/092(0824)/2011-EMR-I	Suvimal Kumar Sindhu

The funders had no role in study design, data collection and interpretation, or the
decision to submit the work of publication.

____

____

Author contributions
S.K.S., and J.S. designed study; S.K.S., A.M., N.U. performed experiments; S.K.S., A.M.,
N.U., and J.S. analyzed data; J.S. Supervised; S.K.S., A.M., and J.S. wrote the paper.

Author ORCIDs
Suvimal Kumar Sindhu ID: 0000-0002-7461-3104
Jonaki Sen ID: 0000-0001-7316-6376

Ethics

553 All experiments conducted in this study were in accordance with the guidelines of Indian
554 Institute of Technology Kanpur, India.
555
556 **Additional files**
557
558 Data availability
559 All data generated or analyzed during this study are including in the manuscript and
560 supporting files.
561

References

Abdel-Mannan, O., Cheung, A. F., & Molnar, Z. (2008). Evolution of cortical neurogenesis. *Brain*
Res Bull, *75*(2-4), 398-404. https://doi.org/10.1016/j.brainresbull.2007.10.047
Abellan, A., Desfilis, E., & Medina, L. (2014). Combinatorial expression of Lef1, Lhx2, Lhx5, Lhx9,
Lmo3, Lmo4, and Prox1 helps to identify comparable subdivisions in the developing
hippocampal formation of mouse and chicken. *Front Neuroanat*, *8*, 59.
https://doi.org/10.3389/fnana.2014.00059
Agarwal, V., Bell, G. W., Nam, J. W., & Bartel, D. P. (2015). Predicting effective microRNA target
sites in mammalian mRNAs. *Elife*, *4*. https://doi.org/10.7554/eLife.05005
Andersson, T., Rahman, S., Sansom, S. N., Alsio, J. M., Kaneda, M., Smith, J., O'Carroll, D.,
Tarakhovsky, A., & Livesey, F. J. (2010). Reversible block of mouse neural stem cell
differentiation in the absence of dicer and microRNAs. *PLoS One*, *5*(10), e13453.
https://doi.org/10.1371/journal.pone.0013453
Atoji, Y., & Wild, J. M. (2012). Afferent and efferent projections of the mesopallium in the pigeon
(Columba livia). *J Comp Neurol*, *520*(4), 717-741. https://doi.org/10.1002/cne.22763
Ball, G. F., & Balthazart, J. (2021). Evolutionary neuroscience: Are the brains of birds and
mammals really so different? *Curr Biol*, *31*(13), R840-R842.
https://doi.org/10.1016/j.cub.2021.05.004
Bartel, D. P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*, *116*(2),
281-297. https://doi.org/10.1016/s0092-8674(04)00045-5
Bian, S., Hong, J., Li, Q., Schebelle, L., Pollock, A., Knauss, J. L., Garg, V., & Sun, T. (2013).
MicroRNA cluster miR-17-92 regulates neural stem cell expansion and transition to
intermediate progenitors in the developing mouse neocortex. *Cell Rep*, *3*(5), 1398-1406.
https://doi.org/10.1016/j.celrep.2013.03.037
Boutin, C., Hardt, O., de Chevigny, A., Core, N., Goebbels, S., Seidenfaden, R., Bosio, A., &
Cremer, H. (2010). NeuroD1 induces terminal neuronal differentiation in olfactory
neurogenesis. *Proc Natl Acad Sci U S A*, *107*(3), 1201-1206.
https://doi.org/10.1073/pnas.0909015107
Butler, A. B., Reiner, A., & Karten, H. J. (2011). Evolution of the amniote pallium and the origins of
mammalian neocortex. *Ann N Y Acad Sci*, *1225*, 14-27. https://doi.org/10.1111/j.1749-
6632.2011.06006.x
Cardenas, A., Villalba, A., de Juan Romero, C., Pico, E., Kyrousi, C., Tzika, A. C., Tessier-Lavigne,
M., Ma, L., Drukker, M., Cappello, S., & Borrell, V. (2018). Evolution of Cortical
Neurogenesis in Amniotes Controlled by Robo Signaling Levels. *Cell*, *174*(3), 590-606
e521. https://doi.org/10.1016/j.cell.2018.06.007
Chen, B., Wang, S. S., Hattox, A. M., Rayburn, H., Nelson, S. B., & McConnell, S. K. (2008). The
Fezf2-Ctip2 genetic pathway regulates the fate choice of subcortical projection neurons

in the developing cerebral cortex. *Proc Natl Acad Sci U S A*, *105*(32), 11382-11387.
https://doi.org/10.1073/pnas.0804918105
Chen, Y., & Wang, X. (2020). miRDB: an online database for prediction of functional microRNA
targets. *Nucleic Acids Res*, *48*(D1), D127-D131. https://doi.org/10.1093/nar/gkz757
Cheung, A. F., Pollen, A. A., Tavare, A., DeProto, J., & Molnar, Z. (2007). Comparative aspects of
cortical   neurogenesis   in   vertebrates.   *J   Anat*,   *211*(2),   164-176.
https://doi.org/10.1111/j.1469-7580.2007.00769.x
Christensen, J., Cloos, P., Toftegaard, U., Klinkenberg, D., Bracken, A. P., Trinh, E., Heeran, M., Di
Stefano, L., & Helin, K. (2005). Characterization of E2F8, a novel E2F-like cell-cycle
regulated repressor of E2F-activated transcription. *Nucleic Acids Res*, *33*(17), 5458-5470.
https://doi.org/10.1093/nar/gki855
D'Amico, L. A., Boujard, D., & Coumailleau, P. (2013). The neurogenic factor NeuroD1 is
expressed in post-mitotic cells during juvenile and adult Xenopus neurogenesis and not
in   progenitor   or   radial   glial   cells.   *PLoS   One*,   *8*(6),   e66487.
https://doi.org/10.1371/journal.pone.0066487
De Pietri Tonelli, D., Pulvers, J. N., Haffner, C., Murchison, E. P., Hannon, G. J., & Huttner, W. B.
(2008). miRNAs are essential for survival and differentiation of newborn neurons but not
for expansion of neural progenitors during early neurogenesis in the mouse embryonic
neocortex. *Development*, *135*(23), 3911-3921. https://doi.org/10.1242/dev.025080
Deng, C., Zhang, P., Harper, J. W., Elledge, S. J., & Leder, P. (1995). Mice lacking p21CIP1/WAF1
undergo normal development, but are defective in G1 checkpoint control. *Cell*, *82*(4),
675-684. https://doi.org/10.1016/0092-8674(95)90039-x
Diaz, J. L., Siththanandan, V. B., Lu, V., Gonzalez-Nava, N., Pasquina, L., MacDonald, J. L.,
Woodworth, M. B., Ozkan, A., Nair, R., He, Z., Sahni, V., Sarnow, P., Palmer, T. D., Macklis,
J. D., & Tharin, S. (2020). An evolutionarily acquired microRNA shapes development of
mammalian cortical projections. *Proc Natl Acad Sci U S A*, *117*(46), 29113-29122.
https://doi.org/10.1073/pnas.2006700117
Dugas-Ford, J., Rowell, J. J., & Ragsdale, C. W. (2012). Cell-type homologies and the origins of the
neocortex.   *Proc   Natl   Acad   Sci   U   S   A*,   *109*(42),   16974-16979.
https://doi.org/10.1073/pnas.1204773109
Ebert, M. S., & Sharp, P. A. (2010). MicroRNA sponges: progress and possibilities. *RNA*, *16*(11),
2043-2050. https://doi.org/10.1261/rna.2414110
Fan, Y., Yin, S., Hao, Y., Yang, J., Zhang, H., Sun, C., Ma, M., Chang, Q., & Xi, J. J. (2014). miR-19b
promotes tumor growth and metastasis via targeting TP53. *RNA*, *20*(6), 765-772.
https://doi.org/10.1261/rna.043026.113
Gao, Z., Ure, K., Ables, J. L., Lagace, D. C., Nave, K. A., Goebbels, S., Eisch, A. J., & Hsieh, J. (2009).
Neurod1 is essential for the survival and maturation of adult-born neurons. *Nat
Neurosci*, *12*(9), 1090-1092. https://doi.org/10.1038/nn.2385
Gupta, S., Maurya, R., Saxena, M., & Sen, J. (2012). Defining structural homology between the
mammalian and avian hippocampus through conserved gene expression patterns
observed   in   the   chick   embryo.   *Dev   Biol*,   *366*(2),   125-141.
https://doi.org/10.1016/j.ydbio.2012.03.027
Hamburger, V., & Hamilton, H. L. (1951). A series of normal stages in the development of the
chick   embryo.   *J   Morphol*,   *88*(1),   49-92.
https://www.ncbi.nlm.nih.gov/pubmed/24539719

Harper, J. W., Adami, G. R., Wei, N., Keyomarsi, K., & Elledge, S. J. (1993). The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell*, *75*(4), 805-816. https://doi.org/10.1016/0092-8674(93)90499-g

Haubensak, W., Attardo, A., Denk, W., & Huttner, W. B. (2004). Neurons arise in the basal neuroepithelium of the early mammalian telencephalon: a major site of neurogenesis. *Proc Natl Acad Sci U S A*, *101*(9), 3196-3201. https://doi.org/10.1073/pnas.0308600100

Karten, H. J. (1997). Evolutionary developmental biology meets the brain: the origins of mammalian cortex. *Proc Natl Acad Sci U S A*, *94*(7), 2800-2804. https://doi.org/10.1073/pnas.94.7.2800

Kawase-Koga, Y., Low, R., Otaegi, G., Pollock, A., Deng, H., Eisenhaber, F., Maurer-Stroh, S., & Sun, T. (2010). RNAase-III enzyme Dicer maintains signaling pathways for differentiation and survival in mouse cortical neural stem cells. *J Cell Sci*, *123*(Pt 4), 586-594. https://doi.org/10.1242/jcs.059659

Kim, J. W., You, Y. H., Jung, S., Suh-Kim, H., Lee, I. K., Cho, J. H., & Yoon, K. H. (2013). miRNA-30a-5p-mediated silencing of Beta2/NeuroD expression is an important initial event of glucotoxicity-induced beta cell dysfunction in rodent models. *Diabetologia*, *56*(4), 847-855. https://doi.org/10.1007/s00125-012-2812-x

Lange, C., Huttner, W. B., & Calegari, F. (2009). Cdk4/cyclinD1 overexpression in neural stem cells shortens G1, delays neurogenesis, and promotes the generation and expansion of basal progenitors. *Cell Stem Cell*, *5*(3), 320-331. https://doi.org/10.1016/j.stem.2009.05.026

Lee, J. E., Hollenberg, S. M., Snider, L., Turner, D. L., Lipnick, N., & Weintraub, H. (1995). Conversion of Xenopus ectoderm into neurons by NeuroD, a basic helix-loop-helix protein. *Science*, *268*(5212), 836-844. https://doi.org/10.1126/science.7754368

Lee, J. K., Cho, J. H., Hwang, W. S., Lee, Y. D., Reu, D. S., & Suh-Kim, H. (2000). Expression of neuroD/BETA2 in mitotic and postmitotic neuronal cells during the development of nervous system. *Dev Dyn*, *217*(4), 361-367. https://doi.org/10.1002/(SICI)1097-0177(200004)217:4<361::AID-DVDY3>3.0.CO;2-8

Li, J., Ran, C., Li, E., Gordon, F., Comstock, G., Siddiqui, H., Cleghorn, W., Chen, H. Z., Kornacker, K., Liu, C. G., Pandit, S. K., Khanizadeh, M., Weinstein, M., Leone, G., & de Bruin, A. (2008). Synergistic function of E2F7 and E2F8 is essential for cell survival and embryonic development. *Dev Cell*, *14*(1), 62-75. https://doi.org/10.1016/j.devcel.2007.10.017

Lim, Y. J., Park, S. A., Wang, D., Jin, W., Ku, W. L., Zhang, D., Xu, J., Patino, L. C., Liu, N., Chen, W., Kazmi, R., Zhao, K., Zhang, Y. E., Sun, L., & Chen, W. (2024). MicroRNA-19b exacerbates systemic sclerosis through promoting Th9 cells. *Cell Rep*, *43*(8), 114565. https://doi.org/10.1016/j.celrep.2024.114565

Liu, K., Liu, Y., Mo, W., Qiu, R., Wang, X., Wu, J. Y., & He, R. (2011). MiR-124 regulates early neurogenesis in the optic vesicle and forebrain, targeting NeuroD1. *Nucleic Acids Res*, *39*(7), 2869-2879. https://doi.org/10.1093/nar/gkq904

Maiti, B., Li, J., de Bruin, A., Gordon, F., Timmers, C., Opavsky, R., Patil, K., Tuttle, J., Cleghorn, W., & Leone, G. (2005). Cloning and characterization of mouse E2F8, a novel mammalian E2F family member capable of blocking cellular proliferation. *J Biol Chem*, *280*(18), 18211-18220. https://doi.org/10.1074/jbc.M501410200

McConnell, S. K., & Kaznowski, C. E. (1991). Cell cycle dependence of laminar determination in developing neocortex. *Science*, *254*(5029), 282-285. https://doi.org/10.1126/science.254.5029.282

Mendell, J. T. (2008). miRiad roles for the miR-17-92 cluster in development and disease. *Cell*,
133(2), 217-222. https://doi.org/10.1016/j.cell.2008.04.001
Miyata, T., Kawaguchi, A., Saito, K., Kawano, M., Muto, T., & Ogawa, M. (2004). Asymmetric
production of surface-dividing and non-surface-dividing cortical progenitor cells.
Development, *131*(13), 3133-3145. https://doi.org/10.1242/dev.01173
Molyneaux, B. J., Arlotta, P., Menezes, J. R., & Macklis, J. D. (2007). Neuronal subtype
specification in the cerebral cortex. *Nat Rev Neurosci*, *8*(6), 427-437.
https://doi.org/10.1038/nrn2151
Nguyen, L., Besson, A., Heng, J. I., Schuurmans, C., Teboul, L., Parras, C., Philpott, A., Roberts, J.
M., & Guillemot, F. (2006). p27kip1 independently promotes neuronal differentiation
and migration in the cerebral cortex. *Genes Dev*, *20*(11), 1511-1524.
https://doi.org/10.1101/gad.377106
Noctor, S. C., Martinez-Cerdeno, V., Ivic, L., & Kriegstein, A. R. (2004). Cortical neurons arise in
symmetric and asymmetric division zones and migrate through specific phases. *Nat
Neurosci*, *7*(2), 136-144. https://doi.org/10.1038/nn1172
Nomura, T., Ohtaka-Maruyama, C., Yamashita, W., Wakamatsu, Y., Murakami, Y., Calegari, F.,
Suzuki, K., Gotoh, H., & Ono, K. (2016). The evolution of basal progenitors in the
developing non-mammalian brain. *Development*, *143*(1), 66-74.
https://doi.org/10.1242/dev.127100
Nowakowski, T. J., Mysiak, K. S., Pratt, T., & Price, D. J. (2011). Functional dicer is necessary for
appropriate specification of radial glia during early development of mouse
telencephalon. *PLoS One*, *6*(8), e23013. https://doi.org/10.1371/journal.pone.0023013
Ochocinska, M. J., & Hitchcock, P. F. (2009). NeuroD regulates proliferation of photoreceptor
progenitors in the retina of the zebrafish. *Mech Dev*, *126*(3-4), 128-141.
https://doi.org/10.1016/j.mod.2008.11.009
Ohtaka-Maruyama, C., & Okado, H. (2015). Molecular Pathways Underlying Projection Neuron
Production and Migration during Cerebral Cortical Development. *Front Neurosci*, *9*, 447.
https://doi.org/10.3389/fnins.2015.00447
Olive, V., Bennett, M. J., Walker, J. C., Ma, C., Jiang, I., Cordon-Cardo, C., Li, Q. J., Lowe, S. W.,
Hannon, G. J., & He, L. (2009). miR-19 is a key oncogenic component of mir-17-92. *Genes
Dev*, *23*(24), 2839-2849. https://doi.org/10.1101/gad.1861409
Park, S. A., Lim, Y. J., Ku, W. L., Zhang, D., Cui, K., Tang, L. Y., Chia, C., Zanvit, P., Chen, Z., Jin, W.,
Wang, D., Xu, J., Liu, O., Wang, F., Cain, A., Guo, N., Nakatsukasa, H., Wu, C., Zhang, Y.
E.,…Chen, W. (2022). Opposing functions of circadian protein DBP and atypical E2F
family E2F8 in anti-tumor Th9 cell differentiation. *Nat Commun*, *13*(1), 6069.
https://doi.org/10.1038/s41467-022-33733-8
Pataskar, A., Jung, J., Smialowski, P., Noack, F., Calegari, F., Straub, T., & Tiwari, V. K. (2016).
NeuroD1 reprograms chromatin and transcription factor landscapes to induce the
neuronal program. *EMBO J*, *35*(1), 24-45. https://doi.org/10.15252/embj.201591206
Puelles, L., Kuwana, E., Puelles, E., Bulfone, A., Shimamura, K., Keleher, J., Smiga, S., &
Rubenstein, J. L. (2000). Palial and subpallial derivatives in the embryonic chick and
mouse telencephalon, traced by the expression of the genes Dlx-2, Emx-1, Nkx-2.1, Pax-
6, and Tbr-1. *J Comp Neurol*, *424*(3), 409-438. https://doi.org/10.1002/1096-
9861(20000828)424:3<409::aid-cne3>3.0.co;2-7
Roybon, L., Hjalt, T., Stott, S., Guillemot, F., Li, J. Y., & Brundin, P. (2009). Neurogenin2 directs
granule neuroblast production and amplification while NeuroD1 specifies neuronal fate

during hippocampal neurogenesis. *PLoS One*, *4*(3), e4779. https://doi.org/10.1371/journal.pone.0004779

Schwab, M. H., Bartholomae, A., Heimrich, B., Feldmeyer, D., Druffel-Augustin, S., Goebbels, S., Naya, F. J., Zhao, S., Frotscher, M., Tsai, M. J., & Nave, K. A. (2000). Neuronal basic helix-loop-helix proteins (NEX and BETA2/Neuro D) regulate terminal granule cell differentiation in the hippocampus. *J Neurosci*, *20*(10), 3714-3724. https://doi.org/10.1523/JNEUROSCI.20-10-03714.2000

Shen, Q., Wang, Y., Dimos, J. T., Fasano, C. A., Phoenix, T. N., Lemischka, I. R., Ivanova, N. B., Stifani, S., Morrisey, E. E., & Temple, S. (2006). The timing of cortical neurogenesis is encoded within lineages of individual progenitor cells. *Nat Neurosci*, *9*(6), 743-751. https://doi.org/10.1038/nn1694

Shu, P., Wu, C., Ruan, X., Liu, W., Hou, L., Fu, H., Wang, M., Liu, C., Zeng, Y., Chen, P., Yin, B., Yuan, J., Qiang, B., Peng, X., & Zhong, W. (2019). Opposing Gradients of MicroRNA Expression Temporally Pattern Layer Formation in the Developing Neocortex. *Dev Cell*, *49*(5), 764-785 e764. https://doi.org/10.1016/j.devcel.2019.04.017

Sindhu, S. K., Udaykumar, N., Zaidi, M. A. A., Soni, A., & Sen, J. (2019). MicroRNA-19b restricts Wnt7b to the hem, which induces aspects of hippocampus development in the avian forebrain. *Development*, *146*(20). https://doi.org/10.1242/dev.175729

Stacho, M., Herold, C., Rook, N., Wagner, H., Axer, M., Amunts, K., & Gunturkun, O. (2020). A cortex-like canonical circuit in the avian forebrain. *Science*, *369*(6511). https://doi.org/10.1126/science.abc5534

Suzuki, I. K., & Hirata, T. (2014). A common developmental plan for neocortical gene-expressing neurons in the pallium of the domestic chicken Gallus gallus domesticus and the Chinese softshell turtle Pelodiscus sinensis. *Front Neuroanat*, *8*, 20. https://doi.org/10.3389/fnana.2014.00020

Suzuki, I. K., Kawasaki, T., Gojobori, T., & Hirata, T. (2012). The temporal sequence of the mammalian neocortical neurogenetic program drives mediolateral pattern in the chick pallium. *Dev Cell*, *22*(4), 863-870. https://doi.org/10.1016/j.devcel.2012.01.004

Thurlings, I., Martinez-Lopez, L. M., Westendorp, B., Zijp, M., Kuiper, R., Tooten, P., Kent, L. N., Leone, G., Vos, H. J., Burgering, B., & de Bruin, A. (2017). Synergistic functions of E2F7 and E2F8 are critical to suppress stress-induced skin cancer. *Oncogene*, *36*(6), 829-839. https://doi.org/10.1038/onc.2016.251

Todorov, H., Weissbach, S., Schlichtholz, L., Mueller, H., Hartwich, D., Gerber, S., & Winter, J. (2024). Stage-specific expression patterns and co-targeting relationships among miRNAs in the developing mouse cerebral cortex. *Commun Biol*, *7*(1), 1366. https://doi.org/10.1038/s42003-024-07092-7

Ventura, A., Young, A. G., Winslow, M. M., Lintault, L., Meissner, A., Erkeland, S. J., Newman, J., Bronson, R. T., Crowley, D., Stone, J. R., Jaenisch, R., Sharp, P. A., & Jacks, T. (2008). Targeted deletion reveals essential and overlapping functions of the miR-17 through 92 family of miRNA clusters. *Cell*, *132*(5), 875-886. https://doi.org/10.1016/j.cell.2008.02.019

Wang, Y., Brzozowska-Prechtl, A., & Karten, H. J. (2010). Laminar and columnar auditory cortex in avian brain. *Proc Natl Acad Sci U S A*, *107*(28), 12676-12681. https://doi.org/10.1073/pnas.1006645107

Xu, D., Zhong, L. T., Cheng, H. Y., Wang, Z. Q., Chen, X. M., Feng, A. Y., Chen, W. Y., Chen, G., & Xu, Y. (2023). Overexpressing NeuroD1 reprograms Muller cells into various types of retinal

neurons. *Neural Regen Res*, *18*(5), 1124-1131. https://doi.org/10.4103/1673-
5374.355818
Yan, R. T., & Wang, S. Z. (2004). Requirement of neuroD for photoreceptor formation in the chick
retina. *Invest Ophthalmol Vis Sci*, *45*(1), 48-58. https://doi.org/10.1167/iovs.03-0774
Zhang, P., Wong, C., Liu, D., Finegold, M., Harper, J. W., & Elledge, S. J. (1999). p21(CIP1) and
p57(KIP2) control muscle differentiation at the myogenin step. *Genes Dev*, *13*(2), 213-
224. https://doi.org/10.1101/gad.13.2.213
Zhang, Z. W., Zhang, L. Q., Ding, L., Wang, F., Sun, Y. J., An, Y., Zhao, Y., Li, Y. H., & Teng, C. B.
(2011). MicroRNA-19b downregulates insulin 1 through targeting transcription factor
NeuroD1. *FEBS Lett*, *585*(16), 2592-2598. https://doi.org/10.1016/j.febslet.2011.06.039
Zheng, H., Ying, H., Yan, H., Kimmelman, A. C., Hiller, D. J., Chen, A. J., Perry, S. R., Tonon, G., Chu,
G. C., Ding, Z., Stommel, J. M., Dunn, K. L., Wiedemeyer, R., You, M. J., Brennan, C.,
Wang, Y. A., Ligon, K. L., Wong, W. H., Chin, L., & DePinho, R. A. (2008). p53 and Pten
control neural and glioma stem/progenitor cell renewal and differentiation. *Nature*,
455(7216), 1129-1133. https://doi.org/10.1038/nature07443
Zheng, H., Zeng, Y., Zhang, X., Chu, J., Loh, H. H., & Law, P. Y. (2010). mu-Opioid receptor agonists
differentially regulate the expression of miR-190 and NeuroD. *Mol Pharmacol*, *77*(1),
102-109. https://doi.org/10.1124/mol.109.060848
799

Figure legends

Figure 1. **Expression patterns of NEUROD1 and miR-19b in the developing chick pallium and their spatial distribution relative to PH3-positive cells.** (A) Screening of NEUROD1 (IHC) and miR-19b (RNA-ISH) expression within chicken pallium at different stages (A) A cross-section of HH24 chicken pallium showing NeuroD1 and miR-19b with a merged image in the inset. Scale bar- 500μm, Inset (100μm), n=3 (B) miR-19b and NeuroD1 expression dynamics in wulst (Red inset) and DVR (Yellow inset) region at stage HH29 chicken pallium. Scale bar- 500μm, Inset (250μm), n=3 (C) a similar comparison of miR-19b and NeuroD1 expression at stage HH29 chicken pallium. Scale bar - 800μm, inset (250μm), n=3 (D) Distribution of PH3 and NeuroD1- positive cells at all above-mentioned stages domain of PH3 positive cells is marked by green arrows wherein NeuroD1 is represented by red arrows. The intensity plot shows mutually exclusive expression domains of NeuroD1 and miR-19b. Insets depict magnified images. n=3, Scale bar: 800μm. White arrowheads show NeuroD1 positive cells in SVZ. A.U.= arbitrary unit; VZ= ventricular zone, SVZ= sub-ventricular zone.

Figure 2. miR-19b regulates E2f8 expression and cell-cycle progression.   (A) Venn diagram showing miR-19b targets predicted by miRDB and TargetScan in chicken (*Gallus gallus*) and gene ontology analysis for predicted targets. (B) List of functionally validated targets. The red square indicates the role in the mentioned process/pathway. Potential targets and associated processes highlighted in magenta. (C) Sensor-assay for in vitro demonstration of miR-19b binding to its target region of E2f8. Respective 3'UTRs fused with the coding sequence of mCherry were subjected to miR-19b manipulation. mCherry intensity was further quantified and plotted. Scale bar: 100μm, Plot bar: mean ± s.e.m., N=3 wells, t-test performed, **p≤0.01 and ****p≤0.0001. A.U.= arbitrary unit. (D) represent the conservation of mi19b binding sequences in E2f8 3' UTR

across all species. Effect of miR-19b manipulation on E2f8 (RNA-ISH) (E) and (F), scale
bar-300µm. (G) Flow cytometry experiment to analyze cell-cycle distribution of DF1 cells
upon manipulation of miR-19b. Go/G1, S and G2/M are presented in population group
named as P6, P7 and P8, respectively. Representative images of EdU labeling assay(H)
and PH3 immunohistochemistry (I)to demonstrate effect of miR-19b manipulation on cell
proliferation. Scale bar: 200µm (H), 50µm (I). n=3, Plot bar: mean ± s.e.m.; A.U.=
arbitrary unit. ns= not significant; VZ= ventricular zone, SVZ= sub-ventricular zone. t-test
was performed, *p≤0.1, **p≤0.01 and ***p≤0.001, ns = non-significant.

Figure 3. Expression Patterns of Fezf2 and Mef2c in the Embryonic Chick Pallium
and the Effects of CDKN1A Overexpression and miR-19b Loss of Function(A)
Expression patterns of Fezf2 and NeuroD1 in the SVZ and Mef2c in the dorsal pallium
layer at HH28. The inset highlights double-positive cells for Fezf2 and NeuroD1. Scale
bar = 300 µm; 25 µm (inset). n = 3. (B) Fezf2 expression is restricted to the APH, while
Mef2c expression extends into the mesopallium (Msp). The inset shows magnified
views, and the intensity plot depicts spatial expression profiles of Fezf2 and Mef2c.
Scale bar=150µm; n=3; DP= Dorsal Pallium. (C) Effect of ectopic CDKN1A expression
on Fezf2(ISH) and NeuroD1(IHC) expression. Electroporation of CDKN1A-
overexpressing and control constructs demonstrates altered expression of Fezf2.
Magnified regions highlight colocalization of Fezf2 and NeuroD1 (circled regions). Bar
graphs display signal intensity quantification; n = 4. Scale bar: 300µm (D) Impact of
CDKN1A overexpression on Mef2c expression (IHC). White arrows indicate double-
positive cells for Mef2c, and bar graphs quantify their abundance. n=3. (E) Loss of miR-
19b affects Fezf2 expression. Signal intensity of Fezf2 per unit electroporated region is
quantified and plotted. Bar graph plotted for quantification. n=4, Scale bar: 150µm; Plot
bar: mean ± s.e.m. A.U.= arbitrary unit. ns= not significant; VZ= ventricular zone, SVZ=
sub-ventricular zone; t-test was performed; *p≤0.1, **p≤0.01 and ***p≤0.001; ns = non-
significant.

Figure 4. miR-19b Mediated Regulation of NeuroD1 and Its Impact on Patterning in
the Developing Chick Pallium (A) mRNA in situ hybridization (ISH) screening of miR-
19b and NeuroD1 immunostaining in the chick pallium at HH29. The intensity plot
highlights mutually exclusive expression domains of NeuroD1 and miR-19b. Insets show
magnified views of the expression patterns. Scale bar = 100µm; 30µm (insets), n = 3. (B)
Conservation analysis of miR-19b binding sequences in the NeuroD1 3′ UTR across
species. (C) Sensor assay demonstrating in vitro binding of miR-19b to the target region
of NeuroD1. The 3′ UTR of NeuroD1 was fused with mCherry and subjected to miR-19b
manipulation. mCherry intensity was quantified and plotted. Scale bar = 100 µm, with n =
3 wells. A.U. = arbitrary units. (D, E) miR-19b regulates NeuroD1 expression in *in vivo* as
shown by immunostaining in gain-of-function (D) and loss-of-function (E) experiments.
Scale bar = 100 µm, n=4. (F) Expression patterns of NeuroD1 and Mef2c in the chick
pallium at HH29. Scale bar = 500µm. (G, H) NeuroD1 loss-of-function experiment using
NeuroD1-DBD-EnR electroporation shows loss of Mef2c ISH expression. (G) and IHC
expression (H). The yellow asterisk (control) and white asterisk (test) mark the Mef2c
mRNA expression domains. Scale bar:100µm, n=3 (I-J) NeuroD1 gain-of-function led to
ectopic Mef2c mRNA (I) and Proteins (J) shown by ISH and immunostaining
(arrowheads show ectopic expression of Mef2c). n=3, scale bar: 100µm. (K) Effect of
miR-19b overexpression and Sponge-19b expression on Mef2c expression. Yellow
asterisk is showing Mef2c ISH in control and white asterisk show decreased Mef2c

expression using ISH; n=3; Scale bar:100µm; Plot bar: mean ± s.e.m. A.U.= arbitrary
unit. ns= not significant; VZ= ventricular zone, SVZ= sub-ventricular zone; Msp =
Mesopallium; t-test was performed; *p≤0.1, **p≤0.01 and ***p≤0.001; ns = non-
significant.
878
879

Figure 1

Figure 2

Figure 3

Figure 4