Dot1L interacts with Zc3h10 to activate UCP1 transcription and promote thermogenesis

Danielle Yi[1,2], Hai P. Nguyen[1,2], Jennie Dinh[1], Jose A. Viscarra[1], Ying Xie[1], Jon M. Dempersmier[1], Yuhui Wang[1], and Hei Sook Sul[1,2,3]

Department of Nutritional Sciences & Toxicology[1], Endocrinology Program[2], University of California, Berkeley, CA 94720

[3]Corresponding author: hsul@berkeley.edu

ABSTRACT

Brown adipose tissue is a metabolically beneficial organ capable of dissipating chemical energy into heat, thereby increasing energy expenditure. Here, we identify Dot1L, the only known H3K79 methyltransferase, as an interacting partner of Zc3h10 that transcriptionally activates the UCP1 promoter and other BAT genes. Through a direct interaction, Dot1L is recruited by Zc3h10 to the promoter regions of thermogenic genes to function as a coactivator by methylating H3K79. We also show that Dot1L is induced during brown fat cell differentiation and by cold exposure and that Dot1L and its H3K79 methyltransferase activity is required for thermogenic gene program. Furthermore, we demonstrate that Dot1L ablation in mice using UCP1-Cre prevents activation of UCP1 and other target genes to reduce thermogenic capacity and energy expenditure, promoting adiposity. Hence, Dot1L plays a critical role in the thermogenic program and may present as a future target for obesity therapeutics.

INTRODUCTION

Adipose tissue has a central role in controlling mammalian energy metabolism in that, while white adipose tissue (WAT) is to store excess calories, brown adipose tissue (BAT) is to dissipate energy via non-shivering thermogenesis (Cannon and Nedergaard, 2004). Classic brown adipocytes contain a high density of mitochondria that contain Uncoupling protein 1 (UCP1), which uncouples respiration from ATP synthesis and generates heat instead (Farmer, 2008, Cannon and Nedergaard, 2004). In addition, beige/brite adipocytes in WAT depot are recruited upon cold exposure or β_3-adrenergic stimulation (Seale et al., 2008, Wang and Seale, 2016, Sanchez-Gurmaches et al., 2016, Cypess et al., 2009, Lichtenbelt et al., 2009, Virtanen et al., 2009, Chondronikola et al., 2014). Based on several cross-sectional studies, adult human brown fat or brown fat-like tissue is inversely correlated with body mass index and visceral fat (Lichtenbelt et al., 2009, Hibi et al., 2016, Jimenez et al., 2007). Therefore, unraveling the mechanisms underlying the thermogenic gene program has drawn growing attention in obesity research as a promising avenue to combat obesity and associated metabolic diseases.

One of the major advances in BAT biology has been understanding the transcription network that governs the thermogenic gene program and finding critical factors that activate the UCP1 gene. A multitude of transcriptional regulators have been implicated in the transcription of UCP1, including transcription factors, Zfp516, IRF4 and EBF2, and transcriptional coregulators, PRDM16, PGC1α and LSD1 (Seale et al., 2008, Rajakumari et al., 2013, Dempersmier et al., 2015, Puigserver et al., 1998, Sambeat et al., 2016, Kong et al., 2014). Recently, we identified a BAT-enriched and cold-induced transcription factor, Zc3h10 that activates the UCP1 and other target genes, such as Tfam and Nrf1 for mitochondrial biogenesis (Yi et al., 2019). Zc3h10 activates the UCP1 promoter by directly binding to the -4.6 kb region. Upon sympathetic stimulation, Zc3h10 is phosphorylated at S126 by p38 mitogen-activated protein kinase (MAPK) to increase binding to this distal region of the UCP1 promoter. Consequently, Zc3h10 ablation in

mice impairs the thermogenic gene program, while Zc3h10 overexpression in adipose tissue
enhances the thermogenic capacity and energy expenditure, protecting mice from diet-induced
obesity (Yi et al., 2019).
As in most biological processes, BAT as well as beige fat, rely heavily on
environmental cues for its full activation of the thermogenic gene program. Integrating
epigenetic effectors into the thermogenic network may provide a comprehensive understanding
in the regulation of thermogenic gene program (Yi et al., 2020). Here, we identify Dot1L
(disruptor of telemetric silencing 1-like) as an interacting partner of Zc3h10 and a critical
coactivator of thermogenic genes. Dot1L is the only known methyltransferase that catalyzes the
sequential mono-, di- and tri-methylation of H3K79, which, unlike major epigenetic sites, is
located at the globular domain of nucleosome (Min et al., 2003, van Leeuwen et al., 2002,
Frederiks et al., 2008). Differing from other histone methyltransferases, Dot1L does not contain
a SET domain but uniquely has an AdoMET motif (Feng et al., 2002, van Leeuwen et al., 2002).
Dot1L is broadly known to play roles in telomere silencing, cell cycle regulations and is
particularly well studied in mixed lineage leukemia (MLL)-related leukemogenesis (Schulze et
al., 2009, Okada et al., 2005, Nguyen and Zhang, 2011, Ng et al., 2002). However, not much is
known about Dot1L recruitment to specific sites by specific transcription factors. Nor the role of
Dot1L is thermogenic gene program is known.
We show here that Dot1L is recruited to the UCP1 promoter region via its direct
interaction to Zc3h10 for Zc3h10-mediated transcriptional activation at the UCP1 and other
target genes. By using the specific and potent chemical inhibitor of Dot1L-H3K79
methyltransferase activity, pinometostat (EPZ-5676), we demonstrate that Dot1L
methyltransferase activity is required for thermogenic gene expression in vitro and in vivo.
Moreover, Dot1L ablation in brown adipocytes impairs the thermogenic program, whereas
ectopic Dot1L expression enhances and thermogenic gene program. Dot1L ablation in UCP1$^+$
cells in mice impairs the thermogenic capacity and lowers oxygen consumption, leading to
weight gain. In this regard, the GWAS database reveals multiple SNPs of Dot1L associated with
a waist-hip ratio and body mass index, further supporting a potential role of Dot1L in human
obesity (GWAS Central identifier: HGVPM1111, HGVPM1114).
81
82   **RESULTS**
83
Dot1L is recruited by Zc3h10 for activation of BAT gene program
We previously reported Zc3h10 as a BAT-enriched transcription factor that promotes the
BAT gene program (Yi et al., 2019). Most transcription factors do not work alone but rather form
a complex to recruit other cofactors for transcription. Thus, as a DNA-binding protein, Zc3h10
may interact with coregulators to activate BAT gene transcription. Therefore, we searched for
potential interacting partners of Zc3h10. The Zc3h10 with the streptavidin- and calmodulin-
binding epitope-tag were incubated with nuclear extracts from BAT and the Zc3h10 complex
after sequential purification were subjected to mass spectrometry. We identified multiple
potential Zc3h10 interacting proteins, among which were methyltransferases and chromatin
remodelers (Supplementary Figure 1A). For further studying as a Zc3h10 interacting protein, we
selected Dot1L (disruptor of telemetric silencing 1-like), the H3K79 methyltransferase, as it
directly interacted with Zc3h10 and was the only BAT enriched gene out of the candidates
tested.
First, for validation of interaction between Dot1L and Zc3h10, we performed Co-IP
experiments with lysates of HEK293FT cells transfected with HA-Dot1L and Flag-Zc3h10 by
using FLAG and HA antibodies. Indeed, we detected FLAG-Zc3h10 upon immunoprecipitation
with HA antibodies. Conversely, HA-Dot1L was detected upon immunoprecipitation with FLAG
antibodies (Figure 1A, left). We then tested an interaction of the two endogenous proteins using
BAT from mice. Again, we detected the presence of endogenous Zc3h10 when nuclear extracts

of BAT from mice were immunoprecipitated with Dot1L antibody. By reverse Co-IP, we
confirmed the interaction between endogenous Dot1L and Zc3h10 (Figure 1A, right). We next
asked whether Dot1L can directly bind Zc3h10, by using glutathione S-transferase (GST) fused
to Dot1L expressed and purified from *E. Coli* (Supplementary Figure 1B). Incubation of GST-
Dot1L fusion protein immobilized on glutathione beads with *in vitro* transcribed and translated
[^{35}S]-Zc3h10, but not control GST-alone, detected the presence of Zc3h10 (Figure 1B). Overall,
these results demonstrate that Dot1L directly interacts with Zc3h10. We next examined domains
of Dot1L for its interaction with Zc3h10. We generated three Dot1L truncated constructs; N-
terminal Dot1L (1-500aa) containing the catalytic domain, the middle fragment (501-1000aa)
containing the coiled-coil domain, and the C-terminal domain (1001-1540aa); All constructs
were N-terminally tagged with HA. We then cotransfected these constructs with full-length Flag-
tagged Zc3h10. By Co-IP, we detected the middle fragment (501 -1000) of Dot1L, that contains
coiled coil motifs, interacting with Zc3h10, but not the other two regions of Dot1L (Figure 1C).
We conclude that the middle fragments of Dot1L contains the Zc3h10 interacting domain.
Since Zc3h10 activates UCP1 and other target genes for BAT gene program and Dot1L
interacts with Zc3h10, Dot1L expression pattern could be similar to Zc3h10 and thus BAT-
enriched. Indeed, tissue expression profiling by RT-qPCR and immunoblotting showed that
Dot1L was enriched in mouse BAT compared to other tissues tested, such as WAT, liver and
stomach (Figure 1D). We next compared Dot1L mRNA levels between the stromal vascular
fraction (SVF) that contains preadipocytes with adipocyte fraction of BAT from 10-wk-old wild-
type (WT) mice. As expected, UCP1 was enriched in the adipocyte fraction, whereas PDGFRα
expression was higher in the SVF fraction. We found Dot1L to be enriched in the brown
adipocyte fraction by over 2-fold compared to the SVF (Figure 1E, left). Moreover, similar to
Zc3h10, expression of Dot1L in BAT was induced upon cold exposure (Figure 1E, right). We
next examined Dot1L expression during BAT cell differentiation in vitro. During the course of
brown adipocyte differentiation, as expected, UCP1 expression was induced, whereas
expression of preadipocyte gene, Pref-1, was suppressed. More importantly, similar to Zc3h10,
Dot1L mRNA level was increased during BAT cell differentiation (Figure 1F). Overall, we
conclude that Dot1L expression pattern is similar to Zc3h10 in terms of enrichment in mature
brown adipocytes and induction by cold exposure, which may allow potential a cooperative
function of Dot1L and Zc3h10 for the BAT gene program.
Next, to examine the functional significance of Dot1L and Zc3h10 interaction in the
activation of UCP1 promoter, we performed the luciferase (Luc) reporter assay using the -5.5 kb
UCP1-Luc promoter. Along with the UCP1 promoter-Luc construct, Zc3h10 and Dot1L were
cotransfected into HEK293 cells. As expected, compared to empty vector control, Zc3h10 alone
activated the UCP1 promoter over 3-fold. Dot1L alone could not activate the UCP1 promoter.
When co-transfected with Zc3h10, Dot1L was able to synergistically activate the UCP1
promoter, resulting in a robust 11-fold increase in the UCP1 promoter activity (Figure 1G, left).
These results demonstrate the cooperative function of Dot1L and Zc3h10 in UCP1 promoter
activation. Next, because Dot1L interacts with Zc3h10 to enhance the UCP1 promoter activity,
we predicted that Dot1L should occupy the same UCP1 promoter region where Zc3h10 binds.
We performed chromatin immunoprecipitation (ChIP) using BAT cells transduced with
adenovirus containing Flag- tagged Zc3h10 and HA-tagged Dot1L. By using Flag (for Zc3h10)
and HA (for Dot1L) antibodies, we detected strong enrichment of both Zc3h10 and Dot1L at the
- 4.6 kb region of the UCP1 promoter, a region corresponding to the Zc3h10 binding site (Yi et
al., 2019) (Figure 1G, right). This illustrates co-occupancy of Dot1L and Zc3h10 at the same
region of the UCP1 promoter. Collectively, we demonstrate that Dot1L, as an interacting partner
of Zc3h10, is a critical coactivator of the UCP1 promoter.
151

Figure 1. Dot1L directly interacts with Zc3h10 for UCP1 promoter activation

Figure 1. Dot1L directly interacts with Zc3h10 for UCP1 promoter activation
(A) (Left) CoIP using αFlag for Zc3h10 or αHA for Dot1L after IP with either αHA- or αFlag,
respectively, using lysates from HEK293FT cells transfected with Flag-Zc3h10 and HA-Dot1L.
(Right) CoIP of endogenous Zc3h10 and Dot1L using BAT tissue of C57BL/6 using αZc3h10
and αDot1L.
(B) Autoradiograph of GST pull-down using GST-Dot1L and ^{35}S-labeled in vitro transcribed/
translated Zc3h10.
(C) CoIP using αFlag for Zc3h10 after IP with αHA for Dot1L using lysates from HEK293FT
cells transfected with Flag-Zc3h10 and various HA-Dot1L constructs.
(D) RT-qPCR and immunoblotting of Dot1L in various tissues from 10-wk-old C57BL/6 mice.
(E) (Left) RT-qPCR for indicated genes in the adipocyte fraction and SVF from BAT. (Right) RT-
qPCR for Dot1L mRNA of BAT and iWAT from mice housed at either 30°C or 4°C (n=4).
(F) RT-qPCR for indicated genes during BAT cell differentiation.

166   **(G)** (Left) HEK293FT cells were cotransfected with the -5.5 kb UCP1-Luc promoter with Zc3h10
167   or Dot1L alone or together. (Right) ChIP-qPCR for Zc3h10 and Dot1L enrichment at the -4.6 kb
168   region of UCP1 promoter using Flag or HA antibodies. Differentiated BAT cells transduced with
169   either Flag-Zc3h10 or HA-Dot1L were used.
170
Dot1L promotes the thermogenic gene program in vitro
We next tested whether Dot1L is required for BAT gene program in cultured BAT cells.
BAT cells at Day 2 of differentiation were transduced with adenovirus expressing short hairpin
RNAs targeting Dot1L for knockdown (Dot1L KD). Transduction of shDot1L adenovirus in BAT
cells caused a decrease in Dot1L mRNA levels by approximately 60% (Figure 2A, left). While
there were no apparent changes in expression of transcription factors critical for brown
adipocyte differentiation, such as PPARγ, there was a 50% reduction of UCP1 expression at
mRNA and protein levels (Figure 2A). Expression of other Zc3h10 target genes, including Tfam,
Nrf1 and Elovl3, was also decreased significantly (Figure 2A, middle). Moreover, Dot1L KD BAT
cells had decreased oxygen consumption rate (OCR) in oligomycin-, FCCP-, and
rotenone/antimycin A-treated conditions (Figure 2B, left). Notably, the uncoupled respiration of
Dot1L KD cells was significantly lower than the control BAT cells in both basal and in forskolin-
treated stimulated conditions (Figure 2B, right). These results suggest that Dot1L is critical for
full activation of the BAT gene program and mitochondriogenesis, and thus thermogenic
function in BAT cells.
Next, in order to further establish the role for Dot1L in BAT gene program, we treated
BAT cells with a specific inhibitor of Dot1L H3K79 methyltransferase, EPZ5676. We predicted
that if Dot1L is required for the transcriptional activation of UCP1 and other BAT genes,
inhibition of Dot1L activity should prevent BAT gene expression. Dot1L inhibition resulted in a
significant reduction in mRNA and protein levels for UCP1 by 90% (Figure 2C). Expression of
other BAT-enriched Zc3h10 target genes, such as Tfam, Nrf1, Cox7a and Cox8b, was
decreased by 40-80%, whereas PPARγ expression was not affected (Figure 2C). These results
show that Dot1L methyltransferase activity is required for thermogenic gene expression and
other Zc3h10 target genes for the BAT gene program.
We then tested the ability of Dot1L to induce UCP1 and other thermogenic genes by
gain of function experiment. We overexpressed Dot1L or Zc3h10 alone or together via
adenoviral transduction in 3T3-L1 cells and the cells were induced to browning by forskolin
treatment. We first verified Zc3h10 and Dot1L overexpression by RT-qPCR and immunoblotting
(Figure 2D, top). Interestingly, Dot1L overexpression alone increased UCP1 mRNA levels over
2-fold, probably due to the presence of endogenous Dot1L in 3T3-L1 cells. Co-overexpression
of both Dot1L and Zc3h10 resulted in a further increase in UCP1 mRNA levels by 3-fold.
(Figure 2D, bottom). Furthermore, overexpression of Dot1L and Zc3h10 together in
differentiated 3T3- L1 cells further increased mRNA levels for Tfam and Nrf1 in comparison to
Dot1L or Zc3h10 alone. Expression of a terminal adipose gene, FABP4, was not significantly
different in all conditions (Figure 2D, bottom). With increased expression of UCP1 and other
BAT-enriched genes upon overexpression of Dot1L and Zc3h10 (Figure 2D and 2E), we next
assessed the functional consequence of the oxygen consumption. Indeed, forskolin-treated
Dot1L alone overexpressing cells had somewhat increased OCR. Co-overexpression of both
Dot1L and Zc3h10 further increased the total OCR as well as uncoupled OCR, suggesting
cooperative function of Dot1L and Zc3h10 in browning of 3T3-L1 cells (Figure 2F). Altogether,
these results demonstrate that Dot1L plays a critical role in promoting thermogenic program.
212

Figure 2. Dot1L promotes thermogenic gene program in vitro

213
Figure 2. Dot1L promotes thermogenic gene program in vitro
(A) (Left) RT-qPCR for Dot1L and thermogenic genes in BAT cells infected either scrambled
(Control) or adenovirus expressing short hairpin targeting Dot1L (Dot1L KD) after Day 2 of
adipogenic differentiation. (Right) Immunoblotting for UCP1.
(B) (Left) OCR measured in Dot1L KD cells using Seahorse XF24 analyzer. (Right) Uncoupled
OCR in BAT cells infected with control or shDot1L under oligomycin (0.5uM).
(C) (Left) RT-qPCR for indicated genes in SGC0496 treated BAT cells. BAT cells treated with
either DMSO or Dot1L inhibitor EPZ5676 (5nM). (Right) Immunoblotting for UCP1.
(D) RT-qPCR for indicated genes and immunoblotting for indicated proteins in 3T3-L1 cells that
were transduced with either AdGFP or AdZc3h10 or AdDot1L individually or in combination for

overexpression (OE) of Zc3h10 and Dot1L. The differentiated cells were treated with forskolin
(10uM) for 6 hr to induce beiging.
(E) Immunoblotting for UCP1.
(F) (Left) OCR measured in Zc3h10 OE and Dot1L OE, individually and in combination in
differentiated 3T3-L1 adipocytes that were induced to beiging. (Right) Uncoupled OCR under
oligomycin (0.5 uM).
230
Inhibition of Dot1L activity impairs BAT gene program in vivo
To study Dot1L function in vivo, we first used a selective and potent small molecule
inhibitor of Dot1L, EPZ5676. We administered either saline (control) or EPZ5676 to WT mice via
daily subcutaneous injection for 8 days (Figure 3A). By immunoblotting with H3K79me3
antibody, we confirmed inhibition of H3K79 methylation in BAT of animals treated with EPZ5676
(Figure 3B, left). Indeed, BAT from the inhibitor treated mice showed significantly decreased
mRNA levels for UCP1, Nrf1, and Tfam, as well as other thermogenic genes, such as PGC1α
and Elovl3, while no changes in adipogenic transcription factor, such as PPARγ (Figure 3B,
middle). Also, the immunoblotting detected significantly lower UCP1 protein levels in BAT of
EPZ5676 treated mice compared to the control mice (Figure 3B, middle). Similarly, EPZ5676
treatment also decreased thermogenic gene expression in iWAT, WAT depot known to undergo
browning (Figure 3B, right), but not in pWAT or in liver (Supplementary Figure 2A).
Next, to investigate the physiological outcome from decreased expression of
thermogenic genes, we subjected these EPZ5676 treated mice to acute cold exposure at 4°C.
Both groups of mice had body temperatures around 37°C prior to the cold exposure. After 4 hrs
of cold exposure, the body temperature of the EPZ5676 treated mice was 4°C lower compared
to their non-treated littermates, demonstrating the significantly reduced thermogenic capacity
(Figure 3C). Next, we assessed the energy expenditure by measuring the whole body O_2
consumption using CLAMS. Indeed, EPZ5676 treated mice had significantly lower VO_2 than the
control mice at 4°C (Figure 3D), whereas food intake and locomotive activity were similar
between the two groups (Supplementary Figure 2B). Altogether, these results support that
Dot1L enzymatic activity is critical for activating the thermogenic gene program in vivo.
253

Figure 3. Inhibition of Dot1L activity impairs BAT gene program in vivo

Figure 3. Inhibition of Dot1L activity impairs BAT gene program in vivo

(A) Schematic diagram of the strategy used to inject either Saline for control or Dot1L chemical inhibitor, EPZ5676.

(B) (Left) Immunoblotting for H3K79me3 and total H3. (Middle) RT-qPCR for indicated genes in BAT and iWAT from either PBS or EPZ5676 injected mice (n=4 mice per group) and immunoblotting for UCP1 protein.

(C) Rectal temperature of 14-wk old mice maintained at 4°C at indicated time points (hr) (n=4 mice per group)

(D) VO$_2$ assayed in mice that were housed at 4°C by indirect calorimetry using CLAMS.

Dot1L is required for cold-induced thermogenesis in mice

To further evaluate whether Dot1L is critical for the BAT gene program in vivo, we performed Dot1L ablation in UCP1$^+$ cells in mice (Dot1L-BKO) by crossing Dot1L floxed mice with UCP1-Cre mice (Figure 4A, left). We validated our mouse model by genotyping and by comparing Dot1L mRNA levels in BAT and iWAT of Dot1L-BKO mice and Dot1L f/f control mice (WT). Dot1L-BKO mice had a 70% reduction in Dot1L mRNA level in BAT. Dot1L-BKO mice also showed a 60% decrease in Dot1L mRNA level in iWAT, probably due to the mild cold exposure at room temperature that can induce UCP1$^+$ adipocytes in iWAT (Figure 4A, right). Importantly, UCP1 expression was significantly reduced by 40% and 60% in BAT and iWAT, respectively (Figure 4B). UCP1 protein level was reduced significantly, as detect in BAT (Figure 4B right). In addition, expression of other BAT enriched genes, such as Dio2 and CideA, and other Zc3h10 target genes, such as Tfam and Nrf1, were all significantly decreased in BAT and iWAT of Dot1L-BKO mice, compared to WT littermates (Figure 4C).

Dot1L is known to be the only known H3K79 methyltransferase (VanLeeuwen, 2002). Hence, we performed ChIP-qPCR, to assess methylation status of H3K79 methylation at the Zc3h10 binding region of the UCP1 promoter using BAT of mice. Indeed, we detected a 50% reduction in di-methylation and a 70% reduction in tri-methylation of H3K79 at the -4.6 kb UCP1 promoter region (Figure 4D). Thus, the decreased UCP1 mRNA level upon Dot1L ablation in BAT was correlated with decreased di-methylation and tri-methylation H3K79 at the UCP1 promoter region. Similar to the -4.6 kb UCP1 promoter region, H3K79 methylation status at the -200 bp Nrf1 promoter region and at the +50 bp Tfam region was also decreased (Figure 4D). We previous reported these regions to contain Zc3h10 binding sites. The reduced H3K79 methylation at Zc3h10 binding sites in BAT of Dot1L-BKO mice suggest that the recruitment of Dot1L by Zc3h10 to the target genes is important for BAT gene expression. Moreover, these observations support the notion that Dot1L participates in the transcriptional activation of UCP1 and other Zc3h0 target genes by modifying H3K9 methylation status.

To examine the effect of Dot1L deficiency on the thermogenic capacity, we subjected these Dot1L-BKO mice to an acute cold exposure at 4°C. While body temperatures of mice were similar initially, Dot1L-BKO mice were severely cold intolerant as their body temperature started to drop significantly after 2 hrs of cold challenge. After 3 hrs of cold exposure, the body temperature of Dot1L-BKO mice was 5°C lower than that of control mice (Figure 4E), an in vivo evidence of the requirement of Dot1L for cold-induced thermogenesis. Considering such lower thermogenic gene expression and decreased thermogenic capacity, we assessed the metabolic effect in the Dot1L-BKO mice by measuring whole body OCR using CLAMS. Indeed, the Dot1L-BKO mice had significantly reduced VO$_2$ compared to WT littermates at room temperature and at 4°C, while locomotor activity and food consumption were similar (Figure 4F and Supplementary Figure 3A). To examine the contribution of BAT to the altered energy expenditure in Dot1L-BKO mice, by Seahorse assay, we measured OCR in BAT dissected out from these mice. We found OCR in BAT from Dot1L-BKO mice to be decreased by 20%. (Figure 4F, right). In line with these results, we then asked whether decreased energy expenditure and impaired BAT function in Dot1L-BKO mice would be reflected in changes in

adiposity. The body weights of mice started to diverge starting on wk 10, then they were maintained on a high-fat-diet (HFD) from wk 11 to wk 13 of age. By 13 wks, the Dot1L-BKO mice were approximately 8 g heavier relative to the control mice (Figure 4G), and the Dot1L-BKO had significantly larger iWAT and pWAT depots, which primarily accounted for the higher total body weights of Dot1L-BKO mice (Supplementary Figure 3B). There were no differences in food/energy intake between the two genotypes despite differences in body weights. Collectively, these results establish a defective BAT function and impaired thermogenic capacity that affects adiposity in Dot1L-BKO mice and that Dot1L is required for the BAT thermogenic program in vivo.

Figure 4. Dot1L is required for cold-induced thermogenesis in mice

Figure 4. Dot1L is required for cold-induced thermogenesis in mice

 (A) (Left) Schematic diagram of the strategy used to generate BAT specific Dot1L conditional KO mice. PCR genotyping of the mice: Top gel, Dot1L allele; bottom gel, Cre. (Right) RT-qPCR for Dot1L in BAT, iWAT and pWAT from Dot1L f/f (WT) and Dot1L-BKO mice (n=6 mice per group).

(B) RT-qPCR and immunoblotting for UCP1 in BAT and iWAT from Dot1L f/f (WT) and Dot1L-BKO mice.

(C) RT-qPCR for indicated genes in BAT and iWAT from Dot1L f/f (WT) and Dot1L-BKO mice.

(D) ChIP-qPCR of H3K79me2 and H3K79me3 at Zc3h10 binding regions of BAT tissue from control (f/f) or Dot1L-BKO mice (n=5).

(E) Rectal temperature measured in 13 wk-old mice at 4°C at indicated time points (hours) (n=6 mice per group).

(F) (Left) whole body VO_2 assayed in WT and Dot1L-BKO mice, housed at indicated ambient temperatures (n=6 per group) by indirect calorimetry using CLAMS. (Right) OCR measured in BAT of WT and Dot1L-BKO mice using Seahorse XF24 Analyzer (n=5 per group).

(G) Representative photograph of 14 wk-old control and body weight of WT and Dot1L-BKO mice fed HFD from wk 12.

(H) RT-qPCR for indicated genes from WT injected with GFP or Zc3h10 adenovirus or Dot1L-BKO mice injected with Zc3h10 adenovirus for overexpression.

(I) Rectal temperature measured at 4°C at indicated time points (hours) (n=4 mice per group).

To test the relevance of Dot1L-Zc3h10 interaction to thermogenesis in vivo, we next asked whether Dot1L is required for Zc3h10-induced thermogenesis. By direct injection of Zc3h10 adenovirus into BAT, we overexpressed Zc3h10 in WT and in Dot1L-BKO mice. We detected a 4-fold increase in Zc3h10 mRNA levels in BAT of Zc3h10 adenovirus-injected mice (Zc3h10 OE). As expected, Dot1L-BKO mice showed a 60% reduction in Dot1L mRNA levels in BAT (Figure 4H). We previously have reported that overexpression Zc3h10 in adipose tissue enhances thermogenic gene expression (Yi et al., 2019). As predicted Zc3h10 overexpressing mice had increased expression of BAT-enriched genes, such as UCP1, Cox8b, and CideA, compared to the control mice. More importantly, expression BAT-enriched genes was significantly reduced in Dot1L-BKO even upon Zc3h10 overexpression, while adipogenic markers, such as PPARγ, remained unchanged (Figure 4H). These results show that Dot1L is critical and required for activation of the thermogenic program by Zc3h10 in vivo. Moreover, when these mice were subjected to cold exposure, Zc3h10 OE mice had higher core body temperature compared to the control mice after 3 hrs (Figure 4I). However, Dot1L-BKO Zc3h10 OE mice were severely cold sensitive, and their body temperature was significantly lower than the control and Zc3h10 OE mice. Collectively, these results demonstrate the requirement of Dot1L-Zc3h10 interaction for activating the thermogenic program in brown fat.

Dot1L increases chromatin accessibility for activation of thermogenic genes

Thus far, we have shown that Dot1L-BKO mice have severely reduced thermogenic capacity due to significantly decreased BAT gene expression accompanied with reduced H3K79me2/3 at promoter regions of thermogenic genes. To study the underlying mechanism of how Dot1L ablation in BAT decreases H3K79me2/3 to decrease chromatin accessibility, thereby decreasing gene transcription, we used nuclei isolated from cold exposed Dot1L-BKO mice and their littermates to assess the chromatin landscape during thermogenesis by Assay for Transposase-Accessible Chromatin (ATAC-seq). Strikingly, Dot1L-BKO had significantly reduced peaks genome wide near the transcription start sites (TSS), while WT had concentrated open chromatin regions at TSS, as H3K79 methylation is reported as a gene activation marker. (Fig 5A left). With our primary interest in assessing Dot1L's role in the regulation of thermogenesis, we compared relative ATAC-seq peaks for thermogenic genes,

such as PRDM16, Zfp516, Cox8b and for mitochondria/oxidative phosphorylation, such as
Cox7a1, Tfam, and Cyc1, in response to Dot1L ablation (Figure 5, middle). Also, Aifm2, Dot1L-
BKO mice had much smaller peaks around promoters of UCP1, Cox8b and Elovl3 compared to
WT mice (Figure 4, right), consistent with significantly decreased thermogenic gene expression
in the Dot1L-BKO mice. Also, we found Dot1L-BKO had decreased expression of mitochondrial
genes as well as Aifm2, a recently reported BAT-specific enzyme that converts NADH to NAD to
sustain robust glycolysis to fuel thermogenesis (Nguyen et al., 2020). These data suggest that
H3K79me2/3 by Dot1L is important for creating a chromatin landscape permissive to
transcription in brown fat upon cold-stimulation.
376

Figure 5. Dot1L increases chromatin accessibility for activation of thermogenic genes

Figure 5. Dot1L increases chromatin accessibility for activation of thermogenic genes
(A) ATAC-seq using BAT from WT and Dot1L-BKO mice after 2 hr cold exposure n = 2 per
group. (Left) Heatmaps showing open chromatin regions focused at the transcription start site
(TSS). Color scale shows peaks detected. (Right) UCSC genome browser screenshot of
representative peaks at a subset of promoter regions of thermogenic genes.
(B) RNA-seq using BAT from WT and Dot1L-BKO mice, n = 2 pooled RNA samples per group.
(Left) Heatmap showing changes in gene expression. Color scale shows changes in gene
expression as determined by Z-score, green is –2 and red is 2. (Middle) Representative top GO
terms of upregulated and downregulated genes identified by differential expression analysis.
(C) Venn diagrams showing number of unique or shared genes between ATAC- and RNA-seq
datasets and charts for representative top gene ontology (GO) terms.
389
Next, to examine gene expression changes upon Dot1L ablation in UCP1$^+$ cells as a
result of chromatin compaction, we performed RNA-seq using BAT from Dot1L-BKO mice and
their littermates. Gene ontology of genes that were at least 2-fold downregulated in Dot1L-BKO
mice revealed that mitochondrion organization, oxidative phosphorylation and cold-induced
thermogenesis were some of the most affected processes (Figure 5B, middle). Notably, the
downregulated thermogenic genes include UCP1, Cox8b, Elov3 and PGC1α as well as those
related to mitochondrion organization and oxidative phosphorylation such as Nrf1, Cox20,
Tomm13, and Immp21 (Figure 5B, left). With ATAC-seq results, these data suggest a
mechanistic link between Dot1L-mediated chromatin remodeling and the global landscape of
genome accessibility in brown fat for thermogenic gene expression. In addition, regulation of
triglyceride and carbohydrate catabolism were compromised by Dot1L ablation. Conversely,
chromosome organization, negative regulation of telomere maintenance, response to DNA
damage were upregulated in the Dot1L knockout. In fact, we detected upregulated genes
involved in chromatin compaction and gene repression, such as HDAC3, HDAC4 and DNMT1,
further supporting that Dot1L increases chromatin accessibility (Figure 5B, right).
We then compared RNA-seq data to ATAC-seq obtained to identify common
downregulated pathways found at lower levels using ATACs upon Dot1L ablation in UCP1+
cells, and we found about 40% of the total of downregulated genes were also detected from the
ATAC-seq dataset (Figure 5C, left). Some of similar processes between the two datasets
include regulation of cold-induced thermogenesis, transcription initiation/elongation factors,
mitochondrial membrane organization as well as G-protein receptor signaling pathway (Figure
5C, right). Remarkably, the cluster of shared genes included thermogenic genes, such
as UCP1, PGC1α, Cox8b, CideA, further underscoring the importance of Dot1L in brown
adipose tissue. This suggest that Dot1L ablation results in reduced chromatin accessibility,
leading to downregulated gene expression for those specific pathways. Altogether, we conclude
that H3K79me2/3 by Dot1L plays a critical role in the activation of transcription of thermogenic
genes by increasing chromatin accessibility.
417
418   **DISCUSSION**
419
Numerous studies have supported the concept that a network of transcription factors
and epigenetic coregulators must work together to fine-tune thermogenic program in response
to environmental conditions (Sambeat et al., 2017). Here, we identify Dot1L, the only known
H3K79 methyltransferase, as an interacting partner of Zc3h10, a transcriptional factor that
activates UCP1, as well as Tfam and Nrf1, for thermogenesis. We show that Dot1L is recruited
by Zc3h10 to the UCP1 and other thermogenic genes. We demonstrate that Dot1L directly
interacts with Zc3h10 to co-occupy the same region of the UCP1 promoter. Zc3h10 binds to
Dot1L via the 501-1000aa fragment, which is largely composed of coiled coil domains,

evolutionarily conserved and widely-known for protein-protein interactions (Mier et al., 2017).
This same coiled coil domain has been reported to interact with some of mixed lineage leukemia
(MLL) oncogenic fusion proteins, such as AF10, to induce H3K79 methylation and constitutively
activate a leukemic transcriptional program (Song et al., 2019). Here, we show that Dot1L is
required for Zc3h10's function for thermogenic gene program. In fact, Zc3h10 overexpression
could not rescue the decreased thermogenic gene expression upon Dot1L ablation in mice,
demonstrating that Dot1L is critical in Zc3h10 mediated activation of thermogenesis.
We establish that H3K79me2/3, catalyzed by Dot1L, as an activation mark for
thermogenic gene program. In this regard, previous studies reported that H3K79 methylation is
strongly associated with active transcription (Steger et al., 2008, Wood et al., 2018). Our Dot1L-
BKO mice showed a markedly less open chromatin regions in promoter regions of thermogenic
genes. The ATAC-seq data showed significantly reduced peaks genome-wide near the
transcription start sites (TSS) in BAT of Dot1L-BKO mice, compared to control mice that had
concentrated open chromatin regions at TSS upon cold exposure. In addition, studies reported
that H3K79me2/3 may be active enhancer marks and may even be required for enhancer-
promoter interactions to increase transcription (Markenscoff-Papadimitriou et al., 2014, Gilan et
al., 2016, Godfrey et al., 2019). Godfrey et al. found H3K79me2/3 to be abundant at a subset of
super-enhancers in MLL-related leukemia cells and that a loss of H3K79me2/3 by Dot1L
inhibition led to significantly reduced enhancer-promoter interaction coupled with a decrease in
transcription (Godfrey et al., 2019). We show that Dot1L is recruited by Zc3h10 to a UCP1
upstream region of -4.6 kb, a region suggested to be one of so-called super enhancers (SEs)
(Whyte et al., 2013; Harms et al., 2015). In fact, Dot1L ablation in BAT of mice resulted in a
significantly reduction of H3K79me2/3 at the -4.6 kb UCP1 region, correlating with decreased
expression of UCP1. It is possible that H3K79me2/3 catalyzed by Dot1L may be important in
maintaining enhancer/promoter association for transcriptional activation.
Due to limited accessibility of the H3K79 residue, located in the nucleosome core, how
H3K79 methylation promotes gene activation is not clearly understood (van Leeuwen et al.,
2002, Lu et al., 2008). Several studies support the concept that H3K79me2/3 and other well-
profiled gene activation marks are interdependent (Steger et al., 2008). Steger et al.
demonstrated that H3K79me2/3 at gene promoters highly correlates with H3K4me1/2/3 and
H3K36me3, known gene activation marks in mammalian cells. Furthermore, Chen et al.
reported that Dot1L may inhibit the localization of SIRT1 and SUV39H1, a H3K9ac demethylase
and a H3K9me2 methyltransferase, respectively, at MLL fusion target genes correlating with
elevated H3K9ac and low H3K9me2, to maintain an open chromatin state (Chen et al., 2015).
These authors proposed that H3K79me2/3 mark, in part, may function by inhibiting the histone
deacetylase activity (Chen et al., 2015, Kang et al., 2018). In line with these findings, Dot1L
probably works with other histone modifiers to maintain activation marks or to suppress
repressive marks. Regardless, we establish that H3K79me2 and H3K79me3, in particular,
function as activation marks for thermogenic gene program.
Overall, we demonstrate that Dot1L is critical for thermogenic program and is required
for Zc3h10-induced thermogenesis through its H3K79 methyltransferase activity. With combined
ATAC-seq and RNA-seq data, we provide molecular evidence that the decreased thermogenic
gene expression is due to decreased H3K79me2/3 on promoter regions of thermogenic genes,
as well as genes involved in chromatin remodeling. Consequently, our mouse line with Dot1L
ablated in UCP1[+] cells had significantly decreased thermogenic gene expression including
UCP1, resulting in cold intolerance with decreased oxygen consumption (Figure 4C-E), leading
to increased adiposity.
475
476 **MATERIALS AND METHODS**
477
Animals

Dot1L conditional knock-out mice (Dot1ltm1a(KOMP)Wtsi) for this research project were
generated by the trans-NIH Knock-Out Mouse Project (KOMP) and obtained from the KOMP
Repository (www.komp.org). NIH grants to Velocigene at Regeneron Inc (U01HG004085) and
the CSD Consortium (U01HG004080) funded the generation of gene-targeted ES cells for 8500
genes in the KOMP Program and archived and distributed by the KOMP Repository at UC Davis
and CHORI (U42RR024244). Mice were first mated with FLPe mice from Jackson Lab. FLP
mediated recombination was confirmed by PCR and resultant progeny were mated with UCP1-
cre mice (B6.FVB-Tg(Ucp1-cre)1Evdr/J) also from Jackson Lab.Unless otherwise stated, mice
used in experiments were male and between 10 -14 weeks of age. Mice were fed a chow diet or
a palatable high fat diet (45% fat derived calories- Dyets) ad libitum. EPZ-5676 was purchased
from MedChemExpress (Cat. No.: HY-15593) and injected 20mg/kg/day for 8 days into BAT. All
protocols for mice studies were approved from the University of California at Berkeley Animal
Care and Use Committee.
492

GST-pulldown

493
GST-Dot1L was purchased from epicypher. GST and GST-Dot1L plasmids were
transformed into BL21(DE3) E. coli (NEB) and production was induced by induction by 0.1M
IPTG. Resultant proteins were purified using Glutathione Sepharose 4B (GE) according to the
manufacturer recommended protocol. [35S]-labeled Zc3h10 protein was produced by using TNT
coupled transcription/translation kit (Promega). 20ug of GST fusion proteins were incubated for
2hrs at 4°C with in vitro translated Zc3h10 and glutathione sepharose beads. The beads were
washed 3 times with binding buffer, and bound proteins were eluted by boiling in Laemmli
sample buffer, separated by SDS-PAGE and analyzed by autoradiography.
502

TAP assay and mass spectrometry Analysis

503
10ug of Zc3h10-CTAP or CTAP vector plasmid were transfected in 293FT cells using
lipofectamine 2000 (Invitrogen). Cells were lysed and IPed using buffers from the InterPlay
Mammalian TAP system manufacturer recommended with the noted exception: after binding of
the cell lysates to the Streptavidin resin, the resin was re-incubated with 200ug of BAT nuclear
extracts diluted to 1mL in SBB overnight. TAP eluates were boiled in SDS, ran on an SDS-
PAGE gel, and stained with Coomassie Brilliant Blue (Sigma). Bands that were identified to be
specific to the Zc3h10-TAP lane were excised from both the Zc3h10-TAP and TAP vector lanes
and protein were identified by mass spectrometry performed by the Vincent J. Coates
Proteomics/ Mass Spectrometry Laboratory (P/MSL) at UC Berkeley.
513

Cold-induced Thermogenesis

514
515   Core body temperature was determined using a Physitemp BAT-12 probe at 4°C.
516

Indirect Calorimetry

517
518   Oxygen Consumption was measured using the Comprehensive Laboratory Animal
519   Monitoring System (CLAMS). Data were normalized to lean body mass determined by
520   EchoMRI. Mice were individually caged and maintained under a 12 hr light/12 hr dark cycle.
521   Food consumption and locomotor activity were tracked.
522

Cell culture

523
HEK293FT and 3T3-L1 cell lines were obtained from UCB Cell Culture Facility which is
supported by The University of California Berkeley. The immortalized BAT cell line was from Dr.
Shingo Kajimura (Harvard). Cells were grown in standard condition with 5% CO2, at 37°C. BAT
cells, 3T3-L1 and 293FT cells were maintained in DMEM containing 10% FBS and 1%
pen/strep prior to differentiation/ transfection. Brown adipocyte differentiation was performed as
described in (Yi et al., 2019). Ad-m-Dot1L-shRNA, Ad-Zc3h10-flag adenovirus and Ad-GFP-m-

DOT1L were purchased from Vector Biolabs. Knockdown of Dot1L accomplished using
adenoviral transduction of MOI of 250 at day 4 of brown adipocyte differentiation. Differentiation
of 3T3-L1 was induced by treating confluent cells with DMEM containing 10% FBS, 850nM
insulin, 0.5mM isobutyl-methylxanthine, 1μM dexamethasone, 1nM T3, 125 nM indomethacin.
After 48h of induction, cells were switched to a maintenance medium containing 10% FBS,
850nM insulin, 1nM T3. 3T3-L1 cells were infected on day 4 of differentiation using either GFP,
Dot1L or Zc3h10 adenoviruses. Then, viral medium was replaced by maintenance medium the
following day. To stimulate thermogenesis, differentiated 3T3-L1 cells were treated 6h with
10μM forskolin on day 6. Inhibition of Dot1l was accomplished by addition of either 5nM
EPZ5676 (MedChemExpress, Cat. No.: HY-15593) or DMSO.
540
Seahorse oxygen consumption assay
BAT cell line cells were differentiated in 12-well plates, trysinized, and reseeded in XF24
plates at 50K cells per well at day 4 of differentiation and assayed on day 5 of differentiation. On
the day of experiments, the cells were washed three times and maintained in XF-DMEM
(Sigma-Aldrich) supplemented with 1 mM sodium pyruvate and 17.5 mM glucose. Oxygen
consumption was blocked by 1 μM oligomycin. Maximal respiratory capacity was assayed by
the addition of 1 μM FCCP. Tissues were incubated for 1 hour at 37°C without CO_2 prior to
analysis on the XF24 Analyzer. Uncoupled respiration was calculated as OCR under oligomycin
treatment minus OCR under antimycin A/rotenone.
550
RT-qPCR Analysis and Western blotting
Reverse transcription was performed with 500ng of total RNA using SuperScript III
(Invitrogen). RT-qPCR was performed in triplicate with an ABI PRISM 7500 sequence detection
system (Applied Biosystems) to quantify the relative mRNA levels for various genes. Statistical
analysis of the qPCR was obtained using the ΔΔCt (2^-ddCT) method with Eef1a1 or 18s as the
control. RT-qPCR primer sets are listed in Table 1. Library generation for RNA sequencing was
carried out at the Functional Genomics Laboratory at UC Berkeley.
For western blot analysis, total cell lysates were prepared using RIPA buffer and nuclear
extracts were isolated using the NE-PER Nuclear and Cytoplasmic Extraction kit (Thermo).
Proteins were separated by SDS-PAGE, transferred to nitrocellulose membrane and probed
with the indicated antibodies.
562
Antibodies
The following antibodies were used where indicated: UCP1 (Abcam, 10983), Zc3h10
(Thermo Fisher, PA5-31814; for immunoblotting), Zc3h10 (Abnova, H00084872-B01P; for
ChIP), GAPDH (Cell signaling, 5174S), Dot1L (Bethyl, A300-953A), FLAG (Cell signaling,
14793S), HA (Cell signaling, 3724S), Tubulin (Abcam, Ab52866), Histone H3 (Cell signaling,
4499S), H3K79me3 (Abcam, ab2621), H3K79me2 (Abcam, ab3594)
569
570   **Luciferase-reporter Assay**
571   293FT cells were transfected with 300ng Zc3h10 and/or Dot1L, together with 100 ng of
572   indicated luciferase reporter construct and 0.5ng pRL-CMV in 24-well plates. Cells were lysed
573   48h post-transfection and assayed for luciferase activity using the Dual-Luciferase Kit
574   (Promega) according to manufacturer recommended protocol.
575
576   **Plasmid Constructs**
577   The HA-Dot1L expression vector was purchased from Genecopeiea. The Dot1L
578   sequence was subcloned by PCR amplifying and inserting into CTAP vectors from Agilent as
579   well as pGEX-4T-3 from GE.
580

Coimmunoprecipitation

For Co-IP experiments using tagged constructs, 293FT cells were transfected using Lipofectamine2000 to express FLAG-tagged Zc3h10 and HA-tagged Dot1L. Cells were lysed in IP buffer containing 20 mM Tris, pH 7.4, 150 mM NaCl, 1mM EDTA, 10% glycerol, 1% NP-40 supplemented with proteases inhibitors. Total cell lysates were incubated 2 hours at 4°C with either anti-FLAG M2. For Co-IP in brown adipose tissue of wild type mice, nuclear extraction was carried out using the NE-PER Nuclear and Cytoplasmic Extraction kit (Thermo). Equal amounts of nuclear extracts were incubated with specific antibodies and protein A/G agarose beads overnight at 4°C. Agarose beads were washed 3 times and bound proteins were eluted by boiling in Laemmli sample buffer and analysed by immunoblotting using the indicated antibodies.

ChIP-qPCR

ChIP was performed using ChIP kit (Cell signaling, 57976s). Briefly, BAT cells overexpressing Dot1L or Zc3h10 were fixed with disuccinimidyl glutarate (DSG) at 2 mM concentration in PBS for 45 min at room temperature before cross-linking with 1% methanol-free formaldehyde in PBS for 10 min. For ChIP experiments in BAT, tissues from 3 BATs were combined and minced on ice. The reaction was stopped by incubating with 125 mM glycine for 10 min. Cells or tissues were rinsed with ice-cold phosphate-buffered saline (PBS) for three times, and lysed in IP lysis buffer containing 500 mM HEPES-KOH, pH 8, 1 mM EDTA, 0.5 mM EGTA, 140 mM NaCl, 0.5% NP-40, 0.25% Triton X-100, 10% glycerol, and protease inhibitors for 10 min at 4 °C. Nuclei were collected by centrifugation at 600 × g for 5 min at 4 °C. Nuclei were released by douncing on ice and collected by centrifugation. Nuclei were then lysed in nuclei lysis buffer containing 50 mM Tris, pH 8.0, 1% SDS 10 mM EDTA supplemented with protease inhibitors, and sonicated three times by 20 s burst, each followed by 1 min cooling on ice. Chromatin samples were diluted 1:10 with the dilution buffer containing 16.7 mM Tris pH 8.1, 0.01% SDS 1.1 % Triton X-100 1.2 mM EDTA, 1.67 mM NaCl, and proteinase inhibitor cocktail. Soluble chromatin was quantified by absorbance at 260 nm, and equivalent amounts of input DNA were immunoprecipitated using 10 μg of indicated antibodies or normal mouse IgG (Santa Cruz) and protein A/G magnetic beads (Thermo-Fisher). After the beads were washed and cross-linking reversed, DNA fragments were purified using the Simple ChIP kit (cell signaling). Samples were analyzed by qPCR using the primer sets in Table 1 for enrichment in target areas. The promoter occupancy for UCP1 and other indicated genes was confirmed by RT-qPCR. The fold enrichment values were normalized to input.

RNA-seq

WT and Dot1L-BKO mice were cold exposed for 2 hr. Total RNA from BAT was prepared using RNEasy kit (Qiagen). Strand-specific libraries were generated from 500 ng total RNA using the TruSeq Stranded Total RNA Library Prep Kit (Illumina). cDNA libraries were single-end sequenced (50 bp) on an Illumina HiSeq 2000 or 4000. Using Partek software, reads were aligned to the mouse genome (NCBI38/mm10). A gene was included in the analysis if it met all the following criteria: the maximum RPKM reached four at any time point, the gene length was >100 bp. Selected genes were induced at least +/- 1.8-fold, and the expression was significantly different from the basal (p<0.05).

ATAC-seq

For ATAC-seq, we used two mice per condition and processed nuclei extracted from BAT separately, thus each sample analyzed was a separate biological replicate. ATAC-seq was performed using Nextera DNA library Preparation kit (Illumina, 15028212) according to Chen et al. 50. In all, 5×10^4 nuclei from tissues were collected in lysis buffer containing 10 mM Tris pH 7.4, 10 mM NaCl, 3 mM MgCl2, and 1% NP-40, and spun at 500 × g at 4 °C for 10 min. The

pellets were resuspended in the transposase reaction mixture containing 25 µL
2 × Tagmentation buffer, 2.5 µL transposase, and 22.5 µL nuclease-free water, and incubated at
37 °C for 30 min. The samples were purified using MinElute PCR Purification kit (Qiagen,
28006) and amplification was performed in 1 × next PCR master mix (NEB, M0541S) and
1.25 µM of custom Nextera PCR primers 1 and 2 with the following PCR conditions: 72 °C for
5 min; 98 °C for 30 s; and thermocycling at 98 °C for 10 s, 63 °C for 30 s, and 72 °C for 1 min.
Samples were amplified for five cycles and 5 µL of the PCR reaction was used to determine the
required cycles of amplification by qPCR. The remaining 45 µL reaction was amplified with the
determined cycles and purified with MinElute PCR Purification kit (Qiagen, 28006) yielding a
final library concentration of ~30 nM in 20 µL. Libraries were subjected to pair-end 50 bp
sequencing on HiSeq4000 with 4–6 indexed libraries per lane. Partek Flow Genomics Suite was
used to analyze sequencing data. Reads were aligned to mm10 genome assembly using BWA-
backtrack. MACS2 in ATAC mode was then used to identify peaks from the aligned reads with a
q-value cutoff of 0.05 and fold enrichment cutoff of 2.0. Quantify regions tool was used to
quantify the peaks identified by MACS2 from each sample to generate a union set of regions.
Regions were then annotated to RefSeq mRNA database and analysis of variance (ANOVA)
was performed to determine significant peak differences between groups. UCSC genome
browser was used to visualize peaks. Gene set enrichment and pathway analysis was used to
highlight biological processes with the highest q-values among those identified.
651
Statistical analysis
Statistical comparisons were made using a two-tailed unpaired t-test using GraphPad
Prism 8 software (GraphPad Software Inc., La Jolla, CA, USA). For genome-wide analyses, we
employed Partek Genomics Suite (Partek Inc., St. Louis, Missouri, USA) using ANOVA for
ATAC-seq comparisons, and DESeq2 for RNA-seq differential expression comparisons, and
subsequently used Gene set enrichment tool for gene ontology to identify the significantly
affected pathways presented in Data are expressed as means ± standard errors of the means
(SEM). The statistical differences in mean values were assessed by Student's *t* test. All
experiments were performed at least twice and representative data are shown.
661

662 <div align="center">**AUTHOR CONTRIBUTIONS**</div>
663
D.Y. performed loss-of-function/gain-of-function studies in vitro/in vivo experiments.
H.P.N. performed the interaction studies and assisted in animal studies. J.D. assisted in animal
studies. J.V. made plasmid constructs and performed bioinformatics analysis and edited the
manuscript. Y.X. assisted in RT-qPCR/immunoblotting. J.M.D. found and characterized Dot1L.
Y.W. performed ChIP and ATAC-seq. H.S.S. designed the project and guided experiments.
D.Y., H.P.N., and H.S.S wrote the manuscript.
670

671 <div align="center">**ACKLOWLDGEMENTS**</div>
672
673   We thank M. Zhu for technical assistance. The work was supported by NIH grant
674 DK120075 to H.S.S.
675

676 <div align="center">**COMPETING INTERESTS**</div>
677
678   The authors declare no competing interests.
679

680 <div align="center">**SUPPLEMTARY FIGURES AND TABLES**</div>
681

S1. Dot1L directly interacts with Zc3h10 for UCP1 promoter activation

Candidate proteins identified by TAP-Mass spec:
-Dot1L
-Phb
-Smarca4
-Smarca5
-Snw1
-Fus

CBB stained TAP

(A) Gel stained with Coomassie Blue after cTAP purification assay and a list of potential Zc3h10 interacting proteins identified by Mass spec

(B) Coomassie staining for GST alone and GST-Dot1L fusion protein

S2. Inhibition of Dot1L activity impairs BAT gene program in vivo

(A) RT-qPCR for indicated genes in pWAT and liver of control and EPZ5676 injected mice.
(B) Food intake and locomotor activity of control and EPZ5676 injected mice.

S3.Dot1L is required for cold-induced thermogenesis in mice

(A) Food intake and locomotor activity of WT and Dot1L-BKO mice.
(B) Body weight and mass of various tissues taken from WT and Dot1L-BKO mice at 14-week of age.

684
685 **Table 1: Primer Sets used for RT-qPCR**

	Forward Primer	Reverse Primer
Dot1L	CGA CTA ATG CTG CAC CTC CT	AGG AGT AGT GGT GTG GCT CA
UCP1	ACT GCC ACA CCT CCA GTC ATT	CTT TGC CTC ACT CAG GAT TGG
Tfam	GTC CAT AGG CAC CGT ATT GC	CCC ATG CTG GAA AAA CAC TT
Zc3h10	CGA CTA ATG CTG CAC CTC CT	AGG AGT AGT GGT GTG GCT CA
Nrf1	GAC AAG ATC ATC AAC CTG CCT GTA G	GCT CAC TTC CTC CGG TCC TTT G
Cox8b	TGT GGG GAT CTC AGC CAT AGT	AGT GGG CTA AGA CCC ATC CTG
Cox7a1	CAG CGT CAT GGT CAG TCT GT	AGA AAA CCG TGT GGC AGA GA
Eef1a1	ACG AGG CAA TGT TGC TGG TGA C	GTG TGA CAA TCC AGA ACA GGA GC
Elovl3	TCC GCG TTC TCA TGT AGG TCT	GGA CCT GAT GCA ACC CTA TGA
PGC1α	CCC TGC CAT TGT TAA GAC C	TGC TGC TGT TCC TGT TTT C
PPARγ	GTG CCA GTT TCG ATC CGT AGA	GGC CAG CAT CGT GTA GAT GA
CideA	TGC TCT TCT GTA TCG CCC AGT	GCC GTG TTA AGG AAT CTG CTG
Dio2	CAG TGT GGT GCA CGT CTC CAA TC	TGA ACC AAA GTT GAC CAC CAG
FABP4	ACA CCG AGA TTT CCT TCA AAC TG	CCA TCT AGG GTT ATG ATG CTC TTC A
Pdgfrα	CAAACCCTGAGACCACAATG	TCCCCCAACAGTAACCCAAG
UCP1-4.6kb	CTC AGA GTG CAA CCC CTC AC	GAG GTC GCA GAT CTG TTC CA

Nrf1 -200	GGT CCA ATC AGA TAG CCG GG	CCT CAC TCA TTG GTG GGT CC
Tfam +50	CTC GGG CCG ACG AAT GAT G	CGG GCT TCC CAC AGT ACC C
Ins1 -400	TGA CCA ATG AGT GGG CTA CG	CCA TTG ATA GCT GGG CCC TT

686
687
688 **REFERENCES**
689

CANNON, B. & NEDERGAARD, J. 2004. Brown adipose tissue: function and physiological
significance. *Physiol Rev,* 84**,** 277-359.
CHEN, C. W., KOCHE, R. P., SINHA, A. U., DESHPANDE, A. J., ZHU, N., ENG, R., DOENCH, J. G., XU,
H., CHU, S. H., QI, J., WANG, X., DELANEY, C., BERNT, K. M., ROOT, D. E., HAHN, W. C.,
BRADNER, J. E. & ARMSTRONG, S. A. 2015. DOT1L inhibits SIRT1-mediated epigenetic
silencing to maintain leukemic gene expression in MLL-rearranged leukemia. *Nat Med,*
21**,** 335-43.
CHONDRONIKOLA, M., VOLPI, E., BORSHEIM, E., PORTER, C., ANNAMALAI, P., ENERBACK, S.,
LIDELL, M. E., SARAF, M. K., LABBE, S. M., HURREN, N. M., YFANTI, C., CHAO, T.,
ANDERSEN, C. R., CESANI, F., HAWKINS, H. & SIDOSSIS, L. S. 2014. Brown adipose tissue
improves whole-body glucose homeostasis and insulin sensitivity in humans. *Diabetes,*
63**,** 4089-99.
CYPESS, A. M., LEHMAN, S., WILLIAMS, G., TAL, I., RODMAN, D., GOLDFINE, A. B., KUO, F. C.,
PALMER, E. L., TSENG, Y., DORIA, A., KOLODNY, G. M. & KAHN, C. R. 2009. Identification
and Importance of Brown Adipose Tissue in Adult Humans. *New England Journal of*
Medicine, 360**,** 1509-1517.
DEMPERSMIER, J., SAMBEAT, A., GULYAEVA, O., PAUL, S. M., HUDAK, C. S., RAPOSO, H. F.,
KWAN, H. Y., KANG, C., WONG, R. H. & SUL, H. S. 2015. Cold-inducible Zfp516 activates
UCP1 transcription to promote browning of white fat and development of brown fat.
Mol Cell, 57**,** 235-46.
FARMER, S. R. 2008. Molecular determinants of brown adipocyte formation and function.
Genes Dev, 22**,** 1269-75.
FENG, Q., WANG, H., NG, H. H., ERDJUMENT-BROMAGE, H., TEMPST, P., STRUHL, K. & ZHANG,
Y. 2002. Methylation of H3-lysine 79 is mediated by a new family of HMTases without a
SET domain. *Curr Biol,* 12**,** 1052-8.
FREDERIKS, F., TZOUROS, M., OUDGENOEG, G., VAN WELSEM, T., FORNEROD, M., KRIJGSVELD,
J. & VAN LEEUWEN, F. 2008. Nonprocessive methylation by Dot1 leads to functional
redundancy of histone H3K79 methylation states. *Nat Struct Mol Biol,* 15**,** 550-7.
GILAN, O., LAM, E. Y., BECHER, I., LUGO, D., CANNIZZARO, E., JOBERTY, G., WARD, A., WIESE,
M., FONG, C. Y., FTOUNI, S., TYLER, D., STANLEY, K., MACPHERSON, L., WENG, C. F.,
CHAN, Y. C., GHISI, M., SMIL, D., CARPENTER, C., BROWN, P., GARTON, N., BLEWITT, M.
E., BANNISTER, A. J., KOUZARIDES, T., HUNTLY, B. J., JOHNSTONE, R. W., DREWES, G.,
DAWSON, S. J., ARROWSMITH, C. H., GRANDI, P., PRINJHA, R. K. & DAWSON, M. A. 2016.

723   Functional interdependence of BRD4 and DOT1L in MLL leukemia. *Nat Struct Mol Biol,*
724    23**,** 673-81.
725 GODFREY, L., CRUMP, N. T., THORNE, R., LAU, I. J., REPAPI, E., DIMOU, D., SMITH, A. L.,
726    HARMAN, J. R., TELENIUS, J. M., OUDELAAR, A. M., DOWNES, D. J., VYAS, P., HUGHES, J.
727    R. & MILNE, T. A. 2019. DOT1L inhibition reveals a distinct subset of enhancers
728    dependent on H3K79 methylation. *Nat Commun,* 10**,** 2803.
729 HIBI, M., OISHI, S., MATSUSHITA, M., YONESHIRO, T., YAMAGUCHI, T., USUI, C., YASUNAGA, K.,
730    KATSURAGI, Y., KUBOTA, K., TANAKA, S. & SAITO, M. 2016. Brown adipose tissue is
731    involved in diet-induced thermogenesis and whole-body fat utilization in healthy
732    humans. *International Journal of Obesity,* 40**,** 1655-1661.
733 JIMENEZ, M. A., AKERBLAD, P., SIGVARDSSON, M. & ROSEN, E. D. 2007. Critical role for Ebf1 and
734    Ebf2 in the adipogenic transcriptional cascade. *Mol Cell Biol,* 27**,** 743-57.
735 KANG, J. Y., KIM, J. Y., KIM, K. B., PARK, J. W., CHO, H., HAHM, J. Y., CHAE, Y. C., KIM, D., KOOK,
736    H., RHEE, S., HA, N. C. & SEO, S. B. 2018. KDM2B is a histone H3K79 demethylase and
737    induces transcriptional repression via sirtuin-1-mediated chromatin silencing. *FASEB J,*
738    32**,** 5737-5750.
739 KONG, X., BANKS, A., LIU, T., KAZAK, L., RAO, R. R., COHEN, P., WANG, X., YU, S., LO, J. C.,
740    TSENG, Y. H., CYPESS, A. M., XUE, R., KLEINER, S., KANG, S., SPIEGELMAN, B. M. &
741    ROSEN, E. D. 2014. IRF4 is a key thermogenic transcriptional partner of PGC-1alpha. *Cell,*
742    158**,** 69-83.
743 LICHTENBELT, W. D. V., VANHOMMERIG, J. W., SMULDERS, N. M., DROSSAERTS, J. M. A. F. L.,
744    KEMERINK, G. J., BOUVY, N. D., SCHRAUWEN, P. & TEULE, G. J. J. 2009. Cold-Activated
745    Brown Adipose Tissue in Healthy Men (vol 360, pg 1500, 2009). *New England Journal of*
746    *Medicine,* 360**,** 1917-1917.
747 LU, X., SIMON, M. D., CHODAPARAMBIL, J. V., HANSEN, J. C., SHOKAT, K. M. & LUGER, K. 2008.
748    The effect of H3K79 dimethylation and H4K20 trimethylation on nucleosome and
749    chromatin structure. *Nat Struct Mol Biol,* 15**,** 1122-4.
750 MARKENSCOFF-PAPADIMITRIOU, E., ALLEN, W. E., COLQUITT, B. M., GOH, T., MURPHY, K. K.,
751    MONAHAN, K., MOSLEY, C. P., AHITUV, N. & LOMVARDAS, S. 2014. Enhancer interaction
752    networks as a means for singular olfactory receptor expression. *Cell,* 159**,** 543-57.
753 MIER, P., ALANIS-LOBATO, G. & ANDRADE-NAVARRO, M. A. 2017. Protein-protein interactions
754    can be predicted using coiled coil co-evolution patterns. *J Theor Biol,* 412**,** 198-203.
755 MIN, J., FENG, Q., LI, Z., ZHANG, Y. & XU, R. M. 2003. Structure of the catalytic domain of
756    human DOT1L, a non-SET domain nucleosomal histone methyltransferase. *Cell,* 112**,**
757    711-23.
758 NG, H. H., FENG, Q., WANG, H., ERDJUMENT-BROMAGE, H., TEMPST, P., ZHANG, Y. & STRUHL,
759    K. 2002. Lysine methylation within the globular domain of histone H3 by Dot1 is
760    important for telomeric silencing and Sir protein association. *Genes Dev,* 16**,** 1518-27.
761 NGUYEN, A. T. & ZHANG, Y. 2011. The diverse functions of Dot1 and H3K79 methylation. *Genes*
762    *Dev,* 25**,** 1345-58.
763 NGUYEN, H. P., YI, D., LIN, F., VISCARRA, J. A., TABUCHI, C., NGO, K., SHIN, G. W., LEE, A. Y. F.,
764    WANG, Y. H. & SUL, H. S. 2020. Aifm2, a NADH Oxidase, Supports Robust Glycolysis and
765    Is Required for Cold- and Diet-Induced Thermogenesis. *Molecular Cell,* 77**,** 600-+.

OKADA, Y., FENG, Q., LIN, Y., JIANG, Q., LI, Y., COFFIELD, V. M., SU, L., XU, G. & ZHANG, Y. 2005.
hDOT1L links histone methylation to leukemogenesis. *Cell,* 121**,** 167-78.
PUIGSERVER, P., WU, Z., PARK, C. W., GRAVES, R., WRIGHT, M. & SPIEGELMAN, B. M. 1998. A
cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell,*
92**,** 829-39.
RAJAKUMARI, S., WU, J., ISHIBASHI, J., LIM, H. W., GIANG, A. H., WON, K. J., REED, R. R. & SEALE,
P. 2013. EBF2 determines and maintains brown adipocyte identity. *Cell Metab,* 17**,** 562-
74.
SAMBEAT, A., GULYAEVA, O., DEMPERSMIER, J. & SUL, H. S. 2017. Epigenetic Regulation of the
Thermogenic Adipose Program. *Trends Endocrinol Metab,* 28**,** 19-31.
SAMBEAT, A., GULYAEVA, O., DEMPERSMIER, J., THARP, K. M., STAHL, A., PAUL, S. M. & SUL, H.
S. 2016. LSD1 Interacts with Zfp516 to Promote UCP1 Transcription and Brown Fat
Program. *Cell Rep,* 15**,** 2536-49.
SANCHEZ-GURMACHES, J., HUNG, C. M. & GUERTIN, D. A. 2016. Emerging Complexities in
Adipocyte Origins and Identity. *Trends in Cell Biology,* 26**,** 313-326.
SCHULZE, J. M., JACKSON, J., NAKANISHI, S., GARDNER, J. M., HENTRICH, T., HAUG, J.,
JOHNSTON, M., JASPERSEN, S. L., KOBOR, M. S. & SHILATIFARD, A. 2009. Linking cell
cycle to histone modifications: SBF and H2B monoubiquitination machinery and cell-
cycle regulation of H3K79 dimethylation. *Mol Cell,* 35**,** 626-41.
SEALE, P., BJORK, B., YANG, W., KAJIMURA, S., CHIN, S., KUANG, S., SCIME, A., DEVARAKONDA,
S., CONROE, H. M., ERDJUMENT-BROMAGE, H., TEMPST, P., RUDNICKI, M. A., BEIER, D.
R. & SPIEGELMAN, B. M. 2008. PRDM16 controls a brown fat/skeletal muscle switch.
Nature, 454**,** 961-7.
SONG, X., YANG, L., WANG, M., GU, Y., YE, B., FAN, Z., XU, R. M. & YANG, N. 2019. A higher-
order configuration of the heterodimeric DOT1L-AF10 coiled-coil domains potentiates
their leukemogenenic activity. *Proc Natl Acad Sci U S A,* 116**,** 19917-19923.
STEGER, D. J., LEFTEROVA, M. I., YING, L., STONESTROM, A. J., SCHUPP, M., ZHUO, D., VAKOC, A.
L., KIM, J. E., CHEN, J., LAZAR, M. A., BLOBEL, G. A. & VAKOC, C. R. 2008. DOT1L/KMT4
recruitment and H3K79 methylation are ubiquitously coupled with gene transcription in
mammalian cells. *Mol Cell Biol,* 28**,** 2825-39.
VAN LEEUWEN, F., GAFKEN, P. R. & GOTTSCHLING, D. E. 2002. Dot1p modulates silencing in
yeast by methylation of the nucleosome core. *Cell,* 109**,** 745-756.
VIRTANEN, K. A., LIDELL, M. E., ORAVA, J., HEGLIND, M., WESTERGREN, R., NIEMI, T.,
TAITTONEN, M., LAINE, J., SAVISTO, N. J., ENERBACK, S. & NUUTILA, P. 2009. Functional
brown adipose tissue in healthy adults. *N Engl J Med,* 360**,** 1518-25.
WANG, W. S. & SEALE, P. 2016. Control of brown and beige fat development. *Nature Reviews*
Molecular Cell Biology, 17**,** 691-702.
WOOD, K., TELLIER, M. & MURPHY, S. 2018. DOT1L and H3K79 Methylation in Transcription and
Genomic Stability. *Biomolecules,* 8.
YI, D., DEMPERSMIER, J. M., NGUYEN, H. P., VISCARRA, J. A., DINH, J., TABUCHI, C., WANG, Y. &
SUL, H. S. 2019. Zc3h10 Acts as a Transcription Factor and Is Phosphorylated to Activate
the Thermogenic Program. *Cell Rep,* 29**,** 2621-2633 e4.
YI, D., NGUYEN, H. P. & SUL, H. S. 2020. Epigenetic dynamics of the thermogenic gene program
of adipocytes. *Biochem J,* 477**,** 1137-1148.