Artificial light at night amplifies seasonal relapse of haemosporidian parasites in a

widespread songbird

Daniel J. Becker[1*], Devraj Singh[1,2], Qiuyun Pan[1], Jesse D. Montoure[1], Katherine M. Talbott[1],

Sarah Wanamaker[1], Ellen D. Ketterson[1,2]

[1]Department of Biology, Indiana University, Bloomington, IN

[2]Environmental Resilience Institute, Indiana University, Bloomington, IN

[*]corresponding author: danbeck@iu.edu

Running head: ALAN and parasite relapse

Keywords: avian malaria; ecoimmunology; generalized additive models; *Junco hyemalis*;

photoperiod; urbanization

Abstract

Urban habitats can shape interactions between hosts and parasites by altering not only exposure

rates but also within-host processes. Artificial light at night is common in urban environments,

and chronic exposure can impair host immunity in ways that may increase infection. However,

studies of causal links between this stressor, immunity, and infection dynamics are rare,

particularly in migratory animals. Here, we experimentally tested how artificial light at night

affects cellular immunity and haemosporidian parasite intensity across the annual cycle of

migrant and resident subspecies of the dark-eyed junco (*Junco hyemalis*). We monitored an

experimental group exposed to light at night and a control group under natural light/dark cycles

as they passed through short days simulating early spring to longer days simulating the breeding

season, followed by fall migration. Using generalized additive models, we show that artificial

light at night increased inflammation, and leukocyte counts were greatest in early spring and fall.

At the start of the experiment, few birds had active infections based on microscopy, but PCR

revealed many birds had chronic infections. Artificial light at night increased parasitemia across

the annual cycle, with strong peaks in spring and fall that were largely absent in control birds. As

birds were kept in indoor aviaries to prevent vector exposure, this increased parasitemia indicates

relapse of chronic infection during costly life history stages (i.e., reproduction). Although the

immunological and parasitological time series were in phase for control birds, cross-correlation

analyses also revealed artificial light at night desynchronized leukocyte profiles and parasitemia,

which could suggest a general exaggerated inflammatory response. Our study shows how a

common anthropogenic influence can shape within-host processes to affect infection dynamics.

Introduction

As suburban and urban habitats expand globally, certain species are attracted to these environments owing to a combination of altered microclimate, predation risk, and resource availability [1,2]. Such changes in biotic and abiotic conditions can have cascading and complex effects on wildlife [3,4]. When considering infectious disease, urban habitats can shape the interactions between hosts and parasites by altering within-host processes such as resistance or tolerance [5]. Understanding how urbanization affects these processes is especially important, as it can facilitate novel cross-species transmission opportunities between various host species [6].

One aspect of urban habitats that could have strong effects on these within-host processes is artificial light at night (ALAN). By altering animal circadian rhythms, ALAN can cause a suite of physiological changes relevant to infection [7]. In particular, chronic exposure to realistic urban levels of ALAN (e.g., 5 lux) can suppress host immune function, including cellular response and microbial killing ability [8,9]. Prolonged ambient light exposure at similar or greater intensities (e.g., 5 to 400 lux) can also dysregulate the inflammatory response, causing inflammation with pathological effects [10,11]. Ultimately, these changes likely alter host infection outcomes and parasite transmission [12]. In a rare test of causal relationships, similar levels of ALAN (i.e., 8 lux) altered gene regulatory networks and extended the infectious period for West Nile virus in house sparrows (*Passer domesticus*) [13]. This suggests that ecologically relevant exposure to ALAN could enhance the transmission of parasites to arthropod vectors and thus increase epidemic potential; however, the consequences of ALAN on infectious disease dynamics in wild hosts generally remain poorly understood.

Although the impacts of ALAN on immunity have been studied in various captive and free-living animals, these consequences have been less explored in migratory species. For birds in particular, many species migrate at night [14], facilitating exposure to light pollution along major migration routes [15]. Many birds also use suburban and urban habitats as stopover sites [16], and widespread ALAN in these environments can further affect habitat selection during migration [17,18]. Because physiological preparation for long-distance migration can itself suppress immune function [19,20], any effects of ALAN on within-host processes could be especially pronounced in migrants. If such changes amplify parasite transmission, urban habitats could become ecological traps for migratory species and in turn hotspots of spillover risk [21].

Here, we experimentally tested how ALAN exposure affects immunity and infection

across the annual cycle of migratory and resident dark-eyed juncos (*Junco hyemalis*). Juncos are

broadly distributed across North America [22], composed of subspecies that diverged 15,000

years ago and vary in migratory behavior, reproductive timing, and morphology [23]. Migratory

slate-colored juncos (*J. h. hyemalis*) breed across Canada and the northern United States,

migrating south in autumn. In the Appalachian Mountains, resident juncos (*J. h. carolinensis*) are

sympatric with overwintering migrants from fall through early spring, when *J. h. hyemalis*

initiate northern migration and *J. h. carolinensis* transition into breeding condition. To recreate

the annual cycle of residents and migrants captured during sympatry, we modified photoperiod,

which provides a consistent and predictive cue for animals to coordinate seasonal events such as

reproduction and migration [24]. Mirroring natural changes in daylength across six months

allowed us to capture these key life history events from early spring through late fall.

To determine effects of ALAN on within-host infection processes across these annual

cycles in migrants and residents, we quantified cellular immunity and parasitemia. We focused

on haemosporidians (i.e., causative agents of avian malaria) [25], which are common in juncos

and are transmitted by arthropod vectors from spring through autumn [26,27]. Haemosporidians

cause a short, acute infection followed by a chronic phase characterized by low parasitemia and

latency in peripheral host tissue [25,28]. Parasites present in hosts sampled in winter therefore

signal chronic infections [27,29]. However, seasonal relapse of chronic infections commonly

occurs from physiological changes in preparation for bird reproduction or migration [29–31].

Manipulating photoperiod in an indoor aviary free of potential vectors allowed us to isolate

effects of ALAN and seasonality on within-host processes, whereas field comparisons may also

have greater exposure to vectors in urban habitats [32]. We predicted ALAN would promote

inflammation and hematological indices of physiological stress (i.e., ratios of heterophils to

lymphocytes [33]). Under control conditions, we expected these cellular immune profiles and

parasitism to generally be synchronized; however, immune dysregulation from ALAN, as

reflected by desynchronization (i.e., a phase lag) of these relationships, could cause chronic

infections to reactivate. Because stages of the annual cycle carry distinct energetic costs, we also

expected these relationships would vary by photoperiod and migratory strategy. Parasitemia

could be greatest under ALAN in late spring, when migrants prepare for northern migration and

residents start summer breeding. However, migrants could display greater parasitemia than

residents in early autumn, when residents complete breeding while migrants initiate their

southern migration.

Methods

Junco sampling and housing

In December 2017, we captured adult male migrant and resident juncos at the Mountain Lake

Biological Station (Pembroke, VA). Each bird was fitted with an aluminum U.S. Fish & Wildlife

Service leg band. Subspecies were differentiated by bill color, plumage, and size [22]. Residents

($n = 18$) and migrants ($n = 24$) were housed with *ad libitum* food and water in an indoor aviary at

Indiana University in Bloomington, Indiana. During a prior experiment, we gradually increased

then decreased photoperiod to simulate a full annual cycle through 2018 [34], and birds were

maintained in a photosensitive state at 9 hours daylength (i.e., 9L) until our experiment in 2019.

Experimental design

We randomly housed equal numbers of residents and migrants in mixed-flock, free-flying rooms

in two separate treatments. Per treatment, birds were held in two interconnected and equal-sized

rooms (2.59 by 2.06 meters). To minimize room artifacts, all rooms were maintained at equal

temperature ($18 \pm 2°C$), were on the same side of the aviary, had the same door and window

placement, and had equal numbers of perches and food and water dishes. Starting in February

2019, we simulated another full annual photoperiodic cycle while exposing control birds to

natural light/dark (LD) cycles and treatment birds to constant ~2.5 ± 0.5 lux average intensity

during night (ALAN). ALAN intensity was determined from a dose-dependent study in great tits,

in which 0.5–5 lux increased reproductive activity [35], and such levels reflect ALAN exposure

of urban birds [36]. All rooms blocked natural sunlight and provided photoperiod light with floor

lamps via fluorescent bulbs (Phillips, 32W; full spectrum white light). Daylight was maintained

at 140 ± 5 lux average intensity, and LD birds were maintained at total darkness at night.

We increased photoperiod every fourteen days by 0.3–1 hour from February 16 to

September 2 (9L:15D to 16L:8D), continued for eight weeks at 16L:8D, and then decreased

photoperiod until November 22 (16L:8D to 9L:15D). At each timepoint, we captured free-flying

indoor birds 30 minutes after the onset of morning photoperiod light. For a complementary study

of how ALAN affects reproductive physiology, birds received a standardized intramuscular

challenge of gonadotropin releasing hormone at each timepoint [37]. Thirty minutes after

injection, we collected 50–100 µL blood from the brachial vein using sterile 26G needles and

made thin blood smears for hematology. Birds were kept in opaque bags between injections to

reduce stress, and the 30 minutes of holding time between challenge and bleeding was uniform

across birds and is unlikely to affect circulating leukocyte counts [33]. We collected repeated

blood smears across nine timepoints (Table 1), which we chose from photoperiodic thresholds in

junco reproductive condition and timing of the junco annual cycle [22,34]. Residents and

migrants initiate elevation of testosterone at 11L and 12L, respectively, followed by gonadal

recrudescence at 12.4L and 13L. These photoperiods correspond to early spring daylengths and

were followed by increasing daylengths simulating the breeding season, for which maintenance

at 16L promotes a photorefractory state to terminate reproduction. Declining daylengths end this

refractory state and simulate onset of fall migration to complete a full annual reproductive cycle.

Some birds were removed from the study due to injury or mortality, resulting in 21 migrants

(LD=10, ALAN=11) and 12 residents (LD=5, ALAN=7) repeatedly sampled for analyses.

Hematology and PCR

Blood smears were stained with buffered Wright–Geimsa (Astral Diagnostics Quick III). From

each sample ($n = 297$), we evaluated leukocyte profiles and parasite intensity using light

microscopy (AmScope, B120C-E1). A single observer, blind to the treatment and migratory

strategy of each bird, first recorded the number of white blood cells under 40X magnification

across 10 random fields to estimate inflammatory state and investment in cellular immunity.

Next, a differential count was performed by recording the identity of the first 100 leukocytes at

1000X magnification [38]. Approximately 100 fields of view at 1000X magnification were

screened for *Plasmodium*, *Haemoproteus*, and *Leucocytozoon* [25]. We derived the mean total

white blood cell count (total leukocytes), ratio of heterophils to lymphocytes (HL ratios), and

total number of haemosporidian parasites (parasite intensity) as three separate response variables.

As microscopy primarily detects active haemosporidian infection [25,28], we used PCR

to detect chronic infections at the start of our experiment (i.e., 9L). We extracted genomic DNA

from erythrocytes in Longmire's solution using RSC Whole Blood DNA Kits (Promega) run on

a Maxwell RSC at Indiana University, and DNA was eluted in 70 µL buffer. We used previously

published primers and PCR protocols to amplify a conserved fragment of the haemosporidian

mitochondrial rRNA gene common to *Plasmodium*, *Haemoproteus*, and *Leucocytozoon* [39,40].

We re-screened any negative birds after initial PCR to reduce possible risks of false negatives.

Statistical analysis

We analyzed all response variables using R [41]. We first used generalized additive mixed

models (GAMMs) with the *mgcv* package to account for potential nonlinear effects of

seasonality (i.e., photoperiod) and treatment [42]. We analyzed data from migrants and residents

separately, as initial analyses suggested no difference in any of these three outcomes (i.e., total

leukocytes, HL ratios, parasitemia) by migratory strategy or its interaction with treatment (Table

S1). For each response variable per migrants and residents, we fit a GAMM with treatment, a

smoothed effect of photoperiod, and their interaction using thin plate splines [43]. GAMMs

included a random effect of individual band number to account for repeated measures. We

modeled both total leukocytes and HL ratios with Tweedie distributions, whereas we used a

zero-inflated Poisson distribution for parasitemia.

To test temporal dependence between cellular immunity and infection, we estimated the

cross-correlation between log-transformed parasitemia and (*i*) log-transformed total leukocytes

and (*ii*) square-root HL ratios for all photoperiod time lags between each time series [44]. We

estimated cross-correlation coefficients for each treatment within migrants and residents. We

then fit an additional set of GAMs with a smoothed effect of photoperiod lag, treatment, and

their interaction with cyclic cubic splines and a Gaussian distribution for each strata of the data.

Strong, positive correlations around the zero lag would support synchrony between the time

series (i.e., in phase), whereas weaker correlations across broader photoperiod lags would

suggest desynchronization of immunity and parasitemia and possibly immune dysregulation.

Results

ALAN effects on leukocyte profiles

ALAN increased total leukocytes compared to LD birds (Fig. 1A), and effect sizes were similar

for residents and migrants (Table 2). Regardless of migratory strategy, we also found strong

seasonal (i.e., photoperiodic) variation in total leukocytes, with greater inflammation at the start

and end of our experiment (i.e., early spring and late fall). Although GAMMs explained up to

59% of the deviance in total leukocytes, seasonal patterns did not vary by treatment. In contrast,

we found weaker effects of treatment and photoperiod on HL ratios (Table S2). ALAN did not

broadly affect HL ratios (Fig. 1B). Only migrants showed an effect of photoperiod on HL ratios,

for which this measure of physiological stress was highest in ALAN birds during early spring.

Haemosporidian intensity under ALAN

At our first blood smear collection (10L), the prevalence of active infection (i.e., infected

erythrocytes detectable with microscopy) was 13.3% for LD birds ($n = 15$) and 16.7% for ALAN

birds ($n = 18$); only ALAN residents had no microscopy-detectable infection (Fig. 2A). PCR

confirmed that 53.3% of LD birds and 66.7% of ALAN birds had chronic infections at the start

of the study (i.e., 9L; Fig. S1), and the odds of PCR-detectable infection did not differ by

treatment or between migrants and residents (Table S3). Most PCR-negative birds at 9L did not

show microscopically detectable infection at our first blood smear collection (10L, 92%; Fig.

S2). Diagnostic mismatches suggest PCR-negative birds likely harbored latent infections (i.e.,

those present in peripheral tissue) that relapsed later in the experiment, although false negatives

are also possible [25,45]. However, by the end of the study, active infections were common in all

experimental groups, ranging from 30% to 100% prevalence ($\overline{x} = 63\%$; Fig. 2A).

Our GAMMs explained up to 63% of the deviance in haemosporidian intensity and

revealed marked effects of ALAN on seasonal parasitemia (Fig. 2B). Haemosporidian intensity

in juncos was higher under ALAN compared to natural LD conditions, although this effect was

greater in residents when compared to migrants (Table 3). For all ALAN birds, parasitemia

peaked around 12L, corresponding to spring, followed by a secondary but smaller parasite

intensity peak in photoperiods corresponding to autumn (i.e., 12–10L). Control residents had

relatively minor seasonal changes in parasitemia, whereas control migrants also increased

haemosporidian intensity in photoperiods corresponding to both spring and fall (Fig. 2B).

Cross-correlation between immunity and infection

ALAN affected the cross-correlation between cellular immunity measures and haemosporidian intensity across our experiment, with our GAMMs explaining up to 84% of the deviance in cross-correlation coefficients (Tables S4 and S5). For control residents, correlations between total leukocytes and parasitemia showed strong nonlinearity ($F=5.43$, $p<0.01$) and were most positive near the zero-photoperiod lag, whereas HL ratios had weak effects on parasitemia ($F=0.20$, $p=0.25$; Fig. 3). However, ALAN removed this synchrony, resulting in positive correlations between total leukocytes and haemosporidian intensity across all lags and a non-significant trend ($F=0$, $p=0.13$). For migrants, LD birds instead showed negative correlations around the zero lag (i.e., when the time series were in phase) between total leukocytes and parasitemia ($F=18.9$, $p<0.01$) and positive correlations between HL ratios and parasitemia ($F=24.6$, $p<0.01$; Fig. 3). Similar to residents, ALAN weakened migrant immunity–intensity relationships for total leukocytes ($F=2.63$, $p=0.02$) and HL ratios ($F=7.06$, $p<0.01$; Fig. 3).

Discussion

Understanding how anthropogenic habitats affect host–parasite interactions is important to forecast how these environments will shape infection risks for wildlife and humans. Here, we experimentally showed that ecologically relevant levels of ALAN, a pervasive component of suburban and urban habitats, increased total leukocytes and haemosporidian intensity in resident and migratory songbirds. Treatment birds had pronounced parasitemia peaks in the spring and fall that were largely absent in control birds, indicating that ALAN caused relapse of chronic infection during energetically costly life history stages. Further, although our immunological and parasitological time series were in phase for control birds, ALAN desynchronized leukocyte profiles and haemosporidian intensity, which could suggest general immune dysregulation.

Because our study occurred in an indoor aviary and PCR confirmed many birds had chronic infections, elevated total leukocytes and haemosporidian intensity in birds exposed to ALAN reflect altered within-host infection dynamics rather than acute infections resulting from vector exposure [32]. In some contexts, ALAN can be immunosuppressive [8,9]; however, the

elevated leukocyte counts observed here more likely indicate inflammation in response to
haemosporidians [46]. In mice, ALAN can induce an exaggerated inflammatory response driven
by proinflammatory cytokines, excessive production of which causes immunopathology and can
limit control of infection [10,11]. Similarly, zebra finches (*Taeniopygia guttata*) exposed to
ALAN also displayed neuroinflammation with these cytokines [47]. The mechanisms by which
the avian immune system controls haemosporidians remains unclear [48], but adaptive responses
are likely also important [49,50]. Related to a possibly exaggerated inflammatory response, total
leukocytes were positively associated with parasite intensity regardless of time lag, whereas
control birds showed more acute synchrony with parasitemia. Notably, we observed these
immunological effects at a lower light intensity (2.5 lux) than similar studies (e.g., 5 to 400 lux)
[8–11,13,47], suggesting that chronic exposure to even dim ALAN could cause immune
dysregulation. However, further quantification of immune gene expression would help identify
the specific pathways by which ALAN affects reactivation of haemosporidian infection [51].

Although our results agree with the prediction that ALAN would promote inflammation,
we observed generally weak effects on HL ratios. HL ratios can signal elevated glucocorticoids
in response to stressors [33], although increases occur more slowly than plasma corticosterone
and can be caused by not only intense stressors but also more mild and chronic challenges [52].
Our results thus suggest chronic stimulation of the hypothalamic–pituitary–adrenal axis may not
be the underlying mechanism linking ALAN with parasite relapse. This finding is similar to
other work on ALAN and within-host processes (e.g., infectious periods for West Nile virus in
house sparrows), which suggested relationships between ALAN and increased viremia were not
mediated by corticosterone [13]. Alternatively, relationships between ALAN, immunity, and
parasitemia could be mediated by other seasonally varying hormones such as testosterone or
melatonin [53]. In particular, melatonin is secreted for a longer duration in short days and can
facilitate immunostimulation [54]. Additionally, fluctuations in host melatonin can influence life
cycle periodicity of some haemosporidians [25]. Suppression of melatonin with ALAN could
explain seasonal relapse during periods in which immunity would otherwise be enhanced [55].
Future work could experimentally assess the hormonal basis of ALAN-induced parasite relapse.

Residents and migrants generally responded in a similar fashion to ALAN, with elevated
leukocytes and increased parasitemia in spring and fall. However, both control and treatment

migrants displayed reactivation of chronic infections in spring, likely signaling physiological

changes in preparation for migration combined with those for reproduction [29,30]. In contrast,

whereas residents exposed to ALAN also experienced dramatic spring relapse, control residents

showed only weak increases in parasitemia in late autumn. These patterns suggest that effects of

ALAN on parasite relapse may be greater during periods of reproduction rather than preparation

for migration, especially as complementary work demonstrated that ALAN also advanced timing

of junco reproduction [37]. Mounting an immune response against haemosporidians can be

costly, which could drive weaker parasite control during energetic trade-offs with reproductive

activity [48,56]. However, our study only assessed transitions into migratory preparedness and

not long-distance migration itself, which could impose stronger effects on parasite control and

therefore on cycles of latency and reactivation [57,58]. Lastly, fall increases in parasitemia for

both residents and migrants under ALAN could again signal melatonin suppression [53,55].

Although the mechanisms linking ALAN with parasite relapse require further study,

particularly for non-migratory hosts, our findings also have implications for the population-level

consequences of infection. For haemosporidian parasites, fitness costs for birds can be highly

variable [25]. In some cases, haemosporidians have caused dramatic population declines in hosts

such as some Hawaiian honeycreepers (e.g., *Vestiaria coccinea*) and possibly house sparrows

(*Passer domesticus*) in Europe [59,60]. Relapse of chronic infections under ALAN in

anthropogenic habitats could accordingly either facilitate infection-related population declines in

less tolerant avian hosts or increase parasitemia in tolerant species that instead serve as sources

of infection for generalist vectors, thereby increasing transmission in the local avian community.

To reduce such negative impacts, alternative lighting schemes in suburban and urban

environments, such as warmer light spectra, minimizing upwards-facing lighting structures, and

restricting use of broad-spectrum light during demanding stages of avian annual cycles (e.g.,

migration), warrant further investigation as ecologically friendly interventions [61–63].

Interdisciplinary collaborations between physiologists and urban planners could enable testing

such interventions in field experiments and facilitate their adoption into these landscapes.

More generally, our study highlights reactivation of chronic or latent infection as a novel

mechanism by which urbanization can affect infectious disease dynamics. Examining how

ALAN affects cycles of latency and reactivation in other host and parasite systems, such as

Borrelia burgdorferi and arboviruses in songbirds [64,65] or henipaviruses and herpesviruses in
bats [66,67], could inform diseases risks relevant for wildlife conservation, domestic animal
health, and human health. Such work could be especially informative for synanthropic wildlife
and migratory species that are increasingly becoming sedentary in anthropogenic habitats [68].

Author contributions

308 DS, DJB, and EDK designed the study; DS, DJB, JDM, KMT, and SW collected samples; QP
309 analyzed blood smears; KMT conducted PCR; and DJB analyzed data and wrote the manuscript.
310 All authors contributed to revisions.

Acknowledgements

313 We thank members of the Ketterson lab and two reviewers for helpful feedback.

Ethics

316 All procedures were approved by the Indiana University Institutional Animal Care and Use
317 Committee (18-030) and conducted under permits issued by the Virginia Department of Game
318 and Inland Fisheries (052971) and U.S. Fish and Wildlife Service (20261).

Funding

This work was supported by the Environmental Resilience Institute, funded by Indiana
University's Prepared for Environmental Change Grand Challenge Initiative. DJB was supported
by an appointment to the Intelligence Community Postdoctoral Research Fellowship Program,
administered by Oak Ridge Institute for Science and Education through an interagency
agreement between the U.S. Department of Energy and Office of the Director of National
Intelligence.

Data availability

329 Data are available in the Dryad Digital Repository:
330 https://datadryad.org/stash/share/IxJYbB61kJJifIIXuEC72ZcIPEpblW6TjIII6sQ8rQM

Tables

Table 1. Experimental sampling timepoints and corresponding photoperiod.

Week	Photoperiod
3/18/2019	10L:14D
4/15/2019	11.4L:12.6D
5/27/2019	12.5L:11.5D
7/8/2019	15L:9D
9/2/2019	16L:8D
10/10/2019	12L:12D
10/24/2019	11L:13D
11/7/2019	10L:14D
11/21/2019	9L:15D

Table 2. Effects of ALAN treatment, photoperiod, and their interaction on total leukocytes. GAMMs were stratified by migratory strategy and fit using restricted maximum likelihood. Fixed effects are presented with model coefficients (categorical) or the estimated degrees of freedom (EDF) and test statistics. Deviance explained (%) is presented for both GAMMs.

Term	Residents (deviance explained=52%)					Migrants (deviance explained=59%)				
	β	t	EDF	F	p	β	t	EDF	F	p
Intercept	-0.30	-1.88			0.06	-0.28	-2.51			0.01
ALAN	0.55	2.67			0.01	0.50	3.18			<0.01
s(photoperiod)			3.27	5.32	<0.01			2.90	7.03	<0.01
s(photoperiod, LD)			0.50	0.34	0.60			0.00	0.07	0.99
s(photoperiod, ALAN)			1.00	2.34	0.13			3.13	2.96	0.07

339   Table 3. Effects of ALAN treatment, photoperiod, and their interaction on parasite intensity.

340   GAMMs were stratified by migratory strategy and fit using restricted maximum likelihood.

341   Fixed effects are presented with model coefficients (categorical) or the estimated degrees of

342   freedom (EDF) and test statistics. Deviance explained (%) is presented for both GAMMs.

Term	Residents (deviance explained=63%)					Migrants (deviance explained=54%)				
	β	t	EDF	F	p	β	t	EDF	F	p
Intercept	-5.14	-2.47			0.01	0.04	0.11			0.91
ALAN	6.90	3.27			<0.01	2.12	3.92			<0.01
s(photoperiod)			1.00	1.05	0.31			2.12	10.3	0.01
s(photoperiod, LD)			1.92	17.2	<0.01			2.33	10.2	0.01
s(photoperiod, ALAN)			3.98	276	<0.01			3.18	20.7	<0.01

Figures

Figure 1. Effects of ALAN on (A) total leukocytes and (B) HL ratios across the simulated annual
cycle (i.e., photoperiod, L) of resident and migrant dark-eyed juncos. Points show medians with
jackknife estimates of standard error and are jittered to reduce overlap. Lines and shaded bands
show the fitted values and 95% confidence interval from each GAMM. The vertical axes are
displayed with a modulus transformation to accommodate lower bounds at zero.

Figure 2. Effects of ALAN on haemosporidian infection dynamics across the simulated annual

cycle (i.e., photoperiod, L) of resident and migrant dark-eyed juncos. (A) Prevalence of

microscopy-detectable infection is displayed with points and 95% confidence intervals. (B)

Parasite intensity is shown as medians with jackknife estimates of standard error. Lines and

shaded bands show the fitted values and 95% confidence interval from the GAMM. The vertical

axes are displayed with a modulus transformation to accommodate lower bounds at zero. All

points are jittered to reduce overlap.

Figure 3. Cross-correlation between immune measures (total leukocytes and HL ratios) and

haemosporidian intensity throughout the duration of the experiment. Points display correlation

coefficients (*r*) at each photoperiod lag and migratory strategy and are colored by treatment.

Lines and shaded bands display the fitted values and 95% confidence interval from each GAM.

References

1. Faeth SH, Warren PS, Shochat E, Marussich WA. 2005 Trophic Dynamics in Urban Communities. *Bioscience* **55**, 399–407.

2. Grimm NB, Faeth SH, Golubiewski NE, Redman CL, Wu J, Bai X, Briggs JM. 2008 Global change and the ecology of cities. *Science* **319**, 756–760.

3. Birnie-Gauvin K, Peiman KS, Gallagher AJ, de Bruijn R, Cooke SJ. 2016 Sublethal consequences of urban life for wild vertebrates. *Environmental Reviews*. **24**, 416–425. (doi:10.1139/er-2016-0029)

4. Murray MH, Sánchez CA, Becker DJ, Byers KA, Worsley-Tonks KEL, Craft ME. 2019 City sicker? A meta-analysis of wildlife health and urbanization. *Front. Ecol. Environ.* **17**, 575–583.

5. Becker DJ, Albery GF, Kessler MK, Lunn TJ, Falvo CA, Czirják GÁ, Martin LB, Plowright RK. 2020 Macroimmunology: The drivers and consequences of spatial patterns in wildlife immune defence. *J. Anim. Ecol.* **89**, 972–995.

6. Hassell JM, Begon M, Ward MJ, Fèvre EM. 2017/1 Urbanization and Disease Emergence: Dynamics at the Wildlife–Livestock–Human Interface. *Trends Ecol. Evol.* **32**, 55–67.

7. Navara KJ, Nelson RJ. 2007 The dark side of light at night: physiological, epidemiological, and ecological consequences. *J. Pineal Res.* **43**, 215–224.

8. Bedrosian TA, Fonken LK, Walton JC, Nelson RJ. 2011 Chronic exposure to dim light at night suppresses immune responses in Siberian hamsters. *Biol. Lett.* **7**, 468–471.

9. Aubrecht TG, Weil ZM, Nelson RJ. 2014 Dim light at night interferes with the development of the short-day phenotype and impairs cell-mediated immunity in Siberian hamsters (Phodopus sungorus). *J. Exp. Zool. A Ecol. Genet. Physiol.* **321**, 450–456.

10. Fonken LK, Weil ZM, Nelson RJ. 2013 Mice exposed to dim light at night exaggerate inflammatory responses to lipopolysaccharide. *Brain Behav. Immun.* **34**, 159–163.

11. Castanon-Cervantes O *et al.* 2010 Dysregulation of inflammatory responses by chronic circadian disruption. *J. Immunol.* **185**, 5796–5805.

12. Kernbach ME, Hall RJ, Burkett-Cadena ND, Unnasch TR, Martin LB. 2018 Dim light at night: physiological effects and ecological consequences for infectious disease. *Integr. Comp. Biol.* **58**, 995–1007.

13. Kernbach ME *et al.* 2019 Light pollution increases West Nile virus competence of a ubiquitous passerine reservoir species. *Proc. Biol. Sci.* **286**, 20191051.

14. Newton I. 2010 *The Migration Ecology of Birds*. Elsevier.

15. Cabrera-Cruz SA, Smolinsky JA, Buler JJ. 2018 Author Correction: Light pollution is
greatest within migration passage areas for nocturnally-migrating birds around the world.
Sci. Rep. **8**, 4799.

16. Rodewald PG, Matthews SN. 2005 Landbird use of Riparian and Upland Forest Stopover
Habitats in an Urban Landscape. *Condor* **107**, 259–268.

17. La Sorte FA, Fink D, Buler JJ, Farnsworth A, Cabrera-Cruz SA. 2017 Seasonal associations
with urban light pollution for nocturnally migrating bird populations. *Glob. Chang. Biol.* **23**,
4609–4619.

18. McLaren JD, Buler JJ, Schreckengost T, Smolinsky JA, Boone M, van Loon EE, Dawson
DK, Walters EL. 2018 Artificial light at night confounds broad-scale habitat use by
migrating birds. *Ecology Letters.* **21**, 356–364. (doi:10.1111/ele.12902)

19. Eikenaar C, Hegemann A. 2016 Migratory common blackbirds have lower innate immune
function during autumn migration than resident conspecifics. *Biol. Lett.* **12**, 20160078.

20. Owen JC, Moore FR. 2008 Swainson's thrushes in migratory disposition exhibit reduced
immune function. *J. Ethol.* **26**, 383–388.

21. Battin J. 2004 When Good Animals Love Bad Habitats: Ecological Traps and the
Conservation of Animal Populations: Ecological Traps. *Conserv. Biol.* **18**, 1482–1491.

22. Nolan V, Ketterson ED, Cristol DA, Rogers CM, Clotfelter ED, Titus RC, Schoech SJ,
Snajdr E. 2002 *Dark-eyed junco (Junco hyemalis)*. American Ornithologists' Union.

23. Milá B, McCormack JE, Castañeda G, Wayne RK, Smith TB. 2007 Recent postglacial
range expansion drives the rapid diversification of a songbird lineage in the genus Junco.
Proc. Biol. Sci. **274**, 2653–2660.

24. Wingfield JC, Hahn TP, Levin R, Honey P. 1992 Environmental predictability and control
of gonadal cycles in birds. *J. Exp. Zool.* **261**, 214–231.

25. Valkiunas G. 2004 *Avian Malaria Parasites and other Haemosporidia*. CRC Press.

26. Deviche P, Greiner EC, Manteca X. 2001 Seasonal and age-related changes in blood
parasite prevalence in dark-eyed juncos (Junco hyemalis, Aves, Passeriformes). *J. Exp.*
Zool. **289**, 456–466.

27. Becker DJ, Talbott KM, Smiley TM, Clark KL, Sauer PE, Ketterson ED. 2019 Leukocyte
profiles vary with breeding latitude and infection status in a seasonally sympatric songbird.
Animal Migration. **6**, 28–40. (doi:10.1515/ami-2019-0004)

28. Valkiunas G, Iezhova TA, Krizanauskiene A, Palinauskas V, Sehgal RNM, Bensch S. 2008
A comparative analysis of microscopy and PCR-based detection methods for blood
parasites. *J. Parasitol.* **94**, 1395–1401.

29. Beaudoin RL, Applegate JE, Davis DE, McLean RG. 1971 A model for the ecology of
avian malaria. *J. Wildl. Dis.* **7**, 5–13.

30. Applegate JE. 1970 Population changes in latent avian malaria infections associated with
season and corticosterone treatment. *J. Parasitol.* **56**, 439–443.

31. Cornelius JM, Zylberberg M, Breuner CW, Gleiss AC, Hahn TP. 2014 Assessing the role of
reproduction and stress in the spring emergence of haematozoan parasites in birds. *J. Exp.*
Biol. **217**, 841–849.

32. Ouyang JQ, de Jong M, van Grunsven RHA, Matson KD, Haussmann MF, Meerlo P, Visser
ME, Spoelstra K. 2017 Restless roosts: Light pollution affects behavior, sleep, and
physiology in a free-living songbird. *Glob. Chang. Biol.* **23**, 4987–4994.

33. Davis AK, Maney DL, Maerz JC. 2008 The use of leukocyte profiles to measure stress in
vertebrates: a review for ecologists. *Funct. Ecol.* **22**, 760–772.

34. Singh D, Reed SR, Kimmitt AA, Alford KA, Ketterson ED. 2019 Breeding at higher
latitude as measured by stable isotope is associated with higher photoperiod threshold and
delayed reproductive development in a songbird. *bioRxiv*

35. Dominoni DM, de Jong M, Bellingham M, O'Shaughnessy P, van Oers K, Robinson J,
Smith B, Visser ME, Helm B. 2018 Dose-response effects of light at night on the
reproductive physiology of great tits (Parus major): Integrating morphological analyses with
candidate gene expression. *J Exp Zool A Ecol Integr Physiol* **329**, 473–487.

36. Dominoni D, Quetting M, Partecke J. 2013 Artificial light at night advances avian
reproductive physiology. *Proc. Biol. Sci.* **280**, 20123017.

37. Singh D, Montoure J, Ketterson ED. 2020 Exposure to artificial light at night accelerates
but does not override latitude-dependent seasonal reproductive response in dark-eyed
juncos. *bioRxiv.* , 2020.06.21.163774. (doi:10.1101/2020.06.21.163774)

38. Campbell TW, Others. 1995 *Avian hematology and cytology*. Iowa State University Press.

39. Fallon SM, Ricklefs RE, Swanson BL, Bermingham E. 2003 Detecting avian malaria: an
improved polymerase chain reaction diagnostic. *J. Parasitol.* **89**, 1044–1047.

40. Arriero E, Pérez-Tris J, Ramírez A, Remacha C. 2018 Trade-off between tolerance and
resistance to infections: an experimental approach with malaria parasites in a passerine bird.
Oecologia **188**, 1001–1010.

41. Team RC, Others. 2013 R: A language and environment for statistical computing.

42. Wood SN. 2017 *Generalized Additive Models: An Introduction with R, Second Edition*.
CRC Press.

43. Wood SN. 2003 Thin plate regression splines. *J. R. Stat. Soc. Series B Stat. Methodol.* **65**,

95–114.

44. Shumway RH, Stoffer DS. 2010 *Time Series Analysis and Its Applications: With R*
Examples. Springer New York.

45. Valkiūnas G, Iezhova TA. 2017 Exo-erythrocytic development of avian malaria and related
haemosporidian parasites. *Malar. J.* **16**, 101.

46. Ots I, Hõrak P. 1998 Health impact of blood parasites in breeding great tits. *Oecologia* **116**,
441–448.

47. Mishra I, Knerr RM, Stewart AA, Payette WI, Richter MM, Ashley NT. 2019 Light at night
disrupts diel patterns of cytokine gene expression and endocrine profiles in zebra finch
(Taeniopygia guttata). *Sci. Rep.* **9**, 15833.

48. Delhaye J, Jenkins T, Glaizot O, Christe P. 2018 Avian malaria and bird humoral immune
response. *Malar. J.* **17**, 77.

49. Videvall E, Cornwallis CK, Palinauskas V, Valkiūnas G, Hellgren O. 2015 The avian
transcriptome response to malaria infection. *Mol. Biol. Evol.* **32**, 1255–1267.

50. Atkinson CT, Dusek RJ, Lease JK. 2001 Serological responses and immunity to
superinfection with avian malaria in experimentally-infected Hawaii amakihi. *J. Wildl. Dis.*
37, 20–27.

51. Fassbinder-Orth CA. 2014 Methods for Quantifying Gene Expression in Ecoimmunology:
From qPCR to RNA-Seq. *Integrative and Comparative Biology*. **54**, 396–406.
(doi:10.1093/icb/icu023)

52. Davis AK, Maney DL. 2018 The use of glucocorticoid hormones or leucocyte profiles to
measure stress in vertebrates: what's the difference? Methods Ecol Evol 9: 1556--1568.

53. Nelson RJ, Demas GE. 1996 Seasonal changes in immune function. *Q. Rev. Biol.* **71**, 511–
548.

54. Trivedi AK, Singh D, Dixit AS, Kumar V. 2017 Pineal Gland, Melatonin, and Timekeeping
in Nonmammalian Vertebrates: Avian Perspective. In *Biological Timekeeping: Clocks,*
Rhythms and Behaviour (ed V Kumar), pp. 521–541. New Delhi: Springer India.

55. Jones TM, Durrant J, Michaelides EB, Green MP. 2015 Melatonin: a possible link between
the presence of artificial light at night and reductions in biological fitness. *Philos. Trans. R.*
Soc. Lond. B Biol. Sci. **370**. (doi:10.1098/rstb.2014.0122)

56. Sheldon BC, Verhulst S. 1996 Ecological immunology: costly parasite defences and trade-
offs in evolutionary ecology. *Trends Ecol. Evol.* **11**, 317–321.

57. Becker DJ, Ketterson ED, Hall RJ. 2020 Reactivation of latent infections with migration
shapes population-level disease dynamics. *bioRxiv*. , 2020.05.12.091736.

499   (doi:10.1101/2020.05.12.091736)

500 58. Nebel S, Bauchinger U, Buehler DM, Langlois LA, Boyles M, Gerson AR, Price ER,
501   McWilliams SR, Guglielmo CG. 2012 Constitutive immune function in European starlings,
502   Sturnus vulgaris, is decreased immediately after an endurance flight in a wind tunnel. *J.*
503   *Exp. Biol.* **215**, 272–278.

504 59. Dadam D, Robinson RA, Clements A, Peach WJ, Bennett M, Rowcliffe JM, Cunningham
505   AA. 2019 Avian malaria-mediated population decline of a widespread iconic bird species. *R*
506   *Soc Open Sci* **6**, 182197.

507 60. Van Riper C III, Van Riper SG, Goff ML, Laird M. 1986 The epizootiology and ecological
508   significance of malaria in Hawaiian land birds. *Ecol. Monogr.* **56**, 327–344.

509 61. Kernbach ME, Cassone VM, Unnasch TR, Martin LB. 2020 Broad-spectrum light pollution
510   suppresses melatonin and increases West Nile virus–induced mortality in House Sparrows
511   (Passer domesticus). *Condor* (doi:10.1093/condor/duaa018)

512 62. Ouyang JQ, de Jong M, Hau M, Visser ME, van Grunsven RHA, Spoelstra K. 2015
513   Stressful colours: corticosterone concentrations in a free-living songbird vary with the
514   spectral composition of experimental illumination. *Biol. Lett.* **11**.
515   (doi:10.1098/rsbl.2015.0517)

516 63. Gaston KJ, Davies TW, Bennie J, Hopkins J. 2012 Reducing the ecological consequences
517   of night-time light pollution: options and developments. *J. Appl. Ecol.* **49**, 1256–1266.

518 64. Gylfe A, Bergström S, Lundström J, Olsen B. 2000 Reactivation of Borrelia infection in
519   birds. *Nature* **403**, 724–725.

520 65. Reisen WK, Chiles RE, Green EN, Fang Y, Mahmood F, Martinez VM, Laver T. 2003
521   Effects of immunosuppression on encephalitis virus infection in the house finch,
522   Carpodacus mexicanus. *J. Med. Entomol.* **40**, 206–214.

523 66. Glennon EE *et al.* 2019 What is stirring in the reservoir? Modelling mechanisms of
524   henipavirus circulation in fruit bat hosts. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **374**,
525   20190021.

526 67. Gerow CM, Rapin N, Voordouw MJ, Elliot M, Misra V, Subudhi S. 2019 Arousal from
527   hibernation and reactivation of Eptesicus fuscus gammaherpesvirus (EfHV) in big brown
528   bats. *Transbound. Emerg. Dis.* **66**, 1054–1062.

529 68. Satterfield DA, Marra PP, Sillett TS, Altizer S. 2018 Responses of migratory species and
530   their pathogens to supplemental feeding. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **373**.
531   (doi:10.1098/rstb.2017.0094)

532