Mycobacterium tuberculosis **complex lineage 5 exhibits high levels of within-lineage**

genomic diversity and differing gene content compared to the type strain H37Rv

C. N'Dira Sanoussi[1,2,3], Mireia Coscolla[4], Boatema Ofori-Anyinam[5], Isaac Darko Otchere[6], Martin Antonio[7],

Stefan Niemann[8,9], Julian Parkhill[10,11], Simon Harris[10], Dorothy Yeboah-Manu[6], Sebastien Gagneux[12,13], Leen

Rigouts[2,3], Dissou Affolabi[1], Bouke C. de Jong[2], Conor J. Meehan[2,14]

[1] Laboratoire de Référence des Mycobactéries, Cotonou, Benin

[2] Mycobacteriology Unit, Institute of Tropical Medicine, Antwerp, Belgium

[3] Department of Biomedical Sciences, University of Antwerp, Antwerp, Belgium

[4] I2SysBio, University of Valencia-FISABIO joint Unit, Valencia, Spain

[5] Center for Global Health Security and Diplomacy, Ottawa, Canada

[6] Noguchi Memorial Institute for Medical Research, University of Ghana, Legon, Accra, Ghana

[7] Medical Research Council Unit in The Gambia at the London School of Hygiene and Tropical Medicine,

Fajara, The Gambia

[8] Research Center Borstel, Molecular and Experimental Mycobacteriology, Borstel, Germany

[9] German Center for Infection Research, partner site Borstel-Hamburg-Lübeck-Riems, Borstel Germany

[10] Wellcome Sanger Institute, United Kingdom

[11] Department of Veterinary Medicine, University of Cambridge, Cambridge, UK

[12] Swiss Tropical and Public Health Institute, Basel, Switzerland

[13] University of Basel, Basel, Switzerland

[14] School of Chemistry and Biosciences, University of Bradford, Bradford, UK

Abstract

Pathogens of the *Mycobacterium tuberculosis* complex (MTBC) are considered monomorphic,

with little gene content variation between strains. Nevertheless, several genotypic and

phenotypic factors separate the different MTBC lineages (L), especially L5 and L6 (traditionally

termed *Mycobacterium africanum*), from each other. However, genome variability and gene

content especially of L5 and L6 strains have not been fully explored and may be potentially

important for pathobiology and current approaches for genomic analysis of MTBC isolates,

including transmission studies.

We compared the genomes of 358 L5 clinical isolates (including 3 completed genomes and

355 Illumina WGS (whole genome sequenced) isolates) to the L5 complete genomes and

H37Rv, and identified multiple genes differentially present or absent between H37Rv and L5

strains. Additionally, considerable gene content variability was found across L5 strains,

including a split in the L5.3 sublineage into L5.3.1 and L5.3.2. These gene content differences

had a small knock on effect on transmission cluster estimation, with clustering rates

influenced by the selection of reference genome, and with potential over-estimation of recent

transmission when using H37Rv as the reference genome.

Our data show that the use of H37Rv as reference genome results in missing SNPs in genes

unique for L5 strains. This potentially leads to an underestimation of the diversity present in

the genome of L5 strains and in turn affects the transmission clustering rates. As such, a full

capture of the gene diversity, especially for high resolution outbreak analysis, requires a

variation of the single H37Rv-centric reference genome mapping approach currently used in

most WGS data analysis pipelines. Moreover, the high within-lineage gene content variability

suggests that the pan-genome of *M. tuberculosis* is at least several kilobases larger than

previously thought, implying a concatenated or reference-free genome assembly (*de novo*)

approach may be needed for particular questions.

Keywords: gene content, *M. africanum*, Lineage 5, H37Rv, reference genome, genomic

diversity, within-lineage variability

Data summary

Sequence data for the Illumina dataset are available at European Genome-phenome Archive

(EGA; https://www.ebi.ac.uk/ega/) under the study accession numbers PRJEB38317 and

PRJEB38656. Individual runs accession numbers are indicated in Table S8.

PacBio raw reads for the L5 Benin genome are available on the ENA accession SAME3170744.

The assembled L5 Benin genome is available on NCBI with accession PRJNA641267. To ensure

naming conventions of the genes in the three L5 genomes can be followed, we have uploaded

these annotated GFF files to figshare at https://doi.org/10.6084/m9.figshare.12911849.v1.

Custom python scripts used in this analysis can be found at

https://github.com/conmeehan/pathophy.

Introduction

Tuberculosis (TB) is caused by pathogenic bacteria of the *Mycobacterium tuberculosis*

complex (MTBC) that consists of 9 human-adapted lineages and several animal-adapted

lineages (Mireia Coscolla and Gagneux 2014; Ngabonziza et al. 2020; M Coscolla et al. 2020;

Brites et al. 2018). This group is highly clonal with no horizontal gene transfer (Boritsch et al.

2016; Chiner-Oms et al. 2019). Strains of particular lineages are primarily defined by large

sequence polymorphisms (LSP, the presence or deletion of genomic regions) such as the TbD1

region (MTBC specific deletion 1), other regions of difference (RD) (Boritsch et al. 2016; Brosch

et al. 2002; Gordon et al. 1999; Behr et al. 1999; Gagneux et al. 2006), and signature single

nucleotide polymorphisms (SNPs)(Coll et al. 2014). RDs are particular long sequence genomic

region deleted in some groups of MTBC strains but present in others; so that MTBC strains

can be classified in groups using that LSP (Gordon et al. 1999; Brosch et al. 2002; Mostowy et

al. 2002; Gagneux et al. 2006). Broadly, the lineages of the MTBC occur within three major

clades: (1) L1-L4 and L7 form one group, (2) L5, L6, L9 and the animal lineages form another,

and (3) L8 sits within its own clade (Coscolla and Gagneux 2014; Ngabonziza et al. 2020;

Coscolla et al. 2020), based on the presence/absence of specific RDs, especially TbD1

(Firdessa et al. 2013; Brosch et al. 2002; Nebenzahl-Guimaraes et al. 2016). L5 and L6 are also

called *M. tuberculosis* var. africanum (or historically *M. africanum* West-african 1 and 2

respectively) (Riojas et al. 2018; Niemann et al. 2004) and are restricted to West Africa where

they cause up to 40% of human TB (de Jong, Antonio, and Gagneux 2010). Reasons for the

geographical restriction of L5 and L6 remain unclear, however, adaptation to particular

human subpopulation has been suggested (Intemann et al. 2009; Thye et al. 2011; Comas et

al. 2013; Gagneux et al. 2006).

Several phenotypic and genotypic features separate L5 and L6 from the other human-adapted

lineages. Some TB diagnostics have a lower performance for L5 and L6 strains, compared to

other MTBC lineages (Sanoussi et al. 2018; Ofori-Anyinam et al. 2016) and these lineages are

less likely to grow in culture (Sanoussi et al. 2017), with dysgonic appearance on solid medium

(Leao et al. 2004; Pattyn et al. 1970; Sanoussi et al. 2018; Niemann et al. 2004). Mutations in

genes essential for growth in culture were identified for L6 strains (Gehre et al. 2013), yet for

L5 strains the reasons for the difficulty in growth remain unclear but may be related to the

presence of the TbD1 region and the absence of RD9 (Brosch et al. 2002). L6 strains and those

of the closely related animal strains/lineages such as *M. bovis* are reported to be less virulent

in humans than those from other human-adapted MTBC lineages in population-based studies

(de Jong et al. 2008; Magnus 1966; Reiling et al. 2013) and in genome studies of mutations in

virulence regulation genes/systems (Gonzalo-Asensio et al. 2014; Ofori-Anyinam et al. 2020).

Infection with an L6 strain progresses slowly to TB disease, and is associated with impaired

immunity in some settings but not all (e.g. HIV infection (de Jong et al. 2008)). In contrast,

little is known about the genomics, virulence and disease progression of L5 strains.

Whole genome sequencing (WGS) based on next generation sequencing (NGS) of MTBC

strains often involves data analysis by mapping of short sequence reads to a reference

genome, usually H37Rv (Cole et al. 1998; Camus et al. 2019; Meehan et al. 2019) or a

reconstructed ancestor with the same gene content as H37Rv (Comas et al. 2013; Goig et al.

2019, 2020). The resulting SNPs are then used for drug resistance determination, subtyping

and transmission analyses (Meehan et al. 2019). However, since H37Rv is a L4 strain, it might

not be representative of other MTBC lineages, including L5 strains. Thus, if H37Rv is used as

basis for genome analysis, several genes or larger genomic regions may be missed resulting

in an underestimation of genome diversity among strains of other lineages that may,

however, provide additional information e.g. for transmission analysis.

The members of the MTBC evolved from an environmental organism to an obligate pathogen

through genome reduction and acquisition of new genes (Gagneux 2018) and it is known that

some differences in gene content exist between lineages (Meehan et al. 2019; Bifani et al.

2000; Periwal et al. 2015; Kato-Maeda et al. 2001; O'Toole and Gautam 2017; Gagneux et al.

2006; Tsolaki et al. 2005). Furthermore, it was reported that genes had a higher genetic

diversity among L6 strains compared to L5 ones (Darko Otchere et al. 2018), but the difference

in gene content between these two lineages was not investigated. Also, little is known about

the gene differences between L4 and L5 and the potential limitations this may impose for in-

depth analysis of L5 genome studies (e.g. sub-lineage detection and transmission tracking).

Similarly, little is known about within-lineage diversity in terms of gene content.

To address these questions, we assessed the gene content diversity of L5 by analyzing in the

context of whole genome sequencing and reference selection. To this end, we analyzed 358

genomes of L5 strains including three completed genomes and compared them to H37Rv and

closely related lineages (L6, *M. tuberculosis* var. bovis (hereafter called *M. bovis*)(Riojas et al.

2018)). Our main focus was to determine particularities in the genomes of L5 strains and to

define the level of within-lineage gene content differences among strains of this lineage of

the MTBC.

Materials and Methods

Genomes

Complete genomes (PacBio, completed long reads). Three completed L5 genomes from

clinical isolates, all sequenced with the PacBio SMRT technology, were analyzed. One genome

was from a Benin isolate (PcbL5Ben; sequenced in this study), one from an isolate from The

Gambia (PcbL5Gam, WBB1453_11-00429-1) (Phelan et al. 2018), and one from Nigeria

(PcbL5Nig, WBB1454_IB091-1) (Phelan et al. 2018). These genomes are also from disparate

parts of the L5 diversity, representing a broad range of L5 sub-lineages, as were recently

defined (Coscolla et al. 2020). The previously published reference/complete genomes H37Rv

(L4) (NC_000962.3)(Cole et al. 1998; Camus et al. 2019; Lew et al. 2011), L6 (GM041182,

Genbank accession n°: FR878060.1, GCF_0001593225.1_ASM159322v1) (Bentley et al. 2012)

and *M. bovis* (AF2122/97, accession: LT708304.1) (Malone et al. 2017) were also included.

Whole genomic DNA extraction. Genomic DNA extraction was performed on growth from

fresh Löwenstein-Jensen slants using the semi-automated Maxwell 16 Cell DNA purification

kit in the Maxwell 16 machine, or from the late exponential phase of growth in 7H9 medium,

using the CTAB method (Belisle and Sonnenberg 1998; Coscolla et al. 2020)

Assembly and annotation of the completed PacBio genomes. The Benin Pacbio genome was

assembled using HGAP (Chin et al. 2013) and Quiver (Chin et al. 2013) and checked for

sufficient quality and coverage (a 10,000 bp sliding window coverage was always above 65x

and average coverage across entire genome is 107x). The genome sequences of all three

completed PacBio genomes were annotated using Prokka-1.12 (Seemann 2014) based on the

reference genome H37Rv annotation.

Gene presence-absence analysis

Gene content differences were assessed with an all-vs-all BLASTn approach, using BLAST+

version+2.8.1 (Altschul et al. 1990; Camacho et al. 2009) (Fig.1). For a specific genome-

genome comparison, the following procedure was used: the gene sequences of genome 1 (ffn

file; coding regions only) were compared to those of genome 2 using the default cut-offs to

look for any homology for each gene. Those genes found in genome 1 and not in genome 2

were then compared to the completed PacBio genome (fna file; coding and non-coding

regions) of genome 2 to look for pseudogenes (herein qualified as ''suspected pseudogenes'')

using BLASTn. These ''suspected pseudogenes'' were then confirmed using tBLASTn of the

genome 1 protein sequences (faa file) compared to the completed PacBio genome (fna file)

of genome 2. This procedure was used to compare all three L5 completed PacBio genomes to

each other as well as each to H37Rv. Those genes present or absent in H37Rv (or pseudogenes

in either) were compared in a similar manner to the L6 and *M. bovis* reference strains to

determine if these genes/pseudogenes are L5-specific (i.e. present in L5, but not in H37Rv, L6

and *M. bovis*).

The ffn files of H37Rv, the three L5 completed PacBio genomes and the L6 complete genomes,

were also aligned using progressive Mauve (ver 20150226)(Darling et al. 2004) to identify

rearrangements and examine synteny.

L5 Illumina (short reads) genomes. In total 355 L5 isolates genomes sequenced on the

Illumina platform from various countries were included in the study. These genomes were

derived from a larger study on genetic diversity of L5 and L6 (Coscolla et al. 2020). After

reducing isolate redundancy (i.e. one representative retained for those that were extremely

closely related), 205 L5 isolates genomes formed a non-redundant dataset. These 205 isolates

(genomes) originated from West, South, East and Central Africa (Table S8), but primarily

(n=155) from two regions within West Africa: the Western part of West Africa (including The

Gambia, Sierra Leone, Ivory Coast, Liberia, Guinea and Mali) and the Eastern part of West

Africa (including Ghana, Benin, and Nigeria) (Coscolla et al. 2020). In general, L5 and L6 are

geographically restricted to West and Central-Africa, making this L5 genome selection

representative of all the geographical regions affected.

Mapping of L5 Illumina reads to H37Rv and L5 completed PacBio genomes.

Raw reads (fastq files) of the 205 non-redundant Illumina genomes were mapped respectively to H37Rv and each of the 3 L5 completed PacBio genomes, using the MTBseq pipeline (Kohl et al. 2018). The depth mapping coverage of the samples in the clinical WGS data Illumina genomes against the reference genomes (percent read mapping, unambiguous coverage mean) was compared between the three L5 completed PacBio genomes and between each L5 completed PacBio genome and H37Rv. Mapping statistics parameters such as percent unambiguous total base, uncovered bases, SNPs, deletions, insertions, substitutions, percent genes mapped to reference, were also compared using each L5 completed PacBio genome or H37Rv as a reference.

Checking unique versus missing genes in Illumina L5 genomes, and SNPs in those confirmed as L5 specific. The genes found present or absent in L5 based on genome comparisons of the three L5 completed PacBio genomes with H37Rv were checked for their expected presence or absence in the L5 Illumina genomes. Using the Position Tables produced by the MTBseq pipeline, a gene was considered absent if 95% of its position in the genome had less than 8 reads covering them. From this data, a gene presence/absence matrix was generated for L5 Illumina genomes mapped to H37Rv and each of the three L5 completed PacBio genomes. Genes found to be L5-specific (present in completed PacBio and Illumina genomes) were also checked for SNPs in these genes using each of the L5 completed PacBio genomes as a reference. The script used for undertaking this analysis can be found at https://github.com/conmeehan/pathophy.

Calculating the effect of reference genome selection on pairwise SNP distances

Transmission analysis of MTBC isolates often involves clustering isolates together based on

specific SNP cut-offs (Meehan et al. 2019, 2018; Walker et al. 2013). We assessed whether

the selection of reference genome (H37Rv, PcbL5Ben, PcbL5Gam or PcbL5Nig) changed the

clustering rate of L5 isolates. For this we used the non-redundant set of L5 genomes from a

previous study of L5 diversity (Coscolla et al. 2020), which included the 205 detailed above

and a further 145 isolates that were closely related to at least one of these 205 isolates.

Additionally, the TB-Profiler online SRA search tool (https://tbdr.lshtm.ac.uk/sra) was used to

identify a further five isolates that were not in this dataset, resulting in 355 isolates included

in this clustering study.

SNP alignments of the Illumina-sequenced genomes were created by first mapping to each of

the four reference genomes (H37Rv and three PcbL5). The Amend function of MTBseq does

this automatically for H37Rv, including masking of repetitive regions, accounting for 10% of

the genome (Cole et al. 1998), and exclusion of columns with 95% ambiguous calls. To

undertake this for the three L5 reference genomes, repetitive regions were first determined

from the annotations of the genes. All genes whose description contained one of the

following words was excluded: integrase, PE family, PE-PGRS, phage, transposase. SNP

alignments were then created using MTBseq as was done for H37Rv. Pairwise distance

matrices, one for each alignment based on each reference genome, were created using a

python script which can be found at https://github.com/conmeehan/pathophy. Loose

transmission clusters were created from these matrices at a cut-off of one, five and twelve

SNPs, as previously described (Meehan et al. 2018). Clustering rates and the presence of a

strain in a transmission cluster was then compared across the four reference genome

mapping approaches.

Determination of putative function of genes differentially present or absent in the L5 completed PacBio genomes

To find the (putative) function of genes present or absent only in L5 strains, and genes present in L5 strains but pseudogenes in H37Rv/L6/*M. bovis* or vice-versa, the fasta sequence of the genes were searched against the NR database of NCBI using BLASTX-2.8.1+ (Altschul et al. 1997) as well as against the Tuberculist database (http://tuberculist.epfl.ch), Mycobrowser (https://mycobrowser.epfl.ch/) (Kapopoulou, Lew, and Cole 2011) and literature. Furthermore, the gene function group/class was found using the COG database (Tatusov et al. 2000) and Mycobrowser.

Determination of the sublineage of the L5 strains

The sublineage of the strains was determined by looking for sublineage defining SNP in the L5 genomes as previously described (Coscolla et al. 2020; Ates et al. 2018). The sublineage defining SNP were searched in the SNP data generated using the MTBseq pipeline with H37Rv as the reference genome.

Results

Comparative analysis of the completed L5 PacBio genomes and other lineages reveal differences in gene content

The number of genes, including paralogs, in the completed L5 PacBio genomes was: 4189 in the PcbL5Ben genome, 4162 in PcbL5Gam, and 4134 in PcbL5Nig versus 4126 in H37Rv, 4126 in the reference L6 genome and 4059 in the reference *M. bovis* genome. Hence, the maximum gene count difference was 55 genes among the three L5 isolates from this study, 63 genes

among the human-adapted lineages (H37Rv, L5, L6), and 130 genes among the human- and

animal adapted lineage (H37Rv, L5, L6 and *M. bovis*).

The visualization of the structure of the genomes showed that all three L5 genomes had a

region (herein called region A) that was absent in H37Rv (gap between 1683331-1716665 on

Fig.2). Furthermore, the Nigerian L5 completed PacBio genome (PcbL5Nig) was missing an

additional region (herein called region B, between 1966666-2000000 in Fig.2) present in the

Benin and Gambian L5 genomes (Fig.2) and also present in H37Rv.

The Benin genome contained 5 genes (a block of 3 genes (1148 bp, 1104 bp, 2840 bp) and the

other 2 (152 bp, 368 bp) randomly spread in the genome) that were neither present in the

Nigerian or Gambian genomes, while the others contained respectively 1 (150 bp) and 2 (152

bp, 366 bp) unique genes randomly spread in the genome. These increased the pangenome

of L5 by at least 6,280 bp. The Benin and Gambian strains each contained 33 genes that were

not present in the Nigerian strain (Fig.3, Table 1, Table S1). Interestingly, 32 of these genes

were also present in H37Rv (Fig.4, Table 1, Table S1). These 32 genes were thus missing in the

Nigerian L5 strain exclusively, meaning that the default/ancestral state have the genes and

then is loosing them. Those 32 genes- absent in the Nigerian L5 strain only- were sequentially

contiguous and formed three blocks of 18, 12 and 2 genes (Table S1). We found that 30 of

those genes, the blocks of 18 and 12 genes were separated by one gene and represented the

region B (mentioned in the section above, Fig 2) found missing in the Nigerian L5 genome

during another analysis (comparison of the structure of the 3 L5 completed PacBio genomes

using Mauve pipeline). Ten genes (region A mentioned above) that were shared by the three

L5 genomes were absent in H37Rv, while 9 genes (spread across the genome) present in

H37Rv were not present in any of the three L5 genomes. Two genes (*Rv2073c*, *Rv2074*) of

those 9 genes were only present in H37Rv but completely absent in the L5, L6 and *M. bovis*

complete genomes.

Six of the genes shared by the three L5 completed PacBio genomes were confirmed

pseudogenes in H37Rv (Table S2). Three genes present in H37Rv were confirmed to be

pseudogenes in the three L5 completed PacBio genomes (Table S3).

Gene presence/absence and related SNPs in lineage specific genes in a wider set of clinical

isolates

The mapping estimates of the Illumina genomes are presented in Table 2. Mapping quality

and coverage against a completed L5 reference was superior to the H37Rv reference

approach (Table 2, Fig.5), as expected. Using the completed PacBio genomes from Benin and

The Gambia had similar mapping estimates that were better than those of the Nigerian L5

genome, likely due to the large deletion in this genome. The genes specific to PcbL5Ben (5),

PcbL5Gam (2), PcbL5Nig (1) were each found at various rates among the 205 L5 Illumina

genomes (77.6%-97.1%, 159-199/205, Table 3).

Interestingly, 2.9% of the Illumina L5 genomes (6/205) had similar patterns of large gene loss

as the Nigerian completed PacBio genome as they missed 30 of the 33 genes present in the

Benin and Gambian genomes and H37Rv. These six PcbL5Nig-like Illumina L5 genomes formed

a monophyletic group, within the L5 clade (Fig. 6), suggesting a single loss of these gene

clusters, although those six L5 genomes originated from several different countries including

Benin, Ghana and Nigeria. The two blocks of 18 (14835 bp, *Rv1493* through *Rv1509*) and 12

genes (10209 bp, *Rv1511* through *Rv1521*), separated by one gene (3441 bp, *Rv1510*)

amounted in total to 25044 bp which is around 0.6% of the length (around 4 million bp) of a

typical MTBC (H37Rv) genome. The two blocks contain genes whose annotations include

mutB, *mazE4*, *mazF4* and others with various putative functions (Table S4).

Four of the ten genes present in the three completed L5 PacBio genomes but absent in H37Rv

were found in all the 205 L5 Illumina genomes while the other were found in variable amounts

81-98% (165-201) (Table S5). These four gene amounted to a total of 2382 bp (Table S5),

representing around 0.05% of the length of a typical MTBC genome. These four genes include:

Mb2048c, PE35, a hypothetical protein- possibly an IS*256* transposase, and a hypothetical

protein possibly related to the CAAX conserved domain. Two of these four genes were present

in L5 genomes only (PE35 and hypothetical protein possibly CAAX conserved domain) while

the remaining two (*Mb2048c* and hypothetical protein possibly IS256 transposase) were

found in L6 and *M. bovis* as well (Table S5). Importantly for phylogenetic purposes, SNPs were

detected in all four genes in 1.5-3.4% of the 205 L5 Illumina genomes (Table S5). Predicted

functions for all L5-specific genes are listed in table S5.

Five (*Rv1977*, *Rv1979c*, *Rv1993c*, *Rv1995*, *Rv2073c*) of the 9 genes present in H37Rv and

absent in the three completed L5 PacBio genomes were absent in all the 205 L5 Illumina

genomes. These five genes amounted to a total of 4,245 bp (Table S6), representing around

0.1% of the length of a typical MTBC genome. Four of these genes (*Rv1977*, *Rv1979c*, *Rv1993c*,

Rv1995) were absent in L5 strains only (i.e. present in L6 and *M. bovis* reference genomes)

while the fifth (*Rv2073c*) was absent in L6 and *M. bovis* as well (Table S6). The four other

genes absent from the completed PacBio genomes were present in a minority of the L5

Illumina genomes (Table S6, Fig. 5). Predicted functions for all nine genes are listed in table

S6.

Sublineage of the strains (PacBio completed and Illumina genomes)

The determination of the strains sublineage revealed that the PcbL5Ben is an L5.3 strain that

contains the 30 genes (two blocks of 18 and 12 genes: *Rv1493* through *Rv1509* and *Rv1511*

through *Rv1521*) whereas PcbL5Nig is also a L5.3 strain, but situated in a different clade (Fig.

6). Based on this region (a new RD), we classified PcbL5Ben as an L5.3.1 strain and the

PcbL5Nig as a L5.3.2 strain; the PcbL5Gam isolate is an L5.1.5 strain. The Illumina genomes

were distributed as follows (Table S7): 90 L5.1.1; 17 L5.1.2; 12 L5.1.3; 21 L5.1.4; 9 L5.1.5

amounting to a total of 149 L5.1 strains (72.7%); 13 (6.3%) L5.2 strains; 8 L5.3.1; 6 L5.3.2

amounting to a total of 14 L5.3 strains (6.8%) and 29 (14.2%) strains of unknown sublineages

(potentially new sublineages). All L5 strains including L5.1, L5.2, L5.3.1, unknown sublineages,

except L5.3.2 strains have that 30 genes region.

Changes to transmission clustering rates based on selection of reference genome

Transmission analysis was undertaken on an expanded set of 355 L5 isolates using each of the

four reference genomes (H37Rv, PcbL5Ben, PcbL5Gam and PcbL5Nig) for the SNP mapping.

The current gold standard is to use H37Rv for calling SNPs and then creating transmission

clusters at a specific SNP cut-off. For this dataset, using a 12 SNP cut-off, it was found that

40.6% (n=144) isolates were within a transmission cluster with at least one other isolate (Table

4), for a otal of 55 clusters. Substituting PcbL5Ben instead of H37Rv as the reference genome

reduced the number of clusters to 54, removing 3 isolates from transmission clusters, resulting

in a clustering rate of 39.7% at a 12 SNP cut-off (Table 4). A similar reduction in transmission

clustering was observed when using the PcbL5Nig genome as a reference but not when using

the PcbL5Gam. When a more conservative SNP cut-off was used, such as 1 SNP or 5 SNPs,

the reduction in transmission clustering rates was more pronounced and was observed for all

the PcbL5 genomes (Table 4). For example, at a 1 SNP cut-off, H37Rv-based mapping resulted

in 100 isolates (28.2%) being placed inside transmission clusters, but this reduced to 94 or 95

isolates (26.5-26.8%) when using a L5 genome as the reference. This demonstrates that even

on this small dataset, transmission cluster estimations are affected by the selection of reference

genome, with potential over-estimation of recent transmission when using H37Rv as the

reference genome.

Discussion

Comparison of completed PacBio and Illumina short-reads genomes from clinical isolates

revealed gene content diversity within L5 isolates as well as between L5 and other lineages.

We found that four genes, present in all the L5 isolates of our multi-country collection, were

absent in the H37Rv genome. Those genes represent 2,382 bp which is around 0.05% of the

total length (base-pair) of a typical MTBC genome. Furthermore, those genes contain SNPs

that can be used to identify additional diversity between L5 strains. This has implications

when using precise SNP cut-offs to define transmission clusters (Walker et al. 2013; Meehan

et al. 2018) as SNPs in these genes would be missed with H37Rv mapping, resulting in

underestimation of genome diversity of L5 strains. Our findings suggest, that using an L5

genome as a reference would likely increase the resolution of L5 genome analyses over the

H37Rv approach and can potentially increase the discrimination power, leading to a reduction

of the amount of recent L5 transmission inferred.

Our findings are in line with other reports indicating that some genes present in genomes of

clinical MTBC strains were absent in the H37Rv genome (Periwal et al. 2015; O'Toole and

Gautam 2017), and with other recommendations to use additional reference genomes

different to H37Rv (Norman et al. 2019; O'Toole and Gautam 2017; Lee et al. 2020). Although

Lee *et al*. (Lee and Behr 2016) - after comparing various lineages as reference genome for L4

genome analysis - concluded that there is no need to use a lineage-specific reference genome,

their observation was only based on the analysis of L4 clinical isolates and focused on SNPs

and short indels, not larger gene deletions or SNPs within these regions. In contrast, our

findings indicate that mapping of NGS data from L5 strains to L4 reference will have an impact,

in terms of both reference genome coverage and coverage of lineage-specific genes.

Although the use of an L5 genome as reference would have many benefits over H37Rv for

particular study questions, several gene content differences were still observed within the L5

lineage. Each of the three L5 PacBio genomes had genes unshared with the two others that

were all found in L5 Illumina genomes (thus excluding exogenous contamination). The

absence of 30 genes in the Nigerian L5 completed PacBio genome (but shared by the Benin

and Gambian long read genomes), also found in six L5 genomes from different countries,

confirmed the large deletions in the PcbL5Nig genome and showed that it is representative

of some L5 strains in some countries. Deletions in the genomes of strains of MTBC lineages

were previously reported (Tsolaki et al. 2005; Gagneux et al. 2006; Kato-Maeda et al. 2001).

The fact that, on the phylogenetic tree the six isolates formed a monophyletic group of the

(SNP defined) L5.3 sub-lineage (Coscolla et al. 2020) suggests that the loss of these 30 genes

(*Rv1493* through *Rv1509*, and *Rv1511* through *Rv1521*) is a marker of the L5.3.2 sub-lineage,

compared to the sublineage L5.3.1 strains that have these genes intact. Thus, we find that

regions of difference (RDs) are present both between lineages and between sub-lineages.

We found that the genes *Rv2073c* and *Rv2074* that were absent in L5, L6 and *M. bovis*. It was

previously reported that L5, L6 and *M. bovis* all lack the RD9 region (Brosch et al. 2002;

Mostowy et al. 2002; Gagneux et al. 2006), which contains *Rv2073c*, *Rv2074* and partly

Rv2075 (Gordon et al. 1999). The lack of RD9 by L5, L6 and *M. bovis* is thus confirmed by our

data, however one (1/205) of the L5 genomes had *Rv2074* present. Of the five genes (*Rv1977*,

Rv1979c, *Rv1993c*, *Rv1995*, *Rv2073c*) absent in all L5 strains, besides *Rv2073c* included in RD9,

one (*Rv1977*) was part of the RD7 region (Gordon et al. 1999), another (*Rv1979c*) was part of

the RD2 region (Gordon et al. 1999), and *Rv1993c* and *Rv1995* potentially included in other

RD regions. These findings show that gene presence/absence approach sheds more light on

the differences between lineages than the RD approach alone would do.

The classification of the L5-specific genes into functional groups based on Mycobrowser and

Tuberculist databases (Table 4) revealed a variety of functions including: a PE/PPE protein

family (possibly PE35) which may be linked to antigenic variation (NCBI annotation) and

virulence (STRING database annotation (Szklarczyk et al. 2017)); two ABC-transporters; an

IS*256*-like transposase, whose homology (IS256) is a marker of specific virulent lineages of

staphylococci (Murugesan et al. 2018; Gu et al. 2005); and a gene with a CAAX conserved

domain, implicated in post-translation modification or signal transduction (Gao, Liao, and

Yang 2009a, 2009b). Only one (*Rv1979c*) of the four genes absent in only L5 strains had a

predicted function, labelled as "cell wall and cell processes" (Table S6). While *Rv1979c* is

associated with clofazimine and bedaquiline resistance (Zhang et al. 2015; Ghodousi et al.

2019), two of the drugs used for the treatment of RIF-resistant/MDR-TB (WHO 2016, 2019),

minimal inhibitory concentration testing of clofazimine in five L5 strains did not find the

deletion conferred resistance (Merker et al. 2020). Further studies and protein function

discovery is needed to investigate the consequences of the absence of those genes in L5.

The genes *Rv1978* and *Rv1994c* present in H37Rv and the completed reference genomes of

L6 and *M. bovis*, were absent in the 3 completed Pacbio genomes of L5 strains, yet present in

few L5 isolates Illumina genomes (3/205, 1.5% for *Rv1978*; 1/205, 0.5% for *Rv1994c*). *Rv1978*

is required for bacterial survival in macrophages, and non-essential for *in vitro* growth of

H37Rv (Rengarajan, Bloom, and Rubin 2005). *Rv1994c* is involved in the regulation and

transport (efflux) of toxic metals, especially copper, which is toxic in excess (Samanovic et al.

2012) and may hamper *in vitro* growth, and survival during the chronic phase of infection,

similar to the effect of disruption of *csoR* (Marcus et al. 2016; Rowland and Niederweis 2012;

Ward, Hoye, and Talaat 2008; Cavet et al. 2003) (Table S6).

The functional annotation for the H37Rv-specific genes *Rv2073c* (short chain dehydrogenase)

and *Rv2074* (pyridoxamine-5-phosphate oxydase: vitamin B6 (pyridoxine)) absent in L5, L6

and *M. bovis* strains is "intermediary metabolism and respiration". Those two genes are part

of the RD9 region, which confirmed their absence in L5, L6 and *M. bovis* that have that region

deleted (Gordon et al. 1999). *Rv2073c* is an NAD(P)-dependent oxidoreductase (NCBI) that

catalyzes a wide range of reactions and is involved in redox sensor mechanisms (Kavanagh et

al. 2008; Sellés Vidal et al. 2018). *Rv2074* was previously thought to be a pyridoxine (vitamin

B6) oxidase, but is now known to be a F420 dependent biliverdin reductase, a cofactor of

vitamin B6 synthesis (UniProt) (Ahmed et al. 2016; Selengut and Haft 2010). Vitamin B6 is

essential for survival and virulence of *M. tuberculosis* (Dick et al. 2010) and its cofactor (F420

dependent biliverdin reductase) is implicated in immune-evasive mechanisms to allow

bacterial persistence (Ahmed et al. 2016; Selengut and Haft 2010)(Table S6). Recently, it has

been reported that the synthesis of F420 might be stimulated by phosphoenolpyruvate

(Grinter et al. 2020), which is the precursor to pyruvate, a supplement often added to culture

medium to improve the *in vitro* growth of L5, L6 and *M. bovis*. Furthermore, there is a

suggestion that F420 is needed for the activation of the antituberculosis drugs pretomanid

and delamanid (Grinter et al. 2020).

The absence of these genes in most L5 isolates suggests that L5 strains would be less likely to

survive in macrophages (*Rv1978*), have reduced growth *in vitro* (*Rv1994c* , as previously found

by Sanoussi et al (Sanoussi et al. 2017)), be less immune-evasive, less persistent and less

virulent (*Rv2074*) than L4 strains (at least H37Rv; as previously suggested for L5 and L6 strains

(Coscolla and Gagneux 2014; Reiling et al. 2013)). In addition, the absence of *Rv2074* in most

L5 (and in the complete genomes of L6 and *M. bovis* as well) suggests that L6 and *M. bovis*

strains would also be less likely to be immune-evasive, to be persistent and to survive than

their L4 counterparts. Despite the presence of vitamin B6 in boiled egg (thus Lowenstein-

Jensen medium), its quantity in the L.J medium is probably insufficient to allow a high yield of

L5, L6 and *M. bovis* growth in culture (survival). Further studies, including vitamin B6

supplementation, could investigate the consequences of the absence of those genes in most

L5 strains (and L6 and *M. bovis* for *Rv2074*).

Transmission clustering analyses revealed differences in clustering rates and inclusion of

isolates in clusters based on the selection of the reference genome (Table 4). Overall, H37Rv-

based mapping was found to place a small percentage more isolates in transmission clusters

than any L5-based mapping approach. This is due to the additional genes found in these

genomes also being present in the clinical isolates and these genes containing differentiating

SNPs. Thus, with an increased discovery of SNPs in non-H37Rv genes, some isolates that

previously satisfied the SNP distance criterion for clustering were now excluded as they are

less similar than they appear to be when using H37Rv as a reference. This was most

pronounced at the lower SNP cut-offs (Table 4). This has strong implications for molecular

epidemiology in West Africa, where most L5 strains are found. Additionally, if similar

scenarios exist for other lineages, this may result in changes to all non-L4 transmission

analyses.

The within-lineage differences in both gene content and potential functionality indicates a

need for more closed genomes of MTBC sub-lineages and that perhaps a sub-lineage-specific

reference genome approach would be appropriate for MTBC WGS and molecular

epidemiology. However, although an *ad hoc* (e.g. outbreak-specific) reference genome

provides a more accurate and precise comparison of strains involved in a specific situation,

that reference is only specific to that particular situation/outbreak/population/lineage and

cannot be used in another context, making comparisons between lineages and settings

difficult.

As an alternative to using a single MTBC-wide, lineage or sub-lineage specific reference

genome approach, the variable presence of genes within one lineage suggests that an L5

pangenome-based reference genome capturing all the known diversity may be required

instead. Such a genome would be a composite of all the genes found in all lineages and sub-

lineages of the MTBC (i.e. both the core and accessory genome) (McInerney et al. 2020) and

is usually represented as a graph instead of a single sequence (Rakocevic et al. 2019; Paten

et al. 2017; Martiniano et al. 2019; Marschall et al. 2018; Church et al. 2015). Alternatively, it

can be a selection of reference genomes, with mapping to all or a subset undertaken, as has

been done with *M. chimaera* and other pathogens (van Ingen et al. 2017; Garimella et al.

2019). Other authors have also reported that because of the genetic variability between

strains, using a single strain genome as reference genome lacks accuracy (Ballouz, Dobin, and

Gillis 2019; Sherman et al. 2019; Li et al. 2010). This MTBC-wide pangenome approach has

been suggested before, including for other organisms (Periwal et al. 2015; Meehan et al.

2019; Jandrasits et al. 2019; Ballouz, Dobin, and Gillis 2019; Medini et al. 2005; Tettelin et al.

2005; Sherman et al. 2019; Li et al. 2010). However, such an approach also has its own

drawbacks, including difficulty in mapping reads that bridge the boundaries between

accessory genes and the rest of the genome, comparing strains of different lineages including

phylogenetic analyses and retention of gene names and codes in clinical use, where H37Rv is

deeply embedded (Marschall et al. 2018). The specific sequence of each gene would also need

to be chosen for such a reference, with the inferred ancestral genome representative of MTBC

lineages approach being the most likely method (Goig et al. 2020, 2018; Comas et al. 2013)

Another alternative approach is a *de novo* assembly reference free approach (Maretty et al.

2017; Iqbal et al. 2012; Meehan et al. 2019; Li et al. 2010) where strains are either assembled

into contigs without the use of a reference genome, or compared to each other without first

calling SNPs. This would allow for clustering of samples regardless of lineage, e.g. for

genotyping or transmission analyses using Mash (software for fast/meta -genome distance

estimation technique) (Ondov et al. 2016) but would require new cut-offs for defining clusters

and many additional steps for further gene annotation and between sample comparisons,

making its clinical use potentially confusing.

A limitation of this study is that the completed (PacBio) and short-read (Illumina) genomes

were derived from positive cultures, excluding possible minority L5 strain diversity as L5

strains are overrepresented in negative cultures (Sanoussi et al. 2017). WGS applied directly

on sputum is more and more needed, especially for ancestral lineages (including L5 and L6)

where growth in culture is problematic (negative culture or dysgonic isolates that are a

challenge to grow for DNA extraction (Sanoussi et al. 2018, 2017)). Also, a limitation regarding

the comparison of gene presence/absence in L5 versus L4 (including H37Rv), L6, *M. bovis*

complete genomes, is that our study included only a single complete genome of L4, L6 and

M. bovis, while these lineages may also display variability similar to the intra-L5 variability we

observed in our study.

In conclusion, the use of a (sub-)lineage reference genome can increase the resolution for

strains comparison in comparison to a H37Rv based mapping approach for L5 genome

analyses for epidemiology (transmission), phylogeny and sub-lineage determination. Still, the

use of a (sub-)lineage reference genome may miss some within-lineage gene differences. For

drug-resistance detection in clinical L5 strains or strains of other lineages, H37Rv could still be

used as reference genome as resistance-related mutations are usually among core genes

(shared across all lineages). The high within-lineage gene content variability suggests the

pangenome of MTBC strains may be larger (at least by 6,280 bp) than previously thought,

implying a reference-free genome assembly (*de novo assembly*) approach may be needed.

Authors and contributors

CNS was involved in conceptualization, methodology, formal analysis, investigation, data
curation, writing of the original draft, review/editing the manuscript and visualization.

MC was involved in conceptualization, methodology, investigation, data curation, review and
editing the manuscript and visualization.

BO was involved in methodology, investigation and review/editing the manuscript.

IDO was involved in resources and review/editing the manuscript

MA was involved in resources.

SN was involved in conceptualization, reviewing/editing the manuscript and supervision.

JP was involved in investigation, data curation and reviewing/editing the manuscript.

SH was involved in conceptualization, investigation, data curation and reviewing/editing the
manuscript.

DY was involved in conceptualization, resources and reviewing/editing the manuscript.

SG was involved in conceptualization and reviewing/editing the manuscript.

LR was involved in conceptualization, resources, review/editing the manuscript and
supervision.

DA was involved in resources, review/editing the manuscript and supervision.

BCdJ was involved in conceptualization, methodology, writing of the original draft,
review/editing the manuscript and supervision.

CJM was involved in conceptualization, methodology, formal analysis, investigation, writing
of the original draft, review/editing the manuscript, supervision and visualization.

Conflict of interest

536 The authors declare no conflicts of interest

Funding information

This work was supported by funds from the Directorate General for Development (DGD),
Belgium (FA4 to C.N.S., B.C.D.J., D.A., and L.R.); the European Research Council-INTERRUPTB
starting grant (number 311725 to B.C.D.J., C.J.M., and L.R.)

M.C. is supported by ESCMID, Ministerio de Ciencia (RYC-2015-18213 and RTI2018-094399-
A-I00) and Generalitat Valenciana (SEJI/2019/011). S.G. is supported by the Swiss National
Science Foundation (grants 310030_188888, IZRJZ3_164171, IZLSZ3_170834 and
CRSII5_177163) and the European Research Council (883582-ECOEVODRTB).

The funders had no role in study design, data collection and interpretation or the decision to
submit the work for publication.

Ethical approval

550 The PacBio sequenced L5 strain from Benin was isolated during the multicentric OFLOTUB
551 study approved by the National Ethics Committee in Cotonou, Benin and each of the
552 participating countries (Merle et al. 2012).

Acknowledgements

554 We would like to acknowledge input from Pim de Rijk Willem and Patrick Beckert for their
555 technical inputs.

Reference

Ahmed, F Hafna, A Elaaf Mohamed, Paul D Carr, Brendon M Lee, Karmen Condic-Jurkic, Megan L
O'mara, and Colin J Jackson. 2016. "Rv2074 Is a Novel F 420 H 2-Dependent Biliverdin
Reductase in Mycobacterium Tuberculosis." https://doi.org/10.1002/pro.2975.

Altschul, Stephen F., Warren Gish, Webb Miller, Eugene W. Myers, and David J. Lipman. 1990. "Basic
Local Alignment Search Tool." *Journal of Molecular Biology* 215 (3): 403–10.
https://doi.org/10.1016/S0022-2836(05)80360-2.

Altschul, Stephen F, Thomas L Madden, Alejandro A Schäffer, Jinghui Zhang, Zheng Zhang, Webb

Miller, and David J Lipman. 1997. "Gapped BLAST and PSI-BLAST: A New Generation of Protein
Database Search Programs." *Nucleic Acids Research*. Vol. 25. Oxford University Press.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC146917/pdf/253389.pdf.

Ates, Louis S, Anzaan Dippenaar, Fadel Sayes, Alexandre Pawlik, Christiane Bouchier, Laurence Ma,
Robin M Warren, et al. 2018. "Unexpected Genomic and Phenotypic Diversity of
Mycobacterium Africanum Lineage 5 Affects Drug Resistance, Protein Secretion, and
Immunogenicity." *Genome Biology and Evolution* 10 (8): 1858–74.
https://doi.org/10.1093/gbe/evy145.

Ballouz, Sara, Alexander Dobin, and Jesse A. Gillis. 2019. "Is It Time to Change the Reference
Genome?" *Genome Biology*. BioMed Central Ltd. https://doi.org/10.1186/s13059-019-1774-4.

Behr, M. A., M. A. Wilson, W. P. Gill, H. Salamon, G. K. Schoolnik, S. Rane, and P. M. Small. 1999.
"Comparative Genomics of BCG Vaccines by Whole-Genome DNA Microarray." *Science* 284
(5419): 1520–23. https://doi.org/10.1126/science.284.5419.1520.

Belisle, John T., and Michael G. Sonnenberg. 1998. "Isolation of Genomic DNA from Mycobacteria."
In *Mycobacteria Protocols*, 31–44. New Jersey: Humana Press. https://doi.org/10.1385/0-
89603-471-2:31.

Bentley, Stephen D., Iñaki Comas, Josephine M. Bryant, Danielle Walker, Noel H. Smith, Simon R.
Harris, Scott Thurston, et al. 2012. "The Genome of Mycobacterium Africanum West African 2
Reveals a Lineage-Specific Locus and Genome Erosion Common to the M. Tuberculosis
Complex." Edited by Pamela L. C. Small. *PLoS Neglected Tropical Diseases* 6 (2): e1552.
https://doi.org/10.1371/journal.pntd.0001552.

Bifani, P, S Moghazeh, B Shopsin, J Driscoll, A Ravikovitch, and B N Kreiswirth. 2000. "Molecular
Characterization of Mycobacterium Tuberculosis H37Rv/Ra Variants: Distinguishing the
Mycobacterial Laboratory Strain." *Journal of Clinical Microbiology* 38 (9): 3200–3204.
https://www.ncbi.nlm.nih.gov/pubmed/10970357.

Boritsch, Eva C., Varun Khanna, Alexandre Pawlik, Nadine Honoré, Victor H. Navas, Laurence Ma,
Christiane Bouchier, et al. 2016. "Key Experimental Evidence of Chromosomal DNA Transfer
among Selected Tuberculosiscausing Mycobacteria." *Proceedings of the National Academy of
Sciences of the United States of America* 113 (35): 9876–81.
https://doi.org/10.1073/pnas.1604921113.

Brites, Daniela, Chloé Loiseau, Fabrizio Menardo, Sonia Borrell, Maria Beatrice Boniotti, Robin
Warren, Anzaan Dippenaar, et al. 2018. "A New Phylogenetic Framework for the Animal-
Adapted Mycobacterium Tuberculosis Complex." *Frontiers in Microbiology* 9 (NOV).
https://doi.org/10.3389/fmicb.2018.02820.

Brosch, R, S V Gordon, M Marmiesse, P Brodin, C Buchrieser, K Eiglmeier, T Garnier, et al. 2002. "A
New Evolutionary Scenario for the Mycobacterium Tuberculosis Complex." *Proceedings of the
National Academy of Sciences* 99 (6): 3684–89. https://doi.org/10.1073/pnas.052548299.

Camacho, Christiam, George Coulouris, Vahram Avagyan, Ning Ma, Jason Papadopoulos, Kevin
Bealer, and Thomas L Madden. 2009. "BLAST+: Architecture and Applications." *BMC
Bioinformatics* 10 (December): 421. https://doi.org/10.1186/1471-2105-10-421.

Camus, Jean-Christophe, Melinda J Pryor, Claudine Me, and Stewart T Cole. 2019. "Re-Annotation of
the Genome Sequence of Mycobacterium Tuberculosis H37Rv." Vol. 24. www.sanger.ac.uk.

Cavet, Jennifer S., Alison I. Graham, Wenmao Meng, and Nigel J. Robinson. 2003. "A Cadmium-Lead-
Sensing ArsR-SmtB Repressor with Novel Sensory Sites. Complementary Metal Discrimination

by NmtR and CmtR in a Common Cytosol." *Journal of Biological Chemistry* 278 (45): 44560–66. https://doi.org/10.1074/jbc.M307877200.

Chin, Chen Shan, David H. Alexander, Patrick Marks, Aaron A. Klammer, James Drake, Cheryl Heiner, Alicia Clum, et al. 2013. "Nonhybrid, Finished Microbial Genome Assemblies from Long-Read SMRT Sequencing Data." *Nature Methods* 10 (6): 563–69. https://doi.org/10.1038/nmeth.2474.

Chiner-Oms, L. Sánchez-Busó, J. Corander, S. Gagneux, S. R. Harris, D. Young, F. González-Candelas, and I. Comas. 2019. "Genomic Determinants of Speciation and Spread of the Mycobacterium Tuberculosis Complex." *Science Advances* 5 (6). https://doi.org/10.1126/sciadv.aaw3307.

Church, Deanna M., Valerie A. Schneider, Karyn Meltz Steinberg, Michael C. Schatz, Aaron R. Quinlan, Chen Shan Chin, Paul A. Kitts, et al. 2015. "Extending Reference Assembly Models." *Genome Biology* 16 (1): 1–5. https://doi.org/10.1186/s13059-015-0587-3.

Cole, S. T., R. Brosch, J. Parkhill, T. Garnier, C. Churcher, D. Harris, S. V. Gordon, et al. 1998. "Deciphering the Biology of Mycobacterium Tuberculosis from the Complete Genome Sequence." *Nature* 393 (6685): 537–44. https://doi.org/10.1038/31159.

Coll, Francesc, Ruth McNerney, José Afonso Guerra-Assunção, Judith R. Glynn, João Perdigão, Miguel Viveiros, Isabel Portugal, Arnab Pain, Nigel Martin, and Taane G. Clark. 2014. "A Robust SNP Barcode for Typing Mycobacterium Tuberculosis Complex Strains." *Nature Communications* 5 (1): 4812. https://doi.org/10.1038/ncomms5812.

Comas, Iñaki, Mireia Coscolla, Tao Luo, Sonia Borrell, Kathryn E Holt, Midori Kato-Maeda, Julian Parkhill, et al. 2013. "Out-of-Africa Migration and Neolithic Coexpansion of Mycobacterium Tuberculosis with Modern Humans." *Nature Genetics* 45 (10). https://doi.org/10.1038/ng.2744.

Coscolla, M, Daniela Brites, Fabrizio Menardo, Chloe Loiseau, Isaac Darko Otchere, Adwoa Asante-Poku, Prince Asare, et al. 2020. "Phylogenomics of Mycobacterium Africanum Reveals a New Lineage and a Complex Evolutionary History." *BioRxiv* 17 (June): 19. https://doi.org/10.1101/2020.06.10.141788.

Coscolla, Mireia, and Sebastien Gagneux. 2014. "Consequences of Genomic Diversity in Mycobacterium Tuberculosis." *Seminars in Immunology* 26 (6): 431–44. https://doi.org/10.1016/j.smim.2014.09.012.

Darko Otchere, Isaac, Mireia Coscollá, Leonor Sánchez-Busó, Adwoa Asante-Poku, Daniela Brites, Chloe Loiseau, Conor Meehan, et al. 2018. "Comparative Genomics of Mycobacterium Africanum Lineage 5 and Lineage 6 from Ghana Suggests Distinct Ecological Niches." *SCIENTIfIC RepoRtS* | 8: 11269. https://doi.org/10.1038/s41598-018-29620-2.

Darling, Aaron C E, Bob Mau, Frederick R. Blattner, and Nicole T. Perna. 2004. "Mauve: Multiple Alignment of Conserved Genomic Sequence with Rearrangements." *Genome Research* 14 (7): 1394–1403. https://doi.org/10.1101/gr.2289704.

DeJong, Bouke C, Philip C Hill, Alex Aiken, Timothy Awine, Martin Antonio, Ifedayo M Adetifa, Dolly J Jackson-Sillah, et al. 2008. "Progression to Active Tuberculosis, but Not Transmission, Varies by M. Tuberculosis Lineage in The Gambia." *J Infect Dis* 198 (7): 1037–43. https://doi.org/10.1086/591504.

Dick, Thomas, Ujjini Manjunatha, Barbara Kappes, and Martin Gengenbacher. 2010. "Vitamin B6 Biosynthesis Is Essential for Survival and Virulence of Mycobacterium Tuberculosis." *Molecular Microbiology* 78 (4): 980–88. https://doi.org/10.1111/j.1365-2958.2010.07381.x.

Firdessa, Rebuma, Stefan Berg, Elena Hailu, Esther Schelling, Balako Gumi, Girume Erenso, Endalamaw Gadisa, et al. 2013. "Mycobacterial Lineages Causing Pulmonary and

Extrapulmonary Tuberculosis, Ethiopia." *Emerging Infectious Diseases* 19 (3): 460–63. https://doi.org/10.3201/eid1903.120256.

Gagneux, Sebastien. 2018. "Ecology and Evolution of Mycobacterium Tuberculosis." *Nature Reviews Microbiology*. https://doi.org/10.1038/nrmicro.2018.8.

Gagneux, Sebastien, Kathryn Deriemer, Tran Van, Midori Kato-Maeda, Bouke C De Jong, Sujatha Narayanan, Mark Nicol, et al. 2006. "Variable Host-Pathogen Compatibility in Mycobacterium Tuberculosis." www.pnas.orgcgidoi10.1073pnas.0511240103.

Gao, Juehua, Jie Liao, and Guang-Yu Yang. 2009a. "Review Article CAAX-Box Protein, Prenylation Process and Carcinogenesis." *Am J Transl Res*. Vol. 1. www.ajtr.org/AJTR905003.

Gao, Juehua, Jie Liao, and Guang Yu Yang. 2009b. "CAAX-Box Protein, Prenylation Process and Carcinogenesis." *American Journal of Translational Research*.

Garimella, Kiran V, Zamin Iqbal, Michael A. Krause, Susana Campino, Mihir Kekre, Eleanor Drury, Dominic Kwiatkowski, Juliana M. Sa, Thomas E. Wellems, and Gil McVean. 2019. "Detection of Simple and Complex de Novo Mutations without, with, or with Multiple Reference Sequences." *BioRxiv*, July, 698910. https://doi.org/10.1101/698910.

Gehre, Florian, Jacob Otu, Kathryn DeRiemer, Paola Florez de Sessions, Martin L. Hibberd, Wim Mulders, Tumani Corrah, Bouke C. de Jong, and Martin Antonio. 2013. "Deciphering the Growth Behaviour of Mycobacterium Africanum." *PLoS Neglected Tropical Diseases* 7 (5). https://doi.org/10.1371/journal.pntd.0002220.

Ghodousi, Arash, Alamdar Hussain Rizvi, Aurangzaib Quadir Baloch, Abdul Ghafoor, Faisal Masood, Mehmood Qadir, Emanuele Borroni, Alberto Trovato, Sabira Tahseen, and Daniela Maria. 2019. "Acquisition of Cross-Resistance to Bedaquiline and Clofazimine Following Treatment for Tuberculosis in 2 Pakistan." https://doi.org/10.1101/630012.

Goig, Galo A., Silvia Blanco, Alberto L. Garcia-Basteiro, and Iñaki Comas. 2020. "Contaminant DNA in Bacterial Sequencing Experiments Is a Major Source of False Genetic Variability." *BMC Biology* 18 (1): 24. https://doi.org/10.1186/s12915-020-0748-z.

Goig, Galo A, Silvia Blanco, Alberto L. Garcia-Basteiro, and Iñaki Comas. 2019. "Contaminant DNA in Bacterial Sequencing Experiments Is a Major Source of False Genetic Variability." *BioRxiv*.

Goig, Galo A, Silvia Blanco, Alberto L Garcia-Basteiro, and Iñaki Comas. 2018. "Pervasive Contaminations in Sequencing Experiments Are a Major Source of False Genetic Variability: A Mycobacterium Tuberculosis Meta-Analysis." https://doi.org/10.1101/403824.

Gonzalo-Asensio, Jesús, Wladimir Malaga, Alexandre Pawlik, Catherine Astarie-Dequeker, Charlotte Passemar, Flavie Moreau, Françoise Laval, et al. 2014. "Evolutionary History of Tuberculosis Shaped by Conserved Mutations in the PhoPR Virulence Regulator." *Proceedings of the National Academy of Sciences of the United States of America* 111 (31): 11491–96. https://doi.org/10.1073/pnas.1406693111.

Gordon, Stephen V., Roland Brosch, Alain Billault, Thierry Garnier, Karin Eiglmeier, and Stewart T. Cole. 1999. "Identification of Variable Regions in the Genomes of Tubercle Bacilli Using Bacterial Artificial Chromosome Arrays." *Molecular Microbiology* 32 (3): 643–55. https://doi.org/10.1046/j.1365-2958.1999.01383.x.

Grinter, Rhys, Blair Ney, Rajini Brammananth, Christopher K. Barlow, Paul R.F. Cordero, David L. Gillett, Thierry Izoré, et al. 2020. "Cellular and Structural Basis of Synthesis of the Unique Intermediate Dehydro-F420-0 in Mycobacteria." *BioRxiv*, February, 2020.02.27.968891. https://doi.org/10.1101/2020.02.27.968891.

Gu, J, H Li, M Li, C Vuong, M Otto, Y Wen, and Q Gao. 2005. "Bacterial Insertion Sequence IS256 as a
Potential Molecular Marker to Discriminate Invasive Strains from Commensal Strains of
Staphylococcus Epidermidis." *The Journal of Hospital Infection* 61 (4): 342–48.
https://doi.org/10.1016/j.jhin.2005.04.017.

Herrmann, Jean Louis, Robin Delahay, Alex Gallagher, Brian Robertson, and Douglas Young. 2000.
"Analysis of Post-Translational Modification of Mycobacterial Proteins Using a Cassette
Expression System." *FEBS Letters* 473 (3): 358–62. https://doi.org/10.1016/S0014-
5793(00)01553-2.

Ingen, Jakko van, Thomas A. Kohl, Katharina Kranzer, Barbara Hasse, Peter M. Keller, Anna Katarzyna
Szafrańska, Doris Hillemann, et al. 2017. "Global Outbreak of Severe Mycobacterium Chimaera
Disease after Cardiac Surgery: A Molecular Epidemiological Study." *The Lancet Infectious
Diseases* 17 (10): 1033–41. https://doi.org/10.1016/S1473-3099(17)30324-9.

Intemann, Christopher D., Thorsten Thye, Stefan Niemann, Edmund N. L. Browne, Margaret Amanua
Chinbuah, Anthony Enimil, John Gyapong, et al. 2009. "Autophagy Gene Variant IRGM −261T
Contributes to Protection from Tuberculosis Caused by Mycobacterium Tuberculosis but Not by
M. Africanum Strains." Edited by William Bishai. *PLoS Pathogens* 5 (9): e1000577.
https://doi.org/10.1371/journal.ppat.1000577.

Iqbal, Zamin, Mario Caccamo, Isaac Turner, Paul Flicek, and Gil Mcvean. 2012. "De Novo Assembly
and Genotyping of Variants Using Colored de Bruijn Graphs." https://doi.org/10.1038/ng.1028.

Jandrasits, Christine, Stefan Kröger, Walter Haas, and Bernhard Y. Renard. 2019. "Computational
Pan-Genome Mapping and Pairwise SNP-Distance Improve Detection of Mycobacterium
Tuberculosis Transmission Clusters." Edited by Sergei L. Kosakovsky Pond. *PLOS Computational
Biology* 15 (12): e1007527. https://doi.org/10.1371/journal.pcbi.1007527.

Jong, Bouke C De, Martin Antonio, and Sebastien Gagneux. n.d. "Mycobacterium Africanum-Review
of an Important Cause of Human Tuberculosis in West Africa." Accessed May 21, 2019.
https://doi.org/10.1371/journal.pntd.0000744.

Kalyaanamoorthy, Subha, Bui Quang Minh, Thomas K F Wong, Arndt von Haeseler, and Lars S
Jermiin. 2017. "ModelFinder: Fast Model Selection for Accurate Phylogenetic Estimates."
Nature Methods 14 (6): 587–89. https://doi.org/10.1038/nmeth.4285.

Kapopoulou, Adamandia, Jocelyne M. Lew, and Stewart T. Cole. 2011. "The MycoBrowser Portal: A
Comprehensive and Manually Annotated Resource for Mycobacterial Genomes." *Tuberculosis*
91 (1): 8–13. https://doi.org/10.1016/j.tube.2010.09.006.

Kato-Maeda, Midori, Jeanne T Rhee, Thomas R Gingeras, Hugh Salamon, Jorg Drenkow, Nat
Smittipat, and Peter M Small. 2001. "Comparing Genomes within the Species Mycobacterium
Tuberculosis." https://doi.org/10.1101/gr166401.

Kavanagh, K L, H Jçrnvall, B Persson, and U Oppermann. 2008. "The SDR Superfamily: Functional and
Structural Diversity within a Family of Metabolic and Regulatory Enzymes" 171: 77.
https://doi.org/10.1007/s00018-008-8588-y.

Kohl, Thomas Andreas, Christian Utpatel, Viola Schleusener, Maria Rosaria De Filippo, Patrick
Beckert, Daniela Maria Cirillo, and Stefan Niemann. 2018. "MTBseq: A Comprehensive Pipeline
for Whole Genome Sequence Analysis of Mycobacterium Tuberculosis Complex Isolates."
https://doi.org/10.7717/peerj.5895.

Leao, Sylvia, Anandi Martin, Gloria Isabel Mejia, Francoise Portaels, and Por- Taels F Leao SC, Martin
A, Mejia GI, Palomino JC, Robledo J, Telles MAS. 2004. "Practical Handbook for the Phenotypic

and Genotypic Identification of Mycobacteria.," 77–126. https://doi.org/10.1007/BF01045823.

Lee, Robyn S., Jean François Proulx, Fiona McIntosh, Marcel A. Behr, and William P. Hanage. 2020.
"Previously Undetected Super-Spreading of Mycobacterium Tuberculosis Revealed by Deep
Sequencing." *ELife* 9 (February). https://doi.org/10.7554/eLife.53245.

Lee, Robyn S, and Marcel A Behr. 2016. "Does Choice Matter? Reference-Based Alignment for
Molecular Epidemiology of Tuberculosis." *Journal of Clinical Microbiology* 54 (7): 1891–95.
https://doi.org/10.1128/JCM.00364-16.

Lew, Jocelyne M., Adamandia Kapopoulou, Louis M. Jones, and Stewart T. Cole. 2011. "TubercuList -
10 Years After." *Tuberculosis* 91 (1): 1–7. https://doi.org/10.1016/j.tube.2010.09.008.

Li, Ruiqiang, Yingrui Li, Hancheng Zheng, Ruibang Luo, Hongmei Zhu, Qibin Li, Wubin Qian, et al.
2010. "Building the Sequence Map of the Human Pan-Genome." *Nature Biotechnology* 28 (1):
57–62. https://doi.org/10.1038/nbt.1596.

Magnus, K. 1966. "Epidemiological Basis of Tuberculosis Eradication 3. Risk of Pulmonary
Tuberculosis after Human and Bovine Infection *." *Bull. Org. Mond. Sante*. Vol. 35.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2476032/pdf/bullwho00607-0027.pdf.

Malone, Kerri M, Damien Farrell, Tod P Stuber, Olga T Schubert, Ruedi Aebersold, Suelee Robbe-
Austerman, and Stephen V Gordon. 2017. "Updated Reference Genome Sequence and
Annotation of Mycobacterium Bovis AF2122/97." *Genome Announcements* 5 (14): e00157-17.
https://doi.org/10.1128/genomeA.00157-17.

Marcus, Sarah A, Sarah W Sidiropoulos, Howard Steinberg, and Adel M Talaat. 2016. "CsoR Is
Essential for Maintaining Copper Homeostasis in Mycobacterium Tuberculosis."
https://doi.org/10.1371/journal.pone.0151816.

Maretty, Lasse, Jacob Malte Jensen, Bent Petersen, Jonas andreas sibbesen, siyang Liu, Palle Villesen,
Laurits skov, et al. 2017. "Sequencing and de Novo Assembly of 150 Genomes from Denmark as
a Population Reference." *Nature Publishing Group* 548. https://doi.org/10.1038/nature23264.

Marschall, Tobias, Manja Marz, Thomas Abeel, Louis Dijkstra, Bas E. Dutilh, Ali Ghaffaari, Paul
Kersey, et al. 2018. "Computational Pan-Genomics: Status, Promises and Challenges." *Briefings
in Bioinformatics* 19 (1): 118–35. https://doi.org/10.1093/bib/bbw089.

Martiniano, Rui, Erik Garrison, Eppie R Jones, Andrea Manica, and Richard Durbin. 2019. "Removing
Reference Bias in Ancient DNA Data Analysis by Mapping to a Sequence Variation Graph."
BioRxiv, September, 782755. https://doi.org/10.1101/782755.

McInerney, James O., Fiona J. Whelan, Maria Rosa Domingo-Sananes, Alan McNally, and Mary J.
O'Connell. 2020. "Pangenomes and Selection: The Public Goods Hypothesis." In *The
Pangenome*, 151–67. Springer International Publishing. https://doi.org/10.1007/978-3-030-
38281-0_7.

Medini, Duccio, Claudio Donati, Hervé Tettelin, Vega Masignani, and Rino Rappuoli. 2005. "The
Microbial Pan-Genome." *Current Opinion in Genetics and Development*. Elsevier Current
Trends. https://doi.org/10.1016/j.gde.2005.09.006.

Meehan, Conor J., Pieter Moris, Thomas A. Kohl, Jūlija Pečerska, Suriya Akter, Matthias Merker,
Christian Utpatel, et al. 2018. "The Relationship between Transmission Time and Clustering
Methods in Mycobacterium Tuberculosis Epidemiology." *EBioMedicine* 37 (November): 410–
16. https://doi.org/10.1016/j.ebiom.2018.10.013.

Meehan, Conor J, Galo A Goig, Thomas A Kohl, Lennert Verboven, Anzaan Dippenaar, Matthew

Ezewudo, Maha R Farhat, et al. 2019. "Whole Genome Sequencing of Mycobacterium
Tuberculosis: Current Standards and Open Issues." *Nature Reviews Microbiology*.
https://doi.org/10.1038/s41579-019-0214-5.

Merker, Matthias, Thomas A Kohl, Ivan Barilar, Sönke Andres, Philip W Fowler, Erja Chryssanthou,
Kristian Ängeby, et al. 2020. "Phylogenetically Informative Mutations in Genes Implicated in
Antibiotic Resistance in Mycobacterium Tuberculosis Complex." *Genome Medicine*.
https://doi.org/10.1186/s13073-020-00726-5.

Merle, Corinne S.C., Charalambos Sismanidis, Oumou B. Sow, Martin Gninafon, John Horton, Olivier
Lapujade, Mame B. Lo, et al. 2012. "A Pivotal Registration Phase III, Multicenter, Randomized
Tuberculosis Controlled Trial: Design Issues and Lessons Learnt from the Gatifloxacin for TB
(OFLOTUB) Project." *Trials* 13 (May). https://doi.org/10.1186/1745-6215-13-61.

Mostowy, Serge, Debby Cousins, Jacqui Brinkman, Alicia Aranaz, and Marcel A Behr. 2002. "Genomic
Deletions Suggest a Phylogeny for the Mycobacterium Tuberculosis Complex." *JID*.

Murugesan, Saravanan, Stalin Mani, Indhumathy Kuppusamy, and Padma Krishnan. 2018. "Role of
Insertion Sequence Element Is256 as a Virulence Marker and Its Association with Biofilm
Formation among Methicillin-Resistant Staphylococcus Epidermidis from Hospital and
Community Settings in Chennai, South India." *Indian Journal of Medical Microbiology* 36 (1):
124. https://doi.org/10.4103/ijmm.IJMM_17_276.

Nebenzahl-Guimaraes, Hanna, Solomon A. Yimer, Carol Holm-Hansen, Jessica De Beer, Roland
Brosch, and Dick Van Soolingen. 2016. "Genomic Characterization of Mycobacterium
Tuberculosis Lineage 7 and a Proposed Name: 'Aethiops Vetus.'" *Microbial Genomics* 2 (6).
https://doi.org/10.1099/mgen.0.000063.

Ngabonziza, Jean Claude Semuto, Chloé Loiseau, Michael Marceau, Agathe Jouet, Fabrizio Menardo,
Oren Tzfadia, Esdras Belamo Niyigena, et al. 2020. *A Sister Lineage of the Mycobacterium
Tuberculosis Complex Discovered in the African Great Lakes Region*. *BioRxiv*.
https://doi.org/10.1101/2020.01.20.912998.

Niemann, S., T. Kubica, F. C. Bange, O. Adjei, E. N. Browne, M. A. Chinbuah, R. Diel, et al. 2004. "The
Species Mycobacterium Africanum in the Light of New Molecular Markers." *Journal of Clinical
Microbiology* 42 (9): 3958–62. https://doi.org/10.1128/JCM.42.9.3958-3962.2004.

Norman, Anders, Dorte Bek Folkvardsen, Søren Overballe-Petersen, and Troels Lillebaek. 2019.
"Complete Genome Sequence of Mycobacterium Tuberculosis DKC2, the Predominant Danish
Outbreak Strain." https://doi.org/10.1128/MRA.

O'Toole, Ronan F., and Sanjay S. Gautam. 2017. "Limitations of the Mycobacterium Tuberculosis
Reference Genome H37Rv in the Detection of Virulence-Related Loci." *Genomics* 109 (5–6):
471–74. https://doi.org/10.1016/j.ygeno.2017.07.004.

Ofori-Anyinam, Boatema, Fatoumatta Kanuteh, Schadrac C. Agbla, Ifedayo Adetifa, Catherine Okoi,
Gregory Dolganov, Gary Schoolnik, et al. 2016. "Impact of the Mycobaterium Africanum West
Africa 2 Lineage on TB Diagnostics in West Africa: Decreased Sensitivity of Rapid Identification
Tests in The Gambia." *PLoS Neglected Tropical Diseases* 10: 1–12.
https://doi.org/10.1371/journal.pntd.0004801.

Ofori-Anyinam, Boatema, Abi Janet Riley, Tijan Jobarteh, Ensa Gitteh, Binta Sarr, Tutty Isatou Faal-
Jawara, Leen Rigouts, et al. 2020. "Comparative Genomics Shows Differences in the Electron
Transport and Carbon Metabolic Pathways of Mycobacterium Africanum Relative to
Mycobacterium Tuberculosis and Suggests an Adaptation to Low Oxygen Tension." *Tuberculosis*
120 (January): 101899. https://doi.org/10.1016/j.tube.2020.101899.

Ondov, Brian D., Todd J. Treangen, Páll Melsted, Adam B. Mallonee, Nicholas H. Bergman, Sergey
Koren, and Adam M. Phillippy. 2016. "Mash: Fast Genome and Metagenome Distance
Estimation Using MinHash." *Genome Biology* 17 (1): 132. https://doi.org/10.1186/s13059-016-
0997-x.

Paten, Benedict, Adam M. Novak, Jordan M. Eizenga, and Erik Garrison. 2017. "Genome Graphs and
the Evolution of Genome Inference." *Genome Research*. Cold Spring Harbor Laboratory Press.
https://doi.org/10.1101/gr.214155.116.

Periwal, Vinita, Ashok Patowary, Shamsudheen Karuthedath Vellarikkal, Anju Gupta, Meghna Singh,
Ashish Mittal, Shamini Jeyapaul, et al. 2015. "Comparative Whole-Genome Analysis of Clinical
Isolates Reveals Characteristic Architecture of Mycobacterium Tuberculosis Pangenome."
https://doi.org/10.1371/journal.pone.0122979.

Phelan, Jody, Paola Florez De Sessions, Leopold Tientcheu, Joao Perdigao, Diana Machado, Rumina
Hasan, Zahra Hasan, et al. 2018. "Methylation in Mycobacterium Tuberculosis Is Lineage
Specific with Associated Mutations Present Globally." *Scientific Reports* 8 (1): 1–7.
https://doi.org/10.1038/s41598-017-18188-y.

Rademacher, Corinna, and Bernd Masepohl. 2012. "Copper-Responsive Gene Regulation in
Bacteria." https://doi.org/10.1099/mic.0.058487-0.

Rakocevic, Goran, Vladimir Semenyuk, Wan Ping Lee, James Spencer, John Browning, Ivan J.
Johnson, Vladan Arsenijevic, et al. 2019. "Fast and Accurate Genomic Analyses Using Genome
Graphs." *Nature Genetics* 51 (2): 354–62. https://doi.org/10.1038/s41588-018-0316-4.

Reiling, Norbert, Susanne Homolka, Kerstin Walter, Julius Brandenburg, Lisa Niwinski, Martin Ernst,
Christian Herzmann, et al. 2013. "Clade-Specific Virulence Patterns of Mycobacterium
Tuberculosis Complex Strains in Human Primary Macrophages and Aerogenically Infected
Mice." *MBio* 4 (4). https://doi.org/10.1128/mBio.00250-13.

Rengarajan, Jyothi, Barry R Bloom, and Eric J Rubin. 2005. "Genome-Wide Requirements for
Mycobacterium Tuberculosis Adaptation and Survival in Macrophages." *PNAS June*. Vol. 7.
www.pnas.orgcgidoi10.1073pnas.0503272102.

Riojas, Marco A, Katya J Mcgough, Cristin J Rider-Riojas, Nalin Rastogi, and Manzour Hernando
Hazbón. 2018. "Phylogenomic Analysis of the Species of the Mycobacterium Tuberculosis
Complex Demonstrates That Mycobacterium Africanum, Mycobacterium Bovis, Mycobacterium
Caprae, Mycobacterium Microti and Mycobacterium Pinnipedii Are Later Heterotypic
Synonyms of Mycob." https://doi.org/10.1099/ijsem.0.002507.

Rowland, Jennifer L, and Michael Niederweis. 2012. "Resistance Mechanisms of Mycobacterium
Tuberculosis against Phagosomal Copper Overload."
https://doi.org/10.1016/j.tube.2011.12.006.

S.R. Pattyn, F. Portaels, L. Spanoghe, and J. Magos. 1970. "Further Studies on African Strains of
Mycobacterium Tuberculosis. Comparison with M. Bovis and M. Microti." Ann. Soc. belge Méd.
Trop. https://www.ncbi.nlm.nih.gov/pubmed/4996738.

Samanovic, Marie I., Chen Ding, Dennis J. Thiele, and K. Heran Darwin. 2012. "Copper in Microbial
Pathogenesis: Meddling with the Metal." *Cell Host & Microbe* 11 (2): 106–15.
https://doi.org/10.1016/j.chom.2012.01.009.

Sanoussi, C. N'Dira, Dissou Affolabi, Leen Rigouts, Séverin Anagonou, and Bouke de Jong. 2017.
"Genotypic Characterization Directly Applied to Sputum Improves the Detection of
Mycobacterium Africanum West African 1, under-Represented in Positive Cultures." *PLOS*

Neglected Tropical Diseases 11 (9): e0005900. https://doi.org/10.1371/journal.pntd.0005900.

Sanoussi, C. N'Dira, Bouke C De Jong, Mathieu Odoun, Karamatou Arekpa, Moulikatou Ali Ligali, Ousman Bodi, Simon Harris, et al. 2018. "Low Sensitivity of the MPT64 Identification Test to Detect Lineage 5 of the Mycobacterium Tuberculosis Complex." https://doi.org/10.1099/jmm.0.000846.

Seemann, T. 2014. "Prokka: Rapid Prokaryotic Genome Annotation." *Bioinformatics* 30 (14): 2068–69. https://doi.org/10.1093/bioinformatics/btu153.

Selengut, Jeremy D, and Daniel H Haft. 2010. "Unexpected Abundance of Coenzyme F 420-Dependent Enzymes in Mycobacterium Tuberculosis and Other Actinobacteria †." *JOURNAL OF BACTERIOLOGY* 192 (21): 5788–98. https://doi.org/10.1128/JB.00425-10.

Sellés Vidal, Lara, Ciarán L. Kelly, Paweł M. Mordaka, and John T. Heap. 2018. "Review of NAD(P)H-Dependent Oxidoreductases: Properties, Engineering and Application." *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* 1866 (2): 327–47. https://doi.org/10.1016/j.bbapap.2017.11.005.

Sherman, Rachel M., Juliet Forman, Valentin Antonescu, Daniela Puiu, Michelle Daya, Nicholas Rafaels, Meher Preethi Boorgula, et al. 2019. "Assembly of a Pan-Genome from Deep Sequencing of 910 Humans of African Descent." *Nature Genetics*. Nature Publishing Group. https://doi.org/10.1038/s41588-018-0273-y.

Szklarczyk, Damian, John H. Morris, Helen Cook, Michael Kuhn, Stefan Wyder, Milan Simonovic, Alberto Santos, et al. 2017. "The STRING Database in 2017: Quality-Controlled Protein-Protein Association Networks, Made Broadly Accessible." *Nucleic Acids Research* 45 (D1): D362–68. https://doi.org/10.1093/nar/gkw937.

Tatusov, Roman L, Michael Y Galperin, Darren A Natale, and Eugene V Koonin. 2000. "The COG Database: A Tool for Genome-Scale Analysis of Protein Functions and Evolution." *Nucleic Acids Research*. Vol. 28. http://www.ncbi.nlm.nih.gov/COG.

Tettelin, Hervé, Vega Masignani, Michael J. Cieslewicz, Claudio Donati, Duccio Medini, Naomi L. Ward, Samuel V. Angiuoli, et al. 2005. "Genome Analysis of Multiple Pathogenic Isolates of Streptococcus Agalactiae: Implications for the Microbial 'Pan-Genome.'" *Proceedings of the National Academy of Sciences of the United States of America* 102 (39): 13950–55. https://doi.org/10.1073/pnas.0506758102.

Thye, Thorsten, Stefan Niemann, Kerstin Walter, Susanne Homolka, Christopher D. Intemann, Margaret Amanua Chinbuah, Anthony Enimil, et al. 2011. "Variant G57E of Mannose Binding Lectin Associated with Protection against Tuberculosis Caused by Mycobacterium Africanum but Not by M. Tuberculosis." *PLoS ONE* 6 (6). https://doi.org/10.1371/journal.pone.0020908.

Tsolaki, Anthony G., Sebastien Gagneux, Alexander S. Pym, Yves Olivier L. Goguet De La Salmoniere, Barry N. Kreiswirth, Dick Van Soolingen, and Peter M. Small. 2005. "Genomic Deletions Classify the Beijing/W Strains as a Distinct Genetic Lineage of Mycobacterium Tuberculosis." *Journal of Clinical Microbiology* 43 (7): 3185–91. https://doi.org/10.1128/JCM.43.7.3185-3191.2005.

Walker, Timothy M, Camilla LC Ip, Ruth H Harrell, Jason T Evans, Georgia Kapatai, Martin J Dedicoat, David W Eyre, et al. 2013. "Whole-Genome Sequencing to Delineate Mycobacterium Tuberculosis Outbreaks: A Retrospective Observational Study." *The Lancet Infectious Diseases* 13 (2): 137–46. https://doi.org/10.1016/S1473-3099(12)70277-3.

Ward, Sarah K, Elizabeth A Hoye, and Adel M Talaat. 2008. "The Global Responses of Mycobacterium Tuberculosis to Physiological Levels of Copper †." *JOURNAL OF BACTERIOLOGY* 190 (8): 2939–

46. https://doi.org/10.1128/JB.01847-07.

WHO. 2016. "THE SHORTER MDR-TB REGIMEN FEATURES OF THE SHORTER MDR-TB REGIMEN
REGIMEN COMPOSITION 4-6 Km-Mfx-Pto-Cfz-Z-H High-Dose-E / 5 Mfx-Cfz-Z-E."
www.who.int/tb.

———. 2019. *WHO Consolidated Guidelines on Drug-Resistant Tuberculosis Treatment*.
http://apps.who.int/bookorders.

Zhang, Shuo, Jiazhen Chen, Peng Cui, Wanliang Shi, Wenhong Zhang, and Ying Zhang. 2015.
"Identification of Novel Mutations Associated with Clofazimine Resistance in Mycobacterium
Tuberculosis." https://doi.org/10.1093/jac/dkv150.

944 **Figure 1. Methodology for finding gene presence-absence between two genomes**

Locus region A　　**Locus region B**

Figure 2. Visualization of possible rearrangements using Mauve pipeline: H37Rv (L4) genome and three completed PacBio genomes of *M. africanum* L5 isolates from Benin, The Gambia and Nigeria.

The region circled in red (one on the H37Rv genome (missing region A), and one on the Nigerian completed PacBio genome (missing region B)) represents the regions absent in that strain genome but present in the other three strains complete genomes.

Figure 3. Gene count difference between three completed PacBio genomes of *M. africanum* L5 from
Benin, The Gambia and Nigeria: L5Ben, L5Gam and L5Nig. The numbers are the count/number of
gene difference (present or absent) between genomes. The sign (+) indicates the presence of genes
while (-) indicates the absence of genes. For example the "- 2 +" gene difference between L5Gam and
L5Ben, indicates that 2 genes are present in the L5Ben genome but absent in the L5Gam genome. 5,
2, 1 are the number of genes only found in respectively L5Ben, L5Gam, L5Nig.

Figure 4. Gene count difference between *M. tuberculosis* H37Rv (L4) genome and three completed PacBio genomes of *M. africanum* L5 isolates from Benin, The Gambia and Nigeria. * These 10 genes are present in each of the three L5 completed PacBio genomes, and absent in H37RV.

H37Rv (L4) genes [A*]

PcbL5Ben genes [B*]

L5 Illumina genomes

1003

PcbL5Gam genes [C*]

PcbL5Nig genes [D]

L5 Illumina genomes

Figure 5. Mapping of 205 clinical (Illumina platform sequenced) genomes (L5_ig) of *M. africanum* L5 from various countries to *M. tuberculosis* H37Rv (L4)

and three *M. africanum* L5 isolates (completed PacBio genomes) from Benin, The Gambia and Nigeria as reference (PcbL5Ben; PcbL5Gam; PcbL5Nig).

(A) **5**/9 genes present in H37Rv but absent in the three L5_cg were **absent in all 205 L5_ig.**

(B) **5**/5 genes **present in PcbL5Ben but absent in the two other L5 completed PacBio genomes** were **present in 159-199/205 L5_ig.**

(C) **2**/2 genes **present in PcbL5Gam but absent in the two other L5 completed PacBio genomes** were **present in 195 & 199/205 L5_ig.**

(D) **1**/1 gene **present in PcbL5Nig but absent in the two other L5 completed PacBio genomes** was **present in 195/205 L5_ig.**

* The 32-33 genes absent in PcbL5Nig but present in PcbL5Ben, PcbL5Gam and *H37Rv* were also absent in 6/205 L5_ig (Fig.9).

.... **or** **....** gene absent

1021

1022

1023

1024

1025

1026

1027

1028

1029

* Region B= region of 30 genes (two blocks of 18 and 12 genes: *Rv1493* through *Rv1509* and *Rv1511* through *Rv1521*) missing in L5.3.2 strains but present in all other L5 strains (L5.1, L5.2, L5.3.1 and new sublineages).

Figure 6. Phylogenetic tree showing the 6 L5 isolates Illumina genomes (L5.3.2) similar to the completed PacBio genome of the Nigerian L5 strain (PcbL5Nig) and the position of the other two L5 PacBio genomes (PcbL5Ben and PcbL5Gam)

1063

Table 1. Gene content difference between *M. tuberculosis* H37Rv (L4) genome and completed PacBio genomes of three L5 isolates from Benin, The Gambia and Nigeria.

		Present in			
		PcbL5Ben	**PcbL5Gam**	**PcbL5Nig**	**H37Rv**
Absent in	**PcbL5Ben**		2^B	1^C	9^D
	PcbL5Gam	$2 + 5^A$		1^C	$2 + 9^D$
	PcbL5Nig	$33 + 5^A$	$33 + 2^B$		$32 + 9^D$
	H37Rv	$10^* + 5^A$	$10^* + 2^B$	$10^* + 1^C$	
		4189	4162	4134	4126

1066

1067 **A:** include 5 only present in PcbL5Ben

1068 **B:** includes 2 only present in PcbL5Gam

1069 **C:** includes 1 only present in PcbL5Nig

1070 **D:** includes 9 only present in H37Rv

1071 *: includes 10 genes shared by the PcbL5 (Ben, Gam, Nig) and absent in H37Rv

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

Table 2. Mapping of 205 L5 isolates Illumina genomes to the *M. tuberculosis* H37Rv (L4) genome and completed genomes of three L5 isolates from Benin, The Gambia and Nigeria (mapping statistics/estimates). The best mapping results (numbers) are written in bold. When the best mapping result is obtained for PcbL5Nig as the reference, the next best result is also written in bold (as PcbL5Nig compared to the other two PcbL5 genomes missed a 30 genes region).

	H37Rv	PcbL5Ben	PcbL5Gam	PcbL5Nig
Reads				
Mean of Percent L5 reads mapped to	96.9	**97.5**	97.3	96.5
Mean of unambiguous coverage mean	121.9	122.7	**122.8**	122.5
Bases				
Mean of Percent unambiguous total bases	0.983	0.988	**0.989**	0.982
Mean uncovered	30444.8	18801.7	**14657.5**	35215.4
Mean SNP	2210	529.9	**513.2**	**503.7**
Mean deletions	373.3	95.9	**95.7**	97.2
Mean insertions	239.2	**77.3**	107.6	**60.8**
Mean substitutions (including stop codons)	1193	**0**	**0**	**0**
Genes				
Mean of Percent gene mapped (presence)	99.59	99.72	**99.79**	99.76
L5 illumina having all the ref. genome genes % (n=205)	0	2 (4)	7.3 (15)	0
Mean gene count difference between L5 Illumina genomes (gene count per Illumina-sequenced genome minus minimum gene count)	29.8	**38.3**	33.4	33

Table 3. Presence in 205 L5 isolates Illumina genomes of genes detected in only one of the three L5 completed genomes from Benin, Nigeria or The Gambia

	Gene	Present in L5 Illumina genomes % (n=205)
PcbL5Ben	PcbL5_01893	**77.6** (159)
	PcbL5_01894	**77.6** (159)
	PcbL5_01895	**77.6** (159)
	PcbL5_02043	**95.1** (195)
	PcbL5_03043	**97.1** (199)
PcbL5Gam	PcbL5_02028	**95.1** (195)
	PcbL5_03020	**97.1** (199)
PcbL5Nig	PcbL5_02001	**95.1** (195)

1099 **Table 4. Comparison of transmission clustering rates based on choice of reference genome.** Short read data from 355 L5 isolates were mapped against
1100 each of the four reference genomes for SNP calling. Distance matrices between all isolates were constructed per reference approach and transmission
1101 clusters were defined based on specific SNP cut-offs

1 SNP

Reference used	In transmission cluster	Percentage of total dataset	Number of clusters
H37rv	100	28.17	44
PcbL5Ben	94	26.48	43
PcbL5Gam	95	26.76	43
PcbL5Nig	95	26.76	43

5 SNP

Reference used	In transmission cluster	Percentage of total dataset	Number of clusters
H37rv	129	36.34	53
PcbL5Ben	124	34.93	51
PcbL5Gam	124	34.93	51
PcbL5Nig	124	34.93	51

12 SNP

Reference used	In transmission cluster	Percentage of total dataset	Number of clusters
H37rv	144	40.56	55
PcbL5Ben	141	39.72	54
PcbL5Gam	144	40.56	55
PcbL5Nig	141	39.72	54

1102

1103