Functional characterization of SARS-CoV-2 infection suggests a complex inflammatory

response and metabolic alterations

Lucía Trilla-Fuertes [1], Ricardo Ramos [2], Natalia Blanca-López [3], Elena López-Camacho [1], Laura

Martín-Pedraza [3], Pablo Ryan Murua [5], Mariana Díaz-Almirón [6], Carlos Llorens [7], Toni

Gabaldón[8,9,10], Andrés Moya [11,12,13], Juan Ángel Fresno Vara[1,14,15], and Angelo Gámez-Pozo [1,14*]

[1] Biomedica Molecular Medicine SL, Madrid, Spain.

[2] Genomics Unit, Parque Científico de Madrid, Madrid, Spain.

[3] Allergy Service, Infanta Leonor University Hospital, 28031, Madrid, Spain.

[4] Allergy Laboratory, Infanta Leonor University Hospital, 28031, Madrid, Spain

[5] Internal Medicine Service, Infanta Leonor University Hospital, 28031, Madrid, Spain

[6] Biostatistics Unit, Hospital Universitario La Paz, Madrid, Spain.

[7] Biotechvana SL, Valencia, Spain.

[8] Barcelona Supercomputing Centre (BSC-CNS). Jordi Girona, 29. 08034. Barcelona, Spain.

[9] Institute for Research in Biomedicine (IRB Barcelona), The Barcelona Institute of Science and

Technology, Baldiri Reixac, 10, 08028 Barcelona, Spain

[10] Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain

[11] Institute for Integrative Systems Biology, University of València and Consejo Superior de

Investigaciones Científicas, València, Spain.

[12] Fundación para el Fomento de la Investigación Sanitaria y Biomédica de la Comunidad

Valenciana, (FISABIO), València, Spain.

[13] CIBER en Epidemiología y Salud Pública (CIBEResp), Madrid, Spain.

[14] Molecular Oncology and Pathology Lab, Hospital Universitario La Paz, IDIPAZ, Madrid, Spain.

[15] Biomedical Research Networking Center on Oncology-CIBERONC, ISCIII, Madrid, Spain.

*Corresponding author

Abstract

Covid-19, caused by the SARS-CoV-2 virus, has reached the category of a worldwide pandemic.
Even though intensive efforts, no effective treatments or a vaccine are available. Molecular
characterization of the transcriptional response in Covid-19 patients could be helpful to
identify therapeutic targets. In this study, RNAseq data from peripheral blood mononuclear
cell samples from Covid-19 patients and healthy controls was analyzed from a functional point
of view using probabilistic graphical models. Two networks were built: one based on genes
differentially expressed between healthy and infected individuals and another one based on
the 2,000 most variable genes in terms of expression in order to make a functional
characterization. In the network based on differentially expressed genes, two inflammatory
response nodes with different tendencies were identified, one related to cytokines and
chemokines, and another one related to bacterial infections. In addition, differences in
metabolism, which were studied in depth using Flux Balance Analysis, were identified. SARS-
CoV2- infection caused alterations in glutamate, methionine and cysteine, and
tetrahydrobiopterin metabolism. In the network based on 2,000 most variable genes, also two
inflammatory nodes with different tendencies between healthy individuals and patients were
identified. Similar to the other network, one was related to cytokines and chemokines.
However, the other one, lower in Covid-19 patients, was related to allergic processes and self-
regulation of the immune response. Also, we identified a decrease in T cell node activity and
an increase in cell division node activity. In the current absence of treatments for these
patients, functional characterization of the transcriptional response to SARS-CoV-2 infection
could be helpful to define targetable processes. Therefore, these results may be relevant to
propose new treatments.

Author Summary

SARS-CoV-2 infection caused Covid-19 which has reached the category of a worldwide
pandemic. However, no treatments or vaccines are still available. For this reason, it is still
necessary the molecular study of this disease. In this study, we reanalyzed data from
peripheral blood mononuclear cells from Covid-19 patients and healthy controls using
computational techniques that allow the study of differential biological processes and
metabolic pathways. The results suggested a complex inflammatory response, involving genes
related to response to bacterial infection and allergic processes, and alterations in metabolic
pathways such as glutamate metabolism, cysteine and methionine metabolism or

tetrahydrobiopterin metabolism. These processes could be used in the future as therapeutic

targets in Covi-19 infection.

Introduction

The emerging coronavirus SARS-CoV-2 has rapidly expanded from its origin in Wuhan, China, to become a worldwide pandemic only after four months since its first identification. At June 2nd of 2020, 6,194,533 cases and 376,320 deaths have been reported worldwide, according to the World Health Organization [1].

The most common symptoms are fever, cough, fatigue, shortness of breath, accompanied by elevated inflammatory biomarkers and pulmonary infiltrates. However, during the SARS-CoV-2 infection, a fraction of patients will develop severe pneumonia, pulmonary oedema, severe acute respiratory syndrome (SARS) or multiple organ failure, ending in death [2]. These severe symptoms are associated with systemic inflammation related to an overproduction of macrophagic cytokines. Different treatments focused on these inflammatory processes are being investigated [3].

Recently, Xiong et al. analyzed the transcriptional response in samples from peripheral blood mononuclear cells (PBMCs) from patients diagnosed with SARS-CoV-2 and compared them with healthy controls. Based on the results, the authors suggested that patient's lymphopenia may be caused by an activation of apoptosis in lymphocytes, and also that SARS-Cov-2 induced excessive cytokine production, which correlates with lung tissue injury [4]. However, these conclusions were based only in the functional enrichment analysis of differentially expressed genes.

Probabilistic graphical models (PGMs) have demonstrated their utility in analyzing gene expression data by identifying relevant biological processes [5, 6]. These models allow making associations between genes according to their expression patterns across a series. Interestingly, the PGM networks have functional structure, allowing study expression data from a functional point of view. We thus set out to re-analyze Xiong. et. al. data using PGMs, aiming for a deeper understanding of biological processes involved in SARS-CoV-2 pathogenesis.

Results

Processing of RNA sequencing data

After alignment of raw files, 13,398 expressed genes were identified. After applying the quality criteria of a detectable reading in at least 50% of the samples, 13,182 genes were used for the subsequent analyses.

Analysis of differential genes between healthy controls and patients

Using CuffDiff, 1,569 differentially expressed genes were determined between SARS-CoV-2 patients and healthy controls. After applying the quality criteria of detectable measurements in at least 50% of the samples, 1,234 genes remained as differential ones. These genes were mostly related to inflammatory response, innate immune response, T cells, lysosomes, apoptotic processes and angiogenesis, among others.

A PGM was used to organize these genes according to their biological functions. The resulting network was composed by eight functional nodes: metabolism, lysosomes, T cells, two nodes related to inflammatory response, two nodes related to response to virus, and one node with no overrepresented function (Fig 1, S1 File).

Fig 1: Probabilistic graphical model based on the expression of 1,234 differential genes between healthy individuals and patients.

Inflammatory response A, lysosome ad metabolism functional nodes activities were significantly differential between healthy controls and patients in the data were shown. Strikingly, one of the nodes of inflammatory response had a higher functional node activity in healthy controls than in patients, and the other node of inflammatory response had a functional node activity higher in patients than healthy controls. The same tendencies were shown in response to virus nodes. Lysosome and metabolism had a higher functional node activity in patients than in controls. Finally, T cell functional node activity was higher in healthy individuals than in patients (Fig 2).

Fig 2: Functional node activities from the network based on the expression of the 1,234 genes defined as significantly differential by CuffDiff. In the Y axis the activity of the functional node in arbitrary units, understanding as the mean expression of those genes in each node that were related to the overrepresented function in the node. In the x axis, healthy controls and Covid-19 patients. **, ≤ 0.01 ; * ≤ 0.05.

Metabolism functional node

Metabolism node was composed of 102 genes, 19 of them related with metabolism pathways. This node contained the genes PKM (pyruvate kinase) and PDHB (pyruvate dehydrogenase), both implicated on glycolysis, and several ATPases from the mitochondrial complex, responsible for H+ transporting. There were also genes related to drug metabolism such as PAPSS1 or CES1.

Inflammatory response A functional node

This node included 236 genes, of which 18 were implicated in an inflammatory response. This node mostly comprised chemokines.

Inflammatory response B functional node

This node included 129 genes, of which 17 were involved in inflammatory response. This functional node included three toll-like receptors, TLR2, TLR5 and TLR4. Among other functions, TLR2 promotes apoptosis in response to bacterial lipoproteins. TLR5 protein recognizes bacterial flagellin, the principal component of bacterial flagella. Additionally, TLR4 has been implicated in signal transduction events induced by lipopolysaccharide found in most gran negative bacteria. This node also contained AOAH and CD14, both genes implicated in response to bacterial lipopolysaccharides as well. Finally, this functional node included VNN1 which plays a suppressive role in influenza virus replication in human alveolar epithelial cells [7]. Therefore, this node is mostly related to the response to bacterial infections.

Functional characterization based on the 2,000 most variable genes

We obtained an alternative PGM network, now based on the 2,000 most variable genes according to their SD and functionally characterized. The resulting network was divided into nine functional nodes: apoptosis, oxygen binding, blood coagulation, response to the virus, T cell, cell division, and three nodes related to an inflammatory response (Fig 3, S2 File).

Fig 3: Probabilistic graphical model based on the expression of the 2,000 most variable genes.

Cell division and inflammatory response B functional node activities were differential between controls and patients. On the one hand, patients had a higher activity of response to virus, cell division, and one of the inflammatory functional nodes. On the other hand, healthy individuals had a higher activity of T cell and two out of the three functional nodes related to inflammatory processes (Fig 4).

146 **Fig 4: Functional node activities from the network based on the expression of the 2,000 most**

147 **variable genes.** In the Y axis the activity of the functional node in arbitrary units,

148 understanding as the mean expression of those genes in each node that were related to the

149 overrepresented function. In the X axis, healthy controls and Covid-19 patients. ***, ≤ 0.001;

150 ** , ≤ 0.01 ; * ≤ 0.05.

Inflammatory response A functional node

152 This node is composed of 77 genes, of which 28 were related to an inflammatory response.

153 Most of these genes were cytokines, chemokines and toll-like receptors, whose function is the

154 modulation of the inflammatory response. This functional node included toll-like receptors

155 TLR6, TLR8, TLR5, TLR1, and TLR4.

Inflammatory response B functional node

157 This node was formed by 69 genes, 6 of them related to the inflammatory response. These six

158 genes were CCR3, CCL4L2, TNFRSF18, NCR3, CCL5, and MS4A2. CCL5 and CCL4L2 are

159 chemokine ligands, and CCR3 and MS4A2 are implicated in an allergic response.

Inflammatory response C functional node

161 This functional node had 77 genes, 6 of them related to inflammatory processes. These genes

162 were CCL4, CXCR2, FPR2, IL1RAP, CXCL8, and ORM1, mostly of them chemokines.

T cell functional node

164 This functional node was composed of 210 genes, 6 of them related to T cells, more concretely

165 with T cell receptors, including GATA3 gene, which plays a vital role in nasopharyngeal virus

166 detection.

Cell division functional node

168 This node was composed of 141 genes, 5 of them involved in cell division. These genes were

169 CDK1, CENPW, CCNB1, UBE2C, and CCNB2, mostly related to M-phase promoting factor

170 complex and microtubules.

Response to virus functional node

172 This node had 130 genes, nine related to response to virus ontology. This node contained two

173 genes whose proteins are induced by interferon, IFI44L and IFITM3.

Metabolic modeling

FBA was performed to characterize metabolic alterations caused by SARS-CoV-2 infection (S3

File). Glutamate metabolism, methionine and cysteine metabolism, and tetrahydrobiopterin

metabolism flux activities were differential between healthy controls and Covid-19 patients

(Fig 5). In addition, several tendencies in other metabolic pathways such as TCA cycle and

steroid metabolism that need to be confirmed were shown (S1 Figure).

Fig 5: Differential flux activities between healthy controls and patients. a.u. = arbitrary units.

* $p < 0.05$.

Discussion

SARS-CoV-2 infection has reached the category of a pandemic. Tremendous efforts have been made to find a suitable vaccine and to determine effective treatments but till the date there are neither of them [8].

We have re-analyzed the work of Xiong et al. [4] with a different functional inference approach. Coincidences between both results were expected. Xiong et al. described an up-regulation of genes related to cell cycle and cytokines in SARS-CoV-2 patients, which agreed with the higher functional node activity that we observed in cell division node and one of the inflammatory nodes, mostly composed by cytokines and chemokines. They also described a reduction of immune cells in blood patient samples, which may be related to the lower activity of T cell node in patients than in healthy controls.

Additionally, our analysis offered complementary information. For instance, in the network that characterizes differences between healthy individuals and SARS-CoV-2 patients, there were two functional nodes related to inflammatory response. Strikingly, inflammatory response A functional node activity was higher in patients than in healthy controls. This node was composed by cytokines and chemokines. However, inflammatory response B node activity was higher in healthy controls than in patients and it was related to response to bacterial infections. SARS-CoV-2 can carry with a bacterial co-infection of *Mycoplasma pneumoniae* so the study of those genes in relation to the presence of a bacterial co-infection may be relevant [9]. Metabolism node had also a higher functional node activity in Covid-19 patients than in healthy controls. The increase in glycolysis reactions implies an increase in Krebs cycle reactions as well and therefore in ATP production, essential for the virus replication [10, 11]. The differences on metabolism functional node suggested that a deeper analysis of metabolism could supply more detail information. Glutamate metabolism showed differences between controls and Covid-19 patients. Interestingly, an alteration caused by another RNA virus, the HIV-1, has been previously described [11]. Moreover, it has been previously suggested that methionine plays a relevant role in viral replication of other coronaviruses [12]. No alterations in tetrahydrobiopterin metabolism have been previously described.

On the other hand, in the network obtained for the 2,000 most variable genes, two functional nodes related to inflammatory response were also identified. Inflammatory response A functional node was again mostly composed by cytokines and chemokines. Inflammatory response B functional node had a lower activity in SARS-CoV-2 patients. Interestingly, the inflammatory response B node was composed of genes related to allergic response and

regulation of immunological self-tolerance. This fact may be related to the severe acute

respiratory syndrome, associated with a dysregulation of the immune response [2].

In this inflammatory response B node is included CCR3, a chemokine highly expressed in

eosinophils and basophils, and is also detected in TH1 and TH2 cells, as well in airway epithelial

cells [13, 14]. This receptor may contribute to the accumulation and activation of inflammatory

cells in allergic airway and it is also known to be an entry co-receptor for HIV-1. MS4A2 is also

implicated in allergic processes [15]. Therefore, this node seems to be more related to the self-

control of the inflammatory response instead of the other inflammatory functional nodes,

more related to chemokines and cytokines.

Cell division functional node had a significantly higher activity in Covid-19 patients than in

controls. This node is mainly composed by genes related to M-phase and mitosis process,

which may be related to viral infection. An accumulation of G2/M phase cells in other

coronaviruses has been previously described in order to promote favorable conditions for viral

replication [16].

As expected, functional nodes related to response to the virus were relevant in both networks.

In the case of the network built based on the 2,000 most variable genes, this functional node

was mainly related to interferon response. Remarkably, this node included IFITM3 gene,

whose genetics is associated with immunity to other well-known viruses such as influenza A or

dengue virus [17, 18]. IFITM3 protein has been described as related to the entry of MERS- CoV

and SARS-CoV[19]. The first response to a viral infection of the immune system is mediated by

interferons so it seems logical that these genes were overexpressed in patients infected by

SARS-CoV-2.

In addition, in the T cell functional node appeared GATA3 gene which has been previously

related to nasopharyngeal virus infections [20]. Since SARS-CoV-2 presents mainly respiratory

tropism, GATA3 may play an essential role.

Our study had some limitations. Most importantly, the reduced number of samples severely

limited the statistical power and the information that could be obtained by functional

analyses. A larger number of samples will be useful to deepen in the molecular

characterization of this disease. Also, studying a larger cohort stratified according the severity

of the disease could be of much interest as it may help define how functional modules vary in

relation to the virulence of the infection.

In this study, some previously not described relevant processes in SARS-CoV-2 pathogenesis

such as bacterial inflammatory response processes, metabolism or allergic processes, were

proposed. In the absence of treatments for these patients, molecular characterization of the

disease could be helpful to define targetable processes. Therefore, these results may be

relevant to propose new therapeutic treatments.

Materials and Methods

Patient cohort

Three samples from PBMCs from three patients infected with SARS-CoV-2 and three samples from healthy controls were analyzed. These samples are all from the work of Xiong et al. [4].

Processing of RNA sequencing data

Before processing fragments per kilobase of exon model per million of reads (FPKM) data, we checked their quality using FastQC (v0.11.9, Brabaham, UK). Reads longer than 100 nt showed the presence of Illumina adapter sequences which were removed by trimming using Prinseq [21]so all samples were matched to 2x100 format. Then, reads were mapped against the human genome (GRCh38.96) using TopHat, using an estimated paired-end inner size of 25 and finally FPKM data were obtained using CuffDiff. All these programs were accessed using the integrated GPRO suite (Biotechvana, Valencia, Spain) [22].

After FPKM processing, Perseus v1.6.5 software was used to filter RNAseq data [23]. Log2 was calculated and only those genes with at least 50% of the detectable readings were used for the subsequent analyses.

Probabilistic graphical models

2,000 most variable genes were selected according to their standard deviation (SD) of expression across the series and used to build a PGM network. RNAseq expression data was used without other a priori information.

The resulting network was divided into functional nodes by gene ontology analyses. These gene ontology analyses were performed in DAVID webtool v8 using "Homo sapiens" as background and KEGG, Biocarta and GOTERM-FAT as categories [24].

The same analysis pipeline was used to characterize the differential genes defined by CuffDiff, i.e. a network was built using the genes defined as significantly differential between healthy controls and patients.

These analyses was done using *grapHD* package [25] and R v3.2.5. Network visualization was done in Cytoscape [26]. PGMs were built in two steps, first, the spanning tree with the maximum likelihood was found, and then, the edges were refined based on the minimization of the Bayesian Information Criterion (BIC) [27].

Statistical analysis

Functional node activities were calculated as previously described [6]. Briefly, the mean

expression of those genes of each node related to the overrepresented function in this node

was calculated. Then, functional node activities were compared between healthy individuals

and patients using a T-test.

Flux Balance Analysis and metabolic models

Flux Balance Analysis (FBA) allows metabolic modeling from gene expression data. It is widely

used in microbiology and cancer [28]. The complete human metabolic reconstruction Recon 3D

was used to perform these analyses. It contains 10,600 reactions, 5,835 metabolites and 5,939

Gene-Protein-Reaction rules (GPRs), which contain information in the form of Boolean

expressions about what genes are involved in each metabolic reaction. GPRs were solved using

a modification of Barker et al. algorithm [29, 30], solving "AND" expressions as the minimum

and "OR" expressions as the sum. Then, the obtained values were introduced as the reaction

bounds by a modified E-flux algorithm based on the Max-min function [30, 31]. Finally, FBA

was solved using COBRA Toolbox library v2.0 [32] and MATLAB.

The 10,600 metabolic reactions are grouped into 103 metabolic pathways or subsystems. In

order to compare metabolic activity between controls and Covid-19 patients, flux activities

were calculated as previously described as the sum of fluxes of the reactions contained in a

concrete metabolic pathway [5, 33]. To compare flux activities between control and patients a

T-test was used.

References

1.	WHO Reports https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/.

2.	Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. Lancet. 2020;395(10223):507-13. Epub 2020/01/30. doi: 10.1016/S0140-6736(20)30211-7. PubMed PMID: 32007143; PubMed Central PMCID: PMCPMC7135076.

3.	Li G, Fan Y, Lai Y, Han T, Li Z, Zhou P, et al. Coronavirus infections and immune responses. J Med Virol. 2020;92(4):424-32. Epub 2020/02/07. doi: 10.1002/jmv.25685. PubMed PMID: 31981224; PubMed Central PMCID: PMCPMC7166547.

4.	Xiong Y, Liu Y, Cao L, Wang D, Guo M, Jiang A, et al. Transcriptomic characteristics of bronchoalveolar lavage fluid and peripheral blood mononuclear cells in COVID-19 patients. Emerg Microbes Infect. 2020;9(1):761-70. doi: 10.1080/22221751.2020.1747363. PubMed PMID: 32228226; PubMed Central PMCID: PMCPMC7170362.

5.	Trilla-Fuertes L, Gámez-Pozo A, Arevalillo JM, Díaz-Almirón M, Prado-Vázquez G, Zapater-Moros A, et al. Molecular characterization of breast cancer cell response to metabolic drugs. Oncotarget. 2018;9(11):9645-60. Epub 2018/01/08. doi: 10.18632/oncotarget.24047. PubMed PMID: 29515760; PubMed Central PMCID: PMCPMC5839391.

6.	Gámez-Pozo A, Berges-Soria J, Arevalillo JM, Nanni P, López-Vacas R, Navarro H, et al. Combined label-free quantitative proteomics and microRNA expression analysis of breast cancer unravel molecular differences with clinical implications. Cancer Res; 2015. p. 2243-53.

7.	Yamashita N, Yashiro M, Ogawa H, Namba H, Nosaka N, Fujii Y, et al. Metabolic pathway catalyzed by Vanin-1 pantetheinase plays a suppressive role in influenza virus replication in human alveolar epithelial A549 cells. Biochem Biophys Res Commun. 2017;489(4):466-71. Epub 2017/05/30. doi: 10.1016/j.bbrc.2017.05.172. PubMed PMID: 28576495.

8.	Kruse RL. Therapeutic strategies in an outbreak scenario to treat the novel coronavirus originating in Wuhan, China. F1000Res. 2020;9:72. Epub 2020/01/31. doi: 10.12688/f1000research.22211.2. PubMed PMID: 32117569; PubMed Central PMCID: PMCPMC7029759.

9.	Fan BE, Lim KGE, Chong VCL, Chan SSW, Ong KH, Kuperan P. COVID-19 and mycoplasma pneumoniae coinfection. Am J Hematol. 2020;95(6):723-4. Epub 2020/04/03. doi: 10.1002/ajh.25785. PubMed PMID: 32173883.

10.	Gerresheim GK, Roeb E, Michel AM, Niepmann M. Hepatitis C Virus Downregulates Core Subunits of Oxidative Phosphorylation, Reminiscent of the Warburg Effect in Cancer Cells. Cells. 2019;8(11). Epub 2019/11/08. doi: 10.3390/cells8111410. PubMed PMID: 31717433; PubMed Central PMCID: PMCPMC6912740.

11.	Hegedus A, Kavanagh Williamson M, Khan MB, Dias Zeidler J, Da Poian AT, El-Bacha T, et al. Evidence for Altered Glutamine Metabolism in Human Immunodeficiency Virus Type 1 Infected Primary Human CD4. AIDS Res Hum Retroviruses. 2017;33(12):1236-47. Epub 2017/10/04. doi: 10.1089/AID.2017.0165. PubMed PMID: 28844150; PubMed Central PMCID: PMCPMC5709700.

12.	Chen Y, Su C, Ke M, Jin X, Xu L, Zhang Z, et al. Biochemical and structural insights into the mechanisms of SARS coronavirus RNA ribose 2'-O-methylation by nsp16/nsp10 protein complex. PLoS Pathog. 2011;7(10):e1002294. Epub 2011/10/13. doi: 10.1371/journal.ppat.1002294. PubMed PMID: 22022266; PubMed Central PMCID: PMCPMC3192843.

13.	Beck LA, Tancowny B, Brummet ME, Asaki SY, Curry SL, Penno MB, et al. Functional analysis of the chemokine receptor CCR3 on airway epithelial cells. J Immunol. 2006;177(5):3344-54. doi: 10.4049/jimmunol.177.5.3344. PubMed PMID: 16920975.

351 14. Khanolkar A, Burden SJ, Hansen B, Wilson AR, Philipps GJ, Hill HR. Evaluation of CCR3
352 as a basophil activation marker. Am J Clin Pathol. 2013;140(3):293-300. doi:
353 10.1309/AJCPLSN0RQKHJX1A. PubMed PMID: 23955446.

354 15. Guo H, Peng T, Luo P, Li H, Huang S, Li S, et al. Association of FcεRIβ polymorphisms
355 with risk of asthma and allergic rhinitis: evidence based on 29 case-control studies. Biosci Rep.
356 2018;38(4). Epub 2018/07/31. doi: 10.1042/BSR20180177. PubMed PMID: 29654163; PubMed
357 Central PMCID: PMCPMC6066650.

358 16. Dove B, Brooks G, Bicknell K, Wurm T, Hiscox JA. Cell cycle perturbations induced by
359 infection with the coronavirus infectious bronchitis virus and their effect on virus replication. J
360 Virol. 2006;80(8):4147-56. doi: 10.1128/JVI.80.8.4147-4156.2006. PubMed PMID: 16571830;
361 PubMed Central PMCID: PMCPMC1440480.

362 17. Prabhu SS, Chakraborty TT, Kumar N, Banerjee I. Association between IFITM3 rs12252
363 polymorphism and influenza susceptibility and severity: A meta-analysis. Gene. 2018;674:70-9.
364 Epub 2018/06/22. doi: 10.1016/j.gene.2018.06.070. PubMed PMID: 29940276.

365 18. Allen EK, Randolph AG, Bhangale T, Dogra P, Ohlson M, Oshansky CM, et al. SNP-
366 mediated disruption of CTCF binding at the IFITM3 promoter is associated with risk of severe
367 influenza in humans. Nat Med. 2017;23(8):975-83. Epub 2017/07/17. doi: 10.1038/nm.4370.
368 PubMed PMID: 28714988; PubMed Central PMCID: PMCPMC5702558.

369 19. Zhao X, Sehgal M, Hou Z, Cheng J, Shu S, Wu S, et al. Identification of Residues
370 Controlling Restriction versus Enhancing Activities of IFITM Proteins on Entry of Human
371 Coronaviruses. J Virol. 2018;92(6). Epub 2018/02/26. doi: 10.1128/JVI.01535-17. PubMed
372 PMID: 29263263; PubMed Central PMCID: PMCPMC5827390.

373 20. Jartti T, Palomares O, Waris M, Tastan O, Nieminen R, Puhakka T, et al. Distinct
374 regulation of tonsillar immune response in virus infection. Allergy. 2014;69(5):658-67. Epub
375 2014/03/29. doi: 10.1111/all.12396. PubMed PMID: 24684577; PubMed Central PMCID:
376 PMCPMC7159333.

377 21. Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets.
378 Bioinformatics. 2011;27(6):863-4. Epub 2011/01/28. doi: 10.1093/bioinformatics/btr026.
379 PubMed PMID: 21278185; PubMed Central PMCID: PMCPMC3051327.

380 22. Futami R, Muñoz-Pomer A, Viu J, Domínguez-Escribá R, Covelli L, Bernet G, et al. GPRO
381 The professional tool for annotation, management and functional analysis of omic databases.
382 Biotechvana Bioinformatics: SOFT3. 2011.

383 23. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The Perseus
384 computational platform for comprehensive analysis of (prote)omics data. Nat Methods.
385 2016;13(9):731-40. Epub 2016/06/27. doi: 10.1038/nmeth.3901. PubMed PMID: 27348712.

386 24. Huang dW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene
387 lists using DAVID bioinformatics resources. Nat Protoc. 2009;4(1):44-57. doi:
388 10.1038/nprot.2008.211. PubMed PMID: 19131956.

389 25. Abreu G, Edwards D, Labouriau R. High-Dimensional Graphical Model Search with
390 the gRapHD R Package Journal of Statistical Software2010. p. 1-18.

391 26. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a
392 software environment for integrated models of biomolecular interaction networks. Genome
393 Res. 2003;13(11):2498-504. doi: 10.1101/gr.1239303. PubMed PMID: 14597658; PubMed
394 Central PMCID: PMCPMC403769.

395 27. Lauritzen S. Graphical Models. Oxford,UK.: Oxford University Press1996.

396 28. Orth J, Thiele I, Palsson B. What is flux balance analysis? : Nat Biotechnol; 2010. p. 245-
397 8.

398 29. Barker BE, Sadagopan N, Wang Y, Smallbone K, Myers CR, Xi H, et al. A robust and
399 efficient method for estimating enzyme complex abundance and metabolic flux from
400 expression data. Comput Biol Chem. 2015;59 Pt B:98-112. Epub 2015/09/01. doi:
401 10.1016/j.compbiolchem.2015.08.002. PubMed PMID: 26381164; PubMed Central PMCID:
402 PMCPMC4684447.

30.   Gámez-Pozo A, Trilla-Fuertes L, Berges-Soria J, Selevsek N, López-Vacas R, Díaz-Almirón
M, et al. Functional proteomics outlines the complexity of breast cancer molecular subtypes.
Scientific Reports. 2017;7(1):10100. doi: 10.1038/s41598-017-10493-w.
31.   Colijn C, Brandes A, Zucker J, Lun D, Weiner B, Farhat M, et al. Interpreting expression
data with metabolic flux models: Predicting Mycobacterium tuberculosis mycolic acid
production. PLOS Comput Bio; 2009.
32.   Schellenberger J, Que R, Fleming R, Thiele I, Orth J, Feist A, et al. Quantitative
prediction of cellular metabolism with constraint-based models: the COBRA Toolbox v2.0.
Nature Protocols; 2011. p. 1290-307.
33.   Trilla-Fuertes L, Gámez-Pozo A, Díaz-Almirón M, Prado-Vázquez G, Zapater-Moros A,
López-Vacas R, et al. Computational metabolism predicts risk of distant relapse-free survival in
breast cancer patients. Future Oncology. 2019;30:3483-90. doi: 10.2217/fon-2018-0698.

Funding

GPRO is supported by TSI-100903-2019-11 from Secretaría de Estado de Digitalización e

Inteligencia Artificial, Ministerio de Asuntos Económicos y Transformación Digital. This work

was supported by grants to AM from the Spanish Ministry of Economy and Competitiveness

(SAF2015-65878-R) and Generalitat Valenciana (Prometeo/2018/A/133), and co-financed by

the European Regional Development Fund (ERDF). LT-F is supported by the Spanish Economy

and Competitiveness Ministry (DI-15-07614). EL-C is supported by the Spanish Economy and

Competitiveness Ministry (PTQ2018-009760).

Supporting information

S1 File: Genes included in the probabilistic graphical model based on the expression of 1,234

differential genes between healthy individuals and patients.

S2 File: Genes included the probabilistic graphical model based on the expression of the 2,000

most variable genes.

S3 File: Flux Balance Analysis results.

Inflammatory response A

Metabolism

Response to virus A

T cell

Lysosome

Inflammatory response B

Response to virus B

Without function

- ■ Apoptosis
- ■ Oxygen binding
- ■ Blood coagulation
- ■ Response to virus
- ■ Inflammatory response A
- ■ T cell
- ■ Inflammatory response B
- ■ Cell division
- ■ Inflammatory response C

Glutamate metabolism

Methionine metabolism

Tetrahydrobiopterin metabolism

TCA

Steroid metabolism