The COPII components Sec23 and Sec24 form isoform

specific subcomplexes with Sec23D/E and Sec24C/D

essential for tip growth

Mingqin Chang[1,2,3], Shu-Zon Wu[1], Jacquelyn E. O'Sullivan[4,5] and Magdalena

Bezanilla[1]*

[1] Department of Biological Sciences, Dartmouth College, Hanover, NH 03755

[2] Plant Biology Graduate Program, University of MA Amherst, Amherst, MA

01002

[3] Current address: Department of Plant Sciences, UC Davis, Davis, CA 95616

[4] Department of Biology, University of MA Amherst, Amherst, MA 01002

[5] Current address: Tufts School of Medicine, Boston, MA 02111

*Corresponding author

email: magdalena.bezanilla@dartmouth.edu

Abstract

COPII, a coat of proteins that form vesicles on the ER, mediates vesicle

traffic from the ER to the Golgi. In contrast to metazoans that have few genes

encoding each COPII component, plants have expanded these gene families

leading to the hypothesis that plant COPII has functionally diversified. Here, we

analyzed the gene families encoding for the Sec23/24 heterodimer in the moss

Physcomitrium (Physcomitrella) patens. In *P. patens*, *Sec23* and *Sec24* gene

families are each comprised of seven genes. We discovered that *Sec23D* is

functionally redundant with *Sec23E* and together they are essential for

protonemal development. Sec23D is critical for secretion to the Golgi and the

plasma membrane in tip growing protonemata. Sec23D preferentially associates

with Sec24C/D, also essential for protonemata. In contrast, the remaining five

Sec23 genes are dispensable for tip growth, but are redundant with *Sec23D/E* for

gametophore formation. While *Sec23A/B/C/F/G* do not quantitatively affect

Golgi secretion in protonemata, they do contribute to secretion to the plasma

membrane. Furthermore, the three highly expressed *Sec23* genes in

protonemata, *Sec23B/D/G* form morphologically distinct ER exit sites. These data

suggest that *Sec23D/E* form unique ER exit sites contributing to secretion that is

essential for tip growing protonemata.

Introduction

Trafficking from the ER to the Golgi is mediated by the Coat Protein complex II (COPII). COPII facilitates the formation of transport vesicles at ER exit sites promoting the delivery of cargo, including transmembrane proteins, soluble proteins and lipids, to *cis*-Golgi compartments (Brandizzi and Barlowe, 2013)(Barlowe and Helenius, 2016) (Aridor, 2018)(Hanna et al., 2018). The COPII coat comprises two layers of protein complexes. The inner layer consists of a Sar1-Sec23-Sec24 lattice, while the outer layer is made up of Sec31-Sec13 heterotetramers (Stagg et al., 2008; Whittle and Schwartz, 2010; Bi et al., 2002; Fath et al., 2007). Sar1 is a small GTPase of the Ras super family that is the master regulator of COPII vesicle biogenesis (Nakano and Muramatsu, 1989). Assembly of COPII coat is initiated by the conversion of Sar1-GDP to Sar1-GTP, which is mediated by Sec12, a guanine nucleotide exchange factor (GEF) that is an ER resident membrane protein (Barlowe and Schekman, 1993; Nakano and Muramatsu, 1989). At the ER surface, activated Sar1 recruits the Sec23-Sec24 heterodimer, which associates with the ER membrane as an "inner coat". The Sar1-Sec23-Sec24-cargo "pre-budding" complex in turn recruits the Sec13-Sec31 heterotetramer, referred to as an "outer coat" to the nascent bud (Aridor et al., 1998; Salama et al.). The polymerization of Sec13-Sec31 heterotetramers induces membrane curvature and completes vesicle biogenesis (Fath et al., 2007; Stagg et al., 2006; Whittle and Schwartz, 2010).

Selective recruitment of cargos into COPII coated vesicles is driven by ER export signals located on cytoplasmically exposed regions of cargo proteins (Barlowe, 2003). ER export signals of most transmembrane cargo proteins are thought to interact directly with the inner coat components of COPII. However, some transmembrane and most soluble cargo proteins interact with COPII components through ER-resident transmembrane adaptors or receptors (Barlowe, 2003). Sec24 is the COPII component that has been implicated in recognizing these various sorting signals (Mossessova et al., 2003; Nufer, 2003). However, several studies suggested that Sec23 and Sar1 may also play roles in cargo sorting by directly binding to the export signals (Aridor et al., 2001; Mancias and Goldberg, 2007; Giraudo and Maccioni, 2003).

Based on the high sequence similarity among COPII paralogs, and cross-
species complementation studies (De Craene et al., 2014; Khoriaty et al., 2018) it
is assumed that functions of COPII components are conserved across eukaryotes.
While components of the COPII coat are well studied in budding yeast and
mammalian cells, illustrating their molecular functions in plants has been more
challenging. Comparative genomic analyses revealed that gene families encoding
COPII components have undergone gene expansions in plants with Sec23
exhibiting the largest expansion (Aridor et al., 2001; Mancias and Goldberg,
2007; Giraudo and Maccioni, 2003). While there are only one and two Sec23
genes in *S. cerevisiae* and mammals (eg. mice and humans) (Jensen and
Schekman, 2011), respectively, there are seven in both *P. patens* and *A. thaliana*
(Brandizzi, 2018). Since Sec23 and its heterodimeric partner Sec24 are
implicated in cargo sorting, the larger Sec23 gene family could suggest that
plants have more diverse sorting signals requiring specificity at the level of the
Sec23/Sec24 heterodimer. Consistent with this, in contrast to yeast, which has a
single gene for most COPII components, all other components of COPII in plants
are encoded by multiple genes, suggesting that cargo specificity may translate to
formation of distinct COPII complexes and vesicle populations. Alternatively,
larger gene families could simply indicate that plants have higher functional
redundancy in COPII encoded gene families.
Genetic studies in *A. thaliana* have demonstrated that loss of specific
COPII components caused developmental defects. A complete knockout of
AtSec24A was lethal (Faso et al., 2009; Nakano et al., 2009). However, a missense
point mutation in AtSec24A (R693K) resulted in ER morphology defects and
disruption of ER-Golgi integrity. Interestingly, these phenotypes could not be
rescued by either AtSec24B or AtSec24C (Faso et al., 2009; Nakano et al., 2009),
suggesting that AtSec24B and C are functionally distinct from AtSec24A. Even
though AtSec24B and AtSec24C did not rescue the *Atsec24a* missense mutant
and each were not important for plant viability, they were shown to influence the
development of reproductive cells (Tanaka et al., 2013). Functional
diversification was also suggested by analysis of AtSec16 mutants in which
storage protein precursors were accumulated abnormally in *Atsec16a* null
mutants, but not in *Atsec16b* null mutants (Takagi et al., 2013).

While *P. patens* and *A. thaliana* have similar numbers of genes encoding
COPII components, moss has several attributes that make it particularly suited to
study COPII function. *P. patens* juvenile tissues are haploid and comprise
multiple tissue types that establish the plant body. Protonemal tissue, which
emerges from the spore and is also readily propagated vegetatively by tissue
homogenization, is composed of tip-growing cells constituting a two-
dimensional filamentous network. The predictable patterns of growth and cell
division provide an easily tractable system for analysis of mutant phenotypes. As
a single-cell-layer, protonemal tissue also enables high-resolution microscopy of
endogenously tagged fluorescent proteins in live cells (Rensing et al., 2020). In
addition to amenable tissue morphology and haploid lifestyle, *P. patens* exhibits
high rates of efficient homologous recombination and CRISPR Cas9-mediated
genome editing (Rensing et al., 2020)r. Here, we utilized these molecular genetic
and live-cell imaging tools to systematically characterize the *Sec23* gene family.
Analysis of mutant phenotypes and protein localization revealed a high degree of
functional specificity among Sec23 isoforms, differentially influencing secretion
to the Golgi and the plasma membrane as well as the size of presumptive ER exit
sites. We also found that Sec23D only forms heterodimers with a subset of Sec24
isoforms, suggesting that COPII complexes may have fixed isoform compositions,
specified early on by the distinct Sec23/Sec24 heterodimer that forms the inner
coat.
136
Results
Specific Sec23 Isoforms Are Required For Tip Growth
P. patens has seven *Sec23* genes, which group into four subclasses based
on amino acid sequence similarity and gene structure (Fig1A). All seven *Sec23*
genes are expressed in protonemal tissue (Fig1B). With the exception of *Sec23A*,
at least one member of each subclass is highly expressed (Fig 1B). To determine
if Sec23 plays a role in protonemal growth, we used transient RNA interference
(RNAi) to silence the *Sec23* gene family. We selected regions of the coding
sequences from *Sec23A*, *Sec23C*, *Sec23D* and *Sec23G* to silence each of the *Sec23*
subclasses (Fig1A, solid lines under gene models). These sequence regions share
a high degree of sequence similarity to the other gene in each subclass (Fig 1A,

dashed lines under gene models, Table S1) and would effectively simultaneously

silence both genes. By co-transforming the Sec23G RNAi construct together with

a construct containing the target sequences for *Sec23A*, *C* and *D*, we targeted the

entire *Sec23* gene family and discovered that silencing the seven *Sec23* genes

leads to a dramatic reduction in growth (Fig 1C and D). Surprisingly, we found

that silencing subclass III comprised of *Sec23D* and *Sec23E* resulted in a similarly

severe growth defect (Fig 1C and D). In contrast, silencing any of the other

subclasses did not affect growth (Fig 1C and D). These results demonstrated that

there is functional specificity among *Sec23* isoforms, with only subclass III

playing a critical role in protonemal growth, while the remaining five *Sec23*

isoforms appear to be dispensable.

To test whether *Sec23D* and *Sec23E* within subclass III are functionally

redundant, we designed RNAi constructs targeting the untranslated regions

(UTRs) of each gene (Fig 2A).Targeting the untranslated regions of both *Sec23D*

and *Sec23E* resulted in small unpolarized plants with an 85% reduction in plant

area as compared to control RNAi plants (Fig 2B and C). Silencing *Sec23D* alone

resulted in a 59% reduction in plant area, while silencing *Sec23E* did not affect

plant area (Fig 2B and C). Given that *Sec23D* is expressed seven times higher

than *Sec23E* (Fig 1B), it is possible that *Sec23D* contributes to protonemal growth

more than *Sec23E*. To test whether *Sec23E* can replace *Sec23D* function if

expressed at sufficient levels, we performed complementation studies by driving

the full-length coding sequence of *Sec23D* or *Sec23E* with the constitutive maize

ubiquitin promoter. Interestingly, the reduction in plant area resulting from

silencing of *Sec23D* was restored to similar levels when either *Sec23D* or *Sec23E*

full-length cDNAs were expressed (Fig 2B and C), indicating that *Sec23D* and

Sec23E are functionally redundant, and that the highly expressed *Sec23D* is the

predominant group III isoform critical for protonemal growth.

CRISPR-Cas9-mediated deletion of Sec23 isoforms

Using transient RNAi, we were able to deduce that *Sec23* genes display

isoform specific functions during protonemal growth. However transient RNAi

assays are limited to phenotypic characterization of 7-day-old plants. To further

characterize the molecular functions of *Sec23*, we generated stable loss-of-

function mutants of *Sec23* genes using CRISPR-Cas9 mediated genome editing.

We targeted each gene with one protospacer located in the first coding exon (Fig

S1). We obtained single and higher-order mutants of *Sec23* genes carrying out-

of-frame mutations resulting in premature stop codons in their transcripts (Fig

S1A, Table S2). Consistent with the RNAi results, 7-day-old *Δsec23d* plants were

70% smaller than wild type plants while *Δsec23e* plants exhibited no growth

defects (Fig 3A and B). We isolated a number of *sec23d* null alleles (Fig S1A,

Table S2), which had similar growth phenotypes (Figure S1B). We also found

that in three-week-old plants regenerated from protoplasts, *Δsec23d-5* plants

were 62.6% smaller than wild type (Fig 3C and D), and *Δsec23d-5* had fewer

gametophores compared to wild type (Fig 3C and E), suggesting that slow

growing protonemal tissues resulted in a developmental delay in the production

of gametophores.

Although it was possible to isolate many *Δsec23d-5* and *Δsec23e* single

mutants, we were unable to obtain *Δsec23de* double mutants either by targeting

Sec23D and *Sec23E* simultaneously or by targeting *Sec23D* in the *Δsec23e* single

mutant background. This suggests that deleting both *Sec23D* and *Sec23E* is

incompatible with protoplast regeneration, consistent with the severe growth

defects observed by simultaneously silencing *Sec23D* and *Sec23E* using RNAi.

Furthermore, we confirmed that none of the other *Sec23* genes were upregulated

in *Δsec23d-5* plants (Fig S1C), suggesting that the phenotypic consequences

observed were due specifically to loss of *Sec23D*.

Since the RNAi results suggested that all other *Sec23* subclasses were not

required for protonemal growth, we sought to isolate a stable line lacking

Sec23A, *Sec23B*, *Sec23C*, *Sec23F*, and *Sec23G* to determine if these genes

contribute to other aspects of plant growth and development. We successfully

isolated two independent quintuple mutant alleles *Δsec23abcfg-1* and

Δsec23abcfg-2, in which *Sec23A*, *Sec23B*, *Sec23C*, *Sec23F*, and *Sec23G* contained

lesions resulting in null alleles for each gene (Fig S3, Table S2). Most lesions were

small deletions or insertions. However, *Δsec23abcfg-1* contained a three

thousand base pair insertion in the *Sec23G* locus, still resulting in disruption of

the *Sec23G* locus (Fig S3, Table S2). We characterized both quintuple mutants

and found that while seven-day old *Δsec23abcfg-1* plants exhibited no detectable

growth defects, *Δsec23abcfg-2* plants exhibited a mild growth defect (Fig 3B and

C)*. However, three-week-old *Δsec23abcfg-1* plants were larger than wild type,

and *Δsec23abcfg-2* plants were indistinguishable from wild type (Fig 3D and E).

Interestingly, both *Δsec23abcfg-1* and *Δsec23abcfg-2* developed more

gametophores than wild type (Fig 3E), suggesting that these five *Sec23* genes

may negatively regulate gametophore development.

Even though the two quintuple mutant isolates, exhibited statistically

significant differences at different times in development between each other and

wild type, these were minor compared to the differences between wild type and

all isolated *Δsec23d* null mutants. To investigate whether the observed

differences might result from differential upregulation of *Sec23D* of *Sec23E* in the

Δsec23abcfg mutants thereby compensating for the loss of *Sec23ABCFG*

functions, we measured the expression levels of *Sec23D* and *Sec23E* transcripts

in *Δsec23abcfg-1, Δsec23abcfg-2*, and wild type*. In 7-day-old plants, we found

that neither *Sec23D* nor *Sec23E* transcripts were elevated compared to

expression in wild type (Fig 3F), indicating that minor differences observed

between the two quintuple mutants were not readily explained by increased

expression of either *Sec23D* or *E*. Taken together, the phenotypic analyses of

stable *sec23* null plants suggest that *Sec23ABCF* and *G* do not contribute

substantially to establishment of juvenile moss tissues under laboratory growth

conditions. In contrast, *Sec23D* and *E* are essential.

Loss of Sec23D causes ER morphology defects and ER stress

As a predicted member of the COPII complex, Sec23D likely mediates ER-

to-Golgi transport. However, it is unclear whether the *Δsec23d* growth defect is a

result of a general impairment of ER-to-Golgi transport, or the lack of specific

cargo sorted by Sec23D needed for protonemal growth. Studies in other

organisms have shown that blocking COPII function alters ER morphology

(Novick et al., 1980) and results in elevated levels of unfolded proteins in the ER,

leading to activation of the ER unfolded protein response (Belden, 2001; Chung

et al., 2016). To analyze Sec23D's function in ER-to-Golgi transport, we

investigated whether loss of Sec23D affects ER morphology and the ER unfolded

protein response. We used CRISPR-Cas9 mediated genome editing to knock out

Sec23D in a line where the ER is fluorescently labeled by targeting GFP to the ER

lumen (Fig S1, Table S2). When imaged with a confocal microscope, *Δsec23d-B11*

cells exhibited numerous ER dots (Fig 4A, red and cyan boxes), and 65% of

protonemal filaments contained large ER aggregates (Fig 4A, green and yellow

boxes), suggesting that loss of Sec23D causes abnormal accumulations of ER

domains, leading to severe ER morphology defect in *Δsec23d* cells.

If changes in ER morphology result from defective secretion at ER exit

sites, then proteins may be inadvertently accumulating in the ER, resulting in

upregulation of the unfolded protein response. To test this, we analyzed the

expression of an ER heat shock protein (Bip), which is a well-known ER stress

reporter in *Arabidopsis thaliana* (Noh et al., 2003; Srivastava et al., 2013;

Maruyama et al., 2014; Cho and Kanehara, 2017). *P. patens* has two *Bip* paralogs,

Bip1 and *Bip2*. Since *Bip2* expression is low in protonemal tissue (Ortiz-Ramírez

et al., 2017), we tested whether *Bip1* is affected under conditions that induce ER

stress. Brefeldin A (BFA) inhibits protein secretion by disrupting ER-Golgi

integrity (Nebenfuhr, 2002; Niu et al., 2005). We found that after a 24-hour

treatment of BFA, expression of *Bip1* increased by about two-fold (Fig 4B),

indicating that *Bip1* is a plausible ER stress reporter in *P. patens*. Interestingly,

Bip1 expression increased 5-fold in *Δsec23d* compared to in wild type (Fig 4C),

demonstrating that *Δsec23d* suffers from severe ER stress. In contrast,

Δsec23abcfg-1, which didn't exhibit significant growth defects, had normal levels

of *Bip1* expression (Fig 4C), and normal ER morphology (Fig 4A).

Disrupting Sec23D results in secretion defects to the Golgi apparatus and the

plasma membrane

ER morphology defects and elevated ER stress in *Δsec23d* suggest that

transport of secretory cargos might be blocked. We reasoned that proteins

destined to the Golgi and the plasma membrane would accumulate in the ER in

mutants with impaired ER to Golgi transport. To test this, we disrupted *Sec23D*

in lines that either express a YFP fusion of a Golgi resident protein (YFP-Golgi) or

an mCherry fusion of a plasma membrane protein (F-SNAP-mCherry) (Nelson et

al., 2007; Furt et al., 2012; van Gisbergen et al., 2018) (Fig S1, Table S2). Since it

would have been time-consuming to regenerate quintuple mutants in these two

backgrounds, we instead transformed the YFP-Golgi and F-SNAP-mCherry

markers into *Δsec23abcfg-1* plants. We selected lines that had similar expression

levels to control lines by measuring the total amount of fluorescence in cells.

In control YFP-Golgi lines, YFP signal accumulated in the Golgi with very

little fluorescence observed in the rest of the cell (Fig 5A). In contrast in *Δsec23d-*

S2, while YFP fluorescence still accumulated in the Golgi, a large portion of the

fluorescent signal that was presumably retained in the ER was observed

diffusely throughout the cell (Fig 5A). To quantify Golgi secretion efficiency, we

calculated the fluorescence intensity ratio of the signal in the Golgi divided by the

signal in the cell. A high ratio value indicates efficient secretion, as observed in

the control lines (Fig 5B). In contrast, *Δsec23d-S2* exhibited a significantly lower

fluorescence intensity ratio (Fig 5B), indicating that loss of Sec23D impairs

secretion to the Golgi. Imaging of *Δsec23abcfg-1* YFP-Golgi and its control line

revealed that the majority of the YFP signal localizes to the Golgi with the Golgi

to cytoplasm ratios in *Δsec23abcfg-*1 indistinguishable from the control (Fig 5B),

suggesting that secretion to the Golgi is not affected in the quintuple mutant.

Many proteins transported in COPII-coated vesicles ultimately reside at the

plasma membrane. In wild type cells, the fluorescent marker F-SNAP-mCherry

accumulated on the plasma membrane, with only a very small amount of

fluorescence detected inside the cell (Fig 5C). In *Δsec23d,* a large portion of the

mCherry signal was retained in the cell, labeling the ER network (Fig 5C). To

quantify secretion efficiency to the plasma membrane, we calculated the

fluorescence intensity ratio of the plasma membrane to the intensity within the

cell. We found that the ratio in *Δsec23d* was significantly lower than that of its

control line (Fig 5D), indicating that loss of Sec23D inhibits secretion of plasma

membrane bound proteins*.*

Interestingly, we found that the F-SNAP-mCherry signal was also retained

inside the cell in the *Δsec23abcfg-1* mutant, readily observing labeling of the ER

network (Fig 5C). In addition, the fluorescence intensity ratio in *Δsec23abcfg-1*

was lower in comparison to the ratio in its control line (Fig 5D). However, the

ratio in *Δsec23abcfg-1* was still significantly higher than the ratio in *Δsec23d*,

suggesting that secretion of plasma membrane proteins is impaired in

Δsec23abcfg-1 but to a lower extent than in *Δsec23d*. These data together with

the lack of a growth defect in protonomata in the *Δsec23abcfg-1* mutant suggest

that Sec23A, B, C, F, and G contribute to protein secretion particularly to the

plasma membrane, but importantly their cargos are not essential for polarized

growth in protonemata.

Sec23D localizes to discrete puncta associated with the ER

If Sec23 is a bona fide COPII component, then we would expect that Sec23

isoforms should associate with the ER. To test this, we inserted sequences in the

genome of a line expressing GFP in the ER lumen encoding for three tandem

mRuby2 (hereafter, mRuby2) in frame with the coding sequences of the three

mostly highly expressed protonematal *Sec23* genes: *Sec23B*, *Sec23D* and *Sec23G*

(Fig 1B, Fig S3A, B). To ensure that tagging *Sec23D* did not affect its function, we

measured plant area from 7-day old regenerated protoplasts and found that

Sec23D-mRuby2 grew indistinguishably from the control lines (Fig S3C),

suggesting that Sec23D-mRuby2 is functional. By analogy, we reasoned that

tagging *Sec23B* and *G* would also be functional. To ensure we did not observe any

possible dominant negative impacts as a result of the tagging, we performed

growth assays of 7-day old plants regenerated from protoplasts and observed

that *Sec23B-mRuby2* was indistinguishable from the control and *Sec23G-mRuby2*

was slightly smaller (Fig. S3C).

Imaging with confocal microscopy revealed that Sec23D, Sec23B and

Sec23G localized to punctate structures (Fig 6A). Interestingly, Sec23G-mRuby2

puncta were larger than the punctate structures found in Sec23D- and Sec23B-

mRuby2 lines (Fig 6A and C), suggesting that Sec23G compartments maybe

different from Sec23B and D. While Sec23 isoforms labeled structures of

different sizes, all three proteins labeled puncta that were proximal to the ER

(Fig 6A). Co-localization measurements using Pearson's Correlation Coefficient

suggested that Sec23D, B and G puncta positively correlated with the ER. To

ensure that this correlation was significant, we flipped the ER image both

vertically and horizontally and repeated the correlation analysis and found that

the correlation coefficients were significantly lower (Fig 6B).

Sec24 isoforms differentially contribute to protonemal growth

Sec23 forms an obligate heterodimer with Sec24 to build the inner coat of

COPII. We reasoned that if a specific Sec24 isoform dimerizes with Sec23D, then

loss of that Sec24 isoform should result in a phenotype similar to that observed

for *Δsec23d*. *P. patens* has seven *Sec24* genes, which is the same number of genes

found in the *Sec23* gene family. Similar to *Sec23*, there are four subclasses of

Sec24 genes and six of the seven *Sec24* genes exist in closely related pairs (Fig

7A). We found that *Sec24E* was not expressed in 8-day old protonemal tissue,

while the other six *Sec24* genes were expressed (Fig 7B). Of these six genes,

Sec24D exhibited the lowest expression while the other five Sec24 genes were

expressed at similar levels (Fig 7A). Taking advantage of the high sequence

similarity within each *Sec24* subclass, regions of the coding sequences from

Sec24B, *Sec24C*, and *Sec24F* were cloned to generate RNAi constructs (Table S1)

that would target both members of the subclass (Fig 7A). Interestingly, silencing

Sec24C and *Sec24D* resulted in severe growth defects, with a 79% reduction in

plant area (Fig 7C and D). Silencing *Sec24F* and *Sec24G* also impaired growth, but

to a lesser extent resulting in a 48% reduction in plant area (Fig 7C and D). On

the other hand, silencing *Sec24A* and *Sec24B* did not affect growth, although both

genes were expressed at high levels. These data suggest that similar to Sec23,

there is isoform specificity among *Sec24* genes. Due to the similar RNAi

phenotypes observed for Sec23D/E and Sec24C/D, we hypothesized that

Sec23D/E and Sec24C/D form heterodimers.

To test for direct interaction between Sec23D and Sec24 isoforms, we

conducted a yeast two-hybrid assay using Sec23D as bait and Sec24B, Sec24C,

Sec24D, and Sec24F as preys. Full-length cDNAs of these genes were isolated

from protonemal tissue, and with the exception of *Sec24C*, only one transcript

was identified from each locus. For *Sec24C*, we isolated two alternative

transcripts: Sec24C V3.1 and Sec24C V3.3. Sec24C V3.1 is truncated at the N

terminus compared to Sec24C V3.3, lacking the first exon and the first half of the

second exon (Fig S4). Pairwise yeast two-hybrid assays demonstrated that

Sec23D strongly interacts with both versions of Sec24C and with Sec24D. We

observed a weaker interaction between Sec23D and Sec24B, but no detectable

interaction between Sec23D and Sec24F (Fig7E).

If Sec24C/D heterodimerizes with Sec23D, then we would expect

Sec24C/D localization pattern to be similar to Sec23D. To test this, we inserted

mEGFP (Vidali et al., 2009) in frame with the C-terminus of Sec24C at its

endogenous locus. We found that Sec24C-mEGFP localized to punctate structures

(Fig 7F) similar to Sec23D-mRuby (Fig 6A). While we attempted to tag both

Sec23D and Sec24C with a fluorescent protein in the same line, we were unable

to isolate these lines, suggesting that Sec23/Sec24 heterodimerization may be

impaired when both proteins are tagged. To test if Sec24C localization depends

on functional Sec23D, we used CRISPR-Cas9 mediated genome editing to disrupt

Sec23D in the Sec24C-mEGFP line (Fig S1, Table S2). We discovered that when

Sec23D was disrupted, Sec24C-mEGFP no longer localized to puncta and was

diffuse throughout the cytoplasm (Fig 7F), indicating that Sec24C localization

depends on Sec23D.

Discussion

The large number of genes encoding components of COPII in plants

together with functional studies in Arabidopsis (De Craene et al., 2014; Chung et

al., 2016; Barlow and Dacks, 2018; Brandizzi, 2018)(Faso et al., 2009; Qu et al.,

2017; Tanaka et al., 2013) have led to the hypothesis that gene expansion

underlies increased functional diversification required for complex secretory

processes that pattern and build plant cells. Evidence suggests that functional

diversification could stem from tissue/developmental specific expression (Hino

et al., 2011; Lee et al.; Hanton et al., 2009) or from the formation of COPII

complexes comprised of unique isoform compositions (Zeng et al., 2015).

However, it remains unclear whether unique COPII complexes mediate transport

of specific cargos or accumulate at some ER exit sites and not at others. Here, by

analyzing the role of Sec23 isoforms in protonemal development, we have

provided evidence for formation of distinct ER exit sites and the requirement of

only Sec23D/E for transport of cargos uniquely required for tip growth.

In particular, we discovered that one of the Sec23 pairs, Sec23D/E, plays

an essential role in protonemal growth. Several lines of evidence suggest that

Sec23D functions as a bona fide COPII component. First, *Δsec23d* plants had

abnormal ER morphology and elevated ER stress consistent with a block in cargo

secretion from the ER. Second, secretion to both the Golgi and the plasma

membrane was inhibited. Third, Sec23D localized to puncta near the ER. Due to

differential expression levels, we found that Sec23D is more important for

protonemal growth than Sec23E. However, when expressed at sufficient levels,

Sec23E can rescue loss of Sec23D indicating that Sec23D and E are functionally

redundant. The inability to recover a double mutant lacking both Sec23D and E

suggests that together these two genes are essential for moss viability. Given that

viability in moss is measured in the ability to generate protonemata, which is a

tip-growing tissue, we reason that Sec23D and E are essential for tip growth. In

support of this, *Δsec23d* protonemal cells grow significantly slower than wild

type leading to smaller plants that develop gametophores late. However, once

gametophores form they are morphologically normal, suggesting that Sec23D

function is specifically required during tip growth.

Using RNAi and stable knockouts, we found that the remaining five Sec23

genes were dispensable for formation of both protonemata and gametophores.

One possible explanation for this finding is that Sec23A, B, C, F and G have

functionally diversified no longer playing a role in ER to Golgi transport.

However, our data suggest otherwise. In protonemata, we observed secretion

defects of plasma membrane cargo in *Δsec23abcfg* plants. Given that *Sec23B, F,*

and *G* are expressed at similar levels to *Sec23D* in protonemata and Sec23B and G

localize proximal to the ER, it is likely that Sec23B, F and G contribute to COPII

function in protonemata. However, in contrast to Sec23D, their cargos may not

be essential for tip growth. Furthermore, if COPII function is required for

secretion underlying gametophore formation, we would have expected to

observe a mutant combination leading to defective gametophore formation. Yet,

we found that both *Δsec23d* and *Δsec23abcfg* formed morphologically normal

gametophores, suggesting that Sec23D and Sec23A, B, C, F and G are functionally

redundant for COPII-mediated secretion required to shape gametophores. If, on

the other hand, COPII is not required for gametophore formation, then future

experiments conditionally removing Sec23D/E in *Δsec23abcfg* plants would be

expected to form normal gametophores and would be an excellent tool to

identify COPII-independent secretion pathways in plants, which are thought to

play a role in transport of tonoplast resident proteins such as the vacuolar H+-
ATPase, VHA-a3 (Viotti et al., 2013).

While Sec23B and G formed puncta associated with the ER, these puncta
were different from each other and Sec23D. Sec23G formed large dots near the
ER, whereas Sec23D and B were much smaller. Formation of larger domains
suggests that Sec23G may be involved in the secretion of larger cargo from the
ER and provides evidence that moss has morphologically distinct ER exit sites.

Given that Sec23D contributes to both Golgi and plasma membrane
secretion, while Sec23A, B, C, F and G contribute to plasma membrane secretion,
it is plausible to suggest that tip growth requires intact Golgi secretion. In
Δsec23d plants cargos essential for tip growth are either inefficiently carried
from the ER to the Golgi or are not properly processed in the Golgi due to severe
reductions in secretion to the Golgi. Perhaps there is a larger burden for
carbohydrates synthesized by enzymes in the Golgi to generate a tip growing cell
wall as compared to a diffusely growing cell wall. To test this, future studies will
focus on identifying cargos that have been differentially affected in *Δsec23d*
versus *Δsec23abcfg* plants. Comparative mass spectrometry of isolated
microsomes and plasma membrane extracts in these mutant backgrounds could
be very informative. In addition, quantitatively measuring secretion of known
Golgi resident carbohydrate biosynthetic enzymes and plasma membrane
complexes such as cellulose and callose synthases in these mutants could help to
narrow down cargo specificity.

As a COPII component, Sec23D should form a heterodimer with at least
one of the seven *P. patens* Sec24 isoforms. Directed two hybrid assays
demonstrated that Sec23D interacted most strongly with Sec24C and Sec24D.
Furthermore, plants lacking the *Sec24C/D* gene pair phenocopied *Sec23D/E* RNAi
plants. We also found that Sec24C localizes to puncta throughout the cytoplasm
and this localization depends on Sec23D. Taken together these data suggest that
Sec23D and Sec24C, the most highly expressed gene in the *Sec24C/D* pair,
heterodimerize. Due to the strong and unique phenotype associated with loss of
Sec23D/Sec24C, we hypothesize that Sec23D together with its partner Sec24C,
carry cargo required specifically for tip growth. The findings that Sec23D
preferentially binds to Sec24C/D and their loss of function results in the same

defects in tip growth are consistent with studies in *Arabidopsis* showing that

Sar1A and Sec23A form a unique interaction mediating functional specificity

(Zeng et al., 2015).

Taking advantage of the facile molecular genetic manipulation and

imaging in moss protonemal development, our study has shown that the

expanded *Sec23* gene family in moss is multi-faceted. We found evidence that the

Sec23 gene family exhibits aspects of functional redundancy, differential gene

expression, as well as isoform specificity for COPII complex formation, ER exit

site formation, and importantly transport of essential cargos required for tip

growth. Building upon the tools developed in this study, future studies will be

able to characterize the remaining COPII components, thereby providing a

complete picture of how gene expansions in COPII have been deployed in the

evolution of this organism.

Materials and Methods

Expression analyses

To quantify expression of the *Sec23* and *Sec24* genes family members

(Table S3), as well as *Bip1* (Pp3c10_17310), we isolated total RNA from 8-day old

plants (various genotypes, as needed for the particular experiment) regenerated

from protoplasts using the RNeasy Plant mini prep kit (Qiagen). cDNA was

prepared from 1µg of total RNA using SuperScriptIII reverse transcriptase kit,

following the manufacturer's recommendation (Invitrogen). qPCR assays were

conducted with Luna Universal qPCR Master Mix (New England Biolabs).

Primers for qRT-PCR analyses are listed in Table S4. After confirming that

amplification efficiencies (calculated from standard curves) were similar for

each set of primers (Table S4), relative expression levels were calculated, and

expression of Ubiquitin 10 was used for normalization. All experiments had

three technical replicates and three biological replicates.

Constructs

The Sec23 and Sec24 RNAi constructs were generated by PCR

amplification of either the coding sequence or 5' and 3' untranslated region

(UTR) of *Sec23* or *Sec24* from *P. patens* cDNA using primers indicated in Table S4.

PCR fragments were cloned into pENTR-D-TOPO (Invitrogen) following the

manufacturer's recommendations and the resulting vectors were sequenced. The

Sec23ACD-RNAi and Sec23 UTR-RNAi constructs were generated by stitching

together multiple PCR fragments and then directionally cloned into pENTR-D-

TOPO as previously described (Vidali et al., 2007). LR clonase (Invitrogen)

reactions were used to transfer either the coding sequence or 5' and 3' UTR

sequences the RNAi vector pUGGi (Bezanilla et al., 2005) generating the Sec23

UTRi or Sec24-CDSi constructs, respectively. Restriction enzyme digestion was

used to verify these constructs.

Expression constructs were generated by PCR amplifying *Sec23D* and

Sec23E coding sequences from *P. patens* cDNA using a 5' CACC sequence on the

specific primers for oriented cloning into the pENTR-D-TOPO vector

(Invitrogen). LR clonase (Invitrogen) reactions were used to transfer these

coding sequence into pTH-Ubi-Gate vectors (Vidali et al., 2007).

Sec23 and *Sec24* tagging constructs were generated by PCR amplifying

genomic sequences upstream and downstream of the stop codon for *Sec23D*,

Sec23B, *Sec23G* and *Sec24C* using the primers in Table S4. For *Sec23D* and *Sec24C*,

we used homologous recombination. The distance between the 5' and 3'

homology arms was 500bp for both *Sec23D* and *Sec24C*, which when inserted

correctly the tagging construct replaces these 500bp with sequences encoding

for the fluorescent protein and an antibiotic resistance cassette (Fig S3). Using

BP clonase (Invitrogen), homology arms were transferred to vectors, generating

homology arm entry clones. Four-fragment recombination reactions were used

to assemble the homology arms, fluorescent protein sequence, and the antibiotic

resistance cassette into pGEM-gate (Vidali et al., 2009). For *Sec23B* and *Sec23G*

tagging, we used CRISPR Cas9-mediated homology directed repair. 5' and 3'

homology arms upstream and downstream of the stop codon, with a distance

between these two arms of 38bp for *Sec23B* and 131bp for *Sec23G* were PCR

amplified from genomic DNA. Using BP clonase (Invitrogen), homology arms

were transferred to vectors, generating homology arm entry clones, as described

recently (Mallett et al., 2019). Three-fragment recombination reactions were

used to assemble the homology arms and the fluorescent protein sequence into

pGEM-gate. We chose the CRISPR-Cas9 protospacer close to the stop codon.

Protospacers for CRISPR-Cas9 mediated mutagenesis and HDR were

designed with the CRISPOR online software (crispor.tefor.net) (Haeussler et al.,

2016). Protospacers were selected with high specificity scores and low off-target

frequency. Specific overhangs sequences were added to facilitate downstream

cloning. Protospacers were synthesized as oligonucleotides (Table S4) and

annealed as described (Mallett et al., 2019). Annealed protospacer fragments

were transferred to vectors using Instant Sticky-End Ligation Master Mix (New

England Biolabs). *Sec23B* and *G* protospacers designed to tag these genes via

HDR as well as *Sec23A*, *D* and *E* protospacers designed to knock out these genes

were ligated into pENTR-PpU6-sgRNA-L1L2 (Mallett et al., 2019). *Sec23B* and *F*

protospacers designed to knock out these genes were ligated into pENTR-PpU6-

sgRNA-L1L5 (Mallett et al., 2019). Sec23C and G protospacers designed to knock

out these genes were ligated into pENTR-PpU6-sgRNA-R5L2 (Mallett et al.,

2019). LR clonase (Invitrogen) was used to assemble the protospacer containing

vectors into pZeo-Cas9-Gate, pMH-Cas9-Gate, or pMK-Cas9-Gate vectors (Mallett

et al., 2019), as needed. Two-fragment recombination reactions were used to

assembly dual targeting mutagenesis vectors for *Sec23B* and *C*, and for *Sec23F*

and *G*. Restriction enzyme digestion and sequencing was used to verify these

constructs.

Sec23D, *Sec24B*, *CV3.1*, *CV3.3*, *D* and *F* coding sequences were amplified

from *P. patens* cDNA using a 5' CACC sequence on the specific primers for

oriented cloning into pENTR-D-TOPO (Invitrogen). LR clonase (Invitrogen)

reactions were used to transfer these coding sequences into yeast 2-hybrid

vectors. *Sec23D* was transferred to pGBKT7-Gate (Bascom Jr et al., 2018), and all

Sec24 genes were transferred to pGADT7-Gate (Bascom Jr et al., 2018).

Tissue propagation, transformation, and genotyping

All moss lines were propagated weekly by moderate homogenization in

water and pipetting onto a permeable cellophane covered solid media, as

described previously (Wu and Bezanilla, 2014). Tissue was grown at room

temperature in Percival growth chambers, under 85 $\mu mol_{photons}/m^2s$ light with

long-day conditions. PEG-mediated transformations were performed as

previously described to generate stable lines (Wu and Bezanilla, 2014), CRISPR-

Cas9 genome edited lines (Mallett et al., 2019) and perform transient RNAi

(Vidali et al., 2007).

577   To genotype mutants generated by CRISPR-Cas9 mutagenesis, we

extracted DNA from plants that were 3-4 weeks old (0.5-1cm in diameter) using

the protocol described in (Augustine et al., 2011). For editing experiments, we

used primers (Table S4) surrounding the expected Cas9 cleavage site (~300-400

bp on each side). T7 endonuclease assay was used to screen candidate mutants

as described (Mallett et al., 2019). For homology-directed repair tagging and

homologous recombination tagging, we used primers (Table S4) outside of the

homology region to avoid amplification of residual DNA donor template.

Imaging and morphometric analysis of growth assays

587   For transient RNAi, complementation analyses, and growth assays of

stable lines, plants regenerated from protoplasts were photographed seven days

after protoplasting. For transient RNAi and complementation analyses plants

lacking nuclear GFP signal were imaged because these are the plants that are

actively silencing as described previously (Bezanilla et al., 2005; Vidali et al.,

2007). For growth assays of stable lines any regenerated plant was imaged.

Images were acquired with a 1X objective using a stereomicroscope (Leica

MZ16FA or Nikon SMZ25) equipped with a CCD camera (Leica DF300FX or

Nikon digital sight DS-Fi2). Chlorophyll fluorescence and any nuclear GFP signal

were acquired simultaneously using filters (480/40, dichroic 505 (Leica) or

dichroic 510 (Nikon), emission 510 long pass). Exposure settings were

maintained constant throughout an experiment. 3-week old plants regenerated

from protoplasts were imaged with white light on a Nikon SMZ25

stereomicroscope equipped with a Nikon digital sight DS-Fi2 color camera.

601   To quantify plant area, individual plants were cropped from images and

the red channel, representing chlorophyll fluorescence of the plant, was

separated from the RGB image. Threshold settings were set manually for all

plants within an experiment. To quantify brightfield images of 3-week-old plants,

the blue channel, representing chlorophyll fluorescence of the plant, was

separated from the RGB image and thresholded. Total area was calculated from

the largest thresholded object in the selected window. All images analysis was

done using macros (Galotto et al., 2019) written for ImageJ. Analysis of variation

(ANOVA) for multiple comparisons was performed with Kaleidagraph (Synergy)

using the Tukey HSD post hoc tests. The alpha for statistical significance was set

to 0.05.

Laser scanning confocal microscope imaging and quantification

For imaging Sec23D localization and dynamics, 5 to 8-day old plants

regenerated from protoplasts were placed onto an agar pad in Hoagland's buffer

(4mM KNO_3, 2mM KH_2PO_4, 1mM $Ca(NO_3)_2$, 89µM Fe citrate, 300µM $MgSO_4$,

9.93µM H_3BO_3, 220nM $CuSO_4$, 1.966µM $MnCl_2$, 231nM $CoCl_2$, 191nM ZnSO4,

169nM KI, 103nM Na_2MoO_4, and 1% sucrose), covered by a glass cover slip and

sealed with VALAP (1:1:1 parts of Vaseline, lanoline, and paraffin). For imaging

ER morphology, Golgi resident protein secretion, F-SNAP-mCherry secretion, we

used microfluidic imaging chambers (Bascom et al., 2016). Ground protonemal

tissue was gently pipetted into the central part of the device followed by an

infusion of Hoagland's medium. Then the chamber was submerged Hoagland's

medium, and placed under constant 85 $\mu mol_{photons}/m^2s$ light.

Images were acquired on a Nikon A1R confocal microscope system with a

1.4 NA 60X oil immersion objective (Nikon) at room temperature. Image

acquisition was controlled by NIS-Element AR 4.1 software (Nikon). All images

were processed using Image J software with background subtraction, enhanced

contrast and smoothing. All settings were the standard Image J settings.

For quantification of secretion to the Golgi, confocal Z-stacks of individual

cells were converted to a maximum intensity projection. YFP fluorescence

intensity with the same expression pattern was normalize into the same range in

ImageJ. Using thresholding it was possible to isolate the fluorescence intensity of

Golgi dot (Golgi intensity) and the fluorescence intensity of rest of the cell

(background intensity). These values were used to calculate the ratio of Golgi

intensity to background intensity. For quantification of secretion to the plasma

membrane, the medial focal plane was selected for analysis. The mCherry

fluorescence intensity was normalize into the same range using ImageJ. Using

thresholding it was possible to isolate the fluorescence intensity of the plasma

membrane (PM) and the fluorescence intensity of rest of the cell (internal).

These values were used to calculate the ratio of Golgi intensity to background

intensity. To quantify the Pearson's Correlation Coefficient, we analyzed single

focal planes with the Coloc2 plugin in Fiji comparing the original images as well

as horizontally and vertically flipped ER-GFP images. To quantify the size of

Sec23B, D and G puncta, we thresholded single focal plane images to isolate the

dots and then followed with particle analysis in Fiji. Dots resulting from the

merge of two or more puncta were removed from the analysis. For all data,

analysis of variation (ANOVA) for multiple comparisons was performed with

Kaleidagraph (Synergy) using the Tukey HSD post hoc tests. The alpha for

statistical significance was set to 0.05.

2-Hybrid Assays

pGBKT7-Sec23D was transformed into Y2H Gold competent cells (Takara-

Bio) and pGADT7-Sec24 constructs were transformed into Y187 competent cells

(Takara-Bio). Y2H Gold cells and Y187 cells were mated and then diploids were

selected. Diploids were tested for 2-hybrid interactions with serial dilutions of

overnight cultures with the same optical density value equal amount using

minimal medium lacking Tryptophan (selects for pGBKT7 plasmid), Leucine

(selects for pGADT7 plasmid), Histidine and Adenine (promoters driving

expression of genes essential for the synthesis of these amino acids are activated

when there is a positive 2-hybrid interaction).

Author contributions

M.C. and M.B. designed the research. M.C., S.W., and J.E.O. performed the

research. M.C., S.W. and M.B. wrote the article.

Acknowledgements

We thank members of the Bezanilla lab for careful reading of the

manuscript. M.C. received support from the Plant Biology Graduate Program at

the University of Massachusetts, Amherst. This work was supported by

Dartmouth College and grants from the National Science Foundation (MCB-

1330171 and MCB-1715785 to M.B.).

References

675   **Aridor, M.** (2018). COPII gets in shape: Lessons derived from morphological
676   aspects of early secretion. Traffic **19**: 823–839.
677

678   **Aridor, M., Fish, K.N., Bannykh, S., Weissman, J., Roberts, T.H., Lippincott-**
679   **Schwartz, J., and Balch, W.E.** (2001). The Sar1 Gtpase Coordinates
680   Biosynthetic Cargo Selection with Endoplasmic Reticulum Export Site
681   Assembly. Journal of Cell Biology **152**: 213–230.
682

683   **Aridor, M., Weissman, J., Bannykh, S., Nuoffer, C., and Balch, W.E.** (1998).
684   Cargo Selection by the COPII Budding Machinery during Export from the
685   ER. Journal of Cell Biology **141**: 61–70.
686

687   **Augustine, R.C., Pattavina, K.A., Tüzel, E., Vidali, L., and Bezanilla, M.** (2011).
688   Actin interacting protein1 and actin depolymerizing factor drive rapid
689   actin dynamics in Physcomitrella patens. Plant Cell **23**: 3696–3710.
690

691   **Barlow, L.D. and Dacks, J.B.** (2018). Seeing the endomembrane system for the
692   trees: Evolutionary analysis highlights the importance of plants as models
693   for eukaryotic membrane-trafficking. Seminars in Cell & Developmental
694   Biology **80**: 142–152.
695

696   **Barlowe, C.** (2003). Signals for COPII-dependent export from the ER: what's the
697   ticket out? Trends in Cell Biology **13**: 295–300.
698

699   **Barlowe, C. and Helenius, A.** (2016). Cargo Capture and Bulk Flow in the Early
700   Secretory Pathway. Annu. Rev. Cell Dev. Biol. **32**: 197–222.
701

702   **Barlowe, C. and Schekman, R.** (1993). SEC12 encodes a guanine-nucleotide-
703   exchange factor essential for transport vesicle budding from the ER.
704   Nature **365**: 347–349.
705

706   **Bascom, C.S., Wu, S.-Z., Nelson, K., Oakey, J., and Bezanilla, M.** (2016). Long-
707   Term Growth of Moss in Microfluidic Devices Enables Subcellular Studies
708   in Development. Plant Physiology **172**: 28–37.
709

710   **Bascom Jr, C., Burkart, G.M., Mallett, D.R., O'Sullivan, J.E., Tomaszewski, A.J.,**
711   **Walsh, K., and Bezanilla, M.** (2018). Systematic survey of the function of
712   ROP regulators and effectors during tip growth in the moss
713   Physcomitrella patens. J. Exp. Bot. **70**: 447–457.
714

715   **Belden, W.J.** (2001). Role of Erv29p in Collecting Soluble Secretory Proteins into
716   ER-Derived Transport Vesicles. Science **294**: 1528–1531.
717

718   **Bezanilla, M., Perroud, P.-F., Pan, A., Klueh, P., and Quatrano, R.S.** (2005). An
719   RNAi system in Physcomitrella patens with an internal marker for
720   silencing allows for rapid identification of loss of function phenotypes.
721   Plant Biol (Stuttg) **7**: 251–257.
722

723 **Bi, X., Corpina, R.A., and Goldberg, J.** (2002). Structure of the Sec23/24–Sar1
724   pre-budding complex of the COPII vesicle coat. Nature **419**: 271–277.
725

726 **Brandizzi, F.** (2018). Transport from the endoplasmic reticulum to the Golgi in
727   plants: Where are we now? Seminars in Cell & Developmental Biology **80**:
728   94–105.
729

730 **Brandizzi, F. and Barlowe, C.** (2013). Organization of the ER–Golgi interface for
731   membrane traffic control. Nat Rev Mol Cell Biol **14**: 382–392.
732

733 **Cho, Y. and Kanehara, K.** (2017). Endoplasmic Reticulum Stress Response in
734   Arabidopsis Roots. Front. Plant Sci. **8**.
735

736 **Chung, K.P., Zeng, Y., and Jiang, L.** (2016). COPII Paralogs in Plants: Functional
737   Redundancy or Diversity? Trends in Plant Science **21**: 758–769.
738

739 **De Craene, J.-O., Courte, F., Rinaldi, B., Fitterer, C., Herranz, M.C., Schmitt-**
740   **Keichinger, C., Ritzenthaler, C., and Friant, S.** (2014). Study of the Plant
741   COPII Vesicle Coat Subunits by Functional Complementation of Yeast
742   Saccharomyces cerevisiae Mutants. PLoS ONE **9**: e90072.
743

744 **Faso, C., Chen, Y.-N., Tamura, K., Held, M., Zemelis, S., Marti, L., Saravanan,**
745   **R., Hummel, E., Kung, L., Miller, E., Hawes, C., and Brandizzi, F.** (2009).
746   A Missense Mutation in the Arabidopsis COPII Coat Protein Sec24A
747   Induces the Formation of Clusters of the Endoplasmic Reticulum and
748   Golgi Apparatus. The Plant Cell **21**: 3655–3671.
749

750 **Fath, S., Mancias, J.D., Bi, X., and Goldberg, J.** (2007). Structure and
751   Organization of Coat Proteins in the COPII Cage. Cell **129**: 1325–1336.
752

753 **Furt, F., Lemoi, K., Tüzel, E., and Vidali, L.** (2012). Quantitative analysis of
754   organelle distribution and dynamics in Physcomitrella patens protonemal
755   cells. BMC Plant Biol. **12**: 70.
756

757 **Galotto, G., Bibeau, J.P., and Vidali, L.** (2019). Automated Image Acquisition
758   and Morphological Analysis of Cell Growth Mutants in Physcomitrella
759   patens. In Plant Cell Morphogenesis, F. Cvrčková and V. Žárský, eds,
760   Methods in Molecular Biology. (Springer New York: New York, NY), pp.
761   307–322.
762

763 **Giraudo, C.G. and Maccioni, H.J.F.** (2003). Endoplasmic Reticulum Export of
764   Glycosyltransferases Depends on Interaction of a Cytoplasmic Dibasic
765   Motif with Sar1. MBoC **14**: 3753–3766.
766

767 **van Gisbergen, P.A.C., Wu, S.-Z., Chang, M., Pattavina, K.A., Bartlett, M.E., and**
768   **Bezanilla, M.** (2018). An ancient Sec10-formin fusion provides insights
769   into actin-mediated regulation of exocytosis. J. Cell Biol. **217**: 945–957.
770

771 **Haeussler, M., Schönig, K., Eckert, H., Eschstruth, A., Mianné, J., Renaud, J.-B.,**
772   **Schneider-Maunoury, S., Shkumatava, A., Teboul, L., Kent, J., Joly, J.-S.,**
773   **and Concordet, J.-P.** (2016). Evaluation of off-target and on-target
774   scoring algorithms and integration into the guide RNA selection tool
775   CRISPOR. Genome Biol. **17**: 148.
776

777 **Hanna, M.G., Peotter, J.L., Frankel, E.B., and Audhya, A.** (2018). Membrane
778   Transport at an Organelle Interface in the Early Secretory Pathway: Take
779   Your Coat Off and Stay a While: Evolution of the metazoan early secretory
780   pathway. BioEssays **40**: 1800004.
781

782 **Hanton, S.L., Matheson, L.A., Chatre, L., and Brandizzi, F.** (2009). Dynamic
783   organization of COPII coat proteins at endoplasmic reticulum export sites
784   in plant cells. The Plant Journal **57**: 963–974.
785

786 **Hino, T., Tanaka, Y., Kawamukai, M., Nishimura, K., Mano, S., and Nakagawa,**
787   **T.** (2011). Two Sec13p Homologs, AtSec13A and AtSec13B, Redundantly
788   Contribute to the Formation of COPII Transport Vesicles in *Arabidopsis*
789   *thaliana*. Bioscience, Biotechnology, and Biochemistry **75**: 1848–1852.
790

791 **Jensen, D. and Schekman, R.** (2011). COPII-mediated vesicle formation at a
792   glance. Journal of Cell Science **124**: 1–4.
793

794 **Khoriaty, R. et al.** (2018). Functions of the COPII gene paralogs SEC23A and
795   SEC23B are interchangeable in vivo. Proc Natl Acad Sci USA **115**: E7748–
796   E7757.
797

798 **Lee, M.H., Lee, S.H., Kim, H., Jin, J.B., Kim, D.H., and Hwang, I.** A WD40 Repeat
799   Protein, Arabidopsis Sec13 Homolog 1, May Play a Role in Vacuolar
800   Trafficking by Controlling the Membrane Association of AtDRP2A.: 10.
801

802 **Mallett, D.R., Chang, M., Cheng, X., and Bezanilla, M.** (2019). Efficient and
803   modular CRISPR-Cas9 vector system for Physcomitrella patens. Plant
804   Direct **3**: e00168-15.
805

806 **Mancias, J.D. and Goldberg, J.** (2007). The Transport Signal on Sec22 for
807   Packaging into COPII-Coated Vesicles Is a Conformational Epitope.
808   Molecular Cell **26**: 403–414.
809

810 **Maruyama, D., Sugiyama, T., Endo, T., and Nishikawa, S.** (2014). Multiple BiP
811   Genes of Arabidopsis thaliana are Required for Male Gametogenesis and
812   Pollen Competitiveness. Plant and Cell Physiology **55**: 801–810.
813

814 **Mossessova, E., Bickford, L.C., and Goldberg, J.** (2003). SNARE selectivity of
815   the COPII coat. Cell **114**: 483–495.
816

Nakano, A. and Muramatsu, M. (1989). A Novel GTP-binding Protein, Sarlp, Is
Involved in Transport from the Endoplasmic Reticulum to the Golgi
Apparatus. The Journal of Cell Biology **109**: 2677–2691.

Nakano, R.T., Matsushima, R., Ueda, H., Tamura, K., Shimada, T., Li, L.,
Hayashi, Y., Kondo, M., Nishimura, M., and Hara-Nishimura, I. (2009).
GNOM-LIKE1/ERMO1 and SEC24a/ERMO2 Are Required for Maintenance
of Endoplasmic Reticulum Morphology in Arabidopsis thaliana. The Plant
Cell **21**: 3672–3685.

Nebenfuhr, A. (2002). Brefeldin A: Deciphering an Enigmatic Inhibitor of
Secretion. Plant physiology **130**: 1102–1108.

Nelson, B.K., Cai, X., and Nebenführ, A. (2007). A multicolored set of in vivo
organelle markers for co-localization studies in Arabidopsis and other
plants. Plant J. **51**: 1126–1136.

Niu, T.-K., Pfeifer, A.C., Lippincott-Schwartz, J., and Jackson, C.L. (2005).
Dynamics of GBF1, a Brefeldin A-Sensitive Arf1 Exchange Factor at the
Golgi□V. Molecular Biology of the Cell **16**: 1213–1222.

Noh, S.-J., Kwon, C.S., Oh, D.-H., Moon, J.S., and Chung, W.-I. (2003). Expression
of an evolutionarily distinct novel BiP gene during the unfolded protein
response in Arabidopsis thaliana. Gene **311**: 81–91.

Novick, P., Field, C., and Schekman, R. (1980). Identification of 23
complementation groups required for post-translational events in the
yeast secretory pathway. Cell **21**: 205–215.

Nufer, O. (2003). ER export of ERGIC-53 is controlled by cooperation of
targeting determinants in all three of its domains. Journal of Cell Science
116: 4429–4440.

Ortiz-Ramírez, C., Michard, E., Simon, A.A., Damineli, D.S.C., Hernandez-
Coronado, M., Becker, J.D., and Feijó, J.A. (2017). GLUTAMATE
RECEPTOR-LIKE channels are essential for chemotaxis and reproduction
in mosses. Nature **549**: 91–95.

Qu, X., Zhang, R., Zhang, M., Diao, M., Xue, Y., and Huang, S. (2017).
Organizational innovation of apical actin filaments drives rapid pollen
tube growth and turning. Mol Plant **10**: 930–947.

Rensing, S.A., Goffinet, B., Meyberg, R., Wu, S.-Z., and Bezanilla, M. (2020).
The Moss *Physcomitrium* (*Physcomitrella*) *patens* : A Model Organism for
Non-Seed Plants. Plant Cell **32**: 1361–1376.

Salama, N.R., Yeung, T., and Schekman, R.W. The Secl3p complex and
reconstitution of vesicle budding from the ER with purified cytosolic
proteins. The EMBO Journal **12**: 4073–4082.

Srivastava, R., Deng, Y., Shah, S., Rao, A.G., and Howell, S.H. (2013). BINDING
PROTEIN Is a Master Regulator of the Endoplasmic Reticulum Stress
Sensor/Transducer bZIP28 in *Arabidopsis*. Plant Cell **25**: 1416–1429.
870

871   **Stagg, S.M., Gürkan, C., Fowler, D.M., LaPointe, P., Foss, T.R., Potter, C.S.,**
872   **Carragher, B., and Balch, W.E.** (2006). Structure of the Sec13/31 COPII
873   coat cage. Nature **439**: 234–238.
874

875   **Stagg, S.M., LaPointe, P., Razvi, A., Gürkan, C., Potter, C.S., Carragher, B., and**
876   **Balch, W.E.** (2008). Structural Basis for Cargo Regulation of COPII Coat
877   Assembly. Cell **134**: 474–484.
878

879   **Takagi, J., Renna, L., Takahashi, H., Koumoto, Y., Tamura, K., Stefano, G.,**
880   **Fukao, Y., Kondo, M., Nishimura, M., Shimada, T., Brandizzi, F., and**
881   **Hara-Nishimura, I.** (2013). MAIGO5 Functions in Protein Export from
882   Golgi-Associated Endoplasmic Reticulum Exit Sites in Arabidopsis. The
883   Plant Cell **25**: 4658–4675.
884

885   **Tanaka, Y., Nishimura, K., Kawamukai, M., Oshima, A., and Nakagawa, T.**
886   (2013). Redundant function of two Arabidopsis COPII components,
887   AtSec24B and AtSec24C, is essential for male and female gametogenesis.
888   Planta **238**: 561–575.
889

890   **Vidali, L., Augustine, R.C., Kleinman, K.P., and Bezanilla, M.** (2007). Profilin is
891   essential for tip growth in the moss Physcomitrella patens. Plant Cell **19**:
892   3705–3722.
893

894   **Vidali, L., van Gisbergen, P.A.C., Guérin, C., Franco, P., Li, M., Burkart, G.M.,**
895   **Augustine, R.C., Blanchoin, L., and Bezanilla, M.** (2009). Rapid formin-
896   mediated actin-filament elongation is essential for polarized plant cell
897   growth. Proc. Natl. Acad. Sci. U.S.A. **106**: 13341–13346.
898

899   **Viotti, C. et al.** (2013). The Endoplasmic Reticulum Is the Main Membrane
900   Source for Biogenesis of the Lytic Vacuole in *Arabidopsis*. Plant Cell **25**:
901   3434–3449.
902

903   **Whittle, J.R.R. and Schwartz, T.U.** (2010). Structure of the Sec13–Sec16 edge
904   element, a template for assembly of the COPII vesicle coat. The Journal of
905   Cell Biology **190**: 347–361.
906

907   **Wu, S.-Z. and Bezanilla, M.** (2014). Myosin VIII associates with microtubule
908   ends and together with actin plays a role in guiding plant cell division.
909   eLife Sciences **3**: e03498.
910

Zeng, Y., Chung, K.P., Li, B., Lai, C.M., Lam, S.K., Wang, X., Cui, Y., Gao, C., Luo,
M., Wong, K.-B., Schekman, R., and Jiang, L. (2015). Unique COPII
component AtSar1a/AtSec23a pair is required for the distinct function of
protein ER export in *Arabidopsis thaliana*. Proc Natl Acad Sci USA **112**:
14360–14365.

Figure Legends

Figure 1. *Sec23D* and *E* are required for polarized growth. (A) Gene models

of the *P. patens Sec23* genes are shown with exons indicated by boxes and

introns by thin black lines. Coding and untranslated regions are denoted by thick

and thin boxes, respectively. The lines underneath the gene models represent

sequence regions targeted by Sec23 RNAi constructs. Scale bar is 500bp. (B)

Relative expression of *Sec23* genes normalized to UBIQUITIN10 in 8-day-old

wild-type moss plants regenerated from protoplasts. N = 3, Error bars are s.e.m.

(C) Representative chlorophyll autofluorescence images of 7-day-old plants

regenerated from protoplasts expressing the indicated RNAi constructs. Scale

bar, 100 µm. (D) Quantification of plant area is based on the area of the

chlorophyll autofluorescence and is presented normalized to the area of the

control. Means are indicated by horizontal lines. Letters indicate groups with

significantly different means as determined by a one-way ANOVA with a Tukey

post hoc test (α=0.05).

Figure 2. *Sec23D* and *Sec23E* are functionally redundant. (A) *Sec23D* and

Sec23E gene models are shown with exons indicated by boxes and introns by

thin black lines. Coding and untranslated regions are denoted by thick and thin

boxes, respectively. The lines underneath the gene models represent sequence

regions that were targeted by Sec23D or Sec23E UTR RNAi constructs. Scale bar

is 500bp. (B) Representative chlorophyll autofluorescence images of 7-day-old

plants regenerated from protoplasts expressing the indicated constructs. Scale

bar, 100 µm. (C) Quantification of plant area was normalized to the control.

Letters indicate groups with significantly different means as determined by a

one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 3. CRISPR-Cas9-mediated deletion of *Sec23* isoforms demonstrates

that *Sec23D* is critical for protonemal growth. (A) Representative chlorophyll

autofluorescence images of 7-day old plants with the indicated genotype

regenerated from protoplasts. Scale bar, 100µm. (B) Quantification of plant area

normalized to the wild type control. (C) Representative 3-week old moss plants

with the indicated genotype. Scale bar, 1mm. (D) Quantification of the plant area

of 3-week old moss plants normalized to that of the wild type control. (E)

Number of gametophores in 3-week old moss plants. Letters in (B, D-E) indicate

groups with significantly different means as determined by a one-way ANOVA

with a Tukey *post hoc* test (α=0.05). (F) Relative expression of *Sec23D* and *E*

genes normalized to UBIQUITIN10 in 8-day-old wild type and *Δsec23abcfg-1*, and

Δsec23abcfg-2 moss plants regenerated from protoplasts. N=3, Error bars are

s.e.m. No significant difference was determined by a one-way ANOVA with a

Tukey *post hoc* test (α=0.05).

Figure 4. Loss of Sec23D causes ER morphology defects and ER stress. (A)

Laser scanning confocal images of the ER labeled with GFP-KDEL in the indicated

genotype. Colored boxes highlight large and small ER aggregates found in the

Δsec23d-B11 mutant. Scale bar 10 μm. (B, C) Relative expression of *PpBip1* in 8-

day old moss plants regenerated from protoplasts. N=3, p<0.01. Error bars are

s.e.m. In (B) BFA-treated plants were normalized to DMSO-treated plants and in

(C) null mutants were normalized to wild type plants.

Figure 5. Deletion of *Sec23D* results in secretion defects to the Golgi

apparatus and the plasma membrane. (A) Representative maximum

projections of confocal images of the indicated genotypes expressing a YFP-Golgi

marker. Scale bar, 10 μm. (B) Quantification of the fluorescence intensity ratio

within the Golgi to the background. (C) Representative medial sections of

confocal images of the indicated genotypes expressing a plasma membrane

marker (FSnap-mCherry). Scale bar, 10 μm. Letters in (B, D) indicate groups with

significantly different means as determined by a one-way ANOVA with a Tukey

post hoc test (α=0.05).

Figure 6. Sec23 isoforms localize to structures with distinct sizes. (A)

Representative medial sections of confocal images of lines where the indicated

Sec23 gene was endogenously tagged with mRuby in a line that expresses GFP in

the ER. (B) Pearson's correlation coefficient measured between the Sec23

labeled lines and ER-GFP. O, original ER image; V, vertically flipped ER image; H,

horizontally flipped ER image. Letters indicate groups with significantly different

means as determined by a one-way ANOVA with a Tukey *post hoc* test ($\alpha=0.05$)

(C) Quantification of average dot size in the Sec23 images. Asterisks denote

$p<0.001$ as determined by a one-way ANOVA with a Tukey *post hoc* test ($\alpha=0.05$).

Figure 7. *Sec24* isoforms differentially contribute to protonemal growth.

(A) *Sec24* gene models are shown with exons indicated by boxes and introns by

thin black lines. Coding and untranslated regions are denoted by thick and thin

boxes, respectively. Lines underneath the models represent sequence regions

that were targeted by Sec24 RNAi constructs. Scale bar, 1000 bp. (B) Relative

expression of *Sec24* genes normalized to UBIQUITIN10 in 8-day-old wild-type

moss plants regenerated from protoplasts. N=3, Error bars are s.e.m. (C)

Representative chlorophyll autofluorescence images of 7-day old plants

regenerated from protoplasts expressing the indicated RNAi constructs. Scale

bar, 100 µm. (D) Quantification of plant area is based on the area of the

chlorophyll autofluorescence and is presented normalized to the area of the

control. Means are indicated by horizontal lines. Letters indicate groups with

significantly different means as determined by a one-way ANOVA with a Tukey

post hoc test ($\alpha=0.05$). (E) Serial dilutions of yeast strains carrying 2-hybrid

vectors testing for interaction between Sec23D and the indicated Sec24 isoforms.

Growth indicates that tested proteins interact. *Sec24C* V3.1 and V3.3 are two

splice variants isolated from 7-day old protonemal tissue. Positive control:

murine p53 insert and SV40 large T-antigen; Negative control: human lamin and

SV40 large T-antigen (Clontech). (F) Representative maximum projections of

confocal images of endogenously tagged Sec24C-mEGFP in the indicated

genotypes. Large globular structures are chloroplasts, which autofluoresce under

these imaging conditions. Scale bar, 5 µm.

Figure 1. ***Sec23D* and *E* are required for polarized growth.** (A) Gene models of the *P. patens Sec23* genes are shown with exons indicated by boxes and introns by thin black lines. Coding and untranslated regions are denoted by thick and thin boxes, respectively. The lines underneath the gene models represent sequence regions targeted by Sec23 RNAi constructs. Scale bar is 500bp. (B) Relative expression of *Sec23* genes normalized to UBIQUITIN10 in 8-day-old wild-type moss plants regenerated from protoplasts. N = 3, Error bars are s.e.m. (C) Representative chlorophyll autofluorescence images of 7-day-old plants regenerated from protoplasts expressing the indicated RNAi constructs. Scale bar, 100 µm. (D) Quantification of plant area is based on the area of the chlorophyll autofluorescence and is presented normalized to the area of the control. Means are indicated by horizontal lines. Letters indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 2. *Sec23D* and *Sec23E* are functionally redundant. (A) *Sec23D* and *Sec23E* gene models are shown with exons indicated by boxes and introns by thin black lines. Coding and untranslated regions are denoted by thick and thin boxes, respectively. The lines underneath the gene models represent sequence regions that were targeted by Sec23D or Sec23E UTR RNAi constructs. Scale bar is 500bp. (B) Representative chlorophyll autofluorescence images of 7-day-old plants regenerated from protoplasts expressing the indicated constructs. Scale bar, 100 μm. (C) Quantification of plant area was normalized to the control. Letters indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 3. CRISPR-Cas9-mediated deletion of *Sec23* isoforms demonstrates that *Sec23D* is critical for protonemal growth. (A) Representative chlorophyll autofluorescence images of 7-day old plants with the indicated genotype regenerated from protoplasts. Scale bar, 100µm. (B) Quantification of plant area normalized to the wild type control. (C) Representative 3-week old moss plants with the indicated genotype. Scale bar, 1mm. (D) Quantification of the plant area of 3-week old moss plants normalized to that of the wild type control. (E) Number of gametophores in 3-week old moss plants. Letters in (B, D-E) indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05). (F) Relative expression of *Sec23D* and *E* genes normalized to UBIQUITIN10 in 8-day-old wild type and Δ*sec23abcfg-1*, and Δ*sec23abcfg-2* moss plants regenerated from protoplasts. N=3, Error bars are s.e.m. No significant difference was determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 4. Loss of Sec23D causes ER morphology defects and ER stress. (A) Laser scanning confocal images of the ER labeled with GFP-KDEL in the indicated genotype. Colored boxes highlight large and small ER aggregates found in the Δ*sec23d-B11* mutant. Scale bar 10 μm. (B, C) Relative expression of *PpBip1* in 8-day old moss plants regenerated from protoplasts. N=3, p<0.01. Error bars are s.e.m. In (B) BFA-treated plants were normalized to DMSO-treated plants and in (C) null mutants were normalized to wild type plants.

Figure 5. Deletion of *Sec23D* results in secretion defects to the Golgi apparatus and the plasma membrane. (A) Representative maximum projections of confocal images of the indicated genotypes expressing a YFP-Golgi marker. Scale bar, 10 µm. (B) Quantification of the fluorescence intensity ratio within the Golgi to the background. (C) Representative medial sections of confocal images of the indicated genotypes expressing a plasma membrane marker (FSnap-mCherry). Scale bar, 10 µm. Letters in (B, D) indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 6. Sec23 isoforms localize to structures with distinct sizes. (A) Representative medial sections of confocal images of lines where the indicated Sec23 gene was endogenously tagged with mRuby in a line that expresses GFP in the ER. (B) Pearson's correlation coefficient measured between the Sec23 labeled lines and ER-GFP. O, original ER image; V, vertically flipped ER image; H, horizontally flipped ER image. Letters indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05) (C) Quantification of average dot size in the Sec23 images. Asterisks denote $p<0.001$ as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 7. *Sec24* isoforms differentially contribute to protonemal growth. (A) *Sec24* gene models are shown with exons indicated by boxes and introns by thin black lines. Coding and untranslated regions are denoted by thick and thin boxes, respectively. Lines underneath the models represent sequence regions that were targeted by Sec24 RNAi constructs. Scale bar, 1000 bp. (B) Relative expression of *Sec24* genes normalized to UBIQUITIN10 in 8-day-old wild-type moss plants regenerated from protoplasts. N=3, Error bars are s.e.m. (C) Representative chlorophyll autofluorescence images of 7-day old plants regenerated from protoplasts expressing the indicated RNAi constructs. Scale bar, 100 µm. (D) Quantification of plant area is based on the area of the chlorophyll autofluorescence and is presented normalized to the area of the control. Means are indicated by horizontal lines. Letters indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05). (E) Serial dilutions of yeast strains carrying 2-hybrid vectors testing for interaction between Sec23D and the indicated Sec24 isoforms. Growth indicates that tested proteins interact. *Sec24C* V3.1 and V3.3 are two splice variants isolated from 7-day old protonemal tissue. Positive control: murine p53 insert and SV40 large T-antigen; Negative control: human lamin and SV40 large T-antigen (Clontech). (F) Representative maximum projections of confocal images of endogenously tagged Sec24C-mEGFP in the indicated genotypes. Large globular structures are chloroplasts, which autofluoresce under these imaging conditions. Scale bar, 5 µm.

Parsed Citations

Aridor, M. (2018). COPII gets in shape: Lessons derived from morphological aspects of early secretion. Traffic 19: 823–839.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Aridor, M., Fish, K.N., Bannykh, S., Weissman, J., Roberts, T.H., Lippincott-Schwartz, J., and Balch, W.E. (2001). The Sar1 Gtpase Coordinates Biosynthetic Cargo Selection with Endoplasmic Reticulum Export Site Assembly. Journal of Cell Biology 152: 213–230.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Aridor, M., Weissman, J., Bannykh, S., Nuoffer, C., and Balch, W.E. (1998). Cargo Selection by the COPII Budding Machinery during Export from the ER. Journal of Cell Biology 141: 61–70.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Augustine, R.C., Pattavina, K.A, Tüzel, E., Vidali, L., and Bezanilla, M. (2011). Actin interacting protein1 and actin depolymerizing factor drive rapid actin dynamics in Physcomitrella patens. Plant Cell 23: 3696–3710.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Barlow, L.D. and Dacks, J.B. (2018). Seeing the endomembrane system for the trees: Evolutionary analysis highlights the importance of plants as models for eukaryotic membrane-trafficking. Seminars in Cell & Developmental Biology 80: 142–152.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Barlowe, C. (2003). Signals for COPII-dependent export from the ER: what's the ticket out? Trends in Cell Biology 13: 295–300.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Barlowe, C. and Helenius, A. (2016). Cargo Capture and Bulk Flow in the Early Secretory Pathway. Annu. Rev. Cell Dev. Biol. 32: 197–222.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Barlowe, C. and Schekman, R. (1993). SEC12 encodes a guanine-nucleotide-exchange factor essential for transport vesicle budding from the ER. Nature 365: 347–349.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bascom, C.S., Wu, S.-Z., Nelson, K., Oakey, J., and Bezanilla, M. (2016). Long-Term Growth of Moss in Microfluidic Devices Enables Subcellular Studies in Development. Plant Physiology 172: 28–37.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bascom Jr, C., Burkart, G.M., Mallett, D.R., O'Sullivan, J.E., Tomaszewski, A.J., Walsh, K., and Bezanilla, M. (2018). Systematic survey of the function of ROP regulators and effectors during tip growth in the moss Physcomitrella patens. J. Exp. Bot. 70: 447–457.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Belden, W.J. (2001). Role of Erv29p in Collecting Soluble Secretory Proteins into ER-Derived Transport Vesicles. Science 294: 1528–1531.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bezanilla, M., Perroud, P.-F., Pan, A, Klueh, P., and Quatrano, R.S. (2005). An RNAi system in Physcomitrella patens with an internal marker for silencing allows for rapid identification of loss of function phenotypes. Plant Biol (Stuttg) 7: 251–257.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bi, X., Corpina, R.A., and Goldberg, J. (2002). Structure of the Sec23/24–Sar1 pre-budding complex of the COPII vesicle coat. Nature 419: 271–277.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Brandizzi, F. (2018). Transport from the endoplasmic reticulum to the Golgi in plants: Where are we now? Seminars in Cell & Developmental Biology 80: 94–105.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Brandizzi, F. and Barlowe, C. (2013). Organization of the ER–Golgi interface for membrane traffic control. Nat Rev Mol Cell Biol 14: 382–392.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cho, Y. and Kanehara, K. (2017). Endoplasmic Reticulum Stress Response in Arabidopsis Roots. Front. Plant Sci. 8.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Chung, K.P., Zeng, Y., and Jiang, L. (2016). COPII Paralogs in Plants: Functional Redundancy or Diversity? Trends in Plant Science 21: 758–769.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

De Craene, J.-O., Courte, F., Rinaldi, B., Fitterer, C., Herranz, M.C., Schmitt-Keichinger, C., Ritzenthaler, C., and Friant, S. (2014). Study of the Plant COPII Vesicle Coat Subunits by Functional Complementation of Yeast Saccharomyces cerevisiae Mutants. PLoS ONE 9: e90072.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Faso, C., Chen, Y.-N., Tamura, K., Held, M., Zemelis, S., Marti, L., Saravanan, R., Hummel, E., Kung, L., Miller, E., Hawes, C., and Brandizzi, F. (2009). A Missense Mutation in the Arabidopsis COPII Coat Protein Sec24A Induces the Formation of Clusters of the Endoplasmic Reticulum and Golgi Apparatus. The Plant Cell 21: 3655–3671.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fath, S., Mancias, J.D., Bi, X., and Goldberg, J. (2007). Structure and Organization of Coat Proteins in the COPII Cage. Cell 129: 1325–1336.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Furt, F., Lemoi, K., Tüzel, E., and Vidali, L. (2012). Quantitative analysis of organelle distribution and dynamics in Physcomitrella patens protonemal cells. BMC Plant Biol. 12: 70.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Galotto, G., Bibeau, J.P., and Vidali, L. (2019). Automated Image Acquisition and Morphological Analysis of Cell Growth Mutants in Physcomitrella patens. In Plant Cell Morphogenesis, F. Cvrčková and V. Žárský, eds, Methods in Molecular Biology. (Springer New York: New York, NY), pp. 307–322.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Giraudo, C.G. and Maccioni, H.J.F. (2003). Endoplasmic Reticulum Export of Glycosyltransferases Depends on Interaction of a Cytoplasmic Dibasic Motif with Sar1. MBoC 14: 3753–3766.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

van Gisbergen, P.A.C., Wu, S.-Z, Chang, M., Pattavina, K.A, Bartlett, M.E., and Bezanilla, M. (2018). An ancient Sec10-formin fusion provides insights into actin-mediated regulation of exocytosis. J. Cell Biol. 217: 945–957.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Haeussler, M., Schönig, K., Eckert, H., Eschstruth, A, Mianné, J., Renaud, J.-B., Schneider-Maunoury, S., Shkumatava, A, Teboul, L., Kent, J., Joly, J.-S., and Concordet, J.-P. (2016). Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. Genome Biol. 17: 148.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hanna, M.G., Peotter, J.L., Frankel, E.B., and Audhya, A (2018). Membrane Transport at an Organelle Interface in the Early Secretory Pathway: Take Your Coat Off and Stay a While: Evolution of the metazoan early secretory pathway. BioEssays 40: 1800004.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hanton, S.L., Matheson, L.A, Chatre, L., and Brandizzi, F. (2009). Dynamic organization of COPII coat proteins at endoplasmic reticulum export sites in plant cells. The Plant Journal 57: 963–974.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hino, T., Tanaka, Y., Kawamukai, M., Nishimura, K., Mano, S., and Nakagawa, T. (2011). Two Sec13p Homologs, AtSec13A and AtSec13B, Redundantly Contribute to the Formation of COPII Transport Vesicles in Arabidopsis thaliana. Bioscience, Biotechnology, and Biochemistry 75: 1848–1852.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Jensen, D. and Schekman, R. (2011). COPII-mediated vesicle formation at a glance. Journal of Cell Science 124: 1–4.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Khoriaty, R. et al. (2018). Functions of the COPII gene paralogs SEC23A and SEC23B are interchangeable in vivo. Proc Natl Acad Sci

USA 115: E7748–E7757.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Lee, M.H., Lee, S.H., Kim, H., Jin, J.B., Kim, D.H., and Hwang, I. A WD40 Repeat Protein, Arabidopsis Sec13 Homolog 1, May Play a Role in Vacuolar Trafficking by Controlling the Membrane Association of AtDRP2A: 10.

Mallett, D.R., Chang, M., Cheng, X., and Bezanilla, M. (2019). Efficient and modular CRISPR-Cas9 vector system for Physcomitrella patens. Plant Direct 3: e00168-15.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Mancias, J.D. and Goldberg, J. (2007). The Transport Signal on Sec22 for Packaging into COPII-Coated Vesicles Is a Conformational Epitope. Molecular Cell 26: 403–414.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Maruyama, D., Sugiyama, T., Endo, T., and Nishikawa, S. (2014). Multiple BiP Genes of Arabidopsis thaliana are Required for Male Gametogenesis and Pollen Competitiveness. Plant and Cell Physiology 55: 801–810.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Mossessova, E., Bickford, L.C., and Goldberg, J. (2003). SNARE selectivity of the COPII coat. Cell 114: 483–495.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nakano, A. and Muramatsu, M. (1989). A Novel GTP-binding Protein, Sar1p, Is Involved in Transport from the Endoplasmic Reticulum to the Golgi Apparatus. The Journal of Cell Biology 109: 2677–2691.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nakano, R.T., Matsushima, R., Ueda, H., Tamura, K., Shimada, T., Li, L., Hayashi, Y., Kondo, M., Nishimura, M., and Hara-Nishimura, I. (2009). GNOM-LIKE1/ERMO1 and SEC24a/ERMO2 Are Required for Maintenance of Endoplasmic Reticulum Morphology in Arabidopsis thaliana. The Plant Cell 21: 3672–3685.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nebenfuhr, A. (2002). Brefeldin A: Deciphering an Enigmatic Inhibitor of Secretion. Plant physiology 130: 1102–1108.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nelson, B.K., Cai, X., and Nebenführ, A. (2007). A multicolored set of in vivo organelle markers for co-localization studies in Arabidopsis and other plants. Plant J. 51: 1126–1136.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Niu, T.-K., Pfeifer, A.C., Lippincott-Schwartz, J., and Jackson, C.L. (2005). Dynamics of GBF1, a Brefeldin A-Sensitive Arf1 Exchange Factor at the Golgi□V. Molecular Biology of the Cell 16: 1213–1222.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Noh, S.-J., Kwon, C.S., Oh, D.-H., Moon, J.S., and Chung, W.-I. (2003). Expression of an evolutionarily distinct novel BiP gene during the unfolded protein response in Arabidopsis thaliana. Gene 311: 81–91.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Novick, P., Field, C., and Schekman, R. (1980). Identification of 23 complementation groups required for post-translational events in the yeast secretory pathway. Cell 21: 205–215.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nufer, O. (2003). ER export of ERGIC-53 is controlled by cooperation of targeting determinants in all three of its domains. Journal of Cell Science 116: 4429–4440.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ortiz-Ramírez, C., Michard, E., Simon, A.A, Damineli, D.S.C., Hernandez-Coronado, M., Becker, J.D., and Feijó, J.A. (2017). GLUTAMATE RECEPTOR-LIKE channels are essential for chemotaxis and reproduction in mosses. Nature 549: 91–95.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Qu, X., Zhang, R., Zhang, M., Diao, M., Xue, Y., and Huang, S. (2017). Organizational innovation of apical actin filaments drives rapid pollen tube growth and turning. Mol Plant 10: 930–947.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rensing, S.A., Goffinet, B., Meyberg, R., Wu, S.-Z., and Bezanilla, M. (2020). The Moss Physcomitrium (Physcomitrella) patens : A Model Organism for Non-Seed Plants. Plant Cell 32: 1361–1376.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Salama, N.R., Yeung, T., and Schekman, R.W. The Secl3p complex and reconstitution of vesicle budding from the ER with purified cytosolic proteins. The EMBO Journal 12: 4073–4082.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Srivastava, R., Deng, Y., Shah, S., Rao, A.G., and Howell, S.H. (2013). BINDING PROTEIN Is a Master Regulator of the Endoplasmic Reticulum Stress Sensor/Transducer bZIP28 in Arabidopsis. Plant Cell 25: 1416–1429.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Stagg, S.M., Gürkan, C., Fowler, D.M., LaPointe, P., Foss, T.R., Potter, C.S., Carragher, B., and Balch, W.E. (2006). Structure of the Sec13/31 COPII coat cage. Nature 439: 234–238.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Stagg, S.M., LaPointe, P., Razvi, A, Gürkan, C., Potter, C.S., Carragher, B., and Balch, W.E. (2008). Structural Basis for Cargo Regulation of COPII Coat Assembly. Cell 134: 474–484.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Takagi, J., Renna, L., Takahashi, H., Koumoto, Y., Tamura, K., Stefano, G., Fukao, Y., Kondo, M., Nishimura, M., Shimada, T., Brandizzi, F., and Hara-Nishimura, I. (2013). MAIGO5 Functions in Protein Export from Golgi-Associated Endoplasmic Reticulum Exit Sites in Arabidopsis. The Plant Cell 25: 4658–4675.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Tanaka, Y., Nishimura, K., Kawamukai, M., Oshima, A., and Nakagawa, T. (2013). Redundant function of two Arabidopsis COPII components, AtSec24B and AtSec24C, is essential for male and female gametogenesis. Planta 238: 561–575.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Vidali, L., Augustine, R.C., Kleinman, K.P., and Bezanilla, M. (2007). Profilin is essential for tip growth in the moss Physcomitrella patens. Plant Cell 19: 3705–3722.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Vidali, L., van Gisbergen, P.A.C., Guérin, C., Franco, P., Li, M., Burkart, G.M., Augustine, R.C., Blanchoin, L., and Bezanilla, M. (2009). Rapid formin-mediated actin-filament elongation is essential for polarized plant cell growth. Proc. Natl. Acad. Sci. U.S.A. 106: 13341–13346.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Viotti, C. et al. (2013). The Endoplasmic Reticulum Is the Main Membrane Source for Biogenesis of the Lytic Vacuole in Arabidopsis. Plant Cell 25: 3434–3449.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Whittle, J.R.R. and Schwartz, T.U. (2010). Structure of the Sec13–Sec16 edge element, a template for assembly of the COPII vesicle coat. The Journal of Cell Biology 190: 347–361.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Wu, S.-Z. and Bezanilla, M. (2014). Myosin VIII associates with microtubule ends and together with actin plays a role in guiding plant cell division. eLife Sciences 3: e03498.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Zeng, Y., Chung, K.P., Li, B., Lai, C.M., Lam, S.K., Wang, X., Cui, Y., Gao, C., Luo, M., Wong, K.-B., Schekman, R., and Jiang, L. (2015). Unique COPII component AtSar1a/AtSec23a pair is required for the distinct function of protein ER export in Arabidopsis thaliana. Proc Natl Acad Sci USA 112: 14360–14365.
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title