COPII Sec23 proteins form isoform-specific ER exit sites with differential effects on polarized growth

Mingqin Chang[1,2,3], Shu-Zon Wu[1], Samantha E. Ryken[1], Jacquelyn E. O'Sullivan[4,5] and Magdalena Bezanilla[1]*

[1] Department of Biological Sciences, Dartmouth College, Hanover, NH 03755
[2] Plant Biology Graduate Program, University of Massachusetts Amherst, Amherst, MA 01002
[3] Current address: Department of Plant Sciences, University of California Davis, Davis, CA 95616
[4] Department of Biology, University of Massachusetts Amherst, Amherst, MA 01002
[5] Current address: Tufts School of Medicine, Boston, MA 02111

*Corresponding author
email: magdalena.bezanilla@dartmouth.edu

Abstract

COPII, a coat of proteins that form vesicles on the ER, mediates vesicle traffic from the ER to the Golgi. In contrast to metazoans that have few genes encoding each COPII component, plants have expanded these gene families leading to the hypothesis that plant COPII has functionally diversified. Here, we analyzed the gene families encoding for the Sec23/24 heterodimer in the moss *Physcomitrium (Physcomitrella) patens*. In *P. patens*, *Sec23* and *Sec24* gene families are each comprised of seven genes. Silencing the Sec23/24 genes revealed isoform specific contributions to polarized growth, with the closely related *Sec23D/E* and *Sec24C/D* essential for protonemal development. Focusing on the *Sec23* gene family, we discovered that loss of Sec23D alters ER morphology, increases ER stress, inhibits trafficking to the Golgi and to the plasma membrane in tip growing protonemata. In contrast, the remaining five *Sec23* genes are dispensable for tip growth. While *Sec23A/B/C/F/G* do not quantitatively affect ER to Golgi trafficking in protonemata, they do contribute to secretion to the plasma membrane. Of the three highly expressed Sec23 isoforms in protonemata, Sec23G forms ER exit sites that are larger than Sec23B and Sec23D and do not overlap with Sec23D. Furthermore, ER exit sites labeled by Sec23B or Sec23G form in the absence of Sec23D. These data suggest that *Sec23D/E* form unique ER exit sites contributing to secretion that is essential for tip growing protonemata.

Introduction

Trafficking from the ER to the Golgi is mediated by the Coat Protein complex II (COPII). COPII facilitates the formation of transport vesicles at ER exit sites promoting the delivery of cargo, including transmembrane proteins, soluble proteins and lipids, to *cis*-Golgi compartments (Brandizzi and Barlowe, 2013; Barlowe and Helenius, 2016; Aridor, 2018; Hanna et al., 2018; Robinson et al., 2015). The COPII coat comprises two layers of protein complexes. The inner layer consists of a Sar1-Sec23-Sec24 lattice, while the outer layer is made up of Sec31-Sec13 heterotetramers (Stagg et al., 2008; Whittle and Schwartz, 2010; Bi et al., 2002; Fath et al., 2007). Sar1 is a small GTPase of the Ras super family that is the master regulator of COPII vesicle biogenesis (Nakano and Muramatsu, 1989). Assembly of COPII coat is initiated by the conversion of Sar1-GDP to Sar1-GTP, which is mediated by Sec12, a guanine nucleotide exchange factor (GEF) that is an ER resident membrane protein (Barlowe and Schekman, 1993; Nakano and Muramatsu, 1989). At the ER surface, activated Sar1 recruits the Sec23-Sec24 heterodimer, which associates with the ER membrane as an "inner coat". The Sar1-Sec23-Sec24-cargo "pre-budding" complex in turn recruits the Sec13-Sec31 heterotetramer, referred to as an "outer coat" to the nascent bud (Aridor et al., 1998; Salama et al., 1993). The polymerization of Sec13-Sec31 heterotetramers induces membrane curvature and completes vesicle biogenesis (Fath et al., 2007; Stagg et al., 2006; Whittle and Schwartz, 2010).

Selective recruitment of cargos into COPII coated vesicles is driven by ER export signals located on cytoplasmically exposed regions of cargo proteins (Barlowe, 2003). ER export signals of most transmembrane cargo proteins are thought to interact directly with the inner coat components of COPII. However, some transmembrane and most soluble cargo proteins interact with COPII components through ER-resident transmembrane adaptors or receptors (Barlowe, 2003). Sec24 is the COPII component that has been implicated in recognizing these various sorting signals (Mossessova et al., 2003; Nufer, 2003). However, several studies suggested that Sec23 and Sar1 may also play roles in cargo sorting by directly binding to the export signals (Aridor et al., 2001; Mancias and Goldberg, 2007; Giraudo and Maccioni, 2003).

Based on the high sequence similarity among COPII paralogs, and cross-

species complementation studies (De Craene et al., 2014; Khoriaty et al., 2018), it

has been assumed that functions of COPII components are conserved across

eukaryotes. While components of the COPII coat are well studied in budding

yeast and mammalian cells, illustrating their molecular functions in plants has

been more challenging. Comparative genomic analyses revealed that gene

families encoding COPII components have undergone gene expansions in plants

with *Sec23* exhibiting the largest expansion (Aridor et al., 2001; Mancias and

Goldberg, 2007; Giraudo and Maccioni, 2003; Schlacht and Dacks, 2015). While

there are only one and two *Sec23* genes in *S. cerevisiae* and mammals (eg. mice

and humans) (Jensen and Schekman, 2011), respectively, there are seven in both

P. patens and *A. thaliana* (Brandizzi, 2018). Since Sec23 and its heterodimeric

partner Sec24 are implicated in cargo sorting, the larger *Sec23* gene family could

suggest that plants have more diverse sorting signals requiring specificity at the

level of the Sec23/Sec24 heterodimer. Consistent with this, in contrast to yeast,

which has a single gene for most COPII components, all other components of

COPII in plants are encoded by multiple genes, suggesting that cargo specificity

may translate to formation of distinct COPII complexes and vesicle populations.

Alternatively, larger gene families could simply indicate that plants have higher

functional redundancy in COPII encoded gene families.

Genetic studies in *A. thaliana* have demonstrated that loss of specific

COPII components caused developmental defects. A complete knockout of

AtSec24A was lethal (Faso et al., 2009; Nakano et al., 2009). However, a missense

point mutation in AtSec24A (R693K) resulted in ER morphology defects and

disruption of ER-Golgi integrity. Interestingly, these phenotypes could not be

rescued by either AtSec24B or AtSec24C (Faso et al., 2009; Nakano et al., 2009),

suggesting that AtSec24B and C are functionally distinct from AtSec24A. Even

though AtSec24B and AtSec24C did not rescue the *Atsec24a* missense mutant

and each were not important for plant viability, they were shown to influence the

development of reproductive cells (Tanaka et al., 2013). Functional

diversification was also suggested by analysis of AtSec16 mutants in which

storage protein precursors were accumulated abnormally in *Atsec16a* null

mutants, but not in *Atsec16b* null mutants (Takagi et al., 2013).

While *P. patens* and *A. thaliana* have similar numbers of genes encoding

COPII components, moss has several attributes that make it particularly suited to

study COPII function. *P. patens* juvenile tissues are haploid and comprise

multiple tissue types that establish the plant body. Protonemal tissue, which

emerges from the spore and is also readily propagated vegetatively by tissue

homogenization, is composed of tip-growing cells constituting a two-

dimensional filamentous network. The predictable patterns of growth and cell

division provide an easily tractable system for analysis of mutant phenotypes. As

a single-cell-layer, protonemal tissue also enables high-resolution microscopy of

endogenously tagged fluorescent proteins in live cells (Rensing et al., 2020). In

addition to amenable tissue morphology and haploid lifestyle, *P. patens* is

amenable to transient RNA interference for rapid identification of gene function

and exhibits high rates of CRISPR Cas9-mediated genome editing (Rensing et al.,

2020; Mallett et al., 2019). Here, we utilized these molecular genetic and live-cell

imaging tools to systematically characterize the *Sec23* and *Sec24* gene families.

Analysis of mutant phenotypes revealed a high degree of functional specificity

among Sec23 and Sec24 isoforms. Focusing on the *Sec23* gene family, we

discovered that Sec23 isoforms differentially influence ER to Golgi trafficking,

and secretion to the plasma membrane as well as the size of presumptive ER exit

sites.

Results

Sec23 and Sec24 isoforms differentially affect tip growth

P. patens has seven *Sec23* and seven *Sec24* genes. Based on sequence

similarity, we found that there are four subclasses of *Sec23* and *Sec24* genes and

in both cases six of the seven genes exist in closely related pairs (Fig 1A, B). For

Sec23, with the exception of *Sec23A*, at least one member of each subclass is

highly expressed in protonemal cells (Fig 1A). For *Sec24*, we found that one of

the seven *Sec24* genes (*Sec24E)* was not expressed in protonemata (Fig1B). Of

the remaining six genes, *Sec24D* exhibited the lowest expression while the other

five Sec24 genes were expressed at similar levels (Fig 1B). To determine if Sec23

and Sec24 participate in protonemal growth, we used transient RNA interference

(RNAi) to silence the *Sec23* and *Sec24* gene families. We used a region of the

coding sequence from a member of each subclass (Fig1A, B, solid lines under gene models). These sequence regions share a high degree of sequence similarity to the other gene in each subclass (Fig 1A, B, dashed lines under gene models, Table S1) and would effectively simultaneously silence both genes. We did not target *Sec24E* since we were unable to detect a transcript.

By co-transforming the Sec23G RNAi construct together with a construct containing the target sequences for *Sec23A*, *C* and *D*, we targeted the entire *Sec23* gene family and discovered that silencing the seven *Sec23* genes leads to a dramatic reduction in growth (Fig 1C, D). Surprisingly, we found that silencing subclass III comprised of *Sec23D* and *Sec23E* resulted in a similarly severe growth defect (Fig 1C and D). In contrast, silencing any of the other subclasses did not affect growth (Fig 1C, D). These results demonstrated that there is functional specificity among *Sec23* isoforms, with only subclass III playing a critical role in protonemal growth, while the remaining five *Sec23* isoforms appear to be dispensable.

Similar to silencing subclass III *Sec23* genes, silencing *Sec24C* and *Sec24D* resulted in a 79% reduction in plant area (Fig 1E, F). Silencing *Sec24F* and *Sec24G* also impaired growth, but to a lesser extent resulting in a 48% reduction in plant area (Fig 1E, F). On the other hand, silencing *Sec24A* and *Sec24B* did not affect growth, although both genes were expressed at high levels. These data suggest that like Sec23, there is isoform specificity among *Sec24* genes.

Sec23 genes in subclass III are functionally redundant

To test whether *Sec23D* and *Sec23E* within subclass III are functionally redundant, we designed RNAi constructs targeting the untranslated regions (UTRs) of each gene (Fig 2A). Targeting the untranslated regions of both *Sec23D* and *Sec23E* resulted in small unpolarized plants with an 85% reduction in plant area as compared to control RNAi plants (Fig 2B, C). Silencing *Sec23D* alone resulted in a 59% reduction in plant area, while silencing *Sec23E* did not affect plant area (Fig 2B, C). Given that *Sec23D* is expressed seven times higher than *Sec23E* (Fig 1B), it is possible that *Sec23D* contributes to protonemal growth more than *Sec23E*. To test whether *Sec23E* can replace *Sec23D* function if expressed at sufficient levels, we performed complementation studies by driving

the full-length coding sequence of *Sec23D* or *Sec23E* with the constitutive maize

ubiquitin promoter. Interestingly, the reduction in plant area resulting from

silencing of *Sec23D* was restored to similar levels when either *Sec23D* or *Sec23E*

full-length cDNAs were expressed (Fig 2B, C), indicating that *Sec23D* and *Sec23E*

are functionally redundant, and that the highly expressed *Sec23D* is the

predominant group III isoform critical for protonemal growth.

CRISPR-Cas9-mediated deletion of Sec23 isoforms

Using transient RNAi, we were able to deduce that *Sec23* genes display

isoform specific functions during protonemal growth. However transient RNAi

assays are limited to phenotypic characterization of 7-day-old plants. To further

characterize the molecular functions of *Sec23*, we generated stable loss-of-

function mutants of *Sec23* genes using CRISPR-Cas9 mediated genome editing.

We targeted each gene with one protospacer located in the first coding exon (Fig

S1A, S2). We obtained single and higher-order mutants of *Sec23* genes carrying

out-of-frame mutations resulting in premature stop codons in their transcripts

(Fig S1A, S2, Table S2). Consistent with the RNAi results, 7-day-old *Δsec23d*

plants were 70% smaller than wild type plants while *Δsec23e* plants exhibited no

growth defects (Fig 3A, B). We isolated a number of *sec23d* null alleles (Fig S1A,

Table S2), which had similar growth phenotypes (Figure S1B). We also found

that in three-week-old plants regenerated from protoplasts, *Δsec23d-5* plants

were 62.6% smaller than wild type (Fig 3C, D), and Δ*sec23d-5* had fewer

gametophores compared to wild type (Fig 3C, E), suggesting that slow growing

protonemal tissues resulted in a developmental delay in the production of

gametophores.

Although it was possible to isolate many *Δsec23d* and *Δsec23e* single

mutants, we were unable to obtain *Δsec23de* double mutants either by targeting

Sec23D and *Sec23E* simultaneously or by targeting *Sec23D* in the *Δsec23e-6* single

mutant background. This suggests that deleting both *Sec23D* and *Sec23E* is

incompatible with protoplast regeneration, consistent with the severe growth

defects observed by simultaneously silencing *Sec23D* and *Sec23E* using RNAi.

Furthermore, we confirmed that none of the other *Sec23* genes were upregulated

in *Δsec23d-5* plants (Fig S1C), suggesting that the phenotypic consequences

observed were due specifically to loss of *Sec23D*.

Since the RNAi results suggested that all other *Sec23* subclasses were not

required for protonemal growth, we sought to isolate a stable line lacking

Sec23A, *Sec23B*, *Sec23C*, *Sec23F*, and *Sec23G* to determine if these genes

contribute to other aspects of plant growth and development. We successfully

isolated two independent quintuple mutant alleles *Δsec23abcfg-1* and

Δsec23abcfg-2, in which *Sec23A*, *Sec23B*, *Sec23C*, *Sec23F*, and *Sec23G* contained

lesions resulting in null alleles for each gene (Fig S2, Table S2). Most lesions were

small deletions or insertions. However, *Δsec23abcfg-1* contained a three

thousand base pair insertion in the *Sec23G* locus, still resulting in disruption of

the *Sec23G* locus (Fig S2, Table S2). We characterized both quintuple mutants

and found that while seven-day old *Δsec23abcfg-1* plants exhibited no detectable

growth defects, *Δsec23abcfg-2* plants exhibited a mild growth defect (Fig 3B, C).

However, three-week-old *Δsec23abcfg-1* plants were larger than wild type, and

Δsec23abcfg-2 plants were indistinguishable from wild type (Fig 3D, E).

Interestingly, both *Δsec23abcfg-1* and *Δsec23abcfg-2* developed more

gametophores than wild type (Fig 3E), suggesting that these five *Sec23* genes

may negatively regulate gametophore development.

Even though the two quintuple mutant isolates, exhibited statistically

significant differences at different times in development between each other and

wild type, these were minor compared to the differences between wild type and

all isolated *Δsec23d* null mutants. To investigate whether the observed

differences might result from differential upregulation of *Sec23D* or *Sec23E* in the

Δsec23abcfg mutants thereby compensating for the loss of *Sec23ABCFG*

functions, we measured the expression levels of *Sec23D* and *Sec23E* transcripts

in *Δsec23abcfg-1, Δsec23abcfg-2*, and wild type*.* In 8-day-old plants, we found

that neither *Sec23D* nor *Sec23E* transcripts were elevated compared to

expression in wild type (Fig 3F), indicating that minor differences observed

between the two quintuple mutants were not readily explained by increased

expression of either *Sec23D* or *E*. Taken together, the phenotypic analyses of

stable *sec23* null plants suggest that *Sec23A, B, C, F* and *G* do not contribute

substantially to establishment of juvenile moss tissues under laboratory growth

conditions. In contrast, *Sec23D* and *E* are essential.

Loss of Sec23D causes ER morphology defects and ER stress

As a predicted member of the COPII complex, Sec23D likely mediates ER-

to-Golgi transport. However, it is unclear whether the *Δsec23d* growth defect is a

result of a general impairment of ER-to-Golgi transport, or the lack of specific

cargo sorted by Sec23D needed for protonemal growth. Studies in other

organisms have shown that blocking COPII function alters ER morphology

(Novick et al., 1980) and results in elevated levels of unfolded proteins in the ER,

leading to activation of the ER unfolded protein response (Belden, 2001; Chung

et al., 2016). To analyze Sec23D's function in ER-to-Golgi transport, we

investigated whether loss of Sec23D affects ER morphology and the ER unfolded

protein response. We used CRISPR-Cas9 mediated genome editing to knock out

Sec23D in a line where the ER is fluorescently labeled by targeting GFP to the ER

lumen (Fig S1, Table S2). When imaged with a confocal microscope, *Δsec23d-B11*

cells exhibited numerous ER dots (Fig 4A, red and cyan boxes), and 65% of

protonemal filaments contained large ER aggregates (Fig 4A, green and yellow

boxes). Imaging with a super resolution spinning disc confocal system allowed

closer inspection of the aggregates (Fig. 4B). Smaller aggregates were clearly

comprised of accumulated tubules, while large aggregates appeared to have

tubules that wrapped around each other (Fig. 4B, Movie S1).

If changes in ER morphology result from defective secretion at ER exit

sites, then proteins may be inadvertently accumulating in the ER, resulting in

upregulation of the unfolded protein response. To test this, we analyzed the

expression of an ER heat shock protein (Bip), which is a well-known ER stress

reporter in *Arabidopsis thaliana* (Noh et al., 2003; Srivastava et al., 2013;

Maruyama et al., 2014; Cho and Kanehara, 2017). *P. patens* has two *Bip* paralogs,

Bip1 and *Bip2*. Since *Bip2* expression is low in protonemal tissue (Ortiz-Ramírez

et al., 2017), we tested whether *Bip1* is affected under conditions that induce ER

stress. Brefeldin A (BFA) inhibits protein secretion by disrupting ER-Golgi

integrity (Nebenfuhr, 2002; Niu et al., 2005). We found that after a 24-hour

treatment of BFA, expression of *Bip1* increased by about two-fold (Fig 4C),

indicating that *Bip1* is a plausible ER stress reporter in *P. patens*. Interestingly,

Bip1 expression increased 5-fold in *Δsec23d* compared to in wild type (Fig 4D),

demonstrating that *Δsec23d* suffers from severe ER stress. In contrast,

Δsec23abcfg-1, which didn't exhibit significant growth defects, had normal levels

of *Bip1* expression (Fig 4D), and normal ER morphology (Fig 4A).

Disrupting Sec23D results in reduced ER to Golgi trafficking and secretion to

the plasma membrane

ER morphology defects and elevated ER stress in *Δsec23d* suggest that

transport of secretory cargos might be blocked. We reasoned that proteins

destined to the Golgi and the plasma membrane would accumulate in the ER in

mutants with impaired ER to Golgi transport. To test this, we disrupted *Sec23D*

in lines that either express YFP fused to the first 49 amino acids of the soybean

α-1,2-mannosidase, a Golgi resident protein (YFP-MAN), (Nelson et al., 2007;

Furt et al., 2012) or an mCherry fusion of a transmembrane protein (F-SNAP-

mCherry) previously used to measure exocytosis (van Gisbergen et al., 2018)

(Fig S1, Table S2). Since it would have been time-consuming to regenerate

quintuple mutants in these two backgrounds, we instead transformed the YFP-

MAN and F-SNAP-mCherry markers into *Δsec23abcfg-1* plants. We selected lines

that had similar expression levels to control lines by measuring the total amount

of fluorescence in cells.

In control YFP-MAN lines, YFP signal accumulated in the Golgi with very

little fluorescence observed in the rest of the cell (Fig 5A). In contrast in *Δsec23d-*

F12, while YFP fluorescence still accumulated in the Golgi, a large portion of the

fluorescent signal that was presumably retained in the ER was observed

diffusely throughout the cell (Fig 5A). To quantify ER to Golgi trafficking

efficiency, we calculated the fluorescence intensity ratio of the signal in the Golgi

divided by the signal in the cytoplasm. A high ratio value indicates efficient

delivery to the Golgi, as observed in the control lines (Fig 5B). In contrast,

Δsec23d-F12 exhibited a significantly lower fluorescence intensity ratio (Fig 5B),

indicating that loss of Sec23D impairs delivery of YFP-MAN to the Golgi. Imaging

of *Δsec23abcfg-1* YFP-MAN and its control line revealed that the majority of the

YFP signal localizes to the Golgi with the Golgi to cytoplasm ratios in

Δsec23abcfg-1 indistinguishable from the control (Fig 5B), suggesting that

trafficking of YFP-MAN to the Golgi is not affected in the quintuple mutant.

Many proteins transported in COPII-coated vesicles ultimately reside at the

plasma membrane, are secreted, or are trafficked to the vacuole. Without a

fluorescent vacuolar cargo in hand, we instead investigated vacuolar

morphology by staining the vacuole with MDY-64, a dye that stains the tonoplast

in living plant cells (Scheuring et al., 2015). While we found that vacuolar

morphology was similar between wild type, *Δsec23d-5* and *Δsec23abcfg-1* plants

(Fig S3), we can not rule out that vacuolar protein sorting has not been affected.

To investigate trafficking to the plasma membrane, we used an engineered

fluorescent protein cargo, F-SNAP-mCherry, previously used to analyze

exocytosis in protonemata (van Gisbergen et al., 2018). In wild type cells, the F-

SNAP-mCherry accumulated on the plasma membrane, with only a very small

amount of fluorescence detected inside the cell (Fig 5C). In *Δsec23d-S2,* a large

portion of the mCherry signal was retained in the cell, labeling the ER network

(Fig 5C). To quantify secretion efficiency to the plasma membrane, we calculated

the fluorescence intensity ratio of the plasma membrane to the intensity within

the cell. We found that the ratio in *Δsec23d-S2* was significantly lower than that

of its control line (Fig 5D), indicating that loss of Sec23D inhibits secretion of

plasma membrane proteins*.*

Interestingly, we found that the F-SNAP-mCherry signal was also retained

inside the cell in the *Δsec23abcfg-1* mutant, readily observing labeling of the ER

network (Fig 5C). In addition, the fluorescence intensity ratio in *Δsec23abcfg-1*

was lower in comparison to the ratio in its control line (Fig 5D). However, the

ratio in *Δsec23abcfg-1* was still significantly higher than the ratio in *Δsec23d-S2*,

suggesting that secretion of plasma membrane proteins is impaired in

Δsec23abcfg-1 but to a lower extent than in *Δsec23d-S2*. These data together with

the lack of a growth defect in protonomata in the *Δsec23abcfg-1* mutant suggest

that Sec23A, B, C, F, and G contribute to protein secretion particularly to the

plasma membrane, but importantly their cargos are not essential for polarized

growth in protonemata.

Loss of Sec23D does not affect the cytoskeleton during tip growth

To investigate if defects in protein secretion, resulted in alterations to the cytoskeleton, critical for polarized expansion in protonemata, we imaged microtubules and actin in wild type and in cells carrying null mutations in *Sec23D*. Microtubules align along the length of the cell with their plus ends focused just below the cell apex (Hiwatashi et al., 2014), where a dynamic cluster of actin filaments forms predicting the site of cell expansion (Wu et al., 2018). Wild type and *Δsec23d-M4* cells expressing GFP-tubulin exhibit indistinguishable microtubule networks and both have a focal point of microtubules near the cell tip (Fig S4A, arrows). Using lifeact-mCherry to label the actin network, we found that both wild type and *Δsec23d-5* cells form an apical actin cluster that associates with the growing tip (Fig S4B). Time-lapse imaging revealed no discernable differences in the dynamics of the apical actin cluster between wild type and *Δsec23d-5* (Fig S4C, Movie S2). These data suggest that tip growth defects in cells lacking Sec23D are independent of the cytoskeleton.

Sec23 isoforms localize to discrete puncta associated with the ER

If Sec23 is a bona fide COPII component, then we would expect that Sec23 isoforms should associate with the ER. To test this, we inserted sequences in the genome of a line expressing GFP in the ER lumen encoding for three tandem mRuby2 (hereafter, 3XmRuby) in frame with the coding sequences of the three mostly highly expressed protonemal *Sec23* genes: *Sec23B*, *Sec23D* and *Sec23G* (Fig 1B, Fig S5A, B). To ensure that tagging *Sec23D* did not affect its function, we measured plant area from 7-day old regenerated protoplasts and found that *Sec23D-3XmRuby* grew indistinguishably from the control lines (Fig S5C), suggesting that Sec23D-mRuby is functional. By analogy, we reasoned that tagging *Sec23B* and *G* would also be functional. To ensure we did not observe any possible dominant negative impacts as a result of the tagging, we performed growth assays of 7-day old plants regenerated from protoplasts and observed that *Sec23B-3XmRuby* was indistinguishable from the control and *Sec23G-mRuby* was slightly smaller (Fig S5C).

Imaging with confocal microscopy revealed that Sec23D, Sec23B and

Sec23G localized to punctate structures throughout the cytoplasm (Fig 6A). All

three proteins labeled puncta that were often proximal to the ER (Fig 6A). Co-

localization measurements using Pearson's Correlation Coefficient suggested

that Sec23D, B and G puncta positively correlated with the ER. To ensure that

this correlation was significant, we flipped the ER image both vertically and

horizontally and repeated the correlation analysis and found that the correlation

coefficients were significantly lower (Fig 6B). Interestingly, Sec23G-3XmRuby

puncta were larger than the punctate structures found in Sec23D- and Sec23B-

3XmRuby lines (Fig 6A, C). To quantify possible size differences, we imaged all

three tagged Sec23 isoforms with a pixel size that maximizes the resolving

capacity of the imaging system (Fig 6C, see Materials and Methods) and utilized a

segmentation tool in Fiji that identifies particles in the images (Gilles et al.,

2017). We discovered that Sec23G particles are significantly larger than Sec23B

and D. Furthermore, Sec23D is slightly larger than Sec23B. These data suggest

that Sec23 isoforms form distinct compartments on the ER.

Sec23B and Sec23G localization is independent of Sec23D.

To further probe the nature of the Sec23 particles, we analyzed the

localization of Sec23B and Sec23G with respect to Sec23D. We used CRISPR-Cas9

mediated homology directed repair (HDR) to insert sequences encoding for

mNeonGreen (hereafter, mNeon) in-frame with Sec23B or Sec23G in a Sec23D

line endogenously tagged with a single mRuby2 (hereafter, mRuby) (Fig S6). As

expected Sec23B-mNeon formed small puncta whereas Sec23G-mNeon formed

larger puncta (Fig 7A, B). Imaging of Sec23B-mNeon and Sec23D-mRuby

revealed the presence of a distribution of cytoplasmic puncta, some with only

Sec23B-mNeon or only Sec23D-mRuby and some with both (Fig 7A). In contrast,

Sec23G-mNeon and Sec23D-mRuby were largely non-overlapping (Fig 7B). To

quantify the degree of overlap, we utilized the segmentation tool in Fiji that

identifies puncta and then determines the number of segmented particles that

physically overlap (Gilles et al., 2017). We found that 57% of Sec23B-mNeon co-

localized with Sec23-mRuby. In contrast, only 4% of Sec23G-mNeon showed

overlap with Sec23-mRuby (Fig 7C). In addition to identifying co-localizing

particles, the tool also measures the center-to-center distance of overlapping

particles (Gilles et al., 2017). Sec23G-mNeon and Sec23-mRuby had a

significantly larger center-to-center distance as compared to Sec23B-mNeon and

Sec23D-mRuby (Fig 7D), suggesting that any overlap between Sec23G and

Sec23D is coincidental.

While Sec23B and Sec23D exhibited a high degree of co-localization,

COPII vesicles are significantly smaller than the limit of resolution of a light

microscope. Therefore, it is unclear if a single vesicle contains both Sec23B and

Sec23D or if vesicles form with a single Sec23 isoform. Given that Sec23G

structures are distinct from Sec23D, and 40% of Sec23B and Sec23D puncta are

not overlapping, we hypothesized that Sec23B, Sec23D and Sec23G form distinct

ER exit sites. If this is the case, we would expect that Sec23B and Sec23G would

be unaffected by loss of Sec23D. Indeed, we discovered that when we deleted

Sec23D in the Sec23B-3XmRuby and Sec23G-3XmRuby lines, both Sec23B and

Sec23G still localized to puncta (Fig 7E). We quantified the size of the puncta and

found that Sec23G-3XmRuby was unaffected but Sec23B-3XmRuby were slightly

larger in the absence of Sec23D (Fig 7F). However, given that the average

Sec23B-3XmRuby particle size is essentially at the limit of resolution of our

system, it is unclear how significant this increase may be. Nevertheless, these

results demonstrate that even when ER morphology is affected due to loss of

Sec23D, Sec23B and Sec23G are largely unaffected, suggesting that Sec23B and

Sec23G ER exit sites are independent of Sec23D.

Discussion

The large number of genes encoding components of COPII in plants

together with functional studies in Arabidopsis (De Craene et al., 2014; Chung et

al., 2016; Barlow and Dacks, 2018; Brandizzi, 2018; Faso et al., 2009; Qu et al.,

2017; Tanaka et al., 2013) have led to the hypothesis that gene expansion

underlies increased functional diversification required for complex secretory

processes that pattern and build plant cells. Evidence suggests that functional

diversification could stem from tissue/developmental specific expression (Hino

et al., 2011; Lee et al.; Hanton et al., 2009) or from the formation of COPII

complexes comprised of specific isoforms with unique functions (Zeng et al.,

2015; Zeng et al., 2021). However, it remains unclear whether unique COPII

complexes mediate transport of specific cargos or accumulate at some ER exit

sites and not at others. Here, by analyzing the role of Sec23 and Sec24 isoforms

in protonemal development, we have provided evidence for formation of distinct

ER exit sites and the requirement of only Sec23D/E for transport of cargos

uniquely required for tip growth.

Using RNAi, we showed that *Sec23* and *Sec24* isoforms differentially

impact protonemal growth. Silencing of *Sec23D/E* or *Sec24C/D* similarly inhibit

tip growth suggesting that *Sec23D/E* may interact with *Sec24C/D*. However,

Sec24F/G also exhibited reduced growth. Thus, it is possible that a number of

heterodimer pairs may form that are critical for polarized growth. Protein-

protein interactions studies are needed to confirm if isoform specific

heterodimers form in moss. If this is the case, it suggests that isoform specificity

regulates COPII function similar to the finding in Arabidopsis that Sar1A and

Sec23A form a unique interaction mediating functional specificity (Zeng et al.,

2015).

Focusing on *Sec23*, where one of the Sec23 pairs, Sec23D/E, plays an

essential role in protonemal growth, we found several lines of evidence

suggesting that Sec23D functions as a bona fide COPII component. First, *Δsec23d*

plants had abnormal ER morphology and elevated ER stress consistent with a

block in cargo secretion from the ER. Second, trafficking to the Golgi and

secretion to the plasma membrane was reduced in *Δsec23d*. Third, Sec23D

localized to puncta near the ER. Due to differential expression levels, we found

that Sec23D is more important for protonemal growth than Sec23E. However,

when expressed at sufficient levels, Sec23E can rescue loss of Sec23D indicating

that Sec23D and Sec23E are functionally redundant. The inability to recover a

double mutant lacking both Sec23D and E suggests that together these two genes

are essential for moss viability. Given that viability in moss is measured in the

ability to generate protonemata, which is a tip-growing tissue, we reason that

Sec23D and E are essential for tip growth. While *Δsec23d* protonemal cells grow

significantly slower than wild type leading to smaller plants, we were unable to

identify any defects in either the actin or microtubules cytoskeletons, suggesting

that Sec23D contributes to polarized cell expansion independent of the

cytoskeleton. In fact, tip growth in *Δsec23d* resembles the recently identified *P.*

patens Δsabre mutant plants (Cheng & Bezanilla, 2021). SABRE associates with

regions of the ER and its absence has no effect on the cytoskeleton but rather

impacts ER morphology by accumulating aggregates that label with an ER

lumenal marker (Cheng & Bezanilla, 2021). Taken together these data suggest

that abrogated ER function impacts polarized growth without altering

cytoskeletal organization and dynamics.

With respect to gametophore formation, *Δsec23d* develops gametophores

late, but once gametophores form they are morphologically normal. This

suggests that Sec23D function is specifically required during tip growth. In

contrast, we found that the remaining five *Sec23* genes were dispensable for

formation of both protonemata and gametophores. One possible explanation for

this finding is that Sec23A, B, C, F and G have functionally diversified no longer

playing a role in ER to Golgi transport. In support of this hypothesis, unlike

Δsec23d, *Δsec23abcfg* plants did not exhibit ER morphology defects or increased

ER stress characteristic of cargo backing up in the ER often observed with COPII

loss of function mutants. However, we did observe secretion defects of plasma

membrane cargo in *Δsec23abcfg* plants. Given that Sec23B and G localize

proximal to the ER independent of Sec23D, it is possible that Sec23B and G

contribute to COPII function via pathways that may potentially bypass the Golgi

(Cheng et al., 2009; Zhang et al., 2011; Ding et al., 2012; Wang et al., 2017).

Alternatively, Sec23B and G may also function in ER to Golgi trafficking but, in

contrast to Sec23D, their cargos may be fewer and may not be essential for tip

growth.

Interestingly all the mutant combinations that we generated formed

morphologically normal gametophores. However, since we were unable to

generate *Δsec23de* or *Δsec23abcefg* plants, we can not distinguish between

Sec23E playing a critical role in gametophore formation versus Sec23D or

Sec23E acting redundantly with Sec23A, B, C, F and G for COPII secretion

required to shape gametophores. If, on the other hand, COPII is not required for

gametophore formation, then future experiments conditionally removing

Sec23D/E in *Δsec23abcfg* plants would be expected to form normal

gametophores and would be an excellent tool to identify COPII-independent

secretion pathways in plants, which are thought to play a role in transport of

tonoplast resident proteins such as the vacuolar H$^+$-ATPase, VHA-a3 (Viotti et al.,

2013).

While Sec23B, Sec23D, and Sec23G formed puncta associated with the ER,

these puncta were distinct. Sec23G formed large dots near the ER, whereas

Sec23D and Sec23B were smaller. Sec23B and Sec23D puncta exhibited a high

degree of overlap, suggesting that Sec23B and Sec23D may form vesicles in the

same region of the ER. In contrast Sec23G largely did not overlap with Sec23D

and by extension likely not Sec23B. Formation of larger domains on the ER

suggests that Sec23G may be involved in the secretion of bigger cargo and

provides evidence that moss has morphologically distinct ER exit sites, that can

form, even when ER morphology is disrupted, independent of Sec23D.

Given that Sec23D contributes to ER to Golgi trafficking and secretion to

the plasma membrane, while Sec23A, B, C, F and G contributes to plasma

membrane secretion, it is plausible to suggest that tip growth requires intact ER

to Golgi trafficking. In *Δsec23d* plants cargos essential for tip growth are either

inefficiently carried from the ER to the Golgi or are not properly processed in the

Golgi due to severe reductions in trafficking to the Golgi. Perhaps there is a larger

burden for carbohydrates synthesized by enzymes in the Golgi to generate a tip

growing cell wall as compared to a diffusely growing cell wall. To test this, future

studies will focus on identifying cargos that have been differentially affected in

Δsec23d versus *Δsec23abcfg* plants. Comparative mass spectrometry of isolated

microsomes and plasma membrane extracts in these mutant backgrounds could

be very informative. In addition, quantitatively measuring secretion of known

Golgi resident carbohydrate biosynthetic enzymes and plasma membrane

complexes such as cellulose and callose synthases in these mutants could help to

narrow down cargo specificity.

Taking advantage of the facile molecular genetic manipulation and

imaging in moss protonemal development, our study has shown that the

expanded *Sec23* gene family in moss is multi-faceted. We found evidence that the

Sec23 gene family exhibits aspects of functional redundancy, differential gene

expression, as well as isoform specificity for ER exit site formation, ER to Golgi

trafficking, and importantly transport of essential cargos required for tip growth.

Building upon the tools developed in this study, future studies will be able to

characterize the remaining COPII components, thereby providing a complete

picture of how gene expansions in COPII have been deployed.

Materials and Methods

Expression analyses

To quantify expression of the *Sec23* and *Sec24* genes family members

(Table S3), as well as *Bip1* (Pp3c10_17310), we isolated total RNA from 8-day old

plants (various genotypes, as needed for the particular experiment) regenerated

from protoplasts using the RNeasy Plant mini prep kit (Qiagen). cDNA was

prepared from 1µg of total RNA using SuperScriptIII reverse transcriptase kit,

following the manufacturer's recommendation (Invitrogen). qPCR assays were

conducted with Luna Universal qPCR Master Mix (New England Biolabs).

Primers for qRT-PCR analyses are listed in Table S4. After confirming that

amplification efficiencies (calculated from standard curves) were similar for

each set of primers (Table S4), relative expression levels were calculated, and

expression of Ubiquitin 10 was used for normalization. All experiments had

three technical replicates and three biological replicates.

Constructs

The Sec23 and Sec24 RNAi constructs were generated by PCR

amplification of either the coding sequence or 5' and 3' untranslated region

(UTR) of *Sec23* or *Sec24* from *P. patens* cDNA using primers indicated in Table S4.

PCR fragments were cloned into pENTR-D-TOPO (Invitrogen) following the

manufacturer's recommendations and the resulting vectors were sequenced. The

Sec23ACD-RNAi and Sec23 UTR-RNAi constructs were generated by stitching

together multiple PCR fragments and then directionally cloned into pENTR-D-

TOPO as previously described (Vidali et al., 2007). LR clonase (Invitrogen)

reactions were used to transfer either the coding sequence or 5' and 3' UTR

sequences the RNAi vector pUGGi (Bezanilla et al., 2005) generating the Sec23

UTRi or Sec24-CDSi constructs, respectively. Restriction enzyme digestion was

used to verify these constructs.

Expression constructs were generated by PCR amplifying *Sec23D* and

Sec23E coding sequences from *P. patens* cDNA using a 5' CACC sequence on the

specific primers for oriented cloning into the pENTR-D-TOPO vector

(Invitrogen). LR clonase (Invitrogen) reactions were used to transfer these

coding sequence into pTH-Ubi-Gate vectors (Vidali et al., 2007).

Sec23 tagging constructs were generated by PCR amplifying genomic

sequences upstream and downstream of the stop codon for *Sec23D*, *Sec23B*, and

Sec23G using the primers in Table S4. For *Sec23D*, we used homologous

recombination. The distance between the 5' and 3' homology arms was 500bp,

which when inserted correctly the tagging construct replaces these 500bp with

sequences encoding for the fluorescent protein and an antibiotic resistance

cassette (Fig S5, Fig S6). Using BP clonase (Invitrogen), homology arms were

transferred to vectors, generating homology arm entry clones. Four-fragment

recombination reactions were used to assemble the homology arms, fluorescent

protein sequence, and the antibiotic resistance cassette into pGEM-gate (Vidali et

al., 2009). For *Sec23B* and *Sec23G* tagging, we used CRISPR Cas9-mediated HDR.

5' and 3' homology arms upstream and downstream of the stop codon, with a

distance between these two arms of 38bp for *Sec23B* and 131bp for *Sec23G* were

PCR amplified from genomic DNA (Fig S5, Fig S6). Using BP clonase (Invitrogen),

homology arms were transferred to vectors, generating homology arm entry

clones, as described in (Mallett et al., 2019). Three-fragment recombination

reactions were used to assemble the homology arms and the fluorescent protein

sequence into pGEM-gate. We chose the CRISPR-Cas9 protospacer close to the

stop codon.

Protospacers for CRISPR-Cas9 mediated mutagenesis and HDR were

designed with the CRISPOR online software (crispor.tefor.net) (Haeussler et al.,

2016). Protospacers were selected with high specificity scores and low off-target

frequency. Specific overhangs sequences were added to facilitate downstream

cloning. Protospacers were synthesized as oligonucleotides (Table S4) and

annealed as described (Mallett et al., 2019). Annealed protospacer fragments

were transferred to vectors using Instant Sticky-End Ligation Master Mix (New

England Biolabs). *Sec23B* and *G* protospacers designed to tag these genes via

HDR as well as *Sec23A*, *D* and *E* protospacers designed to knock out these genes

were ligated into pENTR-PpU6-sgRNA-L1L2 (Mallett et al., 2019). *Sec23B* and *F*

protospacers designed to knock out these genes were ligated into pENTR-PpU6-

sgRNA-L1L5 (Mallett et al., 2019). Sec23C and G protospacers designed to knock

out these genes were ligated into pENTR-PpU6-sgRNA-R5L2 (Mallett et al.,

2019). LR clonase (Invitrogen) was used to assemble the protospacer containing

vectors into pZeo-Cas9-Gate, pMH-Cas9-Gate, or pMK-Cas9-Gate vectors (Mallett

et al., 2019), as needed. Two-fragment recombination reactions were used to

assemble dual targeting mutagenesis vectors for *Sec23B* and *C*, and for *Sec23F*

and *G*. Restriction enzyme digestion and sequencing was used to verify these

constructs.

Tissue propagation, transformation, and genotyping

All moss lines were propagated weekly by moderate homogenization in

water and pipetting onto a permeable cellophane covered solid media, as

described previously (Wu and Bezanilla, 2014). Tissue was grown at room

temperature in Percival growth chambers, under 85 $\mu mol_{photons}/m^2s$ light with

long-day conditions. PEG-mediated transformations were performed as

previously described to generate stable lines (Wu and Bezanilla, 2014), CRISPR-

Cas9 genome edited lines (Mallett et al., 2019) and perform transient RNAi

(Vidali et al., 2007).

To genotype mutants generated by CRISPR-Cas9 mutagenesis, we

extracted DNA from plants that were 3-4 weeks old (0.5-1cm in diameter) using

the protocol described in (Augustine et al., 2011). For editing experiments, we

used primers (Table S4) surrounding the expected Cas9 cleavage site (~300-400

bp on each side). T7 endonuclease assay was used to screen candidate mutants

as described (Mallett et al., 2019). For homology-directed repair tagging and

homologous recombination tagging, we used primers (Table S4) outside of the

homology region to avoid amplification of residual DNA donor template.

Imaging and morphometric analysis of growth assays

For transient RNAi, complementation analyses, and growth assays of

stable lines, plants regenerated from protoplasts were photographed seven days

after protoplasting. For transient RNAi and complementation analyses plants

lacking nuclear GFP signal were imaged because these are the plants that are

actively silencing as described previously (Bezanilla et al., 2005; Vidali et al.,

2007). For growth assays of stable lines any regenerated plant was imaged.

Images were acquired with a 1X objective using a stereomicroscope (Leica

MZ16FA or Nikon SMZ25) equipped with a CCD camera (Leica DF300FX or

Nikon digital sight DS-Fi2). Chlorophyll fluorescence and any nuclear GFP signal

were acquired simultaneously using filters (480/40, dichroic 505 (Leica) or

dichroic 510 (Nikon), emission 510 long pass). Exposure settings were

maintained constant throughout an experiment. 3-week old plants regenerated

from protoplasts were imaged with white light on a Nikon SMZ25

stereomicroscope equipped with a Nikon digital sight DS-Fi2 color camera.

To quantify plant area, individual plants were cropped from images and

the red channel, representing chlorophyll fluorescence of the plant, was

separated from the RGB image. Threshold settings were set manually for all

plants within an experiment. To quantify brightfield images of 3-week-old plants,

the blue channel, representing chlorophyll fluorescence of the plant, was

separated from the RGB image and thresholded. Total area was calculated from

the largest thresholded object in the selected window. All images analysis was

done using macros (Galotto et al., 2019) written for ImageJ. Analysis of variation

(ANOVA) for multiple comparisons was performed with Kaleidagraph (Synergy)

using the Tukey HSD post hoc tests. The alpha for statistical significance was set

to 0.05.

Laser scanning confocal microscope imaging and quantification

For imaging Sec23B/D/G localization and dynamics, 5 to 8-day old plants

regenerated from protoplasts were placed onto an agar pad in Hoagland's buffer

(4mM KNO_3, 2mM KH_2PO_4, 1mM $Ca(NO_3)_2$, 89μM Fe citrate, 300μM $MgSO_4$,

9.93μM H_3BO_3, 220nM $CuSO_4$, 1.966μM $MnCl_2$, 231nM $CoCl_2$, 191nM $ZnSO4$,

169nM KI, 103nM Na_2MoO_4, and 1% sucrose), covered by a glass cover slip and

sealed with VALAP (1:1:1 parts of Vaseline, lanoline, and paraffin). For imaging

ER morphology, Golgi resident protein secretion, F-SNAP-mCherry secretion, we

used microfluidic imaging chambers (Bascom et al., 2016). Ground protonemal

tissue was gently pipetted into the central part of the device followed by an

infusion of Hoagland's medium. Then the chamber was submerged Hoagland's

medium, and placed under constant 85 $μmol_{photons}/m^2s$ light.

Images were acquired on a Nikon A1R confocal microscope system with a

1.49 NA 60X oil immersion objective (Nikon) at room temperature. Image

acquisition was controlled by NIS-Element AR 4.1 software (Nikon). Laser

illumination at 488 nm was used for exciting mNeon, YFP, GFP and chlorophyll

autofluorescence; 561 nm for mRuby, mCherry. Emission filters were 525/50

nm for mNeon, YFP, and GFP; 595/50 nm for mRuby and mCherry.

Super resolution spinning disc confocal imaging was performed with a

Nikon CSUW1-SoRA system. Images were acquired with a 60x Plan Apo 1.40NA

objective with the 2.8x mag lens. Laser illumination at 488 nm was used for

exciting GFP, and the Emission filter were 525/50. Images were denoised using

the denoise.ai machine learning algorithm and deconvolved using the

Richardson-Lucy algorithm within the NIS-Elements v5.21 software.

For quantification of trafficking to the Golgi, confocal Z-stacks of

individual cells were converted to a maximum intensity projection. For

quantification of secretion to the plasma membrane, the medial focal plane was

selected for analysis. YFP (YFP-MAN) and mCherry (FSnap-mCherry)

fluorescence intensities with the same expression pattern were normalized

using enhanced contrast in Fiji. Images were cropped to isolate regions only

within the cell. Then, using a minimum threshold of 20,000-22,000 it was

possible to isolate and measure the fluorescence intensity of the Golgi structures

(Golgi intensity) or the plasma membrane with analyze particles in Fiji. The

fluorescence intensity of the background was similarly measured using the same

threshold cut-off for each image but unchecking dark background, thereby

selecting all regions below the threshold cut off. These values were used to

calculate the ratio of Golgi/plasma membrane intensity to background intensity.

To quantify the Pearson's Correlation Coefficient, we analyzed single focal planes

with the Coloc2 plugin in Fiji comparing the original images as well as

horizontally and vertically flipped ER-GFP images. To quantify the size and co-

localization of Sec23B, D and G dots, we analyzed single focal planes and used the

diAna tool for ImageJ, which uses object-based identification for distance and co-

localization measurements (Gilles et al., 2017) and was recently used to define

specific subdomains of the trans Golgi network in plant cells (Heinze et al.,

2020). All images were acquired with a pixel size of 0.07 μm and processed by

subtracting background, enhancing contrast with normalization, and smoothing

in Fiji prior to performing segmentation with diAna. For Sec23G-mNeon,

chloroplast autofluorescence was collected in the far-red channel and subtracted

from the green channel. To segment the dots in the image, we used the iterative

option with a minimum threshold of 25,000-27,500 and a step value of 20. After

segmentation, the diAna analyse module was used to measure the size of all

segmented regions of an image, as well as to determine the percent of co-

localizing particles. For particles that did overlap, we also extracted the center-

to-center distance of those overlapping particles. For all data, analysis of

variation (ANOVA) for multiple comparisons was performed with Kaleidagraph

(Synergy) using the Tukey HSD post hoc tests. The alpha for statistical

significance was set to 0.05. For pairwise comparisons, a student-T test was used

using Kaleidagraph (Synergy).

Author contributions

M.C. and M.B. designed the research. M.C., S.W., J.E.O. and S.E.R performed

the research. M.C., S.W. and M.B. wrote the manuscript.

Acknowledgements

We thank members of the Bezanilla lab for careful reading of the

manuscript. M.C. received support from the Plant Biology Graduate Program at

the University of Massachusetts, Amherst. This work was supported by

Dartmouth College and grants from the National Science Foundation (MCB-

1330171 and MCB-1715785 to M.B.).

References

Aridor, M. (2018). COPII gets in shape: Lessons derived from morphological
aspects of early secretion. Traffic 19: 823–839.
737

738 Aridor, M., Fish, K.N., Bannykh, S., Weissman, J., Roberts, T.H., Lippincott-
739 Schwartz, J., and Balch, W.E. (2001). The Sar1 Gtpase Coordinates
740 Biosynthetic Cargo Selection with Endoplasmic Reticulum Export Site
741 Assembly. Journal of Cell Biology 152: 213–230.
742

Aridor, M., Weissman, J., Bannykh, S., Nuoffer, C., and Balch, W.E. (1998). Cargo
Selection by the COPII Budding Machinery during Export from the ER.
Journal of Cell Biology 141: 61–70.

Augustine, R.C., Pattavina, K.A., Tüzel, E., Vidali, L., and Bezanilla, M. (2011). Actin
interacting protein1 and actin depolymerizing factor drive rapid actin
dynamics in Physcomitrella patens. Plant Cell 23: 3696–3710.
749
750 Barlow, L.D. and Dacks, J.B. (2018). Seeing the endomembrane system for the
751   trees: Evolutionary analysis highlights the importance of plants as models
752   for eukaryotic membrane-trafficking. Seminars in Cell & Developmental
753   Biology 80: 142–152.
754
755 Barlowe, C. (2003). Signals for COPII-dependent export from the ER: what's the
756   ticket out? Trends in Cell Biology 13: 295–300.
757
758 Barlowe, C. and Helenius, A. (2016). Cargo Capture and Bulk Flow in the Early
759   Secretory Pathway. Annu. Rev. Cell Dev. Biol. 32: 197–222.
760
761 Barlowe, C. and Schekman, R. (1993). SEC12 encodes a guanine-nucleotide-
762   exchange factor essential for transport vesicle budding from the ER.
763   Nature 365: 347–349.
764
765 Bascom, C.S., Wu, S.-Z., Nelson, K., Oakey, J., and Bezanilla, M. (2016). Long-Term
766   Growth of Moss in Microfluidic Devices Enables Subcellular Studies in
767   Development. Plant Physiology 172: 28–37.
768
769 Belden, W.J. (2001). Role of Erv29p in Collecting Soluble Secretory Proteins into
770   ER-Derived Transport Vesicles. Science 294: 1528–1531.
771
772 Bezanilla, M., Perroud, P.-F., Pan, A., Klueh, P., and Quatrano, R.S. (2005). An RNAi
773   system in Physcomitrella patens with an internal marker for silencing
774   allows for rapid identification of loss of function phenotypes. Plant Biol
775   (Stuttg) 7: 251–257.
776
777 Bi, X., Corpina, R.A., and Goldberg, J. (2002). Structure of the Sec23/24–Sar1 pre-
778   budding complex of the COPII vesicle coat. Nature 419: 271–277.
779
780 Brandizzi, F. (2018). Transport from the endoplasmic reticulum to the Golgi in
781   plants: Where are we now? Seminars in Cell & Developmental Biology 80:
782   94–105.
783
784 Brandizzi, F. and Barlowe, C. (2013). Organization of the ER–Golgi interface for
785   membrane traffic control. Nat Rev Mol Cell Biol 14: 382–392.
786
787 Cheng, F-y., Zamski, E., Guo W-w., Mason Pharr, D., and Williamson, J.D. (2009).
788   Salicylic acid stimulates secretion of the normally symplastic enzyme
789   mannitol dehydrogenase: a possible defense against mannitol-secreting
790   fungal pathogens. Planta 230: 1093–1103.
791
Cheng, X., and Bezanilla M. (2021). SABRE populates ER domains essential for
cell plate maturation and cell expansion influencing cell and tissue
patterning. eLife 10: e65166.

Cho, Y. and Kanehara, K. (2017). Endoplasmic Reticulum Stress Response in
Arabidopsis Roots. Front. Plant Sci. 8.

Chung, K.P., Zeng, Y., and Jiang, L. (2016). COPII Paralogs in Plants: Functional
Redundancy or Diversity? Trends in Plant Science 21: 758–769.

De Craene, J.-O., Courte, F., Rinaldi, B., Fitterer, C., Herranz, M.C., Schmitt-
Keichinger, C., Ritzenthaler, C., and Friant, S. (2014). Study of the Plant
COPII Vesicle Coat Subunits by Functional Complementation of Yeast
Saccharomyces cerevisiae Mutants. PLoS ONE 9: e90072.

Ding, Y., Wang, J., Wang, J., Stierhof, Y-D., Robinson, D.G., and Jiang L. (2012).
Unconventional protein secretion. Trends in Plant Science 17: 606-615.

Faso, C., Chen, Y.-N., Tamura, K., Held, M., Zemelis, S., Marti, L., Saravanan, R.,
Hummel, E., Kung, L., Miller, E., Hawes, C., and Brandizzi, F. (2009). A
Missense Mutation in the Arabidopsis COPII Coat Protein Sec24A Induces
the Formation of Clusters of the Endoplasmic Reticulum and Golgi
Apparatus. The Plant Cell 21: 3655–3671.

Fath, S., Mancias, J.D., Bi, X., and Goldberg, J. (2007). Structure and Organization
of Coat Proteins in the COPII Cage. Cell 129: 1325–1336.

Furt, F., Lemoi, K., Tüzel, E., and Vidali, L. (2012). Quantitative analysis of
organelle distribution and dynamics in Physcomitrella patens protonemal
cells. BMC Plant Biol. 12: 70.

Galotto, G., Bibeau, J.P., and Vidali, L. (2019). Automated Image Acquisition and
Morphological Analysis of Cell Growth Mutants in Physcomitrella patens.
In Plant Cell Morphogenesis, F. Cvrčková and V. Žárský, eds, Methods in
Molecular Biology. (Springer New York: New York, NY), pp. 307–322.

Gilles, J.-F., Dos Santos, M., Boudier, T., Bolte, S., and Heck, N. (2017). DiAna, an
ImageJ tool for object-based 3D co-localization and distance analysis.
Methods 115: 55–64.

Giraudo, C.G. and Maccioni, H.J.F. (2003). Endoplasmic Reticulum Export of
Glycosyltransferases Depends on Interaction of a Cytoplasmic Dibasic
Motif with Sar1. MBoC 14: 3753–3766.

van Gisbergen, P.A.C., Wu, S.-Z., Chang, M., Pattavina, K.A., Bartlett, M.E., and
Bezanilla, M. (2018). An ancient Sec10-formin fusion provides insights
into actin-mediated regulation of exocytosis. J. Cell Biol. 217: 945–957.

Haeussler, M., Schönig, K., Eckert, H., Eschstruth, A., Mianné, J., Renaud, J.-B.,
Schneider-Maunoury, S., Shkumatava, A., Teboul, L., Kent, J., Joly, J.-S., and
Concordet, J.-P. (2016). Evaluation of off-target and on-target scoring

algorithms and integration into the guide RNA selection tool CRISPOR.
Genome Biol. 17: 148.

Hanna, M.G., Peotter, J.L., Frankel, E.B., and Audhya, A. (2018). Membrane
Transport at an Organelle Interface in the Early Secretory Pathway: Take
Your Coat Off and Stay a While: Evolution of the metazoan early secretory
pathway. BioEssays 40: 1800004.

Hanton, S.L., Matheson, L.A., Chatre, L., and Brandizzi, F. (2009). Dynamic
organization of COPII coat proteins at endoplasmic reticulum export sites
in plant cells. The Plant Journal 57: 963–974.

Heinze, L., Freimuth, N., Rößling, A.-K., Hahnke, R., Riebschläger, S., Fröhlich, A.,
Sampathkumar, A., McFarlane, H.E., and Sauer, M. (2020). EPSIN1 and
MTV1 define functionally overlapping but molecularly distinct trans-Golgi
network subdomains in Arabidopsis. PNAS 117: 25880–25889.

Hino, T., Tanaka, Y., Kawamukai, M., Nishimura, K., Mano, S., and Nakagawa, T.
(2011). Two Sec13p Homologs, AtSec13A and AtSec13B, Redundantly
Contribute to the Formation of COPII Transport Vesicles in *Arabidopsis*
thaliana. Bioscience, Biotechnology, and Biochemistry 75: 1848–1852.

Hiwatashi, Y., Sato, Y., and Doonan, J.H. (2014). Kinesins Have a Dual Function in
Organizing Microtubules during Both Tip Growth and Cytokinesis in
Physcomitrella patens. The Plant Cell 26: 1256–1266.

Jensen, D. and Schekman, R. (2011). COPII-mediated vesicle formation at a
glance. Journal of Cell Science 124: 1–4.

Khoriaty, R. et al. (2018). Functions of the COPII gene paralogs SEC23A and
SEC23B are interchangeable in vivo. Proc Natl Acad Sci USA 115: E7748–
E7757.

Lee, M.H., Lee, S.H., Kim, H., Jin, J.B., Kim, D.H., and Hwang, I. A WD40 Repeat
Protein, Arabidopsis Sec13 Homolog 1, May Play a Role in Vacuolar
Trafficking by Controlling the Membrane Association of AtDRP2A.: 10.

Mallett, D.R., Chang, M., Cheng, X., and Bezanilla, M. (2019). Efficient and modular
CRISPR-Cas9 vector system for Physcomitrella patens. Plant Direct 3:
e00168-15.

Mancias, J.D. and Goldberg, J. (2007). The Transport Signal on Sec22 for
Packaging into COPII-Coated Vesicles Is a Conformational Epitope.
Molecular Cell 26: 403–414.

Maruyama, D., Sugiyama, T., Endo, T., and Nishikawa, S. (2014). Multiple BiP
Genes of Arabidopsis thaliana are Required for Male Gametogenesis and
Pollen Competitiveness. Plant and Cell Physiology 55: 801–810.

Mossessova, E., Bickford, L.C., and Goldberg, J. (2003). SNARE selectivity of the
COPII coat. Cell 114: 483–495.
894

895   Nakano, A. and Muramatsu, M. (1989). A Novel GTP-binding Protein, Sarlp, Is
896   Involved in Transport from the Endoplasmic Reticulum to the Golgi
897   Apparatus. The Journal of Cell Biology 109: 2677–2691.
898

899   Nakano, R.T., Matsushima, R., Ueda, H., Tamura, K., Shimada, T., Li, L., Hayashi, Y.,
900   Kondo, M., Nishimura, M., and Hara-Nishimura, I. (2009). GNOM-
901   LIKE1/ERMO1 and SEC24a/ERMO2 Are Required for Maintenance of
902   Endoplasmic Reticulum Morphology in Arabidopsis thaliana. The Plant
903   Cell 21: 3672–3685.
904

905   Nebenfuhr, A. (2002). Brefeldin A: Deciphering an Enigmatic Inhibitor of
906   Secretion. Plant physiology 130: 1102–1108.
907

908   Nelson, B.K., Cai, X., and Nebenführ, A. (2007). A multicolored set of in vivo
909   organelle markers for co-localization studies in Arabidopsis and other
910   plants. Plant J. 51: 1126–1136.
911

912   Niu, T.-K., Pfeifer, A.C., Lippincott-Schwartz, J., and Jackson, C.L. (2005). Dynamics
913   of GBF1, a Brefeldin A-Sensitive Arf1 Exchange Factor at the Golgi□V.
914   Molecular Biology of the Cell 16: 1213–1222.
915

916   Noh, S.-J., Kwon, C.S., Oh, D.-H., Moon, J.S., and Chung, W.-I. (2003). Expression of
917   an evolutionarily distinct novel BiP gene during the unfolded protein
918   response in Arabidopsis thaliana. Gene 311: 81–91.
919

920   Novick, P., Field, C., and Schekman, R. (1980). Identification of 23
921   complementation groups required for post-translational events in the
922   yeast secretory pathway. Cell 21: 205–215.
923

924   Nufer, O. (2003). ER export of ERGIC-53 is controlled by cooperation of targeting
925   determinants in all three of its domains. Journal of Cell Science 116:
926   4429–4440.
927

928   Ortiz-Ramírez, C., Michard, E., Simon, A.A., Damineli, D.S.C., Hernandez-Coronado,
929   M., Becker, J.D., and Feijó, J.A. (2017). GLUTAMATE RECEPTOR-LIKE
930   channels are essential for chemotaxis and reproduction in mosses. Nature
931   549: 91–95.
932

933   Qu, X., Zhang, R., Zhang, M., Diao, M., Xue, Y., and Huang, S. (2017). Organizational
934   innovation of apical actin filaments drives rapid pollen tube growth and
935   turning. Mol Plant 10: 930–947.
936

937   Rensing, S.A., Goffinet, B., Meyberg, R., Wu, S.-Z., and Bezanilla, M. (2020). The
938   Moss *Physcomitrium* (*Physcomitrella*) *patens* : A Model Organism for
939   Non-Seed Plants. Plant Cell 32: 1361–1376.
940

941 Robinson, D.G., Brandizzi, F., Hawes, C., and Nakano, A. (2015). Vesicles versus
942   Tubes: Is Endoplasmic Reticulum-Golgi Transport in Plants
943   Fundamentally Different from Other Eukaryotes? Plant Physiology 168:
944   393–406.
945
946 Salama, N.R., Yeung, T., and Schekman, R.W. The Secl3p complex and
947   reconstitution of vesicle budding from the ER with purified cytosolic
948   proteins. The EMBO Journal 12: 4073–4082.
949
950 Scheuring, D., Schöller, M., Kleine-Vehn, J., and Löfke, C. (2015). Vacuolar staining
951   methods in plant cells. Methods Mol Biol 1242: 83–92.
952
953 Schlacht, A. and Dacks, J.B. (2015). Unexpected Ancient Paralogs and an
954   Evolutionary Model for the COPII Coat Complex. Genome Biology and
955   Evolution 7: 1098–1109.
956
957 Srivastava, R., Deng, Y., Shah, S., Rao, A.G., and Howell, S.H. (2013). BINDING
958   PROTEIN Is a Master Regulator of the Endoplasmic Reticulum Stress
959   Sensor/Transducer bZIP28 in *Arabidopsis*. Plant Cell 25: 1416–1429.
960
961 Stagg, S.M., Gürkan, C., Fowler, D.M., LaPointe, P., Foss, T.R., Potter, C.S.,
962   Carragher, B., and Balch, W.E. (2006). Structure of the Sec13/31 COPII
963   coat cage. Nature 439: 234–238.
964
965 Stagg, S.M., LaPointe, P., Razvi, A., Gürkan, C., Potter, C.S., Carragher, B., and Balch,
966   W.E. (2008). Structural Basis for Cargo Regulation of COPII Coat
967   Assembly. Cell 134: 474–484.
968
969 Takagi, J., Renna, L., Takahashi, H., Koumoto, Y., Tamura, K., Stefano, G., Fukao, Y.,
970   Kondo, M., Nishimura, M., Shimada, T., Brandizzi, F., and Hara-Nishimura,
971   I. (2013). MAIGO5 Functions in Protein Export from Golgi-Associated
972   Endoplasmic Reticulum Exit Sites in Arabidopsis. The Plant Cell 25: 4658–
973   4675.
974
975 Tanaka, Y., Nishimura, K., Kawamukai, M., Oshima, A., and Nakagawa, T. (2013).
976   Redundant function of two Arabidopsis COPII components, AtSec24B and
977   AtSec24C, is essential for male and female gametogenesis. Planta 238:
978   561–575.
979
980 Vidali, L., Augustine, R.C., Kleinman, K.P., and Bezanilla, M. (2007). Profilin is
981   essential for tip growth in the moss Physcomitrella patens. Plant Cell 19:
982   3705–3722.
983
984 Vidali, L., van Gisbergen, P.A.C., Guérin, C., Franco, P., Li, M., Burkart, G.M.,
985   Augustine, R.C., Blanchoin, L., and Bezanilla, M. (2009). Rapid formin-
986   mediated actin-filament elongation is essential for polarized plant cell
987   growth. Proc. Natl. Acad. Sci. U.S.A. 106: 13341–13346.
988

989   Viotti, C. et al. (2013). The Endoplasmic Reticulum Is the Main Membrane Source
990   for Biogenesis of the Lytic Vacuole in *Arabidopsis*. Plant Cell 25: 3434–
991   3449.
992
993   Wang, X., Chung, K. P., Lin, W., and Jiang L. (2017). Protein secretion in plants:
994   conventional and unconventional pathways and new techniques. Journal
995   of Experimental Botany 69: 21-37.
996
997   Whittle, J.R.R. and Schwartz, T.U. (2010). Structure of the Sec13–Sec16 edge
998   element, a template for assembly of the COPII vesicle coat. The Journal of
999   Cell Biology 190: 347–361.
1000
1001   Wu, S.-Z. and Bezanilla, M. (2014). Myosin VIII associates with microtubule ends
1002   and together with actin plays a role in guiding plant cell division. eLife
1003   Sciences 3: e03498.
1004
1005   Wu, S.-Z., Yamada, M., Mallett, D.R., and Bezanilla, M. (2018). Cytoskeletal
1006   discoveries in the plant lineage using the moss Physcomitrella patens.
1007   Biophys Rev 10: 1683–1693.
1008
1009   Zeng, Y., Chung, K.P., Li, B., Lai, C.M., Lam, S.K., Wang, X., Cui, Y., Gao, C., Luo, M.,
1010   Wong, K.-B., Schekman, R., and Jiang, L. (2015). Unique COPII component
1011   AtSar1a/AtSec23a pair is required for the distinct function of protein ER
1012   export in *Arabidopsis thaliana*. Proc Natl Acad Sci USA 112: 14360–14365.
1013
1014   Zeng, Y., Li, B., Ji, C., Feng, L., Niu, F., Deng, C., Chen, S., Lin, Y., Cheung, K.C.P., Shen,
1015   J., Wong, K.-B., and Jiang, L. (2021). A unique AtSar1D-AtRabD2a nexus
1016   modulates autophagosome biogenesis in Arabidopsis thaliana. Proc Natl
1017   Acad Sci U S A 118.
1018
1019   Zhang, H., Zhang, L., Gao, B., Fan, H., Jin, J., Botella, M.A., Jiang, L., and Lin, J.
1020   (2011). Golgi apparatus-localized synaptotagmin 2 is required for
1021   unconventional secretion in Arabidopsis. PLoS ONE 6: e26477.
1022

Figure Legends

Figure 1. *Sec23* and *Sec24* are required for polarized growth. Gene models of

the *P. patens Sec23* (A) and *Sec24* (B) genes are shown with exons indicated by

boxes and introns by thin black lines. Coding and untranslated regions are

denoted by thick and thin boxes, respectively. The lines underneath the gene

models represent sequence regions targeted by RNAi constructs. Scale bar is

500bp. Graphs depict the relative expression of *Sec23* and *Sec24* genes

normalized to UBIQUITIN10 in 8-day-old wild-type moss plants regenerated

from protoplasts. N = 3, Error bars are s.e.m. (C, E) Representative chlorophyll

autofluorescence images of 7-day-old plants regenerated from protoplasts

expressing the indicated RNAi constructs. Scale bar, 100 µm. (D, F)

Quantification of plant area is based on the area of the chlorophyll

autofluorescence and is presented normalized to the area of the control for each

experiment. Means are indicated by horizontal lines. Letters indicate groups with

significantly different means as determined by a one-way ANOVA with a Tukey

post hoc test (α=0.05).

1039

Figure 2. *Sec23D* and *Sec23E* are functionally redundant. (A) *Sec23D* and

Sec23E gene models are shown with exons indicated by boxes and introns by

thin black lines. Coding and untranslated regions are denoted by thick and thin

boxes, respectively. The lines underneath the gene models represent sequence

regions that were targeted by *Sec23D* or *Sec23E* UTR RNAi constructs. Scale bar

is 500bp. (B) Representative chlorophyll autofluorescence images of 7-day-old

plants regenerated from protoplasts expressing the indicated constructs. Scale

bar, 100 µm. (C) Quantification of plant area was normalized to the control for

each experiment. Letters indicate groups with significantly different means as

determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

1050

Figure 3. CRISPR-Cas9-mediated deletion of *Sec23* isoforms demonstrates

that *Sec23D* is critical for protonemal growth. (A) Representative chlorophyll

autofluorescence images of 7-day old plants with the indicated genotype

regenerated from protoplasts. Scale bar, 100µm. (B) Quantification of plant area

normalized to the wild type control. (C) Representative 3-week old moss plants

with the indicated genotype. Scale bar, 1mm. (D) Quantification of the plant area

of 3-week old moss plants normalized to that of the wild type control. (E)

Number of gametophores in 3-week old moss plants. Letters in (B, D-E) indicate

groups with significantly different means as determined by a one-way ANOVA

with a Tukey *post hoc* test (α=0.05). (F) Relative expression of *Sec23D* and *E*

genes normalized to UBIQUITIN10 in 8-day-old wild type and *Δsec23abcfg-1*, and

Δsec23abcfg-2 moss plants regenerated from protoplasts. N=3, Error bars are

s.e.m. No significant difference was determined by a one-way ANOVA with a

Tukey *post hoc* test (α=0.05).

1065

Figure 4. Loss of Sec23D causes ER morphology defects and ER stress. (A) Laser scanning confocal images of the ER labeled with GFP-KDEL in the indicated genotype. Colored boxes highlight large and small ER aggregates found in the *Δsec23d-B11* mutant. Scale bars, 5 μm. (B) Spinning disc confocal super resolution imaging of the ER labeled with GFP-KDEL in a *Δsec23d-B11* cell. Scale bar, 5 μm. Box outlined with yellow dotted line is enlarged to the right and the box outlined with a white dotted line is enlarged below with optical sections spaced every 0.5 μm shown for this region of the cell. Scale bars in boxed regions, 2 μm. Also see Movie S1. (C, D) Relative expression of *PpBip1* in 8-day old moss plants regenerated from protoplasts. N=3, Error bars are s.e.m. Asterisks denote t-probability is <0.01. n.s. denotes no significant difference. In (B) BFA-treated plants were normalized to DMSO-treated plants and in (C) null mutants were normalized to wild type plants.

Figure 5. Deletion of *Sec23D* inhibits Golgi trafficking and secretion to the plasma membrane. (A) Representative maximum projections of confocal images of the indicated genotypes expressing a YFP-MAN marker. Scale bar, 10 μm. (B) Quantification of the fluorescence intensity ratio within the Golgi to the background. (C) Representative medial sections of confocal images of the indicated genotypes expressing a plasma membrane marker (FSnap-mCherry). Scale bar, 10 μm. (D) Quantification of the fluorescence intensity ratio of the plasma membrane to the intensity within the cell. Letters in (B, D) indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 6. Sec23 isoforms localize to ER proximal structures with distinct sizes. (A) Representative confocal Z-sections of lines where the indicated Sec23 gene was endogenously tagged with 3XmRuby in a line that expresses GFP-KDEL labeling the ER lumen (ER-GFP). Scale bars, 5 μm. Boxed region in the mRuby channel is shown to the right of each image. (B) Pearson's correlation coefficient measured between the Sec23 labeled lines and ER-GFP. O, original ER image; V, vertically flipped ER image; H, horizontally flipped ER image. Asterisks denote p<0.001as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05)

(C) Representative confocal Z-sections of *Sec23* genes endogenously tagged with 3XmRuby acquired with the optimal resolution of the confocal imaging system. Scale bars, 5 µm. (D) Quantification of average dot size in the Sec23 images as determined by the diAna tool for Fiji (Gilles et al., 2017). Asterisks denote $p < 0.001$ as determined by a one-way ANOVA with a Tukey *post hoc* test ($\alpha = 0.05$).

Figure 7. Sec23B and Sec23G form ER exit sites independent of Sec23D.
Representative confocal Z-sections of Sec23D-mRuby lines where Sec23B (A) or Sec23G (B) were endogenously tagged with mNeon. Scale bar, 5 µm. Yellow boxed regions are enlarged to the right of the merged image where Sec23D-mRuby is false-colored magenta and in (A) Sec23B-mNeon is false-colored green and in (B) Sec23G-mNeon is false-colored green. (C) Percent of Sec23B-mNeon or Sec23G-mNeon dots that co-localized with Sec23D-mRuby and (D) center-to-center distance of co-localized particles as determined using the diAna tool for Fiji (Gilles et al., 2017). (E) Representative confocal Z-sections of wild type and *Δsec23d-5* cells with either Sec23B or Sec23G endogenously tagged with 3XmRuby. Scale bar, 5 µm. (F) Quantification of average dot size as determined by the diAna tool for Fiji (Gilles et al., 2017). Sec23B and Sec23G dot size in wild type is the same data presented in Figure 6D. Student t-test was performed for data in (C, D, and F). The t-probability is indicated for each data pair in each graph. n.s. denotes no significant difference.

Figure 1. *Sec23* and *Sec24* are required for polarized growth. Gene models of the *P. patens* *Sec23* (A) and *Sec24* (B) genes are shown with exons indicated by boxes and introns by thin black lines. Coding and untranslated regions are denoted by thick and thin boxes, respectively. The lines underneath the gene models represent sequence regions targeted by RNAi constructs. Scale bar is 500bp. Graphs depict the relative expression of *Sec23* and *Sec24* genes normalized to UBIQUITIN10 in 8-day-old wild-type moss plants regenerated from protoplasts. N = 3, Error bars are s.e.m. (C, E) Representative chlorophyll autofluorescence images of 7-day-old plants regenerated from protoplasts expressing the indicated RNAi constructs. Scale bar, 100 µm. (D, F) Quantification of plant area is based on the area of the chlorophyll autofluorescence and is presented normalized to the area of the control for each experiment. Means are indicated by horizontal lines. Letters indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test ($\alpha=0.05$).

Figure 2. *Sec23D* and *Sec23E* are functionally redundant. (A) *Sec23D* and *Sec23E* gene models are shown with exons indicated by boxes and introns by thin black lines. Coding and untranslated regions are denoted by thick and thin boxes, respectively. The lines underneath the gene models represent sequence regions that were targeted by *Sec23D* or *Sec23E* UTR RNAi constructs. Scale bar is 500bp. (B) Representative chlorophyll autofluorescence images of 7-day-old plants regenerated from protoplasts expressing the indicated constructs. Scale bar, 100 μm. (C) Quantification of plant area was normalized to the control for each experiment. Letters indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 3. CRISPR-Cas9-mediated deletion of *Sec23* isoforms demonstrates that *Sec23D* is critical for protonemal growth. (A) Representative chlorophyll autofluorescence images of 7-day old plants with the indicated genotype regenerated from protoplasts. Scale bar, 100μm. (B) Quantification of plant area normalized to the wild type control. (C) Representative 3-week old moss plants with the indicated genotype. Scale bar, 1mm. (D) Quantification of the plant area of 3-week old moss plants normalized to that of the wild type control. (E) Number of gametophores in 3-week old moss plants. Letters in (B, D-E) indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05). (F) Relative expression of *Sec23D* and *E* genes normalized to UBIQUITIN10 in 8-day-old wild type and Δ*sec23abcfg-1*, and Δ*sec23abcfg-2* moss plants regenerated from protoplasts. N=3, Error bars are s.e.m. No significant difference was determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 4. Loss of Sec23D causes ER morphology defects and ER stress. (A) Laser scanning confocal images of the ER labeled with GFP-KDEL in the indicated genotype. Colored boxes highlight large and small ER aggregates found in the Δ*sec23d-B11* mutant. Scale bars, 5 µm. (B) Spinning disc confocal super resolution imaging of the ER labeled with GFP-KDEL in a Δ*sec23d-B11* cell. Scale bar, 5 µm. Box outlined with yellow dotted line is enlarged to the right and the box outlined with a white dotted line is enlarged below with optical sections spaced every 0.5 µm shown for this region of the cell. Scale bars in boxed regions, 2 µm. Also see Movie S1. (C, D) Relative expression of *PpBip1* in 8-day old moss plants regenerated from protoplasts. N=3, Error bars are s.e.m. Asterisks denote t-probability is <0.01. n.s. denotes no significant difference. In (B) BFA-treated plants were normalized to DMSO-treated plants and in (C) null mutants were normalized to wild type plants.

Figure 5. Deletion of *Sec23D* inhibits Golgi trafficking and secretion to the plasma membrane.
(A) Representative maximum projections of confocal images of the indicated genotypes expressing a YFP-MAN marker. Scale bar, 10 µm. (B) Quantification of the fluorescence intensity ratio within the Golgi to the background. (C) Representative medial sections of confocal images of the indicated genotypes expressing a plasma membrane marker (FSnap-mCherry). Scale bar, 10 µm. (D) Quantification of the fluorescence intensity ratio of the plasma membrane to the intensity within the cell. Letters in (B, D) indicate groups with significantly different means as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 6. Sec23 isoforms localize to ER proximal structures with distinct sizes. (A) Representative confocal Z-sections of lines where the indicated Sec23 gene was endogenously tagged with 3XmRuby in a line that expresses GFP-KDEL labeling the ER lumen (ER-GFP). Scale bars, 5 µm. Boxed region in the mRuby channel is shown to the right of each image. (B) Pearson's correlation coefficient measured between the Sec23 labeled lines and ER-GFP. O, original ER image; V, vertically flipped ER image; H, horizontally flipped ER image. Asterisks denote $p<0.001$ as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05) (C) Representative confocal Z-sections of *Sec23* genes endogenously tagged with 3XmRuby acquired with the optimal resolution of the confocal imaging system. Scale bars, 5 µm. (D) Quantification of average dot size in the Sec23 images as determined by the diAna tool for Fiji (Gilles et al., 2017). Asterisks denote $p<0.001$ as determined by a one-way ANOVA with a Tukey *post hoc* test (α=0.05).

Figure 7. Sec23B and Sec23G form ER exit sites independent of Sec23D. Representative confocal Z-sections of Sec23D-mRuby lines where Sec23B (A) or Sec23G (B) were endogenously tagged with mNeon. Scale bar, 5 μm. Yellow boxed regions are enlarged to the right of the merged image where Sec23D-mRuby is false-colored magenta and in (A) Sec23B-mNeon is false-colored green and in (B) Sec23G-mNeon is false-colored green. (C) Percent of Sec23B-mNeon or Sec23G-mNeon dots that co-localized with Sec23D-mRuby and (D) center-to-center distance of co-localized particles as determined using the diAna tool for Fiji (Gilles et al., 2017). (E) Representative confocal Z-sections of wild type and Δ*sec23d-5* cells with either Sec23B or Sec23G endogenously tagged with 3XmRuby. Scale bar, 5 μm. (F) Quantification of average dot size as determined by the diAna tool for Fiji (Gilles et al., 2017). Sec23B and Sec23G dot size in wild type is the same data presented in Figure 6D. Student t-test was performed for data in (C, D, and F). The t-probability is indicated for each data pair in each graph. n.s. denotes no significant difference.

Parsed Citations

Aridor, M. (2018). COPII gets in shape: Lessons derived from morphological aspects of early secretion. Traffic 19: 823–839.
Google Scholar: Author Only Title Only Author and Title

Aridor, M., Fish, K.N., Bannykh, S., Weissman, J., Roberts, T.H., Lippincott-Schwartz, J., and Balch, W.E. (2001). The Sar1 Gtpase Coordinates Biosynthetic Cargo Selection with Endoplasmic Reticulum Export Site Assembly. Journal of Cell Biology 152: 213–230.
Google Scholar: Author Only Title Only Author and Title

Aridor, M., Weissman, J., Bannykh, S., Nuoffer, C., and Balch, W.E. (1998). Cargo Selection by the COPII Budding Machinery during Export from the ER. Journal of Cell Biology 141: 61–70.
Google Scholar: Author Only Title Only Author and Title

Augustine, R.C., Pattavina, K.A, Tüzel, E., Vidali, L., and Bezanilla, M. (2011). Actin interacting protein1 and actin depolymerizing factor drive rapid actin dynamics in Physcomitrella patens. Plant Cell 23: 3696–3710.
Google Scholar: Author Only Title Only Author and Title

Barlow, L.D. and Dacks, J.B. (2018). Seeing the endomembrane system for the trees: Evolutionary analysis highlights the importance of plants as models for eukaryotic membrane-trafficking. Seminars in Cell & Developmental Biology 80: 142–152.
Google Scholar: Author Only Title Only Author and Title

Barlowe, C. (2003). Signals for COPII-dependent export from the ER: what's the ticket out? Trends in Cell Biology 13: 295–300.
Google Scholar: Author Only Title Only Author and Title

Barlowe, C. and Helenius, A. (2016). Cargo Capture and Bulk Flow in the Early Secretory Pathway. Annu. Rev. Cell Dev. Biol. 32: 197–222.
Google Scholar: Author Only Title Only Author and Title

Barlowe, C. and Schekman, R. (1993). SEC12 encodes a guanine-nucleotide-exchange factor essential for transport vesicle budding from the ER. Nature 365: 347–349.
Google Scholar: Author Only Title Only Author and Title

Bascom, C.S., Wu, S.-Z., Nelson, K., Oakey, J., and Bezanilla, M. (2016). Long-Term Growth of Moss in Microfluidic Devices Enables Subcellular Studies in Development. Plant Physiology 172: 28–37.
Google Scholar: Author Only Title Only Author and Title

Belden, W.J. (2001). Role of Erv29p in Collecting Soluble Secretory Proteins into ER-Derived Transport Vesicles. Science 294: 1528–1531.
Google Scholar: Author Only Title Only Author and Title

Bezanilla, M., Perroud, P.-F., Pan, A, Klueh, P., and Quatrano, R.S. (2005). An RNAi system in Physcomitrella patens with an internal marker for silencing allows for rapid identification of loss of function phenotypes. Plant Biol (Stuttg) 7: 251–257.
Google Scholar: Author Only Title Only Author and Title

Bi, X., Corpina, R.A., and Goldberg, J. (2002). Structure of the Sec23/24–Sar1 pre-budding complex of the COPII vesicle coat. Nature 419: 271–277.
Google Scholar: Author Only Title Only Author and Title

Brandizzi, F. (2018). Transport from the endoplasmic reticulum to the Golgi in plants: Where are we now? Seminars in Cell & Developmental Biology 80: 94–105.
Google Scholar: Author Only Title Only Author and Title

Brandizzi, F. and Barlowe, C. (2013). Organization of the ER–Golgi interface for membrane traffic control. Nat Rev Mol Cell Biol 14: 382–392.
Google Scholar: Author Only Title Only Author and Title

Cheng, F-y., Zamski, E., Guo W-w., Mason Pharr, D., and Williamson, J.D. (2009). Salicylic acid stimulates secretion of the normally symplastic enzyme mannitol dehydrogenase: a possible defense against mannitol-secreting fungal pathogens. Planta 230: 1093–1103.
Google Scholar: Author Only Title Only Author and Title

Cheng, X., and Bezanilla M. (2021). SABRE populates ER domains essential for cell plate maturation and cell expansion influencing cell and tissue patterning. eLife 10: e65166.
Google Scholar: Author Only Title Only Author and Title

Cho, Y. and Kanehara, K. (2017). Endoplasmic Reticulum Stress Response in Arabidopsis Roots. Front. Plant Sci. 8.
Google Scholar: Author Only Title Only Author and Title

Chung, K.P., Zeng, Y., and Jiang, L. (2016). COPII Paralogs in Plants: Functional Redundancy or Diversity? Trends in Plant Science 21: 758–769.
Google Scholar: Author Only Title Only Author and Title

De Craene, J.-O., Courte, F., Rinaldi, B., Fitterer, C., Herranz, M.C., Schmitt-Keichinger, C., Ritzenthaler, C., and Friant, S. (2014). Study of the Plant COPII Vesicle Coat Subunits by Functional Complementation of Yeast Saccharomyces cerevisiae Mutants. PLoS ONE 9:

e90072.
Google Scholar: Author Only Title Only Author and Title

Ding, Y., Wang, J., Wang, J., Stierhof, Y-D., Robinson, D.G., and Jiang L. (2012). Unconventional protein secretion. Trends in Plant Science 17: 606-615.
Google Scholar: Author Only Title Only Author and Title

Faso, C., Chen, Y.-N., Tamura, K., Held, M., Zemelis, S., Marti, L., Saravanan, R., Hummel, E., Kung, L., Miller, E., Hawes, C., and Brandizzi, F. (2009). A Missense Mutation in the Arabidopsis COPII Coat Protein Sec24A Induces the Formation of Clusters of the Endoplasmic Reticulum and Golgi Apparatus. The Plant Cell 21: 3655–3671.
Google Scholar: Author Only Title Only Author and Title

Fath, S., Mancias, J.D., Bi, X., and Goldberg, J. (2007). Structure and Organization of Coat Proteins in the COPII Cage. Cell 129: 1325–1336.
Google Scholar: Author Only Title Only Author and Title

Furt, F., Lemoi, K., Tüzel, E., and Vidali, L. (2012). Quantitative analysis of organelle distribution and dynamics in Physcomitrella patens protonemal cells. BMC Plant Biol. 12: 70.
Google Scholar: Author Only Title Only Author and Title

Galotto, G., Bibeau, J.P., and Vidali, L. (2019). Automated Image Acquisition and Morphological Analysis of Cell Growth Mutants in Physcomitrella patens. In Plant Cell Morphogenesis, F. Cvrčková and V. Žárský, eds, Methods in Molecular Biology. (Springer New York: New York, NY), pp. 307–322.
Google Scholar: Author Only Title Only Author and Title

Gilles, J.-F., Dos Santos, M., Boudier, T., Bolte, S., and Heck, N. (2017). DiAna, an ImageJ tool for object-based 3D co-localization and distance analysis. Methods 115: 55–64.
Google Scholar: Author Only Title Only Author and Title

Giraudo, C.G. and Maccioni, H.J.F. (2003). Endoplasmic Reticulum Export of Glycosyltransferases Depends on Interaction of a Cytoplasmic Dibasic Motif with Sar1. MBoC 14: 3753–3766.
Google Scholar: Author Only Title Only Author and Title

van Gisbergen, P.A.C., Wu, S.-Z, Chang, M., Pattavina, K.A, Bartlett, M.E., and Bezanilla, M. (2018). An ancient Sec10-formin fusion provides insights into actin-mediated regulation of exocytosis. J. Cell Biol. 217: 945–957.
Google Scholar: Author Only Title Only Author and Title

Haeussler, M., Schönig, K., Eckert, H., Eschstruth, A, Mianné, J., Renaud, J.-B., Schneider-Maunoury, S., Shkumatava, A, Teboul, L., Kent, J., Joly, J.-S., and Concordet, J.-P. (2016). Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. Genome Biol. 17: 148.
Google Scholar: Author Only Title Only Author and Title

Hanna, M.G., Peotter, J.L., Frankel, E.B., and Audhya, A. (2018). Membrane Transport at an Organelle Interface in the Early Secretory Pathway: Take Your Coat Off and Stay a While: Evolution of the metazoan early secretory pathway. BioEssays 40: 1800004.
Google Scholar: Author Only Title Only Author and Title

Hanton, S.L., Matheson, L.A, Chatre, L., and Brandizzi, F. (2009). Dynamic organization of COPII coat proteins at endoplasmic reticulum export sites in plant cells. The Plant Journal 57: 963–974.
Google Scholar: Author Only Title Only Author and Title

Heinze, L., Freimuth, N., Rößling, A-K., Hahnke, R., Riebschläger, S., Fröhlich, A, Sampathkumar, A, McFarlane, H.E., and Sauer, M. (2020). EPSIN1 and MTV1 define functionally overlapping but molecularly distinct trans-Golgi network subdomains in Arabidopsis. PNAS 117: 25880–25889.
Google Scholar: Author Only Title Only Author and Title

Hino, T., Tanaka, Y., Kawamukai, M., Nishimura, K., Mano, S., and Nakagawa, T. (2011). Two Sec13p Homologs, AtSec13A and AtSec13B, Redundantly Contribute to the Formation of COPII Transport Vesicles in Arabidopsis thaliana. Bioscience, Biotechnology, and Biochemistry 75: 1848–1852.
Google Scholar: Author Only Title Only Author and Title

Hiwatashi, Y., Sato, Y., and Doonan, J.H. (2014). Kinesins Have a Dual Function in Organizing Microtubules during Both Tip Growth and Cytokinesis in Physcomitrella patens. The Plant Cell 26: 1256–1266.
Google Scholar: Author Only Title Only Author and Title

Jensen, D. and Schekman, R. (2011). COPII-mediated vesicle formation at a glance. Journal of Cell Science 124: 1–4.
Google Scholar: Author Only Title Only Author and Title

Khoriaty, R. et al. (2018). Functions of the COPII gene paralogs SEC23A and SEC23B are interchangeable in vivo. Proc Natl Acad Sci USA 115: E7748–E7757.
Google Scholar: Author Only Title Only Author and Title

Lee, M.H., Lee, S.H., Kim, H., Jin, J.B., Kim, D.H., and Hwang, I. A WD40 Repeat Protein, Arabidopsis Sec13 Homolog 1, May Play a Role in Vacuolar Trafficking by Controlling the Membrane Association of AtDRP2A: 10.

Mallett, D.R., Chang, M., Cheng, X., and Bezanilla, M. (2019). Efficient and modular CRISPR-Cas9 vector system for Physcomitrella patens. Plant Direct 3: e00168-15.
  Google Scholar: Author Only Title Only Author and Title

Mancias, J.D. and Goldberg, J. (2007). The Transport Signal on Sec22 for Packaging into COPII-Coated Vesicles Is a Conformational Epitope. Molecular Cell 26: 403–414.
  Google Scholar: Author Only Title Only Author and Title

Maruyama, D., Sugiyama, T., Endo, T., and Nishikawa, S. (2014). Multiple BiP Genes of Arabidopsis thaliana are Required for Male Gametogenesis and Pollen Competitiveness. Plant and Cell Physiology 55: 801–810.
  Google Scholar: Author Only Title Only Author and Title

Mossessova, E., Bickford, L.C., and Goldberg, J. (2003). SNARE selectivity of the COPII coat. Cell 114: 483–495.
  Google Scholar: Author Only Title Only Author and Title

Nakano, A. and Muramatsu, M. (1989). A Novel GTP-binding Protein, Sar1p, Is Involved in Transport from the Endoplasmic Reticulum to the Golgi Apparatus. The Journal of Cell Biology 109: 2677–2691.
  Google Scholar: Author Only Title Only Author and Title

Nakano, R.T., Matsushima, R., Ueda, H., Tamura, K., Shimada, T., Li, L., Hayashi, Y., Kondo, M., Nishimura, M., and Hara-Nishimura, I. (2009). GNOM-LIKE1/ERMO1 and SEC24a/ERMO2 Are Required for Maintenance of Endoplasmic Reticulum Morphology in Arabidopsis thaliana. The Plant Cell 21: 3672–3685.
  Google Scholar: Author Only Title Only Author and Title

Nebenfuhr, A. (2002). Brefeldin A: Deciphering an Enigmatic Inhibitor of Secretion. Plant physiology 130: 1102–1108.
  Google Scholar: Author Only Title Only Author and Title

Nelson, B.K., Cai, X., and Nebenführ, A. (2007). A multicolored set of in vivo organelle markers for co-localization studies in Arabidopsis and other plants. Plant J. 51: 1126–1136.
  Google Scholar: Author Only Title Only Author and Title

Niu, T.-K., Pfeifer, A.C., Lippincott-Schwartz, J., and Jackson, C.L. (2005). Dynamics of GBF1, a Brefeldin A-Sensitive Arf1 Exchange Factor at the Golgi□V. Molecular Biology of the Cell 16: 1213–1222.
  Google Scholar: Author Only Title Only Author and Title

Noh, S.-J., Kwon, C.S., Oh, D.-H., Moon, J.S., and Chung, W.-I. (2003). Expression of an evolutionarily distinct novel BiP gene during the unfolded protein response in Arabidopsis thaliana. Gene 311: 81–91.
  Google Scholar: Author Only Title Only Author and Title

Novick, P., Field, C., and Schekman, R. (1980). Identification of 23 complementation groups required for post-translational events in the yeast secretory pathway. Cell 21: 205–215.
  Google Scholar: Author Only Title Only Author and Title

Nufer, O. (2003). ER export of ERGIC-53 is controlled by cooperation of targeting determinants in all three of its domains. Journal of Cell Science 116: 4429–4440.
  Google Scholar: Author Only Title Only Author and Title

Ortiz-Ramírez, C., Michard, E., Simon, A.A., Damineli, D.S.C., Hernandez-Coronado, M., Becker, J.D., and Feijó, J.A. (2017). GLUTAMATE RECEPTOR-LIKE channels are essential for chemotaxis and reproduction in mosses. Nature 549: 91–95.
  Google Scholar: Author Only Title Only Author and Title

Qu, X., Zhang, R., Zhang, M., Diao, M., Xue, Y., and Huang, S. (2017). Organizational innovation of apical actin filaments drives rapid pollen tube growth and turning. Mol Plant 10: 930–947.
  Google Scholar: Author Only Title Only Author and Title

Rensing, S.A., Goffinet, B., Meyberg, R., Wu, S.-Z., and Bezanilla, M. (2020). The Moss Physcomitrium (Physcomitrella) patens : A Model Organism for Non-Seed Plants. Plant Cell 32: 1361–1376.
  Google Scholar: Author Only Title Only Author and Title

Robinson, D.G., Brandizzi, F., Hawes, C., and Nakano, A. (2015). Vesicles versus Tubes: Is Endoplasmic Reticulum-Golgi Transport in Plants Fundamentally Different from Other Eukaryotes? Plant Physiology 168: 393–406.
  Google Scholar: Author Only Title Only Author and Title

Salama, N.R., Yeung, T., and Schekman, R.W. The Sec13p complex and reconstitution of vesicle budding from the ER with purified cytosolic proteins. The EMBO Journal 12: 4073–4082.
  Google Scholar: Author Only Title Only Author and Title

Scheuring, D., Schöller, M., Kleine-Vehn, J., and Löfke, C. (2015). Vacuolar staining methods in plant cells. Methods Mol Biol 1242: 83–92.
  Google Scholar: Author Only Title Only Author and Title

Schlacht, A. and Dacks, J.B. (2015). Unexpected Ancient Paralogs and an Evolutionary Model for the COPII Coat Complex. Genome Biology and Evolution 7: 1098–1109.

Google Scholar: Author Only Title Only Author and Title

Srivastava, R., Deng, Y., Shah, S., Rao, A.G., and Howell, S.H. (2013). BINDING PROTEIN Is a Master Regulator of the Endoplasmic Reticulum Stress Sensor/Transducer bZIP28 in Arabidopsis. Plant Cell 25: 1416–1429.

Stagg, S.M., Gürkan, C., Fowler, D.M., LaPointe, P., Foss, T.R., Potter, C.S., Carragher, B., and Balch, W.E. (2006). Structure of the Sec13/31 COPII coat cage. Nature 439: 234–238.

Stagg, S.M., LaPointe, P., Razvi, A, Gürkan, C., Potter, C.S., Carragher, B., and Balch, W.E. (2008). Structural Basis for Cargo Regulation of COPII Coat Assembly. Cell 134: 474–484.

Takagi, J., Renna, L., Takahashi, H., Koumoto, Y., Tamura, K., Stefano, G., Fukao, Y., Kondo, M., Nishimura, M., Shimada, T., Brandizzi, F., and Hara-Nishimura, I. (2013). MAIGO5 Functions in Protein Export from Golgi-Associated Endoplasmic Reticulum Exit Sites in Arabidopsis. The Plant Cell 25: 4658–4675.

Tanaka, Y., Nishimura, K., Kawamukai, M., Oshima, A, and Nakagawa, T. (2013). Redundant function of two Arabidopsis COPII components, AtSec24B and AtSec24C, is essential for male and female gametogenesis. Planta 238: 561–575.

Vidali, L., Augustine, R.C., Kleinman, K.P., and Bezanilla, M. (2007). Profilin is essential for tip growth in the moss Physcomitrella patens. Plant Cell 19: 3705–3722.

Vidali, L., van Gisbergen, P.A.C., Guérin, C., Franco, P., Li, M., Burkart, G.M., Augustine, R.C., Blanchoin, L., and Bezanilla, M. (2009). Rapid formin-mediated actin-filament elongation is essential for polarized plant cell growth. Proc. Natl. Acad. Sci. U.S.A. 106: 13341–13346.

Viotti, C. et al. (2013). The Endoplasmic Reticulum Is the Main Membrane Source for Biogenesis of the Lytic Vacuole in Arabidopsis. Plant Cell 25: 3434–3449.

Wang, X., Chung, K. P., Lin, W., and Jiang L. (2017). Protein secretion in plants: conventional and unconventional pathways and new techniques. Journal of Experimental Botany 69: 21-37.

Whittle, J.R.R. and Schwartz, T.U. (2010). Structure of the Sec13–Sec16 edge element, a template for assembly of the COPII vesicle coat. The Journal of Cell Biology 190: 347–361.

Wu, S.-Z. and Bezanilla, M. (2014). Myosin VIII associates with microtubule ends and together with actin plays a role in guiding plant cell division. eLife Sciences 3: e03498.

Wu, S.-Z, Yamada, M., Mallett, D.R., and Bezanilla, M. (2018). Cytoskeletal discoveries in the plant lineage using the moss Physcomitrella patens. Biophys Rev 10: 1683–1693.

Zeng, Y., Chung, K.P., Li, B., Lai, C.M., Lam, S.K., Wang, X., Cui, Y., Gao, C., Luo, M., Wong, K.-B., Schekman, R., and Jiang, L. (2015). Unique COPII component AtSar1a/AtSec23a pair is required for the distinct function of protein ER export in Arabidopsis thaliana. Proc Natl Acad Sci USA 112: 14360–14365.

Zeng, Y., Li, B., Ji, C., Feng, L., Niu, F., Deng, C., Chen, S., Lin, Y., Cheung, K.C.P., Shen, J., Wong, K.-B., and Jiang, L. (2021). A unique AtSar1D-AtRabD2a nexus modulates autophagosome biogenesis in Arabidopsis thaliana. Proc Natl Acad Sci U S A 118.

Zhang, H., Zhang, L., Gao, B., Fan, H., Jin, J., Botella, M.A, Jiang, L., and Lin, J. (2011). Golgi apparatus-localized synaptotagmin 2 is required for unconventional secretion in Arabidopsis. PLoS ONE 6: e26477.