b3galt6 knock-out zebrafish recapitulate β3GalT6-deficiency disorders in human and reveal a trisaccharide proteoglycan linkage region

Sarah Delbaere[a], Adelbert De Clercq[a], Shuji Mizumoto[b], Fredrik Noborn[c,d], Jan Willem Bek[a], Lien Alluyn[a], Charlotte Gistelinck[e], Delfien Syx[a], Phil L. Salmon[f], Paul J. Coucke[a], Göran Larson[c,d], Shuhei Yamada[b], Andy Willaert[a,1], Fransiska Malfait[a,1,2]

[a]Center for Medical Genetics Ghent, Department of Biomolecular Medicine, Ghent University and Ghent University Hospital, Ghent, Belgium.

[b]Department of Pathobiochemistry, Faculty of Pharmacy, Meijo University, 150 Yagotoyama, Tempaku-ku, Nagoya, Aichi 468-8503, Japan.

[c]Department of Laboratory Medicine, Sahlgrenska Academy at the University of Gothenburg, Gothenburg, Sweden.

[d]Laboratory of Clinical Chemistry, Sahlgrenska University Hospital, Gothenburg, Sweden.

[e]Department of Orthopaedics and Sports Medicine, University of Washington, Seattle, WA98195.

[f]Bruker microCT, Kontich, Belgium.

[1]AW and FM contributed equally to this work.

[2]Corresponding author:

Fransiska Malfait

Center for Medical Genetics, Ghent University Hospital, 0K5

Corneel Heymanslaan 10, B-9000 Ghent, Belgium

Tel: +32-9-332 36 03

E-mail: Fransiska.Malfait@UGent.be

Keywords

b3galt6, zebrafish, trisaccharide linkage region, proteoglycans, linkeropathies

Author Contributions

SD, AW and FM designed the study; SD, ADC, SM, FN, JWB, LA and CG performed research; SM, PLS, GL, SY contributed new reagents or analytic tools; SD, ADC, SM, FN, JWB, LA, CG, DS, PLS, PJC, GL, SY, AW and FM analyzed data; SD, AW and FM wrote the paper; all authors reviewed the paper.

Abstract

Proteoglycans are structurally and functionally diverse biomacromolecules found abundantly on cell membranes and in the extracellular matrix. They consist of a core protein linked to glycosaminoglycan chains via a tetrasaccharide linkage region. Here, we show that CRISPR/Cas9-mediated *b3galt6* knock-out zebrafish, lacking galactosyltransferase II, which adds the third sugar in the linkage region, largely recapitulate the phenotypic abnormalities seen in human β3GalT6-deficiency disorders. These comprise craniofacial dysmorphism, generalized skeletal dysplasia, skin involvement and indications for muscle hypotonia. Disturbed collagen fibril organization was the most consistent ultrastructural characteristic throughout affected tissues. Strikingly, despite a strong reduction in glycosaminoglycan content, we identified a small amount of proteoglycans containing a unique linkage region consisting of only three sugars. This implies that formation of glycosaminoglycans with an immature linkage region is possible in a pathogenic context. Our study therefore unveils a novel rescue mechanism for β3GalT6-deficiency, opening new avenues for therapeutic intervention.

Introduction

Proteoglycans (PGs) comprise a group of complex biomacromolecules, consisting of a core protein and one or multiple glycosaminoglycan (GAG) side chains, that are ubiquitously expressed in the extracellular matrix (ECM) and on cell surfaces and basement membranes in vertebrate and invertebrate species (1-4). Besides their structural role, PGs act as mediators between the ECM and intracellular signaling pathways and contribute to development and tissue homeostasis, influencing a variety of cellular processes including cell fate determination, cell proliferation, migration, adhesion, differentiation and survival (5).

The PG superfamily is, depending on the composition of the GAG chains, subdivided into heparan sulfate (HS) and chondroitin/dermatan sulfate (CS/DS) PGs. These GAG chains are linear polysaccharides, composed of a repeated disaccharide unit consisting of an uronic acid and amino sugar. HSPGs, such as perlecan, syndecan and glypican, consist of repeating glucuronic acid (GlcA)- *N*-acetyl-glucosamine (GlcNAc) disaccharides. CSPGs consist of repeating GlcA- *N*-acetyl-galactosamine (GalNAc) disaccharides. DS is a stereoisomer of CS, in which GlcA is epimerized to iduronic acid (IdoA). In mammalian tissues, CS and DS chains are often present as hybrid CS-DS chains on PGs (e.g. decorin, biglycan) (6). Tightly controlled modifications, such as epimerization and sulfation reactions, further increase the structural and functional diversity of the HS and CS/DS PGs (7, 8). GAG biosynthesis of both HSPGs and CS/DSPGs is initiated in the endoplasmic reticulum and Golgi apparatus by the formation of a common tetrasaccharide linkage region (glucuronic acid-galactose-galactose-xylose-*O*--) (GlcA-Gal-Gal-Xyl-*O*-) via *O*-glycosylation of a serine residue of the core protein (Fig. 1A) (1-3). The first step in the biosynthesis of this tetrasaccharide linkage region is catalyzed by xylosyltransferases (XylTs) encoded by the paralogues *XYLT1* [MIM 608124] and *XYLT2* [MIM 608125] (9). Two Gal residues are subsequently added by respectively galactosyltransferase I (GalT-I or β4GalT7 encoded by *B4GALT7* [MIM 604327]) and galactosyltransferase II (GalT-II or β3GalT6 encoded by *B3GALT6* [MIM 615291]) (10, 11). Further addition of a GlcA by glucuronosyltransferase I (GlcAT-I encoded by *B3GAT3)* [MIM 606374] completes the formation of the linkage region (12).

The cellular machinery required for GAG biosynthesis and modification is conserved over a broad range of eukaryotic organisms (13). The particular importance of a correct initiation of GAG synthesis is underscored by the identification of a series of severe, overlapping and multisystemic genetic human disorders that result from deficient activity of any of these "linkage-enzymes", collectively coined as 'linkeropathies' (14-16). The most frequently reported linkeropathy results from galactosyltransferase II deficiency, caused by biallelic pathogenic variants in *B3GALT6.* This leads to two clinically overlapping conditions, spondylodysplastic Ehlers-Danlos syndrome (EDS) (spEDS-*B3GALT6*) and

spondyloepimetaphyseal dysplasia with joint laxity type I (SEMD-JL1) (16, 17), both of which are

characterized by variable degrees of spondyloepimetaphyseal bone dysplasia, (with postnatal growth

restriction, bowing of long bones, hypo- or dysplasia of iliac and long bones, and vertebral body

changes), kyphoscoliosis, bone fragility with spontaneous fractures, joint hypermobility in conjunction

with joint contractures, hypotonia with delayed motor development, skin fragility, and craniofacial

dysmorphisms, including midfacial hypoplasia and dysplastic teeth (16-18). The severe and pleiotropic

phenotype suggests a critical role for *B3GALT6* in the development and homeostasis of various

connective tissues. Pathogenic variants in *B3GALT6* either result in mislocalization and/or reduced

amounts of active galactosyltransferase II (19). Previous studies on several biallelic *B3GALT6* variants

showed that they resulted in a barely detectable *in vitro* galactosyltransferase II enzymatic activity, but

intriguingly HS and CS/DS GAG synthesis was partially preserved *in cellulo* (20). While some variants

may have a greater deleterious effect on the *in vitro* than on the *in cellulo* enzymatic activity, it is

currently unknown whether there are rescue mechanisms at play to (partially) compensate for the

galactosyltransferase II activity.

In recent years, zebrafish have emerged as an excellent tool to model disease and to increase our

understanding of developmental processes and disease mechanisms. Zebrafish share 70% of their

genes with humans (21) and the advantages of their use are reflected by their large breed size, short

lifecycle and low husbandry costs (22). Zebrafish has proven to be a relevant genetic model for the

study of PG biosynthesis since many enzymes involved in this process are strongly conserved between

human and zebrafish (23), and a number of zebrafish models for defective PG synthesis (including

fam20b^{b1125}, xylt1^{b1189}, or b3gat3^{hi307}), provided novel knowledge on PG function in bone and cartilage

formation (23-26).

In this study, we generated *b3galt6* knock-out (KO) zebrafish as a model for spEDS-*B3GALT6* and SEMD-

JL1. These models were used to investigate the structure and amount of different GAG disaccharides

and the PG linkage region composition in these disorders. Furthermore, in-depth phenotypic

characterization of *b3galt6^{-/-}* zebrafish was done to reveal to which extent key features of the human

disorders are recapitulated and to further elucidate the function of galactosyltransferase II in

vertebrate musculoskeletal development.

Results

CRISPR/Cas9-mediated *b3galt6* knock-out in zebrafish leads to reduced HS and CS/DS glycosaminoglycan concentrations in bone, muscle and skin

Similar to its human orthologue, zebrafish *b3galt6* (NM_001045225.1) is a single-exon gene (Fig. 1B).

Human and zebrafish galactosyltransferase II (β3GalT6/B3galt6) protein sequence display 82.5 %

amino acid (AA) similarity, with the highest AA conservation in the luminal domain (Fig. 1C).

In order to investigate the consequences of zebrafish B3galt6-deficiency on GAG biosynthesis in

different tissues, we generated two *b3galt6* KO zebrafish models using CRISPR/Cas9 gene editing. The

first KO model carries the c.181delinsAAAGAGCTCCTAAAGGG indel mutation (cmg20 allele), leading

to a premature termination codon (PTC) (p.(Phe68*)), located upstream to the catalytic domain (Fig.

1D). RT-qPCR showed a 40.3% reduction of *b3galt6 mRNA* expression in homozygous adult *b3galt6* KO

zebrafish compared to wild-type (WT) siblings (Fig. 1E). For the remainder of this manuscript, this

model will be referred to as *b3galt6*$^{-/-}$. The second KO model carries a four base pair deletion

(c.398_401del) (cmg22 allele), predicted to generate a PTC (p.(Asn139*)) in the catalytic domain

(NM_001045225.1). This mutant was solely used for validation purposes (Fig. S1). Both *b3galt6* KO

zebrafish were produced at an expected Mendelian ratio and survived into adulthood. Unfortunately,

commercial antibodies, which were only available against human β3GalT6, did not work in our hands

and precluded the investigation of zebrafish B3galt6 protein levels and localization.

A

GAG chain polymerization | Tetrasaccharide linkage region | Core protein

Golgi | ER

N

Heparan sulfate

Chondroitin sulfate

Dermatan sulfate

B3GAT3
Glucuronosyl-
transferase I

B4GALT7
Galactosyl-
transferase I

B3GALT6
Galactosyl-
transferase II

XYLT1
XYLT2
Xylosyl-
transferase

spEDS-B4GALT7

SEMD-JL1 &
spEDS-B3GALT6

Larsen-like syndrome
B3GAT3 type

Spondylo-ocular
syndrome & Desbuquois
dysplasia type 2

C

★ Xylose (Xyl)   ◆ Glucuronic Acid (GlcA)   ■ N-Acetylgalactosamine (GalNAc)

● Galactose (Gal)   ■ N-Acetylglucosamine (GlcNAc)   ◆ Iduronic acid (IdoA)

● Amino acid   ● Serine residue

B

2.81 kb — Forward strand

H. sapiens
B3GALT6

1.06 kb — Forward strand

D. rerio
b3galt6

C

```
Q96L58  B3GT6_HUMAN    1   MKLLRRAWRRRAALGLGTLALCGAALLYLARCAAEPGDPRAMSGRSP-------PP---P    50
Q256Z7  Q256Z7_DANRE   1   MNLLRLVCRHKTALALGVLLLFAVVLLFLAKCTSETLKPAESPGAAPQAESHQKPPAGPA   60
                          *.****  .  *.::**.** * *  ...**.**.*..*   .*    *  .*    **

Q96L58  B3GT6_HUMAN   51   APARAAAFLAVLVASAPRAAERRSVIRSTWLARRGAPGDVWARFAVGTAGLGAEERRALE  110
Q256Z7  Q256Z7_DANRE  61   RPKELSAFLVILITTGPKYTERRSIIRSTWLTKHDPEVLY--WFVIGTEGLPAEDLQNLA  118
                          *  .  :***.:*:::.*: :****.*******::.       *.:** ** **: : *

Q96L58  B3GT6_HUMAN  111   REQARHGDLLLLPALRDAYENLTAKVLAMLAWLDEHVAFEFVLKADDDSFARLDALLAEL  170
Q256Z7  Q256Z7_DANRE 119   TEQIRHHDLLLLPDLRDSYENLTQKLLHMYSWLDQNVEFKFVLKADDDTFARLDLLKEEL  178
                          ** ** ****** ***.****** *:* *.:***::* *.********:*****  *  **

Q96L58  B3GT6_HUMAN  171   RAREPARRRRLYWGFFSGRGRVKPGGRWREAAWQLCDYYLPYALGGGYVLSADLVHYLRL  230
Q256Z7  Q256Z7_DANRE 179   KVKEPK--ERLYWGFFSGRGRVKTAGKWKESAWELCDYYLPYALGGGYVLSADLVRYIRL  236
                          :..:**   .*************. *.**:.:**:**.********************:**

Q96L58  B3GT6_HUMAN  231   SRDYLRAWHSEDVSLGAWLAPVDVQREHDPRFDTEYRSRGCSNQYLVTHKQSLEDMLEKH  290
Q256Z7  Q256Z7_DANRE 237   NVGFLKIWQSEDVSLGAWLAPVDVKRLHDPRFDTEYKSRGCSNKYLVTHKQSLEDMLEKH  296
                          .  .:*: :******************. *******:*****:*****************

Q96L58  B3GT6_HUMAN  291   ATLAREGRLCKREVQLRLSYVYDWSAPPSQCCQRREGIP  329
Q256Z7  Q256Z7_DANRE 297   QTLQRDGRLCKEEVKLRLSYIYDWSVPPSQCCQRKDGIP  335
                          .**  .*****:**:*****:****.******:.**
```

D

TM
(12-34)

Luminal
(35-335)

WT β3GalT6 | Galactosyltransferase

c.181delinsAAAGAGCTCCTAAAGGG
p.(Arg61Lys)

b3galt6⁻/⁻ | Galactosyltransferase

p.(Phe68*)

E

b3galt6 mRNA
expression level

Relative quantities (± SEM)

**

WT | b3galt6⁻/⁻

Fig. 1. Schematic representation of the GAG biosynthesis and the evolutionary conservation of the zebrafish and human *b3galt6/B3GALT6* gene and B3galt6/β3GalT6 protein. (**A**) Simplified schematic representation of the PG linkage and glycosaminoglycans biosynthesis. PG core proteins are synthesized in the endoplasmic reticulum and undergo further modification in the Golgi apparatus. First, a tetrasaccharide linkage region is synthesized that originates from the addition of a Xyl onto a serine residue of the core protein. Subsequent addition of two Gals and one GlcA residues completes this process. Depending on the addition of GlcNAc or GalNAc to the terminal GlcA residue of the linkage region, HS or CS/DS PGs will be formed, respectively. The elongation of the GAG chain continues by repeated addition of uronic acid and amino sugar residues. The GAG chain is then further modified by epimerization and sulfation (not shown). The main genes encoding for the enzymes involved in the linkage region biosynthesis are indicated in bold and biallelic mutations in these genes result in different linkeropathies. Disorders resulting from deficiency of the enzymes are indicated in black boxes. The deficient enzyme (β3GalT6) and the related disorder studied here are depicted in red (e.g. spEDS-*B3GALT6* and SEMD-JL1). (**B**) Schematic representation of the genomic structure of the *H. sapiens B3GALT6* (NM_080605.4) and *D. rerio b3galt6* (NM_001045225.1) genes. The white and black boxes represent the untranslated regions and the single *B3GALT6/b3galt6* exon, respectively. The zebrafish *b3galt6* gene resides on chromosome 11, spans a region of 1.06 kb and has not been duplicated during teleost evolution. (**C**) Amino acid (AA) sequence alignment (Clustal Omega) between *H. sapiens* β3GalT6 (Uniprot: Q96L58) and *D. rerio* B3galt6 (Uniprot: Q256Z7). The luminal galactosyltransferase domain is highlighted in grey. Conservation of AA sequences is shown below the alignment: "*" residues identical in all sequences in the alignment; ":" conserved substitutions; "." semi-conserved substitutions; "space" no conservation. (**D**) Schematic representation of the cmg20 allele in *b3galt6* mutants. *b3galt6* consists of three parts, a cytosolic domain (AA 1-11), a transmembrane domain (TM, AA 12-34) and a luminal domain (AA 35-335). The location of the CRISPR/Cas9-generated indel variant is indicated by a scissor symbol and the resulting premature termination codon (PTC) is depicted by a red star. (**E**) RT-qPCR analysis of relative *b3galt6* mRNA expression levels in *b3galt6*[−/−] adult zebrafish (0.597 ± 0.03919) (n=5) compared to WT siblings (1 ± 0.05395) (n=6). Data are expressed as mean ± SEM. A Mann-Whitney U test was used to determine significance, **P<0.01.

To examine the activity of B3galt6 in different *b3galt6*[−/−] tissues, concentrations of CS, DS and HS disaccharides extracted from bone, muscle and skin from WT and *b3galt6*[−/−] zebrafish were quantified using a combination of enzymatic digestion and anion-exchange HPLC (Figs. S2-S5 and Tables S1-S4). Overall, a generalized decrease in HS, CS and DS concentration was detected in all examined tissues of *b3galt6*[−/−] zebrafish. In the GAG fractions prepared from bone, the disaccharide concentration of the CS/DS and CS moieties was significantly decreased by respectively 55% and 64%, compared to WT zebrafish (Table 1 and Tables S1 and S2). DS moieties were undetectable in bone of *b3galt6*[−/−] zebrafish, whereas their concentration was 50.5 pmol/mg protein in bone of WT zebrafish (Table 1 and Table

S3). A non-significant decrease in the disaccharide concentration of the HS moiety was observed in *b3galt6*[-/-] mutant bone (Table 1 and Table S4), compared to WT. In GAG fractions prepared from *b3galt6*[-/-] muscle, CS/DS, CS and DS disaccharide moieties were generally lower than in WT muscle (Table 1 and Tables S1-S3). Due to the low concentrations close to the detection limit, a significant reduction could only be detected for the DS moiety (72% reduction) (Table 1 and Table S3). In addition, a significant decrease in HS disaccharide moieties by 56% was noted in *b3galt6*[-/-] muscle compared to WT muscle (Table 1 and Table S4). In the GAG fractions prepared from *b3galt6*[-/-] skin, the concentration of all disaccharide moieties was significantly reduced compared to WT (Table 1 and Tables S1-S4). The disaccharide concentration of the CS/DS, CS, DS and HS moieties were decreased by respectively 77%, 73%, 55% and 62% compared to WT (Table 1 and Tables S1-S4). No altered sulfation patterns were observed in disaccharide moieties when comparing unique GAG chains between WT and *b3galt6*[-/-] zebrafish samples (Tables S1-S4).

Despite the strong reduction in disaccharide concentrations observed in *b3galt6*[-/-] zebrafish, low amounts of GAGs were still produced in the mutants. Similar results were obtained for the second *b3galt6* KO zebrafish (cmg22 allele) (Table S5). RT-qPCR revealed no significant upregulation of other galactosyltransferase(s) member(s) from the *b3galt* family (*b3galt1a*, *b3galt2*, *b3galt4*, *b3galnt2*) or of other linkage-enzymes (*xylt1*, *xylt2*, *b4galt7* and *b3gat3*) (Fig. S6), suggesting that altered expression of these genes did not compensate for the reduced *b3galt6* mRNA levels.

Table 1. Concentrations of GAG disaccharides in wild-type and *b3galt6*[-/-] zebrafish tissues

	pmol/mg protein (Average ± SE)			
	CS/DS	CS	DS	HS
Bone				
Wild-type	774.2 ± 19.9	728.4 ± 148.1	50.5 ± 8.1	178.6 ± 19.8
b3galt6[-/-]	$346.1 \pm 20.0^*$	$279.9 \pm 4.8^{**}$	N.D.	137.1 ± 39.7^{ns}
Muscle				
Wild-type	9.3 ± 0.4	2.1 ± 0.4	14.5 ± 1.1	30.8 ± 1.2
b3galt6[-/-]	5.7 ± 3.1^{ns}	N.D.	$4.1 \pm 1.0^*$	$13.5 \pm 4.0^{**}$
Skin				
Wild-type	236.2 ± 26.5	175.1 ± 25.3	75.9 ± 7.9	55.1 ± 4.5
b3galt6[-/-]	$55.4 \pm 15.2^*$	$47.9 \pm 13.5^{**}$	$33.8 \pm 6.3^{**}$	$21.0 \pm 2.8^{**}$

Student's t-test (vs wild-type) (n=3), N.D.: not detected (<1 pmol/mg protein), *P<0.01, **P<0.05, ns: not significant, SE: standard error.

B3galt6-deficiency leads to the biosynthesis of a non-canonical (GlcA-Gal-Xyl-*O*-) trisaccharide linkage region in zebrafish

In order to explain why *b3galt6*[-/-] zebrafish can still produce GAG chains on PG core proteins, we investigated the composition of the linkage region. To this purpose, whole zebrafish extracts were enriched for all proteoglycans, their GAG chains depolymerized with bacterial lyases, the core proteins digested with trypsin and the remaining glycopeptides subjected to nano-scale liquid chromatography-tandem mass spectrometry (nLC-MS/MS) using a recently developed glycoproteomic approach (27, 28). Potential structural differences in PG linkage regions in WT versus *b3galt6*[-/-] zebrafish are thus elucidated through the combined identities of peptide backbones and GAG linkage region structures. The glycoproteomic analysis identified two CS-glycopeptides derived from biglycan (UniProt: A8BBH0) in both the WT and *b3galt6*[-/-] zebrafish. These CS-glycopeptides displayed different linkage region structures in WT compared with *b3galt6*[-/-] zebrafish (Fig. 2). While the peptide in the WT zebrafish was modified with a hexasaccharide structure (Fig. 2A), the peptide in the *b3galt6*[-/-] zebrafish was modified with a pentasaccharide structure (Fig. 2B), supporting the hypothesis that *b3galt6*[-/-] zebrafish produces a non-canonical trisaccharide linkage region. The precursor ion (m/z 1281.7922; 3+) in the WT zebrafish equated to a mass of a peptide (DQEEGSAVEPYKPEHPTCPFGCR) modified with a hexasaccharide structure, with the peptide containing two carbamidomethyl (CAM) modifications. Furthermore, the hexasaccharide structure was modified with one sulfate group (SO_3^-) on the subterminal GalNAc residue [$GlcAGalNAc-H_2O+SO_3+H$]$^+$, and one phosphate group (HPO_3^-) on the Xyl residue (Fig. S7). In contrast, the precursor ion (*m/z* 1227.7765; 3+) in the *b3galt6*[-/-] zebrafish equated to a mass of the same peptide but modified with a pentasaccharide structure, with two phosphate/sulfate groups, with the peptide containing two CAM modifications. The difference in mass between the WT and *b3galt6*[-/-] precursor ions corresponds to 162.08 Da, equating to the mass of a hexose (galactose) residue (162 Da). Additional HS- and CS- glycopeptides with the expected tetrasaccharide linkage region were found in the WT. In the *b3galt6*[-/-], no HS-modified structures were identified but additional CS-glycopeptides with the non-canonical trisaccharide linkage region were found. Table S6 shows a complete list of all glycopeptides identified in both WT and *b3galt6*[-/-] zebrafish. PGs harboring a tetrasaccharide linkage region were never detected in the *b3galt6*[-/-] adult zebrafish samples and no trisaccharide linkage region was detected in WT zebrafish samples.

These observations suggest that B3GAT3/GlcAT-I can transfer a GlcA residue not only to Gal-Gal-Xyl, but also to a Gal-Xyl substrate. This was confirmed by an *in vitro* GlcAT-I assay, which showed that GlcAT-I is indeed capable of adding a GlcA to a Gal-Xyl-*p*-nitrophynyl, which was chemically synthesized, thus to a substrate ending with a single galactose (Fig. S8).

Fig. 2. The identification of a trisaccharide linkage region in the *b3galt6*[-/-] zebrafish. (**A** and **B**) MS2 fragment mass spectra of CS-glycopeptides derived from biglycan (UniProt: A8BBH0) showed different linkage regions in WT compared to the *b3galt6*[-/-] zebrafish. (**A**) The peptide (DQEEGSAVEPYKPEHPTCPFGCR) in WT zebrafish was modified with a residual hexasaccharide structure, corresponding to the canonical tetrasaccharide linkage region (*m/z* 1281.79; 3+). The hexasaccharide structure was modified with one sulfate group (SO_3^-) at the subterminal GalNAc residue and one phosphate group (HPO_3^-) at the xylose residue (Fig. S7) (**B**) The peptide (DQEEGSAVEPYKPEHPTCPFGCR) in *b3galt6*[-/-] zebrafish was modified with a residual pentasaccharide structure, corresponding to a non-canonical trisaccharide linkage region (*m/z* 1227.78; 3+). The pentasaccharide structure was modified with two phosphate/sulfate groups. Their exact positioning and identity could not be determined due to low intensities of corresponding ion peaks, and the positions are therefore assigned in analogy of the WT structure.

b3galt6^{-/-} zebrafish show early-onset, progressive morphological abnormalities and delay of cartilage and bone formation, reminiscent of human spEDS-*B3GALT6*

One of the clinical hallmarks of human spEDS-*B3GALT6* is the pronounced skeletal involvement with the presence of spondyloepimetaphyseal dysplasia with postnatal growth restriction, bowing of long bones, hypoplasia of iliac bones, femoral head dysplasia, vertebral body changes, kyphoscoliosis and bone fragility with spontaneous fractures (Table 2). To evaluate the development of cartilage and bone associated with B3galt6-deficiency in zebrafish, phenotypic characterization of *b3galt6^{-/-}* zebrafish was performed in juvenile (20 dpf, days post fertilization) and adult stages (four months).

At 20 dpf, the average weight (W), standard length (SL) and Fulton's condition factor (*K*), which is used as a measure for the relative proportion of SL to W (29), were not significantly different between *b3galt6^{-/-}* and WT siblings (Fig. 3A), although for SL there was a trend towards smaller mutants (P=0.0640). Several externally visible morphological anomalies could be observed in 20 dpf *b3galt6^{-/-}* zebrafish as compared to their WT siblings, including a shorter, more compact head, a shorter and rounder frontal region, a shorter parietal/occipital region and a malformed posterior edge of the opercular apparatus (Fig. 3B). Alcian blue (AB) cartilage staining showed no morphological abnormalities, but developmental delay of the head and pectoral and caudal fin endoskeleton was noted (Fig. 3B, Table S7), with cartilage elements being absent or in early development in *b3galt6* KO, while these structures were already fully developed in WT sibling zebrafish. Alizarin red (AR) mineral staining showed delayed mineralization in all regions of the head, the pectoral fin girdle and the caudal fin (Fig. 3B, Table S8), further confirming developmental retardation. In addition, malformations of different mineralized bones, such as the parasphenoid, branchiostegal rays, supracleithrum and the subopercle bone could be detected in the head of *b3galt6^{-/-}* zebrafish larvae (Fig. S9). The vertebral column, i.e. the vertebral centra and their associated elements (neural and haemal arches), did not show delayed mineralization in *b3galt6^{-/-}* zebrafish larvae, but malformation of the associated elements and fusions of preural vertebrae were observed at a much higher frequency in the *b3galt6^{-/-}* mutants. Kyphosis, scoliosis and lordosis were also more frequently observed and pronounced in *b3galt6^{-/-}* larvae compared to WT siblings (Fig. S10 & Table S8). A similar phenotype was observed in the second *b3galt6* KO model (cmg22 allele), confirming the specificity of the displayed phenotype (Fig. S11).

Fig. 3. Developmental delay and morphological abnormalities in 20 dpf *b3galt6⁻/⁻* zebrafish. (**A**) Weight, standard length and Fulton's condition factor of WT and *b3galt6⁻/⁻* zebrafish larvae at 20 dpf (n=10). (**B**) External phenotype of the head of juvenile WT and *b3galt6⁻/⁻* sibling zebrafish. All images are oriented anterior to the left, posterior to the right, dorsal to the top and ventral to the bottom. (**1-1'**) *b3galt6⁻/⁻* zebrafish display a rounder frontal region (black line), shorter parietal/occipital region (white horizontal line) and a malformed posterior edge of the opercular apparatus (dotted black line), compared to WT control. (**2-2'**) Representative images of cartilage AB staining of the head. The olfactory region (indicated by 'o' in the WT and arrowhead in the *b3galt6⁻/⁻* picture) and the otic region (indicated by white asterisk) show delayed development of cartilage elements in *b3galt6⁻/⁻* zebrafish as compared to WT. (**3-3'**) Representative images of AR mineral staining of the head. The olfactory region (indicated by 'o') and the otic region (indicated by asterisk) show severely delayed mineralization in *b3galt6⁻/⁻* zebrafish larvae as compared to WT. (**4-4'**) Representative images of AB staining of the caudal fin. The modified associated elements show developmental delay in *b3galt6⁻/⁻* zebrafish as compared to WT. Abbreviations: E, epural; HS PU2, haemal spine of preural 2; HS PU3, haemal spine of preural 3; Hy1-5, hypural 1-5; NS PU2, neural spine preural 2; opc, opistural cartilage; Ph, parhypural (**5-5'**) Representative images of AR mineral staining of the caudal fin. Caudal vertebrae and their associated elements show no delay in ossification. Malformation or absence of caudal fin associated elements (block arrow indicates a neural arch; arrowhead indicates a missing haemal arch), and malformation of uroneural (uro) and vestigial neural arches and spines (vna/s) are observed in *b3galt6⁻/⁻* zebrafish; hypurals have a bent appearance. Scale bars: images of the head 500 µm, images of the caudal fin 200 µm.

At the adult stage (four months), *b3galt6⁻/⁻* zebrafish had a significantly reduced average SL (1.834 ±

0.077 cm), and W (0.1196 ± 0.014 g), when compared to their WT siblings (2.371 ± 0.049 cm, and

0.2497 ± 0.016 g, respectively), while the Fulton's condition factor (*K*) indicated that SL and W were

normally proportioned to each other (Fig. 4A). A similar phenotype was observed in the second *b3galt6*

KO model (cmg22 allele), confirming the specificity of the displayed phenotype (Fig. S1).

The head malformations, which were already observed in 20 dpf *b3galt6⁻/⁻* zebrafish, showed further

progression, with a more pronounced smaller, rounded head and an abnormal compaction of the

opercular apparatus (Fig. 4B). In addition, a reduced surface area of the fins, with a crenelated postero-

distal fin edge was observed in the mutant animals. The pectoral fins were pointing in an abnormal

upwards angle (Fig. 4B). Kyphoscoliosis was variably present and mainly occurred in the posterior part

of the body (not shown). In addition, an altered pigmentation pattern was observed, mainly on the

ventral part of the body where the dark stripes showed interruptions (Fig. 4B).

Fig. 4. Characterization of the external phenotype of adult *b3galt6*$^{-/-}$ zebrafish. (**A**) *b3galt6*$^{-/-}$ zebrafish (n=16) had a significantly shorter standard length (SL) compared to their WT siblings (n=17). The weight (W) was significantly lower in the mutant zebrafish. The Fulton's condition factor (K= 100*(W/SL^3)) was not significantly different between *b3galt6*$^{-/-}$ zebrafish and WT siblings. Data are expressed as box plots with min-to-max whiskers on which each individual data point is plotted. An unpaired t-test with Welch's correction was used to determine significance. ****$P < 0.0001$, ns: not significant. (**B**) Adult *b3galt6*$^{-/-}$ zebrafish exhibit morphological abnormalities compared to their WT siblings; *b3galt6*$^{-/-}$ zebrafish display interruptions of the blue horizontal stripes (white arrows), smaller fins (arrowhead), aberrations of the head shape (black line), of the shape of the opercular apparatus (white line indicates the posterior edge) and of the pectoral fin (black line and arrow indicate the fin rays at the base of the fin, white line shows the outline of the fin).

Further progression of the larval phenotype was also confirmed by AR staining of the mineralized skeleton in adult (four months) *b3galt6*$^{-/-}$ zebrafish. The olfactory region was underdeveloped (Fig. 5A), the cranial roof bones and sutures were deformed (Fig. S12) and the opercular apparatus was smaller and compacted in KO zebrafish (Fig. 5A and Fig. S12). The fifth ceratobranchial arch and its attached teeth were smaller in *b3galt6*$^{-/-}$ zebrafish (Fig. 5B). In the pectoral fins of *b3galt6*$^{-/-}$ zebrafish the radials were smaller, had an irregular shape, and were often fused (Fig. 5B). The autocentra, i.e. the inner layer of the vertebral centra in the vertebral column and caudal fin (30) of KO zebrafish had a normal hourglass shape, but the associated elements often showed presence of extra intramembranous bone (Fig. 5 and Fig. S12) and extra bony elements (Fig. 5A). Fusions were observed in preural vertebrae (Fig. 5A). Lordosis, kyphosis and scoliosis was variably observed along the vertebral column of *b3galt6*$^{-/-}$ mutant zebrafish (not shown). Extra mineralization was observed at the edge (growth zone) of *b3galt6*$^{-/-}$ scales (Fig. 5B). Notably, variation in the severity of the skeletal malformations in *b3galt6*$^{-/-}$ zebrafish was observed.

Fig. 5. Alizarin red stained mineralized bone in WT and *b3galt6*^{-/-} adult zebrafish. The mineralized bones in the skeleton were studied in four-month-old WT and mutant zebrafish after AR mineral staining. (**A**) *b3galt6*^{-/-} zebrafish display malformation of the cranium, vertebral column and caudal fin. (**A1-1'**) The cranium shows an underdeveloped olfactory region (black line) and a smaller compacted opercular bone (asterisk). (**A2-2'**) The vertebral column shows extra intramembranous outgrowths located where the distal tip of the neural and haemal arches fuse to neural and haemal spine respectively (arrowheads). (**A3-3'**) In the caudal fin fusion of preural vertebrae is observed (asterisk). Associated elements show intramembranous outgrowths (arrowhead) and the presence of extra bony elements (diamond). The distal tip of the hypurals is sometimes split (transparent arrowhead). Structures indicated in the WT caudal fin: Hy1-5, hypural 1-5, Ph, parhypural. (**B**) Details of mineralized structures in *b3galt6*^{-/-} zebrafish. (**B1-1'**) The fifth ceratobranchial arch and its teeth are smaller in KO zebrafish. (**B2-2'**) The proximal (PR) and distal radials (DR) have irregular shapes in mutant zebrafish. Proximal (asterisk) and distal radials are sometimes fused. The bracket indicates a fusion between a proximal and distal

radial, and the first fin ray (FR). (**B3-3'**) Ribs also show extra intramembranous bone (black arrowhead). (**B4-4'**)

More mineralization was observed at the growth zone (bracket) of the scales in mutant zebrafish (arrowhead).

Scale bars = 500 µm.

Subsequent quantitative µCT analysis of four-month-old zebrafish revealed multifaceted effects of

B3galt6-deficiency on vertebral morphology and mineralization. Most notably, the volume of vertebral

centrum, haemal and neural associated elements was significantly reduced throughout the entire

vertebral column in *b3galt6*$^{-/-}$ zebrafish (P<0.01), with more pronounced differences in the abdominal

region (Fig. 6). Tissue mineral density (TMD) was not significantly different in the *b3galt6*$^{-/-}$ vertebral

bodies compared to WT siblings. However, TMD in zebrafish is proportionally related to standard

length (31, 32), and *b3galt6*$^{-/-}$ zebrafish exhibited a significantly reduced body size compared to the

WT siblings. We therefore normalized the µCT data to standard length using the normalization

procedure outlined by Hur *et al.* (33). In this procedure, phenotypic data in WT sibling controls was

scaled using the mean standard length of *b3galt6*$^{-/-}$ zebrafish as the reference length. Remarkably,

after normalization, a significant increase in TMD of centrum, haemal and neural associated elements

was noted in *b3galt6*$^{-/-}$ zebrafish compared to WT controls (P<0.001, P<0.01 and P<0.01, respectively)

(Fig. S13).

Fig. 6. µCT analysis of adult WT and *b3galt6*$^{-/-}$ zebrafish. Quantitative µCT analysis of the vertebral column in

five mutant versus five WT zebrafish at the age of four months. The bone volume, tissue mineral density (TMD)

and bone thickness were calculated for three parts of each vertebra: the vertebral centrum, the neural associated

element (neural arch and neural spine) and the haemal associated element (haemal arch and haemal spine). Plots associated with a significant difference are colored in a lighter coloring scheme. The X-axis represents each vertebra along the anterior-posterior axis. The volume is significantly reduced for each part of the vertebra in *b3galt6*$^{-/-}$ zebrafish (P<0.01, P<0.001 and P<0.000001, respectively). The thickness of the haemal (P<0.01) and neural (P=0.001) associated element is also significantly reduced in the mutants whereas the centrum thickness is not significantly reduced in *b3galt6*$^{-/-}$ zebrafish (P=0.12). TMD is not significantly different between *b3galt6*$^{-/-}$ zebrafish and WT siblings when not normalized for standard length (P>0.2). Data are presented as mean ± SEM. TMD, tissue mineral density; HA, hydroxyapatite.

b3galt6 knock-out zebrafish display a disturbed organization of type I collagen fibrils in skin, bone and intervertebral space

Transmission electron microscopy (TEM) studies on the dermis of spEDS-*B3GALT6* patients revealed an abnormal collagen fibril architecture with loosely packed collagen fibrils of variable size and shape (5). We therefore studied the effect of loss of galactosyltransferase II activity on collagen fibril architecture. TEM sections of skin biopsies of *b3galt6*$^{-/-}$ zebrafish showed an abnormal dermal collagen fibril architecture characterized by loosely packed collagen fibrils and larger and more interfibrillar spaces compared to WT siblings (Fig. 7A). In addition, at the level of the scales a thicker epidermal layer was observed and electron dense structures were present in the interfibrillar spaces (Fig. 7A). Next, we examined the intervertebral region and adjacent vertebral centra in the vertebral column of five-month-old zebrafish by means of TEM. In general, the type I collagen fibers in *b3galt6*$^{-/-}$ zebrafish displayed a lower degree of organization (Fig. 7B). This was observed in immature collagen layers, deposited by osteoblasts in the outer edges of the intervertebral space, but also in the mature type I collagen layers of the intervertebral ligament and in the autocentrum, the inner layer of the vertebral bone, where a lack of the typical plywood-like organization was seen. Detailed investigation of TEM sections of the vertebral bone revealed presence of several electron dense spots of unknown origin in knock-outs. These 'dark spots' may reflect extrafibrillar hydroxyapatite crystals (Fig. 7B).

Mass spectrometry analysis of type I collagen modification in bone derived from the vertebral column showed no significant difference in hydroxylation status of the α1(I) K87 and C-telopeptide lysine residues (Table S9). These lysine residues are conserved in zebrafish and human type I collagen and are crucial for intermolecular crosslinking of collagen fibrils in bone matrix (34).

Fig. 7. Collagen fibrillar architecture in five-month-old adult zebrafish. (**A**) Skin from five-month-old WT and

b3galt6-/- adult zebrafish. The epidermis (Ep), the outer layer of the skin covering the scales (S) of zebrafish,

appears thicker in *b3galt6*-/- zebrafish compared to the epidermis of WT zebrafish. A mature zebrafish scale

consists of several layers. The limiting layer of the scale (LL) is rich in fine granules that in some areas are linearly

arranged, forming dense layers. This matrix contrasts with that of the external layer(s) (EL) below. Structures

resembling extrafibrillar crystals (arrowhead) are noted in the external layer of the mutant scale, which are rare

in WT zebrafish sections. Upon investigation of the dermis of *b3galt6*-/- mutants, increased interfibrillar spaces

are noted (arrows), which are absent in the dermis of WT adults. Scale bars = 500 nm. (**B**) Schematic

representation and TEM images of the zebrafish intervertebral space. Schematic representation of the zebrafish

intervertebral space. Abbreviations: AC, autocentrum; B, bone; Cb, chordoblasts; Cc, chordocytes; ColII, type II

collagen; E, elastin; iMC, immature collagen type I layer; MC, mature collagen type I layer. Starting from the

central body axis, several recognizable structures are present in the intervertebral space along a proximo-distal

axis: (i) notochord tissue consisting of chordocytes and chordoblasts, (ii) the notochord sheath consisting of a

type II collagen layer and an external elastic membrane, (iii) and a connective tissue ligament consistent of a

mature and an immature type I collagen layer. The mature type I collagen layer is organized in a typical plywood-

like pattern and is continuous with the inner layer of the vertebral bone, i.e. the autocentrum, which also shows

a plywood-like organization. The immature type I collagen layers consist of loose collagen fibers connecting the

outer layer of vertebral bones. Notice the higher number of nuclei (N) in the *b3galt6*-/- zebrafish at the

intervertebral space. Mature collagen (MC) displays a typical plywood-like organization, in WT zebrafish but not

in comparable regions for *b3galt6*-/- mutant zebrafish. Cross section and longitudinal type I collagen regions are

less clearly demarcated and smaller in immature collagen (iMC) from *b3galt6*-/- zebrafish compared to WT

siblings. It is more difficult to pinpoint the different layers of the autocentrum (AC) in the KO compared to WT

409 zebrafish. Bone (B) from the *b3galt6*$^{-/-}$ zebrafish shows electron dense structures (arrowhead). Scale bars = 10

410 µm for the intervertebral space and 1 µm for the other images.

b3galt6 knock-out zebrafish display ultrastructural muscle abnormalities and lower endurance

Human spEDS-*B3GALT6* patients often suffer from muscle hypotonia with a delay in motor

development. In view of the observed reduction in the concentration of the different GAG moieties in

muscle of *b3galt6*$^{-/-}$ zebrafish, we investigated the function and the ultrastructural appearance of

muscle.

First, the critical swimming speed (U_{crit}) and endurance of the zebrafish were assessed by means of a

custom-made swim tunnel. The U_{crit}, defined as the maximum velocity that the tested zebrafish could

sustain for at least 5 minutes (35), was significantly lower in *b3galt6*$^{-/-}$ zebrafish (30.1 ± 2.39 cm/s)

compared to length-matched WT (41.61 ± 1.942cm/s) (P<0.01) (Fig. 8A). The stamina of each zebrafish

was determined by an effort test where it was tested how long zebrafish could swim at high speed (37

cm/s) after a three-minute warm-up. This effort test showed that the *b3galt6*$^{-/-}$ zebrafish performed

less well compared to length-matched WT zebrafish, indicating a lower endurance (Fig. 8B).

The muscle tissue in the zebrafish trunk was further investigated at the ultrastructural level. Toluidine

blue-stained semi-thin cross sections revealed that the endomysium, a layer of connective tissue

ensheating the muscle fibers, was enlarged along the horizontal myoseptum of *b3galt6*$^{-/-}$ zebrafish,

although no clear differences in muscle fiber diameter were noted (Fig. 8C). Ultra-thin parasaggital

sections, investigated by TEM, showed that the length of the sarcomeres, the smallest basic units of

striated muscle, as well as the length of the A and I band were significantly increased in the *b3galt6*$^{-/-}$

zebrafish (Fig. 8D).

Fig. 8. Functional and structural analysis of adult WT and *b3galt6*^{-/-} zebrafish muscle. (A) The critical swimming speed and **(B)** the endurance performance was determined for *b3galt6*^{-/-} and length-matched WT zebrafish. **(C)** Semi-thin cross sections, stained with toluidine blue, demonstrate the presence of a thicker endomysium (arrowhead), located between the muscle fibers (Mf) at the horizontal myoseptum of *b3galt6*^{-/-} zebrafish at the age of five months. **(D)** Transmission electron microscope structure of a sarcomere with indications of the Z and M line and A, I and H band. The sarcomere length was determined at 5 different loci per semi-thin section, for 6 semi-thin originating coming from 3 different zebrafish per genotype (N=30). A significant increase in sarcomere length for *b3galt6*^{-/-} is noted. Furthermore, both the A and I band show an increase in length for the mutant zebrafish. Data are expressed as box plots with min-to-max whiskers on which each individual data point is plotted. A Mann-Whitney U test was used to determine significance. ****$P<0.0001$, **$P<0.01$.

Discussion

In this study we developed two viable *b3galt6* KO zebrafish models using CRISPR/Cas9 mutagenesis, and performed an in-depth characterization of the biochemical, ultrastructural and phenotypical consequences of β3GalT6-deficiency on different tissues, including skin, cartilage, mineralized bone and muscle. These models currently represent the only vertebrate models available for the study of the pathogenetic consequences of β3Galt6-deficiency on tissue development and homeostasis. In both models we detected a significant reduction, yet not a complete absence in *b3galt6* mRNA expression (Fig. 1E). One possible explanation is the absence of canonical nonsense-mediated decay (NMD) in the single-exon *b3galt6* gene, due to a lack of exon junction complexes (EJCs) [36]. We were unable to examine the effect of the mutations on B3galt6 protein levels and localization due to the

unavailability of a suitable antibody. Nevertheless, since the cmg20 and cmg22 alleles are predicted to

introduce a PTC respectively upstream and in the catalytic domain, residual translation of some mRNA

into a truncated protein is not expected to lead to the production of an active enzyme. β3galt6 is a key

enzyme in the biosynthesis of the common linkage region of PGs, and reduced levels of active enzyme

are thus expected to equally affect the biosynthesis of HS and CS/DS GAG chains. We therefore

quantified the disaccharide concentrations in bone, muscle and skin of adult *b3galt6*[-/-] zebrafish by

means of a combination of enzymatic digestion and anion-exchange HPLC. This revealed a distinct, and

relatively proportional decrease in the concentration of CS, DS, hybrid CS/DS and HS moieties in all

examined tissues, compared to age-matched WT zebrafish (Table 1). Nonetheless, low amounts of

GAGs were still produced in the KO zebrafish in absence of upregulation of other galactosyltransferase

member(s) from the *b3galt* family or by other linkage-enzymes (Fig. S6). Together with the previous

demonstration that the other five β3GalT human enzymes (e.g. β3GalT1, β3GalT2,

β3GalT3/β3GalNAcT1, β3GalT4 and β3GalT5) are involved in the biosynthesis of glycoproteins and

glycolipids (11), compensation for galactosyltransferase II deficiency in zebrafish by other *b3galt* family

members is unlikely. Chinese hamster ovary *B3galt6* KO cells were also shown to be capable of

producing GAGs, in contrast to *Xylt2*, *B4galt7* or *B3gat3* KO cells (37). The recent identification of the

presence of a non-canonical trisaccharide linkage region, lacking one Gal residue (GlcA-Gal-Xyl-*O*-), as

a minor constituent in the PG bikunin in the urine of healthy human individuals (28), prompted us to

examine the composition of the GAG linkage region in our *b3galt6* KO zebrafish models, by analyzing

protein extracts from adult *b3galt6*[-/-] and WT zebrafish via LC-MS/MS. This led to the striking finding

that the few detected PGs in the *b3galt6*[-/-] zebrafish all presented with a trisaccharide PG linkage

region, consisting of only one Gal (Fig. 2B) (Table S6). Further *in vitro* analysis indicated that GlcAT-

I/B3GAT3 is capable of adding a GlcA to Gal-Xyl-*O*- (Fig. S8). This observation might explain the

phenotypic differences observed between GalT-II/B3GALT6 deficiency and deficiencies of the other

linkage region enzymes, for which such a rescue mechanism might be absent. This hypothesis is

supported by the lack of GAGs in the Chinese hamster ovary *Xylt2*, *B4galt7* or *B3gat3* KO cells and by

the lethal phenotype observed in the *b4galt7* KO zebrafish model (38). At present, we have no

information on the length of the GAG chains attached to the trisaccharide linkage region or the

composition and the possible modifications of these HS and CS/DS chains. As such, the GAG structure,

length and modifications in the *b3galt6*[-/-] zebrafish might be different from the ones in WT zebrafish.

HSPGs and CS/DSPGs are essential macromolecules for the development, signaling and homeostasis

of many tissues. It therefore comes as no surprise that defects in linkage region biosynthesis can affect

many organs. Endochondral ossification of cartilage (forming chondral bones) for example is a multi-

staged process that must be coordinated in space and time. CS/DSPGs and HSPGS, which have key

roles in modulating signaling events during vertebrate skeletal development, are expressed in a spatially and temporally highly controlled manner (24), thus regulating the timing of endochondral ossification. Genetic defects disrupting PG synthesis or structure have been shown to disrupt the timing of skeletal development. Two zebrafish models with abnormal PG linkage region biosynthesis, i.e. the *fam20b* and *xylt1*-deficient models, revealed partial decrease in CSPGs, which are predominantly present in cartilage, and accelerated endochondral ossification, showing that CSPGs can negatively regulate skeletogenic timing (24). On the other hand, a strong reduction in CSPGs and partial decrease in HSPGs was noted in *b3gat3* mutant zebrafish, but this led to a delay in endochondral ossification and short and disorganized cartilaginous structures (39). Our B3galt6-deficient zebrafish models also show a delay in skeletal development at juvenile stages. At 20 dpf, a delay was noticed in the development of cartilage and mineralized bony elements of the craniofacial structures, the pectoral and caudal fin endoskeleton (Fig. 3B) (Tables S7 and S8), and these abnormalities had progressed in adult zebrafish. The reduced body length, misshapen cranial bones and smaller teeth observed in the head skeleton, and general skeletal dysplasia with kyphosis and scoliosis seen in the vertebral column of *b3galt6*[-/-] adult zebrafish is reminiscent of the skeletal phenotype observed in human β3GalT6-deficient individuals (Table 2).

In addition to skeletal dysplasia, we also observed abnormalities in bone mineralization. Adult mutant zebrafish stained with AR showed the presence of extra intramembranous bone and extra bony elements on the associated elements of the vertebral centra. On the other hand, quantitative μCT analysis of the vertebral column and associated elements revealed a generalized reduction in bone volume and thickness, and a relative increase in TMD after correction for reduced body length of the mutant zebrafish (Fig. 6 and Fig. S13). Although it is not clear so far if the increased fracture risk in spEDS-*B3GALT6* patients can be attributed to a lower bone volume combined with a higher TMD, such hypothesis has been put forward for the brittle bone disorder Osteogenesis Imperfecta (OI) based on observations in both patients and animal models (40, 41). In the Chihuahua (*col1a1a*[chi/+]) OI zebrafish model the intrafibrillar mineral platelets were shown to be smaller and more densely packed, leading to an increased mineral to matrix ratio, and consequently a reduced elastic modulus-to-hardness ratio, a measure for fracture toughness (41-43). In addition, in both OI patients and the mouse model *oim,* reduced collagen matrix organization was shown to lead to the formation of extrafibrillar crystals, which are larger compared to intrafibrillar crystals (43). Eventually, additional minerals stabilize the tissue, but they do so at the cost of increased brittleness of the bone (44). In accordance, in *b3galt6*[-/-] zebrafish we observed an irregular collagen organization interspersed with large interfibrillar spaces and electron dense spots on ultra-thin sections of skin and bone via TEM. The 'dark spots' observed in

the vertebral bone could be remnants of extrafibrillar mineral aggregates, contributing to increased

mineralization but also increased brittleness of the skeleton.

Abnormalities in collagen fibril organization were observed in many mutant zebrafish tissues. Besides

vertebral bone, irregular collagen organization was also observed in the intervertebral space,

intervertebral ligament, and the dermis. (Fig. 7). This strongly resembles the previously reported

abnormal dermal collagen fibril architecture in the skin from a β3GalT6-deficient individual (5). This

could be attributed to abnormalities in GAG chains of small leucine-rich PGs (SLRPs), such as the DSPGs

decorin and biglycan, which are known to be of particular importance for regulating collagen fibril

organization in connective tissues through their core protein and GAG chains (45). SLRPs are also

thought to influence collagen cross-linking patterns, but the mechanism by which they do so remain

largely unknown (45). Although similar hydroxylation levels were observed at K87 and the C-

telopeptide of the type I collagen α1 chain in $b3galt6^{-/-}$ and WT adult zebrafish (Table S9), we cannot

rule out that type I collagen cross-linking might still be affected.

In view of the observation that human β3GalT6-deficient individuals often present quite severe muscle

hypotonia with delayed gross motor development and reduced endurance, we also studied the effect

of B3galt6-deficiency on GAG synthesis in muscle, and on muscle function and ultrastructure (Fig. 8).

Similar to the other examined tissues, HS and CS/DS GAG concentrations were severely reduced in

muscle tissue. It is known that the ECM in skeletal muscle plays an important role in muscle fiber force

transmission, maintenance and repair. Numerous PGs, including the HSPGs syndecans 1-4, glypican-1,

perlecan and agrin, and CS/DSPGs, such as decorin and biglycan, are present in muscle ECM. Besides

their structural role, many of them are known to bind growth factors (e.g. TGFβ). Those growth factors

can be stored and released by the negatively charged GAGs. Certain enzymes in the ECM can cleave

GAG chains, thereby releasing associated growth factors, allowing their interaction in cell signaling and

mechanotransduction (46). The significant decrease in critical swimming speed (U_{crit}) and endurance

observed in the $b3galt6^{-/-}$ zebrafish suggest decreased muscle function (Fig. 8A-B). Further evidence

for an intrinsic muscle dysfunction in the $b3galt6^{-/-}$ model comes from semi-thin muscle sections,

showing an enlargement of the connective tissue sheet surrounding the muscle fibers (Fig. 8C).

Nevertheless, we cannot exclude that the observed skeletal malformations and/or other

abnormalities, such as respiratory or neuronal dysfunction, could also influence the outcome of these

tests.

Table 2. Clinical features and ultrastructural and biochemical abnormalities observed in spEDS-

***B3GALT6* and SEMD-JL1 individuals compared to the $b3galt6^{-/-}$ zebrafish model.** n.a.: not available.

GAG: glycosaminoglycan. The columns 'spEDS-*B3GALT6*' and 'SEMD-JL1' summarize the most important clinical features of all hitherto reported patients for whom this data was available (20, 47).

	Human		Zebrafish
	spEDS-*B3GALT6*	SEMD-JL1	*b3galt6*$^{-/-}$ mutant
General morphology			
Short stature	22/30	11/14	Reduced body length
Craniofacial dysmorphisms (e.g. midfacial hypoplasia, micrognatia)	25/30	12/14	Deformed, small and round head with shorter parietal/occipital region and underdeveloped olfactory region
Dental abnormalities	14/30	n.a.	Smaller teeth
Musculoskeletal involvement			
Bowing of limbs, metaphyseal widening or vertebral body changes	20/30	14/14	Deformities of the fin skeleton
Bone fragility with spontaneous fractures	19/30	0/3	Extra intramembranous bone, lower degree of collagen type I fibril organization and higher TMD after correction for smaller size
Joint contractures (especially hands)	23/30	5/14	Pectoral fins point upwards
Joint hypermobility	26/30	6/14	n.a.
Kyphoscoliosis	23/30	13/14	Kyphosis and scoliosis
Muscle hypotonia	18/30	1/14	Reduced critical swimming speed and endurance
Skin features			
Skin hyperextensibility	19/30	2/14	n.a.
Biochemical findings			
Reduced GAG synthesis	9/9	4/4 (3/4; < HS but > CS/DS)	Reduced GAG production
Ultrastructural findings			
Loosely packed collagen fibrils	1/1	n.a.	Disturbed collagen fibril organization in bone and loosely packed collagen fibrils in the dermis

Taken together, we generated and characterized the first *b3galt6* KO zebrafish models and show that their phenotype largely mimics that of human β3Galt6-deficiency. A strong reduction in HS, CS and DS GAG concentration was observed in *b3galt6*$^{-/-}$ zebrafish. Yet, we did observe a small amount of GAGs being produced with a unique trisaccharide PG linkage region. Furthermore, we showed that a lack of B3galt6 causes disturbances in collagen fibril organization in skin, ligament and bone, with the latter

558 also showing abnormalities in mineralization, and leads to structural and functional abnormalities of
559 the muscle. These KO zebrafish can serve as useful models to further unravel the underlying
560 pathogenic mechanisms of this complex multisystemic disorder and serve as a tool for the
561 development and evaluation of possible therapeutic interventions, such as the stimulation of intrinsic
562 compensational mechanisms for the production of PGs.

Material and Methods

Zebrafish Maintenance

567 Wild-type (AB strain) and *b3galt6* knock-out zebrafish lines were reared and maintained by standard
568 protocols (48). All animal studies were performed in agreement with EU Directive 2010/63/EU for
569 animals, permit number: ECD 16/18. All zebrafish used in a single experiment were bred at the same
570 density. All efforts were made to minimize pain, distress and discomfort.

Alcian blue and alizarin red staining

572 Twenty-day-old larvae (n=10) and five-month-old WT and *b3galt6*$^{-/-}$ (n=5) adult zebrafish were
573 stained as previously described (38). See Supplementary Information for details.

μCT scanning analysis

Whole-body μCT scans of four-month-old WT (n=5) and *b3galt6*$^{-/-}$ (n=5) siblings were acquired on a
SkyScan 1275 (Bruker, Kontich, Belgium) using the following scan parameters: 0.25 mm aluminum
filter, 50 kV, 160 μA, 65 ms integration time, 0.5° rotation step, 721 projections/360°, 5 m 57 s scan
duration and 21 μm voxel size. DICOM files of individual zebrafish were segmented in MATLAB using
custom FishCuT software and data were analyzed and corrected for standard length in the R statistical
environment, as previously described (33).

Histology and transmission electron microscopy

582 Zebrafish bone, consisting of the last transitional vertebra and two consecutive caudal vertebrae (49,
583 50), skin, at the lateral line below the dorsal fin, and muscle samples, located dorsolateral underneath
584 the dorsal fin, dissected from five-month-old WT (n=5) and *b3galt6*$^{-/-}$ (n=5) zebrafish, were fixed and
585 embedded in epon following the procedures outlined by Huysseune and Sire (51).

Swim tunnel experiments

Ten WT (five months) and ten *b3galt6*[-/-] zebrafish (eight months), were selected based on their similar

(not significantly different) length and weight. The critical swimming speed (U_{crit}) was measured in a

swim tunnel for both WT and *b3galt6*[-/-] zebrafish as previously described (52). Furthermore, the

endurance of each zebrafish was tested by means of an effort test in the swim tunnel. Each zebrafish

had a three minute warm-up and after the third minute, the speed increased until 37 cm/s for

maximum 7 minutes. When the zebrafish was not able to swim anymore, the test was stopped.

Collagen crosslinking analysis

The collagen crosslinking was analyzed in the dissected vertebral column (excluding Weberian

apparatus and urostyle) from five-month-old wild-type (n=5) and *b3galt6*[-/-] (n=5) siblings as described

before in Gistelinck *et al.* (40).

Disaccharide composition analysis of CS, DS, and HS chains

The level of total disaccharides from CS, DS, and HS in bone, muscle and skin tissues from nine-month-

old zebrafish (n=5) were determined as described previously (53, 54).

Glycopeptide preparation and analysis

LC-MS/MS sample preparation - CS- and HS-glycopeptides were purified from six-month-old WT (n=3)

and *b3galt6*[-/-] (n=3) zebrafish using a combination of extraction, trypsin digestion and anion exchange

chromatography, slightly modified (see Supplementary Information) from previously described

protocols (27, 55).

LC-MS/MS analysis - The samples were analyzed on an Orbitrap Fusion mass spectrometer coupled to

an Easy-nLC 1200 liquid chromatography system (Thermo Fisher Scientific., Waltham, MA) (see

Supplementary Information). Nanospray Flex ion source was operated in positive ionization mode and

three separate higher-energy collision-induced dissociation (HCD) MS/MS spectra were recorded for

each precursor ion (27, 55).

LC-MS/MS data analysis - The glycopeptide files were analyzed using both manual interpretation and

automated Mascot searches for GAG-glycopeptides. The files were manually interpreted using the

Xcalibur software (Thermo Fisher Scientific). Database searches were performed against *Danio rerio* in

the UniProtKB/Swiss-Prot database and NCBI using Mascot Distiller (version 2.6.1.0, Matrix Science,

London, U.K) and an in-house Mascot server (version 2.5.1). Glycan structures with very small mass

differences were manually evaluated at the MS2-level.

GlcAT-I assay

GlcA-transferase assay was carried out by the methodology as previously described (12, 56, 57), with
slight modifications.

Detailed information regarding the procedures used is provided in SI Appendix, Supplementary
Material and Methods.

Competing interests

623　The authors declare no competing interests.

Acknowledgements

We would like to thank the TEM facility of the Nematology Research Unit, member of the UGent TEM-
Expertise center (life sciences). We are grateful to Hanna De Saffel and Petra Vermassen for their
technical assistance on processing the TEM samples. The LC-MS/MS analyses were performed at the
BioMS node at the Proteomics Core Facility, Sahlgrenska Academy, University of Gothenburg with
support from the Swedish National Infrastructure for Biological Mass Spectrometry (BioMS), funded
by the Swedish Research Council.

Funding

DS and FM are a postdoctoral fellow and a senior clinical investigator, respectively, of the Research
Foundation Flanders (FWO), Belgium. This work was supported by a Methusalem Grant
(BOFMET2015000401) from the Ghent University to Prof. Anne De Paepe, by grants from the FWO to
DS (12Q5917N) and to FM (1842318N) and by grant (2017-00955) from the Swedish Research Council
to FN and GL. CG is supported at the UW by a post-doctoral fellowship of the Belgian American
Educational Fund (BAEF). This work was further supported by a Grant-in-Aid for Scientific Research (C)
19K07054 (to SM) from the Japan Society for the Promotion of Science, Japan; by the Takeda Science
Foundation (to SM); Grant-in Aid for Research Center for Pathogenesis of Intractable Diseases from
the Research Institute of Meijo University (SM and SY).

References

References

1.   Prydz K, Dalen KT. Synthesis and sorting of proteoglycans - Commentary. J Cell Sci.
2000;113(2):193-205.

2.   Couchman JR, Pataki CA. An introduction to proteoglycans and their localization. J Histochem
Cytochem. 2012;60(12):885-97.

3.   Lindahl U, Couchman J, Kimata K, Esko JD. Proteoglycans and Sulfated Glycosaminoglycans.
In: rd, Varki A, Cummings RD, Esko JD, Stanley P, Hart GW, et al., editors. Essentials of Glycobiology.
Cold Spring Harbor (NY)2015. p. 207-21.

4.   Selleck SB. Proteoglycans and pattern formation: sugar biochemistry meets developmental
genetics. Trends Genet. 2000;16(5):206-12.

5.   Malfait F, Kariminejad A, Van Damme T, Gauche C, Syx D, Merhi-Soussi F, et al. Defective
Initiation of Glycosaminoglycan Synthesis due to B3GALT6 Mutations Causes a Pleiotropic Ehlers-
Danlos-Syndrome-like Connective Tissue Disorder. Am J Hum Genet. 2013;92(6):935-45.

6.   Sugahara K, Mikami T, Uyama T, Mizuguchi S, Nomura K, Kitagawa H. Recent advances in the
structural biology of chondroitin sulfate and dermatan sulfate. Curr Opin Struct Biol. 2003;13(5):612-
20.

7.   Kusche-Gullberg M, Kjellen L. Sulfotransferases in glycosaminoglycan biosynthesis. Curr Opin
Struct Biol. 2003;13(5):605-11.

8.   Bui C, Ouzzine M, Talhaoui I, Sharp S, Prydz K, Coughtrie MW, et al. Epigenetics: methylation-
associated repression of heparan sulfate 3-O-sulfotransferase gene expression contributes to the
invasive phenotype of H-EMC-SS chondrosarcoma cells. FASEB J. 2010;24(2):436-50.

9.   Gotting C, Kuhn J, Zahn R, Brinkmann T, Kleesiek K. Molecular cloning and expression of
human UDP-d-Xylose:proteoglycan core protein beta-d-xylosyltransferase and its first isoform XT-II. J
Mol Biol. 2000;304(4):517-28.

10.   Almeida R, Levery SB, Mandel U, Kresse H, Schwientek T, Bennett EP, et al. Cloning and
expression of a proteoglycan UDP-galactose:beta-xylose beta1,4-galactosyltransferase I. A seventh
member of the human beta4-galactosyltransferase gene family. J Biol Chem. 1999;274(37):26165-71.

11.   Bai X, Zhou D, Brown JR, Crawford BE, Hennet T, Esko JD. Biosynthesis of the linkage region of
glycosaminoglycans: cloning and activity of galactosyltransferase II, the sixth member of the beta 1,3-
galactosyltransferase family (beta 3GalT6). J Biol Chem. 2001;276(51):48189-95.

12.   Kitagawa H, Tone Y, Tamura J, Neumann KW, Ogawa T, Oka S, et al. Molecular cloning and
expression of glucuronyltransferase I involved in the biosynthesis of the glycosaminoglycan-protein
linkage region of proteoglycans. J Biol Chem. 1998;273(12):6615-8.

13.   Wen J, Xiao J, Rahdar M, Choudhury BP, Cui J, Taylor GS, et al. Xylose phosphorylation

functions as a molecular switch to regulate proteoglycan biosynthesis. Proc Natl Acad Sci U S A.

2014;111(44):15723-8.

14.   Jones KL, Schwarze U, Adam MP, Byers PH, Mefford HC. A homozygous B3GAT3 mutation

causes a severe syndrome with multiple fractures, expanding the phenotype of linkeropathy

syndromes. Am J Med Genet A. 2015;167A(11):2691-6.

15.   Guo MH, Stoler J, Lui J, Nilsson O, Bianchi DW, Hirschhorn JN, et al. Redefining the progeroid

form of Ehlers-Danlos syndrome: report of the fourth patient with B4GALT7 deficiency and review of

the literature. Am J Med Genet A. 2013;161A(10):2519-27.

16.   Nakajima M, Mizumoto S, Miyake N, Kogawa R, Iida A, Ito H, et al. Mutations in B3GALT6,

which encodes a glycosaminoglycan linker region enzyme, cause a spectrum of skeletal and

connective tissue disorders. Am J Hum Genet. 2013;92(6):927-34.

17.   Malfait F, Kariminejad A, Van Damme T, Gauche C, Syx D, Merhi-Soussi F, et al. Defective

initiation of glycosaminoglycan synthesis due to B3GALT6 mutations causes a pleiotropic Ehlers-

Danlos-syndrome-like connective tissue disorder. Am J Hum Genet. 2013;92(6):935-45.

18.   Malfait F, Francomano C, Byers P, Belmont J, Berglund B, Black J, et al. The 2017 international

classification of the Ehlers-Danlos syndromes. Am J Med Genet C Semin Med Genet. 2017;175(1):8-

26.

19.   Ritelli M, Chiarelli N, Zoppi N, Dordoni C, Quinzani S, Traversa M, et al. Insights in the

etiopathology of galactosyltransferase II (GalT-II) deficiency from transcriptome-wide expression

profiling of skin fibroblasts of two sisters with compound heterozygosity for two novel B3GALT6

mutations. Mol Genet Metab Rep. 2015;2:1-15.

20.   Van Damme T, Pang X, Guillemyn B, Gulberti S, Syx D, De Rycke R, et al. Biallelic B3GALT6

mutations cause spondylodysplastic Ehlers-Danlos syndrome. Hum Mol Genet. 2018;27(20):3475-87.

21.   Santoriello C, Zon LI. Hooked! Modeling human disease in zebrafish. J Clin Invest.

2012;122(7):2337-43.

22.   Dooley K, Zon LI. Zebrafish: a model system for the study of human disease. Curr Opin Genet

Dev. 2000;10(3):252-6.

23.   Eames BF, Singer A, Smith GA, Wood ZA, Yan YL, He X, et al. UDP xylose synthase 1 is

required for morphogenesis and histogenesis of the craniofacial skeleton. Dev Biol. 2010;341(2):400-

15.

24.   Eames BF, Yan YL, Swartz ME, Levic DS, Knapik EW, Postlethwait JH, et al. Mutations in

fam20b and xylt1 reveal that cartilage matrix controls timing of endochondral ossification by

inhibiting chondrocyte maturation. PLoS Genet. 2011;7(8):e1002246.

25. Amsterdam A, Nissen RM, Sun Z, Swindell EC, Farrington S, Hopkins N. Identification of 315

genes essential for early zebrafish development. Proc Natl Acad Sci U S A. 2004;101(35):12792-7.

26. Nissen RM, Amsterdam A, Hopkins N. A zebrafish screen for craniofacial mutants identifies

wdr68 as a highly conserved gene required for endothelin-1 expression. BMC Dev Biol. 2006;6:28.

27. Noborn F, Gomez Toledo A, Sihlbom C, Lengqvist J, Fries E, Kjellen L, et al. Identification of

chondroitin sulfate linkage region glycopeptides reveals prohormones as a novel class of

proteoglycans. Mol Cell Proteomics. 2015;14(1):41-9.

28. Persson A, Nilsson J, Vorontsov E, Noborn F, Larson G. Identification of a non-canonical

chondroitin sulfate linkage region trisaccharide. Glycobiology. 2019;29(5):366-71.

29. Nash RDM, Valencia AH, Geffen AJ. The origin of Fulton's condition factor - Setting the record

straight. Fisheries. 2006;31(5):236-8.

30. Arratia G, Schultze HP. Reevaluation of the caudal skeleton of certain actinopterygian fishes:

III. Salmonidae. Homologization of caudal skeletal structures. J Morphol. 1992;214(2):187-249.

31. McMenamin SK, Chandless MN, Parichy DM. Working with zebrafish at postembryonic

stages. Methods Cell Biol. 2016;134:587-607.

32. Parichy DM, Elizondo MR, Mills MG, Gordon TN, Engeszer RE. Normal table of postembryonic

zebrafish development: staging by externally visible anatomy of the living fish. Dev Dyn.

2009;238(12):2975-3015.

33. Hur M, Gistelinck CA, Huber P, Lee J, Thompson MH, Monstad-Rios AT, et al. MicroCT-based

phenomics in the zebrafish skeleton reveals virtues of deep phenotyping in a distributed organ

system. Elife. 2017;6.

34. Gistelinck C, Gioia R, Gagliardi A, Tonelli F, Marchese L, Bianchi L, et al. Zebrafish Collagen

Type I: Molecular and Biochemical Characterization of the Major Structural Protein in Bone and Skin.

Sci Rep. 2016;6:21540.

35. Plaut I. Effects of fin size on swimming performance, swimming behaviour and routine

activity of zebrafish Danio rerio. J Exp Biol. 2000;203(Pt 4):813-20.

36. Wittkopp N, Huntzinger E, Weiler C, Sauliere J, Schmidt S, Sonawane M, et al. Nonsense-

mediated mRNA decay effectors are essential for zebrafish embryonic development and survival. Mol

Cell Biol. 2009;29(13):3517-28.

37. Chen YH, Narimatsu Y, Clausen TM, Gomes C, Karlsson R, Steentoft C, et al. The GAGOme: a

cell-based library of displayed glycosaminoglycans. Nat Methods. 2018;15(11):881-8.

38. Delbaere S, Van Damme T, Syx D, Symoens S, Coucke P, Willaert A, et al. Hypomorphic

zebrafish models mimic the musculoskeletal phenotype of β4GalT7-deficient Ehlers-Danlos

syndrome. Matrix Biol. 2019.

39.   Holmborn K, Habicher J, Kasza Z, Eriksson AS, Filipek-Gorniok B, Gopal S, et al. On the roles

and regulation of chondroitin sulfate and heparan sulfate in zebrafish pharyngeal cartilage

morphogenesis. J Biol Chem. 2012;287(40):33905-16.

40.   Gistelinck C, Witten PE, Huysseune A, Symoens S, Malfait F, Larionova D, et al. Loss of Type I

Collagen Telopeptide Lysyl Hydroxylation Causes Musculoskeletal Abnormalities in a Zebrafish Model

of Bruck Syndrome. J Bone Miner Res. 2016;31(11):1930-42.

41.   Fiedler IAK, Schmidt FN, Wolfel EM, Plumeyer C, Milovanovic P, Gioia R, et al. Severely

Impaired Bone Material Quality in Chihuahua Zebrafish Resembles Classical Dominant Human

Osteogenesis Imperfecta. Journal of Bone and Mineral Research. 2018;33(8):1489-99.

42.   Bishop N. Bone Material Properties in Osteogenesis Imperfecta. J Bone Miner Res.

2016;31(4):699-708.

43.   Viguet-Carrin S, Garnero P, Delmas PD. The role of collagen in bone strength. Osteoporos Int.

2006;17(3):319-36.

44.   Grabner B, Landis WJ, Roschger P, Rinnerthaler S, Peterlik H, Klaushofer K, et al. Age- and

genotype-dependence of bone material properties in the osteogenesis imperfecta murine model

(oim). Bone. 2001;29(5):453-7.

45.   Kalamajski S, Oldberg A. The role of small leucine-rich proteoglycans in collagen

fibrillogenesis. Matrix Biol. 2010;29(4):248-53.

46.   Gillies AR, Lieber RL. Structure and function of the skeletal muscle extracellular matrix.

Muscle Nerve. 2011;44(3):318-31.

47.   Caraffi SG, Maini I, Ivanovski I, Pollazzon M, Giangiobbe S, Valli M, et al. Severe Peripheral

Joint Laxity is a Distinctive Clinical Feature of Spondylodysplastic-Ehlers-Danlos Syndrome (EDS)-

B4GALT7 and Spondylodysplastic-EDS-B3GALT6. Genes (Basel). 2019;10(10).

48.   Gistelinck C, Kwon RY, Malfait F, Symoens S, Harris MP, Henke K, et al. Zebrafish type I

collagen mutants faithfully recapitulate human type I collagenopathies. Proc Natl Acad Sci U S A.

2018;115(34):E8037-E46.

49.   Bird NC, Mabee PM. Developmental morphology of the axial skeleton of the zebrafish, Danio

rerio (Ostariophysi: Cyprinidae). Dev Dyn. 2003;228(3):337-57.

50.   Morin-Kensicki EM, Melancon E, Eisen JS. Segmental relationship between somites and

vertebral column in zebrafish. Development. 2002;129(16):3851-60.

51.   Huysseune A, Sire JY. Bone and cartilage resorption in relation to tooth development in the

anterior part of the mandible in cichlid fish: a light and TEM study. Anat Rec. 1992;234(1):1-14.

52.   Brett JR. The Respiratory Metabolism and Swimming Performance of Young Sockeye Salmon.

J Fish Res Board Can. 1964;21(5):1183-226.

53.   Mizumoto S, Mikami T, Yasunaga D, Kobayashi N, Yamauchi H, Miyake A, et al. Chondroitin 4-

O-sulfotransferase-1 is required for somitic muscle development and motor axon guidance in

zebrafish. Biochem J. 2009;419(2):387-99.

54.   Mizumoto S, Sugahara K. Glycosaminoglycan chain analysis and characterization

(glycosylation/epimerization). Methods Mol Biol. 2012;836:99-115.

55.   Noborn F, Gomez Toledo A, Green A, Nasir W, Sihlbom C, Nilsson J, et al. Site-specific

identification of heparan and chondroitin sulfate glycosaminoglycans in hybrid proteoglycans. Sci

Rep. 2016;6:34537.

56.   Baasanjav S, Al-Gazali L, Hashiguchi T, Mizumoto S, Fischer B, Horn D, et al. Faulty initiation of

proteoglycan synthesis causes cardiac and joint defects. Am J Hum Genet. 2011;89(1):15-27.

57.   Job F, Mizumoto S, Smith L, Couser N, Brazil A, Saal H, et al. Functional validation of novel

compound heterozygous variants in B3GAT3 resulting in severe osteopenia and fractures: expanding

the disease phenotype. BMC Med Genet. 2016;17(1):86.