A whole-cortex probabilistic diffusion tractography connectome

Burke Q. Rosen[a] and Eric Halgren[a,b]

[a] Neurosciences graduate program, University of California, San Diego
 La Jolla, CA, 92093, United States
[b] Departments of Radiology and Neurosciences, University of California, San Diego
 La Jolla, CA, 92093, United States

Correspondence:
bqrosen@ucsd.edu
Burke Rosen
UC San Diego Radiology Imaging Laboratory
3510 Dunhill St. San Diego, CA 92121
United States

Abstract

The WU-Minn Human Connectome Project (HCP) is a publicly-available dataset containing

state-of-art structural, functional, and diffusion-MRI for over a thousand healthy subjects. While

the planned scope of the HCP included an anatomical connectome, resting-state functional-

MRI forms the bulk of the HCP's current connectomic output. We address this by presenting a

full-cortex connectome derived from probabilistic diffusion tractography and organized into the

HCP-MMP1.0 atlas. Probabilistic methods and large sample sizes are preferable for whole-

connectome mapping as they increase the fidelity of traced low-probability connections. We

find that overall, connection strengths are lognormally distributed and decay exponentially with

tract length, that connectivity reasonably matches macaque histological tracing in homologous

areas, that contralateral homologs and left-lateralized language areas are hyperconnected, and

that hierarchical similarity influences connectivity. We compare the diffusion-MRI connectome

to existing resting-state fMRI and cortico-cortico evoked potential connectivity matrices and

find that it is more similar to the latter. This work helps fulfill the promise of the HCP and will

30 make possible comparisons between the underlying structural connectome and functional

31 connectomes of various modalities, brain states, and clinical conditions.

33 **Keywords:** Diffusion MRI, structural connectome, tractography, Human Connectome Project

Significance Statement

The tracts between cortical parcels can be estimated from diffusion MRI, but most studies

concentrate on only the largest connections. Here we present an atlas, the largest and most

detailed of its kind, showing connections among all cortical parcels. Connectivity is relatively

enhanced between frontotemporal language areas and homologous contralateral locations. We

find that connectivity decays with fiber tract distance more slowly than predicted by brain

volume and that structural and stimulation-derived connectivity are more similar to each other

than to resting-state functional MRI correlations. The connectome presented is publicly

available and organized into a commonly used scheme for defining brain areas in order to

enable ready comparison to other brain imaging datasets of various modalities.

Introduction

In the 21st century, advances in computation, theory, and neuroimaging have spurred a broad

and intense interest in the anatomical connections and physiological correlations among

human brain areas. Bivariate functional connectivity has given way to full functional

connectomes, the most comprehensive of which may be the WU-Minn Human Connectome

Project's (HCP) resting-state fMRI dense connectome (Van Essen et al., 2013). The planned

scope of WU-Minn HCP also included a full anatomical connectome (Van Essen and Ugurbil,

2017), and the project has collected, curated, and preprocessed diffusion imaging (dMRI) data

for 1,065 subjects. However, a structural connectome has to-date not been released for these

data. This report seeks to address this omission by presenting a full-cortex anatomical

connectome derived from local, probabilistic tractography.

dMRI techniques detect white matter by registering the orientation biases of water

molecule diffusion within myelinated axons. The majority of dMRI studies focus on differences

in specific connections between treatment groups. In contrast, we seek here to present a

robust, densely populated average connectivity matrix for the entire cortex using data from a

large, healthy sample. Local dMRI fiber tract tracing algorithms can be broadly organized into

two classes: deterministic e.g. dsi-studio (Yeh et al., 2013), and probabilistic e.g. probtrackX

(Behrens et al., 2007). Deterministic tractography considers the most likely orientation at each

voxel yielding the maximum likelihood tracts whereas probabilistic tractography considers the

entire distribution of possible orientations, yielding a probability cloud of connections. As our

goal is instead to explore all possible connections between regions, we employed local,

probabilistic tractography (Behrens et al., 2007). This method has been validated against

macaque retrograde tracers within-species (Donahue et al., 2016) and the dMRI protocol and

equipment used for the WU-Minn HCP database were optimized in anticipation of this analysis

(Sotiropoulos et al., 2013).

The physiological relevance of a connectome is maximized if its nodes form functionally

distinct areas. Within the scope of cortex, this amounts to selecting a parcellation scheme. The

HCP multi-modal parcellation (HCP-MMP1.0) (Glasser et al., 2016) has several advantages: it's

boundaries are both functionally and anatomically guided, it has sufficient parcels (360) to

generate a rich connectome while few enough that the parcels' extents comfortably exceed the

dMRI voxel size, and mechanisms exist (Fischl et al., 2004) for it to be readily applied to

individuals. Most importantly, the HCP-MMP1.0 parcellation is publicly available and widely

adopted, facilitating the comparison of the generated matrices to other structural and

functional connectomes.

Given the computational intensity of dMRI fiber tractography and the field's inclination

towards elucidating specific connections, it is not surprising that the number of existing

publicly available dMRI datasets exceeds that of finished, readily applicable connectivity

matrices. However, there do exist some prior examples. The USC Multimodal connectivity

database (http://umcd.humanconnectomeproject.org), contains two dMRI tractography

connectomes with standard surface-based parcellations: Hagmann (Hagmann et al., 2008) and

ICBM (Mori et al., 2008), with sample-sizes of 5 and 138, respectively. A third is available at

http://www.dutchconnectomelab.nl which contains 114 controls. All of these use the Desikan-

Killiany atlas (Desikan et al., 2006) which consists of 68 cortical parcels and were produced

with deterministic tractography. An atlas of major fiber tracts for the HCP 1200 cohort has

recently released at http://brain.labsolver.org, (Yeh et al., 2018). However, this deterministic

tractography connectome is spatially coarse, consisting of only 54 cortical parcels, and lacks

dynamic range and statistical dispersion, as weaker connections are unrepresented, rendering

the connectivity matrix nearly binary. The HCP-MMP1.0 atlas employed here has more than

five times as many parcels while retaining the functional distinctness of areas. In contrast to the

relatively sparse existing deterministic matrices, the probabilistic approach may better resolve

weak or low probability connections leading to densely populated connectivity matrices like

those found non-human primate tracing studies (Markov et al., 2014). Furthermore, the cohort

studied is large and many other types of data are available for the same individuals including

the NIH neuropsychological toolbox (Gershon et al., 2013), as well as fMRI and MEG data for

resting-state and cognitive tasks, permitting within-cohort comparison to functional

connectivity.

The following report presents a novel structural connectome of the human neocortex based

on probabilistic diffusion tractography. The connectome is partially validated against

retrograde tracing in macaques and the relationship between tract length and connection

strength is quantified. Further validation is provided by reasonable connectivity properties

between contralateral homologous parcels, within language cortex, and between parcels lying

at similar levels of the cortical hierarchy. Finally, the dMRI connectome is compared to cortico-

cortico evoke potential (CCEP) and resting-state fMRI derived connectivity.

Materials & Methods

Subjects & data sources

No new data was collected for this study, and the existing data used was gathered

from publicly available databases. Individual subject's high-resolution T1-weighted

structural magnetic resonance volumes (MRI), Diffusion images (dMRI), and group

average grayordinate resting-state function MRI (rs-fMRI) connectivity were gathered

from the Human Connectome Project's (HCP) WU-Minn 1200 release (Van Essen et al.,

2013) at https://db.humanconnectome.org. The diffusion imaging dataset consists of

1065 individuals (575 women), aged 22-36+ years old. These datasets include some

twin and non-twin siblings. However, individuals' family structure, as well as exact age,

handedness, and ethnicity are access-restricted to protect the privacy of the subjects

and these data were not requested as they are not critical to this study. Group-average

dense T1w/T2w myelination index were gathered from the same source. Macaque

retrograde tracer connectivity was sourced from supplementary table 6 of (Markov et

al., 2014). Parcel-by-parcel values were averaged across monkey and hemisphere.

Group average, parcellated cortco-cortico evoked potential (CCEP) connectivity was

gathered from the v1903 release of the Functional Brain Tractography project (F-

TRACT) (David et al., 2013; Trebaul et al., 2018) at https://f-tract.eu.

Cortical parcellation & functional networks

The HCP multimodal parcellation scheme (HCP-MMP1.0), consisting of 180 cortical parcels

per hemisphere, was projected from the Workbench (Marcus et al., 2011) 32k grayordinate

template brain to the FreeSurfer (Fischl, 2012) ico5 fsaverage template as per (Coalson et al.,

2016). Using the FreeSurfer reconstruction directories gathered from the database, surface-

based fsaverage parcel labels were mapped onto each individual's white matter surface using

spherical landmark registration (`fs_label2label`), (Fischl et al., 1999). Grayordinate rs-fMRI

connectivity values were morphed to the ico5 fsaverage template then averaged within each

parcel. Finally, individual's surface-based parcel labels were converted to binary volumes

marking the gray matter — white matter boundary (`mri_label2vol`) to serve as seed and

target regions for probabilistic tractography. Workbench and FreeSurfer functions were

sourced from releases 1.2.3 and 6.0, respectively.

To facilitate interpretation of the connectome, parcels were ordered and grouped into

functional networks adapted from (Ji et al., 2019), which applied iterative Louvain clustering

(Blondel et al., 2008; Rubinov and Sporns, 2010) and other criteria to a resting-state fMRI

connectivity. These functional groupings and parcel order were selected as they were also

generated using (a subset of) the WU-Minn HCP dataset and the HCP-MMP1.0 parcellation

scheme. For this study the parcels of the left and right hemispheres were separated and the

order and groupings of the left hemisphere in (Ji et al., 2019) were used for homologous

parcels in the both right and left hemisphere, respectively. Two pairs of the original networks

(primary and secondary visual, ventral and posterior multimodal) contained too few parcels for

effective analysis and were highly inter-related. These network pairs were simplified by

combining them into visual and multimodal groups, yielding 10 functional networks per

hemisphere, see table **2**.

Probabilistic tractography

All analysis of diffusion imaging data was performed with FSL (Behrens et al., 2007; Jenkinson

et al., 2012) release 6.0.1. Analyses were performed identically for each subject and broadly

follow (Burns, 2014). The diffusion and bedpostX precursor directories made available from the

HCP database were used as inputs without modification. The WU-Minn HCP diffusion data are

is correction for eddy currents and movement with FSL eddy (Andersson and Sotiropoulos,

2016). Subjects' estimated displacement over time from their initial position is written to the

eddy_restricted_movement_rms output. Using these data , a scalar index of each subject's

motion was derived by integrating their displacement over time.

Fractional anisotropy (FA) analysis was performed using `dtifit`. The resulting FA volumes

were not analyzed but only used for registering the FreeSurfer and dMRI volumes (`flirt`), as is

necessary to map the parcel masks into dMRI space (`probtrackx2` arguments `--xfm --`

`seedref`). Non-invasive probabilistic tractography was performed with `probtrackx2` in voxel-

by-parcel mode (`--os2t --s2tastext`). In this configuration, the number and length of

streamlines (`--ompl --opd`) is estimated from each voxel in the seed parcel to each target

parcel as a whole. To aid parallelization of these computationally intensive processes, the list

of target parcels (`--targetmasks`) was quartered into four sub-lists. Therefore `probtrackx2`

was invoked 1440 times per subject, estimating the connectivity between 1 seed parcel and 90

target parcels in each invocation. The default ½ voxel step length, 5000 samples and 2000

steps were used (`--steplength 0.5 -P 5000 -S 2000`). To avoid artifactual loops,

streamlines that loop back on themselves were discarded (`-l`) and tractography was

constrained by a 90° threshold (`-c 0`) for maximal curvature between successive steps. Within-

parcel connectivity and cotico-subcortical connectivity were not examined in this study. All

post-hoc analyses and visualization of connectivity data were performed in Matlab 2019b

(Mathworks) except for figure **1C** which was rendered in fsleyes.

Normalization & symmetrization

Raw streamline counts were averaged across all subjects, then normalized and symmetrized

following procedure developed for non-human primate histological tracing (Donahue et al.,

2016; Theodoni et al., 2020). Briefly, fractionally scaled values are defined as the ratio of the

number of streamlines originating at parcel A and terminating at parcel B to the total number of

streamlines that either originate at parcel A or terminate at parcel B while excluding within-

parcel connections.

Eq. 1 $F\left(DTI_{i,j}\right) = \dfrac{DTI_{i,j}}{\sum_{x=1}^{N} DTI_{i,x} + \sum_{y=1}^{N} DTI_{y,j}}, where\ x \neq i\ \&\ y \neq j$

Fractional scaling is one of several plausible normalization strategies. Because we used

5000 samples (-P 5000) and voxel-by-parcel mode (--os2t) in our probtrackX invocation, the

maximum possible raw streamline count between any two parcels is 5000N where N is the # of

voxels in the seed parcel. Note that because, for probtrackX, all parcels were defined as a

single layer of 1mm isotropic voxels at the white matter — gray matter interface, Ni is also

equivalent to the area of the seed parcel, in mm^2. As shown in extended data figure **1-1,** We

examined four strategies for normalizing the raw streamline counts: (1) dividing by the number

of samples, 5000, (2) dividing by the number of samples and seed area, $5000N_i$, (3) dividing by

the number of samples and the areas of both the seed and target parcels, $5000N_i^{0.5}N_j^{0.5}$, and (4)

fractional scaling, see Eq. 1. These approaches yield similar connectivity matrices, distributions

of pairwise connectivity, and rates of connectivity fall-off with fiber tract distance. The choice of

normalization does shift the absolute scale of pairwise connectivity strengths, but as this effect

is mostly homogenous across all connections, subsequent analyses are not greatly affected.

The correlation coefficient of connectivity strengths between normalization techniques exceeds

0.97 for all pairwise comparisons, and exceeds 0.99 if the samples-only normalization

approach is excluded (data not shown).

While diffusion tractography is not sensitive to the directionality of connections, because

parcel A to B and parcel B to A streamlines are computed separately minor asymmetries arise.

Connectivity matrix symmetry is enforced by taking the arithmetic mean of the A-B and B-A

fractionally scaled connection weights.

Eq. 2 $F_{i,j} = \frac{F_{i,j}+F_{j,i}}{2}$

Because probabilistic tractography values span several orders of magnitude, and are

approximately log-normally distributed (Fig. **1-1 B**), data were log-transformed (log_{10}) prior to

subsequent analyses. The CCEP and rs-fMRI connectivity matrices were (re)normalized

following the same procedure. However the rsMRI connectivity values were not log-

transformed because these data are already approximately normally distributed, if bimodal, in

linear space, see figure **9B**.

Network theory metrics

All network theoretic measures were computed in matlab using the Brain Connectivity Toolbox,

2019-03-03 release (Rubinov and Sporns, 2010). It is available at http://www.brain-

connectivity-toolbox.net or https://www.nitrc.org/projects/bct. The definitions for the metrics

used (for binary and undirected networks) are repeated below.

Precursor measures

Eq. 3 $d_{i,j} = \sum_{a_{u,v} \in g_{i \leftrightarrow j}} a_{u,v}$

Where $d_{i,j}$ is the shortest path length, a basis for measuring integration, between nodes i and j,

N is the set of all nodes in the network, n is the number of nodes, and $a_{u,v}$ is the binarized

connectivity between nodes u and v.

Eq. 4 $t_i = \frac{1}{2} \sum_{j,h \in N} a_{i,j} a_{i,h} a_{j,h}$

Where t_i is the number of triangles, a basis for measuring integration, around node i.

Eq. 5 $k_i = \sum_{j \in N} a_{i,j}$

Where k_i is the number of degree, or number of links, connected to node i.

Mean Clustering Coefficient (MCC)

Eq. 6 $C_i = \frac{1}{n}\Sigma_{i \in N}\frac{2t_i}{k_i(k_i-1)}$,

Where C_i is the clustering coefficient of node i. ($C_i = 0$ for $k_i < 2$), (Watts and Strogatz, 1998).

Eq. 7 $MCC = \frac{1}{n}\Sigma_{i \in N} C_i$

Characteristic Path Length (CPL)

Eq. 8 $L_i = \frac{1}{n}\Sigma_{i \in N}\frac{\Sigma_{j=1,j \neq i}^{n} d_{i,j}}{n-1}$

Where L_i is the number of the average distance between node i and all other nodes, (Watts and

Strogatz, 1998).

Eq. 9 $CPL = \frac{1}{n}\Sigma_{i \in N} L_i$

Global Efficiency

Eq. 10 $E_i = \frac{1}{n}\Sigma_{i \in N}\frac{\Sigma_{j \in N,j \neq i}^{n} d_{i,j}^{-1}}{n-1}$

Where E_i is the efficiency of node i.

Eq. 11 $E = \frac{1}{n}\Sigma_{i \in N} L_i$

Where E is the global efficiency of the network, (Latora and Marchiori, 2001).

<u>Modularity</u>

Eq. 12 $Q = \frac{1}{l}\sum_{i,k\in\mathrm{N}}\left(a_{i,j} - \frac{k_i k_j}{l}\right)\delta_{m_i,m_j}$

Where l is the number of links in the network, m_i is module containing node i, δ_{m_i,m_j} = 1 if m_i =

m_j, and 0 otherwise, and Q is the global efficiency of the network, (Newman, 2004).

<u>Gamma (normalized MCC)</u>

Eq. 12 $\gamma = \frac{MCC}{MCC_{rand}}$

Where MCC_{rand} is the MCC of a random network of the same statistical makeup.

<u>Lambda (normalized CPL)</u>

Eq. 13 $\lambda = \frac{CPL}{CPL_{rand}}$

Where CPL_{rand} is the CPL of a random network of the same statistical makeup. Note that this

measure is unrelated to the length constant λ.

<u>Small Worldness</u>

Eq. 14 $S = \frac{\gamma}{\lambda}$

Where S is the network small-worldness (Humphries and Gurney, 2008).

<u>Transitivity</u>

Eq. 15 $T = \frac{\sum_{i\in N} 2t_i}{\sum_{i\in N} k_i(k_i-1)}$

Where T is the transitivity of the network (Newman, 2003).

<u>Assortativity</u>

Eq. 16 $r = \frac{l^{-1}\sum_{(i,j)\in L} k_i k_j - \left[l^{-1}\sum_{(i,j)\in L}\frac{1}{2}(k_i+k_j)\right]^2}{l^{-1}\sum_{(i,j)\in L}\frac{1}{2}(k_i^2+k_j^2) - \left[l^{-1}\sum_{(i,j)\in L}\frac{1}{2}(k_i+k_j)\right]^2}$

Where L is the set of all links and r is the assortativity coefficient of the network (Newman,

2003).

<u>Network Density</u>

Eq. 17 $D = \frac{l}{n^2-n}$

Where D is the density of the network before thresholding and binarization.

Results

A whole-cortex structural connectome

Figure **1A** shows the group average parcel to parcel and probabilistic diffusion tractography

connectome. This matrix consists of connectivity among 360 cortical parcels and is further

organized into 10 functional groups (Ji et al., 2019) per hemisphere. The raw probabilistic

tractography streamline counts have been normalized by fractionally scaling (Eq. 1) into log

probabilities (F_{pt}) following procedures developed for tracing non-human primate connectivity.

As dMRI reveals structural connections, the network is undirected and therefore symmetric.

The main diagonal is masked as intra-parcel connectivity was not examined in this study. The

upper left quadrant shows connectivity among the 180 parcels of the left hemisphere, the

lower right quadrant the connectivity within the right hemisphere. The upper right and lower left

quadrants are duplicates and show the inter-hemispheric, or callosal, connections. The 180^{th}

(or half-) diagonal is clearly visible (white arrows); this shows the connectivity between

homologous parcels in the right and left hemispheres, which is greater than non-homologous

callosal connectivity for most parcels.

After $\log_{10}$ transformation, F_{pt} connectivity among all parcel pairs is approximately Gaussian

in distribution with a mean -3.903 ($CI_{95\%}$ = [-3.910 -3.897]), standard deviation 0.8111 ($CI_{95\%}$ =

[0.806 0.816]), skewness 0.627 ($CI_{95\%}$ = [0.6082 0.6440]), and kurtosis 3.605 ($CI_{95\%}$ = [3.5603

3.6498]). In addition to bringing the range of F_{pt} values into the same order of magnitude, $\log_{10}$

transformation is justified as it brings the distribution's skewness significantly closer to zero

(pre-$\log_{10}$: 9.047, $CI_{95\%}$ = [8.719 9.469]), and kurtosis significantly closer to three, pre-$\log_{10}$:

103.684 ($CI_{95\%}$ = [93.991 117.026]) thus bringing the distribution closer to normality. See

extended data figure **1-1 B**,**C** for a graphical comparison. Empirical confidence intervals were

estimated via bootstrapping with 2000 iterations. The values of the group average and

individual probabilistic dMRI connectivity matrices, as well as all other figure source data can

be found at https://doi.org/10.5281/zenodo.4060485.

Tract length strongly predicts connectivity strength, with exponential decay

In addition to the connection strength, diffusion tractography estimates the fiber tract length

between all pairs of parcels. As shown in figure **2**, structural connectivity ($10^\wedge F_{pt}$) falls off as an

exponential function of fiber tract length with the form $10^\wedge F_{pt} = \alpha * e^{-d/\lambda}$ where λ is the length

constant, α the scaling coeffect, and d the tract length. Alternative functional forms were

examined (see figure **2-2),** but the exponential was selected for parsimony, goodness-of-fit,

and concordance with histological tracing data (see Discussion). Note that λ is sometimes

reported in inverted units of mm^{-1}, e.g. (Markov et al., 2013; Theodoni et al., 2020), but we here

use the λ convention from neuronal cable theory (Dayan and Abbott, 2001) which has more

intuitive units (mm); the conventions are conceptually equivalent. For the group-average

connectome, $\lambda = 23.4$ mm and the least squares exponential fit explains 84% of the variance in

$10^\wedge F_{pt}$ across all parcel pairs. Callosal connectivity, when isolated, decays more slowly with

respect to tract length, $\lambda = 32.8$, and hews to the exponential expectation less consistently $r^2 =$

0.62. Because the tracing of long fiber tracts may be hampered by poor scan quality, we

investigated the effects of subjects' motion on λ. For each subject, λ was calculated for non-

zero connections in the same manner as the group average. While subjects' motion within the

scanner does reduce λ, this effect is modest, only explaining 1.96% of the inter-subject

variance, see figure **2-2**.

Inter-individual variability

The inter-individual variability of connectivity was assessed by deriving the across-subject

coefficient of variation, CV, for each pairwise connection F_{pt}, see figure **3**. For this analysis, the

normalization, symmetrization, and $\log_{10}$-transformation of raw connectivity values was

performed on each subject. Pairwise connections with zero streamlines were not log-

transformed in order to avoid infinities. While there is no clear relationship between fiber tract

distance and inter-individual variability, the most consistent connection appear in two clusters

of around 50-100 mm and 170-225 mm (Fig. **3B**) . When the most consistent quintile of

connections is isolated (Roberts et al., 2017), connectivity falls off more slowly with tract

distance, with λ increasing to ~28 mm (Fig. **3D**). Since the proportional size of V1/V2 varies ~3-

fold across individuals and is highly heritable (Yoon et al., 2019), we hypothesized that the

ipsilateral V1-V2 connection would also be highly variable, with that variability being correlated

across hemispheres. Indeed, we find that the ipsilateral V1 – V2 connection is very strong, with

~1.8 fold variability which is strongly correlated across hemispheres (r=0.70). The scatter-plot

of right vs. left F_{pt} values for this connection across subjects (Fig. **3F**) does not reveal obvious

outliers which would be indicative of subject-specific artifacts. This analysis of inter-individual

variability should be considered preliminary. The WU-Minn HCP dataset is rich in individual

data, including the NIH neuropsychological toolbox (Gershon et al., 2013), twin and non-twin

siblings subsets, and genotypic data (dbGaP phs001364.v1.p1), though the latter two data

types are only available by application in order to ensure subject anonymity. With access to

these data, a full examination of inter-individual variability, including assessing the heritability

and genetic correlates of the strength of specific connections could be made.

Probabilistic dMRI tract tracing in humans reasonably corresponds with histological fiber tracing in macaques

The development of both the HCP-MMP1.0 human cortical atlas (Glasser et al., 2016) and FV91 macaque parcellation scheme (Felleman and Van Essen, 1991) were led by David Van Essen and the parcel definitions of the human atlas were informed by human-macaque homology. As such, the parcel names of these atlases have considerable overlap, particularly for visual and visual association areas as well as the non-visual parcels 1, 2, 25, and 44. We therefore assumed that parcels with the same name were roughly homologous and limited the scope of the inter-species comparison to these parcels. Furthermore, the macaque FLne values found in (Markov et al., 2014) are directly comparable to fractionally scaled Fpt values (Donahue et al., 2016). Comparing the pairwise connectivity between species, we found a Pearson correlation of r = 0.35 (p = 0.0013), see figure 4. Considering that for macaques, Donahue and colleages (2016) found a *within-species*, between-technique correlation of r = 0.59 when comparing retrograde tracing and probabilistic diffusion tractography, we find the magnitude of *between-species* correlation to be reasonable supporting evidence for the efficacy of the technique.

Contralateral connectivity exceeds ipsilateral connectivity in some regions

On the whole, cortical connectivity is dominated by ipsilateral connections. This effect is readily-observed by comparing the ipsilateral and contralateral quadrants of figure **1A**. However, there are exceptions to this rule. The differential connectome of ipsi- vs. contra-lateral connections is shown in figure **5**. This is achieved by subtracting the mean of left-right and right-left contralateral connectivity from the mean of the right and left ipsilateral connectivity, i.e. subtracting the mean of the first and third quadrants from the mean of the second and fourth. A cingulo-parietal somatomotor region (parcels 5m, 5L, 24dd, and 24dv)

are more strongly connected to most contralateral cortex than ipsilateral cortex. Lateromedial

connectivity in select prefrontal (a10p, a9-46v, a10p, p10p, p47r, p9-46v, 11l, IFSa, IFJp, a24,

d32, p32, 10r) and postcentral – superior parietal lobule (LIPv, VIP, 7AL, 7PC, 1, 2, 3a, 6d, 31a,

31pd, PCV) regions is stronger between hemispheres than within them. We speculate that a

possible commonality between these three regions is that they have been broadly implicated in

the unitary processes of somatosensory object recognition, emotion, and spatial cognition,

respectively. Conversely, the entire auditory network and superior temporal cortices (STGa,

STSda, DTDdp, A5, and TPOJ1) as well as the operculum and temporoparietal junction (Ig, MI,

FOP1-FOP5, OP1-OP4, PF, PFcm, PFop, PI, PoI1, PoI2, and 43) have pronounced hyper-

ipsilateral connectivity, consistent with the low transmission latency required for auditory

processing, the left-lateralization of language, and the right lateralization of attention.

With the exception of some language areas, most parcels are disproportionately

connected to their contralateral homologs

The two hemispheres of the cortex have a high degree of functional and anatomical symmetry.

It follows then that most regions will have greater connectivity to their contralateral homologs

than other contralateral areas, in order to coordinate their overlapping processing tasks. This is

hinted at by the visibility of the 180^{th}, (or half-) diagonal in figure **1A**. To further quantify this

effect, for all 180 parcels we compared the connectivity between interhemispheric homologs to

the mean of all other callosal connectivity. Bonferroni corrected, empirical 95% confidence

intervals were estimated via bootstrapping with 2000 iterations. As detailed in extended data

figure **6-1** and visualized in Figure **6**, 147 parcels are hyperconnected to their contralateral

homologs, 18 are hypoconnected, and 15 have homologous callosal connectivity not

significantly different than their callosal mean connectivity. Interestingly, parcels that are not

hyper-connected to their contralateral homologs are concentrated within and adjacent to the

language network, consistent with the greater degree of lateralization in these areas.

The language network is hyper-connected at long distances and left lateralized

In order to investigate distance-resolved left laterality in connections among language-

implicated cortex, pairwise connections were binned by fiber tract length in 15 mm increments.

Within each bin, connections were grouped as being within the combined language and

auditory network, or between the combined networks and the rest of the cortex. For each

subject, the F_{pt} of grouped connections within each bin was averaged before being log-

transformed. The grand-averages of these within- and between- language/auditory cortex in

each distance bin for each hemisphere are shown in figure **7A**. Bonferroni corrected, empirical

95% confidence intervals for these grand-averages were estimated via bootstrapping with

2000 iterations. Within-language connectivity is slightly attenuated at distances less that 100

mm, but strongly amplified at distances above 100 mm, especially ~100-140 mm connections

in the left hemisphere. A plurality of these are between frontal and temporoparietal language

areas (18/45 connections between 100 and 140mm). The differential traces of between- vs.

within-language connectivity (Fig. **7B**) clearly show the left-hemisphere dominance of this

effect.

Connectivity is influenced by the cortical hierarchy

Hierarchy is a central organizing principle of the cortex (Burt et al., 2018; Felleman and Van

Essen, 1991; Markov et al., 2014; Theodoni et al., 2020). Higher order areas, e.g. supporting

abstract processing, have low myelination, and lower order areas, e.g. supporting unimodal

sensory processing, have high myelination. Furthermore, areal myelination is indexed by the

ratio between T1- and T2-wieghted MRI contrast (Glasser and Van Essen, 2011). The WU-Minn

HCP 1200 release includes smoothed group-average myelination indices for all vertices in the

32k grayordinate template brain. These values were averaged for each parcel in the HCP-

MMP1.0 atlas (Glasser et al., 2016) to yield a group-average parcel-wise index of myelination.

The relationship between cortical hierarchy and connectivity was assessed in two ways. We

first examined whether regions of similar level in the cortical hierarchy are better connected, as

predicted by (Barbas, 2015). An index of hierarchical similarity, $F_{|\Delta \text{ myelination}|}$, was obtained for

each pair of parcels by computing the pairwise difference in myelination between parcels and

fractionally scaling it in the same manner as F_{pt}, with smaller values indicating hierarchical

closeness. The similarity matrix created by this derivation is shown in figure **8-1**. Correlations

were obtained for the left and right hemisphere as a whole as well as the colossal connections,

figure **8A**. In addition, for each of the twenty functional networks (10 per hemisphere) the

Pearson correlation between the $F_{|\Delta \text{ myelination}|}$ and F_{pt} for pairwise within-network connections

was computed, see figure **8B**. With the exception of the interhemispheric connections,

calculations were performed on the hemispheres separately to avoid the collinearity introduced

by hemispheric homology.

With the exceptions of the bilateral visual and somatomotor networks and right language

network, for which there is convincingly no relationship, the preponderance of coefficients are

negative, indicating that, on average, areas at similar levels of the cortical hierarchy are better

connected. However, quantified in this way, the influence of hierarchy is modest, explaining

about 1% of the variance in F_{pt} overall, though perhaps 10-30% in certain subsets of parcels,

such as the left auditory and language networks. The left lateralization of the influence of

hierarchy in these networks is striking, as is the right-lateralization of the dorsal attention

network.

Secondly, we investigated whether a cortical region's hierarchical level affected its overall

connectivity. For each parcel, the Pearson correlation between the parcel's F_{pt} to all other

parcels and the parcel-wise index of myelination was computed. In other words, correlation

between each row of the connectome matrix and the vector of myelination indices was obtained.

After Bonferroni correction for multiple comparisons, 74 of 360 parcels (see extended data

figures **8-2, 8-3**) have connectivity significantly correlated to their myelination index and of these

the vast majority (70) are negatively correlated, indicating that low myelination predicts high

connectivity, see figure **8C**. These areas form a contiguous bilateral prefrontal network as

shown in figure **8D**, indicating that prefrontal areas are more connected with higher cortical

regions. The rare positively correlated exceptions are the left and right DVT and V6A.

Probabilistic dMRI connectivity more closely resembles CCEPs than resting-state fMRI

In order to further contextualize the dMRI connectome, we compared it to existing

connectivity matrices generated from two other brain mapping modalities: cortico-cortico

evoked potential probability (CCEP) and resting-state fMRI correlation magnitude (rs-fMRI). As

shown figure **9A**, the qualitative pattern of rs-fMRI markedly differs from the other two

modalities with proportionally stronger ipsilateral across-network connections and especially

non-homologous contralateral connections, though the latter is somewhat obscured for CCEPs

due to sparse spatial sampling. Over all connections, pairwise probabilistic dMRI connectivity

values are nearly twice as linearly correlated to pairwise CCEP connectivity than to rs-fMRI

connectivity (fig. **9B)**, and this contrast is equally evident in the ipsilateral connection within

each hemisphere, see figure **9-1**. Contralateral connections were not examined in isolation as

contralateral sampling for the CCEP modality is relatively rare.

When comparing the distributions of pairwise connectivity strength (fig. **9C**), rs-fMRI again

exhibits properties different than the other two modalities. While both dMRI and CCEP

distributions skew in opposite directions (0.63 and -0.43, respectively), their strengths form

unimodal log-normal distributions and thus shown with log-transformed values. In contrast, rs-

fMRI connectivity values form a bimodal Gaussian-mixture distribution in linear space. The two

modes were characterized by obtaining the maximum-likelihood fit (`fitgmdist`) of a 2-

component Gaussian-mixture to the data, yielding a left mode ($\mu = 0.0011$, $\sigma = 8.1e-8$) forming

63% of the distribution and a right mode ($\mu = 0.0017$, $\sigma = 8.1e-8$) forming 37%, respectively.

Splitting the rs-fMRI modes at the midpoint between their means (0.0014) and plotting their

respective connectivity matrices (fig. **9D**) reveals that the low-connectivity (left) mode consists

primarily of connections between the default mode / frontoparietal networks and other regions

of the cortex.

To further contrast the three connectivity modalities we computed six network theoretic

metrics for each of the connectivity matrices: mean clustering coefficient (MCC), characteristic

path length (CPL), global efficiency, gamma (normalized MCC), lambda (normalized CPL), small

worldness, transitivity, and assortativity (see Appendix). Binarized network metrics were

assessed after thresholding by edge weight (connectivity strength) at intervals of 0.1. Note that

this lambda is unrelated to the exponential length constant reported above. To account for the

order-based arbitrary treatment of equal edge weights when thresholding, the node (parcel)

order was randomized 1000 times, and the mean metric values are shown. Empirical 95%

confidence intervals for these means are too small to be shown at scale. Networks densities

above 0.6 were not examined as the un-thresholded network density of CCEP connectivity

matrix, treating missing data as non-connections, is less than 0.7. However, all measures

appear to converge as binary network density approaches 1. As shown in figure 10, the MCC,

CPL, global efficiency, small worldness, transitivity, and assortativity are markedly different for

rs-fMRI connectivity than for CCEP and probabilistic dMRI tractography, whose metrics as a

function of network density are more similar to each other. Normalizing by metrics computed

for a random network with the same statistical makeup changes this pattern. For gamma the

rs-fMRI and CCEP networks are more similar than either is to probabilistic dMRI tractography,

and lambda rs-fMRI and probabilistic dMRI tractography are more similar than either is to the

CCEP network. The high MCC, transitivity, and assortativity and low global efficiency of rs-

fMRI relative to the other modalities may be indicative of strong, long-range correlativity

beyond that predicted by anatomical connections.

Discussion

In this study we compiled a whole-cortex structural connectome by applying probabilistic

tractography to the diffusion MR volumes of 1065 subjects from the WU-Minn Human

Connectome Project. We report a novel, complete, and high-dynamic-range connectivity

matrix discretized into the 360 parcels of the HCP-MMP1.0 atlas and further arranged into 10

functional networks. It is shown that connectivity strength exponentially decays with fiber tract

length, that the parts of the connectome with clear homology to macaques correspond

reasonably to retrograde tracer mappings in that species, that contralateral homologs are

hyperconnected, and that some connections within language-implicated cortex are stronger

than expected and left-lateralized. While ipsilateral connectivity generally dominates, some

regions have stronger contralateral connections. Inter-individual variability is relatively high for

early visual cortex, whose connectivity co-varies across hemispheres. Cortical areas tend to be

more connected with areas at similar levels of the cortical hierarchy, as indexed by their

estimated myelination, particularly in prefrontal areas. Lastly, it is shown that probabilistic

tractography connectivity more closely resembles that of CCEPs than rs-fMRI. In sum, we

quantify a dMRI-based estimate of medium- to long-range anatomical cortico-cortical

connectivity in a large normative sample.

Diffusion MR imaging and automated post-hoc tractography are powerful tools for the

elucidation of cerebral connectivity. The defining advantages of these techniques are non-

invasiveness and large field-of-view, enabling whole-brain mapping in humans. However, dMRI

does have significant limitations when compared to histological fiber tracing, EM microscopy,

or stimulation. The most obvious of these is insensitivity to whether underlying axons are

anterograde or retrograde, as evidenced by the symmetry of the connectivity matrix. The

anisotropic diffusion of water molecules occurs in both anterograde and retrograde directions.

Thus, the true one-way connectivity between two areas could be anywhere between none to all

of the symmetric diffusion connectivity. Another important limitation is spatial resolution. While

the 1.25 mm isotropic voxels achieved by the WU-Minn dMRI protocol are smaller than those

of most studies (Jeurissen et al., 2019), they are still more than three orders-of-magnitude

larger than the typical submicron axon diameter (Liewald et al., 2014; von Keyserlingk Graf and

Schramm, 1984). This discrepancy is particularly impactful when fiber orientations are not

consistent within a voxel, i.e. crossing fibers. Probabilistic diffusion tractography (Behrens et

al., 2007) partially ameliorates the issue by modeling the probability distribution of orientations

and accounting for uncertainty, but ultimately dMRI with current technology is a meso- to

macroscale technique. Direct histological validation of dMRI techniques is uncommon, but has

been performed for probabilistic tractography *in vitro* in pigs (Dyrby et al., 2007) and macaques

(Donahue et al., 2016; Jbabdi et al., 2013), with the latter two studies using the same

probtrackX algorithm as the current study (Behrens et al., 2007). We have extending these

validations with a between-species comparison (Fig. **4**).

Of the several families of dMRI tractography algorithms available, we selected local,

probabilistic tractography (Behrens et al., 2007). The WU-Minn HCP makes available the

bedpostX precursor files and creating a probabilistic tractography connectome was always a

stated component of the WU-Minn HCP project (Van Essen et al., 2013; Van Essen and

Ugurbil, 2017). That such a connectome has not yet been released for these data may be due

to the immense computational challenge of performing these analyses at the scale of the HCP.

An advantage of probabilistic tractography is its sensitivity to minor, or low-probability

connections. Deterministic dMRI tractography connectomes typically have low network

densities, e.g. 0.18 (Mori et al., 2008) or 0.23 (Cui et al., 2019), when compared to histological

fiber tracing in macaques, 0.66 (Markov et al., 2014), and this is likely a lower bound as such

tracing is subject to false-negatives due to imperfect dye uptake and incomplete cortical

sampling. This suggests the deterministic dMRI connectomes are missing weaker connections.

On the other hand, dMRI in general and probabilistic tractography in particular has been found

vulnerable to false-positive connections (Maier-Hein et al., 2017). This exchange of specificity

for sensitivity (Sarwar et al., 2019; Zalesky et al., 2016) is consistent with our very high group-

average network density of 1.0 and the likely presence false-positive connections, and is thus

an important caveat to the data presented here. In cases where false-negative connections are

less concerning than false-positive connections, such as topological analyses (Zalesky et al.,

2016), subsequent users of these data may opt to threshold the connectivity matrix by either

connection strength or consistency (Roberts et al., 2017), see figure **3**.

When constructing this connectome, we divided the cortex into 180 parcels per

hemisphere following the HCP-MMP1.0 atlas (Glasser et al., 2016). To ease interpretation, we

further organized the parcels into 10 functional networks modified from (Ji et al., 2019). These

networks were created by applying iterative Louvain clustering (Blondel et al., 2008; Rubinov

and Sporns, 2010) and other criteria to HCP resting state fMRI data. While these fMRI-defined

network definitions correspond reasonably to the structural connections reported here, there

are exceptions. The operculum and temporoparietal junction, in particular, appears to be a

structurally distinct area that has been folded into several functional networks (Ji et al., 2019).

However, this contiguous region forms the lateral salience network in (Barnett et al., 2020)

which similarly applied a very similar methodology to a non-HCP cohort. Like many cortically-

focused studies, we used a surface-based methodology to define these areas, with seed and

target regions constrained to the white-matter – gray-matter interface. This approach reduces

the overrepresentation of major bundles (Jeurissen et al., 2019), enables the automated

assessment based on inter-subject homology (Fischl et al., 1999), facilitates comparison to

other cortical datasets, and is true to the anatomical nature of the cortical ribbon.

Unfortunately, the subcortex and cerebellum are omitted in this analysis, as are short-range,

often unmyelinated, intra-parcel connections. While the inclusion of the thalamic radiations, in

particular, is a merited future extension of this connectome, the small size of subcortical

structures relative to diffusion imaging voxels, the nuclear (as opposed the sheet-like)

organization of subcortical structures, and complex geometry of the subcortical white matter

— gray matter interface (e.g. the internal medullary lamina of the thalamus), all render the

challenges and methods for obtaining subcortical tractography substantially distinct from those

of cortico-cortico tractography.

The HCP-MMP1.0 atlas used was selected because of its wide adoption, symmetry, and

high parcel count. Furthermore, the parcels are based on multiple functional and anatomical

criteria and are consistent with previous functional parcellations in human and non-human

primates (Felleman and Van Essen, 1991; Glasser et al., 2016). Because the parcels are

relatively small and informed by function, erroneous averaging of disparate connections, a

connectomic extension of the partial volume artifact, is minimized. However, this comes at the

cost of non-uniformity in both parcel area and shape. Methodologically, parcels are assembled

from vertices on the tessellated cortical surface. A future vertex- or voxel-based connectome,

while computationally challenging, would have the distinct advantage of being readily

reformulated into any arbitrary surface-based parcellation scheme.

We found that pairwise connectivity between cortical parcels exhibits an exponential decay

rule with respect to fiber tract distance with a length constant λ of ~23 mm (~33 mm for

callosal connections). While a tight exponential relationship between probabilistic diffusion

tractography strength and fiber length has been previously reported, (Roberts et al., 2016), this

study did not report the observed λ or release its data. Histological studies in non-human

primates (Donahue et al., 2016; Markov et al., 2013; Theodoni et al., 2020) consistently show

exponential connectivity decay with distance. Such a rule when combined with a roughly

Gaussian distribution of interareal distances explains the observed log-normal distribution of

connectivity strength (Markov et al., 2013). Histological data indicate a λ of about 3.33 mm for

marmosets (Theodoni et al., 2020) and 5.55 mm for macaques (Markov et al., 2013). Across

species, there appears to be a linear relationship between the logs of λ and total gray matter

volume, predicting a human λ of 10 mm (Theodoni et al., 2020). While methodological

differences between diffusion and histological tractography cannot be completely ruled out,

Donahue and colleagues found similar λ for the two methods in macaques (Donahue et al.,

2016). Our results suggest that, compared to other species, human cortical areas are

exceptionally well connected relative to their cortical volume, reflected in a disproportionately

long λ. Conservatively restricting the exponential fit to only the most consistent quintile of

connections (Fig. **3D**) yields a λ of ~28 mm, further accentuating the proportional long-range

hyperconnectivity of humans.

Geometric scaling strongly constrains cortico-cortical connectivity in humans. Considering

primate brains increasing in diameter d, volume and number of cortical neurons increases by

d^3,(Ventura-Antunes et al., 2013), so arriving at a constant probability of connection between

any two neurons would require d^6 axons, and since they would need to be about d times as

long, this would require a volume proportional to d^7, or more if axonal diameter is increased to

maintain a relatively constant latency of communication (Wang et al., 2008). However, the

actual white matter volume is less than d^4 (Zhang and Sejnowski, 2000), and consequently the

probability of cortico-cortical connectivity must be highly limited in humans. The relatively long

λ in humans we report reduces even further the number of connections which can be

accommodated within the available white matter volume. A consequence of fewer but longer

connections would be reduced metabolic cost, inasmuch the cost of an action potential is 1/3

axonal transmission (proportional to length) and 2/3 synaptic transmission (Lennie, 2003). The

low firing rate of human pyramidal cells (Chan et al., 2014) would also reduce the metabolic

cost of their axons. These observations are consistent with the proposal that the metabolic

costs of cortico-cortical connections may help constrain their organization in the primate brain

(Ercsey-Ravasz et al., 2013). Given this strong correlation of connection strength with distance,

as well as the bias of tract-tracing techniques toward shorter, less geometrically complex

connections (Jeurissen et al., 2019), there may be some merit in regressing out the effect of

tract length when evaluating the relative connectivity of different cortical areas. However, the

considerations enumerated above imply a strong evolutionary selection to place cortical

parcels which require high connectivity to perform their calculations to be situated in direct

physical proximity to each other. The patterns of relatively long distance connectivity identified

here thus must be viewed as minor deviations from an overall strong tendency favoring local

connectivity, a conceptualization consistent with the view of the cortex as a spatially

embedded small world network.

One striking deviation from the distance-based connectivity was the left-lateralized hyper-

connectivity between language areas, and specifically between posterior and anterior language

areas. This connectivity presumably passes, completely or in part, through the classical

language pathways (reviewed in (Dick and Tremblay, 2012)). The lateralization we observed

may then reflect that of the arcuate and inferior longitudinal fasciculi which connect the same

structures and show significant left lateralization in humans but not macaques (Eichert et al.,

2019; Panesar et al., 2018). Left-lateralization of the arcuate fasciculus develops late (Lebel and

Beaulieu, 2011), and is sensitive to the presence, quality and quantity of early language

experience (Cheng et al., 2019; Romeo et al., 2018). More generally, many of the connectivity

patterns observed here could be the indirect result of co-activation of the connected parcels

(Mount and Monje, 2017). The left-lateralized ipsilateral connectivity may be compensated by a

relative lack of callosal connections from the same areas, under the hypothesis that the total

connectivity is constrained.

A more general factor that might induce deviations from a distance-based connectivity rule

may be the principle of hierarchical organization. It has been proposed that distant areas with

similar laminar properties, and thus of similar hierarchical order may have privileged

connections (Barbas, 2015). Across the entire cortex we find that myelination similarity explains

a significant but small amount of the overall variance. However, there are regions where the

influence of hierarchical position is more pronounced including the right dorsal attention and

left auditory/language networks. The observed hyperconnectivity and high degree of

lateralization in these regions may be a consequence of the low-latencies necessary for the

functions they underly. More broadly, the effects of transmission latency constraints on

neuroanatomy and conduction delay on large-scale physiological recordings are an emerging

area of study in human neuroscience (Muller et al., 2018). Latency is a hybrid structural–

functional property of connectivity, and might in future be quantified using the latency of

cortico-cortical evoked potentials (CCEP).

By emphasizing the cortical connectivity matrix over the white matter bundles *per se* and

organizing the matrix into the widely adopted HCP-MMP1.0 atlas (Glasser et al., 2016), the

structural connectome reported here enables ready comparison to other structural, functional,

and hybrid connectomes. As an example, we compared the probabilistic tractography

connectivity to exist resting-state fMRI (rs-fMRI) (Van Essen et al., 2013) and CCEP (Trebaul et

al., 2018) connectivity matrices and found that our dMRI-inferred structural connectivity better

reflects CCEP probability than rs-fMRI connectivity in both linear and network-theoretic

comparisons, despite the dMRI and rs-fMRI cohorts being highly overlapping. Although

resting-state functional connectivity is constrained by anatomical networks and can be partially

predicted by them (Honey et al., 2009), indirect connections or parallel processing of stimuli in

different areas can produce correlated activity even in the absence of direct anatomical

connections. One notable example of the latter may be inter-hemispheric connectivity. While

we did find hyperconnectivity between inter-hemispheric homologs when compared to other

callosal connections, anatomical interhemispheric connectivity on the whole is much weaker

than found in rs-fMRI. CCEPs, being directed by clinical requirements, have poor inter-

hemispheric sampling, but we found that even among ipsilateral connections, rs-fMRI is still

less similar to CCEP than probabilistic tractography. These inter-modal connectivity

comparisons are not intended to be comprehensive. The HCP cohort also includes source-

localized resting-state magneto-encephalography (MEG) (Larson-Prior et al., 2013), which

could be used to examine the degree to which the functional connectivity of various frequency

bands corresponds to anatomical connectivity. Furthermore, neuropsychological metrics,

including the NIH toolbox (Gershon et al., 2013), and genotypic data (dbGaP phs001364.v1.p1)

are also available for this cohort, enabling future studies of the interplay between cortical

connectivity, cognition, and genetics.

The Human Connectome Project was a scientific undertaking of visionary scope and

ambition. Its commitment to open science and accessibility of data by the public enabled this

study and will continue to facilitate further studies for years to come. Emerging clinical

applications of brain connectomics will be underpinned by a strong base of normative data for

comparison. The whole-cortex probabilistic diffusion tractography connectome reported here

fulfills a key goal outlined in the project's conception and we hope it will empower yet further

study of the myriad and beautiful web of connectivity that the human brain embodies.

____

Data availability

Individual and group average connectivity matrices as well as all other figure source data can

be found at https://doi.org/10.5281/zenodo.4060485. The preprocessed HCP data using in this

study was retrieved from https://db.humanconnectome.org and the preprocessing code used

to create these files is availed at https://github.com/Washington-University/HCPpipelines. The

source code for FSL, including probtrackx2 is available from

https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FSL. Network theory measures were computed with the

brain connectivity Matlab toolbox whose source code is available from http://www.brain-

connectivity-toolbox.net.

Acknowledgements

This work was supported by NIMH 1RF1MH117155-01 and 5T32MH020002-19, NINDS

1R01NS109553-01A1, and ONR 00014-16-1-2415. Data were provided, in part, by the Human

Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and

Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the

NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems

Neuroscience at Washington University. Data were provided, in part, by the Functional Brain

Tractography Project, funded by the European Research council. We thank Adam Niese and

the San Diego Supercomputer Center for providing technical support.

References

Andersson JLR, Sotiropoulos SN. 2016. An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. *Neuroimage*. doi:10.1016/j.neuroimage.2015.10.019

Barbas H. 2015. General Cortical and Special Prefrontal Connections: Principles from Structure to Function. *Annu Rev Neurosci* **38**:269–289. doi:10.1146/annurev-neuro-071714-033936

Barnett AJ, Reilly W, Dimsdale-zucker H, Mizrak E, Reagh Z, Barnett AJ. 2020. Organization of cortico-hippocampal networks in the human brain 1–30.

Behrens TEJ, Berg HJ, Jbabdi S, Rushworth MFS, Woolrich MW. 2007. Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? *Neuroimage* **34**:144–155. doi:10.1016/j.neuroimage.2006.09.018

Blondel VD, Guillaume JL, Lambiotte R, Lefebvre E. 2008. Fast unfolding of communities in large networks. *J Stat Mech Theory Exp*. doi:10.1088/1742-5468/2008/10/P10008

Burns S. 2014. A Recipe for Cortical Tractography Using Freesufer Labels. http://sburns.org/2014/05/03/cortical-tractography-recipe.html

Burt JB, Demirtaş M, Eckner WJ, Navejar NM, Ji JL, Martin WJ, Bernacchia A, Anticevic A, Murray JD. 2018. Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. *Nat Neurosci* **21**. doi:10.1038/s41593-018-0195-0

Chan AM, Dykstra AR, Jayaram V, Leonard MK, Travis KE, Gygi B, Baker JM, Eskandar E, Hochberg LR, Halgren E, Cash SS. 2014. Speech-specific tuning of neurons in human superior temporal gyrus. *Cereb Cortex* **24**:2679–2693. doi:10.1093/cercor/bht127

Cheng Q, Roth A, Halgren E, Mayberry RI. 2019. Effects of Early Language Deprivation on Brain Connectivity: Language Pathways in Deaf Native and Late First-Language Learners of American Sign Language. *Front Hum Neurosci* **13**:1–12. doi:10.3389/fnhum.2019.00320

Coalson T, Van Essen D, Glasser M. 2016. hcp-users FAQ #9: How do I map data between FreeSurfer and HCP?

Cui LB, Wei Y, Xi Y Bin, Griffa A, De Lange SC, Kahn RS, Yin H, Van Den Heuvel MP. 2019. Connectome-Based Patterns of First-Episode Medication-Naïve Patients with Schizophrenia. *Schizophr Bull* **45**:1291–1299. doi:10.1093/schbul/sbz014

David O, Job AS, De Palma L, Hoffmann D, Minotti L, Kahane P. 2013. Probabilistic functional tractography of the human cortex. *Neuroimage* **80**:307–317. doi:10.1016/j.neuroimage.2013.05.075

Dayan P, Abbott LF. 2001. Theoretical neuroscience: computational and mathematical modeling of neural systems.

Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM, Maguire RP, Hyman BT, Albert MS, Killiany RJ. 2006. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* **31**:968–980. doi:10.1016/j.neuroimage.2006.01.021

Dick AS, Tremblay P. 2012. Beyond the arcuate fasciculus: Consensus and controversy in the connectional anatomy of language. *Brain*. doi:10.1093/brain/aws222

Donahue CJ, Sotiropoulos SN, Jbabdi S, Hernandez-Fernandez M, Behrens TE, Dyrby TB, Coalson T, Kennedy H, Knoblauch K, Van Essen DC, Glasser MF. 2016. Using Diffusion Tractography to Predict Cortical Connection Strength and Distance: A Quantitative Comparison with Tracers in the Monkey. *J Neurosci* **36**:6758–6770. doi:10.1523/JNEUROSCI.0493-16.2016

Dyrby TB, Søgaard L V., Parker GJ, Alexander DC, Lind NM, Baaré WFC, Hay-Schmidt A,
Eriksen N, Pakkenberg B, Paulson OB, Jelsing J. 2007. Validation of in vitro probabilistic
tractography. *Neuroimage* **37**:1267–1277. doi:10.1016/j.neuroimage.2007.06.022
Eichert N, Verhagen L, Folloni D, Jbabdi S, Khrapitchev AA, Sibson NR, Mantini D, Sallet J,
Mars RB. 2019. What is special about the human arcuate fasciculus? Lateralization,
projections, and expansion. *Cortex*. doi:10.1016/j.cortex.2018.05.005
Ercsey-Ravasz M, Markov NT, Lamy C, VanEssen DC, Knoblauch K, Toroczkai Z, Kennedy H.
2013. A Predictive Network Model of Cerebral Cortical Connectivity Based on a Distance
Rule. *Neuron* **80**:184–197. doi:10.1016/j.neuron.2013.07.036
Felleman DJ, Van Essen DC. 1991. Distributed Hierarchical Processing in the Primate Cerebral
Cortex. *Cereb Cortex* **1**:1–47. doi:10.1093/cercor/1.1.1
Fischl B. 2012. FreeSurfer. *Neuroimage*. doi:10.1016/j.neuroimage.2012.01.021
Fischl B, Sereno MI, Tootell RBH, Dale AM. 1999. High-resolution intersubject averaging and a
coordinate system for the cortical surface. *Hum Brain Mapp* **8**:272–284.
doi:10.1002/(SICI)1097-0193(1999)8:4<272::AID-HBM10>3.0.CO;2-4
Fischl B, Van Der Kouwe A, Destrieux C, Halgren E, Ségonne F, Salat DH, Busa E, Seidman LJ,
Goldstein J, Kennedy D, Caviness V, Makris N, Rosen B, Dale AM. 2004. Automatically
Parcellating the Human Cerebral Cortex. *Cereb Cortex* **14**:11–22.
doi:10.1093/cercor/bhg087
Gershon RC, Wagster M V, Hendrie HC, Fox NA, Cook KF, Nowinski CJ. 2013. NIH toolbox for
assessment of neurological and behavioral function. *Neurology* **80**:S2–S6.
Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K, Andersson
J, Beckmann CF, Jenkinson M, Smith SM, Van Essen DC. 2016. A multi-modal
parcellation of human cerebral cortex. *Nature*. doi:10.1038/nature18933
Glasser MF, Van Essen DC. 2011. Mapping Human Cortical Areas In Vivo Based on Myelin
Content as Revealed by T1- and T2-Weighted MRI. *J Neurosci* **31**:11597–11616.
doi:10.1523/JNEUROSCI.2180-11.2011
Hagmann P, Cammoun L, Gigandet X, Meuli R, Honey CJ, Van Wedeen J, Sporns O. 2008.
Mapping the structural core of human cerebral cortex. *PLoS Biol* **6**:1479–1493.
doi:10.1371/journal.pbio.0060159
Honey CJ, Sporns O, Cammoun L, Gigandet X, Thiran JP, Meuli R, Hagmann P. 2009.
Predicting human resting-state functional connectivity from structural connectivity. *Proc
Natl Acad Sci U S A* **106**:2035–2040. doi:10.1073/pnas.0811168106
Humphries MD, Gurney K. 2008. Network "small-world-ness": A quantitative method for
determining canonical network equivalence. *PLoS One*. doi:10.1371/journal.pone.0002051
Jbabdi S, Lehman JF, Haber SN, Behrens TE. 2013. Human and monkey ventral prefrontal
fibers use the same organizational principles to reach their targets: Tracing versus
tractography. *J Neurosci* **33**:3190–3201. doi:10.1523/JNEUROSCI.2457-12.2013
Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. 2012. FSL. *Neuroimage*
62:782–790. doi:10.1016/j.neuroimage.2011.09.015
Jeurissen B, Descoteaux M, Mori S, Leemans A. 2019. Diffusion MRI fiber tractography of the
brain. *NMR Biomed* **32**:1–22. doi:10.1002/nbm.3785
Ji JL, Spronk M, Kulkarni K, Repovš G, Anticevic A, Cole MW. 2019. Mapping the human
brain's cortical-subcortical functional network organization. *Neuroimage* **185**:35–57.
doi:10.1016/j.neuroimage.2018.10.006
Larson-Prior LJ, Oostenveld R, Della Penna S, Michalareas G, Prior F, Babajani-Feremi A,
Schoffelen JM, Marzetti L, de Pasquale F, Di Pompeo F, Stout J, Woolrich M, Luo Q,
Bucholz R, Fries P, Pizzella V, Romani GL, Corbetta M, Snyder AZ. 2013. Adding
dynamics to the Human Connectome Project with MEG. *Neuroimage* **80**:190–201.

doi:10.1016/j.neuroimage.2013.05.056

Latora V, Marchiori M. 2001. Efficient behavior of small-world networks. *Phys Rev Lett*.
doi:10.1103/PhysRevLett.87.198701

Lebel C, Beaulieu C. 2011. Longitudinal development of human brain wiring continues from
childhood into adulthood. *J Neurosci* **31**:10937–10947.

Lennie P. 2003. The cost of cortical computation. *Curr Biol* **13**:493–497.

Liewald D, Miller R, Logothetis N, Wagner HJ, Schüz A. 2014. Distribution of axon diameters in
cortical white matter: an electron-microscopic study on three human brains and a
macaque. *Biol Cybern* **108**:541–557. doi:10.1007/s00422-014-0626-2

Maier-Hein KH, Neher PF, Houde JC, Côté MA, Garyfallidis E, Zhong J, Chamberland M, Yeh
FC, Lin YC, Ji Q, Reddick WE, Glass JO, Chen DQ, Feng Y, Gao C, Wu Y, Ma J, Renjie H,
Li Q, Westin CF, Deslauriers-Gauthier S, González JOO, Paquette M, St-Jean S, Girard G,
Rheault F, Sidhu J, Tax CMW, Guo F, Mesri HY, Dávid S, Froeling M, Heemskerk AM,
Leemans A, Boré A, Pinsard B, Bedetti C, Desrosiers M, Brambati S, Doyon J, Sarica A,
Vasta R, Cerasa A, Quattrone A, Yeatman J, Khan AR, Hodges W, Alexander S,
Romascano D, Barakovic M, Auría A, Esteban O, Lemkaddem A, Thiran JP, Cetingul HE,
Odry BL, Mailhe B, Nadar MS, Pizzagalli F, Prasad G, Villalon-Reina JE, Galvis J,
Thompson PM, Requejo FDS, Laguna PL, Lacerda LM, Barrett R, Dell'Acqua F, Catani M,
Petit L, Caruyer E, Daducci A, Dyrby TB, Holland-Letz T, Hilgetag CC, Stieltjes B,
Descoteaux M. 2017. The challenge of mapping the human connectome based on
diffusion tractography. *Nat Commun* **8**. doi:10.1038/s41467-017-01285-x

Marcus DS, Harwell J, Olsen T, Hodge M, Glasser MF, Prior F, Jenkinson M, Laumann T,
Curtiss SW, Van Essen DC. 2011. Informatics and data mining tools and strategies for the
human connectome project. *Front Neuroinform* **5**:1–12. doi:10.3389/fninf.2011.00004

Markov, Ercsey-Ravasz MM, Ribeiro Gomes AR, Lamy C, Magrou L, Vezoli J, Misery P,
Falchier A, Quilodran R, Gariel MA. 2014. A weighted and directed interareal connectivity
matrix for macaque cerebral cortex. *Cereb cortex* **24**:17–36.

Markov NT, Ercsey-Ravasz M, Van Essen DC, Knoblauch K, Toroczkai Z, Kennedy H. 2013.
Cortical High-Density Counterstream Architectures. *Science (80-)* **342**:1238406–1238406.
doi:10.1126/science.1238406

Mori S, Oishi K, Jiang H, Jiang L, Li X, Akhter K, Hua K, Faria A V., Mahmood A, Woods R,
Toga AW, Pike GB, Neto PR, Evans A, Zhang J, Huang H, Miller MI, van Zijl P, Mazziotta
J. 2008. Stereotaxic white matter atlas based on diffusion tensor imaging in an ICBM
template. *Neuroimage* **40**:570–582. doi:10.1016/j.neuroimage.2007.12.035

Mount CW, Monje M. 2017. Wrapped to adapt: experience-dependent myelination. *Neuron*
95:743–756.

Muller L, Chavane F, Reynolds J, Sejnowski TJ. 2018. Cortical travelling waves: mechanisms
and computational principles. *Nat Rev Neurosci*. doi:10.1038/nrn.2018.20

Newman MEJ. 2004. Fast algorithm for detecting community structure in networks. *Phys Rev E*
69:66133.

Newman MEJ. 2003. The structure and function of complex networks. *SIAM Rev*.
doi:10.1137/S003614450342480

Panesar SS, Yeh F-C, Jacquesson T, Hula W, Fernandez-Miranda JC. 2018. A quantitative
tractography study into the connectivity, segmentation and laterality of the human inferior
longitudinal fasciculus. *Front Neuroanat* **12**:47.

Roberts JA, Perry A, Lord AR, Roberts G, Mitchell PB, Smith RE, Calamante F, Breakspear M.
2016. The contribution of geometry to the human connectome. *Neuroimage* **124**:379–393.
doi:10.1016/j.neuroimage.2015.09.009

Roberts JA, Perry A, Roberts G, Mitchell PB, Breakspear M. 2017. Consistency-based

thresholding of the human connectome. *Neuroimage* **145**:118–129.
doi:10.1016/j.neuroimage.2016.09.053

Romeo RR, Segaran J, Leonard JA, Robinson ST, West MR, Mackey AP, Yendiki A, Rowe ML,
Gabrieli JDE. 2018. Language exposure relates to structural neural connectivity in
childhood. *J Neurosci* **38**:7870–7877.

Rubinov M, Sporns O. 2010. Complex network measures of brain connectivity: Uses and
interpretations. *Neuroimage*. doi:10.1016/j.neuroimage.2009.10.003

Sarwar T, Ramamohanarao K, Zalesky A. 2019. Mapping connectomes with diffusion MRI:
deterministic or probabilistic tractography? *Magn Reson Med* **81**:1368–1384.
doi:10.1002/mrm.27471

Sotiropoulos SN, Jbabdi S, Xu J, Andersson JL, Moeller S, Auerbach EJ, Glasser MF,
Hernandez M, Sapiro G, Jenkinson M, Feinberg DA, Yacoub E, Lenglet C, Van Essen DC,
Ugurbil K, Behrens TEJ. 2013. Advances in diffusion MRI acquisition and processing in
the Human Connectome Project. *Neuroimage* **80**:125–143.
doi:10.1016/j.neuroimage.2013.05.057

Sporns O, Tononi G, Kötter R. 2005. The human connectome: A structural description of the
human brain. *PLoS Comput Biol* **1**:0245–0251. doi:10.1371/journal.pcbi.0010042

Theodoni P, Majka P, Reser DH, Wójcik DK, Rosa MGP, Wang X-J. 2020. Structural attributes
and principles of the neocortical connectome in the marmoset monkey. *bioRxiv*.

Trebaul L, Deman P, Tuyisenge V, Jedynak M, Hugues E, Rudrauf D, Bhattacharjee M, Tadel F,
Chanteloup-Foret B, Saubat C, Reyes Mejia GC, Adam C, Nica A, Pail M, Dubeau F,
Rheims S, Trébuchon A, Wang H, Liu S, Blauwblomme T, Garcés M, De Palma L, Valentin
A, Metsähonkala EL, Petrescu AM, Landré E, Szurhaj W, Hirsch E, Valton L, Rocamora R,
Schulze-Bonhage A, Mindruta I, Francione S, Maillard L, Taussig D, Kahane P, David O.
2018. Probabilistic functional tractography of the human cortex revisited. *Neuroimage*
181:414–429. doi:10.1016/j.neuroimage.2018.07.039

Van Essen DC, Smith SM, Barch DM, Behrens TEJ, Yacoub E, Ugurbil K. 2013. The WU-Minn
Human Connectome Project: An overview. *Neuroimage* **80**:62–79.
doi:10.1016/j.neuroimage.2013.05.041

Van Essen DC, Ugurbil K. 2017. Components of the Human Connectome Project - Diffusion
Tractography. https://www.humanconnectome.org/study/hcp-young-adult/project-
protocol/diffusion-tractography

Ventura-Antunes L, Mota B, Herculano-Houzel S. 2013. Different scaling of white matter
volume, cortical connectivity, and gyrification across rodent and primate brains. *Front
Neuroanat* **7**:1–12. doi:10.3389/fnana.2013.00003

von Keyserlingk Graf D, Schramm U. 1984. Diameter of axons and thickness of myelin sheaths
of the pyramidal tract fibres in the adult human medullary pyramid. *Anat Anz* **157**:97–111.

Wang SSH, Shultz JR, Burish MJ, Harrison KH, Hof PR, Towns LC, Wagers MW, Wyatt KD.
2008. Functional trade-offs in white matter axonal scaling. *J Neurosci* **28**:4047–4056.
doi:10.1523/JNEUROSCI.5559-05.2008

Watts DJ, Strogatz SH. 1998. Collective dynamics of 'small-world'networks. *Nature* **393**:440.

Yeh FC, Panesar S, Fernandes D, Meola A, Yoshino M, Fernandez-Miranda JC, Vettel JM,
Verstynen T. 2018. Population-averaged atlas of the macroscale human structural
connectome and its network topology. *Neuroimage* **178**:57–68.
doi:10.1016/j.neuroimage.2018.05.027

Yeh FC, Verstynen TD, Wang Y, Fernández-Miranda JC, Tseng WYI. 2013. Deterministic
diffusion fiber tracking improved by quantitative anisotropy. *PLoS One* **8**:1–16.
doi:10.1371/journal.pone.0080713

Yendiki A, Panneck P, Srinivasan P, Stevens A, Zöllei L, Augustinack J, Wang R, Salat D,

Ehrlich S, Behrens T, Jbabdi S, Gollub R, Fischl B. 2011. Automated probabilistic
947   reconstruction of white-matter pathways in health and disease using an atlas of the
948   underlying anatomy. *Front Neuroinform* **5**:1–12. doi:10.3389/fninf.2011.00023
949   Yoon JM, Benson NC, Forenzo D, Winawer J, Engel SA, Kay KN. 2019. Heritability of V1/V2/V3
950   surface area in the HCP 7T Retinotopy Dataset. *J Vis* **19**:41b-41b.
951   Zalesky A, Fornito A, Cocchi L, Gollo LL, van den Heuvel MP, Breakspear M. 2016.
952   Connectome sensitivity or specificity: which is more important? *Neuroimage* **142**:407–420.
953   doi:10.1016/j.neuroimage.2016.06.035
954   Zhang K, Sejnowski TJ. 2000. A universal scaling law between gray matter and white matter of
955   cerebral cortex. *Proc Natl Acad Sci U S A* **97**:5621–5626. doi:10.1073/pnas.090504197
956
957

Legends

Figure 1. Probabilistic diffusion tractography structural connectome of the human cortex (**A**)

Group average (N = 1065) structural connectivity matrix consisting of the 360 HCP-MMPS1.0

atlas parcels organized into ten functional networks. Raw streamline counts are fractionally

scaled yielding the log probability F_{pt}. The white arrows highlight the diagonal which contains

contralateral homologs. (**B**) The first row of the connectivity matrix, showing connection

probabilities from left V1 to all other parcels, projected onto the fsaverage template cortex. (**C**)

Single subject (100307) volume ray casting visualization of left V1-originating streamline

probabilities within the skull-stripped T1-weighted structural MR volume. (**D**) Ten functional

networks, adapted from (Ji et al., 2019), within HCP-MMPS1.0 atlas. These are indicated by

red boxes in panel **A**.

Figure 1-1. Comparison of normalization methods. Shown are the (**A**) connectivity matrices,

(**B**) distributions of pairwise connectivity, (**C**) the pre-log distribution of F_{pt} (**D**) relationships

between connectivity and fiber tract length for four normalization methods.

Figure 2. . Connectivity strength exponential decays with fiber tract length. (**A**) and (**B**)

connections within the right and left hemispheres, respectively. (**C**) Connections between the

right and left hemisphere. (**D**) All connections. Each marker represents a pair of parcels. Red

traces show the least-squares exponential fit; inset are the length constant λ and r^2 of this fit.

Note that F_{pt} is log-transformed making these axes effectively semi-log.

Figure 2-1. Alternative models for fitting connectivity strength as a function of fiber tract

length. Each gray marker shows the average pair-wise Fpt between two parcels and fiber tract

length between them, as also shown in figure 2D. The colored traces show maximum likelihood

estimates for several listed functional forms. The AIC, AICc, aBIC columns contain the Akaike,

corrected Akaike, and Bayesian information criteria, respectively. While the Gaussian fits

explain slightly more variance and have a slightly lower AIC than the exponential fit, the

exponential has fewer parameters and is consistent with histological non-human primate

evidence (Donahue et al., 2016; Markov et al., 2013; Theodoni et al., 2020).

Figure 2-2. Effect of motion during the dMRI scan. (**A**) Time-course of displacement relative to

initial position for one subject (996782). The six runs of the HCP dMRI protocol can be seen.

(**B**) Exponential fall-off coefficient λ is only modestly affected by motion, r = 0.140, p = 4.6E-6.

Each marker represents a subject.

Figure 3. Inter-individual variability. Shown are (**A**) the matrix of connectivity coefficients of

variation (CV) across subjects (**B**) pairwise CV vs. fiber tract length, (**C**) the distribution of CV

across all connections, (**D**) the F_{pt} vs. fiber tract length for the connections in the highest

quintile of inter-individual consistency, and (**F**) the F_{pt} of right hemisphere V1 – V2 connection in

all subjects vs. left hemisphere V1 – V2 connection. In panels **B** and **D** each marker represents

a sample statistic for a connection between two parcels. In panel **F** each marker represents an

individual subject. In panel **D** the red trace show the least-squares exponential fit and inset are

the length constant λ and r^2 of this fit. Note that F_{pt} is log-transformed making this panel's axes

effectively semi-log. In panel F, the r^2 of the least squares linear fit is reported.

1003

1004   **Figure 4.** Comparison of human diffusion tractography and macaque retrograde tracing

1005   connectomes. Subset of homologous parcels in the human HCP-MMPS1.0 and macaque fv91

1006   atlas. (**A**) Macaque group-average retrograde tracer derived structural connectome, gray

1007   indicates missing data. (**B**) Human probabilistic diffusion tractography connectome. (**C**)

1008   Pairwise correlation between macaque and human structural connectivity, r = 0.35, p = 0.0013.

1009

1010   **Figure 5.** Interhemispheric connectivity. Differential connectivity between ipsilateral and

1011   contralateral connectivity. Greater Ipsilateral connectivity dominates and is indicated in red.

1012   Parcel-pairs with greater *contra*lateral connectivity than ipsilateral are blue. The green cortical

1013   patches show anatomic extent of parcel groups of notable contrast.

1014

1015   **Figure 6.** Contralateral homologs. Differential connectivity between contralateral homologous

1016   parcels vs the mean of all other contralateral parcels. Red indicates contralateral homologous

1017   connectivity greater than mean contralateral connectivity. Note that many language-implicated

1018   regions have relatively weak connectivity with their contralateral homologs.

1019

1020   **Figure 6-1.** Differential connectivity between contralateral homologous parcels vs the mean of

1021   all other contralateral parcels. Confidence intervals are Bonferroni-corrected for multiple

1022   comparisons.

1023

Figure 7. Language/Auditory network hyperconnectivity and left-lateralization. (**A**) Distance-

binned connectivity within the language and auditory networks compared to connectivity

between the language and auditory networks and other networks, separately for the left and

right hemispheres (**B**) The differential trace for the within- and between- connectivity in both

hemispheres. In both panels, gray patches show Bonferroni-corrected bootstrapped 95%

confidence intervals across subjects.

1030

Figure 8. Connectivity is influenced by the cortical hierarchy. (**A**, **B**) Connectivity is strongly

predicted by hierarchical similarity in some networks and modestly predicted overall. (**A**) All

connectivity vs. myelination difference, including within- and across- network connections, for

the left, right, and callosal connections. For both panels, each marker represents a parcel pair.

(**B**) Within-network connectivity vs. myelination difference for 10 functional networks. Linear fits

and correlation coefficients computed independently for the left and right hemisphere. A

negative correlation indicates that parcels at similar hierarchical levels tend to be more

connected. (**C**, **D**) Higher order prefrontal areas are better connected. (**C**) Histogram of

correlation coefficients between areal myelination and F_{pt} connectivity to each parcel. Only

significant coefficients after Bonferroni correction are shown. Most coefficients are negative

indicating high connectivity to low-myelination (i.e., higher-order) areas. (**D**) Significant negative

coefficients (red) map onto bilateral prefrontal cortex. Only the bilateral DVT and V6A are show

positive significant correlations (blue).

1044

1045   **Figure 8-1.** Myelination difference connectivity matrix. This provides an estimate for the

1046   difference in hierarchical level between cortical parcels. Values have been fractionally scaled.

1047 Note that the color scale has been reversed when compared to figure 1, as |Δmyelination| is

1048 inversely proportional to connectivity.

1049

1050 **Figure 8-2.** Pearson correlations between the F_{pt} from each left hemisphere parcel to all others

1051 and the target parcels' myelination indices. p values are Bonferroni-corrected for multiple

1052 comparisons.

1053

1054 **Figure 8-3.** Pearson correlations between the F_{pt} from each right hemisphere parcel to all

1055 others and the target parcels' myelination indices. p values are Bonferroni-corrected for

1056 multiple comparisons.

1057

Figure 9. Probabilistic dMRI more closely resembles CCEPs than resting-state fMRI. (**A**)

Connectivity matrices for probabilistic dMRI tractography, CCEP, and rs-fMRI. For CCEPs

missing data has been colored grey and pre-log zero-strength connections black. (**B**)

Correlations among the three modalities. The least-squares linear fit is shown in red. (**C**) Non-

zero pairwise connection strength distributions. Note that rs-fMRI connectivity values, which

are not log-transformed, display two modes, separated at 0.0014.(**D**) Cortical parcels

displaying lower (left) and higher (right) modes of rs-fMRI connectivity.

1065

1066 **Figure 9-1.** Within-hemisphere comparison of probabilistic dMRI tractography, CCEP, and rs-

1067 fMRI connectivity. For the left and right hemisphere, the distribution of pairwise non-zero

1068 connection strengths and correlations among the three modalities are shown. The least-

1069 squares linear fit is shown in red. All within-hemisphere findings are concordant with the overall

1070 findings, shown in figure **9**.

1071

1072   **Figure 10.** Network theoretic differences between the connectivity modalities. Binarized

1073   network metrics after thresholding by edge weight (connectivity strength).

1074   **Table 1.** Connectome Features

1075

1076   **Table 2.** Parcel order and network assignment. The emboldened indices refer to the parcel

1077   order in figure 1A. The Orig. indices refer to the original parcel order presented in (Glasser et

1078   al., 2016). All indices refer to the left hemisphere, adding 180 yields the homologous right

1079   hemisphere indices.

1080

Table 3. Statistics and uncertainty. Where multiple uncertainties are listed for a figure panel,

they correspond to the statistics read left-to-right, top-to-bottom in that panel. For figure 8B

only uncertainties for significant correlations are listed. Uncertainties for figures 6, 7, 8 and 10

are not shown. Figure 6-1 contains bootstrapped 95% confidence intervals for the 180 means

shown in figure 6, n = 179. Figure 7 shows Bootstrapped 95% confidence intervals in gray; the

values of these intervals for all distance bins are available in the figure source data at

https://doi.org/10.5281/zenodo.4060485. For figure 10 means across shuffled matrices are

only necessary to account for arbitrary ordering among tied edge weights and the

bootstrapped 95% confidence intervals for these means are vanishingly small. The values of

these intervals at all network densities are also included in the figure source data. For nonlinear

regressions confidence intervals are estimated using R^{-1}, the inverse R factor from QR

decomposition of the Jacobian, the degrees of freedom for error, and the root mean squared

error. For linear correlations the confidence intervals are based on an asymptotic normal

distribution of 0.5*log((1+r)/(1-r)), with an approximate variance equal to 1/(N-3). For descriptive

statistics, e.g. means, empirical 95% confidence intervals are estimated by bootstrapping with

2000 iterations.

1097

1098

1099

1100

1101

1102

Figure 1. Probabilistic diffusion tractography structural connectome of the human cortex (**A**) Group average (N = 1065) structural connectivity matrix consisting of the 360 HCP-MMPS1.0 atlas parcels organized into ten functional networks. Raw streamline counts are fractionally scaled yielding the log probability F_{pt}. The white arrows highlight the diagonal which contains contralateral homologs. (**B**) The first row of the connectivity matrix, showing connection probabilities from left V1 to all other parcels, projected onto the fsaverage template cortex. (**C**) Single subject (100307) volume ray casting visualization of left V1-originating streamline probabilities within the skull-stripped T1-weighted structural MR volume. (**D**) Ten functional networks, adapted from (Ji et al., 2019), within HCP-MMPS1.0 atlas. These are indicated by red boxes in panel **A**.

Figure 1-1. Comparison of normalization methods. Shown are the (**A**) connectivity matrices, (**B**) distributions of pairwise connectivity, (**C**) the pre-log distribution of F_{pt} (**D**) relationships between connectivity and fiber tract length for four normalization methods.

1106
1107
1108
1109

Figure 2. Connectivity strength exponential decays with fiber tract length. (**A**) and (**B**) connections within the right and left hemispheres, respectively. (**C**) Connections between the right and left hemisphere. (**D**) All connections. Each marker represents a pair of parcels. Red traces show the least-squares exponential fit; inset are the length constant λ and r^2 of this fit. Note that F_{pt} is log-transformed making these axes effectively semi-log.

Function	Equation	# parameters	r²	AIC	AICc	BIC
exponential	$\alpha^* \exp(-d/\lambda)$	2	0.837	-773853	-773857	-773826
power law	$c^* d^\alpha$	2	0.717	-736752	-736756	-736725
truncated power law	$\exp(-d/\lambda)^* d^\alpha$	2	0.818	-766810	-766814	-766783
Gaussian	$\alpha^* \exp(-(d-b)^2/c^2)$	3	0.840	-775050	-775055	-775013
2 term Gaussian	$\alpha_1^* \exp(-(d-b_1)^2/c_1^2) + \alpha_2^* \exp(-(d-b_2)^2/c_2^2)$	6	0.844	-776436	-776444	-776373

Figure 2-1. Alternative models for fitting connectivity strength as a function of fiber tract length. Each gray marker shows the average pair-wise Fpt between two parcels and fiber tract length between them, as also shown in figure 2D. The colored traces show maximum likelihood estimates for several listed functional forms. The AIC, AICc, aBIC columns contain the Akaike, corrected Akaike, and Bayesian information criteria, respectively. While the Gaussian fits explain slightly more variance and have a slightly lower AIC than the exponential fit, the exponential has fewer parameters and is consistent with histological non-human primate evidence (Donahue et al., 2016; Markov et al., 2013; Theodoni et al., 2020).

Figure 2-2. Effect of motion during the dMRI scan. (**A**) Time-course of displacement relative to initial position for one subject (996782). The six runs of the HCP dMRI protocol can be seen. (**B**) Exponential fall-off coefficient λ is only modestly affected by motion, r = 0.140, p = 4.6E-6. Each marker represents a subject.

Figure 3. Inter-individual variability. Shown are (**A**) the matrix of connectivity coefficients of variation (CV) across subjects (**B**) pairwise CV vs. fiber tract length, (**C**) the distribution of CV across all connections, (**D**) the F_{pt} vs. fiber tract length for the connections in the highest quintile of inter-individual consistency, and (**F**) the F_{pt} of right hemisphere V1 – V2 connection in all subjects vs. left hemisphere V1 – V2 connection. In panels **B** and **D** each marker represents a sample statistic for a connection between two parcels. In panel **F** each marker represents an individual subject. In panel **D** the red trace show the least-squares exponential fit and inset are the length constant λ and r² of this fit. Note that F_{pt} is log-transformed making this panel's axes effectively semi-log. In panel F, the r² of the least squares linear fit is

Figure 4. Comparison of human diffusion tractography and macaque retrograde tracing connectomes. Subset of homologous parcels in the human HCP-MMPS1.0 and macaque fv91 atlas. (**A**) Macaque group-average retrograde tracer derived structural connectome, gray indicates missing data. (**B**) Human probabilistic diffusion tractography connectome. (**C**) Pairwise correlation between macaque and human structural connectivity. R = 0.35 p = 0.0013

Figure 5. Interhemispheric connectivity. Differential connectivity between ipsilateral and contralateral connectivity. Greater Ipsilateral connectivity dominates and is indicated in red. Parcel-pairs with greater *contra*lateral connectivity than ipsilateral are blue. The green cortical patches show anatomic extent of parcel groups of notable contrast.

Figure 6. Contralateral homologs. Differential connectivity between contralateral homologous parcels vs the mean of all other contralateral parcels. Red indicates contralateral homologous connectivity greater than mean contralateral connectivity. Note that many language-implicated regions have relatively weak connectivity with their contralateral homologs.

Idx	Parcel	F_{pt} to contra-lateral homolog	Mean [$CI_{95\%}$] F_{pt} to non-homologous contra-lateral parcels	Idx	Parcel	F_{pt} to contra-lateral homolog	Mean [$CI_{95\%}$] F_{pt} to non-homologous contra-lateral parcels	Idx	Parcel	F_{pt} to contra-lateral homolog	Mean [$CI_{95\%}$] F_{pt} to non-homologous contra-lateral parcels
1	V1	-1.69	-3.72 [-3.88 -3.57]	61	46	-1.57	-4.15 [-4.39 -3.90]	121	IP1	-2.92	-4.14 [-4.31 -3.99]
2	ProS	-4.87	-4.71 [-4.81 -4.60]	62	9-46d	-1.66	-3.92 [-4.12 -3.70]	122	PFm	-3.63	-4.09 [-4.19 -3.99]
3	DVT	-2.49	-3.97 [-4.11 -3.83]	63	43	-1.38	-3.65 [-3.82 -3.46]	123	p10p	-4.75	-4.51 [-4.59 -4.43]
4	MST	-1.98	-3.86 [-4.01 -3.73]	64	PFcm	-1.70	-4.23 [-4.43 -3.99]	124	p47r	-3.80	-4.33 [-4.46 -4.19]
5	V6	-2.31	-3.95 [-4.10 -3.81]	65	Pol2	-1.35	-4.09 [-4.30 -3.87]	125	A1	-4.16	-4.49 [-4.59 -4.38]
6	V2	-3.18	-4.25 [-4.39 -4.09]	66	FOP4	-5.19	-5.06 [-5.19 -4.91]	126	52	-4.07	-4.37 [-4.50 -4.24]
7	V3	-3.90	-4.62 [-4.74 -4.49]	67	MI	-2.18	-3.94 [-4.11 -3.75]	127	RI	-4.26	-4.50 [-4.62 -4.38]
8	V4	-1.66	-3.80 [-4.00 -3.61]	68	FOP1	-3.14	-4.45 [-4.62 -4.30]	128	TA2	-4.60	-4.62 [-4.74 -4.52]
9	V8	-2.48	-3.89 [-4.04 -3.72]	69	FOP3	-1.38	-3.60 [-3.78 -3.43]	129	PBelt	-4.84	-4.80 [-4.92 -4.70]
10	V3A	-2.14	-4.17 [-4.35 -4.00]	70	PFop	-2.06	-3.84 [-3.99 -3.67]	130	MBelt	-4.68	-4.67 [-4.77 -4.52]
11	V7	-3.17	-4.58 [-4.74 -4.38]	71	PF	-2.52	-4.09 [-4.23 -3.93]	131	LBelt	-3.10	-3.91 [-4.04 -3.80]
12	IPS1	-2.43	-4.14 [-4.28 -3.99]	72	Pol1	-1.67	-3.78 [-3.93 -3.62]	132	A4	-4.46	-4.47 [-4.60 -4.37]
13	FFC	-2.64	-4.05 [-4.18 -3.89]	73	FOP5	-2.67	-4.18 [-4.33 -4.00]	133	7m	-3.89	-4.21 [-4.32 -4.10]
14	V3B	-1.88	-3.69 [-3.83 -3.52]	74	PI	-2.93	-4.17 [-4.32 -4.01]	134	POS1	-3.94	-4.29 [-4.40 -4.19]
15	LO1	-2.11	-3.51 [-3.64 -3.37]	75	a32pr	-3.53	-4.20 [-4.31 -4.08]	135	23d	-4.01	-4.28 [-4.41 -4.17]
16	LO2	-3.71	-4.43 [-4.55 -4.32]	76	p24	-4.24	-4.43 [-4.56 -4.30]	136	v23ab	-4.25	-4.42 [-4.52 -4.29]
17	PIT	-3.59	-4.30 [-4.43 -4.18]	77	PEF	-2.49	-4.04 [-4.17 -3.88]	137	d23ab	-4.26	-4.34 [-4.44 -4.24]
18	MT	-3.83	-4.37 [-4.49 -4.27]	78	7PL	-2.85	-4.23 [-4.37 -4.08]	138	31pv	-4.08	-4.28 [-4.38 -4.18]
19	LIPv	-4.33	-4.60 [-4.70 -4.46]	79	MIP	-4.13	-4.86 [-5.00 -4.69]	139	a24	-4.84	-4.80 [-4.91 -4.70]
20	VIP	-3.77	-4.56 [-4.69 -4.42]	80	LIPd	-4.77	-5.06 [-5.21 -4.91]	140	d32	-5.15	-4.81 [-4.91 -4.70]
21	PH	-3.62	-4.39 [-4.51 -4.26]	81	6a	-3.03	-4.41 [-4.57 -4.27]	141	p32	-5.17	-4.71 [-4.80 -4.61]
22	V6A	-3.72	-4.46 [-4.59 -4.34]	82	PFt	-3.09	-4.31 [-4.47 -4.16]	142	10r	-2.54	-3.96 [-4.09 -3.80]
23	VMV1	-4.55	-4.60 [-4.71 -4.50]	83	AIP	-2.38	-4.13 [-4.31 -3.98]	143	47m	-3.55	-4.15 [-4.25 -4.03]
24	VMV3	-4.69	-4.75 [-4.88 -4.63]	84	PHA3	-2.53	-4.12 [-4.27 -3.95]	144	8Av	-4.12	-4.56 [-4.69 -4.44]
25	V4t	-3.63	-4.30 [-4.42 -4.18]	85	TE2p	-2.68	-4.13 [-4.29 -3.97]	145	8Ad	-3.30	-4.14 [-4.26 -4.02]
26	FST	-1.86	-3.65 [-3.81 -3.49]	86	PHT	-2.64	-4.21 [-4.34 -4.03]	146	9m	-3.76	-4.35 [-4.47 -4.24]
27	V3CD	-1.75	-3.53 [-3.67 -3.38]	87	PGp	-2.30	-3.98 [-4.12 -3.83]	147	8BL	-3.46	-4.42 [-4.54 -4.27]
28	LO3	-4.28	-4.52 [-4.64 -4.40]	88	IP0	-1.38	-3.89 [-4.06 -3.70]	148	9p	-2.89	-4.00 [-4.14 -3.89]
29	VMV2	-2.00	-3.57 [-3.68 -3.44]	89	55b	-2.57	-4.18 [-4.33 -4.06]	149	10d	-3.06	-4.00 [-4.14 -3.89]
30	VVC	-1.73	-3.58 [-3.75 -3.41]	90	PSL	-2.31	-3.99 [-4.13 -3.82]	150	47l	-4.84	-4.69 [-4.81 -4.58]
31	4	-2.67	-3.96 [-4.10 -3.80]	91	SFL	-3.67	-4.39 [-4.49 -4.27]	151	9a	-3.42	-4.06 [-4.16 -3.96]
32	3b	-1.64	-4.06 [-4.26 -3.87]	92	STV	-4.67	-4.85 [-5.01 -4.72]	152	10v	-2.96	-4.13 [-4.25 -4.00]
33	5m	-1.84	-3.90 [-4.07 -3.72]	93	44	-1.99	-4.15 [-4.30 -3.92]	153	10pp	-3.53	-4.67 [-4.81 -4.53]
34	5L	-1.64	-3.98 [-4.17 -3.77]	94	45	-4.67	-4.70 [-4.82 -4.56]	154	OFC	-4.15	-4.79 [-4.92 -4.67]
35	24dd	-2.12	-4.05 [-4.23 -3.81]	95	IFJa	-4.00	-4.53 [-4.65 -4.42]	155	47s	-4.32	-4.61 [-4.73 -4.50]
36	24dv	-1.94	-3.82 [-3.98 -3.65]	96	IFSp	-2.78	-4.24 [-4.41 -4.07]	156	EC	-4.04	-4.49 [-4.61 -4.38]
37	7AL	-2.49	-3.85 [-3.98 -3.69]	97	STGa	-2.52	-4.22 [-4.37 -4.04]	157	PreS	-4.41	-4.44 [-4.55 -4.34]
38	7PC	-2.47	-3.86 [-4.04 -3.70]	98	A5	-2.82	-4.37 [-4.54 -4.21]	158	H	-3.82	-4.48 [-4.60 -4.33]
39	1	-2.02	-3.59 [-3.75 -3.46]	99	STSda	-4.16	-4.62 [-4.76 -4.49]	159	PHA1	-3.97	-4.43 [-4.56 -4.32]
40	2	-1.92	-3.90 [-4.06 -3.67]	100	STSdp	-4.44	-4.65 [-4.76 -4.53]	160	STSvp	-4.14	-4.75 [-4.87 -4.62]
41	3a	-2.40	-4.17 [-4.36 -3.97]	101	TPOJ1	-4.95	-4.80 [-4.92 -4.68]	161	TGd	-2.32	-3.85 [-4.01 -3.66]
42	6d	-2.06	-3.50 [-3.60 -3.38]	102	TGv	-5.82	-5.08 [-5.21 -4.97]	162	TE1a	-1.99	-3.89 [-4.07 -3.68]
43	6mp	-1.57	-3.67 [-3.84 -3.42]	103	RSC	-4.49	-4.43 [-4.52 -4.33]	163	TE2a	-3.62	-4.48 [-4.61 -4.37]
44	6v	-2.03	-3.82 [-3.97 -3.66]	104	POS2	-4.43	-4.67 [-4.80 -4.54]	164	PGi	-1.55	-4.22 [-4.41 -4.02]
45	OP4	-1.97	-3.41 [-3.55 -3.27]	105	7Pm	-4.01	-4.48 [-4.59 -4.34]	165	PGs	-1.51	-4.34 [-4.55 -4.07]
46	OP1	-2.65	-3.76 [-3.87 -3.65]	106	8BM	-4.18	-4.21 [-4.33 -4.09]	166	PHA2	-2.16	-4.17 [-4.33 -4.02]
47	OP2-3	-2.69	-3.90 [-4.01 -3.78]	107	8C	-4.46	-4.46 [-4.57 -4.36]	167	31pd	-4.05	-4.16 [-4.25 -4.05]
48	FOP2	-3.04	-4.07 [-4.17 -3.94]	108	a47r	-4.41	-4.52 [-4.67 -4.39]	168	31a	-5.76	-5.07 [-5.17 -4.94]
49	Ig	-2.55	-3.74 [-3.85 -3.64]	109	IFJp	-4.91	-4.65 [-4.77 -4.52]	169	25	-5.26	-4.79 [-4.89 -4.66]
50	FEF	-3.14	-4.16 [-4.26 -4.05]	110	IFSa	-3.04	-4.09 [-4.23 -3.94]	170	s32	-2.53	-4.13 [-4.26 -3.97]
51	5mv	-2.32	-3.83 [-3.96 -3.69]	111	p9-46v	-5.51	-4.88 [-5.00 -4.76]	171	STSva	-2.92	-4.24 [-4.40 -4.07]
52	23c	-2.37	-3.82 [-4.00 -3.67]	112	a9-46v	-4.75	-4.48 [-4.59 -4.36]	172	TE1m	-3.81	-4.16 [-4.25 -4.04]
53	SCEF	-2.75	-4.18 [-4.31 -4.03]	113	a10p	-4.71	-4.76 [-4.90 -4.60]	173	PCV	-4.43	-4.49 [-4.60 -4.37]
54	6ma	-1.95	-4.03 [-4.17 -3.88]	114	11l	-5.68	-5.03 [-5.17 -4.88]	174	TPOJ2	-4.09	-4.50 [-4.60 -4.37]
55	7Am	-1.68	-3.79 [-3.93 -3.64]	115	13l	-6.03	-5.14 [-5.26 -5.04]	175	TPOJ3	-3.63	-4.26 [-4.36 -4.14]
56	p24pr	-2.72	-4.32 [-4.49 -4.13]	116	i6-8	-3.63	-4.61 [-4.74 -4.48]	176	PeEc	-4.70	-4.66 [-4.77 -4.55]
57	33pr	-1.78	-4.08 [-4.26 -3.85]	117	s6-8	-4.15	-4.68 [-4.81 -4.56]	177	TF	-4.35	-4.48 [-4.57 -4.37]
58	a24pr	-2.02	-4.11 [-4.35 -3.87]	118	AVI	-3.84	-4.27 [-4.40 -4.15]	178	Pir	-4.41	-4.34 [-4.43 -4.23]
59	p32pr	-1.92	-4.06 [-4.27 -3.84]	119	TE1p	-3.35	-4.09 [-4.23 -3.96]	179	AAIC	-1.93	-4.01 [-4.19 -3.82]
60	6r	-1.97	-3.98 [-4.21 -3.76]	120	IP2	-3.10	-4.00 [-4.11 -3.88]	180	pOFC	-1.76	-4.03 [-4.25 -3.80]

Figure 6-1. Differential connectivity between contralateral homologous parcels vs the mean of all other contralateral parcels. Confidence intervals are Bonferroni-corrected for multiple comparisons.

1118

Figure 7. Language/Auditory network hyperconnectivity and left-lateralization. (**A**) Distance-binned connectivity within the language and auditory networks compared to connectivity between the language and auditory networks and other networks, separately for the left and right hemispheres (**B**) The differential trace for the within- and between- connectivity in both hemispheres. In both panels, gray patches show Bonferroni-corrected bootstrapped 95% confidence intervals across subjects.

Figure 8. Connectivity is influenced by the cortical hierarchy. (**A**, **B**) Connectivity is strongly predicted by hierarchical similarity in some networks and modestly predicted overall. (**A**) All connectivity vs. myelination difference, including within- and across- network connections, for the left, right, and callosal connections. For both panels, each marker represents a parcel pair. (**B**) Within-network connectivity vs. myelination difference for 10 functional networks. Linear fits and correlation coefficients computed independently for the left and right hemisphere. A negative correlation indicates that parcels at similar hierarchical levels tend to be more connected. (**C**, **D**) Higher order prefrontal areas are better connected. (**C**) Histogram of correlation coefficients between areal myelination and F_{pt} connectivity to each parcel. Only significant coefficients after Bonferroni correction are shown. Most coefficients are negative indicating high connectivity to low-myelination (i.e., higher-order) areas. (**D**) Significant negative coefficients (red) map onto bilateral prefrontal cortex. Only the bilateral DVT and V6A are show positive significant correlations (blue).

Figure 8-1. Myelination difference connectivity matrix. This provides an estimate for the difference in hierarchical level between cortical parcels. Values have been fractionally scaled. Note that the color scale has been reversed when compared to figure **1**, as |Δmyelination| is inversely proportional to connectivity.

Idx.	Parcel	r	p	Idx.	Parcel	r	p	Idx.	Parcel	r	p
1	V1	0.07	n.s.	61	46	-0.31	7.18E-07	121	IP1	0.12	n.s.
2	ProS	0.15	n.s.	62	9-46d	-0.31	4.04E-07	122	PFm	0.04	n.s.
3	DVT	0.23	3.99E-03	63	43	0.00	n.s.	123	p10p	-0.33	2.77E-08
4	MST	0.06	n.s.	64	PFcm	0.03	n.s.	124	p47r	-0.26	2.51E-04
5	V6	0.20	n.s.	65	PoI2	-0.03	n.s.	125	A1	0.05	n.s.
6	V2	0.09	n.s.	66	FOP4	-0.14	n.s.	126	52	0.06	n.s.
7	V3	0.12	n.s.	67	MI	-0.12	n.s.	127	RI	0.01	n.s.
8	V4	0.12	n.s.	68	FOP1	-0.06	n.s.	128	TA2	0.05	n.s.
9	V8	0.08	n.s.	69	FOP3	-0.10	n.s.	129	PBelt	0.10	n.s.
10	V3A	0.15	n.s.	70	PFop	0.05	n.s.	130	MBelt	0.08	n.s.
11	V7	0.16	n.s.	71	PF	0.05	n.s.	131	LBelt	0.07	n.s.
12	IPS1	0.17	n.s.	72	PoI1	-0.01	n.s.	132	A4	0.10	n.s.
13	FFC	0.12	n.s.	73	FOP5	-0.10	n.s.	133	7m	-0.05	n.s.
14	V3B	0.16	n.s.	74	PI	-0.02	n.s.	134	POS1	0.07	n.s.
15	LO1	0.15	n.s.	75	a32pr	-0.30	2.60E-06	135	23d	-0.12	n.s.
16	LO2	0.12	n.s.	76	p24	-0.27	6.33E-05	136	v23ab	-0.08	n.s.
17	PIT	0.12	n.s.	77	PEF	-0.16	n.s.	137	d23ab	-0.10	n.s.
18	MT	0.05	n.s.	78	7PL	0.13	n.s.	138	31pv	-0.13	n.s.
19	LIPv	0.06	n.s.	79	MIP	0.14	n.s.	139	a24	-0.31	1.12E-06
20	VIP	0.08	n.s.	80	LIPd	0.08	n.s.	140	d32	-0.34	1.23E-08
21	PH	0.10	n.s.	81	6a	-0.11	n.s.	141	p32	-0.35	4.09E-09
22	V6A	0.20	4.92E-02	82	PFt	0.07	n.s.	142	10r	-0.36	4.05E-10
23	VMV1	0.09	n.s.	83	AIP	0.07	n.s.	143	47m	-0.16	n.s.
24	VMV3	0.08	n.s.	84	PHA3	0.11	n.s.	144	8Av	-0.21	2.27E-02
25	V4t	0.09	n.s.	85	TE2p	0.08	n.s.	145	8Ad	-0.30	3.27E-06
26	FST	0.08	n.s.	86	PHT	0.03	n.s.	146	9m	-0.38	4.11E-11
27	V3CD	0.17	n.s.	87	PGp	0.13	n.s.	147	8BL	-0.35	2.80E-09
28	LO3	0.11	n.s.	88	IP0	0.17	n.s.	148	9p	-0.33	4.18E-08
29	VMV2	0.06	n.s.	89	55b	-0.10	n.s.	149	10d	-0.37	1.90E-10
30	VVC	0.13	n.s.	90	PSL	0.03	n.s.	150	47l	-0.15	n.s.
31	4	0.02	n.s.	91	SFL	-0.29	1.00E-05	151	9a	-0.33	3.62E-08
32	3b	0.04	n.s.	92	STV	0.03	n.s.	152	10v	-0.34	2.60E-08
33	5m	0.01	n.s.	93	44	-0.21	2.25E-02	153	10pp	-0.26	3.45E-04
34	5L	0.04	n.s.	94	45	-0.18	n.s.	154	OFC	-0.28	3.28E-05
35	24dd	-0.09	n.s.	95	IFJa	-0.17	n.s.	155	47s	-0.19	n.s.
36	24dv	-0.16	n.s.	96	IFSp	-0.24	1.01E-03	156	EC	0.02	n.s.
37	7AL	0.04	n.s.	97	STGa	-0.03	n.s.	157	PreS	0.01	n.s.
38	7PC	0.06	n.s.	98	A5	0.07	n.s.	158	H	0.06	n.s.
39	1	0.06	n.s.	99	STSda	0.04	n.s.	159	PHA1	0.07	n.s.
40	2	0.06	n.s.	100	STSdp	-0.01	n.s.	160	STSvp	-0.04	n.s.
41	3a	0.05	n.s.	101	TPOJ1	0.00	n.s.	161	TGd	-0.11	n.s.
42	6d	-0.05	n.s.	102	TGv	-0.03	n.s.	162	TE1a	-0.04	n.s.
43	6mp	-0.03	n.s.	103	RSC	-0.03	n.s.	163	TE2a	-0.02	n.s.
44	6v	-0.09	n.s.	104	POS2	0.11	n.s.	164	PGi	0.01	n.s.
45	OP4	0.06	n.s.	105	7Pm	0.00	n.s.	165	PGs	0.06	n.s.
46	OP1	0.03	n.s.	106	8BM	-0.37	1.14E-10	166	PHA2	0.08	n.s.
47	OP2-3	-0.01	n.s.	107	8C	-0.22	1.25E-02	167	31pd	-0.13	n.s.
48	FOP2	-0.06	n.s.	108	a47r	-0.25	4.75E-04	168	31a	-0.09	n.s.
49	Ig	-0.02	n.s.	109	IFJp	-0.17	n.s.	169	25	-0.27	9.02E-05
50	FEF	-0.07	n.s.	110	IFSa	-0.25	3.91E-04	170	s32	-0.34	2.28E-08
51	5mv	-0.07	n.s.	111	p9-46v	-0.28	2.50E-05	171	STSva	0.02	n.s.
52	23c	-0.12	n.s.	112	a9-46v	-0.30	1.33E-06	172	TE1m	-0.05	n.s.
53	SCEF	-0.20	n.s.	113	a10p	-0.30	2.32E-06	173	PCV	-0.07	n.s.
54	6ma	-0.13	n.s.	114	11l	-0.18	n.s.	174	TPOJ2	0.02	n.s.
55	7Am	0.01	n.s.	115	13l	-0.19	n.s.	175	TPOJ3	0.04	n.s.
56	p24pr	-0.18	n.s.	116	i6-8	-0.16	n.s.	176	PeEc	0.02	n.s.
57	33pr	-0.18	n.s.	117	s6-8	-0.27	5.76E-05	177	TF	0.08	n.s.
58	a24pr	-0.26	2.99E-04	118	AVI	-0.13	n.s.	178	Pir	-0.16	n.s.
59	p32pr	-0.26	1.58E-04	119	TE1p	0.01	n.s.	179	AAIC	-0.18	n.s.
60	6r	-0.14	n.s.	120	IP2	0.06	n.s.	180	pOFC	-0.24	1.58E-03

Figure 8-2. Pearson correlations between the F_{pt} from each left hemisphere parcel to all others and the target parcels' myelination indices. p values are Bonferroni-corrected for multiple comparisons.

Idx.	Parcel	r	p	Idx.	Parcel	r	p	Idx.	Parcel	r	p
181	R_V1	0.08	n.s.	241	R_46	-0.29	1.20E-05	301	R_IP1	0.13	n.s.
182	R_ProS	0.12	n.s.	242	R_9-46d	-0.29	7.49E-06	302	R_PFm	0.02	n.s.
183	R_DVT	0.23	2.52E-03	243	R_43	0.00	n.s.	303	R_p10p	-0.30	2.07E-06
184	R_MST	0.11	n.s.	244	R_PFcm	0.05	n.s.	304	R_p47r	-0.21	1.69E-02
185	R_V6	0.20	n.s.	245	R_PoI2	0.03	n.s.	305	R_A1	0.06	n.s.
186	R_V2	0.10	n.s.	246	R_FOP4	-0.12	n.s.	306	R_52	0.12	n.s.
187	R_V3	0.13	n.s.	247	R_MI	-0.08	n.s.	307	R_RI	0.03	n.s.
188	R_V4	0.14	n.s.	248	R_FOP1	-0.07	n.s.	308	R_TA2	0.09	n.s.
189	R_V8	0.09	n.s.	249	R_FOP3	-0.08	n.s.	309	R_PBelt	0.12	n.s.
190	R_V3A	0.18	n.s.	250	R_PFop	0.07	n.s.	310	R_MBelt	0.09	n.s.
191	R_V7	0.19	n.s.	251	R_PF	0.05	n.s.	311	R_LBelt	0.08	n.s.
192	R_IPS1	0.20	n.s.	252	R_PoI1	0.09	n.s.	312	R_A4	0.15	n.s.
193	R_FFC	0.15	n.s.	253	R_FOP5	-0.05	n.s.	313	R_7m	-0.05	n.s.
194	R_V3B	0.18	n.s.	254	R_PI	0.08	n.s.	314	R_POS1	0.06	n.s.
195	R_LO1	0.19	n.s.	255	R_a32pr	-0.29	6.54E-06	315	R_23d	-0.10	n.s.
196	R_LO2	0.17	n.s.	256	R_p24	-0.27	9.60E-05	316	R_v23ab	-0.08	n.s.
197	R_PIT	0.15	n.s.	257	R_PEF	-0.15	n.s.	317	R_d23ab	-0.10	n.s.
198	R_MT	0.11	n.s.	258	R_7PL	0.16	n.s.	318	R_31pv	-0.11	n.s.
199	R_LIPv	0.09	n.s.	259	R_MIP	0.17	n.s.	319	R_a24	-0.28	3.54E-05
200	R_VIP	0.11	n.s.	260	R_LIPd	0.08	n.s.	320	R_d32	-0.32	2.51E-07
201	R_PH	0.15	n.s.	261	R_6a	-0.08	n.s.	321	R_p32	-0.35	5.11E-09
202	R_V6A	0.23	4.12E-03	262	R_PFt	0.07	n.s.	322	R_10r	-0.37	1.63E-10
203	R_VMV1	0.11	n.s.	263	R_AIP	0.05	n.s.	323	R_47m	-0.07	n.s.
204	R_VMV3	0.10	n.s.	264	R_PHA3	0.12	n.s.	324	R_8Av	-0.20	n.s.
205	R_V4t	0.14	n.s.	265	R_TE2p	0.12	n.s.	325	R_8Ad	-0.26	1.58E-04
206	R_FST	0.14	n.s.	266	R_PHT	0.09	n.s.	326	R_9m	-0.37	2.04E-10
207	R_V3CD	0.19	n.s.	267	R_PGp	0.16	n.s.	327	R_8BL	-0.35	1.95E-09
208	R_LO3	0.17	n.s.	268	R_IP0	0.20	n.s.	328	R_9p	-0.31	8.63E-07
209	R_VMV2	0.08	n.s.	269	R_55b	-0.08	n.s.	329	R_10d	-0.36	3.48E-10
210	R_VVC	0.14	n.s.	270	R_PSL	0.05	n.s.	330	R_47l	-0.07	n.s.
211	R_4	0.04	n.s.	271	R_SFL	-0.22	1.06E-02	331	R_9a	-0.30	1.63E-06
212	R_3b	0.06	n.s.	272	R_STV	0.04	n.s.	332	R_10v	-0.34	1.05E-08
213	R_5m	0.04	n.s.	273	R_44	-0.20	n.s.	333	R_10pp	-0.24	9.97E-04
214	R_5L	0.06	n.s.	274	R_45	-0.12	n.s.	334	R_OFC	-0.24	1.10E-03
215	R_24dd	-0.06	n.s.	275	R_IFJa	-0.18	n.s.	335	R_47s	-0.07	n.s.
216	R_24dv	-0.15	n.s.	276	R_IFSp	-0.24	1.75E-03	336	R_EC	0.03	n.s.
217	R_7AL	0.07	n.s.	277	R_STGa	0.03	n.s.	337	R_PreS	0.01	n.s.
218	R_7PC	0.07	n.s.	278	R_A5	0.11	n.s.	338	R_H	0.07	n.s.
219	R_1	0.07	n.s.	279	R_STSda	0.09	n.s.	339	R_PHA1	0.09	n.s.
220	R_2	0.07	n.s.	280	R_STSdp	0.04	n.s.	340	R_STSvp	0.03	n.s.
221	R_3a	0.07	n.s.	281	R_TPOJ1	0.03	n.s.	341	R_TGd	-0.07	n.s.
222	R_6d	-0.03	n.s.	282	R_TGv	0.03	n.s.	342	R_TE1a	0.01	n.s.
223	R_6mp	-0.01	n.s.	283	R_RSC	-0.05	n.s.	343	R_TE2a	0.02	n.s.
224	R_6v	-0.08	n.s.	284	R_POS2	0.12	n.s.	344	R_PGi	0.03	n.s.
225	R_OP4	0.07	n.s.	285	R_7Pm	0.03	n.s.	345	R_PGs	0.08	n.s.
226	R_OP1	0.04	n.s.	286	R_8BM	-0.35	1.71E-09	346	R_PHA2	0.08	n.s.
227	R_OP2-3	0.00	n.s.	287	R_8C	-0.21	1.51E-02	347	R_31pd	-0.12	n.s.
228	R_FOP2	-0.05	n.s.	288	R_a47r	-0.23	3.49E-03	348	R_31a	-0.09	n.s.
229	R_Ig	0.00	n.s.	289	R_IFJp	-0.15	n.s.	349	R_25	-0.24	1.61E-03
230	R_FEF	-0.05	n.s.	290	R_IFSa	-0.21	3.22E-02	350	R_s32	-0.32	1.37E-07
231	R_5mv	-0.04	n.s.	291	R_p9-46v	-0.27	8.34E-05	351	R_STSva	0.08	n.s.
232	R_23c	-0.09	n.s.	292	R_a9-46v	-0.28	2.20E-05	352	R_TE1m	0.02	n.s.
233	R_SCEF	-0.16	n.s.	293	R_a10p	-0.25	8.16E-04	353	R_PCV	-0.05	n.s.
234	R_6ma	-0.08	n.s.	294	R_11l	-0.13	n.s.	354	R_TPOJ2	0.06	n.s.
235	R_7Am	0.04	n.s.	295	R_13l	-0.11	n.s.	355	R_TPOJ3	0.08	n.s.
236	R_p24pr	-0.17	n.s.	296	R_i6-8	-0.13	n.s.	356	R_PeEc	0.04	n.s.
237	R_33pr	-0.18	n.s.	297	R_s6-8	-0.22	9.03E-03	357	R_TF	0.13	n.s.
238	R_a24pr	-0.26	2.39E-04	298	R_AVI	-0.03	n.s.	358	R_Pir	-0.11	n.s.
239	R_p32pr	-0.25	4.93E-04	299	R_TE1p	0.07	n.s.	359	R_AAIC	-0.09	n.s.
240	R_6r	-0.14	n.s.	300	R_IP2	0.03	n.s.	360	R_pOFC	-0.20	3.99E-02

Figure 8-3. Pearson correlations between the F_{pt} from each right hemisphere parcel to all others and the target parcels' myelination indices. p values are Bonferroni-corrected for multiple comparisons.

Figure 9. Probabilistic dMRI more closely resembles CCEPs than resting-state fMRI. (**A**) Connectivity matrices for probabilistic dMRI tractography, CCEP, and rs-fMRI. For CCEPs missing data has been colored grey and pre-log zero-strength connections black. (**B**) Correlations among the three modalities. The least-squares linear fit is shown in red. (**C**) Non-zero pairwise connection strength distributions. Note that rs-fMRI connectivity values, which are not log-transformed, display two modes, separated at 0.0014.(**D**) Cortical parcels displaying lower (left) and higher (right) modes of rs-fMRI connectivity.

Figure 9-1. Within-hemisphere comparison of probabilistic dMRI tractography, CCEP, and rs-fMRI connectivity. For the left and right hemisphere, the distribution of pairwise non-zero connection strengths and correlations among the three modalities are shown. The least-squares linear fit is shown in red. All within-hemisphere findings are concordant with the overall findings, shown in figure **9**.

Figure 10. Network theoretic differences between the connectivity modalities. Binarized network metrics after thresholding by edge weight (connectivity strength).

1126

Connectome Features
Probabilistic methodology sensitive to weak connections yielding a fully-populated, un-thresholded connectome
Cortex parcellated into the standardized, relatively dense, and functionally relevant HCP-MMP1.0 atlas
Large normative sample size (N = 1,065)
Enables comparison with other measures in the WU-Minn HCP and other cohorts

Table 1. Connectome Features

Idx.	Parcel	Orig.	Network	Idx.	Parcel	Orig.	Network	Idx.	Parcel	Orig.	Network
1	V1	1	Cingulo-Opercular	61	46	84	Cingulo-Opercular	121	IP1	145	Frontoparietal
2	ProS	121	Visual	62	9-46d	86	Cingulo-Opercular	122	PFm	149	Frontoparietal
3	DVT	142	Visual	63	43	99	Cingulo-Opercular	123	p10p	170	Frontoparietal
4	MST	2	Visual	64	PFcm	105	Cingulo-Opercular	124	p47r	171	Frontoparietal
5	V6	3	Visual	65	PoI2	106	Cingulo-Opercular	125	A1	24	Auditory
6	V2	4	Visual	66	FOP4	108	Cingulo-Opercular	126	52	103	Auditory
7	V3	5	Visual	67	MI	109	Cingulo-Opercular	127	RI	104	Auditory
8	V4	6	Visual	68	FOP1	113	Cingulo-Opercular	128	TA2	107	Auditory
9	V8	7	Visual	69	FOP3	114	Cingulo-Opercular	129	PBelt	124	Auditory
10	V3A	13	Visual	70	PFop	147	Cingulo-Opercular	130	MBelt	173	Auditory
11	V7	16	Visual	71	PF	148	Cingulo-Opercular	131	LBelt	174	Auditory
12	IPS1	17	Visual	72	PoI1	167	Cingulo-Opercular	132	A4	175	Auditory
13	FFC	18	Visual	73	FOP5	169	Cingulo-Opercular	133	7m	30	Default Mode
14	V3B	19	Visual	74	PI	178	Cingulo-Opercular	134	POS1	31	Default Mode
15	LO1	20	Visual	75	a32pr	179	Cingulo-Opercular	135	23d	32	Default Mode
16	LO2	21	Visual	76	p24	180	Cingulo-Opercular	136	v23ab	33	Default Mode
17	PIT	22	Visual	77	PEF	11	Dorsal Attention	137	d23ab	34	Default Mode
18	MT	23	Visual	78	7PL	46	Dorsal Attention	138	31pv	35	Default Mode
19	LIPv	48	Visual	79	MIP	50	Dorsal Attention	139	a24	61	Default Mode
20	VIP	49	Visual	80	LIPd	95	Dorsal Attention	140	d32	62	Default Mode
21	PH	138	Visual	81	6a	96	Dorsal Attention	141	p32	64	Default Mode
22	V6A	152	Visual	82	PFt	116	Dorsal Attention	142	10r	65	Default Mode
23	VMV1	153	Visual	83	AIP	117	Dorsal Attention	143	47m	66	Default Mode
24	VMV3	154	Visual	84	PHA3	127	Dorsal Attention	144	8Av	67	Default Mode
25	V4t	156	Visual	85	TE2p	136	Dorsal Attention	145	8Ad	68	Default Mode
26	FST	157	Visual	86	PHT	137	Dorsal Attention	146	9m	69	Default Mode
27	V3CD	158	Visual	87	PGp	143	Dorsal Attention	147	8BL	70	Default Mode
28	LO3	159	Visual	88	IP0	146	Dorsal Attention	148	9p	71	Default Mode
29	VMV2	160	Visual	89	55b	12	Language	149	10d	72	Default Mode
30	VVC	163	Visual	90	PSL	25	Language	150	47l	76	Default Mode
31	4	8	Somatomotor	91	SFL	26	Language	151	9a	87	Default Mode
32	3b	9	Somatomotor	92	STV	28	Language	152	10v	88	Default Mode
33	5m	36	Somatomotor	93	44	74	Language	153	10pp	90	Default Mode
34	5L	39	Somatomotor	94	45	75	Language	154	OFC	93	Default Mode
35	24dd	40	Somatomotor	95	IFJa	79	Language	155	47s	94	Default Mode
36	24dv	41	Somatomotor	96	IFSp	81	Language	156	EC	118	Default Mode
37	7AL	42	Somatomotor	97	STGa	123	Language	157	PreS	119	Default Mode
38	7PC	47	Somatomotor	98	A5	125	Language	158	H	120	Default Mode
39	1	51	Somatomotor	99	STSda	128	Language	159	PHA1	126	Default Mode
40	2	52	Somatomotor	100	STSdp	129	Language	160	STSvp	130	Default Mode
41	3a	53	Somatomotor	101	TPOJ1	139	Language	161	TGd	131	Default Mode
42	6d	54	Somatomotor	102	TGv	172	Language	162	TE1a	132	Default Mode
43	6mp	55	Somatomotor	103	RSC	14	Frontoparietal	163	TE2a	134	Default Mode
44	6v	56	Somatomotor	104	POS2	15	Frontoparietal	164	PGi	150	Default Mode
45	OP4	100	Somatomotor	105	7Pm	29	Frontoparietal	165	PGs	151	Default Mode
46	OP1	101	Somatomotor	106	8BM	63	Frontoparietal	166	PHA2	155	Default Mode
47	OP2-3	102	Somatomotor	107	8C	73	Frontoparietal	167	31pd	161	Default Mode
48	FOP2	115	Somatomotor	108	a47r	77	Frontoparietal	168	31a	162	Default Mode
49	Ig	168	Somatomotor	109	IFJp	80	Frontoparietal	169	25	164	Default Mode
50	FEF	10	Cingulo-Opercular	110	IFSa	82	Frontoparietal	170	s32	165	Default Mode
51	5mv	37	Cingulo-Opercular	111	p9-46v	83	Frontoparietal	171	STSva	176	Default Mode
52	23c	38	Cingulo-Opercular	112	a9-46v	85	Frontoparietal	172	TE1m	177	Default Mode
53	SCEF	43	Cingulo-Opercular	113	a10p	89	Frontoparietal	173	PCV	27	Multimodal
54	6ma	44	Cingulo-Opercular	114	11l	91	Frontoparietal	174	TPOJ2	140	Multimodal
55	7Am	45	Cingulo-Opercular	115	13l	92	Frontoparietal	175	TPOJ3	141	Multimodal
56	p24pr	57	Cingulo-Opercular	116	i6-8	97	Frontoparietal	176	PeEc	122	Multimodal
57	33pr	58	Cingulo-Opercular	117	s6-8	98	Frontoparietal	177	TF	135	Multimodal
58	a24pr	59	Cingulo-Opercular	118	AVI	111	Frontoparietal	178	Pir	110	Orbito-Affective
59	p32pr	60	Cingulo-Opercular	119	TE1p	133	Frontoparietal	179	AAIC	112	Orbito-Affective
60	6r	78	Cingulo-Opercular	120	IP2	144	Frontoparietal	180	pOFC	166	Orbito-Affective

Table 2. Parcel order and network assignment. The emboldened indices refer to the parcel order in figure 1A. The Orig. indices refer to the original parcel order presented in (Glasser et al., 2016). All indices refer to the left hemisphere, adding 180 yields the homologous right hemisphere indices.

Location	Data structure	Test or analysis	N	Uncertainty [CI$_{95\%}$]
Fig 1-1D	Gaussian predictor Exponential response	Nonlinear regression (iterative optimization)	64,620 64,620 64,620 64,620	λ = 23.8 [23.5 24.0] λ = 22.8 [22.7 22.9] λ = 22.2 [22.1 22.2] λ = 23.4 [23.3 23.6]
Fig 2A	Gaussian predictor Exponential response	Nonlinear regression (iterative optimization)	16,110	λ = 23.1 [22.8 23.3]
Fig 2B	Gaussian predictor Exponential response	Nonlinear regression (iterative optimization)	16,110	λ = 23.9 [23.7 24.2]
Fig 2C	Gaussian predictor Exponential response	Nonlinear regression (iterative optimization)	32,400	λ = 32.8 [32.5 33.0]
Fig 2D	Gaussian predictor Exponential response	Nonlinear regression (iterative optimization)	64,620	λ = 23.4 [23.3 23.6]
Fig 2-2B	Gaussian predictor Gaussian response	Linear correlation	1,065	r = -0.14 [-0.20 -0.08]
Fig 3D	Gaussian predictor Exponential response	Nonlinear regression (iterative optimization)	12,924	λ = 27.8 [27.4 28.2]
Fig 3F	Gaussian predictor Gaussian response	Linear correlation	1,065	r = 0.70 [0.67 0.73]
Fig 4C	Gaussian predictor Gaussian response	Linear correlation	80	r = 0.35 [0.14 0.53]
Fig 8A	Gaussian predictor Gaussian response	Linear correlation	16,110 16,110 32,400	r = -0.10 [-0.12 -0.09] r = -0.12 [-0.13 -0.10] r = -0.11 [-0.12 -0.10]
Fig 8B	Gaussian predictor Gaussian response	Linear correlation	351 351 66 91 231 231 28 780 780 10	r = -0.17 [-0.27 -0.06] r = -0.13 [-0.23 -0.02] r = -0.41 [-0.60 -0.19] r = -0.26 [-0.44 -0.06] r = -0.30 [-0.42 -0.18] r = -0.30 [-0.40 -0.17] r = -0.56 [-0.77 -0.24] r = -0.12 [-0.19 -0.05] r = -0.17 [-0.24 -0.10] r = -0.74 [-0.93 -0.20]
Fig 9B	Gaussian predictor Gaussian response	Linear correlation	19,667 19,667 64,620	r = 0.43 [0.42 0.44] r = 0.23 [0.21 0.24] r = 0.06 [0.05 0.07]
Figure 9-1	Gaussian predictor Gaussian response	Linear correlation	8,483 8,483 16,110 8,370 8,370 16,110	r = 0.42 [0.40 0.44] r = 0.22 [0.20 0.24] r = 0.06 [0.05 0.07] r = 0.40 [0.38 0.42] r = 0.22 [0.20 0.24] r = 0.11 [0.10 0.13]

Table 3. Statistics and uncertainty. Where multiple uncertainties are listed for a figure panel, they correspond to the statistics read left-to-right, top-to-bottom in that panel. For figure 8B only uncertainties for significant correlations are listed. Uncertainties for figures 6, 7, 8 and 10 are not shown. Figure 6-1 contains bootstrapped 95% confidence intervals for the 180 means shown in figure 6, n = 179. Figure 7 shows Bootstrapped 95% confidence intervals in gray; the values of these intervals for all distance bins are available in the figure source data at https://doi.org/10.5281/zenodo.4060485. For figure 10 means across shuffled matrices are only necessary to account for arbitrary ordering among tied edge weights and the bootstrapped 95% confidence intervals for these means are vanishingly small. The values of these intervals at all network densities are also included in the figure source data. For nonlinear regressions confidence intervals are estimated using R^{-1}, the inverse R factor from QR decomposition of the Jacobian, the degrees of freedom for error, and the root mean squared error. For linear correlations the confidence intervals are based on an asymptotic normal distribution of 0.5*log((1+r)/(1-r)), with an approximate variance equal to 1/(N-3). For descriptive statistics, e.g. means, empirical 95% confidence intervals are estimated by bootstrapping with