Rapid molecular evolution of *Spiroplasma* symbionts of *Drosophila*

Michael Gerth[1*], Humberto Martinez-Montoya[2], Paulino Ramirez[3], Florent Masson[4], Joanne S. Griffin[1], Rodolfo Aramayo[5], Stefanos Siozios[1], Bruno Lemaitre[4], Mariana Mateos[6] & Gregory D.D. Hurst[1]

[1] Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK

[2] Laboratory of Genetics and Comparative Genomics, Unidad Académica Multidisciplinaria Reynosa Aztlán, Universidad Autónoma de Tamaulipas, Reynosa, Mexico

[3] Department of Cell Systems and Anatomy, University of Texas Health San Antonio, San Antonio, Texas, USA

[4] Global Health Institute, School of Life Sciences, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

[5] Department of Biology, Texas A&M University, College Station, Texas, USA

[6] Department of Ecology and Conservation Biology, Texas A&M University, College Station, Texas, USA

*Author for correspondence.

Current address: Department of Biological and Medical Sciences, Oxford Brookes University, Oxford, UK mgerth@brookes.ac.uk

Abstract

Spiroplasma are a group of Mollicutes whose members include plant pathogens, insect pathogens, and endosymbionts of animals. In arthropods, *Spiroplasma* are found across a broad host range, but typically with lower incidence than other bacteria with similar ecology, such as *Wolbachia* or *Rickettsia*. *Spiroplasma* symbionts of *Drosophila* are best known as male-killers and protective symbionts, and both phenotypes are mediated by *Spiroplasma*-encoded toxins. *Spiroplasma* phenotypes have been repeatedly observed to be spontaneously lost in *Drosophila* cultures, and several studies have documented a high genomic turnover in *Spiroplasma* symbionts and plant pathogens. These observations suggest that *Spiroplasma* evolves quickly. Here, we systematically assess evolutionary rates and patterns of *Spiroplasma poulsonii*, a natural symbiont of *Drosophila*. We analysed genomic evolution of *s*Hy within flies, and *s*Mel within *in vitro* culture over several years. We observed that *S. poulsonii* substitution rates are among the highest reported for any bacteria, and markedly increased compared with other symbionts. The absence of mismatch repair loci *mutS* and *mutL* is conserved across *Spiroplasma* and likely contributes to elevated substitution rates. Further, the closely related strains *s*Mel and *s*Hy (>99.5% sequence identity in shared loci) show extensive structural genomic differences, which may be explained by a higher degree of host adaptation in *s*Hy, a protective symbiont of *Drosophila hydei*. Finally, comparison across diverse *Spiroplasma* lineages confirms previous reports of dynamic evolution of toxins, and identifies loci similar to the male-killing toxin Spaid in several *Spiroplasma* lineages and other endosymbionts. Overall, our results highlight the peculiar nature of *Spiroplasma* genome evolution, which may explain unusual features of its evolutionary ecology.

Introduction

Many bacterial lineages have evolved to become associates of animal hosts (McFall-Ngai et al., 2013). In arthropods, such associations are the rule, and maternally inherited, endosymbiotic bacteria are especially common and diverse (Duron et al., 2008; Weinert et al., 2015). Prominent examples include obligate nutritional symbionts of blood and sap feeders (Douglas, 2009), reproductive manipulators (Drew et al., 2019), and protective symbionts (Oliver et al., 2014). Collectively, inherited symbionts of arthropods are highly diverse with respect to potential benefits and costs induced, and their degree of adaptation to hosts. For example, even within a single lineage of inherited symbionts - *Wolbachia* - there are nutritional mutualists (Hosokawa et al., 2010), protective symbionts (Teixeira et al., 2008), and reproductive manipulators (Dyson et al., 2002). Importantly, these symbiont-conferred traits may be modulated by environmental factors (Corbin et al., 2017) and host genetic background (Hornett et al., 2008).

Spiroplasma are small, helical bacteria that lack cell-walls. Like other Mollicutes (Mycoplasma-like organisms), they are exclusively found in host association and are often pathogenic (Razin, 2006). Among the first discovered spiroplasmas were plant pathogens (reviewed in Saglio and Whitcomb, 1979) and an inherited symbiont of *Drosophila* with the ability to shift sex ratios towards females (Poulson and Sakaguchi, 1961). Subsequently, *Spiroplasma* symbionts were found in a number of different *Drosophila* species (Haselkorn, 2010), as well as many other arthropods (Gasparich, 2002). More recently, molecular evidence was brought forward for *Spiroplasma* symbionts in non-arthropod animals (Duperron et al., 2013; He et al., 2018). In insects, two phenotypes induced by inherited *Spiroplasma* have been studied in detail (reviewed in Anbutsu and Fukatsu, 2011; Ballinger

and Perlman, 2019). Firstly, *Spiroplasma* can enhance survival rates of its hosts under parasite or parasitoid attack. In *Drosophila neotestecea*, *Spiroplasma* inhibits growth and reproduction of the parasitic nematode *Howardula*, and thus reverses nematode-induced sterility (Jaenike et al., 2010). Further, *Spiroplasma* symbionts may greatly enhance *Drosophila* survival when attacked by parasitoids (J. Xie et al., 2014; Xie et al., 2010). Both protective phenotypes are mediated by *Spiroplasma* encoded RIP toxins (ribosome-inactivating proteins), which target the attackers' ribosomes (Ballinger and Perlman, 2017; Hamilton et al., 2015, 2014). These processes may be complementary to protection through competition between *Spiroplasma* and parasitoids for lipids in the host hemolymph (Paredes et al., 2016). Secondly, *Spiroplasma* kills males early in development in several species of *Drosophila*, planthoppers (Sanada-Morimura et al., 2013), ladybird beetles (Tinsley and Majerus, 2006), lacewings (Hayashi et al., 2016), and butterflies (Jiggins et al., 2000). The male-killing phenotype is also linked to a toxin: in *Drosophila melanogaster*, a *Spiroplasma* protein containing OTU (Ovarian Tumour-like deubiquitinase) and ankyrin domains was demonstrated to kill male embryos, and thus termed *Spiroplasma poulsonii* androcidin ("Spaid", Harumoto and Lemaitre, 2018).

Spiroplasma induced phenotypes were commonly observed to be dynamic compared with other symbiont traits, with repeated observation of phenotype change over relatively short time frames. For example, spontaneous loss of male killing was found in *Spiroplasma* symbionts of *Drosophila nebulosa* (Yamada et al., 1982), and *Drosophila willistoni* (Ebbert, 1991), and spontaneous emergence of non male-killing *Spiroplasma* in *Drosophila melanogaster* has occurred at least twice in a single culture (Harumoto and Lemaitre, 2018; Masson et al., 2020). In addition, *Spiroplasma* symbionts artificially transferred from their native host *Drosophila hydei* into *D. melanogaster* initially caused pathogenesis, but evolved

to become benign over just a few host generations (Nakayama et al., 2015). Genomic analysis

has further documented dynamic *Spiroplasma* evolution driven by viral proliferation (Carle et

al., 2010; Ku et al., 2013; Ye et al., 1996) and by extensive transfer of genetic material

between plasmids, and from plasmids to chromosomes (Joshi et al., 2005; Mouches et al.,

1984). Notably, plasmids in *Spiroplasma* commonly encode the systems that establish their

phenotypic effect on their host (Berho et al., 2006). Examples include Spaid (Harumoto and

Lemaitre, 2018) and RIP genes (Ballinger and Perlman, 2017), and rapid plasmid evolution

may therefore contribute to the high phenotypic evolvability of *Spiroplasma*. Furthermore,

elevated rates of evolution were found in various *Mycoplasma* species (Delaney et al., 2012;

Woese et al., 1984), which are pathogens closely related to *Spiroplasma* (Gupta et al., 2019).

Bacteria adapting to hosts often follow similar evolutionary trajectories, and both increased

mutational rates and proliferation of mobile genetic elements are commonly observed in

diverse symbiotic taxa (McCutcheon and Moran, 2011). Therefore, spontaneous loss of

Spiroplasma phenotypes and high genomic turnover could be explained by evolutionary

mechanisms common to all symbiotic taxa. Although independent findings discussed above

suggest that *Spiroplasma* symbionts may evolve quickly, this has never been quantified, and it

has remained unclear how substitution rates compare to those of other symbionts. However,

rates and patterns of evolutionary change may have important implications for *Spiroplasma*

evolutionary ecology, and for its potential use in biological control programmes (Clark and

Whitcomb, 1984; Schneider et al., 2019).

In this study, we systematically investigate rates and patterns of *Spiroplasma* evolution. To

this end, we employed two strains of *Spiroplasma poulsonii*: 1) *s*Hy is a natural associate of

Drosophila hydei (Ota et al., 1979) and confers protection against parasitic wasps (Xie et al.,

2010). We monitored its evolution over ~10 years in fly culture, and determined changes

through denovo reference genome sequencing and re-sequencing of multiple isolates. 2) *s*Mel

occurs naturally in *Drosophila melanogaster*, where it kills male offspring (Montenegro et al.,

2005) and protects from parasitoids (J. Xie et al., 2014). Using the previously established

complete genome (Paredes et al., 2015) and cell-free culture (Masson et al., 2018), we

re-sequenced isolates spanning ~2.5 years of evolution. This approach enabled tracing

Spiroplasma evolution on various levels: within-strain (<10 years divergence), between

strains, and through comparison with other *Spiroplasma* genomes, across a whole clade of

arthropod symbionts. It further allowed comparison between symbionts evolving in a host and

in axenic culture. We observed that rates of molecular evolution and chromosomal

rearrangements in *Spiroplasma poulsonii* are substantially faster than in other inherited

bacteria, and that this has resulted in markedly different genomic organisation in *s*Hy and

*s*Mel. Our work thus highlights *Spiroplasma poulsonii* as an amenable, highly evolvable, but

potentially unpredictable model insect symbiont.

Materials and Methods

Spiroplasma strains and genome sequencing

We documented the evolution of the *Spiroplasma poulsonii* strains *s*Hy and *s*Mel in fly hosts

and in cell free culture, respectively (Tab. 1). sHy is one of the *Spiroplasma* symbionts

naturally associated with *Drosophila hydei (Mateos et al., 2006)*. Our evolution experiments

covered ca. ten years and three variants of *s*Hy, all of which all originated from a single

isolate (Fig. 1). Over the course of the study, *Drosophila hydei* hosts were maintained on

standard corn meal agar diet (1% agarose, 8.5% sugar, 6% maize meal, 2% autolysed yeast,

0.25% nipagin), at 25°C, and a 12h:12h light:dark cycle. To isolate *Spiroplasma* *s*Hy-Liv

DNA from *D. hydei*, we followed a protocol by Paredes et al. (2015): We collected ca. 300 *D.*

hydei virgins and aged them for 4 weeks. The flies were then anesthetized with CO_2 and

pricked into the anterior sternopleurum using a sterile needle. Hemolymph was extracted via

centrifugation (10,000g, 5 minutes, at 4°C) through a 0.45μm filter (Corning Costar Spin-X).

To avoid hemolymph coagulation, we only extracted batches of 20 flies at a time and

immediately transferred extracted hemolymph into ice cold phosphate buffered saline solution

(1X PBS buffer; 137 mmol NaCl, 2.7 mmol KCl, 10 mmol Na2HPO4, 1.8 mmol KH2PO4).

Extracts of all batches were combined, and the PBS solution with *Spiroplasma* cells was

centrifuged (12,000g, 5 minutes, at 4°C). Supernatant PBS was discarded and the cell pellet

was subjected to DNA extraction using a phenol-chloroform protocol and subsequent ethanol

precipitation. DNA was eluted in water, and sequenced at the Centre for Genomics Research

at University of Liverpool. Libraries were prepared using the Nextera XT kit and sequenced

as 2x250bp run on an Illumina MiSeq sequencer.

DNA extraction for the *Spiroplasma* strain *s*Hy-Tx followed the protocol described above,

with the following modifications: Immediately after the piercing, 35-40 flies were placed into

0.5 ml microcentrifuge tubes previously pierced in the bottom, which was placed within a 1.5

ml microcentrifuge tube containing 20μl PBS, and centrifuged at 4,500g for 10 seconds to

collect hemolymph. DNA was recovered using a chloroform-ethanol procedure

(Supplementary protocol 1) and diluted in AE buffer (Qiagen). A pair-ended library was

constructed by Eureka Genomics (Hercules, CA). DNA was fragmented, end repaired, A'

tagged, ligated to adaptors, size-selected, and enriched with 25 cycles of PCR during which

an index was incorporated to the sample. Sample preparation was performed according to

Illumina's Multiplexing Sample Preparation Guide and Eureka Genomics' proprietary

158 method. The resulting library was subjected to Illumina HiSeq 2500 sequencing at the Texas

159 AgriLife Genomics and Bioinformatics Services Facility (College Station, TX).

160 **Tab. 1:** *Spiroplasma poulsonii* strains investigated in this study. Note that we follow the
161 naming scheme suggested by Ballinger & Perlman (2017).

Name used in the current study		Natural host	Establishment in fly or cell-free culture	DNA extracted from, date	Also known as
sHy	sHy-Tx12	*Drosophila hydei*	01/2004 (Mateos et al., 2006)	Hemolymph extract, 03/2012	Hap_1, TEN104-106
	sHy-Liv18a	*Drosophila hydei*	06/2008 (copy of sHy-Tx12)	Hemolymph extract, 04/2018	-
	sHy-Liv18b	*Drosophila hydei*	03/2014 (copy of sHy-Liv18A)	Hemolymph extract, 04/2018	-
sMel	sMel-Ug16	*Drosophila melanogaster*	10/2016 (Masson et al., 2018)	Cell free culture, 4 time points 2016–18 (Tab. 2)	MSRO-Uganda
	sMel-Br18	*Drosophila melanogaster*	07/1997 (Montenegro et al., 2000)	Whole fly, 03/2018	MSRO-RED42

162 **Fig. 1:** Overview on *Spiroplasma poulsonii* sHy symbionts sampled from *Drosophila*
163 *hydei* cultures. The evolution of three strains was investigated over a total timespan of
164 ten years. Blue dates denote splitting of *D. hydei* cultures, red dates mark
165 *Spiroplasma* extraction for sequencing.

sMel is a naturally occurring *Spiroplasma* symbiont of *D. melanogaster* best known for its male-killing phenotype (Harumoto and Lemaitre, 2018; Montenegro et al., 2000). Cell free culture of *s*Mel-Ug was recently accomplished (Masson et al., 2018), and we used a time series of this culture to investigate *s*Mel molecular rates of evolution outside of host tissues. The time series covered 29 months in total, and sequencing was performed at four time points (Table 2). Illumina DNA sequencing was performed at MicrobesNG (Birmingham, UK) using the "standard service". Culturing, passaging, and DNA extraction was performed as described in Masson et al. (2018).

Tab 2: Timeline for *Spiroplasma poulsonii s*Mel culturing and sequencing.

ID	Date of isolation for sequencing	Age (months)
HL	11/10/2016	0 (start of culture)
P14	09/02/2017	4
P57	23/05/2018	19
P79	25/03/2019	29

We further sequenced a single isolate of sMel-Br deriving from a strain established in 1997 (Tab. 1). For Illumina sequencing, 1-2 grams of whole body flies from *Spiroplasma*-positive *D. melanogaster* were collected for CTAB-Phenol based DNA extraction. Illumina libraries were prepared with Illumina Truseq DNA kit and sequenced on an Illumina NovaSeq-6000 S2 150 paired end flow cell by Texas AgriLife Genomics and Bioinformatics Services Facility (College Station, TX). For Oxford Nanopore MinIon sequencing, heads were first removed by immersing whole flies in liquid Nitrogen, and vigorous shaking on a metal sieve. The headless specimens were immediately subjected to phenol-chloroform extraction. Nanopore library preparation was performed using the Ligation Sequencing kit 1D (SQK-LSK108), following precautionary measures to ensure long read lengths. The library was then sequenced on MinION sequencing device (MIN-101B) with an SpotON Flow Cell

Mark 1 (R9.5- FLO-MIN107.1). Base Calling was performed using Albacore version 2.0

(Oxford Nanopore, Oxford, UK) and the reads were processed with Porevhop version 0.24

(Wick et al., 2017a) to remove adapter sequences. A draft assembly of sMel-Br was created

using Illumina and Nanopore reads and Unicycler version 0.4.7 (Wick et al., 2017b). The

assembly was fragmented, but did contain 3 circular contigs with sequence similarity to other

Spiroplasma contigs.

*s*Hy Reference genome sequencing and annotation

In order to reconstruct a high quality reference genome for *s*Hy, we extracted DNA from 30

Drosophila hydei virgin females carrying the symbiont. To limit the impact of potentially

heterogeneous *Spiroplasma* populations within the flies, we used F3 individuals from a single

isofemale line for DNA extraction. The flies were anesthetized at -20°C, washed with 70%

ethanol, and briefly dried. The flies were then homogenized in G2 lysis buffer from the

Qiagen Genomic DNA Buffer Set using a 1ml dounce tissue grinder (DWK Life Science),

and DNA was extracted from the homogenate using a Qiagen Genomic-Tip 20/G following

the protocol modifications from Miller et al. (2018). A total of 4μg DNA was then used in the

"1D gDNA long reads without BluePippin protocol" library preparation protocol with the 1D

Ligation Sequencing Kit (SQK-LSK108, Oxford Nanopore, Oxford, UK). Sequencing was

performed on a MinION sequencer (Oxford Nanopore, Oxford, UK) using a single

FLO-MIN106 R9 flow cell for 48 hours, and the raw Nanopore signals were base-called

using Albacore version 2.3.1.

All reads passing the quality checks were used to create an assembly with Unicycler version

0.4.7 (Wick et al., 2017b). The resulting assembly contained 6 circular chromosomes (~15 kb

− ~1.6 Mbp) with high similarity to the previously published *s*Mel genome as determined by

BLAST+ version 2.9 searches (Camacho et al., 2009) in Bandage (Wick et al., 2015). We also

created a hybrid assembly with Unicycler using all long reads and the *s*Hy-Liv18a Illumina

reads. The hybrid assembly was more fragmented, but did contain 3 contigs which spanned

almost the entire 1.6 Mbp circular contig of the long-read only assembly. Because hybrid

assemblies are typically more accurate than assemblies based on long reads only (De Maio et

al., 2019), we corrected the long read assembly using the hybrid contigs. This was achieved

by calling differences between the assemblies with the 'dnadiff' function implemented in

MUMmer version 3.18 (Kurtz et al., 2004), and correcting the assembly accordingly. Finally,

Pilon version 1.23 (Walker et al., 2014) was used to polish the corrected assembly, again

using the *s*Hy-Liv18a Illumina reads. After nine rounds of polishing with Pilon, no further

improvements could be observed and the assembly was considered final.

Rates of molecular evolution in *s*Hy and *s*Mel

We determined changes in *s*Hy and *s*Mel chromosomes over time by employing the Snippy

pipeline version 4.1.0 (Seemann, 2015). We used our newly assembled *s*Hy genome and the

previously sequenced *s*Mel genome (Masson et al., 2018) as references, annotated using the

Prokka pipeline version 1.13 with standard parameters and the *Spiroplasma* genetic code

(translation table 4, Seemann, 2014). *s*Hy and *s*Mel variants determined by Snippy were

filtered to only include positions which were covered by all *s*Hy or all *s*Mel Illumina libraries

with at least 5x coverage, respectively. Further, to compare rates of molecular evolution

between *Spiroplasma* and other microbial species, we counted the number of changes in third

codon positions of coding sequences (CDS). For both strains, we only considered CDS that

were entirely covered by sequencing depth of at least 5x in all Illumina libraries. Rates of

molecular evolution were calculated by: number of observed changes at 3rd codon positions /

number of considered 3rd codon positions / time in years over which the changes were

observed. For *s*Hy, using *s*Hy-Liv18a as reference, we calculated two rate estimates, one

based on comparison with *s*Hy-Liv18b, and one by comparing *s*Hy-Tx12 to the reference

(Fig. 1). For *s*Mel, the hemolymph extract from which the culture was established was used

as the reference ('HL', Tab. 2), and three rate estimates were calculated based on the changes

observed in the three sampled time points of the culture (Tab. 2).

<u>Comparative genomics</u>

We first compared the newly assembled *s*Hy genome with the most closely related, fully

sequenced *Spiroplasma* genome, *s*Mel. Both genomes were annotated using the Kyoto

Encyclopedia of Genes and Genomes (KEGG) and BlastKOALA (Kanehisa et al., 2016,

2006) and orthologous protein sequences determined using OrthoFinder version 2.2.3 (Emms

and Kelly, 2015). Synteny between the genomes was assessed through aligning the genomes

with Minimap2 version 2.15 (Li, 2018). Genome degradation was investigated by

documenting the number of unique KEGG numbers for *s*Hy and *s*Mel, all of which were

verified by BLAST+ searches. For each strain, we considered genes to be likely

pseudogenized or truncated if the longest CDS within an orthogroup spanned at most 60% of

the CDS length from the other strain.

Secondly, we compared *s*Hy with 17 *Spiroplasma* genomes from the

Citri-Chrysopicola-Mirum clade (Table S1, Supplementary material for full list of names and

accession numbers). All genomes were annotated using Prokka version 1.4.0, and orthology

between predicted CDS was established using OrthoFinder. We extracted single-copy

orthologs present in all strains, and aligned each locus separately with the L-INS-i method

implemented in Mafft version 7.450 (Katoh and Standley, 2013). Recombination was

evaluated with pairwise homoplasy index and window sizes of 100, 50, and 20 amino acids

for each locus (Bruen et al., 2006), and any locus showing evidence for recombination was

excluded. We used IQ-TREE version 1.6.10 (Nguyen et al., 2015) to reconstruct a core

genome phylogeny from the remaining 96 loci (covering 26,019 amino acid positions). Each

locus was treated as a separate partition with a distinct evolutionary rate (Chernomor et al.,

2016), and optimal models and number of partitions was estimated with IQ-TREE

(Kalyaanamoorthy et al., 2017; Lanfear et al., 2012). Branch support was assessed using

1,000 replicates of ultrafast bootstraps (Hoang et al., 2018), and approximate likelihood ratio

test (Guindon et al., 2010) in IQ-TREE.

For all genomes, insertion sequence elements were annotated with Prokka. Prophage regions

were predicted with PhiSpy version 3.7.8, which uses typical prophage features (e.g., gene

length, strand directionality, AT/GC skews, insertion points, Akhter et al., 2012) for its

predictions (Akhter et al., 2012). We also used the Phaster web server, which uses sequence

similarities and other criteria to predict prophages (Arndt et al., 2016). We screened for the

presence of toxin genes implied in protective phenotypes (Ballinger and Perlman, 2017;

Hamilton et al., 2015) and male killing (Harumoto and Lemaitre, 2018). To this end, we

downloaded UniProt sequence alignments from the Pfam database (Finn et al., 2016) for the

protein families RIP (PF00161, ribosome inactivating protein) and OTU (PF02328, OTU-like

cysteine protease), respectively. We used these alignments as databases for searches with

HMMER version 3.2.1 (Eddy, 2011), and protein sequences from all genomes as queries.

Domain architecture of all matching proteins was then determined using PfamScan with an

e-value of 0.001 (Madeira et al., 2019), SignalP 5.0 (Almagro Armenteros et al., 2019), and

TMHMM 2.0 (Krogh et al., 2001). RIP domains showed a high degree of divergence, and

were therefore aligned to the reference RIP HMM profile using hmmalign from the HMMER

software package. From this alignment, positions present in fewer than three sequences were

removed, and a phylogeny reconstructed with IQ-TREE. OTU domains were aligned using

MAFFT, and a phylogeny reconstructed with IQ-TREE.

Thirdly, we determined plasmid synteny across *Spiroplasma poulsonii* strains. In addition to

the plasmids of *s*Mel and *s*Hy, we also included a plasmid of *s*Neo and all circular contigs

from our *s*Mel-Br assembly. Plasmids were annotated using Prokka version 1.13 with a

protein database composed of all plasmid proteins available on NCBI GenBank. Plasmid

alignment and visualisation was performed with AliTV version 1.06 (Ankenbrand et al.,

2017).

<u>Data visualisation</u>

Figures were prepared in R version 3.6.2 (R Core Team, 2016) using the packages 'ape'

(Paradis and Schliep, 2019), 'cowplot' (Wilke, 2019), 'ggalluvial' (Brunson, 2019), 'ggplot2'

(Wickham, 2009), 'ggtree' (Yu et al., 2017), and 'ggridges' (Wilke, 2020). Phylogenetic trees

and domain architecture of toxin loci were visualised with EvolView version 3.0

(Subramanian et al., 2019).

Results

<u>Rates and patterns of evolutionary change in *Spiroplasma poulsonii*</u>

The *Spiroplasma poulsonii s*Hy reference genome comprises a single circular chromosome

(1,625,797bp) and five circular contigs (23,069bp – 15,710bp) with sequence similarities to

plasmids from other *Spiroplasma*. The chromosome contains 1,584 predicted coding

sequences (CDS), a single ribosomal RNA cluster, and 30 tRNAs. Overall, genome content

and metabolic capacities are highly similar to the previously sequenced, very closely related

*s*Mel genome (Paredes et al., 2015), and a detailed comparison follows further below.

To estimate rates of molecular evolution in *Spiroplasma poulsonii*, we measured

chromosome-wide changes in coding sequences of *Spiroplasma* from fly hosts (*s*Hy) and

axenic culture (*s*Mel) over time. Our estimates for *s*Hy and *s*Mel are overlapping and range

from $6.4 \times 10^{-6} - 1.8 \times 10^{-5}$ changes per position per year. This rate exceeds the rate reported for

other symbiotic bacteria such as *Wolbachia* and *Buchnera* by at least two orders of magnitude

(Fig. 2). Our estimate overlaps with rates calculated from some of fastest evolving human

pathogens (e.g., *Enterococcus faecium* and *Acinetobacter baumannii*), and with evolutionary

rates observed in the poultry pathogen *Mycoplasma gallisepticum* (which is closely related to

Spiroplasma). Indeed, *Spiroplasma* substitution rates fall at around the lower estimates for

viruses (Fig. 2).

Fig 2. Comparison of estimated evolutionary rates across various microbes. For details
on how evolutionary rates were calculated, please refer to the original studies and the
Materials and Methods section.

Over ~10 years of evolution, we observed a similar absolute number of variants on the

chromosome (~1.6 Mbp) and plasmids (~0.1 Mbp) of *s*Hy, i.e., relatively more changes on

the plasmids (Fig. 3, Table S2, supplementary material). Most variants in CDS affected

hypothetical proteins, but were enriched in prophage-associated loci and adhesin related

proteins (ARP, Fig. 3). Overall, *s*Hy variants in CDS were about equally often found to be

synonymous as non-synonymous (Fig. 3) and changes were biased towards GC > AT

substitutions (Fig. S1, Supplementary material). The changes in *s*Mel over ~2.5 years in

culture affected only 15 different CDS in total, of which four were ARPs, and three

lipoproteins (Table S2, supplementary material).

Fig. 3: Overview of all observed variants in *s*Hy (100% = 570 variants). All annotations
(Type, Feature, Product, Effect) are taken from results of the snippy pipeline.

Comparing the genomes of *s*Hy and *s*Mel revealed a notable contrast between the high degree

of nucleotide sequence identity on the one hand, and striking structural and gene content

differences on the other hand. In 604 single copy orthologues shared between the genomes,

average nucleotide identity was >99.5%. However, compared with *s*Mel, *s*Hy is reduced in

gene content (1,584 vs 2,388 genes), chromosome size (1.61 vs 1.88 Mbp), and coding

density (65% vs 82%). Further, *s*Hy contains fewer predicted insertion sequences (9 vs 89 in

*s*Mel) and intact prophages as predicted with Phaster (*s*Mel: 16 such regions spanning 426

Kb; *s*Hy: 2 regions, 14 Kb), but not PhiSpy (Fig. S2, Supplementary material). Both *s*Mel and

*s*Hy have a number of missing or truncated (i.e., potentially pseudogenized) genes when

compared with each other, but the level of genomic deterioration is higher in *s*Hy, and covers

a range of different genes (Fig. 4a, Table S3, supplementary material). Notable pseudogenised

or absent loci in *s*Hy include parts of the phosphotransferase system, in particular the loci

required for the uptake of N-Acetylmuramic Acid and N-acetylglucosamine (see Table S3,

Supplementary material for a full list).

Strikingly, there is evidence for a history of extensive chromosomal rearrangements since the

last common ancestor of *s*Hy and *s*Mel. The levels of synteny between the strains is much

lower than expected at the observed sequence similarity (Fig. 4b). Similarly, a comparison of

plasmids across different *Spiroplasma poulsonii* strains (*s*Hy, *s*Mel-Ug, *s*Mel-Br, and *s*Neo)

revealed similar gene content across the plasmids of different strains, but large differences in

arrangement and number (Fig. S3, Supplementary material).

Fig 4: Comparison of *s*Hy and *s*Mel. **a)** Number of genes with KEGG annotation that are present in one strain, but absent or truncated in the other. See Table S3, supplementary material for a complete list. **b)** Synteny as determined with minimap2.

Genome and toxin evolution in the genus *Spiroplasma*

Comparing *s*Hy with other sequenced strains of the *Spiroplasma* clades Citri, Poulsonii, Chrysopicola, and Mirum showed dynamic genome evolution within this genus of symbionts (Fig. 5). All investigated spiroplasmas have reduced genomes (~1.1–1.9Mbp), and the poulsonii strains *s*Hy, *s*Mel and *s*Neo are among the strains with the largest main chromosomes (Fig. 5). Plasmid numbers range from 0–5, and *s*Hy has the highest number of plasmids of any of the investigated strains, although plasmid number is unclear for some of the genomes in draft status, and *S. citri* may have seven plasmids (Saillard et al., 2008). Prophage regions of varying sizes were predicted in all of the genomes (despite the lack of clear homologues to viral sequences in some of these, Ku et al., 2013), and strains with especially high numbers of prophage associated loci show lower median CDS length when compared to strains with very small predicted prophage regions. Further, in the bigger *Spiroplasma* genomes, the number of insertion sequences was typically higher (Fig. 5). The comparison indicates that *s*Hy may be more degraded than its closest relatives *s*Neo and

*s*Mel, with a smaller chromosome size, fewer predicted CDS and mobile genetic elements,

and the lowest coding density of these three (Fig. 5).

Fig 5: Genome properties of *Spiroplasma* strains from Citri, Poulsonii, Chrysopicola, and
Mirum clades. Genome size and plasmid number was taken from the literature (Table S1,
supplementary material). Coding density, CDS length and number of IS elements was
determined with Prokka, and proportion of prophage was estimated with PhiSpy.
Spiroplasma phylogeny is based on partitioned maximum likelihood analysis of 96
concatenated single copy genes present in all of the genomes (26,019 amino acid
positions). All nodes were maximally supported by UFB and aLRT.

Spiroplasmas' most striking phenotypes in insects have been mechanistically linked to toxin

genes. For example, ribosome inactivating proteins (RIP) may protect *Spiroplasma* hosts by

cleaving ribosomal RNA of parasites and parasitoids (Hamilton et al., 2015). Five RIP loci

are present in *s*Mel (RIP1–5, of which 3–5 are almost identical copies). As expected, the

protective *s*Hy also encodes RIPs, but only has a single orthologue for RIP1. However, it

contains an additional RIP gene in three copies that appears to be absent in *s*Mel, and one of these copies also contains ankyrin repeats (Fig. S4, Supplementary material).

Further, the male killing phenotype of *s*Mel was recently established to be caused by *Spiroplasma* androcidin (Spaid), and both ankyrin repeats and a deubiquitinase domain (OTU) of this gene are necessary to induce male killing (Harumoto and Lemaitre, 2018). HMMER searches using OTU domain profiles revealed a number of *Spiroplasma* loci similar to Spaid (Fig. 6) which however lack its characteristic domain composition. For example, *s*Hy encodes three loci similar to Spaid: one lacks a signal peptide, one has no ankyrin repeats, and another encodes an epsilon-toxin like domain in addition to OTU (Fig. 6). Other bacterial loci with notable similarities to the *Spiroplasma* OTU could only be detected in the symbiotic taxa *Rickettsia* and *Wolbachia*. In the phylogenetic reconstruction of OTU domains rooted with the eukaryotic sequences (from the ciliate *Stentor coeruleus*), *Rickettsia* and *Wolbachia* loci are nested within the *Spiroplasma* sequences. This topology suggests lateral gene transfer of OTU domain containing proteins from *Spiroplasma* to the other intracellular taxa, although differences in genetic code between *Spiroplasma* and other bacteria likely limit the probability for transfer of functional protein coding genes.

Fig. 6: Spaid-like proteins in *Spiroplasma* and other symbionts. Maximum likelihood tree was reconstructed from an alignment of OTU domains (147 amino acid positions), and domain predictions are based on PfamScan, SignalP, and TMHMM. Abbreviations (corresponding to PFAM domains): Ank - ankyrin repeats, OTU - Ovarian tumor like deubiquitinase, ETX_MTX2 - Clostridium epsilon toxin ETX/Bacillus mosquitocidal toxin MTX2, Tox-SGS - Salivary gland secreted protein domain toxin, SEC-C - SEC-C nucleic acid binding domain.

Discussion

Spiroplasma, an exceptionally fast evolving symbiont

The substitution rates we observed in *Spiroplasma* symbionts are among the highest reported for any bacteria (Fig. 1). In a study comparing genome-wide evolutionary rates in bacterial human pathogens (Duchêne et al., 2016), most taxa showed considerably slower rates, and our estimate of *Spiroplasma* evolutionary rates overlaps with substitution rates of RNA viruses (Holmes, 2009). A number of reasons have been proposed for elevated substitution rates in bacterial symbionts and pathogens with small genome sizes: Firstly, host-associated microbes may acquire essential metabolic intermediates from their hosts and thus need not synthesise them (Moran, 2003). Obsolete metabolic genes subsequently accumulate now neutral substitutions, resulting in pseudogenization. Secondly, reduced population sizes of

symbionts, together with bottlenecks at transmission events limit purifying selection of

deleterious mutations (Moran, 1996) and mobile genetic elements (Lynch, 2006) – both in

turn create further genomic deterioration. Thirdly, genomes of host-associated microbes are

AT rich (Rocha and Danchin, 2002), which makes introduction of errors through replication

slippage more likely (Moran et al., 2009). Fourthly, many symbionts lack DNA repair genes

(Moran et al., 2008), which may result in increased substitution rates.

All of these points very likely apply for *Spiroplasma poulsonii*: its metabolic capacities are

reduced compared with free living microbes, it has a high proportion of pseudogenes and

prophages, very high AT content, and lacks several DNA repair genes. However, many

symbiotic bacteria show similar trends of genome evolution (Burger and Lang, 2003), and it

is therefore somewhat puzzling that *Spiroplasma* stands out with this exceptionally high

substitution rate. Loss of DNA repair genes has been shown to co-occur with elevated

mutation rates in intracellular bacteria (Itoh et al., 2002; Koonin et al., 1996; Massey, 2008),

and may explain our observations. The loci encoding the mismatch repair proteins *mutS* and

mutL are universally lacking in *Spiroplasma*, but are present in the slower evolving symbionts

*Wolbachia w*Mel, and *Buchnera aphidicola* (Richardson et al., 2012; Fig. S6, supplementary

material, Shigenobu et al., 2000; Wu et al., 2004). In *E. coli*, hypermutator strains that have

lost such loci have a ~10–300 fold increase of spontaneous mutation rates (LeClerc et al.,

1996; Lee et al., 2012), which is comparable to the ~two orders of magnitude difference

between substitution rates of *Buchnera* and *Spiroplasma* (Fig. 2). Like *Spiroplasma*,

Mycoplasma universally lacks the mismatch repair loci *mutS*, *mutL*, and *mutH* (Carvalho et

al., 2005; Chen et al., 2012). In line with this observation, elevated evolutionary rates were

hypothesized for mycoplasmas (Woese et al., 1984), and in our comparison, *M. gallisepticum*

appears to be the only bacterial taxon with substitution rates similar to *Spiroplasma* (Delaney

et al., 2012). Absence of DNA mismatch repair pathway may thus be ancestral to Entomoplasmatales (Spiroplasmatacea + Entomoplasmataceae) and contribute to the dynamic genome evolution across this taxon (Lo et al., 2016; Rocha and Blanchard, 2002).

Notably, substitution rates often appear larger when estimated over brief time periods (Ho et al., 2005), and this may have biased our estimates. Duchene et al. (2016) found that substitution rates measured over 10 years can be up to one order of magnitude larger than those measured over 100 years. In agreement with such bias, we found more variants in our *s*Mel culture after 19 months than for the 29 months isolate (Table S2, supplementary material). The back mutations that likely have happened between 19 and 29 months of our *s*Mel culture would go unnoticed when comparing genomes over larger time scales. More generally, it is difficult to estimate divergence times between *Spiroplasma* strains. The substitution rates estimated in *s*Hy and *s*Mel would suggest the two strains have diverged 1,120–2,260 years ago. However, this estimate is unreliable due to a number of factors that we cannot control for. For example, the number of generations per year for *Drosophila* hosts of *Spiroplasma* differ depending on species and location, and is expected to be lower in the wild compared with laboratory strains. Further, *Spiroplasma* may move between species, for example via ectoparasitic mites (Jaenike et al., 2007). A number of partially sympatric *Drosophila* species carry similar *Spiroplasma* strains (*D. hydei*, *D. melanogaster*, *D. willistoni*, *D. neotestacea*) (Haselkorn, 2010), and with our data it is impossible to determine the number and direction of potential *Spiroplasma* transfers between these species.

Reductive genome evolution in *Spiroplasma*

According to a widely supported model of genome reduction in symbiotic bacteria (McCutcheon and Moran, 2011), the first stages of host restriction involve accumulation of

pseudogenes and mobile genetic elements, chromosomal rearrangements, increased

substitution rates, and excess deletions. Advanced stages of adaptation to a host are

accompanied by further genome shrinkage, and purging of mobile genetic elements, which

overall result in more stable chromosomes. Using these characteristics, different levels of

genome reduction are apparent in the investigated *Spiroplasma* genomes, across which

genome sizes correlate positively with number of mobile genetic elements (plasmids,

prophages, insertion sequences, Fig. 5). On this spectrum, *S. syrphidicola* and *S. chrysopicola*

have the most reduced genomes, without homologues of prophage loci, and high levels of

synteny between the two strains (Ku et al., 2013). It was therefore argued that phages have

likely invaded *Spiroplasma* only after the split of the Syrphidicola and Citri+Poulsonii clades

(Ku et al., 2013). However, several observations support the conclusion that the reduced

Spiroplasma genomes have lost their prophage regions, rather than never having been

susceptible to viruses: 1) Genomes lacking prophage loci are also found within the

Citri+Poulsonii clade (TU-14 & NBRC100-390, Fig. 5), illustrating that prophage loss after

viral invasion is possible in *Spiroplasma*. 2) Strains lacking prophages show other hallmarks

of reduced symbiont genomes (small size, high coding density, lack of plasmids and

transposons, Fig. 5), which is in line with the model of genome reduction discussed above. 3)

Prophage prediction using an algorithm that does not mainly rely on sequence similarities to

known viral sequences (PhiSpy, Akhter et al., 2012) identified at least very small prophage

regions in all of the investigated genomes. In summary, while the genomic instability in

Spiroplasma genomes is likely driven by prophages (Ku et al., 2013; Lo et al., 2013; Ye et al.,

1996), several lineages display reduced genomes from which the phages have largely been

purged.

Using the model of genome reduction during host adaptation introduced above, *Spiroplasma poulsonii*'s genome characteristics suggest that host restriction has happened relatively recently. Although very closely related (99.5% sequence identity) and found in very similar hosts, *s*Mel and *s*Hy differ markedly in coding density, genome size, and proportion of prophage regions. Interestingly, PHASTER predicted 426 Kb of intact phage regions in *s*Mel, but only 14 Kb for *s*Hy (Fig. S2, supplementary material). Both phage proliferation in *s*Mel, and prophage loss in *s*Hy could have contributed to this. However, when using the similarity agnostic tool PhiSpy, the differences in predicted prophage regions were less pronounced (Fig. 5). This observation is compatible with degradation and/or pseudogenization of prophage regions in *s*Hy, which would lead to reduced sequence similarity to viral loci (and thus reduced detectability by PHASTER), but not entirely blur prophage characteristics employed by PhiSpy (e.g., gene length, strand directionality, AT/GC skews, insertion points, Akhter et al., 2012). Since the split of the lineages, *s*Hy has not only lost prophage regions and insertion sequences compared with *s*Mel, but also several genes that are often lost in adaptation to a host, such as parts of the phosphotransferase system for the uptake of carbohydrates, one tRNA locus, and *uvrD* which plays a role in mismatch repair, and nucleotide excision repair (Table S2, supplementary material). Using genome degradation as a proxy for host adaptation, our findings collectively suggest that host adaptation is more advanced in *s*Hy than in *s*Mel. This may be a result of the fitness benefits associated with *s*Hy under parasitoid pressure, and the absence of detectable costs of *s*Hy infection in *Drosophila hydei* (Osaka et al., 2013; Jialei Xie et al., 2014; Xie et al., 2010). However, the *Spiroplasma* symbiont of *Drosophila neotestacea* *s*Neo is also protective, does not cause obvious fitness costs (Jaenike et al., 2010), but has a less reduced genome (Fig.5, Ballinger and Perlman, 2017). Further, we cannot exclude a pathway in which the genome reduction in *s*Hy was

mainly driven by stochastic effects or even by adaptation to laboratory conditions, as we have

not investigated contemporary *s*Hy from wild *D. hydei* populations.

<u>Implications for *Spiroplasma* evolutionary ecology</u>

Spiroplasma evolutionary ecology shows several parallels to that of the most widely

distributed arthropod symbiont *Wolbachia*: both symbionts are found in a range of different

hosts (Duron et al., 2008), have the ability to invade novel hosts (Conner et al., 2017;

Haselkorn et al., 2009), may confer protection (Teixeira et al., 2008; Xie et al., 2010) but also

kill males (Hurst and Jiggins, 2000), and can spread across host populations swiftly

(Cockburn et al., 2013; Turelli and Hoffmann, 1991). However, there are also pronounced

differences between the symbionts: *Spiroplasma* rarely reaches the high infection frequencies

often observed in *Wolbachia* (Hilgenboecker et al., 2008; Watts et al., 2009), and is arguably

found in a more diverse host range that encompasses arthropods and plants (Regassa and

Gasparich, 2006), but also molluscs (Duperron et al., 2013), echinoderms (He et al., 2018),

and cnidarians (Viver et al., 2017). Further, *Wolbachia* show greatly reduced spontaneous

mutation rates compared with *Spiroplasma*, likely caused by a more complete set of DNA

repair genes (Fig. S5, supplementary material).

In theory, fast evolutionary rates should enable *Spiroplasma* to adapt to novel hosts quickly

(i.e., to reduce pathogenicity, and to maximise vertical transmission efficiency), and

experimental studies have found high horizontal transmission efficiency of *Spiroplasma*

(Haselkorn et al., 2013; Haselkorn and Jaenike, 2015; Nakayama et al., 2015). Consistent

with this, we found that genes implicated in host-symbiont compatibility and virulence (e.g.,

adhesion-related proteins) have evolved especially fast in our evolution experiments (Fig. 2).

In addition, we documented dynamic evolution and turnover of toxin loci, which are

important for host fitness and symbiont compatibility (Fig. 6, Fig. S4, supplementary material). Such genomic flexibility may contribute to *Spiroplasma*'s broader host range when compared with *Wolbachia*. However, elevated evolutionary rates also make deleterious changes more likely, and, in the absence of strong selection, may result in faster loss of symbionts (Jaenike, 2012). This may explain the generally low abundance of *Spiroplasma* symbionts (Haselkorn, 2010), which seems to increase only when carrying the symbiont is associated with a large fitness benefit (Jaenike et al., 2010). In contrast, virtually identical *Wolbachia* strains ("superspreaders") are found in many different host species at very high frequencies (Turelli et al., 2018) – demonstrating that stationary genomes may be evolutionary advantageous. In summary, the nature of *Spiroplasma* genomic evolution likely contributes to its peculiar evolutionary ecology.

From a practical perspective, *Spiroplasma poulsonii* has many features of a desirable model for symbiont-host interactions: fast rates of evolution make it more likely that adaptation and spontaneous changes in phenotypes can be determined over short time scales, as has been observed previously (Harumoto and Lemaitre, 2018; Masson et al., 2020; Nakayama et al., 2015). On the other hand, fast evolutionary changes make experiments less predictable, and because stochastic effects become more pronounced, links of genomic changes with phenotypes may be obscured. Further, the generalisability of experimental results may be limited for extremely fast evolving symbionts. Our findings therefore also underline the importance of regular validation of laboratory symbiont strains through re-sequencing.

Author contributions

Designed and conceived project: MG, GDDH, MM. Experimental passage of *Spiroplasma* in vivo: MG, JG, GDDH, MM. Experimental passage of *Spiroplasma* in vitro: FM, BL. Symbiont DNA purification and sequencing: MG, PR, SS, RA, HMM, FM. Assembly and annotation: MG, HMM, PR, RA. Molecular evolution analyses: MG. Wrote first draft of paper: MG, MM, GDDH. All authors contributed in revising the manuscript and have read and approved the final manuscript.

Acknowledgements

MG has received funding from the European Union's Horizon 2020 Research and Innovation Program under Marie Sklodowska-Curie grant agreement 703379 and from EMBO through a long term fellowship (ALTF 48-2015) co-funded by the European Commission (LTFCOFUND2013, GA-2013-609409). Sequencing was supported by an award from the University of Liverpool Technology Directorate Voucher Scheme, and by a Texas A&M Genomics Seed Grant.

Data accessibility

Raw reads for sHy-Tx can be found under NCBI BioProject Accession PRJNA274591 (Biosample SAMN08102256, SRA SRR6326389). Reads from sMel-Br are stored under PRJNA507275 (SAMN10488582). All other reads are available under NCBI BioProject PRJNA640980. Alignments and supplementary protocol are available from Zenodo under the DOI 10.5281/zenodo.3903208.

References

Akhter S, Aziz RK, Edwards RA. 2012. PhiSpy: a novel algorithm for finding prophages in bacterial genomes that combines similarity- and composition-based strategies. *Nucleic Acids Res* **40**:e126.

Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, von Heijne G, Nielsen H. 2019. SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nat Biotechnol* **37**:420–423.

Anbutsu H, Fukatsu T. 2011. *Spiroplasma* as a model insect endosymbiont. *Environ Microbiol Rep* **3**:144–153.

Ankenbrand MJ, Hohlfeld S, Hackl T, Förster F. 2017. AliTV—interactive visualization of whole genome comparisons. *PeerJ Computer Science* **3**:e116.

Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* **44**:W16–21.

Ballinger MJ, Perlman SJ. 2019. The defensive *Spiroplasma*. *Curr Opin Insect Sci* **32**:36–41.

Ballinger MJ, Perlman SJ. 2017. Generality of toxins in defensive symbiosis: Ribosome-inactivating proteins and defense against parasitic wasps in *Drosophila*. *PLoS Pathog* **13**:e1006431.

Berho N, Duret S, Danet J-L, Renaudin J. 2006. Plasmid pSci6 from *Spiroplasma citri* GII-3 confers insect transmissibility to the non-transmissible strain *S. citri* 44. *Microbiology* **152**:2703–2716.

Bruen TC, Philippe H, Bryant D. 2006. A simple and robust statistical test for detecting the presence of recombination. *Genetics* **172**:2665–2681.

Brunson JC. 2019. ggalluvial: Alluvial Plots in "ggplot2."

Burger G, Lang BF. 2003. Parallels in genome evolution in mitochondria and bacterial symbionts. *IUBMB Life* **55**:205–212.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* **10**:421.

Carle P, Saillard C, Carrère N, Carrère S, Duret S, Eveillard S, Gaurivaud P, Gourgues G, Gouzy J, Salar P, Verdin E, Breton M, Blanchard A, Laigret F, Bové J-M, Renaudin J, Foissac X. 2010. Partial chromosome sequence of *Spiroplasma citri* reveals extensive viral invasion and important gene decay. *Appl Environ Microbiol* **76**:3420–3426.

Carvalho FM, Fonseca MM, Batistuzzo De Medeiros S, Scortecci KC, Blaha CAG, Agnez-Lima LF. 2005. DNA repair in reduced genome: the *Mycoplasma* model. *Gene* **360**:111–119.

Chen L-L, Chung W-C, Lin C-P, Kuo C-H. 2012. Comparative analysis of gene content evolution in phytoplasmas and mycoplasmas. *PLoS One* **7**:e34407.

Chernomor O, von Haeseler A, Minh BQ. 2016. Terrace Aware Data Structure for Phylogenomic Inference from Supermatrices. *Syst Biol* **65**:997–1008.

Clark TB, Whitcomb RF. 1984. Pathogenicity of mollicutes for insects: possible use in biological control. *Ann Microbiol* **135A**:141–150.

Cockburn SN, Haselkorn TS, Hamilton PT, Landzberg E, Jaenike J, Perlman SJ. 2013. Dynamics of the continent-wide spread of a *Drosophila* defensive symbiont. *Ecol Lett* **16**:609–616.

Conner WR, Blaxter ML, Anfora G, Ometto L, Rota-Stabelli O, Turelli M. 2017. Genome comparisons indicate recent transfer of *w*Ri-like *Wolbachia* between sister species *Drosophila suzukii* and *D. subpulchrella*. *Ecol Evol* **7**:9391–9404.

Corbin C, Heyworth ER, Ferrari J, Hurst GDD. 2017. Heritable symbionts in a world of varying temperature. *Heredity* **118**:10–20.

Delaney NF, Balenger S, Bonneaud C, Marx CJ, Hill GE, Ferguson-Noel N, Tsai P, Rodrigo A, Edwards SV. 2012. Ultrafast Evolution and Loss of CRISPRs Following a Host Shift in a Novel Wildlife Pathogen, *Mycoplasma gallisepticum*. *PLoS Genetics* **8**:e1002511.

De Maio N, Shaw LP, Hubbard A, George S, Sanderson ND, Swann J, Wick R, AbuOun M, Stubberfield E, Hoosdally SJ, Crook DW, Peto TEA, Sheppard AE, Bailey MJ, Read DS, Anjum MF, Walker AS, Stoesser N, On Behalf Of The Rehab Consortium. 2019. Comparison of

long-read sequencing technologies in the hybrid assembly of complex bacterial genomes. *Microb*
Genom **5**:e000294.
Douglas AE. 2009. The microbial dimension in insect nutritional ecology. *Funct Ecol* **23**:38–47.
Drew GC, Frost CL, Hurst GDD. 2019. Reproductive Parasitism and Positive Fitness Effects of
Heritable Microbes. *eLS* **2**:1–8.
Duchêne S, Holt KE, Weill F-X, Le Hello S, Hawkey J, Edwards DJ, Fourment M, Holmes EC. 2016.
Genome-scale rates of evolutionary change in bacteria. *Microb Genom* **2**:e000094.
Duperron S, Pottier M-A, Léger N, Gaudron SM, Puillandre N, Le Prieur S, Sigwart JD, Ravaux J,
Zbinden M. 2013. A tale of two chitons: is habitat specialisation linked to distinct associated
bacterial communities? *FEMS Microbiol Ecol* **83**:552–567.
Duron O, Bouchon D, Boutin S, Bellamy L, Zhou L, Engelstädter J, Hurst GDD. 2008. The diversity
of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC Biol* **6**:27.
Dyson EA, Kamath MK, Hurst G. 2002. *Wolbachia* infection associated with all-female broods in
Hypolimnas bolina (Lepidoptera: Nymphalidae): evidence for horizontal transmission of a
butterfly male killer. *Heredity* **88**:166–171.
Ebbert MA. 1991. The interaction phenotype in the *Drosophila willistoni-Spiroplasma* symbiosis.
Evolution **45**:971–988.
Eddy SR. 2011. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**:e1002195.
Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biol* **16**:157.
Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi M,
Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016. The Pfam protein families database:
towards a more sustainable future. *Nucleic Acids Res* **44**:D279–85.
Gasparich GE. 2002. Spiroplasmas: evolution, adaptation and diversity. *Front Biosci* **7**:d619–d640.
Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms and
methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0.
Syst Biol **59**:307–321.
Gupta RS, Son J, Oren A. 2019. A phylogenomic and molecular markers based taxonomic framework
for members of the order Entomoplasmatales: proposal for an emended order Mycoplasmatales
containing the family Spiroplasmataceae and emended family Mycoplasmataceae comprised of
six genera. *Antonie Van Leeuwenhoek* **112**:561–588.
Hamilton PT, Leong JS, Koop BF, Perlman SJ. 2014. Transcriptional responses in a *Drosophila*
defensive symbiosis. *Mol Ecol* **23**:1558–1570.
Hamilton PT, Peng F, Boulanger MJ, Perlman SJ. 2015. A ribosome-inactivating protein in a
Drosophila defensive symbiont. *Proc Natl Acad Sci U S A* **112**:350–355.
Harumoto T, Lemaitre B. 2018. Male-killing toxin in a bacterial symbiont of *Drosophila*. *Nature*
557:252–255.
Haselkorn TS. 2010. The *Spiroplasma* heritable bacterial endosymbiont of *Drosophila*. *Fly* **4**:80–87.
Haselkorn TS, Cockburn SN, Hamilton PT, Perlman SJ, Jaenike J. 2013. Infectious adaptation:
potential host range of a defensive endosymbiont in *Drosophila*. *Evolution* **67**:934–945.
Haselkorn TS, Jaenike J. 2015. Macroevolutionary persistence of heritable endosymbionts:
acquisition, retention, and expression of adaptive phenotypes in *Spiroplasma*. *Mol Ecol*
14:3752–3765.
Haselkorn TS, Markow TA, Moran NA. 2009. Multiple introductions of the *Spiroplasma* bacterial
endosymbiont into *Drosophila*. *Molecular Ecology* **18**:1294–1305.
Hayashi M, Watanabe M, Yukuhiro F, Nomura M, Kageyama D. 2016. A Nightmare for Males? A
Maternally Transmitted Male-Killing Bacterium and Strong Female Bias in a Green Lacewing
Population. *PLoS ONE* **11**:e0155794.
He L-S, Zhang P-W, Huang J-M, Zhu F-C, Danchin A, Wang Y. 2018. The Enigmatic Genome of an
Obligate Ancient *Spiroplasma* Symbiont in a Hadal Holothurian. *Appl Environ Microbiol*
84:e01965–17.

Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH. 2008. How many species are infected with *Wolbachia*?- a statistical analysis of current data. *FEMS Microbiol Lett* **281**:215–220.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol* **35**:518–522.

Holmes EC. 2009. The Evolution and Emergence of RNA Viruses. Oxford University Press.

Hornett EA, Duplouy AMR, Davies N, Roderick GK, Wedell N, Hurst GDD, Charlat S. 2008. You Can't Keep a Good Parasite Down: Evolution of a Male-Killer Suppressor Uncovers Cytoplasmic Incompatibility. *Evolution* **62**:1258–1263.

Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. 2010. *Wolbachia* as a bacteriocyte-associated nutritional mutualist. *Proc Natl Acad Sci U S A* **107**:769–774.

Ho SYW, Phillips MJ, Cooper A, Drummond AJ. 2005. Time dependency of molecular rate estimates and systematic overestimation of recent divergence times. *Mol Biol Evol* **22**:1561–1568.

Hurst GDD, Jiggins FM. 2000. Male-killing bacteria in insects: mechanisms, incidence, and implications. *Emerg Infect Dis* **6**:329–336.

Itoh T, Martin W, Nei M. 2002. Acceleration of genomic evolution caused by enhanced mutation rate in endocellular symbionts. *Proc Natl Acad Sci U S A* **99**:12944–12948.

Jaenike J. 2012. Population genetics of beneficial heritable symbionts. *Trends Ecol Evol* **27**:226–232.

Jaenike J, Polak M, Fiskin A, Helou M, Minhas M. 2007. Interspecific transmission of endosymbiotic *Spiroplasma* by mites. *Biol Lett* **3**:23–25.

Jaenike J, Unckless R, Cockburn SN, Boelio LM, Perlman SJ. 2010. Adaptation via Symbiosis: Recent Spread of a *Drosophila* Defensive Symbiont. *Science* **329**:212–215.

Jiggins FM, Hurst GDD, Jiggins CD, von der Schulenburg JHG, Majerus MEN. 2000. The butterfly *Danaus chrysippus* is infected by a male-killing *Spiroplasma* bacterium. *Parasitology* **120**:439–446.

Joshi BD, Berg M, Rogers J, Fletcher J, Melcher U. 2005. Sequence comparisons of plasmids pBJS-O of *Spiroplasma citri* and pSKU146 of *S. kunkelii*: implications for plasmid evolution. *BMC Genomics* **6**:175.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**:587–589.

Kanehisa M, Goto S, Hattori M, Aoki-Kinoshita KF, Itoh M, Kawashima S, Katayama T, Araki M, Hirakawa M. 2006. From genomics to chemical genomics: new developments in KEGG. *Nucleic Acids Res* **34**:D354–7.

Kanehisa M, Sato Y, Morishima K. 2016. BlastKOALA and GhostKOALA: KEGG Tools for Functional Characterization of Genome and Metagenome Sequences. *J Mol Biol* **428**:726–731.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* **30**:772–780.

Koonin EV, Mushegian AR, Rudd KE. 1996. Sequencing and analysis of bacterial genomes. *Current Biology* **6**:404–416.

Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* **305**:567–580.

Ku C, Lo W-S, Chen L-L, Kuo C-H. 2013. Complete genomes of two dipteran-associated spiroplasmas provided insights into the origin, dynamics, and impacts of viral invasion in spiroplasma. *Genome Biol Evol* **5**:1151–1164.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome Biol* **5**:R12.

Lanfear R, Calcott B, Ho SYW, Guindon S. 2012. Partitionfinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Mol Biol Evol* **29**:1695–1701.

LeClerc JE, Li B, Payne WL, Cebula TA. 1996. High mutation frequencies among *Escherichia coli* and *Salmonella* pathogens. *Science* **274**:1208–1211.

Lee H, Popodi E, Tang H, Foster PL. 2012. Rate and molecular spectrum of spontaneous mutations in the bacterium *Escherichia coli* as determined by whole-genome sequencing. *Proc Natl Acad Sci U S A* **109**:E2774–83.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**:3094–3100.

Lo W-S, Chen L-L, Chung W-C, Gasparich GE, Kuo C-H. 2013. Comparative genome analysis of Spiroplasma melliferum IPMB4A, a honeybee-associated bacterium. *BMC Genomics* **14**:22.

Lo W-S, Huang Y-Y, Kuo C-H. 2016. Winding paths to simplicity: genome evolution in facultative insect symbionts. *FEMS Microbiol Rev* **40**:855–874.

Lynch M. 2006. Streamlining and simplification of microbial genome architecture. *Annu Rev Microbiol* **60**:327–349.

Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN, Potter SC, Finn RD, Lopez R. 2019. The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Res* **47**:W636–W641.

Massey SE. 2008. The proteomic constraint and its role in molecular evolution. *Mol Biol Evol* **25**:2557–2565.

Masson F, Calderon-Copete S, Schüpfer F, Vigneron A, Rommelaere S, Garcia-Arraez MG, Paredes JC, Lemaitre B. 2020. Blind killing of both male and female *Drosophila* embryos by a natural variant of the endosymbiotic bacterium *Spiroplasma poulsonii*. *Cell Microbiol* **22**:e13156.

Masson F, Copete SC, Schüpfer F, Garcia-Arraez G, Lemaitre B. 2018. In Vitro Culture of the Insect Endosymbiont *Spiroplasma poulsonii* Highlights Bacterial Genes Involved in Host-Symbiont Interaction. *mBio* **9**. doi:10.1128/mbio.00024-18

Mateos M, Castrezana SJ, Nankivell BJ, Estes AM, Markow T a., Moran NA. 2006. Heritable endosymbionts of *Drosophila*. *Genetics* **174**:363–376.

McCutcheon JP, Moran NA. 2011. Extreme genome reduction in symbiotic bacteria. *Nat Rev Microbiol* **10**:13–26.

McFall-Ngai M, Hadfield MG, Bosch TCG, Carey HV, Domazet-Lošo T, Douglas AE, Dubilier N, Eberl G, Fukami T, Gilbert SF, Hentschel U, King N, Kjelleberg S, Knoll AH, Kremer N, Mazmanian SK, Metcalf JL, Nealson K, Pierce NE, Rawls JF, Reid A, Ruby EG, Rumpho M, Sanders JG, Tautz D, Wernegreen JJ. 2013. Animals in a bacterial world, a new imperative for the life sciences. *Proc Natl Acad Sci U S A* **110**:3229–3236.

Miller DE, Staber C, Zeitlinger J, Hawley RS. 2018. Highly Contiguous Genome Assemblies of 15 Species Generated Using Nanopore Sequencing. *G3- Genes Genom Genet* **8**:3131–3141.

Montenegro H, Solferini VN, Klaczko LB, Hurst GDD. 2005. Male-killing *Spiroplasma* naturally infecting *Drosophila melanogaster*. *Insect Mol Biol* **14**:281–287.

Montenegro H, Souza WN, da Silva Leite D, Klaczko LB. 2000. Male-killing selfish cytoplasmic element causes sex-ratio distortion in *Drosophila melanogaster*. *Heredity* **85 Pt 5**:465–470.

Moran NA. 2003. Tracing the evolution of gene loss in obligate bacterial symbionts. *Curr Opin Microbiol* **6**:512–518.

Moran NA. 1996. Accelerated evolution and Muller's rachet in endosymbiotic bacteria. *Proc Natl Acad Sci U S A* **93**:2873–2878.

Moran NA, McCutcheon JP, Nakabachi A. 2008. Genomics and evolution of heritable bacterial symbionts. *Annu Rev Genet* **42**:165–190.

Moran NA, McLaughlin HJ, Sorek R. 2009. The dynamics and time scale of ongoing genomic erosion in symbiotic bacteria. *Science* **323**:379–382.

Mouches C, Barroso G, Gadeau A, Bové JM. 1984. Characterization of two cryptic plasmids from *Spiroplasma citri* and occurrence of their DNA sequences among various spiroplasmas. *Ann Inst Pasteur Microbiol* **135**:17–24.

Nakayama S, Parratt SR, Hutchence KJ, Lewis Z, Price TAR, Hurst GDD. 2015. Can maternally inherited endosymbionts adapt to a novel host? Direct costs of *Spiroplasma* infection, but not vertical transmission efficiency, evolve rapidly after horizontal transfer into *D. melanogaster*. *Heredity* **114**:539–543.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* **32**:268–274.

Oliver KM, Smith AH, Russell JA. 2014. Defensive symbiosis in the real world - advancing ecological studies of heritable, protective bacteria in aphids and beyond. *Funct Ecol* **28**:341–355.

Osaka R, Ichizono T, Kageyama D, Nomura M, Watada M. 2013. Natural variation in population densities and vertical transmission rates of a *Spiroplasma* endosymbiont in *Drosophila hydei*. *Symbiosis* **60**:73–78.

Ota T, Kawabe M, Oishi K, Poulson DF. 1979. Non-male-killing spiroplasmas in *Drosophila hydei*. *J Hered* **70**:211–213.

Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**:526–528.

Paredes JC, Herren JK, Schüpfer F, Lemaitre B. 2016. The Role of Lipid Competition for Endosymbiont-Mediated Protection against Parasitoid Wasps in *Drosophila*. *mBio* **7**:e01006–16.

Paredes JC, Herren JK, Schüpfer F, Marin R, Claverol S, Kuo C-H, Lemaitre B, Béven L. 2015. Genome Sequence of the *Drosophila melanogaster* Male-Killing *Spiroplasma* Strain MSRO Endosymbiont. *MBio* **6**:e02437–14.

Poulson DF, Sakaguchi B. 1961. Nature of "sex-ratio" agent in *Drosophila*. *Science* **133**:1489–1490.

Razin S. 2006. The Genus Mycoplasma and Related Genera (Class Mollicutes) In: Dworkin M., Falkow S., Rosenberg E., Schleifer KH., Stackebrandt E., editor. The Prokaryotes. New York: Springer. pp. 836–904.

R Core Team. 2016. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Regassa LB, Gasparich GE. 2006. Spiroplasmas: evolutionary relationships and biodiversity. *Front Biosci* **11**:2983–3002.

Richardson MF, Weinert LA, Welch JJ, Linheiro RS, Magwire MM, Jiggins FM, Bergman CM. 2012. Population Genomics of the *Wolbachia* Endosymbiont in *Drosophila melanogaster*. *PLoS Genet* **8**:e1003129.

Rocha EPC, Blanchard A. 2002. Genomic repeats, genome plasticity and the dynamics of *Mycoplasma* evolution. *Nucleic Acids Res* **30**:2031–2042.

Rocha EPC, Danchin A. 2002. Base composition bias might result from competition for metabolic resources. *Trends Gene* **18**:291–294.

Saglio PHM, Whitcomb RF. 1979. Diversity of wall-less prokaryotes in plant vascular tissue, fungi, and invertebrate animals In: Whitcomb RF, Tully JG, editors. The Mycoplasmas. New York: Academic Press. pp. 1–36.

Saillard C, Carle P, Duret-Nurbel S, Henri R, Killiny N, Carrère S, Gouzy J, Bové J-M, Renaudin J, Foissac X. 2008. The abundant extrachromosomal DNA content of the *Spiroplasma* citri GII3-3X genome. *BMC Genomics* **9**:195.

Sanada-Morimura S, Matsumura M, Noda H. 2013. Male Killing Caused by a *Spiroplasma* Symbiont in the Small Brown Planthopper, *Laodelphax striatellus*. *J Hered* **104**:821–829.

Schneider DI, Saarman N, Onyango MG, Hyseni C, Opiro R, Echodu R, O'Neill M, Bloch D, Vigneron A, Johnson TJ, Dion K, Weiss BL, Opiyo E, Caccone A, Aksoy S. 2019. Spatio-temporal distribution of *Spiroplasma* infections in the tsetse fly (*Glossina fuscipes fuscipes*) in northern Uganda. *PLoS Neglect Trop D* **13**:e0007340.

Seemann T. 2015. snippy: fast bacterial variant calling from NGS reads.

Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**:2068–2069.

Shigenobu S, Watanabe H, Hattori M, Sakaki Y, Ishikawa H. 2000. Genome sequence of the endocellular bacterial symbiont of aphids *Buchnera* sp. APS. *Nature* **407**:81–86.

Subramanian B, Gao S, Lercher MJ, Hu S, Chen W-H. 2019. Evolview v3: a webserver for visualization, annotation, and management of phylogenetic trees. *Nucleic Acids Res* **47**:W270–W275.

Teixeira L, Ferreira A, Ashburner M. 2008. The bacterial symbiont *Wolbachia* induces resistance to

RNA viral infections in *Drosophila melanogaster*. *PLoS Biol* **6**:e1000002.

Tinsley MC, Majerus MEN. 2006. A new male-killing parasitism: *Spiroplasma* bacteria infect the ladybird beetle *Anisosticta novemdecimpunctata* (Coleoptera: Coccinellidae). *Parasitology* **132**:757–765.

Turelli M, Cooper BS, Richardson KM, Ginsberg PS, Peckenpaugh B, Antelope CX, Kim KJ, May MR, Abrieux A, Wilson DA, Bronski MJ, Moore BR, Gao J-J, Eisen MB, Chiu JC, Conner WR, Hoffmann AA. 2018. Rapid Global Spread of *w*Ri-like *Wolbachia* across Multiple *Drosophila*. *Curr Biol* **28**:963–971.e8.

Turelli M, Hoffmann AA. 1991. Rapid spread of an inherited incompatibility factor in California *Drosophila*. *Nature* **353**:440–442.

Viver T, Orellana LH, Hatt JK, Urdiain M, Díaz S, Richter M, Antón J, Avian M, Amann R, Konstantinidis KT, Rosselló-Móra R. 2017. The low diverse gastric microbiome of the jellyfish *Cotylorhiza tuberculata* is dominated by four novel taxa. *Environ Microbiol* **19**:3039–3058.

Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* **9**:e112963.

Watts T, Haselkorn TS, Moran NA, Markow TA. 2009. Variable Incidence of *Spiroplasma* Infections in Natural Populations of *Drosophila* Species. *PLoS One* **4**:e5703.

Weinert LA, Araujo-Jnr EV, Ahmed MZ, Welch JJ. 2015. The incidence of bacterial endosymbionts in terrestrial arthropods. *Proc R Soc B* **282**:20150249.

Wickham H. 2009. ggplot2: Elegant Graphics for Data Analysis. Springer Science & Business Media.

Wick RR, Judd LM, Gorrie CL, Holt KE. 2017a. Completing bacterial genome assemblies with multiplex MinION sequencing. *Microb Genom* **3**:e000132.

Wick RR, Judd LM, Gorrie CL, Holt KE. 2017b. Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**:e1005595.

Wick RR, Schultz MB, Zobel J, Holt KE. 2015. Bandage: interactive visualization of de novo genome assemblies. *Bioinformatics* **31**:3350–3352.

Wilke CO. 2020. ggridges: Ridgeline Plots in "ggplot2."

Wilke CO. 2019. cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."

Woese CR, Stackebrandt E, Ludwig W. 1984. What are mycoplasmas: the relationship of tempo and mode in bacterial evolution. *J Mol Evol* **21**:305–316.

Wu M, Sun LV, Vamathevan J, Riegler M, Deboy R, Brownlie JC, McGraw EA, Martin W, Esser C, Ahmadinejad N, Wiegand C, Madupu R, Beanan MJ, Brinkac LM, Daugherty SC, Durkin AS, Kolonay JF, Nelson WC, Mohamoud Y, Lee P, Berry K, Young MB, Utterback T, Weidman J, Nierman WC, Paulsen IT, Nelson KE, Tettelin H, O'Neill SL, Eisen JA. 2004. Phylogenomics of the reproductive parasite *Wolbachia pipientis w*Mel: a streamlined genome overrun by mobile genetic elements. *PLoS Biol* **2**:e69.

Xie J, Butler S, Sanchez G, Mateos M. 2014. Male killing *Spiroplasma* protects *Drosophila melanogaster* against two parasitoid wasps. *Heredity* **112**:399–408.

Xie JL, Vilchez I, Mateos M. 2010. *Spiroplasma* Bacteria Enhance Survival of *Drosophila hydei* Attacked by the Parasitic Wasp *Leptopilina heterotoma*. *PLoS One* **5**. doi:10.1371/journal.pone.0012149

Xie J, Winter C, Winter L, Mateos M. 2014. Rapid spread of the defensive endosymbiont *Spiroplasma* in *Drosophila hydei* under high parasitoid was pressure. *FEMS Microbiol Ecol* **91**:1–11.

Yamada M-A, Nawa S, Watanabe TK. 1982. A mutant of SR organism (SRO) in *Drosophila* that does not kill the host males. *Jpn J Genet* **57**:301–305.

Ye F, Melcher U, Rascoe JE, Fletcher J. 1996. Extensive chromosome aberrations in *Spiroplasma citri* strain BR3. *Biochem Genet* **34**:269–286.

Yu G, Smith DK, Zhu H, Guan Y, Lam TT. 2017. ggtree : an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol*

870   *Evol* **8**:28–36.

Supplementary material

Figure S1. Mutational bias in *s*Hy evolution. Rates were calculated by counting the corresponding numbers of snps and dividing by the total number of AT or GC positions in the *s*Hy genome.

Figure S2. Predicted prophage regions in *s*Hy and *s*Mel by PHASTER (blue) and PhiSpy (yellow).

Figure S3. Plasmid synteny in *Spiroplasma poulsonii* strains. Numbers on ticks correspond to nucleotide position on plasmids in kb. Sequence similarities (>=60% and spanning >= 2kb) between plasmids are indicated by ribbons. Abbreviations: ARP - adhesion related protein, OTU - ovarian tumor domain containing protein (in *s*Mel: Spaid), ParA - plasmid partitioning protein like, RIP - ribosome inactivating domain containing protein, Spiralin - spiralin like protein. Figure was created with AliTV.

Figure S4. RIP loci in different *Spiroplasma* genomes. Maximum likelihood tree was calculated with IQ-TREE based on an alignment of RIP domains (288 amino acid positions), created using the hmmalign function of the HMMER software, and manually trimmed to exclude positions present in < 3 sequences. Domain prediction is based on PfamScan, SignalP, and TMHMM. Clades identified by Ballinger & Perlmann (2017) are indicated, and sequences from *s*Hy highlighted in bold font. UFB Bootstrap values ≥90 are shown on nodes. Abbreviations: SP - signal peptide, TM - transmembrane helix, Ank - ankyrin repeats, RIP - ribosome inactivating domain containing protein

Figure S5. Absence (light colors) and presence (dark colors) of genes involved in DNA repair in *Buchnera aphidicola*, *Wolbachia w*Mel, *Mycoplasma gallisepticum*, and *Spiroplasma poulsonii s*Hy and *s*Mel. *Arsenophonus nasoniae* is included as an example for a symbiont with a relatively large genome and complete DNA repair pathways. Genes involved in more than one pathway are listed only once, and those missing in all of the taxa are not displayed. Data is taken from the KEGG pathway database (Kanehisa et al., 2006), and annotation of *s*Mel and *s*Hy genomes as described in Materials and Methods section.

Table S1. *Spiroplasma* genomes included in comparative genomic analyses, with accession numbers, genome properties, and phenotypes induced in host.

Table S2. List of changes observed in *s*Hy and *s*Mel over the course of the evolution experiments.

Table S3. List of missing and truncated genes in *s*Hy and *s*Mel with KEGG annotation.