Nuclear lamin A/C promotes cancer cell survival and lung metastasis without

restricting transendothelial migration

Francesco Roncato[1], Ofer Regev[1], Sara W. Feigelson[1], Sandeep Kumar Yadav[1], Lukasz

Kaczmarczyk[2], Nehora Levi[2], Diana Drago-Garcia[3], Samuel Ovadia[1], Marina Kizner[1], Yoseph

Addadi[4], João C. Sabino[5], Yossi Ovadya[6], Sérgio F. de Almeida[5], Ester Feldmesser[4], Gabi

Gerlitz[2] and Ronen Alon[1]*

1 Department of Immunology, Weizmann Institute of Science, Rehovot, 76100, Israel

2 Department of Molecular Biology, Ariel University, Kiryat Hamada, Ariel, 40700, Israel

3 Department of Biological Regulation, Weizmann Institute of Science, Rehovot 76100, Israel

4 Life Sciences Core Facilities, Weizmann Institute of Science, Rehovot, 76100, Israel

5 Instituto de Medicina Molecular João Lobo Antunes, Faculdade de Medicina da

Universidade de Lisboa, Lisboa, 1649-028, Portugal.

6 Department of Molecular Cell Biology, Weizmann Institute of Science, 76100 Rehovot, Israel

*For correspondence: ronen.alon@weizmann.ac.il

Abstract

The mechanisms by which the nuclear lamina of tumor cells controls their migration and

survival are poorly understood. Lamin A and its variant lamin C are key nuclear lamina proteins

that control nucleus stiffness and chromatin conformation. Downregulation of lamin A/C levels

in two metastatic lines, B16F10 melanoma and E0771 breast carcinoma, facilitated cell

squeezing through rigid pores, elevated nuclear deformability and reduced heterochromatin.

Unexpectedly, the transendothelial migration of both cancer cells in vitro and in vivo, through

lung capillaries, was not elevated by lamin A/C knockdown. Both cancer cells with lamin A/C

knockdown grew normally in primary tumors and in vitro on rigid surfaces. Strikingly, however,

both lamin A/C deficient melanoma and breast cancer cells grew poorly in 3D spheroids

expanded in soft agar cultures. Experimental lung metastasis of both lamin A/C knockdown

cells was also markedly reduced. Taken together, our results suggest that high content of

lamin A/C in multiple cancer cells promotes cancer cell survival and ability to generate lung

metastasis without compromising cancer cell emigration from lung vessels.

Introduction

The nucleus is the largest and stiffest organelle in all cells and therefore imposes the main

barrier for cell crossing of cellular and mechanically resistant extracellular barriers (Burke &

Stewart, 2013; Friedl, Wolf, & Lammerding, 2011; Fruleux & Hawkins, 2016). The nucleus

must undergo various shape changes during cell migration through cellular and extracellular

barriers (Friedl et al., 2011). Lamin A and its splice variant lamin C are key nuclear lamina

intermediate filament proteins that control nucleus stiffness (Burke & Stewart, 2013;

Lammerding et al., 2006; Roman et al., 2017; Shin et al., 2013), and regulate chromatin

conformation and accessibility (Bronshtein et al., 2015; Towbin, Meister, & Gasser, 2009). A-

type lamins also control nuclear crosstalk with all types of the cell cytoskeleton, including

microtubules and actin filaments (Chambliss et al., 2013; Chang, Folker, Worman, &

Gundersen, 2013; Etienne-Manneville & Lammerding, 2017; Khatau et al., 2009; Lombardi &

Lammerding, 2011), and thereby regulate nuclear location and response to mechanical

signals from the extracellular environment (Borrego-Pinto et al., 2012; Gonzalez-Granado et

al., 2014; Graham et al., 2018; Kim & Wirtz, 2015; Kirby & Lammerding, 2018; Swift et al.,

2013).

Alterations in nuclear lamina stiffness can take place by various DNA damaging

processes including the recently described nuclear autophagy, nucleophagy (Dou et al., 2015)

or by genetic changes (J. L. V. Broers & Ramaekers, 2014). The latter changes can be

experimentally introduced by controlled suppression or overexpression of lamin A/C and such

changes have been studied in different types of cells migrating through variably rigid

confinements in vitro and in vivo (C. M. Denais et al., 2016). The soft nuclei of most leukocytes

contain low levels of lamin A/C and high levels of other lamins, primarily of the B type (Shin et

al., 2013). The low ratio of A and B lamins allows leukocytes to undergo massive and rapid

deformation during fast squeezing through vascular endothelial junctions and collagenous

interstitial spaces (Yadav et al., 2018). How nuclear squeezing is regulated in solid cancer

cells migrating through variable interstitial ECM barriers and constricted vascular spaces is

only partially understood (C. M. Denais et al., 2016; Raab et al., 2016; Wolf et al., 2013). In

contrast to leukocyte nuclei which express very low levels of lamins A/C, the nuclei of variably

invasive metastatic tumor cells of mesenchymal origin is predicted to contain higher levels of

these lamins and are therefore generally more stiff, potentially imposing higher restrictions on

the ability of these cells to cross endothelial junctions and interstitial barriers (Cao et al., 2016).

The cytoskeleton of endothelial cells which comprise the major barriers for nuclear squeezing

is, however, fairly elastic and contractile (Barzilai et al., 2017; Heemskerk et al., 2016). In

contrast, although collagenous barriers can undergo extensive remodeling by cell-generated

forces and proteolysis (Infante et al., 2018) solid tumor cells with high lamin A/C content might

less efficiently cope with these barriers than with endothelial barriers (Yadav et al., 2018).

Support for this idea was recently provided by experiments with leukocytes overexpressing

lamin A: a 10-fold increase in the ratio of lamin A to lamin B dramatically restricted leukocyte

nuclear squeezing through rigid pores and dense collagenous barriers but was largely

permissive for leukocyte transendothelial migration (Yadav et al., 2018).

Lamins A/C are also involved in chromatin conformation and epigenetics via their

interactions with heterochromatin, transcriptionally repressed tightly folded chromatin tethered

to the nuclear lamina (Becker, Nicetto, & Zaret, 2016; Dechat, Adam, Taimen, Shimi, &

Goldman, 2010; Harr et al., 2015). Nevertheless, the direct contributions of lamins to cancer

cell migration, growth, and malignancy have been in debate (J. L. V. Broers & Ramaekers,

2014). On one hand, lamin A/C expression is reduced in several solid cancers and cancer

progression was suggested to correlated with lower lamin A/C expression (Bell & Lammerding,

2016; C. Denais & Lammerding, 2014; Kaufmann, Mabry, Jasti, & Shaper, 1991), it is absent

in around 40% of human breast cancer tissues (Capo-chichi et al., 2011), and decreased lamin

A/C expression is a sign of poor prognosis in skin, breast, lung and colon cancers (J. L. Broers

et al., 1993; Capo-chichi et al., 2011; Venables et al., 2001). Since lamin A/C knockdown

nuclei can be more easily deformed and squeeze through rigid confinements, it has been

postulated that tumor cells with low expression levels of these lamins can more readily invade

tissues (Davidson, Denais, Bakshi, & Lammerding, 2014). Nevertheless, the direct in vivo

evidence for this assumption has never been provided. Furthermore, overexpression of A-

type lamins in some cancer cells was shown to correlate with enhanced growth and faster

migration via the activation of the PI3K/AKT/PTEN pathway (Kong et al., 2012). Part of these

discrepancies are attributed to the complex roles of A-type lamins in protecting the nuclei from

mechanical nuclear rupture, DNA damage, and cell growth arrest (Cho et al., 2019). These

discrepancies have motivated us to address these standing questions by controlled

downregulation of lamin A/C expression introduced into multiple bona fide metastatic cells.

We specifically wished to address whether extravasation of circulating cancer cells from

different blood vessels, and in particular from the relatively impermeable lung capillaries, are

favored by the softening of tumor cell nuclei via downregulation of lamin A/C. We hence

systematically addressed both in vitro and in vivo in syngeneic mice models if the forced

downregulation of lamins A/C expression with retained levels of B lamins alters both invasive

and proliferative properties of two prototypic metastatic cell lines, namely, B16 melanoma and

E0771 breast carcinoma. In vitro, as expected, downregulated lamin A/C levels introduced by

ectopic expression of lamin A/C specific shRNA dramatically facilitated the squeezing of both

cells through rigid pores, reduced their heterochromatin content, and altered their

transcriptional signature. Surprisingly, however, circulating lamin A/C knockdown cells

normally emigrated from the pulmonary circulation and normally accumulated inside the lung

parenchyma. Nevertheless, the ability of lamin A/C low melanoma and breast cancer cells to

generate metastatic lesions in the lungs was markedly compromised. In vitro, both B16

melanoma and E0771 breast carcinoma cells with lamin A/C knockdown exhibited specialized

proliferation defects when grown in spheroids within 3D environment whereas their intrinsic

growth on solid 2D surfaces remained intact. Our results collectively suggest that high nuclear

lamin A/C content does not compromise the squeezing ability of melanoma and breast cancer

cells through physiological endothelial barriers. High content of lamin A/C is necessary,

however, for optimal growth of these multiple cancer cells in multi-cellular assemblies in vitro

and for their ability to generate lung metastasis in vivo.

Results

Lamin A downregulation in B16F10 melanoma increases nucleus deformability and

squeezing through rigid pores but does not affect transendothelial migration.

To gain insight into the contribution of type A lamins to the ability of solid tumor cells to

translocate their nuclei across endothelial barriers, we downregulated lamin A and lamin C

expression in the nuclei of the B16F10 melanoma cells by stably introducing a lamin A/C

shRNA construct targeting exon 8 of the *Lmna* gene (Figure 1A, B). Since the nuclear lamina

stiffness is sensitive to the ratio between type A and type B lamins in both mesenchymal and

hematopoietic cells (Shin et al., 2013), we expected that lamin A/C downregulation would also

increase nuclear deformability and squeezing capacities in our melanoma cell model.

Downregulating lamin A and lamin C expression in the B16 melanoma line by 90%, but leaving

lamin B1 levels intact, altered the ratio of type A to type B lamins by approximately 10-fold

(Figure 1A). As expected, lamin A/C downregulation resulted in a dramatic enhancement of

B16 squeezing through small rigid pores but much less so through large pores (Figure 1C, D).

A similar gain of squeezing of lamin A/C downregulated B16F10 cells was observed with

another lamin A/C shRNA construct (Figure 1−figure supplement 1A) and in cells subjected to

transient *Lmna* gene exon-4 targeted siRNA mediated downregulation of lamin A/C

transcription (Figure 1−figure supplement 1B).

To assess if and how these changes in the composition and deformability of the tumor

nucleus promotes tumor cell squeezing through physiological endothelial barriers, we

established a new in vitro video-microscopy based assay in which nuclear squeezing of tumor

cell crossing confluent monolayers of bEnd.3 murine endothelial cells can be compared.

Nucleus location, deformation and squeezing in individual transmigrating tumor cells could be

readily tracked in real time by fluorescence and phase contrast microscopy of tumor cells

whose nuclei were prelabeled with the nuclear dye Hoechst (Video 1-2). This assay allowed

us to follow how individual tumor cells complete an entire sequence of TEM steps immediately

after settling on confluent endothelial monolayers, including protrusion, generation of large

sub-endothelial pseudopodia (lamellipodia), squeezing their nuclei, tail detachment, and

locomotion underneath the endothelial monolayer (Figure 1E, Video 2). Interestingly, out of

three murine cancer cell lines compared in this new assay, namely B16F10, E0771, and LL/2,

B16 melanoma exhibited the highest extent of TEM (Figure 1F). Notably, the TEM of B16F10

as well as of other cancer cells compared in this system was over 30-fold slower than that of

leukocytes (Barzilai et al., 2017; Yadav et al., 2018) due to a very slow formation of the sub-

endothelial leading edge (t= 26±22 min for B16F10 as compared to 30±15 s for T cells), and

the inability of the tumor nucleus to translocate into this leading edge as during leukocyte TEM

(Barzilai et al., 2017; Yadav et al., 2018).

Strikingly, and in contrast to the squeezing results across rigid pores (Figure 1C, D),

the extent of B16F10 TEM was not increased by lamin A/C knockdown (Figure 2A, B, Video

3). However, the extent of nuclear deformation for lamin A/C downregulated nuclei was

significantly greater than that of the control tumor cells (Figure 2C). The nuclei of the lamin

A/C knockdown B16F10 deformed more readily also when these cells spread on a non-

confined 2D substrate coated with the basement membrane deposited by the endothelial

monolayer, whereas the motility of lamin A/C knockdown cells remained normal (Figure 2D,

Video 4). These results collectively suggest that the reduced nuclear stiffness and the

increased nuclear deformability of lamin A/C knockdown B16F10 cells (Figure 2C, D) do not

provide a migratory advantage for the transmigration of these cells through endothelial

barriers. These observations also suggest that differences in tumor cell squeezing through

infinitely rigid barriers do not correlate with squeezing through physiological cellular barriers.

This finding could be due to the high mechanical flexibility of the endothelial cytoskeletal

barriers maintained by their rapid actin turnover (Ofer, Mogilner, & Keren, 2011) and dynamic

contractility (Heemskerk et al., 2016).

Lamin A downregulation does not increase B16F10 extravasation across lung vessels in vivo and does not accelerate melanoma apoptosis in the lung parenchyma

Our in vitro results thus indicated that lamin A/C downregulation in B16F10 cells enhance their squeezing through rigid confinement without affecting ability to transmigrate across endothelial barriers. To assess the distinct migratory outcomes of lamin A/C downregulation in B16 melanoma cells in vivo, we introduced an experimental lung metastasis model based on i.v. injection of minute numbers of fluorescently labeled cells (Figure 2–figure supplement 1), in order to minimize non-physiological inflammatory responses associated with a bolus of cancer cells that simultaneously enter the lung vasculature. Tumor metastasis in lung spreads nearly exclusively through the pulmonary capillaries (Miles, Pruitt, van Golen, & Cooper, 2008), an extensive network of relatively impermeable capillaries (Chambers, Groom, & MacDonald, 2002) considered to be poorly permeable compared with vessels targeted by metastatic cells at other organs like the bone marrow and liver (Valastyan & Weinberg, 2011). I.V. injections of tumor cells are extensively used for studying hematogenous dissemination and expansion in the lung (Gorelik & Flavell, 2001). Since the tumor cells are introduced i.v. in a single event, their arrival at the lung vasculature is synchronized which allows accurate temporal dissection of the earliest extravasation steps taken by individual circulating metastatic cells entering the lung vasculature several hours post injection. To determine the effects of lamin A/C downregulation on earliest tumor cell extravasation across the lung capillaries, we compared the cancer cell partition inside and outside lung vessels with newly developed 3D imaging of the injected fluorescently labeled cancer cells in relation to CD31 stained lung vessels (Figure 3A, B, Video 5). Strikingly, lamin A/C downregulated B16F10 cells extravasated at similar efficiencies to normal B16F10 cells (Figure 3C, Videos 6-8). Lamin A/C downregulation also did not affect the total number of B16F10 cells accumulated inside the recipient lung at this early time points (Figure 3D). These results suggest that unlike its dramatic effects on tumor squeezing in vitro, but consistent with our TEM results in vitro (Figure 2A), lamin A/C suppression does not facilitate melanoma extravasation across lung vessels in vivo.

Lamin A/C downregulation reduces the content of H3K9Me3 heterochromatin but does not alter DNA stability

Lmna knockout (KO) was shown to reduce the major marker of constitutive heterochromatin H3K9me3 and the methyltransferases that generate it in primary cells (Liu et al., 2013). As predicted, lamin A/C downregulation led to lower levels of H3K9Me3 and significantly reduced levels of SUV39H2 and SETDB1, two main histone methyltransferases that maintain this suppressive epigenetic marker (Figure 4A, B). In contrast, facultative heterochromatin levels probed by the H3K27me3 marker remained unchanged (Figure 4B). This global H3K9me3 reduction was not associated with global repression of transcription as determined by a transcription run-on experiment (Figure 4C). Notably, transcription of specific genes may be altered by reduced H3K9me3 levels and lamin A/C deficiency was reported to increase chromatin dynamics, which could also affect transcription (Bronshtein et al., 2015; Solovei et al., 2013; Sullivan et al., 1999). To evaluate these possibilities, we performed RNA-seq analysis on control and lamin A/C shRNA transduced B16F10 cells. We found that the transcript levels of LINC complex components such as nesprins 1-4, emerin, as well as of other lamin A/C interactors involved in heterochromatin content and stability (e.g. ,Trim28, Tpr or Nup153 (Krull et al., 2010; Kubben et al., 2010)) were all normally transcribed in lamin A/C knockdown B16 cells (Supplementary file 1). Lamin A/C downregulated B16F10 cells exhibited, however, alterations in several hundred genes, including nuclear matrix genes, nuclear body genes, as well as in lamin B receptor (Figure 4D, E and Figure 4–figure supplement 1A-C; Supplementary file 1). Consistent with their similar transmigration properties in vitro and similar capacity to extravasate lung capillaries in vivo, our transcript analysis did not indicate any differences in vascular permeability factors such as VEGF family member levels, MMP levels, or canonical and abundantly expressed cytoskeletal machineries (e.g., Rho GTPases and myosin) (Fig. 4D and Supplementary file 1).

Lamin A/C downregulation does not alter cell proliferation on rigid 2D surfaces but impairs 3D cell growth in spheroids

Our transcriptional analysis did not detect changes in genes involved in melanoma growth (Supplementary file 1). Indeed, melanoma cells knockdown in lamin A/C expression exhibited similar growth rates when seeded at low densities on tissue culture plates (Figure 5A). Interestingly, deficiency in lamin A/C content also did not increase the susceptibility of the nuclei of these melanoma cells to DNA damage: lamin A/C knockdown B16 cells remained as resistant as control melanoma cells to apoptosis and growth arrest induced by the DNA damage and cell cycle arrest inducer etoposide (Dai et al., 2017) (Figure 5B). An additional incubation period without etoposide (Figure 5–figure supplement 1), drove the growth arrested B16 cells into senescence with a slightly higher senescence acquisition exhibited by the lamin A/C knockdown melanoma cells (Figure 5C). Nevertheless, the quantity of R-loops, transient RNA-DNA hybrids which correlate with DNA damage (Crossley, Bocek, & Cimprich, 2019) were comparable for control and lamin A/C knockdown B16 cells (Figure 5–figure supplement 2). These results collectively suggest that lamin A/C knockdown B16F10 cells expand normally in conventional 2D culture conditions, do not exhibit any spontaneous DNA damage, and undergo normal growth arrest induced by a chemical DNA damaging signal.

The soft agar colony formation assay has been a hallmark of cancer survival analysis, as it measures the ability of cells to proliferate in semi-solid matrices (Horibata, Vo, Subramanian, Thompson, & Coonrod, 2015). This readout also allows a direct comparison of cancer cell growth in spheroids vs. in colonies expanded in the same soft agar, while remaining in direct contact with the culture dish 2D surface (Figure. 5D). Interestingly, whereas in the first 3 days both lamin A/C knockdown and control B16F10 cells gave rise to similar sized spheroids (Figure 5E), the ability of lamin A/C high B16 spheroids to further expand during the next 3 days significantly surpassed that of lamin A/C knockdown cells (Figure 5E). In contrast, the growth rate of both B16F10 cell groups settled on the culture plate in the presence of identical soft agar remained similar (Figure 5E, inset). These results were

reproduced with a second lamin A/C knockdown B16F10 line expressing a distinct lamin A/C

shRNA (Figure 5–figure supplement 3). Remarkably, even in the presence of nutrient rich agar

(i.e., a 5-fold higher serum concentration) that accelerated the growth of both lamin A/C high

and lamin knockdown B16 cells, the mean size of lamin A/C high B16 cells spheroids was

significantly higher than that of lamin A/C low B16 cells (Figure 5F). Interestingly, a negligible

fraction of cells isolated from spheroids (9 vs 12 %) underwent apoptosis as depicted by

annexin V and propidium iodide staining (Figure 5–figure supplement 4), ruling out the

possibility that cells growing at the core of these spheroids have poor accessibility to nutrients.

Collectively, our results suggest that whereas initially, lamin A/C knockdown B16F10 normally

proliferate under both 2D and 3D conditions, the ability of lamin A/C knockdown cells to

expand within 3D spheroids is significantly impaired, possibly due to slower growth of these

cells inside large spheroids embedded in 3D environments.

Lamin A/C deficiency does not affect melanoma growth in distinct primary transplants

but results in poorer survival in the lungs and reduced metastasis

In order to validate our in vitro observations, lamin A/C knockdown B16F10 cells or their

control counterparts were subcutaneously implanted at 10-20 fold lower numbers than in

commonly used orthotopic melanoma models in order to avoid the masking of any potential

growth differences. Interestingly, both lamin A/C knockdown and control B16F10 cells grew at

similar rates in vivo (Figure 6A). Histological analysis of the orthotopic tumors also failed to

show any noticeable differences in the vascular microenvironment (data not shown).

Furthermore, both lamin A/C high and low melanoma cells shared similar growth properties

when implanted in the mammary fat pad (Figure 6B). Notably, although lamin A/C knockdown

B16F10 accumulated at comparable levels to control B16F10 inside recipient lungs within the

first 3 days after injection the number of viable lamin A/C knockdown B16F10 recovered from

recipient lungs 7 days after injection was significantly reduced relative to control B16F10 cells

(Figure 6C). Furthermore, the ability of lamin A/C B16F10 cells to generate metastatic lesions

14 days after injection into recipient mice was dramatically reduced by lamin A/C

downregulation (Figure 6D). These results collectively suggest that lamin A/C knockdown

B16F10 cells poorly survive in the lungs, in spite of their normal apparent extravasation

potential when circulating and entering the blood vasculature.

Reduced lamin A/C levels in E0771 breast cancer cells recapitulate the in vitro and in

vivo migratory properties and survival deficiencies of lamin A/C knockdown B16 cells

We next reasoned that reduced lamin A/C levels could differentially impact distinct types of

cancer cells. We therefore addressed how reduced lamin A/C expression in the bona fide

breast cancer cells, the E0771 line, affects their squeezing, migration, epigenetics and growth

properties both in vitro and in vivo. Downregulation of lamin A/C levels with conserved lamin

B content (Figure 7A), introduced either by shRNA-expressing lentiviral vectors or by siRNA,

dramatically facilitated cancer cell squeezing through rigid pores in vitro (Figure 7B and Figure

7−figure supplement 1A, B), in agreement with our findings with B16F10 cells. While the

nuclear circularity index of control E0771 cells was higher than that of control B16F10 cells

(Figure 7−figure supplement 1C and Figure 2D), the nuclei of these cells underwent significant

increases in deformability (i.e. reduced circularity) upon lamin A/C downregulation (Figure

7−figure supplement 1D), reminiscent of the effect of lamin A/C downregulation on B16 nuclei.

Nevertheless, the transendothelial migration capacity of these cells in vitro and in vivo was

insensitive to reduced lamin A/C expression (Figure 7C and Figure 7−figure supplement 2A-

B). Although downregulated lamin A/C expression resulted in reduced H3K9Me3

heterochromatin content and in altered gene expression (Figure 7D and Figure 7−figure

supplement 3A-C), it did not increase global RNA transcription rates (Figure 7−figure

supplement 4). Furthermore, primary breast cancer growth in vitro or in vivo in the mammary

fat pad or in non-orthotopic skin implants was unaffected by lamin A/C downregulation (Figure

7E, F and Figure 7−figure supplement 5). DNA damage-induced growth arrest was also

insensitive to downregulated lamin A/C expression (Figure 7G). Nevertheless, and

reminiscent of the proliferation results of lamin A/C knockdown B16F10 cells lamin A/C low,

E0771 poorly proliferated in spheroids grown in 3D conditions compared to control (Figure

7H). Notably, although Lamin A/C knockdown E0771 cells normally accumulated in recipient

lungs when intravenously introduced into syngeneic recipient mice (Figure 7–figure

supplement 6). Breast cancer cells with low lamin A/C content failed to generate any

metastatic lesions in the lungs (Figure 7I, J). Thus, although reduced nuclear content of lamin

A/C does not impair breast cancer growth in the primary tissue and does not alter breast

cancer cell extravasation from lung vessels into the lung parenchyma, it dramatically

compromises the metastatic potential of these breast cancer cells in this organ, reminiscent

of our observations with lamin A/C knockdown melanoma cells.

Discussion

The nucleus is the most bulky organelle in all cells and is protected by a mechanically stable

network underlying the inner nuclear membrane termed the nuclear lamina (Swift et al., 2013;

Katherine L Wilson & Foisner, 2010; Wolf et al., 2013). The main mechanical obstacle for the

extravasation of solid cancer cells across blood vessels in vivo is their stiff nuclei (Shin et al.,

2013). The nuclear lamina is thought to affect the shape and the mechanical properties of the

nucleus, hence to control cell squeezing through different barriers (Lomakin et al., 2020;

Rowat et al., 2013; Wolf et al., 2013; Yadav et al., 2018). The lamina also plays an important

role in the maintenance of the nuclear envelope integrity as well as in the organization of the

nucleus as a whole (Friedl et al., 2011; Kirby & Lammerding, 2018). The nuclear lamina is

also connected to highly condensed chromatin regions (heterochromatin) (Stephens et al.,

2012; K. L. Wilson & Berk, 2010) and can affect chromatin conformation, and epigenetics

(Fernández-Morera, Calvanese, Rodríguez-Rodero, Menéndez-Torre, & Fraga, 2010;

Sullivan et al., 1999). The lamina also controls the entry of key growth control transcriptional

factors including the mechanosensitive transcriptional activator YAP (Elosegui-Artola et al.,

2017). Lamin A/C expression is reduced in several solid cancers but not others, and so the

molecular basis of these changes and their direct link to cancer cell migration, survival and

expansion have been under debate due to the complexity and diversity of tumor growth,

migration and metastasis (Harada et al., 2014; Kong et al., 2012; Wazir et al., 2013).

We chose to address these key standing questions using two prototypic BL/6

metastatic cell lines, B16F10 melanoma and E0771 breast carcinoma. Taking both in vitro

and in vivo reductionist approaches in syngeneic immunocompetent mice, we have

systematically assessed how controlled suppression of the two type A lamins affects the

specific growth and migratory properties of these cells under distinct physiologically relevant

conditions and in different environmental conditions both in vitro and in vivo. Our in vitro

findings of enhanced squeezing capabilities of the lamin A knockdown cells in the transwell

assay are in full agreement with previous results on the critical role of lamin A/C in the ability

of cells of different origins to squeeze through rigid confinements in vitro (Aureille, Belaadi, &

Guilluy, 2017; Rowat et al., 2013). However, we found that nuclear deformability has no impact

on the overall nuclear squeezing kinetics through endothelial junctions and under endothelial

monolayers, indicating that endothelial barriers are highly permissive for nuclear passage and

well adapted to accommodate the squeezing of cells with bulky and stiff nuclei. The

exceptionally slow rates of tumor transendothelial migration, may also provide the endothelial

cytoskeleton with sufficient time to undergo remodeling including activation of contractility

machineries to facilitate the squeezing of the relatively stiff nuclei of most tumor cells though

junctions or transcellular pores (Khuon et al., 2010; Yadav et al., 2018). In order to validate

our in vitro TEM results, we developed a new experimental lung metastasis model to assess

in vivo the intrinsic ability of our lamin A/C suppressed tumor cells to squeeze through the lung

vasculature. Tumor metastasis into lungs occurs nearly exclusively through the pulmonary

capillaries (Miles et al., 2008), an extensive network of relatively impermeable capillaries

(Chambers et al., 2002) considered to be poorly permeable compared with vessels targeted

by metastatic cells at other organs like the bone marrow and liver (Valastyan & Weinberg,

2011). By introducing a new method to accurately measure the relative efficiency of cancer

cell emigration through these vessels, we found that highly invasive cells like melanoma B16

crossed these barriers in vivo independently of their lamin A/C content and irrespectively of

their intrinsic nuclear deformability properties. Although poorly invasive, the ability of E0771

cells to cross identical pulmonary vascular barriers was also irrespective of their intrinsic

nucleus deformability and lamin A/C content. Our in vivo results were therefore fully consistent

with the transmigratory properties of these cells determined in our in vitro setups.

Metastasis involves not only cancer cell extravasation but also intravasation into tumor

engulfed blood vessels (Joyce & Pollard, 2009). Although the extravasation capacity of a given

cancer cell does not predict its intravasation potential across the tumor associated blood

vessels at its original site of dissemination, both our in vitro and in vivo analysis of tumor cell

squeezing through different endothelial barriers predict that tumor cell squeezing through

endothelial junctions is not affected by lamin A/C downregulation. Furthermore, since the lung

capillaries, where the majority of our cancer cell diapedesis took place, are less permeable

than tumor associated vessels and were permissive for the squeezing of lamin A/C rich nuclei,

it is likely that the leaky vessels surrounding the primary tumors also do not pose a barrier for

melanoma or breast cancer intravasation. Likewise, melanoma and breast cancer crossing of

lymphatic vessels nearby tumors, considered highly permeable cell barriers (Dyer &

Patterson, 2010), is probably insensitive to the lamin A/C content of the tumor nuclei.

Our results argue that the squeezing ability of a given tumor cell through non

degradable rigid pores towards a chemotactic or a haptotactic signal does not directly predict

the physiological crossing potential of that cell. Although not addressed in our study, it is

possible that upon entering the interstitial space, the ability of lamin A/C rich tumor cells to

enzymatically degrade glycoprotein and proteoglycan components of collagenous barriers

(Bishop, Schuksz, & Esko, 2007) is much more critical for their interstitial motility than the

mechanical restrictions imposed by the stiffness of their lamin A/C rich nuclei Indeed, a lamin

A/C regulated crosstalk between the nucleus and MT1-MMP has been recently shown to

promote a digest-on-demand program in cells squeezing through constricted spaces (Infante

et al., 2018). This study strongly suggests that high levels of lamin A/C and optimal

connections of these lamins with the cell cytoskeleton (e.g., via the linker of nucleoskeleton

and cytoskeleton (LINC) complex) can be in fact essential for directional proteolysis at the cell

front.

Our epigenetic analysis revealed lower levels of the constitutive heterochromatin

marker H3K9me3 and the methyltransferases that generate it in lamin A/C knockdown

B16F10 and E0771 cells, consistent with previous reports on lamin A/C KO MEFs (Liu et al.,

2013) and human fibroblasts expressing mutated type A lamins (Scaffidi & Misteli, 2006).

Reduced heterochromatin content has been argued to reduce chromatin compaction in B16

melanoma and thereby restrict cell motility (Maizels et al., 2017). The increased squeezing

rate of the lamin A/C knockdown B16 cells and E0771 cells that we find in spite of their lower

H3K9me3 content sheds new light on these earlier findings. Our data therefore suggest that

the altered mechanical properties of our melanoma and breast cancer cell models introduced

by lamin A/C downregulation override any inhibitory effects of reduced heterochromatin

content on chromatin compaction and cell motility (Maizels et al., 2017). Previous observations

showed that depletion of lamin A/C in lung and breast cancer cells, as well as fibrosarcoma

cells, significantly increased the likelihood of transient nuclear envelope rupture events and

cell death especially when cells were forced to migrate through very tight and rigid barriers (C.

M. Denais et al., 2016; Harada et al., 2014). Imaging of HT1080 fibrosarcoma cells invading

the collagen-rich mouse dermis in live tumors after orthotopic implantation confirmed that

migration-induced nuclear envelope rupture occurs in vivo, particularly during cell division

(Vargas, Hatch, Anderson, & Hetzer, 2012) and in individually disseminating cells (C. M.

Denais et al., 2016). Nuclear envelope rupture is less prevalent, however, in cells moving as

multicellular collective strands (C. M. Denais et al., 2016) or on 2D surfaces (Harada et al.,

2014). Notably, both the melanoma and breast cancer cells studied by us grew and survived

normally when orthotopically or non orthotopically implanted in distinct organs. Our results

therefore argue against faster nuclear rupture of these cells in their primary tissues. On the

other hand, lamin A/C knockdown melanoma and breast cancer cells exhibited poor long term

survival in the lungs as well as reduced growth inside spheroid assemblies grown in 3D

environments, two environments that are far less rigid than tissue culture dishes (Barney et

al., 2016; Jaiswal et al., 2017). Therefore, the contribution of lamin A/C to cancer cell growth

within spheroids is unlikely to involve classical lamin A/C regulated mechanotransduction

which predominates cell-extracellular matrix contacts (Donnaloja, Carnevali, Jacchetti, &

Raimondi, 2020).

The contribution of high lamin A/C content to chromatin flexibility and DNA stability is

still disputed. On one hand, high lamin A/C content protects the nuclei from rupture and cell

apoptosis occurring under extreme mechanical stress such as squeezing through rigid

confinements (C. M. Denais et al., 2016; Raab et al., 2016) and also reduces the frequency

of spontaneous aneuploidy (Capo-chichi et al., 2011) and DNA damage (Gonzalo, 2014). A-

type lamins also participate in the maintenance of telomere homeostasis (Gonzalez-Suarez

et al., 2009). Our results extend these findings proposing an additional role of lamin A/C, a

protective function of these lamina proteins in multi-cellular assemblies, in the absence of

apparent squeezing through rigid confinements. Our findings are the first to implicate lamin

A/C as essential protective factors in lung metastasis of multiple types of cancer cells. Our

results predict that lamin A/C deficiency impairs cancer cell survival in the lungs to a greater

extent than the migratory advantages this deficiency may provide to cancer cells. The

relevance of our findings to other types of cancers and metastatic spread in other organs

awaits for future studies.

Materials and methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (*Mus musculus*)	C57BL/6	The Jackson Laboratory		
Cell line (*Homo sapiens*)	HEK293T	ATCC	ATCC CRL-3216	
Cell line (*Mus musculus*)	bEnd.3	Gift from Britta Engelhardt, Theodor Kocher Institute, University of Bern	ATCC CRL-2299	
Cell line (*Mus musculus*)	B16F10	Gift from Prof. Lea Eisenbach, Weizmann Institute of Science	ATCC CRL-6475	
Cell line (*Mus musculus*)	E0771	Gift from Dr. Ravid Straussman, Weizmann Institute of Science	CH3 BioSystems Cat# 94A001	
Cell line (Mus *musculus*)	LL/2 (LLC1)	Gift from Dr. Ravid Straussman, Weizmann Institute of Science	ATCC CRL-1642	
Antibody	anti-Lamin A/C (4C11)	Cell Signaling Technology	Cat# 4777, RRID:AB_10545756	
Antibody	anti-Lamin B1 [EPR8985(B)]	Abcam	Cat# ab133741, RRID:AB_2616597	
Antibody	anti-GAPDH (6C5)	Millipore	Cat# MAB374, RRID:AB_2107445	
Antibody	Alexa Fluor® 647 anti-CD31 (390)	BioLegend	Cat# 102416, RRID:AB_493410	
Antibody	anti-trimethyl-Histone H3 (Lys9) (D4W1U)	Cell Signaling Technology	Cat# 13969, RRID:AB_2798355	
Antibody	anti-KMT1B/SUV39H2 [EPR18495]	Abcam	Cat# ab190870, RRID:AB_2827544	
Antibody	anti-trimethyl Histone H3 (Lys27) (18E9.1)	Millipore	Cat# 05-1951, RRID:AB_11211815	
Antibody	anti-Histone H3 (A3S)	Millipore	Cat# 05-928, RRID:AB_492621	
Antibody	anti- ESET/SETDB1 (D4M8R)	Cell Signaling Technology	Cat# 93212, RRID:AB_2800200	
Antibody	anti-alpha Tubulin (DM1A)	Thermo Fisher Scientific	Cat# 62204, RRID:AB_1965960	
Antibody	anti-DNA-RNA Hybrid (S9.6)	Millipore	Cat# MABE1095, RRID:AB_2861387	
Antibody	anti mouse IgG+IgM+IgA cross-adsorbed (H+L)	Bethyl	Cat# A90-244D2, RRID:AB_10683272	

Antibody	Alexa Fluor® 488 anti-mouse IgG (H+L)	Thermo Fisher Scientific	Cat# A-21202, RRID:AB_141607	
Antibody	Peroxidase-AffiniPure anti-mouse IgG (H+L)	Jackson ImmunoResearch Labs	Cat# 115-035-003, RRID:AB_10015289	
Antibody	Peroxidase-AffiniPure anti-rabbit IgG (H+L)	Jackson ImmunoResearch Labs	Cat# 111-035-003, RRID:AB_2313567	
Recombinant DNA reagent	MISSION® TRC2 pLKO.5-puro Empty Vector (shControl)	Sigma-Aldrich	Cat# SHC201	
Recombinant DNA reagent	MISSION® shRNA plasmid DNA Lamin A/C (shLmna)	Sigma-Aldrich	TRCN0000317672	
Recombinant DNA reagent	MISSION® shRNA plasmid DNA Lamin A/C (shLmna-2)	Sigma-Aldrich	TRCN0000089849	
Sequence-based reagent	Control siRNA (siControl)	Bioneer	Sense: UUCUCCGAACGUG UCACGUtt; anti-sense: ACGUGACACGUUC GGAGAAtt	(Roman et al., 2017)
Sequence-based reagent	Lamin A/C siRNA (siLmna)	Bioneer	Sense: GGCUUGUGGAGAU CGAUAAtt; anti-sense: UUAUCGAUCUCCA CAAGCCgc	(Roman et al., 2017)
Peptide, recombinant protein	Fibronectin	Sigma-Aldrich	Cat# F1141	
Peptide, recombinant protein	Gelatin from bovine skin	Sigma-Aldrich	Cat# G9391	
Peptide, recombinant protein	Collagenase, Type 4	Worthington Biochemical	Cat# LS004188	
Peptide, recombinant protein	DNase I, grade II	Roche	Cat# 10104159001	
Peptide, recombinant protein	Recombinant murine HGF	PeproTech	Cat# 315-23	
Peptide, recombinant protein	Recombinant murine EGF	PeproTech	Cat# 315-09	
Peptide, recombinant protein	Recombinant murine VEGF$_{165}$ (VEGF-A)_	PeproTech	Cat# 450-32	
Peptide, recombinant protein	Recombinant murine SDF-1α (CXCL12)	PeproTech	Cat# 250-20A	
Peptide, recombinant protein	Fibronectin	Sigma-Aldrich	Cat# F1141	
Chemical compound, drug	Etoposide	Sigma-Aldrich	Cat# E1383	
Chemical compound, drug	Puromycin	Sigma-Aldrich	Cat# P8833	
Chemical compound, drug	CellTracker™ Orange CMTMR Dye	Thermo Fisher Scientific	Cat# C2927	
Chemical compound, drug	Hoechst 33342	Thermo Fisher Scientific	Cat# 62249	
Chemical compound, drug	Hoechst 33258	Sigma-Aldrich	Cat# 861405	

Chemical compound, drug	DAPI	Sigma-Aldrich	Cat# D9542	
Chemical compound, drug	X-Gal	Sigma-Aldrich	Cat# 11680293001	
Chemical compound, drug	Ethyl Cinnamate (ECi)	Acros Organics	Cat# 103-36-6	
Chemical compound, drug	cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail	Roche	Cat# 4693159001	
Chemical compound, drug	Poly(ethylene glycol) (PEG8000)	Sigma-Aldrich	Cat# 81268	
Chemical compound, drug	Triton X-100	Sigma-Aldrich	Cat# X100	
Commercial assay or kit	Click-iT™ RNA Alexa Fluor™ 594 Imaging Kit	Thermo Fisher Scientific	Cat# C10330Cat# 4693159001	
Commercial assay or kit	APC Annexin V Apoptosis Detection Kit with PI	BioLegend	Cat# 640932	
Commercial assay or kit	Dynabeads™ mRNA DIRECT™ Purification	Thermo Fisher Scientific	Cat# 61011	
Commercial assay or kit	SuperSignal™ West Pico PLUS Chemiluminescent Substrate	Thermo Fisher Scientific	Cat# 34577	
Software, algorithm	cellSens (v1.16)	Olympus	https://www.olympus-lifescience.com	
Software, algorithm	Imaris (v9.5.1)	Oxford Instruments	https://imaris.oxinst.com/packages	
Software, algorithm	CaseViewer 2.3	3DHISTECH	https://www.3dhistech.com/caseviewer	
Software, algorithm	Fiji	SciJava	https://fiji.sc	
Software, algorithm	CytExpert	Beckman Coulter	www.beckman.com	
Software, algorithm	Flowjo (v10.6.1)	FlowJo	https://flowjo.com	
Software, algorithm	Image Lab (v6.0.1)	Bio-Rad Laboratories	https://www.bio-rad.com	
Software, algorithm	GraphPad Prism 6	GraphPad Software	https://www.graphpad.com	
Software, algorithm	R	R Development Core Team	https://www.r-project.org	
Software, algorithm	UTAP pipeline	(Kohen et al., 2019)	https://utap.readthedocs.io	
Other	Lipofectamine™ 2000	Thermo Fisher Scientific	Cat# 11668	
Other	Lipofectamine™ RNAiMAX	Thermo Fisher Scientific	Cat# 3778075	
Other	Matrigel® Matrix	Corning	Cat# 356234	
Other	Agarose, low gelling temperature	Sigma-Aldrich	Cat# 9045	
Other	Cell Dissociation Solution	Biological Industries	Cat# 03-071-1B	
Other	Red Blood Cell Lysing Buffer Hybri-Max™	Sigma-Aldrich	Cat# R7757	
Other	DMEM, high glucose	Thermo Fisher Scientific	Cat# 41965	

Cells

Murine melanoma (B16F10) and Lewis Lung Carcinoma (LL/2) cells were grown in DMEM supplemented with 10% FBS. Murine breast adenocarcinoma cells (E0771) were grown in DMEM supplemented with 10% FBS, 1 mM sodium pyruvate and 10 mM HEPES. Human embryonic kidney (HEK293T) and murine brain endothelial (bEnd.3) cells were cultured in DMEM medium supplemented with 10% FBS and 2 mM L-glutamine.

Mice

Wild-type mice (WT) on C57BL/6 background were maintained in a pathogen-free facility and all animal procedures were approved by the Animal Care and Use Committee of the Weizmann Institute of Science. Male and female 7- to 8-week-old mice were used in all experiments.

Imaging and Analysis of Tumor Cell Transendothelial Migration

The transmigration assay of tumor cells was performed under shear-free conditions. Murine endothelial bEnd.3 cells (8×10^4) were seeded in a µ-Slide VI0.4 ibiTreat (ibidi), pre-coated with gelatin (1% in DDW) for 30 min at 37°C. A day later, B16F10 or E0771 cells were labeled with 20 µM Hoechst 33342 for 5 min at 37°C and resuspended in binding medium (Hank's balanced-salt solution 1X containing 2 mg/ml BSA and 10 mM HEPES, pH 7.4, supplemented with 1 mM $CaCl_2$ and 1 mM $MgCl_2$) and introduced in the ibidi chamber over a confluent bEnd.3 monolayer. Images were acquired at a rate of one frame every 4-5 min for 4 h using an IX83 inverted microscope (Olympus) equipped with UPlanFLN 20x/0.50 Ph1 ∞/0.17/FN 26.5 objective (Olympus), 49000-ET-DAPI filter set (Chroma). ORCA-Flash4.0LT camera, model: C11440-42U (Hamamatsu). Temperature was maintained at 37°C throughout the assay. For analysis of migratory phenotypes, tumor cells in different fields of view (10-15 cells per field) were individually tracked and categorized using cellSense software 1.16 (Olympus). Close monitoring of individual frames allowed the discrimination of transmigrating tumor cells (TEM) from tumor cells that failed to complete TEM either because of inability to protrude through endothelial junctions (SA) or squeeze their nuclei through these junctions and underneath the endothelial monolayer (SEP). See also Video 2. Notably, we did not observe intercalation of individual tumor cells in between ECs (Reymond et al., 2012). Fiji software was used to determine nuclear

circularity of transmigrating tumor cells at different frames of the time-lapse movies.
This software was also used to incorporate time codes, labels and scale bars into
video segments.

Tumor Cells Transwell Migration Assay
Fibronectin (1.5 µg/ml in PBS) was coated for 30 min at 37°C onto both sides (for
tumor chemotaxis assays) or only on the bottom side (for tumor haptotaxis assays) of
8 or 3 µm hanging cell culture inserts (Millipore, MCEP24H48 and MCSP24H48). After
washing the filters with PBS, B16F10 or E0771 cells (4×10^4) resuspended in DMEM
containing 0.1% BSA were introduced into the top chamber. DMEM with 0.1% BSA
was inserted in the lower chamber in the presence or absence of the chemoattractant.
After 4 or 24 h at 37°C with CO_2, the cells were fixed with paraformaldehyde (4% in
PBS) for 15 min and stained with crystal violet (3% in DDW) for additional 15 min, both
at RT. Cells on the upper side of the filter were scraped using a cotton swab whereas
cells located on the bottom side were imaged using a SZX16 stereo microscope
(Olympus) equipped with SDF PLAPO 1XPF objective (Olympus) set at 10X
magnification. DP73 camera (Olympus).

Generation of Stable shRNA-Expressing Clones
Lentiviruses were produced by co-transfecting HEK293T cells with the shControl or
shLmna vectors and three helper plasmids (Gag-Pol, Rev and VSV-G,) using
Lipofectamine® 2000. The virus-containing medium was harvested 48 or 72 h after
transfection and subsequently precleaned by a brief centrifugation at 600 x g and a
0.45 µm filtration. Viruses were collected and concentrated with a precipitation solution
(40% PEG8000 and 2.5N NaCl) as described (Guo et al., 2012) and stored at -20°C
overnight. A day later, the medium was thawed and centrifuged at 2,400 x g for 30 min
at RT. The viral pseudoparticles were resuspended in 200 µl culture medium and
mixed with 2.5×10^4 of tumor cells for 12 h. 36 h after viral infection, puromycin was
added to the culture medium at a concentration of 2.5 µg/ml (B16F10) or 3 µg/ml
(E0771) and tumor cells were selected and expanded, replacing growth medium every
48 h. The mean knockdown levels were assessed by Western blotting.

Transient siRNA transfection

B16F10 cells ($1x10^5$) were transfected with siRNA (20 nM) using Lipofectamine™

RNAiMAX (Thermo Fisher Scientific) following the manufacturer's instructions. 72 h

after transfection tumor cells were either lysed to quantify the mean knockdown protein

levels or utilized in experimental assays. Lamin A/C siRNA (siLmna) sequence used

in this study was described in (Roman et al., 2017).

Western Blotting

Cells were grown as described above, washed twice with ice-cold PBS, scraped into

lysis buffer (25 mM Tris pH 7.5, 1 mM EDTA, 0.5 mM EGTA, 150 mM NaCl, 1% NP-

40, 0.2% SDS, 2 mM Na_3VO_4, 1 mM NaF, 10 mM Nappi, 80 mM b-glycerol phosphate

and a protease inhibitor tablet), and kept on ice for 30 min with occasional vortexing.

Thereafter, lysates were centrifuged at 14,000 x g for 15 min at 4°C. The supernatant

was collected and protein concentration was determined by BCA protein assay

(Thermo Fisher Scientific). The protein suspension was separated by gel

electrophoresis followed by transfer to nitrocellulose membranes, and blocking with

non-fat milk (5% in PBS-T) for 1 h at RT. Immunoblotting was performed overnight at

4°C according to the manufacturers' guidelines. Antibody binding to membrane blots

was detected using horseradish peroxidase conjugated secondary antibodies for 1 h

at RT, followed by development with a chemiluminescence substrate (Thermo Fisher

Scientific). Chemiluminescence was detected using the ChemiDoc MP (Bio-Rad

Laboratories) imaging system.

Immunofluorescence Staining

B16F10 or E0771 cells ($1.5x10^4$) were seeded into a μ-Slide VI0.4 ibiTreat (ibidi) pre-

coated with fibronectin (10 μg/ml in PBS) for 30 min at 37°C. The next day cells were

rinsed with ice-cold PBS and fixed with paraformaldehyde (4% in PBS) for 15 min at

RT followed by permeabilization with Triton X-100 (0.25% in PBS) for 15 min at RT

and blocking with goat serum (10% in PBS) for 20 min at 37°C. The cells were then

incubated with anti-lamin A/C antibody (1:100) for 1 hour at RT, washed with ice-cold

PBS three times and incubated with an Alexa Fluor 488 conjugated secondary

antibody (1:200) for 1 h at RT. Cells were imaged using an IX83 inverted microscope

(described above) equipped with an UPlanFLN 40X 0.75 Ph2 ∞/0.17/FN 26.5 objective

(Olympus), 49002-ET-EGFP (FITC/Cy2) filter set (Chroma). Alternatively, cells were

fixed with paraformaldehyde (3% in PBS) at RT for 5 min followed by fixation in

methanol at -20°C for another 5 min. H3K9me3 was detected with rabbit monoclonal

anti H3K9me3 and DNA was stained with Hoechst 33258. Immunostaining images

were collected using an Olympus IX81 fluorescent microscope equipped with a

coolSNAP HQ2 CCD camera (Photometrics).

Imaging of Tumor Cell Nuclear Dynamics

B16F10 or E0771 cells ($2x10^4$) were trypsinized, labeled in suspension with 20 µM

Hoechst 33342, resuspended in binding medium (composition described above), and

introduced in a µ-Slide VI0.4 ibiTreat (ibidi) over a bEnd.3-deposited basement

membrane extracellular matrix. Images were acquired at a rate of one frame every 4-

5 min for 2 h using an IX83 Inverted Microscope (described above). Temperature was

kept at 37°C throughout the duration of the assay. Background was subtracted for the

fluorescent channel using cellSense 1.16 (Olympus) software. FiJi (SciJava) software

was used for title and time code labeling and determination of the nuclear circularity.

Light Sheet Fluorescent Microscopy of Tumor Cells and Lung Vasculature

B16F10 ($2x10^4$) or E0771 (10^4) cells labeled with CMTMR dye (Thermo Fisher

Scientific), 10 µM for 30 min according to the manufacturer's instructions, were

injected in the retro-orbital sinus of recipient mice. Euthanasia by administration of

sodium pentobarbital (200 mg/Kg) was practiced 3 hours later. Blood capillaries were

labeled 15 min before the animal sacrifice by intravenous injection of 6 µg of an Alexa

647-conjugated anti-CD31 mAb. Immediately after the sacrifice, mice were

transcardially perfused with PBS and the lungs inflated via the trachea with low gelling

agarose (Sigma-Aldrich), subsequently fixed with paraformaldehyde (4% in PBS) for

2 h, dehydrated and cleared using ethyl cinnamate as described in (Klingberg et al.,

2017). Cleared intact lung lobes, were imaged using an Ultramicroscope II (LaVision

BioTec) operated by the ImspectorPro software (LaVision BioTec). For excitation light

sheet was generated by a Superk Super-continuum white light laser (emission 460 nm

– 800 nm, 1 mW/nm – 3 (NKT photonics), followed by specific excitation filters per

channel. For detection optics microscope was equipped with a single lens

configuration - 4X objective - LVBT 4X UM2-BG, with an adjustable refractive index

collar set to the RI of 1.56. Images were acquired by an Andor Neo sCMOS camera

(2,560 × 2,160, pixel size 6.5 µm x 6.5 µm, Andor). Z stacks were acquired in 3 µm

steps. Channel configuration for GFP and EGFP excitation 470\40 emission 525\50,

for CMTMR, excitation 560\40 emission 630\75, and for CD31-AF647 excitation
640\30 emission 690\50.
567

Image Reconstruction and Analysis
Three-dimensional rendering of LSFM was performed via Imaris software (Oxford
Instruments). Surfaces of CMTMR-labeled tumor cells were created using volume
(comprised between 1000 and 25000 μm^3) and intensity (max of red fluorescent
channel) as defining features to unequivocally separate them from background
signals. Each cell was individually segmented and its distance was measured with
respect to the CD31-labeled blood vessels: intravascular, extravascular or protruding.
See also Videos 6-8.
576

Determination of Tumor cell Accumulation in Washed Lungs
B16F10 ($2x10^4$) or E0771 (10^4) cells labeled with 10 μM CMTMR for 30 min according
to the manufacturer's instructions, were resuspended in PBS and injected into the
retro-orbital sinus of recipient mice. Euthanasia by administration of sodium
pentobarbital (200 mg/Kg) was practiced 3 or 72 h later. Immediately thereafter, mice
were transcardially perfused with PBS and the lungs were extracted, minced and
incubated in RPMI-1640 containing collagenase type 4 (1.5 mg/ml) and DNase I (20
μg/ml) at 37°C for 45 min. Lung cell suspensions were pushed through a 100 μm cell
strainer and centrifuged at 0.2 x g or 5 min at 4°C. RBCs were subsequently lysed
with an RBC lysis buffer (Sigma Aldrich). The cells were resuspended in ice-cold
FACS buffer (PBS with 1% BSA, 0.1% sodium azide and 5 mM EDTA), filtered through
a 70 μm strainer and analyzed using a CytoFLEX flow cytometer (Beckman Coulter).
589

5-Ethynyl Uridine (EU) Labeling of Cultured Cells
B16F10 or E0771 (either shControl or shLmna expressing) cells were grown on serum
coated glass coverslips. 1 mM EU was added to the culture medium and cells were
kept for 1 h at 37°C in 5% CO_2. Cells were then washed with PBS and fixed with 3.7%
formaldehyde for 15 min at RT. The fixative was removed and cells were washed twice
with PBS, followed by permeabilization with 0.5% Triton X-100 for 15 min at RT. Cells
were washed twice with PBS and incorporated EU was detected by click chemistry
using a fluorescent azide following the manufacturer's guidelines of Click-iT RNA
Imaging Kit (Thermo Fisher Scientific). Following the Click-iT reaction (30 min at RT

in the dark), cells were rinsed twice with a rinse buffer. Nuclear staining was performed
with Hoechst 33342. Cells were imaged using a Zeiss LSM 710 point scanning
confocal microscope with stacking acquisition and generation of maximum intensity
projection images. Nucleoplasmic fluorescence intensity measurements were
performed in Fiji, and statistical significance was assessed through Mann-Whitney U,
two-tailed, test.

Bulk MARS-seq protocol and sequencing

RNA was isolated from 10,000 cells from each cell line using Dynabeads® mRNA
Direct Kit (Thermo Fisher Scientific). Libraries for RNA-seq were prepared using a
modified version of TranSeq, as described (Jaitin et al., 2014). Briefly, RNA was
reversed transcribed with MARS-seq barcoded RT primers in a 10 µl volume with the
Affinity Script kit (Agilent). Reverse transcription was analyzed by qRT-PCR and
samples with a similar CT were pooled (up to eight samples per pool). Each pool was
treated with Exonuclease I (NEB) for 30 min at 37°C and cleaned by 1.2× volumes of
SPRI beads (Beckman Coulter). Next, the cDNA was converted to double-stranded
DNA with a second strand synthesis kit (NEB) in a 20 ml reaction, incubating for 2 h at
16 °C. The product was purified with 1.4× volumes of SPRI beads, eluted in 8 µl and
in vitro transcribed (with the beads) at 37 °C overnight for linear amplification using the
T7 High Yield RNA polymerase IVT kit (NEB). Following IVT, the DNA template was
removed with Turbo DNase I (Ambion) 15 min at 37 °C and the amplified RNA (aRNA)
purified with 1.2 volumes of SPRI beads. The aRNA was fragmented by incubating
3 min at 70 °C in Zn^{2+} RNA fragmentation reagents (Ambion) and purified with 2×
volumes of SPRI beads. The aRNA was ligated to the MARS-seq ligation adapter with
T4 RNA Ligase I (NEB). The reaction was incubated at 22 °C for 2 h. After 1.5× SPRI
cleanup, the ligated product was reverse transcribed using Affinity Script RT enzyme
(Agilent) and a primer complementary to the ligated adapter. The reaction was
incubated for 2 min at 42°C, 45 min at 50°C, and 5 min at 85°C. The cDNA was purified
with 1.5× volumes of SPRI beads. The library was completed and amplified through a
nested PCR reaction with 0.5 mM of P5_Rd1 and P7_Rd2 primers and PCR ready mix
(Kappa Biosystems). The amplified pooled library was purified with 0.7× volumes of
SPRI beads to remove primer leftovers. Library concentration was measured with a
Qubit fluorometer (Life Technologies) and mean molecule size was determined with a

2200 TapeStation instrument. RNA-seq libraries were sequenced using the Illumina

NextSeq® 500 High Output v2 Kit (75 cycles).

Bioinformatics analysis

Samples were demultiplexed using the barcode present in the R2 read. The analysis

was performed using the UTAP pipeline (Kohen et al., 2019). In brief, UMI sequences

present in the R2 read were inserted in the read name of R1 sequence file using a

python script. Cutadapt was used to trim low quality, poly A and adapter sequences

(Martin, 2011), (parameters: -a AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC

-a "A{10}" –times 2 -u 3 -u -3 -q 20 -m 25). Sequences were mapped to the UCSC

mm10 mouse genome using STAR (Dobin et al., 2013) v2.4.2a (parameters: –

alignEndsType EndToEnd, –outFilterMismatchNoverLmax 0.05, –twopassMode

Basic, –alignSoftClipAtReferenceEnds No). The pipeline quantified the 3' of RefSeq

annotated genes (1,000 bases upstream of the 3' end and 100 bases downstream)

using HTSeq count (Anders, Pyl, & Huber, 2015) and a modified Refseq gtf file

(downloaded from igenomes UCSC). UMI information was integrated into the BAM

files as tags, using a python script. UMI counting was performed after marking

duplicates (in-house script) using a modified HTSeq-count. DESeq2 (Love, Huber, &

Anders, 2014) was used for normalization and detection of differentially expressed

genes. Raw p-values were adjusted for multiple testing using the procedure of

Benjamini and Hochberg. Genes were considered to be differentially expressed if their

mean normalized expression was greater than 5, the absolute value of the

log2FoldChange was greater than 1, and the adjusted p-value was less than 0.05.

Batch effects were removed from the read counts using the ComBat function from the

sva R package (Johnson, Li, & Rabinovic, 2007). The normalized, batch corrected and

log2 transformed read counts outputted by ComBat were used to draw plots. K-means

clustering of differentially expressed genes was performed using the Partek®

Genomics Suite® software, version 6.6 (Partek Inc., St. Louis, MO, USA). Functional

analysis of the differentially expressed genes was performed using

https://metascape.org/gp/index.html#/main/step1, Metascape (Zhou et al., 2019).

Significantly enriched GO terms for up and downregulated genes were extracted from

this analysis. Sequencing data have been deposited in NCBI's Gene Expression

Omnibus and are accessible through GEO Series accession number GSE146730.

Cell Growth on 2D Culture Plates

B16F10 or E0771 cells were seeded at a low density of 5×10^3/well in a 6-well plate. Puromycin-containing growth medium was replaced every 24 h throughout the duration of the assay (72 h). To determine the number of viable cells proliferating on the plates, cells were trypsinized and counted by flow cytometer every 24 h.

Cell Growth in 3D Soft Agar

B16F10 or E0771 cells were suspended at a density of 4×10^3/ml of 0.3% low gelling agarose in DMEM +10% FBS and equilibrated at 4°C for 15 min in 12 well plate as described (Horibata et al., 2015). The cell agar suspension was warmed to 37°C and cultured in a humidified incubator containing 5% CO_2. Spheroid formation was monitored on the first, third and 6th day using a IX83 inverted microscope (Olympus) equipped with UPlanFLN 4x/0.13 Ph1 ∞/-/FN 26.5 objective (Olympus). In parallel, cancer cell colonies grown directly on the bottom of each well were also images. To isolate individual cells from the spheroids grown in 3D conditions, the layer of agarose containing the spheroids was mechanically removed and solubilized in PBS pre-warmed at 45°C. Spheroids were spun down at 0,8 x g for 5 min at RT, trypsinized with warm trypsin B to recover single cells and stained with annexin V and propidium iodide following the manufacturer instructions.

Primary Tumor Growth in vivo

A suspension of 1.5×10^4 B16F10 or 10^4 E0771 (either shControl or shLmna expressing) cells in 50 µl of Matrigel® Matrix mixed with PBS (at 1:1 v/v), was inoculated subcutaneously in the flank (B16F10) or in the mammary fat pad (E0771) of recipient mice. Tumor size (volume) was assessed throughout the duration of the experiment, by vernier caliper measurements of length (L) and width (W) and the tumor volume (V) was calculated using the formula: V = (L x W x W)/2. After 14 days, the animals were euthanized by CO_2 inhalation and the tumor was extracted, weighted and fixed in paraformaldehyde (4% in PBS).

Experimental Lung Metastases

B16F10 (4×10^4) or E0771 (10^4 or 2×10^4) either shControl or shLmna expressing cells were suspended in 200 µl PBS + 0.25 mM EDTA and injected into the tail vein of recipient mice. Animals were euthanized by administration of sodium pentobarbital

(200 mg/Kg) 14 or 28 days later. Immediately after the sacrifice, mice were
transcardially perfused with PBS and the lungs were extracted and visually analyzed
for the presence of surface metastatic foci, subsequently stored in 4% PFA for 24 h
and 1% PFA at 4°C for long term storage. Paraffin embedding and H&E staining of 5
μm-thick sections were performed by the histology core unit of the Weizmann Institute
of Science. Sections were digitalized using a Pannoramic SCAN II (3DHISTECH) and
analyzed using CaseViewer software (3DHISTECH).
707

Senescence-Associated-β-Galactosidase (SA-β-Gal) Activity
For SA-β-gal activity assay, B16F10 cells (shControl or shLmna) were treated with 5
μM etoposide for 72 h, washed and left in regular culture medium (containing 2.5 μg/ml
of puromycin) for an additional 120 h. Cells were then fixed with 0.5% glutaraldehyde
in PBS for 15 min, washed twice with PBS supplemented with 1 mM $MgCl_2$ (pH 5.5)
and stained with PBS containing 1 mM $MgCl_2$ (pH 5.5) and supplemented with 1 mg/ml
X-gal, 5 mM $K_3Fe[CN]_6$ and 5 mM $K_4Fe[CN]_6$ for 5 h at 37°C in the dark, as described
(Krizhanovsky et al., 2008). Cells were washed with warm PBS, fixed with 4%
paraformaldehyde and imaged using aSZX16 stereo microscope (Olympus) equipped
with SDF PLAPO 1XPF objective (Olympus) set at 10X magnification. DP73 camera
(Olympus).
719

R-loops quantification
B16F10 cells were grown on sterile coverslips and R-loops were detected with the
S9.6 antibody following cell fixation and permeabilization with 100% ice-cold methanol
and acetone for 10 min and 1 min on ice, respectively. Incubation with primary
antibodies was followed by incubation with Dy488-secondary antibody (Bethyl). Nuclei
were stained using DAPI and coverslips were assembled in Vectashield Mounting
Medium (H-1000, Vector Laboratories). All the washing steps were done with PBS
containing 0.05% (v/v) Tween 20. Images were acquired using Confocal Laser Point-
Scanning Microscope Zeiss LSM 710. Fluorescence intensity of the nucleoplasmic
staining of 90 cells from three independent experiments was assessed for each
condition, and statistical significance was assessed through Mann-Whitney U, two-
tailed, test.
732

Statistical analysis

Data in graphs are represented as means ± SEM. All tests of statistical significance were performed using a two-tailed Student's t-test with GraphPad Prism software, with the exception of analysis of Figure 6C and Figure 5–figure supplement 1 for which Mann-Whitney U, two-tailed, test was used. Significance was set to $p < 0.05$. Statistical details of experiments can be found in the figure legends. For figures 6D and 7D, raw p-values were adjusted for multiple testing using the procedure of Benjamini and Hochberg while genes were considered to be differentially expressed if their mean normalized expression was greater than 5, the absolute value of the log2FoldChange was greater than 1, and the adjusted p-value was less than 0.05.

Acknowledgements

We thank Edgar Gomes (Instituto de Medicina Molecular, Lisbon, Portugal), Valery Krizhanovsky and Avri Ben-Ze'Ev (Weizmann Institute) for providing reagents and for helpful discussions. The RNA-Seq work was performed under the guidance of Drs. Merav Kedmi and Hadas Keren-Shaul from the Sandbox unit of the Life Science Core Facility of Weizmann Institute of Science. R.A. is the Incumbent of the Linda Jacobs Chair in Immune and Stem Cell Research. His research is supported by the Israel Science Foundation (grant no. 791/17), the Minerva Foundation, Germany, GIF (grant number I-1470-412.13/2018), as well as grants from the Moross Integrated Cancer Center, Yeda-Sela Center for Basic Research, Helen and Martin Kimmel Institute for Stem Cell Research, the Meyer Henri Cancer Endowment, and from William and Marika Glied and Carol A. Milett. S.N and S.M. are funded by the Wellcome Trust (grant 098291/Z/12/Z to S.N.). G.G's research is supported by an Israel Cancer Association grant no. 20201181.

Competing interests

The authors declare no competing interests.

Author Contributions

Francesco Roncato, Methodology, Validation, Formal analysis, Investigation, Data curation, Writing - original draft, Writing - review and editing, Visualization; Ofer Regev, Methodology, Investigation; Sara W. Feigelson, Validation, Investigation; Writing - original draft, Writing - review and editing, Supervision; Sandeep Kumar Yadav,

Investigation; Lukasz Kaczmarczyk, Validation, Investigation, Formal Analysis,

Visualization; Nehora Levi, Validation, Investigation, Formal Analysis, Visualization;

Diana Drago-Garcia, Software, Formal Analysis, Visualization; Samuel Ovadia,

Investigation; Marina Kizner, Investigation; Yoseph Addadi, Supervision, Writing-

original draft; João C. Sabino, Validation, Investigation, Formal Analysis, Visualization;

Yossi Ovadya, Resources, Investigation; Sérgio F. de Almeida, Resources,

Supervision, Writing-original draft; Ester Feldmesser, Software, Formal analysis,

Visualization; Gabi Gerlitz, Resources, Supervision, Writing-original draft; Ronen Alon,

Conceptualization, Writing - original draft, Writing - review & editing, Supervision,

Project administration, Funding acquisition.

References

Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq-A Python framework to work with high-
throughput sequencing data. *Bioinformatics*, *31*(2), 166–169.
https://doi.org/10.1093/bioinformatics/btu638

Aureille, J., Belaadi, N., & Guilluy, C. (2017). Mechanotransduction via the nuclear envelope:
a distant reflection of the cell surface. *Current Opinion in Cell Biology*, *44*, 59–67.
https://doi.org/10.1016/j.ceb.2016.10.003

Barney, L. E., Jansen, L. E., Polio, S. R., Galarza, S., Lynch, M. E., & Peyton, S. R. (2016,
February 1). The predictive link between matrix and metastasis. *Current Opinion in*
Chemical Engineering, Vol. 11, pp. 85–93. https://doi.org/10.1016/j.coche.2016.01.001

Barzilai, S., Yadav, S. K., Morrell, S., Roncato, F., Klein, E., Stoler-Barak, L., … Alon, R.
(2017). Leukocytes Breach Endothelial Barriers by Insertion of Nuclear Lobes and
Disassembly of Endothelial Actin Filaments. *Cell Reports*, *18*(3), 685–699.
https://doi.org/10.1016/j.celrep.2016.12.076

Becker, J. S., Nicetto, D., & Zaret, K. S. (2016). H3K9me3-Dependent Heterochromatin:
Barrier to Cell Fate Changes. *Trends in Genetics*, *32*(1), 29–41.
https://doi.org/10.1016/J.TIG.2015.11.001

Bell, E. S., & Lammerding, J. (2016). Causes and consequences of nuclear envelope
alterations in tumour progression. *European Journal of Cell Biology*, *95*(11), 449–464.
https://doi.org/10.1016/j.ejcb.2016.06.007

Bishop, J. R., Schuksz, M., & Esko, J. D. (2007). Heparan sulphate proteoglycans fine-tune
mammalian physiology. *Nature*, *446*(7139), 1030–1037. https://doi.org/nature05817
[pii]10.1038/nature05817

Borrego-Pinto, J., Jegou, T., Osorio, D. S., Auradé, F., Gorjánácz, M., Koch, B., … Gomes, E.
R. (2012). Samp1 is a component of TAN lines and is required for nuclear movement.
Journal of Cell Science, *125*(5), 1099–1105. https://doi.org/10.1242/jcs.087049

Broers, J. L., Raymond, Y., Rot, M. K., Kuijpers, H., Wagenaar, S. S., & Ramaekers, F. C.
(1993). Nuclear A-type lamins are differentially expressed in human lung cancer
subtypes. *The American Journal of Pathology*, *143*(1), 211–220. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/8391215

Broers, J. L. V., & Ramaekers, F. C. S. (2014). The role of the nuclear lamina in cancer and
apoptosis. In *Advances in Experimental Medicine and Biology* (Vol. 773, pp. 27–48).
https://doi.org/10.1007/978-1-4899-8032-8_2

Bronshtein, I., Kepten, E., Kanter, I., Berezin, S., Lindner, M., Redwood, A. B., … Garini, Y.
(2015). Loss of lamin A function increases chromatin dynamics in the nuclear interior.
Nature Communications, *6*(1), 8044. https://doi.org/10.1038/ncomms9044

Burke, B., & Stewart, C. L. (2013). The nuclear lamins: flexibility in function. *Nature Reviews*
Molecular Cell Biology, *14*(1), 13–24. https://doi.org/10.1038/nrm3488

Cao, X., Moeendarbary, E., Isermann, P., Davidson, P. M., Wang, X., Chen, M. B., … Shenoy,
V. B. (2016). A Chemomechanical Model for Nuclear Morphology and Stresses during
Cell Transendothelial Migration. *Biophysical Journal*, *111*(7), 1541–1552.
https://doi.org/10.1016/J.BPJ.2016.08.011

Capo-chichi, C. D., Cai, K. Q., Smedberg, J., Ganjei-Azar, P., Godwin, A. K., & Xu, X. X.
(2011). Loss of A-type lamin expression compromises nuclear envelope integrity in
breast cancer. *Chin J Cancer*, *30*(6), 415–425.

Chambers, A. F. A. F., Groom, A. C. A. C., & MacDonald, I. C. I. C. (2002). Dissemination and
growth of cancer cells in metastatic sites. *Nat Rev Cancer*, *2*(8), 563–572.
https://doi.org/10.1038/nrc865

Chambliss, A. B., Khatau, S. B., Erdenberger, N., Robinson, D. K., Hodzic, D., Longmore, G.
D., & Wirtz, D. (2013). The LINC-anchored actin cap connects the extracellular milieu to
the nucleus for ultrafast mechanotransduction. *Sci Rep*, *3*, 1087.
https://doi.org/10.1038/srep01087

Chang, W., Folker, E. S., Worman, H. J., & Gundersen, G. G. (2013). Emerin organizes actin
flow for nuclear movement and centrosome orientation in migrating fibroblasts. *Molecular*
Biology of the Cell, *24*(24), 3869–3880. https://doi.org/10.1091/mbc.e13-06-0307

Cho, S., Vashisth, M., Abbas, A., Majkut, S., Vogel, K., Xia, Y., … Discher, D. E. (2019).
Mechanosensing by the Lamina Protects against Nuclear Rupture, DNA Damage, and
Cell-Cycle Arrest. *Developmental Cell*, *49*(6), 920-935.e5.
https://doi.org/10.1016/j.devcel.2019.04.020

Crossley, M. P., Bocek, M., & Cimprich, K. A. (2019, February 7). R-Loops as Cellular

Regulators and Genomic Threats. *Molecular Cell*, Vol. 73, pp. 398–411.

https://doi.org/10.1016/j.molcel.2019.01.024

Dai, W., Jiang, Y., Chen, K., Qiu, J., Sun, J., Zhang, W., … Li, W. (2017). Effect of etoposide-

induced alteration of the Mdm2-Rb signaling pathway on cellular senescence in A549

lung adenocarcinoma cells. *Oncology Letters*, *14*(4), 3935–3940.

https://doi.org/10.3892/ol.2017.6684

Davidson, P. M., Denais, C., Bakshi, M. C., & Lammerding, J. (2014). Nuclear deformability

constitutes a rate-limiting step during cell migration in 3-D environments. *Cell Mol Bioeng*,

7(3), 293–306. https://doi.org/10.1007/s12195-014-0342-y

Dechat, T., Adam, S. A., Taimen, P., Shimi, T., & Goldman, R. D. (2010). Nuclear Lamins.

Cold Spring Harbor Perspectives in Biology, *2*(11), a000547–a000547.

https://doi.org/10.1101/cshperspect.a000547

Denais, C., & Lammerding, J. (2014). Nuclear mechanics in cancer. *Adv Exp Med Biol*, *773*,

435–470. https://doi.org/10.1007/978-1-4899-8032-8_20

Denais, C. M., Gilbert, R. M., Isermann, P., McGregor, A. L., Te Lindert, M., Weigelin, B., …

Lammerding, J. (2016). Nuclear envelope rupture and repair during cancer cell migration.

Science, *352*(6283), 353–358. https://doi.org/10.1126/science.aad7297

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., … Gingeras, T. R.

(2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.

https://doi.org/10.1093/bioinformatics/bts635

Donnaloja, F., Carnevali, F., Jacchetti, E., & Raimondi, M. T. (2020). Lamin A/C

Mechanotransduction in Laminopathies. *Cells*, *9*(5), 1306.

https://doi.org/10.3390/cells9051306

Dou, Z., Xu, C., Donahue, G., Shimi, T., Pan, J.-A., Zhu, J., … Berger, S. L. (2015). Autophagy

mediates degradation of nuclear lamina. *Nature*, *527*(7576), 105–109.

https://doi.org/10.1038/nature15548

Dyer, L., & Patterson, C. (2010). Development of the Endothelium: An Emphasis on

Heterogeneity. *Seminars in Thrombosis and Hemostasis*, *36*(03), 227–235.

https://doi.org/10.1055/s-0030-1253446

Elosegui-Artola, A., Andreu, I., Beedle, A. E. M., Lezamiz, A., Uroz, M., Kosmalska, A. J., …

Roca-Cusachs, P. (2017). Force Triggers YAP Nuclear Entry by Regulating Transport

across Nuclear Pores. *Cell*, *171*(6), 1397-1410.e14.

https://doi.org/10.1016/j.cell.2017.10.008

Etienne-Manneville, S., & Lammerding, J. (2017). Connecting the plasma membrane to the

nucleus by intermediate filaments. *Molecular Biology of the Cell*, *28*(6), 695–696.

https://doi.org/10.1091/mbc.E16-11-0794

Fernández-Morera, J. L., Calvanese, V., Rodríguez-Rodero, S., Menéndez-Torre, E., & Fraga,

M. F. (2010). Epigenetic regulation of the immune system in health and disease. *Tissue*
Antigens, *76*(6), 431–439. https://doi.org/10.1111/j.1399-0039.2010.01587.x

Friedl, P., Wolf, K., & Lammerding, J. (2011). Nuclear mechanics during cell migration. *Current*
Opinion in Cell Biology, *23*(1), 55–64. https://doi.org/10.1016/j.ceb.2010.10.015

Fruleux, A., & Hawkins, R. J. (2016). Physical role for the nucleus in cell migration. *J Phys*
Condens Matter, *28*(36), 363002. https://doi.org/10.1088/0953-8984/28/36/363002

Gonzalez-Granado, J. M., Silvestre-Roig, C., Rocha-Perugini, V., Trigueros-Motos, L.,
Cibrian, D., Morlino, G., … Andres, V. (2014). Nuclear envelope lamin-A couples actin
dynamics with immunological synapse architecture and T cell activation. *Sci Signal*,
7(322), ra37. https://doi.org/10.1126/scisignal.20048727/322/ra37 [pii]

Gonzalez-Suarez, I., Redwood, A. B., Perkins, S. M., Vermolen, B., Lichtensztejin, D.,
Grotsky, D. A., … Gonzalo, S. (2009). Novel roles for A-type lamins in telomere biology
and the DNA damage response pathway. *The EMBO Journal*, *28*(16), 2414–2427.
https://doi.org/10.1038/emboj.2009.196

Gonzalo, S. (2014). DNA damage and lamins. *Advances in Experimental Medicine and*
Biology, *773*, 377–399. https://doi.org/10.1007/978-1-4899-8032-8_17

Gorelik, L., & Flavell, R. A. (2001). Immune-mediated eradication of tumors through the
blockade of transforming growth factor-β signaling in T cells. *Nature Medicine*, *7*(10),
1118–1122. https://doi.org/10.1038/nm1001-1118

Graham, D. M., Andersen, T., Sharek, L., Uzer, G., Rothenberg, K., Hoffman, B. D., …
Burridge, K. (2018). Enucleated cells reveal differential roles of the nucleus in cell
migration, polarity, and mechanotransduction. *The Journal of Cell Biology*, *217*(3), 895–
914. https://doi.org/10.1083/jcb.201706097

Guo, P., El-Gohary, Y., Prasadan, K., Shiota, C., Xiao, X., Wiersch, J., … Gittes, G. K. (2012).
Rapid and simplified purification of recombinant adeno-associated virus. *Journal of*
Virological Methods, *183*(2), 139–146. https://doi.org/10.1016/j.jviromet.2012.04.004

Harada, T., Swift, J., Irianto, J., Shin, J. W., Spinler, K. R., Athirasala, A., … Discher, D. E.
(2014). Nuclear lamin stiffness is a barrier to 3D-migration, but softness can limit survival.
Proceedings of the IEEE Annual Northeast Bioengineering Conference, NEBEC, *2014-*
Decem(5), 669–682. https://doi.org/10.1109/NEBEC.2014.6972810

Harr, J. C., Luperchio, T. R., Wong, X., Cohen, E., Wheelan, S. J., & Reddy, K. L. (2015).
Directed targeting of chromatin to the nuclear lamina is mediated by chromatin state and
A-type lamins. *The Journal of Cell Biology*, *208*(1), 33–52.
https://doi.org/10.1083/jcb.201405110

Heemskerk, N., Schimmel, L., Oort, C., van Rijssel, J., Yin, T., Ma, B., … van Buul, J. D.
(2016). F-actin-rich contractile endothelial pores prevent vascular leakage during
leukocyte diapedesis through local RhoA signalling. *Nature Communications*, *7*.

https://doi.org/Artn 1049310.1038/Ncomms10493

Horibata, S., Vo, T. V., Subramanian, V., Thompson, P. R., & Coonrod, S. A. (2015). Utilization
of the soft agar colony formation assay to identify inhibitors of tumorigenicity in breast
cancer cells. *Journal of Visualized Experiments*, *2015*(99), 52727.
https://doi.org/10.3791/52727

Infante, E., Castagnino, A., Ferrari, R., Monteiro, P., Agüera-González, S., Paul-Gilloteaux,
P., … Chavrier, P. (2018). LINC complex-Lis1 interplay controls MT1-MMP matrix digest-
on-demand response for confined tumor cell migration. *Nature Communications*, *9*(1), 1–
13. https://doi.org/10.1038/s41467-018-04865-7

Jaiswal, D., Cowley, N., Bian, Z., Zheng, G., Claffey, K. P., & Hoshino, K. (2017). Stiffness
analysis of 3D spheroids using microtweezers. *PLoS ONE*, *12*(11), e0188346.
https://doi.org/10.1371/journal.pone.0188346

Jaitin, D. A., Kenigsberg, E., Keren-Shaul, H., Elefant, N., Paul, F., Zaretsky, I., … Amit, I.
(2014). Massively parallel single-cell RNA-seq for marker-free decomposition of tissues
into cell types. *Science*, *343*(6172), 776–779. https://doi.org/10.1126/science.1247651

Johnson, W. E., Li, C., & Rabinovic, A. (2007). Adjusting batch effects in microarray
expression data using empirical Bayes methods. *Biostatistics*, *8*(1), 118–127.
https://doi.org/10.1093/biostatistics/kxj037

Joyce, J. A., & Pollard, J. W. (2009). Microenvironmental regulation of metastasis. *Nature*
Reviews Cancer, *9*(4), 239–252. https://doi.org/10.1038/nrc2618

Kaufmann, S. H., Mabry, M., Jasti, R., & Shaper, J. H. (1991). Differential expression of
nuclear envelope lamins A and C in human lung cancer cell lines. *Cancer Research*,
51(2), 581–586. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/1985776

Khatau, S. B., Hale, C. M., Stewart-Hutchinson, P. J., Patel, M. S., Stewart, C. L., Searson, P.
C., … Wirtz, D. (2009). A perinuclear actin cap regulates nuclear shape. *Proc Natl Acad*
Sci U S A, *106*(45), 19017–19022. https://doi.org/0908686106
[pii]10.1073/pnas.0908686106

Khuon, S., Liang, L., Dettman, R. W., Sporn, P. H. S., Wysolmerski, R. B., & Chew, T.-L.
(2010). Myosin light chain kinase mediates transcellular intravasation of breast cancer
cells through the underlying endothelial cells: a three-dimensional FRET study. *Journal*
of Cell Science, *123*(3), 431–440. https://doi.org/10.1242/jcs.053793

Kim, D. H., & Wirtz, D. (2015). Cytoskeletal tension induces the polarized architecture of the
nucleus. *Biomaterials*, *48*, 161–172. https://doi.org/10.1016/j.biomaterials.2015.01.023

Kirby, T. J., & Lammerding, J. (2018). Emerging views of the nucleus as a cellular
mechanosensor. *Nature Cell Biology*, *20*(4), 373–381. https://doi.org/10.1038/s41556-
018-0038-y

Klingberg, A., Hasenberg, A., Ludwig-Portugall, I., Medyukhina, A., Männ, L., Brenzel, A., …

Gunzer, M. (2017). Fully Automated Evaluation of Total Glomerular Number and
Capillary Tuft Size in Nephritic Kidneys Using Lightsheet Microscopy. *Journal of the*
American Society of Nephrology : JASN, *28*(2), 452–459.
https://doi.org/10.1681/ASN.2016020232

Kohen, R., Barlev, J., Hornung, G., Stelzer, G., Feldmesser, E., Kogan, K., … Leshkowitz, D.
(2019). UTAP: User-friendly Transcriptome Analysis Pipeline. *BMC Bioinformatics*, *20*(1).
https://doi.org/10.1186/s12859-019-2728-2

Kong, L., Schafer, G., Bu, H., Zhang, Y., Zhang, Y., & Klocker, H. (2012). Lamin A/C protein
is overexpressed in tissue-invading prostate cancer and promotes prostate cancer cell
growth, migration and invasion through the PI3K/AKT/PTEN pathway. *Carcinogenesis*,
33(4), 751–759. https://doi.org/10.1093/carcin/bgs022

Krizhanovsky, V., Yon, M., Dickins, R. A., Hearn, S., Simon, J., Miething, C., … Lowe, S. W.
(2008). Senescence of activated stellate cells limits liver fibrosis. *Cell*, *134*(4), 657–667.
https://doi.org/10.1016/j.cell.2008.06.049

Krull, S., Dörries, J., Boysen, B., Reidenbach, S., Magnius, L., Norder, H., … Cordes, V. C.
(2010). Protein Tpr is required for establishing nuclear pore-associated zones of
heterochromatin exclusion. *The EMBO Journal*, *29*(10), 1659–1673.
https://doi.org/10.1038/emboj.2010.54

Kubben, N., Voncken, J. W., Demmers, J., Calis, C., van Almen, G., Pinto, Y., & Misteli, T.
(2010). Identification of differential protein interactors of lamin a and progerin. *Nucleus*,
1(6), 1–13. https://doi.org/10.4161/nucl.1.6.13512

Lammerding, J., Fong, L. G., Ji, J. Y., Reue, K., Stewart, C. L., Young, S. G., & Lee, R. T.
(2006). Lamins a and C but not lamin B1 regulate nuclear mechanics. *Journal of*
Biological Chemistry, *281*(35), 25768–25780. https://doi.org/10.1074/jbc.M513511200

Liu, B., Wang, Z., Zhang, L., Ghosh, S., Zheng, H., & Zhou, Z. (2013). Depleting the
methyltransferase Suv39h1 improves DNA repair and extends lifespan in a progeria
mouse model. *Nature Communications*, *4*(1), 1–12. https://doi.org/10.1038/ncomms2885

Lomakin, A. J., Cattin, C. J., Cuvelier, D., Alraies, Z., Molina, M., Nader, G. P. F., … Piel, M.
(2020). The nucleus acts as a ruler tailoring cell responses to spatial constraints. *Science*,
370(6514). https://doi.org/10.1126/SCIENCE.ABA2894

Lombardi, M. L., & Lammerding, J. (2011). Keeping the LINC: the importance of
nucleocytoskeletal coupling in intracellular force transmission and cellular function.
Biochemical Society Transactions, *39*(6), 1729–1734.
https://doi.org/10.1042/BST20110686

Love, M. I. M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol*, *15*(12), 550.
https://doi.org/10.1186/s13059-014-0550-8

Maizels, Y., Elbaz, A., Hernandez-Vicens, R., Sandrusy, O., Rosenberg, A., & Gerlitz, G.
(2017). Increased chromatin plasticity supports enhanced metastatic potential of mouse
melanoma cells. *Experimental Cell Research*, *357*(2), 282–290.
https://doi.org/10.1016/J.YEXCR.2017.05.025

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.Journal*, *17*(1), 10. https://doi.org/10.14806/ej.17.1.200

Miles, F. L., Pruitt, F. L., van Golen, K. L., & Cooper, C. R. (2008). Stepping out of the flow:
capillary extravasation in cancer metastasis. *Clinical & Experimental Metastasis*, *25*(4),
305–324. https://doi.org/10.1007/s10585-007-9098-2

Ofer, N., Mogilner, A., & Keren, K. (2011). Actin disassembly clock determines shape and
speed of lamellipodial fragments. *Proc Natl Acad Sci U S A*, *108*(51), 20394–20399.
https://doi.org/1105333108 [pii]10.1073/pnas.1105333108

Raab, M., Gentili, M., de Belly, H., Thiam, H. R., Vargas, P., Jimenez, A. J., … Piel, M. (2016).
ESCRT III repairs nuclear envelope ruptures during cell migration to limit DNA damage
and cell death. *Science*, *352*(6283), 359–362. https://doi.org/science.aad7611 [pii]
10.1126/science.aad7611

Reymond, N., Im, J. H., Garg, R., Vega, F. M., Borda d'Agua, B., Riou, P., … Ridley, A. J.
(2012). Cdc42 promotes transendothelial migration of cancer cells through β1 integrin.
The Journal of Cell Biology, *199*(4), 653–668. https://doi.org/10.1083/jcb.201205169

Roman, W., Martins, J. P., Carvalho, F. A., Voituriez, R., Abella, J. V. G., Santos, N. C., …
Gomes, E. R. (2017). Myofibril contraction and crosslinking drive nuclear movement to
the periphery of skeletal muscle. *Nature Cell Biology*, *19*(10), 1189–1201.
https://doi.org/10.1038/ncb3605

Rowat, A. C., Jaalouk, D. E., Zwerger, M., Ung, W. L., Eydelnant, I. A., Olins, D. E., …
Lammerding, J. (2013). Nuclear envelope composition determines the ability of
neutrophil-type cells to passage through micron-scale constrictions. *J Biol Chem*,
288(12), 8610–8618. https://doi.org/10.1074/jbc.M112.441535

Scaffidi, P., & Misteli, T. (2006). Lamin A-dependent nuclear defects in human aging. *Science*,
312(5776), 1059–1063. https://doi.org/10.1126/science.1127168

Shin, J. W., Spinler, K. R., Swift, J., Chasis, J. A., Mohandas, N., & Discher, D. E. (2013).
Lamins regulate cell trafficking and lineage maturation of adult human hematopoietic
cells. *Proc Natl Acad Sci U S A*, *110*(47), 18892–18897.
https://doi.org/10.1073/pnas.13049961101304996110 [pii]

Solovei, I., Wang, A. S., Thanisch, K., Schmidt, C. S., Krebs, S., Zwerger, M., … Joffe, B.
(2013). LBR and Lamin A/C Sequentially Tether Peripheral Heterochromatin and
Inversely Regulate Differentiation. *Cell*, *152*(3), 584–598.
https://doi.org/10.1016/j.cell.2013.01.009

Stephens, P. J., Tarpey, P. S., Davies, H., Van Loo, P., Greenman, C., Wedge, D. C., … Stratton, M. R. (2012). The landscape of cancer genes and mutational processes in breast cancer. *Nature, 486*(7403), 400–404. https://doi.org/10.1038/nature11017

Sullivan, T., Escalante-Alcalde, D., Bhatt, H., Anver, M., Bhat, N., Nagashima, K., … Burke, B. (1999). Loss of A-type lamin expression compromises nuclear envelope integrity leading to muscular dystrophy. *Journal of Cell Biology*, *147*(5), 913–919. https://doi.org/10.1083/jcb.147.5.913

Swift, J., Ivanovska, I. L., Buxboim, A., Harada, T., Dingal, P. C., Pinter, J., … Discher, D. E. (2013). Nuclear lamin-A scales with tissue stiffness and enhances matrix-directed differentiation. *Science*, *341*(6149), 1240104. https://doi.org/10.1126/science.1240104341/6149/1240104 [pii]

Towbin, B. D., Meister, P., & Gasser, S. M. (2009). The nuclear envelope--a scaffold for silencing? *Current Opinion in Genetics & Development*, *19*(2), 180–186. https://doi.org/10.1016/j.gde.2009.01.006

Valastyan, S., & Weinberg, R. A. (2011). Tumor Metastasis: Molecular Insights and Evolving Paradigms. *Cell*, *147*(2), 275–292. https://doi.org/10.1016/j.cell.2011.09.024

Vargas, J. D., Hatch, E. M., Anderson, D. J., & Hetzer, M. W. (2012). Transient nuclear envelope rupturing during interphase in human cancer cells. *Nucleus*, *3*(1), 88–100. https://doi.org/10.4161/nucl.18954

Venables, R. S., McLean, S., Luny, D., Moteleb, E., Morley, S., Quinlan, R. A., … Hutchison, C. J. (2001). Expression of individual lamins in basal cell carcinomas of the skin. *British Journal of Cancer*, *84*(4), 512–519. https://doi.org/10.1054/bjoc.2000.1632

Wazir, U., Ahmed, M. H., Bridger, J. M., Harvey, A., Jiang, W. G., Sharma, A. K., & Mokbel, K. (2013). The clinicopathological significance of lamin A/C, lamin B1 and lamin B receptor mRNA expression in human breast cancer. *Cellular and Molecular Biology Letters*, *18*(4), 595–611. https://doi.org/10.2478/s11658-013-0109-9

Wilson, K. L., & Berk, J. M. (2010). The nuclear envelope at a glance. *Journal of Cell Science*, *123*(12), 1973–1978. https://doi.org/10.1242/jcs.019042

Wilson, Katherine L, & Foisner, R. (2010). Lamin-binding Proteins. *Cold Spring Harbor Perspectives in Biology*, *2*(4), a000554. https://doi.org/10.1101/cshperspect.a000554

Wolf, K., Te Lindert, M., Krause, M., Alexander, S., Te Riet, J., Willis, A. L., … Friedl, P. (2013). Physical limits of cell migration: control by ECM space and nuclear deformation and tuning by proteolysis and traction force. *J Cell Biol*, *201*(7), 1069–1084. https://doi.org/jcb.201210152 [pii]10.1083/jcb.201210152

Yadav, S. K., Feigelson, S. W., Roncato, F., Antman-Passig, M., Shefi, O., Lammerding, J., & Alon, R. (2018). Frontline Science: Elevated nuclear lamin A is permissive for granulocyte transendothelial migration but not for motility through collagen I barriers. *Journal of*

1058   *Leukocyte Biology*, *104*(2), 239–251. https://doi.org/10.1002/JLB.3HI1217-488R

1059 Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., … Chanda,

1060   S. K. (2019). Metascape provides a biologist-oriented resource for the analysis of

1061   systems-level datasets. *Nature Communications*, *10*(1), 1523.

1062   https://doi.org/10.1038/s41467-019-09234-6

1063

Figure legends

Figure 1. Downregulation of lamin A/C increases melanoma squeezing through small rigid pores but does not facilitate tumor transendothelial migration in vitro

(A) Expression levels of lamin A/C and lamin B1 in B16F10 cells transduced with control or lamin A/C shRNA. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used as loading control. (B) Immunostaining of lamin A/C (green) in B16F10 shControl and shLmna cells. Scale bar, 20 µm. (C) Chemotactic migration of B16F10 shControl and shLmna cells through 8 or 3 µm pores transwell filters in presence (+) or absence (-) of HGF (50 ng/ml) for 4 hours. Values represent the mean ± SEM of five fields in each experimental group. Results shown are from a representative experiment of three. ***p = 0.0002 (D) Haptotactic migration of B16F10 shControl and shLmna cells through 8 or 3 µm pores transwell filters in the presence or absence of fibronectin for 4 hours. Values represent the mean ± SEM of five fields in each experimental group. Results shown are from a representative experiment of three. *p = 0.0107, **p = 0.0024. (E) Representative images of distinct tumor cell categories (referred to as migratory phenotypes) taken from time-lapse videomicroscopy segments of individual B16 cells: round, spread above (SA), forming sub endothelial pseudopodia (SEP), and completing transendothelial migration (TEM). Scale bar represents 20 µm. See also Video 2. (F) Migratory phenotypes of B16F10 melanoma, E0771 breast carcinoma and LL/2 Lewis Lung Carcinoma interacting with bEnd.3 monolayers. Values represent the mean ± SEM of three fields in each experimental group (n= 40 cells). The experiment shown is representative of three.

Figure 2. Downregulation of lamin A/C does not facilitate melanoma TEM but increases tumor nucleus deformability

(A) Migratory phenotypes of control and lamin A/C knockdown B16F10 melanoma (shControl and shLmna, respectively) interacting with unstimulated bEnd.3 cells. Values represent the mean ± SEM of four fields in each experimental group (n = 50). The experiments shown are representative of three. (B) Serial images of representative control and lamin A/C knockdown B16F10 melanoma cells, labeled with Hoechst. Time intervals are depicted in each image. The contours of the melanoma cell's leading edge and nucleus are outlined in each image in

yellow and red respectively. See also Video 3. (C) Nuclear circularity variation during the

distinct indicated phases (1-3) of transendothelial migration of B16F10 shControl and shLmna

cells. Values represent the mean ± SEM of six fields in each experimental group (n = 15). The

experiment shown is representative of three. *p = 0.0105 (shControl); ****p < 0.0001, *p =

0.0288 (shLmna). The right panel depicts the mean nuclear circularity values for each group

of B16F10 cells. The percent changes in mean circularity values are shown on top of the line

plots. *p = 0.0152. (D) Nuclear circularity of B16F10 shControl and shLmna cells spread on a

bEnd.3-derived basement membrane. n = 50 cells per group. Values are mean ± SEM. ****p

< 0.0001. See also Video 4.

1103

Figure 3. In vivo melanoma crossing of lung capillaries is not enhanced by lamin A/C

downregulation

(A) Schematic representation of the LSFM analysis. (B) Visualization of 3D bronchial

structures (green, autofluorescence) together with B16 melanoma cells (red, CMTMR) and

alveolar capillaries (cyan, CD31). Scale bars, 100 µm. See also Video 5. (C) Representative

3D images of intravascular, extravascular, and protrusive tumor cells across the CD31-labeled

lung vasculature together with the percentage of B16F10 shControl and shLmna cells present

in a volume of 5×10^9 µm^3 of the left lung lobe (3 h after injection) counted using Imaris software.

The results shown are representative of 3 experiments. Scale bar, 100 µm. See also Videos

6-8. (D) Number of CMTMR-labeled B16F10 shControl and shLmna present in the lungs of

recipient mice, 3 h after i.v. injection. Data are mean ± SEM. The experiment shown is

representative of two.

1116

Figure 4. Lamin A/C downregulation reduces heterochromatin content and alters gene

transcription

(A) Immunostaining of H3K9me3 (red) in B16F10 shControl and shLmna cells. Hoechst 33258

(cyan) is shown in the left panel. Scale bar, 25 µm. The experiment shown is representative

of three. (B) Equal amounts of proteins from B16F10 shControl or shLmna cells, separated by

SDS-PAGE and analyzed for the indicated proteins by Western blot analysis. The bar graph

represents the mean levels of H3K9me3, H3K27me3 and SUV39H2 normalized to Histone

H3 and of SETDB1 normalized to α-Tubulin ± SEM of at least four independent experiments.

*p < 0.05. (C) Fluorescence microscopy imaging of 5-ethynyl uridine (EU) incorporation (red)

and Hoechst 33342 (blue) in B16F10 shControl and shLmna cells. Cells were grown with 1

mM EU for 1 h. Cells were fixed, permeabilized and treated with Alexa Fluor 594 azide.

Nucleoplasmic fluorescence intensity of the EU staining was measured using ImageJ. Data

from 3 independent experiments are shown in the boxplot. (D) Differentially downregulated

(blue), upregulated (red) and non-differentially expressed genes (grey) are shown (LFC > 1

and adjusted p value < 0.05). The names of the top 20 most significantly expressed genes are indicated in the plot. (E) Gene ontology (GO) enrichment analysis of the top differentially downregulated (blue) and upregulated (red) genes in B16F10 shLmna cells. Biological Process (BP), Molecular Function (MF) and Cellular Component (CC).

Figure 5. Melanoma proliferation, response to DNA damage, and senescence are insensitive to lamin A/C downregulation

(A) In vitro cell growth of B16F10 shControl and shLmna cells cultured with growth medium changed every 24 h. Values represent the mean ± SEM. The experiment shown is representative of three. (B) Growth arrest induced by etoposide at different concentrations on B16F10 shControl and shLmna cells after treatment for 72 h. Values represent the mean ± SEM. The experiment shown is representative of three. (C) Senescence associated β-galactosidase staining of B16F10 shControl and shLmna cells treated with 5 µM etoposide for 72 h, washed and cultured in regular growth medium for an additional 120 h. The experiment shown is representative of two. Scale bar, 20 µm. *p = 0.0470. (D) Schematic representation of the 3D agarose spheroid assay. Spheroids (dark grey) grown inside agarose while few cells migrate and start dividing at the bottom of the culture dish. At day 6, spheroids were recovered. Representative images of the spheroids derived from B16F10 shControl cells grown in 3D agar (supplemented with 10% FBS) imaged on days 3 and 6. Scale bar, 20 µm. An image of a colony of cells proliferating on the bottom of the soft agar well is depicted at the right. Scale bar, 200 µm. (E) Spheroid diameter, measured on days 1, 3 and 6 after seeding. n = 50 spheroids per group. Values represent the mean ± SEM. The experiment shown is representative of three. ****p < 0.0001. Inset depicts the area of individual cell colonies proliferating on the bottom of the soft agar well, measured on day 6. (F) The diameter of individual spheroids grown for 1 and 3 days in soft agar supplemented with rich growth medium (50% FBS). Values represent the mean ± SEM. The experiment shown is representative of two. ****p < 0.0001.

Figure 6. Effects of lamin A/C downregulation on E0771 migration, growth and metastatic potential

(A) 15,000 B16F10 shControl and shLmna cells were implanted subcutaneously in flank or in the mammary fat pad (B) of C57BL/6 mice. Tumor growth was assessed every other day for 14 days post implantation. The experiment shown is representative of three. (C) Number of CMTMR-labeled B16F10 shControl and shLmna present in the lungs of recipient mice, 3 or 7 days after i.v. injection. Data are mean ± SEM. The experiment shown is representative of three. (D) 40,000 B16F10 shControl and shLmna cells were injected in the tail of recipient C57BL/6 mice. After 14 days, animals were euthanized and lungs harvested. Surface

1168 metastatic foci were macroscopically counted, results are mean ± SEM, n = 5 for each
1169 experimental group. *p = 0.0374. A representative lung image from each group is presented.
1170 The experiment shown is representative of three.

1171

Figure 7. Effects of lamin A/C downregulation on E0771 migration, growth and
metastatic potential

(A) Expression levels of lamin A/C and lamin B1 in E0771 cells transduced with control or
lamin A/C shRNA. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used as
loading control. (B) Haptotactic migration of E0771 shControl and shLmna cells through 8 or
3 µm pores transwell filters in the presence (+) or absence (-) of fibronectin for 4 h. Values
represent the mean ± SEM of five fields of view in each experimental group. Results shown
are from a representative experiment of three. *p = 0.0240 (8 µm) *p = 0.0361 (3 µm). (C)
Representative 3D images of intravascular or protrusive tumor cells across the CD31-labeled
lung vasculature together with the percentage of E0771 shControl and shLmna cells present
in a volume of 5×10^9 µm^3 of the left lung lobe (3 h after injection) counted using Imaris software.
The results shown are representative of 3 experiments. Scale bar, 100 µm. (D) Differentially
downregulated (blue), upregulated (red) and non-differentially expressed genes (grey) are
shown (LFC > 1 and adjusted p value < 0.05). The names of the top 20 most significantly
expressed genes are indicated in the plot. (E) In vitro cell growth of E0771 shControl and
shLmna cells cultured with growth medium changed every 24 hours. Values represent the
mean ± SEM. The experiment shown is representative of two. (F) 10,000 E0771 shControl
and shLmna cells were implanted in the mammary fat pad of female C57BL/6 mice. Tumor
growth was assessed every other day for 14 days post implantation. The experiment shown
is representative of three. (G) Growth arrest induced by etoposide at different concentrations
on E0771 shControl and shLmna cells after treatment for 72 hours. Values represent the mean
± SEM. The experiment shown is representative of three. (H) Spheroid diameter, measured
on days 1, 3 and 6 after seeding. n = 50 cells per group. Values represent the mean ± SEM.
The experiment shown is representative of three. ****p < 0.0001. (I) Experimental lung
metastasis of E0771 breast cancer cells. 20,000 E0771 shControl and shLmna cells were
intravenously injected in recipient C57BL/6 mice. After 14 days, animals were euthanized and
lungs harvested. The number of micrometastases present in the left lung lobe of each lung
was determined. The experiment shown is representative of two. **p = 0.0030. (J) 10,000
E0771 shControl and shLmna cells were intravenously injected in recipient C57BL/6 mice.
After 30 days, animals were euthanized and lungs harvested. The number of micrometastases
present in the left lung lobe of each mouse was determined. The experiment shown is
representative of two. *p = 0.0246.

Video Legends

Video 1

Transendothelial migration of B16F10 murine melanoma

Time-lapse movie of an Hoechst-labeled B16F10 crossing a bEnd.3 endothelial monolayer. The contours of the tumor cell leading edges and nucleus are outlined in each image in yellow and red respectively. Elapsed time is designated as h:mm:ss. Scale bar, 20 μm

Video 2

Migratory phenotypes of B16F10 cells over a bEnd.3 endothelium

Time-lapse movie divided in four quadrants in which different B16F10 cells display a unique phenotype when interacting with a bEnd.3 endothelial monolayer. Represented clockwise there are Round, Spread Above (SA), subendothelial pseudopodium (SEP) and transendothelial migration (TEM) phenotypes. Elapsed time is designated as h:mm:ss. Scale bar, 20 μm

Video 3

Transendothelial migration of B16F10 shControl vs shLmna cells

Time-lapse movie depicting a Hoechst labeled B16F10 cell shControl (left) and shLmna (right) crossing a bEnd.3 endothelium. The contours of the tumor cell leading edges and nucleus are outlined in yellow and red respectively. Elapsed time is designated as h:mm:ss. Scale bar, 20 μm.

Video 4

Nuclear deformability of B16F10 shControl vs shLmna cells

Time-lapse movie depicting the Hoechst-labeled nuclei (green) of B16F10 shControl (left) and shLmna (right) cells interacting with a bEnd.3-deposited basement membrane. Elapsed time is designated as h:mm:ss. Scale bar, 20 μm.

Video 5

Light sheet microscopy of tumor cells, bronchial structures and lung vasculature

Three-dimensional animated visualization of a section of murine lung lobe. CMTMR-labeled tumor cells (red) and autofluorescent bronchial structures (green) can be observed from seconds 0 to 21 (movie length). CD31-labeled lung vasculature (cyan) can be observed from seconds 22-40 (movie length).

Video 6

Example of an intravascular B16F10 cell

Three-dimensional animated visualization of a CMTMR-labeled B16F10 cell (red) located inside a CD31-labeled lung vasculature (cyan). Scale bar, 100 μm.

Video 7

Example of a protruding B16F10 cell

Three-dimensional animated visualization of a CMTMR-labeled B16F10 cell (red) protruding through the CD31-labeled lung vasculature (cyan). Scale bar, 100 μm

Video 8

Example of an extravascular B16F10 cell

1241   Three-dimensional animated visualization of a CMTMR-labeled B16F10 cell (red) located outside the

1242   CD31-labeled lung vasculature (cyan). Scale bar, 100 μm

1243   **Supplementary File 1**

1244   DeSeq2 analysis output and gene expression levels in enriched Gene Ontology terms.

Figure 1

A

B

Figure 1–figure supplement 1. Lamin A/C deficiency increases B16 melanoma cell chemotaxis through rigid pores.

(A) Haptotactic of B16F10 shControl vs shLmna-2 through 3 µm pores transwell filters coated (+) or uncoated (-) with fibronectin (1.5 µg/ml) for 4 h. **** $p < 0.0001$. (B) Migration of B16F10 cells towards HGF (50 ng/ml) measured through 8 or 3 µm pore transwell filters after 4 h. Data are mean ± SEM. The results shown are representative of 2 independent experiments. **$p = 0.0076$.

Figure 2

Figure 3

Figure 3–figure supplement 1. Quantification of cancer cells accumulation in lungs.
Scheme depicting the experimental pipeline necessary to quantify single cell accumulation in lungs of recipient mice. Cancer cells were labeled with CMTMR orange cell tracker for 30 min, washed and i.v. injected into recipient WT mice. After 3 hours, 3 or 7 days the mice were euthanized, their lungs harvested, minced and digested into single cell suspension and subsequently analyzed by flow cytometry.

Figure 4

Figure 4–figure supplement 1. Lamin A/C downregulation alters gene transcription.
(A) Heatmap based on clustering of the 300 differentially expressed genes. Red and blue bars represent positive and negative changes, respectively, and the intensity of the color represents the standardized log2 expression level. The top differentially expressed (DE) genes are shown on the left. The full list is available in Table S1. (B, C) Individual gene expression levels of genes belonging to the enriched gene ontology (GO) terms: nuclear matrix (B) and nuclear body (C), summarized in Fig. 6E.

Figure 5

Figure 5–figure supplement 1. DNA damage-induced growth arrest and senescence of cancer cells induced by etoposide treatment in vitro.

Cancer cells were exposed to 5 µM etoposide for 72 hours. The compound was subsequently removed and the cells were cultured with regular growth medium for additional 120 hours. Cells were then fixed in 0.5% glutaraldehyde for 15 min and stained for senescence associated β-Galactosidase as described in Krizhanovsky et al., 2008.

Figure 5–figure supplement 2. Lamin A/C downregulation doesn't alter R-loop (S9.6) formation in B16F10 cells.
R-loop mapping via S9.6-based anti DNA:RNA hybrid antibody immunofluorescence.

Figure 5–figure supplement 3. Soft agar colony formation assay
B16F10 (either shControl or shLmna-2 expressing) cells were embedded in soft agar and their growth was monitored for 6 days. Spheroid diameter was measured at days 1, 3, and 6 after seeding. n = 50 cells per group. Values represent the mean ± SEM. The experiment shown is representative of two. ****p < 0.0001.

Figure 5–figure supplement 4. Representative flow cytometry data to illustrate gating and quadrant strategy for measurement of apoptosis.
Flow cytometry plots showing annexin V (X-axis) and propidium iodide (Y-axis) staining of B16F10 (either shControl or shLmna expressing) cells derived from 3D colonies grown in soft agar and extracted at day 6. The right lower quadrant represents annexin V positive/propidium iodide (PI) negative staining indicating early apoptosis. The right upper quadrant represents high annexin V and high PI staining indicating late apoptosis and the left upper quadrant represents low annexin V and high PI staining indicating necrosis. The left lower quadrant indicates viable cells.

Figure 6

Figure 7

Figure 7–figure supplement 1. Downregulation of lamin A/C increases E0771 breast carcinoma cells squeezing through small rigid pores and alters nuclear shape.

(A) Haptotactic of E0771 shControl vs shLmna-2 through 3 µm pores transwell filters coated (+) or uncoated (-) with Fibronectin (1.5 µg/ml) for 4 h. *** p = 0.0002. (B) Haptotactic migration of E0771 Control and Lmna siRNA cells (72 h post transfection), through 8 or 3 µm pores transwell filters coated (+) or uncoated (-) with fibronectin (1.5 µg/ml) for 4 h. Values represent the mean ± SEM of five fields of view in each experimental group. Results shown are from a representative experiment of three. ***p = 0.0018. (C) Nuclear circularity of E0771 shControl and shLmna cells spread on a bEnd.3-derived basement membrane. ****p <0.0001. (D) Immunostaining of lamin A/C (green) in E0771 shControl and shLmna cells. Scale bar represents 20 µm.

Figure 7–figure supplement 2. In vitro and in vivo breast carcinoma crossing of endothelial barriers is not enhanced by lamin A/C downregulation.

(A) Migratory phenotypes of E0771 breast carcinoma cells TEM. Distinct tumor cell categories (referred to as migratory phenotypes) taken from time lapse videomicroscopy segments of individual E0771 cells: round, spread above (SA), forming sub endothelial pseudopodia (SEP), and completing transendothelial migration (TEM). Values represent the mean ± SEM of three fields in each experimental group. (B) Percentage of E0771 shControl and shLmna cells in a volume of 5×10^9 μm^3 of the left lung lobe (3 days after injection). Values are mean ± SEM of 3 different lung's sections fields of view in each experimental group.

Figure 7–figure supplement 3. Lamin A/C downregulation reduces heterochromatin content and alters gene transcription.

(A) Equal protein amounts from E0771 shControl or shLmna cells, separated by SDS-PAGE and analyzed for the indicated proteins by Western blot analysis. The bar graph represents the mean levels of H3K9me3, H3K27me3 and SUV39H2 normalized to Histone H3 and of SETDB1 normalized to α-Tubulin ± SEM of at least four independent experiments. *p < 0.05. (B) Heatmap based on clustering of 290 differentially expressed genes. Red and blue bars represent positive and negative changes, respectively, and the intensity of the color represents the standardized log2 expression level. (C) Gene ontology (GO) enrichment analysis of the top differentially downregulated (blue) and upregulated (red) genes in E0771 shLmna cells. Biological Process (BP), Molecular Function (MF) and Cellular Component (CC).

Figure 7–figure supplement 4. Fluorescence microscopy imaging of 5-ethynyl uridine (EU) incorporation (red) and Hoechst 33342 (blue) in E0771 shControl and shLmna cells.

Cells were grown with 1 mM EU for 1 h. The cells were fixed, permeabilized and treated with Alexa Fluor 594 azide. Nucleoplasmic fluorescence intensity of the EU staining was measured using ImageJ. Data from 3 independent experiments are shown in the boxplot.

Figure 7–figure supplement 5. Primary E0771 growth in the mammary fat pad or in the skin.

(A) Histology sections of breast primary tumors. A suspension of 10^4 E0771 (either shControl or shLmna expressing) cells in 50 µl of Matrigel® Matrix mixed with PBS (at 1:1 v/v) was inoculated in the mammary fat pad of recipient mice. 14 days later, animals were euthanized, tumors extracted, and fixed in 4% PFA. (B) Number of E0771 shControl and shLmna present in the lungs of recipient mice 3 hours, 3 and 7 days after i.v. injection. n = 3 for each experimental group. Data are mean ± SEM. The experiment shown is representative of three. (B) 10,000 E0771 shControl and shLmna cells were implanted subcutaneously in flank of C57BL/6 mice. Tumor growth was assessed every other day for 14 days post implantation. The experiment shown is representative of two.

Figure 7–figure supplement 6. E0771 accumulation in recipient lungs.
Number of E0771 shControl and shLmna recovered in the lungs of recipient mice 3 hours, 3 and 7 days after i.v. injection of 20,000 cells. n = 3 or 4 for each experimental group. Data are mean ± SEM. The experiment shown is representative of two.