De novo deletion mutations at recombination hotspots in mouse germlines

Agnieszka Lukaszewicz[1], Julian Lange[2], Scott Keeney[2,3], Maria Jasin[1]

[1]Developmental Biology Program, [2]Molecular Biology Program, and [3]Howard Hughes Medical Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA

To whom correspondence should be addressed:
Scott Keeney: s-keeney@ski.mskcc.org
Maria Jasin: m-jasin@ski.mskcc.org, Lead contact

Abstract

Numerous DNA double-strand breaks (DSBs) arise genome-wide during meiosis to ensure recombination between homologous chromosomes, which is required for gamete formation[1,2]. The ATM kinase plays a central role in controlling both the number and position of DSBs[3-5], but the consequences of deregulated DSB formation have not been explored. Here we discovered that an unanticipated type of DNA deletion arises at meiotic recombination hotspots in the absence of ATM. Deletions form via joining of ends from two closely-spaced DSBs at adjacent hotspots or within a single hotspot. Deletions are also detected in normal cells, albeit at much lower frequency, revealing that the meiotic genome has a hidden potential for deletion events. Remarkably, a subset of deletions contain insertions that likely originated from DNA fragments released from hotspots on other chromosomes. Moreover, although deletions form primarily within one chromosome, joining between homologous chromosomes is also observed. This predicts in turn gross chromosome rearrangements, with evidence of damage to multiple chromatids and aborted gap repair. Thus, multiple nearby meiotic DSBs are normally suppressed by ATM to protect genomic integrity. We expect the *de novo* germline mutations we observe to affect human health and genome evolution.

Meiotic recombination is essential for faithful genome transmission in most sexually reproducing
organisms[2]. Recombination is initiated by the programmed formation of DNA double-strand breaks
(DSBs) by the SPO11 protein[6]. SPO11 preferentially cuts within narrow genomic regions called
hotspots[7,8], usage of which is tightly regulated. Of all DSB hotspots (~15,000 defined in mouse), only a
tiny fraction (~250) experiences a DSB in any single meiosis[9-11]. DSBs are also nonrandomly dispersed
along chromosomes to ensure homolog pairing and recombination[3]. This quantitative and spatial
regulation of hotspot usage involves hierarchical and combinatorial levels of control[9,12,13], with the
ATM kinase playing a central role[5,14].
ATM, known for its role in the DNA damage response in mitotic cells[15], is activated by meiotic
DSBs to limit DSB numbers and to control hotspot usage genome-wide[4,9]. Control of DSB numbers by
ATM and its orthologs is conserved among organisms, although ATM-deficient mice have the largest
increase in DSB levels (e.g., ~10 fold compared with ~2 fold in budding yeast deficient for the ATM
ortholog Tel1)[5,16,17]. Control of genome-wide DSB distributions is also conserved[18]. In yeast, Tel1
controls DSB formation locally by suppressing double cutting, that is, simultaneous DSB formation at
adjacent hotspots[14,19]. This suppression is strongest at closely-spaced hotspots (<10 kb), such that in
the absence of Tel1, double cutting at hotspots separated by ~2 kb is more frequent than expected by
chance and ~10-fold higher than in wild type. The loss of this Tel1-mediated DSB suppression is
thought to contribute to the global increase in DSBs[18,19]. In mouse, double cutting at adjacent hotspots
has not been demonstrated, but potentially may be even more likely in the absence of ATM,
considering the extremely elevated DSB numbers. Accordingly, we have previously suggested that
ATM may inhibit multiple nearby DSBs[4,9].
The consequences of deregulated meiotic DSB formation on genomic architecture are not fully
understood. Normally, most meiotic DSBs are repaired by recombination between homologous
chromosomes to promote their segregation in meiosis I. We speculate that in the absence of ATM, too
many DSBs and closely-spaced DSBs interfere with the meiotic recombination program, eliciting the
use of other repair pathways, in particular, mutagenic nonhomologous end-joining (NHEJ), and thus
compromising genome integrity. While NHEJ is a robust mitotic DSB repair pathway in mice[20], it is
thought to normally be suppressed in meiosis[21]. However, NHEJ can occur in mouse spermatocytes
when DSBs are induced by irradiation[22-24], and it is known to operate in *C. elegans* in meiotic mutants
in which DSB processing[25-27] or recombination[28,29] is compromised. In principle, because ATM/Tel1
regulates meiotic DSB resection[30-34], resection defects could favor the use of NHEJ. Moreover, the loss
of ATM may cause DSBs to form at similar locations on multiple chromatids – due to the increase in
DSB numbers and/or the loss of spatial control – and thus promote NHEJ by damaging repair
templates.
In this report, we ask whether multiple DSBs form at adjacent hotspots in the absence of ATM
in mouse spermatocytes and whether they are repaired aberrantly to delete the intervening sequences.
Further, we ask whether multiple DSBs leading to deletions can occur within an individual hotspot. In
the course of these experiments, we uncover novel DSB (mis)repair products that provide insight into
the consequences of loss of DSB control during meiosis.
85
86
87 **Results**
88
89 **Hidden potential of the meiotic genome to form deletions**
90

We first catalogued genomic locations at risk for deletion by identifying pairs of closely-spaced hotspots in nucleotide-resolution DSB maps obtained by sequencing SPO11-bound oligonucleotides (oligos), a by-product of DSB formation[9]. We found that numerous DSB hotspots are separated by a short genomic distance (**Fig. 1A**). For example, ~6% of 13,436 autosomal SPO11-oligo hotspots in wild-type mice are located within 5 kb of another hotspot, and this percentage nearly doubles (~10%) in $Atm^{-/-}$ mice, which have more called hotspots (20,417) (**Fig. S1A**). Most hotspots in $Atm^{-/-}$ mice are shared with those in wild type, although a significant fraction are newly called because the few SPO11 oligos that cluster at these sites in wild type are greatly increased in the mutant[9] (**Fig. S1A**). Newly called hotspots often emerge near shared hotspots, resulting in an increased number of closely-spaced hotspots in $Atm^{-/-}$ (**Fig. S1B**). Moreover, in addition to the number, the strength of the shared hotspot pairs also disproportionately increases in $Atm^{-/-}$ (**Fig. S1C**), at least in part due to the particularly large increase in DSBs at weaker hotspots[9]. Thus, while adjacent DSBs may normally be suppressed, this hidden potential for DSBs at adjacent hotspots may be revealed in $Atm^{-/-}$.

To test for deletions, we selected DSB hotspot pairs that are shared between wild type and mutant and that contain strong hotspots to increase the chance of simultaneous DSB formation. The chosen hotspot pairs on Chr1 and Chr19 are separated by <2 kb, with individual hotspots that rank in the top quartile of all hotspots in both wild type and $Atm^{-/-}$ (**Fig. 1B, S1D, S2A**). We tested for deletions between the two hotspots in each pair by performing nested PCR on testis DNA. Concurrent DSB formation at the hotspot centers with precise joining of the DNA ends would give rise to 1.8-kb deletions, although deletions could vary in size depending on the location of SPO11 cleavage in any particular cell (**Fig. 1B, S2A**). Genomic DNA was digested with restriction enzymes prior to amplification in some experiments to suppress amplification of parental DNA. Nested PCR was performed with input testis DNA equivalent to ~8,000–16,000 haploid genomes per well for a total of ~10–15 million haploid genomes per genotype (**Table S1**).

Smaller PCR products indicative of deletions between the two DSB hotspots were observed in $Atm^{-/-}$ (**Fig. 1C, S2B**). In adult $Atm^{-/-}$ mice, the frequencies of deletions at the Chr1 and Chr19 hotspot pairs were 9.14×10^{-6} and 4.25×10^{-6}, respectively, per haploid genome (**Fig. 1D; Table S1**). Deletions were also detected in ATM-proficient cells but were rare. At the Chr19 hotspot pair, the frequencies were 0.15×10^{-6} and 0.31×10^{-6} in wild type and $Atm^{+/-}$, respectively, reflecting a total of just six events, while no deletions were observed in $Spo11^{-/-}$. For the Chr1 hotspot pair, only two deletions were observed in ATM-proficient cells, although one was also observed in $Spo11^{-/-}$.

Similar results were obtained in juvenile testes at both hotspot pairs at 15 days post partum, when wild-type and mutant testes have a similar distribution of prophase I stages. This finding indicates that the increase in deletion frequency is not due to the different cellular composition of mutant testes caused by prophase I arrest of $Atm^{-/-}$ spermatocytes[35,36] (**Table S1**).

It should be noted that testis DNA is derived from germ cells at different stages of development as well as from some somatic cells, such that the germline deletion frequencies we report are underestimated for all genotypes. Even so, meiotic recombination at a well-studied hotspot that has a similar strength as the individual hotspots examined here occurs at a substantially higher frequency (two to three orders of magnitude[37]) than the deletion frequency we observed at the hotspot pairs in the absence of ATM. However, considering the large number of hotspots located close to one another, the potential to form deletions in meiosis is substantial.

Double cutting in the absence of ATM

To map deletion breakpoints, we sequenced deletion products from $Atm^{-/-}$ adults and juveniles: 134
from the Chr1 hotspot pair and 58 from the Chr19 hotspot pair (**Fig. 2A, S2C; Table S2**). The
distribution of deletion breakpoints resembled the distribution of SPO11 oligos at both hotspots, with
breakpoints clustered around hotspot centers (**Fig. 2B,C, S2D,E**). Smoothed breakpoint and SPO11-
oligo profiles displayed similar shapes, except that breakpoint profiles were slightly wider with more
prominent smaller peaks (**Fig. 2D, S2F**). We defined the breakpoint center within each hotspot as the
position of the smoothed peak in the breakpoint density, in the same manner as hotspot centers were
defined[9]. Breakpoint centers closely matched both hotspot centers (within 13 to 42 bp). These findings
strongly indicate that deletion formation is preceded by concurrent SPO11 cleavage at the two
hotspots, providing the first evidence for double cutting between nearby DSB hotspots in mouse.
The median deletion lengths were similar to the distances between the DSB hotspot centers
(Chr1: 1954 bp vs 1825 bp; Chr19: 1892 bp vs 1763 bp), as expected from SPO11 cleavage of both
hotspots and joining of unprocessed or minimally processed DNA ends. Accordingly, a substantial
number of deletions (Chr1, 21%; Chr19, 19%) had both breakpoints within the narrow (251 bp) central
zones around the hotspot centers that contained the majority (>60%) of SPO11 oligos (**Fig. S3**). The
distribution of events was similar when the restriction digest step was omitted (**Fig. S4**), indicating that
this step does not eliminate other types of deletions. About half of the other deletions had one
breakpoint within the central zone of one hotspot but not the other (Chr1, 41%; Chr19, 45%) (**Fig. S3**).
Breakpoints outside the zone were not evenly distributed, such that some clusters could be observed
that overlapped with, but were more prominent than, the SPO11-oligo distributions (**Fig. 2B–D, S2D–**
F). While the similar distributions of breakpoints and DSBs suggests that DSB ends are usually joined
close to the initial SPO11 cleavage sites, the wider distribution of deletion breakpoints may reflect
preferential deletion formation for DSBs further from the hotspot center, DNA end processing prior to
joining, or even multiple DSBs within the same hotspot (see below).
The few deletions identified in wild type and $Atm^{+/-}$ often had breakpoints at or near the hotspot
centers, including some that were identical to those observed in $Atm^{-/-}$ (**Fig. S5**), consistent with their
being *bona fide* deletions arising from concurrent DSBs. The one deletion obtained from $Spo11^{-/-}$ was
between the two hotspots, with both breakpoints distant from the hotspot centers, which was observed
only once in $Atm^{-/-}$.
165
166 **Deletions occur primarily within one chromosome but can join homologous chromosomes**
167
The deletions we observe predict double cutting at hotspot pairs within a single chromosome, however,
in principle they could also arise from staggered DSBs on homologous chromosomes. To investigate
whether deletions involve one or both homologs, we tested the Chr1 hotspot pair in B6 × DBA/2J
(DBA) F1 hybrid males, which have a high polymorphism density across this region (**Fig. 3A**).
Primers were shifted to amplify a larger fragment to aid in identifying the parental chromosome(s)
involved in the deletion.
We recovered 37 deletions from $Atm^{-/-}$ testis DNA for a frequency of 4.20×10^{-6} per haploid
genome, but none in the presence of ATM ($<0.16 \times 10^{-6}$) (**Table S1**). The distributions of deletion
sizes (median 2046 bp) and breakpoints were broadly similar to those from B6 inbred mice (**Fig. 3A;**
S6A). When we analyzed polymorphisms, 30 of the 34 informative deletions (88%) appeared to have
originated within one chromosome: 10 on B6 and 20 on DBA (**Fig. 3A**). The higher frequency of
deletions on the DBA chromosome might be associated with a polymorphism in the PRDM9 site at the
left hotspot (**Fig. 3A,B**), which could positively affect DSB formation. The preponderance of deletions

within one chromosome indicates double cutting on a single chromosome (same chromatid or sister
chromatids), and supports a role for ATM in suppressing closely-spaced DSBs, like Tel1 in yeast[19].
Strikingly, however, we also captured four deletions that contained sequences from both
homologs (**Fig. 3A**). In one, the left hotspot from DBA was joined to the right hotspot from B6 (DBA-
Δ-B6). In the other three more complex events, a haplotype switch occurred adjacent to one of the
deletion breakpoints (DBA-B6-Δ-B6; DBA-B6-Δ-DBA, and B6-DBA-Δ-?, where the question mark
indicates that the genotype was uninformative at the right end). In these haplotype switches, 6–8
polymorphisms were incorporated, encompassing up to 500 bp (**Fig. 3A**).
The recovery of deletions involving interhomolog interactions shows that DSBs can occur at
nearby locations on both homologs in the same cell, at least in the absence of ATM, and, moreover,
that the DSB ends generated on the two homologs are capable of joining. Thus, although homolog
synapsis is greatly impaired in the absence of ATM[35,36], homolog interactions still occur. Cutting of
both homologs at the same hotspot has been inferred from patterns of strand exchange in yeast
tetrads[38-40].
Further, the three complex interhomolog events suggest that three or more DSBs had occurred
on the two homologs. One possible mechanism for the haplotype switch at a deletion breakpoint is
offset DSB formation on both homologs within the same hotspot, with attempted gap repair that is
aborted by the DSB in the template, followed by joining to a DSB at the other hotspot (**Fig. S6B**).
Alternatively, double cutting within one hotspot on one homolog might release a DNA fragment that
then inserts at the DSB in the other homolog (**Fig. S6C**). Irrespective of the precise mechanism,
however, these results indicate that loss of ATM can cause profound damage to multiple chromatids,
leading to deleterious types of repair.
203
204 **Sites of recurrent deletion breakpoints in the absence of ATM**
205
Most deletions between hotspots from $Atm^{-/-}$ mice were unique: 87% (161/185) of events on Chr1
were observed only once, as were 97% (56/58) for Chr19 (**Table S2**). Microhomologies were present
at the breakpoints in 70% of all deletions but were generally short, with a median of 1 bp for both the
Chr1 and Chr19 deletions (**Fig. S7A**). Thus, substantial microhomology is not required for joining.
However, the 24 recurring deletions on Chr1 had longer microhomologies with a median 6 bp (**Fig.
S7B**). Twelve of the recurring deletions arose within a 22/23-bp imperfect sequence repeat shared
between the two hotspots, and eight of these involved an 8-bp microhomology within the repeat (**Fig.
3C, S7B**). Thus, the availability and distribution of short sequence repeats likely influences deletion
outcomes.
The presence or absence of microhomologies within end-joining products may be indicative of
the end joining pathway used and/or processing at DNA ends. For example, SPO11-bound DSB ends
normally require endonucleolytic processing by MRE11 to release SPO11 oligos[4,41,42]. This may
disfavor canonical nonhomologous end-joining and promote alternative pathways that utilize
microhomology[43]. However, as some DSBs remain unresected in the absence of ATM/Tel1[30-34],
deletions may also arise from joining at or very close to SPO11 cleavage positions, which would
involve alternative SPO11 removal mechanisms, such as TDP2[30,44,45]. Our findings are thus consistent
with multiple end-joining mechanisms being employed to repair adjacent DSBs in the absence of
ATM.
224
225 **Insertion of hotspot-derived fragments**
226

Unexpectedly, we observed insertions in 11% of deletions from $Atm^{-/-}$ mice (27/243; two deletions
with two inserts, for 29 total inserts; **Table S1**). All but one of the inserted DNA fragments mapped to
chromosome positions showing SPO11 activity, either the assayed hotspot pairs themselves or other
hotspots. The insert that mapped to a sequence with no SPO11 oligos contained a CT-rich
microsatellite repeat and shared significant microhomology with both deletion breakpoints (12 and 9
bp) (**Table S3**), suggesting a possible DNA synthesis-based insertion mechanism.
Ectopic insertions from other hotspots. Fifteen inserts of 15–349 bp mapped to other
chromosome positions at hotspots or SPO11-oligo clusters that were below the hotspot-calling
threshold (**Fig. 4A-C; S8, S9A; Table S3**). Microhomologies at both sides of inserts were often short
(0 to 2 bp), similar to deletions without insertions (**Fig. 4D; Table S3**).
Most of these inserts could be uniquely mapped, however, three were derived from the non-
pseudoautosomal region (non-PAR) of the Y chromosome. The mouse non-PAR Y is highly
ampliconic[46], so these inserts cannot be assigned to a single location. Hotspots have been called at
some of these locations based on the assignment of unique SPO11 oligos, although most SPO11 oligos
map to multiple locations that are highly similar (**Fig. 4B,C; S9A; S10; Table S3**). Among all
chromosomes, the non-PAR Y experiences the largest increase in SPO11-oligo density in the absence
of ATM (~30 fold)[9].
Inserts mapped at or close to the hotspot/cluster centers, such that nearly half had an end within
50 bp (**Fig. 4B**). We envision that these inserts derive from DNA fragments released by SPO11 double
cutting. It seems remarkable that inserts can sometimes originate from SPO11-oligo clusters, i.e.,
genomic sites that are much less frequently targeted by SPO11 than called hotspots, as such weak sites
would be unlikely to experience two DSBs in the same cell if each DSB occurred independently at a
population-average frequency. However, these findings are readily explained by the recent observation
of concerted SPO11 cleavage within hotspots (especially in $Atm^{-/-}$)[47]. The ectopic insertion of SPO11-
generated fragments into other SPO11-generated DSB locations represents a novel pathway for
meiosis-specific, nonreciprocal genome modification by translocation of sequences.
Inverted insertions derived locally from the hotspot pair. Twelve deletions had insertions of
16 to 709 bp that mapped to one of the hotspots but were reinserted in opposite orientation (inverted
insertions; **Fig. 4E; S9A,B**), including one with an inverted insertion from each hotspot in the pair.
Reconstructing the insertion breakpoints illuminated potential inversion mechanisms. Specifically, in
each case, one insertion breakpoint shared an inverted microhomology with one deletion breakpoint
(median 5 bp) (**Fig. 4F,G**). The other insertion breakpoint, which was at or near this deletion
breakpoint, had no or little microhomology with the other deletion breakpoint (median 0 bp). Inserts
from the left and right hotspots mirrored each other, i.e., those from the left hotspot had the longer
microhomology at the right breakpoint, whereas those from the right hotspot had the longer
microhomology at the left breakpoint (**Fig. 4E,G; S9B; Table S4**). Insert sequences clustered around
hotspot centers (**Fig. 4H**), indicating that they originated from DSB formation.
We propose two models to explain these events (**Fig. 4F**). In one, DSBs occur at or near the
two inverted microhomologies, creating a small double-cut fragment within one hotspot, and a third
DSB in the other hotspot generates a larger fragment destined for deletion. One end of the small
fragment remains unprocessed, preventing normal recombination at this hotspot, while both ends with
the inverted microhomology are at least partially resected, allowing strand annealing and fragment
inversion. The unprocessed end of the now inverted fragment joins to the third DSB at the other
hotspot. This is an attractive model because it incorporates defects seen in $Atm^{-/-}$ mice, i.e., impaired
DSB processing[30,31], as well as invoking closely spaced DSBs. In the second model, a DSB occurs next
to just one of the inverted microhomologies and resection exposes the second microhomology. A

foldback structure formed by annealing the inverted microhomologies is further processed by a
structure-specific nuclease, generating a 5′ overhang that is filled in prior to joining the DSB end at the
other hotspot. Foldback structures may be common in $Atm^{-/-}$, because the excess of DSBs may exhaust
the pool of RPA[48], which normally prevents formation of secondary structures[49].
Interestingly, four inverted insertions from the right hotspot at the Chr1 hotspot pair involved
the previously identified recurrent deletion site (**Fig. 3B; S9B**). In this case, 13 bp overlapping the
recurrent deletion shares inverted microhomology with the most prominent DSB peak. Thus, this
genomic location often engages in inappropriate repair (9% of all Chr1 deletions), demonstrating that
DSB formation at or nearby such sequences may cause (mis)repair products to arise.
282
283 **Microdeletions within a single hotspot**
284
If more than one DSB can form inside the same hotspot, and not just at adjacent hotspots, we predicted
that small deletions might also arise within single hotspots. The presumed origin of insertions from
hotspot fragments implies that closely spaced DSBs can indeed occur within individual hotspots. We
applied deep sequencing to test for such microdeletions. We selected a hotspot on Chr1 ranked among
the five strongest in both $Atm^{-/-}$ and wild type. The SPO11-oligo profile for this hotspot displays a
major central peak, a secondary peak, and several minor peaks, with SPO11 oligos spreading up to
~500 bp on both sides of the central peak, especially to the right (**Fig. 5A**). We designed four
overlapping amplicons, ~580 bp each, to scan ~900 bp of the hotspot. This strategy would recover
deletions around the major peak as well as those with one or both breakpoints away from the major
peak, as often seen for inserts (**Fig. 4B**). Multiple, independent PCRs were carried out, each with input
testis DNA equivalent to ~16,000 haploid genomes, for a total of ~10 million haploid genomes per
amplicon. From pools of amplified DNA we sequenced ~3.5–5 million molecules per amplicon.
We identified reads containing microdeletions ranging from 11 to 229 bp (**Fig. 5A**). (Reads
with ≤10-bp deletions were not considered; **Fig. S11A**). In $Atm^{-/-}$, 216 reads with microdeletions were
detected from the four amplicons, for a total frequency of 1.58×10^{-5}, significantly greater than in
$Spo11^{-/-}$ (0.19×10^{-5}), wild type (0.54×10^{-5}), and $Atm^{+/-}$ (0.23×10^{-5}) (p<0.0001, Chi square test)
(**Table S4**). For some amplicons, reads from wild type and $Atm^{+/-}$ were also more frequent than in
$Spo11^{-/-}$ (**Table S4**). If we disregard duplicate reads for a specific microdeletion from a given
amplicon, we conservatively estimate that we obtained 149 independent microdeletions in total from
$Atm^{-/-}$ (**Fig. 5A**). This is likely an underestimate since some microdeletions were found in multiple
amplicons (**Fig. S11B**).
Microdeletion breakpoints frequently clustered at SPO11-oligo sites in $Atm^{-/-}$ (**Fig. 5A**).
Prominent clusters were observed at the major SPO11-oligo peak (amplicons 1 to 3) and at the
secondary SPO11-oligo peak (amplicons 1 to 4), indicating that microdeletions arose from SPO11-
generated DSBs. This was even more evident when we pooled the 82 independent microdeletions from
amplicons 1, 2, and 3 for which both breakpoints mapped within the 304-bp overlapping region (i.e.,
between the forward primer of amplicon 3 and the reverse primer of amplicon 1) (**Fig. 5B**). It was
visually striking that many microdeletions had both breakpoints within these two SPO11-oligo peaks
(17/82; 21%, 3 of which were recurring; **Fig. S11B**), implying that they originated from double
cutting. In the smoothed breakpoint profiles, the most prominent left breakpoint peak matched the
major SPO11-oligo peak, while the most prominent right breakpoint peak matched the secondary
SPO11-oligo peak (**Fig. 5C**). These breakpoint peaks became even more prominent when all 116 reads
in this overlap region were considered, something not seen with random sampling of reads (**Fig.
S11C**). Thus, at least some of the duplicated reads likely represent independent microdeletions.

Microdeletions in the overlap region displayed a distinct length distribution (**Fig. 5D**), with two prominent microdeletion classes of ~30 bp and ~60 bp. The ~60-bp microdeletions predominantly mapped between the major SPO11-oligo peak and the secondary SPO11-oligo peak (71%, 17/24 from 49–70 bp; **Fig. 5E**), which are separated by 55 bp (**Fig. 5C**); reads within this class were often duplicated (**Fig. S12A**). A subset of the ~30-bp microdeletions (22%, 4/18 from 27–42 bp; **Fig. 5E**) mapped between a minor SPO11-oligo peak and the secondary SPO11-oligo peak 34 bp away (**Fig. 5C**). These results imply that hotspot substructure predicts double cutting and, hence, microdeletion locations. Conversely, double cutting will affect SPO11-oligo distribution within a hotspot when SPO11 oligos predominantly arise from double cutting.

In addition to microdeletions between two SPO11-oligo peaks, ~30-bp (56%, 10/18) and ~60-bp (21%, 5/24) microdeletions often had only one breakpoint at a SPO11-oligo peak (**Fig. 5E**). Assuming they also arose from two SPO11 breaks, the distinct size classes may reflect a more general underlying mechanism for double cutting, such that the length of double-cut fragments may be constrained. In that case, it remains possible that some of the ~60-bp microdeletions may have arisen from two double cuts (triple cutting), releasing two ~30-bp fragments, rather than a single double cut, although this may be less likely given that very few of the ~30-bp microdeletions have left breakpoints within the major SPO11-oligo peak like the ~60-bp microdeletions.

Microdeletions, especially the ~60-bp class, frequently removed the PRDM9 binding site, which is located between the major and secondary SPO11-oligo peaks (**Fig. 5E**). PRDM9 binding could potentially affect the spacing of double cuts. However, as similar microdeletion lengths are observed outside the PRDM9 motif, another or additional factors must exist that lead to this size class.

Interestingly, the major SPO11-oligo peak and the adjacent minor peak are separated by 21 bp (**Fig. 5C**), but the few small microdeletions (11–24 bp) did not map between these peaks (**Fig. 5D, E**), implying that two SPO11-induced DSBs may be constrained to longer distances. Thus, it is possible that they originated from single DSBs or even from double cuts with partial gap filling.

A less prominent microdeletion class of ~140 bp was also observed (129–153 bp, **Fig. 5D**). In several of these longer microdeletions, the left breakpoint clustered at the major SPO11-oligo peak, whereas the right breakpoint was located at sites in which fewer SPO11 oligos mapped (46%, 6/13; **Fig. 5E**). These longer microdeletions may also arise from double cutting. Long SPO11 oligos are observed in $Atm^{-/-}$ testes[4], but are underrepresented in SPO11-oligo maps, given the size selection during library preparation[9,50]. Even longer microdeletions were also observed that peaked at ~180 bp and ~230 bp (**Fig. 5D**), especially when considering all amplicons where microdeletions were less sized constrained (**Fig. S12B,C**). Some of these long microdeletions may have arisen from multiple double-cut fragments that generated long gaps. However, it is notable that the median insert size from other hotspots in the paired hotspot deletions was quite long (~100 bp), with one insert extending 349 bp, suggesting that long double-cut fragments can be generated.

In $Atm^{+/+}$ and $Atm^{+/-}$, microdeletions did not display obvious spatial patterns (**Fig. 5A,E**). Because the frequency of microdeletions in the presence of ATM is only slightly higher in some amplicons than in $Spo11^{-/-}$, we consider that a portion of these events arose from PCR errors or other genomic damage, as in $Spo11^{-/-}$. Strikingly, however, a ~30-bp (25–43-bp) microdeletion class was also apparent, but not in the $Spo11^{-/-}$ control, although the prominent ~60-bp microdeletion class found in $Atm^{-/-}$ was not observed (**Fig. 5D, S12B**). Microdeletions from the ~30-bp class did not map within SPO11-oligo peaks, although some breakpoints clustered (**Fig. 5E**). Possibly, in the presence of ATM, microdeletions are more likely to occur at DSBs formed outside of SPO11-oligo peaks. In both the presence and absence of ATM, microhomology at breakpoints was generally short (median 1 bp) (**Fig. S12D**), similar to deletions between hotspot pairs.

Our assumption that SPO11 double cutting occurs within individual hotspots, preceding
microdeletion formation, is consistent with recent findings from *S. cerevisiae* in which double-cut
fragments were observed[47,51]. Their minimum size was determined to be ~30 bp but longer fragments
in increments of ~10 bp were abundant as well. The authors suggest models in which Spo11 is
constrained to make the simultaneous cleavages. The recent finding of condensates of the yeast DSB
repair machinery suggests a mechanism whereby centers of DSB activity form which could provide
such a platform[52]. Our single hotspot data fit well with both of these concepts. Further, Johnson et al.
computationally identified potential SPO11 double-cut fragments within mouse hotspots in SPO11-
oligo libraries[9]. Assuming a similar lower size limit in mouse as in yeast, double-cut fragments are
predicted to peak at ~32 bp and extend to ≤60 bp in $Atm^{-/-}$, near the size limit of the SPO11-oligo
library preparation[9]. Our analysis of microdeletions is in line with this finding, given a prominent
microdeletion class of ~30 bp, but also suggests that SPO11-oligo double-cut fragments extend much
longer, i.e., from ≥60 bp to at least ~230 bp, congruent with the observation of long SPO11 oligos in
$Atm^{-/-}$[4].

379
380 **Conclusions**
381
Our study uncovers *de novo* deletions in the mouse germline arising at DNA gaps generated by SPO11
double cutting and repaired by nonhomologous end-joining Although pronounced in the absence of
ATM, deletions and microdeletions also occur at low frequency in the presence of ATM. Deletions in
both our hotspot pairs remove exons, demonstrating that SPO11-mediated deletions can disrupt gene
function and/or produce new alleles. Importantly, short direct and inverted repeats at DSB hotspots
help drive these events. The occurrence of deletions involving two chromosomes also predicts gross
rearrangements. Thus, these deletions reveal a previously hidden potential in meiosis for mutagenesis
and the disruption of genome integrity. Furthermore, double-cut fragments appear to be released from
hotspots and inserted at other genomic sites undergoing DSBs, also with mutagenic consequences.
Because DNA gaps are thought to initiate a subset of meiotic recombination events as part of the
normal meiotic program[47], this raises the question as to how re-insertion of released double-cut
fragments is usually suppressed.
Microdeletions resulted in PRDM9 motif deletion in 38% of events in the absence of ATM and
11% in the presence of ATM. By eliminating PRDM9 motifs, such deletions shape the recombination
landscape, thus influencing genome evolution[53,54] and potentially contributing to speciation[55,56].
Because PRDM9 drives recombination away from promoters[57], deletions may be better tolerated in
PRDM9-dependent meiotic recombination, including in humans, and in turn have a greater potential
impact on human genome evolution.

400
401
402

Acknowledgments
We thank Matthew Neale and Franz Klein for sharing unpublished results; members of the Jasin and
Keeney labs for assistance and discussions, in particular Shintaro Yamada; and MSK core facilities, in
particular the Integrated Genomics Operation (Cassidy Cobbs, Neeman Mohibullah, Agnès Viale) and
the Bioinformatics Core (Nicholas Socci), which are supported by NIH grant P30 CA008748. This
work was supported by March of Dimes grant 1-FY17-799 (A.L.), American Cancer Society
fellowship PF-12-157-01-DMC (J.L.), NIH grant R35 GM118092 (S.K.), and NIH grant R35
GM118175 (M.J.).

Author contributions

A.L., S.K., and M.J. conceived the research and designed the experiments with contributions from J.L. A.L. performed all of the experiments. A.L. and M.J. wrote the manuscript with input from S.K. and J.L.

Ethics declarations

Competing interests

The authors declare no competing interests.

References

de Massy, B. Initiation of meiotic recombination: how and where? Conservation and specificities among eukaryotes. *Annu Rev Genet* **47**, 563-599, doi:10.1146/annurev-genet-110711-155423 (2013).

Hunter, N. Meiotic Recombination: The Essence of Heredity. *Cold Spring Harb Perspect Biol* **7**, doi:10.1101/cshperspect.a016618 (2015).

Keeney, S., Lange, J. & Mohibullah, N. Self-organization of meiotic recombination initiation: general principles and molecular pathways. *Annu Rev Genet* **48**, 187-214, doi:10.1146/annurev-genet-120213-092304 (2014).

Lange, J. *et al.* ATM controls meiotic double-strand-break formation. *Nature* **479**, 237-240, doi:10.1038/nature10508 (2011).

Lukaszewicz, A., Lange, J., Keeney, S. & Jasin, M. Control of meiotic double-strand-break formation by ATM: local and global views. *Cell Cycle* **17**, 1155-1172, doi:10.1080/15384101.2018.1464847 (2018).

Keeney, S. Mechanism and control of meiotic recombination initiation. *Curr Top Dev Biol* **52**, 1-53 (2001).

Petes, T. D. Meiotic recombination hot spots and cold spots. *Nat Rev Genet* **2**, 360-369, doi:10.1038/35072078 (2001).

Baudat, F., Imai, Y. & de Massy, B. Meiotic recombination in mammals: localization and regulation. *Nat Rev Genet* **14**, 794-806, doi:10.1038/nrg3573 (2013).

Lange, J. *et al.* The Landscape of Mouse Meiotic Double-Strand Break Formation, Processing, and Repair. *Cell* **167**, 695-708 e616, doi:10.1016/j.cell.2016.09.035 (2016).

Cole, F. *et al.* Homeostatic control of recombination is implemented progressively in mouse meiosis. *Nat Cell Biol* **14**, 424-430, doi:10.1038/ncb2451 (2012).

Tock, A. J. & Henderson, I. R. Hotspots for Initiation of Meiotic Recombination. *Front Genet* **9**, 521, doi:10.3389/fgene.2018.00521 (2018).

Pan, J. *et al.* A hierarchical combination of factors shapes the genome-wide topography of yeast meiotic recombination initiation. *Cell* **144**, 719-731, doi:10.1016/j.cell.2011.02.009 (2011).

13 Yamada, S. *et al.* Genomic and chromatin features shaping meiotic double-strand break
formation and repair in mice. *Cell Cycle* **16**, 1870-1884, doi:10.1080/15384101.2017.1361065
(2017).
14 Cooper, T. J., Garcia, V. & Neale, M. J. Meiotic DSB patterning: A multifaceted process. *Cell*
Cycle **15**, 13-21, doi:10.1080/15384101.2015.1093709 (2016).
15 Shiloh, Y. The ATM-mediated DNA-damage response: taking shape. *Trends Biochem Sci* **31**,
402-410, doi:10.1016/j.tibs.2006.05.004 (2006).
16 Tian, M. & Loidl, J. A chromatin-associated protein required for inducing and limiting meiotic
DNA double-strand break formation. *Nucleic Acids Res* **46**, 11822-11834,
doi:10.1093/nar/gky968 (2018).
17 Checchi, P. M., Lawrence, K. S., Van, M. V., Larson, B. J. & Engebrecht, J. Pseudosynapsis
and decreased stringency of meiotic repair pathway choice on the hemizygous sex chromosome
of Caenorhabditis elegans males. *Genetics* **197**, 543-560, doi:10.1534/genetics.114.164152
(2014).
18 Mohibullah, N. & Keeney, S. Numerical and spatial patterning of yeast meiotic DNA breaks by
Tel1. *Genome Res* **27**, 278-288, doi:10.1101/gr.213587.116 (2017).
19 Garcia, V., Gray, S., Allison, R. M., Cooper, T. J. & Neale, M. J. Tel1(ATM)-mediated
interference suppresses clustered meiotic double-strand-break formation. *Nature* **520**, 114-118,
doi:10.1038/nature13993 (2015).
20 Chang, H. H. Y., Pannunzio, N. R., Adachi, N. & Lieber, M. R. Non-homologous DNA end
joining and alternative pathways to double-strand break repair. *Nat Rev Mol Cell Biol* **18**, 495-
506, doi:10.1038/nrm.2017.48 (2017).
21 Kim, S., Peterson, S. E., Jasin, M. & Keeney, S. Mechanisms of germ line genome instability.
Semin Cell Dev Biol **54**, 177-187, doi:10.1016/j.semcdb.2016.02.019 (2016).
22 Ahmed, E. A., Philippens, M. E., Kal, H. B., de Rooij, D. G. & de Boer, P. Genetic probing of
homologous recombination and non-homologous end joining during meiotic prophase in
irradiated mouse spermatocytes. *Mutat Res* **688**, 12-18, doi:10.1016/j.mrfmmm.2010.02.004
(2010).
23 Ahmed, E. A., Rosemann, M. & Scherthan, H. NHEJ Contributes to the Fast Repair of
Radiation-induced DNA Double-strand Breaks at Late Prophase I Telomeres. *Health Phys* **115**,
102-107, doi:10.1097/HP.0000000000000852 (2018).
24 Enguita-Marruedo, A. *et al.* Transition from a meiotic to a somatic-like DNA damage response
during the pachytene stage in mouse meiosis. *PLoS Genet* **15**, e1007439,
doi:10.1371/journal.pgen.1007439 (2019).
25 Girard, C., Roelens, B., Zawadzki, K. A. & Villeneuve, A. M. Interdependent and separable
functions of Caenorhabditis elegans MRN-C complex members couple formation and repair of
meiotic DSBs. *Proc Natl Acad Sci U S A* **115**, E4443-E4452, doi:10.1073/pnas.1719029115
(2018).
26 Lemmens, B. B., Johnson, N. M. & Tijsterman, M. COM-1 promotes homologous
recombination during Caenorhabditis elegans meiosis by antagonizing Ku-mediated non-
homologous end joining. *PLoS Genet* **9**, e1003276, doi:10.1371/journal.pgen.1003276 (2013).
27 Yin, Y. & Smolikove, S. Impaired resection of meiotic double-strand breaks channels repair to
nonhomologous end joining in Caenorhabditis elegans. *Mol Cell Biol* **33**, 2732-2747,
doi:10.1128/MCB.00055-13 (2013).
28 Macaisne, N., Kessler, Z. & Yanowitz, J. L. Meiotic Double-Strand Break Proteins Influence
Repair Pathway Utilization. *Genetics* **210**, 843-856, doi:10.1534/genetics.118.301402 (2018).

Martin, J. S., Winkelmann, N., Petalcorin, M. I., McIlwraith, M. J. & Boulton, S. J. RAD-51-dependent and -independent roles of a Caenorhabditis elegans BRCA2-related protein during DNA double-strand break repair. *Mol Cell Biol* **25**, 3127-3139, doi:10.1128/MCB.25.8.3127-3139.2005 (2005).

Paiano, J. *et al.* ATM and PRDM9 regulate SPO11-bound recombination intermediates during meiosis. *Nat Commun* **11**, 857, doi:10.1038/s41467-020-14654-w (2020).

Yamada, S. *et al.* Molecular structures and mechanisms of DNA break processing in mouse meiosis. *Genes Dev*, doi:10.1101/gad.336032.119 (2020).

Mimitou, E. P., Yamada, S. & Keeney, S. A global view of meiotic double-strand break end resection. *Science* **355**, 40-45, doi:10.1126/science.aak9704 (2017).

Cannavo, E. *et al.* Regulatory control of DNA end resection by Sae2 phosphorylation. *Nat Commun* **9**, 4016, doi:10.1038/s41467-018-06417-5 (2018).

Joshi, N., Brown, M. S., Bishop, D. K. & Borner, G. V. Gradual implementation of the meiotic recombination program via checkpoint pathways controlled by global DSB levels. *Mol Cell* **57**, 797-811, doi:10.1016/j.molcel.2014.12.027 (2015).

Barlow, C. *et al.* Atm deficiency results in severe meiotic disruption as early as leptonema of prophase I. *Development* **125**, 4007-4017 (1998).

Xu, Y. *et al.* Targeted disruption of ATM leads to growth retardation, chromosomal fragmentation during meiosis, immune defects, and thymic lymphoma. *Genes Dev* **10**, 2411-2422, doi:10.1101/gad.10.19.2411 (1996).

Cole, F., Keeney, S. & Jasin, M. Comprehensive, fine-scale dissection of homologous recombination outcomes at a hot spot in mouse meiosis. *Mol Cell* **39**, 700-710, doi:10.1016/j.molcel.2010.08.017 (2010).

Marsolier-Kergoat, M. C., Khan, M. M., Schott, J., Zhu, X. & Llorente, B. Mechanistic View and Genetic Control of DNA Recombination during Meiosis. *Mol Cell* **70**, 9-20 e26, doi:10.1016/j.molcel.2018.02.032 (2018).

Anderson, C. M., Oke, A., Yam, P., Zhuge, T. & Fung, J. C. Reduced Crossover Interference and Increased ZMM-Independent Recombination in the Absence of Tel1/ATM. *PLoS Genet* **11**, e1005478, doi:10.1371/journal.pgen.1005478 (2015).

Zhang, L., Kim, K. P., Kleckner, N. E. & Storlazzi, A. Meiotic double-strand breaks occur once per pair of (sister) chromatids and, via Mec1/ATR and Tel1/ATM, once per quartet of chromatids. *Proc Natl Acad Sci U S A* **108**, 20036-20041, doi:10.1073/pnas.1117937108 (2011).

Keeney, S., Giroux, C. N. & Kleckner, N. Meiosis-specific DNA double-strand breaks are catalyzed by Spo11, a member of a widely conserved protein family. *Cell* **88**, 375-384, doi:10.1016/s0092-8674(00)81876-0 (1997).

Neale, M. J., Pan, J. & Keeney, S. Endonucleolytic processing of covalent protein-linked DNA double-strand breaks. *Nature* **436**, 1053-1057, doi:10.1038/nature03872 (2005).

Sfeir, A. & Symington, L. S. Microhomology-Mediated End Joining: A Back-up Survival Mechanism or Dedicated Pathway? *Trends in biochemical sciences* **40**, 701-714, doi:10.1016/j.tibs.2015.08.006 (2015).

Alvarez-Quilon, A. *et al.* ATM specifically mediates repair of double-strand breaks with blocked DNA ends. *Nat Commun* **5**, 3347, doi:10.1038/ncomms4347 (2014).

Johnson, D., Allison, R. M., Cannavo, E., Cejka, P. & Neale, M. J. Removal of Spo11 from meiotic DNA breaks in vitro but not in vivo by Tyrosyl DNA Phosphodiesterase 2. *bioRxiv*, doi:0.1101/527333 (2019).

46   Soh, Y. Q. *et al.* Sequencing the mouse Y chromosome reveals convergent gene acquisition and
amplification on both sex chromosomes. *Cell* **159**, 800-813, doi:10.1016/j.cell.2014.09.052
(2014).
47   Johnson, D. *et al.* Concerted cutting by Spo11 illuminates DNA break mechanisms and initiates
gap repair during meiosis. *bioRxiv*, doi:10.1101/2019.12.18.881268 (2019).
48   Toledo, L. I. *et al.* ATR prohibits replication catastrophe by preventing global exhaustion of
RPA. *Cell* **155**, 1088-1103, doi:10.1016/j.cell.2013.10.043 (2013).
49   Deng, S. K., Chen, H. & Symington, L. S. Replication protein A prevents promiscuous
annealing between short sequence homologies: Implications for genome integrity. *Bioessays*
37, 305-313, doi:10.1002/bies.201400161 (2015).
50   Thacker, D., Mohibullah, N., Zhu, X. & Keeney, S. Homologue engagement controls meiotic
DNA break number and distribution. *Nature* **510**, 241-246, doi:10.1038/nature13120 (2014).
51   Prieler, S. *et al.* Spo11 generates chromosomal gaps through concerted cuts and targets sites of
topological stress. (Submitted).
52   Claeys Bouuaert, C., Pu, S., Wang, J., Patel, D. J. & Keeney, S. DNA-dependent
macromolecular condensation drives self-assembly of the meiotic DNA break machinery.
bioRxiv, doi:10.1101/2020.02.21.960245 (2020).
53   Cole, F. *et al.* Mouse tetrad analysis provides insights into recombination mechanisms and
hotspot evolutionary dynamics. *Nat Genet* **46**, 1072-1080, doi:10.1038/ng.3068 (2014).
54   Baker, C. L. *et al.* PRDM9 drives evolutionary erosion of hotspots in Mus musculus through
haplotype-specific initiation of meiotic recombination. *PLoS Genet* **11**, e1004916,
doi:10.1371/journal.pgen.1004916 (2015).
55   Davies, B. *et al.* Re-engineering the zinc fingers of PRDM9 reverses hybrid sterility in mice.
Nature **530**, 171-176, doi:10.1038/nature16931 (2016).
56   Smagulova, F., Brick, K., Pu, Y., Camerini-Otero, R. D. & Petukhova, G. V. The evolutionary
turnover of recombination hot spots contributes to speciation in mice. *Genes Dev* **30**, 266-280,
doi:10.1101/gad.270009.115 (2016).
57   Brick, K., Smagulova, F., Khil, P., Camerini-Otero, R. D. & Petukhova, G. V. Genetic
recombination is directed away from functional genomic elements in mice. *Nature* **485**, 642-
645, doi:10.1038/nature11089 (2012).

578
579
580

Methods

Mouse husbandry

All experiments were conducted according to US Office of Laboratory Animal Welfare regulatory
standards and were approved by the Memorial Sloan Kettering Cancer Center Institutional Animal
Care and Use Committee. Mice were purchased from The Jackson Laboratory where indicated. $Atm^{+/-}$
males on the C57BL/6J (B6) background (B6.129S6-Atm^{tm1Awb}/J; stock #008536) were crossed with
B6 females (stock #000664) to establish a breeding colony and to obtain experimental animals. The
Atm mutation was previously described[58]. $Atm^{+/-}$ B6 mice were crossed with $Atm^{+/-}$ DBA/2J (DBA)
mice to get F1 hybrid males. $Atm^{+/-}$ mice on the DBA background were obtained by six backcrosses of
$Atm^{+/-}$ mice that were partially on an A/J background with DBA inbred mice (stock #000671). Animals
were fed regular rodent chow with continuous access to food and water.

Isolation of testis genomic DNA

DNA was isolated from testicular single-cell suspensions to enrich for germ cells. Testis dissociation was based on protocols previously described[53,59] with some modifications. Testes from juvenile 15-day-old and adult 2- to 5-month-old mice were decapsulated and seminiferous tubules incubated at 33 °C in 10 ml Gey's Balanced Salt Solution (GBSS) (Sigma) with 0.5 mg/ml collagenase (Sigma) and 2 µg/ml DNase I (Roche) for 15–20 min, shaking at 500 rpm. After 10 min and at the end of the incubation the tubules were gently pipetted up and down using a transfer pipette. The tubules were washed twice with 10 ml GBSS and digested with 0.5 mg/ml trypsin (Sigma) in 10 ml GBSS, supplemented with 2 µg/ml DNase I, for 25–40 min at 33 °C, shaking at 500 rpm. Pipetting up and down was performed at the start, midpoint and end of the incubation. Trypsin was inhibited by adding 10% fetal calf serum. Cell suspension was obtained by repeated pipetting for 2 min and filtration through a 70-µm cell strainer (ThermoFisher). All of the mutant cell suspension and one-third to one-half of the wild-type cell suspension was transferred to a 15 ml tube and centrifuged for 3 min at 1550 × g. The cell pellet was resuspended by tapping the tube, followed by the addition of 600 µl lysis buffer (200 mM NaCl, 100 mM Tris-Cl pH 7.5, 5 mM EDTA, 0.5% SDS, 0.5 mg/ml Proteinase K). The lysate was transferred to a 1.5 ml Eppendorf tube and incubated overnight at 55 °C. Genomic DNA was isolated using phenol-chloroform extraction and alcohol precipitation, with the extraction step repeated twice. DNA was dissolved in 200–400 µl H_2O, incubated for 1 h at 55 °C and overnight at 4 °C. DNA concentration was quantified by NanoDrop (ThermoFisher). DNA was diluted to ~200 ng/µl and incubated for another 2–3 h at 37 °C to reduce the viscosity of the DNA solution. DNA concentration was measured again three times before freezing. DNA samples were stored at -20 °C.

Detection of deletions at hotspot pairs by nested PCR

Deletions were detected by nested PCR using primers flanking the two hotspots of a pair (**Fig. 1B; S2A**). Prior to PCR, genomic DNA was digested with restriction enzymes where indicated to suppress parental DNA amplification (**Fig. S4**). Restriction digestions were performed in a 60 µl volume with 1200 ng of DNA and 4–4.5 µl restriction enzyme(s) (Chr1 hotspot pair: NsiI, NcoI, and BglII or NcoI alone; Chr19 hotspot pair: BamHI or KpnI). Reactions were incubated overnight at 37 °C followed by an additional 3 h of incubation with a 1.5–2 µl of fresh restriction enzyme(s), then placed on ice before PCR. In the majority of PCRs, 2.5 µl of the digestion reaction was used per PCR, equivalent to 50 ng of genomic DNA, with this DNA concentration being optimal for the estimation of deletion frequencies; 1.25 µl and 5 µl, equivalent to 25 ng and 100 ng, respectively, were also tested. In the experiments with no digestion, 50 ng of DNA template was used per PCR in most cases.

Nested PCR was based on the previously described protocol[60] with some adjustments. A first round of PCR was performed using the following primers: Chr1_DH_FW1: 5′-GGTCAGGAGGATGGGGAG-3′ and Chr1_DH_RV1: 5′-CCAAGAAGGTGCTTATCGTCC-3′ for the Chr1 hotspot pair, Chr19_DH_FW2: 5′-TGGAATGTAGCCTCTGGCAG-3′ and Chr19_DH_RV2: 5′-GTGCTGTCAGAAGCCGTGG-3′ for the Chr19 hotspot pair, Chr1_DH_DB_FW1: 5′-CCTGGAGACCAAAGCCAC-3′ and Chr1_DH_RV1: 5′-CCAAGAAGGTGCTTATCGTCC-3′ for the Chr1 hotspot pair in B6/DBA F1 hybrids. A second round of PCR was performed using the following primers: Chr1_DH_FW2: 5′-GCCTTGGCTCAAAGTGTCTG-3′ and Chr1_DH_RV2: 5′-TCCCTTCTGCCATCCTTCTG-3′ for the Chr1 hotspot pair, newChr19_DH_FW2: 5′-

CCCAAAGTTGTCCTTGCTTCC-3′ and newChr19_DH_RV2: 5′-TGGTTTCCAGAGGAGAGGAC-3′ for the Chr19 hotspot pair, Chr1_DH_DB_FW2: 5′-GGCATTGCTTCATTCATGGTCTG-3′ and Chr1_DH_RV2-3: 5′-CATTGTCATCTTGAGCACTCCTC-3′ for the Chr1 hotspot pair in B6/DBA F1 hybrids. The annealing temperature was optimized for each primer pair by gradient PCR with testis genomic DNA of known high quality. DNA samples from different testes were also tested for the appearance of a strong parental PCR band using the primary PCR primers.

For nested PCR, each primary PCR was performed in a 50 μl volume containing: 1x Jeffreys buffer (10x: 450 mM Tris-Cl pH 8.8, 110 mM $(NH_4)_2SO_4$, 45 mM $MgCl_2$, 67 mM 2-mercaptoethanol, 44 μM EDTA, 10 mM each dNTP (Roche), 1.13 mg/ml ultra-pure (non-acetylated) BSA (Ambion), 12.5 mM Tris base), 0.2 μM primers, 0.03 U/μl Taq DNA polymerase (ThermoFisher) and 0.006 U/μl Pfu Turbo DNA polymerase (Agilent). The primary PCR product (0.6 μl) was treated with S1 nuclease (ThermoFisher) at room temperature for 30 min, in a 5 μl volume containing: 0.5x S1 buffer, 0.2 U/μl S1 nuclease and 5 ng/μl sonicated salmon sperm DNA. The reaction was diluted with 45 μl of dilution buffer (10 mM Tris-Cl pH 7.5 and 5 ng/μl sonicated salmon sperm DNA). 4 μl of the diluted S1 reaction was seeded into the secondary PCR, which was performed in a 20 μl volume containing the same components as the primary PCR. Deletion products were identified by electrophoresis on a 1% agarose gel (**Fig. 1C, Fig. S2B**). PCR products smaller than the parental size were excised, purified, and Sanger sequenced using primers from the second round of PCR and in some cases additional primers. Diluted S1 reactions were stored at -20 °C, and used again for secondary PCRs if more PCR product was needed for sequencing.

PCR conditions were tested for each hotspot pair. The best result, that is, the appearance of more intense, smaller PCR products, was obtained when the extension time was shortened (<60 s/kb) in the primary PCR to inhibit amplification of the longer parental PCR product. The extension time in the secondary PCR was slightly shorter.

Primary PCR conditions were denaturation at 96 °C for 2 min, followed by 37 cycles of amplification as follows: denaturation at 96 °C for 20 s, annealing at 59 °C for 30 s, extension at 65 °C for 2 min for the Chr1 and Chr19 hotspot pairs (3.6 kb parental products), and 2 min 30 s for the Chr1 hotspot pair in B6/DBA hybrids (4 kb parental product). Secondary PCR conditions where the same, but with different extension times: 3 min for the Chr1 and Chr19 hotspot pairs (3.4-kb parental products) and 3 min 30 s for the Chr1 hotspot pair in B6/DBA hybrids (3.9-kb parental products).

Sequenced PCR products were analyzed using ApE software (https://jorgensen.biology.utah.edu/wayned/ape/) to identify deletions, microhomologies, and polymorphisms in DNA sequences. Deletion breakpoint profiling was performed using R versions 3.4.4 to 3.6.3 (http://www.r-project.org). Graphing was performed in R and GraphPad Prism version 8.

Detection of microdeletions at a single hotspot by amplicon deep sequencing

Four ~580-bp overlapping amplicons of the hotspot (**Fig. 5A**) were amplified using the following primer pairs: amplicon 1, Chr1_MiSeq_FW1: 5′-TCATGTCCGGTTTGGATGCC-3′ and Chr1_MiSeq_RV1: 5′-CGAATCAAAGACTCCAGAAGAGG-3′; amplicon 2, Chr1_MiSeq_FW2: 5′-AGTAGCAAGAAGGGCTGGAG-3′ and Chr1_MiSeq_RV2: 5′-AGAGACAGATGTGGAGGCTAAC-3′; amplicon 3, Chr1_MiSeq_FW4: 5′-CTGAATCTCTCCTGGTGTGTC-3′ and Chr1_MiSeq_RV4: 5′-AGGACATTGAATGTTTTTATGGTTC-3′; amplicon 4, Chr1_MiSeq_FW5: 5′-AAACAGAAGAAGCAGTAATGCC-3′ and Chr1_MiSeq_RV5: 5′-

GTGTGCAACTGGGTATATTTCAG-3′. For each primer pair the annealing temperature was tested
by gradient PCR with testis genomic DNA of known high quality. To minimize PCR bias and jackpot
amplification of single deletion events, the number of amplification cycles was determined by when a
faint PCR product was first visible, which was 22 cycles for all samples. For each amplicon, PCR
conditions were the same: 96 °C for 2 min followed by 22 cycles at 96 °C for 20 s, 59 °C for 30 s and
65 °C for 35 s. PCR was performed in a 50 μl total containing 1.5 U Taq DNA polymerase
(ThermoFisher) and 0.3 U Pfu Turbo DNA polymerase (Agilent) in the Jeffreys buffer described above
for nested PCR with 50 ng of testis genomic DNA. Testis DNA was derived from 3-5 different mice
for each genotype and amplified independently.
PCRs were carried out in 96-well plates to generate a large number of independently-derived
PCR products for sequencing. For each amplicon, 576 PCRs were performed with a total of 28.8 μg
genomic DNA, equivalent to 9.6 million haploid genomes. PCRs were pooled, and a 1200-μl aliquot
was purified and concentrated on spin columns (Invitrogen PureLink[TM] PCR Purification Kit) as
follows: For each 100 μl PCR aliquot, 400 μl binding buffer from the kit was added, and then the
sample was applied to the column and centrifuged. This step was repeated 6 times in each of two
columns before washing and elution with 37 μl elution buffer. Eluted DNA was combined from the
two columns and a 15-μl aliquot was run on a 2% agarose gel to estimate DNA concentration, which
was ~2 ng/μl. The remaining ~60 μl was submitted to the Integrated Genomics Operation at MSK for
library preparation and deep sequencing. The library was prepared by ligation of indexed adapter
sequences using the KAPA HyperPrep Kit (Roche). Sequencing was performed using Illumina MiSeq
(2 x 300-bp paired-end reads). In a single run, libraries from amplicons 1–4 were mixed with 5% PhiX
DNA control library (Illumina) and sequenced. Because MiSeq can deliver 15–20 million reads per
lane, in our experiment, 3.6–4.8 million reads per amplicon were estimated, which was about 2 times
lower than the 9.6 million input DNA templates. With that calculation, not all deletion events were
recovered, but most recovered deletions should have yielded only a single read.
We performed 4 MiSeq runs with amplicons from different genotypes (**Fig. S11A**). FASTQ
sequencing files were analyzed using CRISPResso2[61,62] by the Bioinformatics Core Facility at MSK.
Reads were aligned to the reference amplicon sequence and 2.9–5.3 million paired-end reads were
mapped per amplicon. Alignments matching the target sequence for the first 20 and last 20 bases and
with one deletion event were filtered. Reads with >1 bp deletion and with no mutations in the sequence
were analyzed in R and ApE. Breakpoint distribution analyses and random sampling (**Fig. 11C**) were
performed in R. Graphs were plotted using R and GraphPad Prism, and the Chi-square test was
performed in GraphPad Prism.

SPO11-oligo maps

SPO11-oligo maps were generated previously (GEO accession number GSE84689)[9]. Mice for the *Atm*[−]
[/−] data set "*Atm* null 1", which was used in all the analyses, were on a mixed B6 and 129/Sv strain
background. Two *Atm*[+/+] data sets are available, "B6", from mice on a pure B6 background, and "*Atm*
wt", from the same colony as *Atm*[−/−] mice. Hotspots selected in this study were called in the "B6" data
set (**Fig. 1B, S1D, S2A**), while analysis of hotspot distributions was done with the "*Atm* wt" data set
(**Fig. 1A, S1A–C**). The "B6" data set was also used to check if SPO11 oligos map to the genomic
locations from which inserts derived (**Fig. S8, S10**).

58   Barlow, C. *et al.* Atm-deficient mice: a paradigm of ataxia telangiectasia. *Cell* **86**, 159-171,
doi:10.1016/s0092-8674(00)80086-0 (1996).

59   Gaysinskaya, V., Soh, I. Y., van der Heijden, G. W. & Bortvin, A. Optimized flow cytometry
isolation of murine spermatocytes. *Cytometry A* **85**, 556-565, doi:10.1002/cyto.a.22463 (2014).
60   Cole, F. & Jasin, M. Isolation of meiotic recombinants from mouse sperm. *Methods Mol Biol*
745, 251-282, doi:10.1007/978-1-61779-129-1_15 (2011).
61   Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing sequence analysis.
Nat Biotechnol **37**, 224-226, doi:10.1038/s41587-019-0032-3 (2019).
62   Pinello, L. *et al.* Analyzing CRISPR genome-editing experiments with CRISPResso. *Nat*
Biotechnol **34**, 695-697, doi:10.1038/nbt.3583 (2016).
63   Schirmer, M. *et al.* Insight into biases and sequencing errors for amplicon sequencing with the
Illumina MiSeq platform. *Nucleic Acids Res* **43**, e37, doi:10.1093/nar/gku1341 (2015).

Figure 1. Meiotic deletions at adjacent DSB hotspots.
(A) Distribution of distances separating adjacent SPO11-oligo hotspots on autosomes. The hotspot calling algorithm[9] merges hotspots separated by edge-to-edge distances of <1 kb. Hotspot pairs separated by >250 kb are not shown. See also **Fig. S1**.

(B) Strategy to detect deletions between adjacent hotspots. Top panel, SPO11-oligo maps for the hotspot pair on Chr1 (RPM, reads per million). Genomic coordinates (MM10 assembly) are shown below the graphs. The estimated fold increase in absolute SPO11-oligo frequency (RPM × 11.3) is indicated for each hotspot. Vertical pink lines indicate hotspot centers. Lower panel, scheme of the PCR assay. Nested PCR across the DSB hotspot pair is carried out with the indicated primer pairs (arrows). Prior to amplification, genomic DNA is optionally digested with restriction enzymes to reduce amplification of parental products. The *Atp1b1* exon located between the two hotspots is removed in deletions.
(C) Detection of deletions by agarose gel electrophoresis. Representative gels for the Chr1 hotspot pair are shown. Each lane is a PCR seeded with the indicated number of haploid genomes.
(D) Deletion frequencies. Three to seven mice of each genotype per hotspot pair were analyzed, listed in **Table S1**.

Figure 2. Deletion breakpoints at the Chr1 hotspot pair.
(A) Deleted DNA sequences at the Chr1 hotspot pair. Deletions (black bars) were combined from adult and juvenile testes (dpp, days post partum).
(B) Arc diagram of deletions in *Atm*⁻/⁻ at the Chr1 hotspot pair. Arcs link the breakpoints of each deletion (134 deletions total). Both deletion breakpoints often mapped close to the hotspot centers. See also **Fig. S3**.
(C) Individual breakpoints cluster around the hotspot centers but are also observed to spread.
(D) Smoothed breakpoint and SPO11-oligo profiles display similar shapes (201-bp Hann filter for smoothing). Offsets between centers of hotspots and breakpoint distributions are indicated.

Figure 3. Meiotic deletions within and between chromosomes.
(**A**) Deletions at the Chr1 hotspot pair in $Atm^{-/-}$ B6 × DBA F1 hybrid males. Nested PCR was performed with the indicated primers (arrows) to identify polymorphisms within the deletion products derived from B6 (red) or DBA (blue). Three deletions were uninformative because the involved chromosome(s) is uncertain.
(**B**) Predicted PRDM9 recognition sequences at the left hotspot in B6 and DBA, aligned to the PRDM9 motif from SPO11-oligo analysis[9].
(**C**) Recurrent deletion within the Chr1 hotspot pair at an imperfect repeat. Twelve simple deletions in $Atm^{-/-}$ map to this repeat at one of the three internal identical repeats of 5, 8, and 6 bp (violet arcs). Four other deletions with inverted insertions also involve this imperfect repeat as well as a 13-bp inverted microhomology at the hotspot center, with the insertions indicated above the deletions (orange arcs). See also **Fig. S7A, S9B**.

Figure 4

Figure 4. Insertions derived from DSB hotspots.
(A–D) Ectopic insertions from other hotspots. **A**, Example of a deletion on Chr1 from $Atm^{-/-}$ containing a 79-bp insertion derived from a hotspot on Chr7. **B**, Inserts from other hotspots map close to their respective hotspot centers. Three inserts map to SPO11-oligo clusters rather than called hotspots (asterisks). Inserts 1 and 3 from the non-PAR ChrY map to multiple hotspots. See also **Fig. S8, S10** and **Table S3**. **C**, Non-PAR ChrY hotspot to which inserts 1 and 2 can be mapped. Some SPO11 oligos can be uniquely mapped to this hotspot, but, as it is one copy of a family of repeats, numerous other SPO11 oligos can be mapped to both this and other copies of the repeat ("Multi"). **D**, Microhomologies at insert breakpoints are often short (median, 1 bp).
(E–H) Inverted insertions derived locally from the hotspot pair. **E**, Deletions with inverted insertions at the Chr19 hotspot pair. Inserts are derived in inverted orientation from the left or the right hotspot. For the Chr1 hotspot pair, see **Fig. S9B**. **F**, Example of an inverted insertion (top) and possible mechanisms (bottom). The right insertion breakpoint shares a 5-bp inverted microhomology with the left deletion breakpoint, while the left insertion breakpoint and the right deletion breakpoint do not share microhomology. Potential models explaining the inversion mechanism involve double cutting at the left hotspot or overhang foldback. **G**, Microhomologies at breakpoints for insertions derived from the Chr1 and Chr19 hotspot pairs. Inserts mapping to the left hotspot in a pair have long microhomologies present at their right breakpoints. Inserts mapping to the right hotspot mirror this pattern. See also **Table S4**. **H**, Inserts from the Chr1 and Chr19 hotspot pairs map close to hotspot centers.

Figure 5. Microdeletions within a single DSB hotspot.
(A) Detection of microdeletions at a strong DSB hotspot on Chr1 using amplicon deep sequencing. The hotspot displays a major SPO11-oligo peak, a secondary peak, and several minor peaks, as shown in the smoothed (21-bp Hann filter) and unsmoothed profiles. Four overlapping amplicons were deep sequenced to identify microdeletions. Arc diagrams of independent microdeletions found in each amplicon for the indicated genotypes are shown. For microdeletion frequencies, see **Table S5**.
(B, C) Microdeletions from $Atm^{-/-}$ in the region overlapping in amplicons 1, 2, and 3. **B**, Arc diagram of independent microdeletions. **C**, Distributions of the left and right breakpoints relative to the SPO11-oligo distribution. Both unsmoothed and smoothed profiles are shown.
(D, E) Microdeletion length distributions in the overlapping region. **D,** Density and scatter plots of microdeletion lengths (>10 bp). **E,** Arc diagrams of microdeletions stratified by length. Microdeletions >160 bp are not shown, as their location may be more restricted by the size of the assessed region. See also **Fig. S12B,C** for larger microdeletion distribution.

A

Dataset	No. hotspots	Hotspot pairs	No. hotspot pairs separated by 1-5 kb	Hotspots in hotspot pairs (%)
Atm$^{+/+}$	13436	*Atm*$^{+/+}$	413	6.1
Atm$^{-/-}$	20417	*Atm*$^{-/-}$	1023	10.0
shared	12172	both shared	304	3.0
new	8245	both new	241	2.4
		new+shared	478	4.7

B

C

Hotspot pairs (1–5 kb)

Atm$^{+/+}$ Shared with *Atm*$^{-/-}$ Median=51

Atm$^{-/-}$ **Both shared**   Median=75

Atm$^{-/-}$ **Both new**   Median=9

Atm$^{-/-}$ **New+shared**   Median=38

D

	Hotspot strength (RPM) [hotspot rank]		Fold increase in *Atm*$^{-/-}$ hotspot strength*
Chr1 pair	*Atm*$^{+/+}$	*Atm*$^{-/-}$	
hotspot left	65 [16%]	53 [16%]	9.2
hotspot right	209 [3%]	93 [6%]	5.0
Chr19 pair			
hotspot left	50 [20%]	41 [21%]	9.3
hotspot right	80 [13%]	69 [11%]	9.7

**Atm$^{-/-}$ RPM divided by *Atm*$^{+/+}$ RPM multiplied by 11.3, the average increase in SPO11 oligos in *Atm*$^{-/-}$.

Figure S1. Higher potential in *Atm*$^{-/-}$ for DSB formation at adjacent hotspots.
Analyses were performed for autosomes.
(A) Hotspot pairs separated by 1–5 kb are observed more frequently in *Atm*$^{-/-}$ than in *Atm*$^{+/+}$. Data are derived from SPO11-oligo maps[9]. "Shared" hotspots are those called in both *Atm*$^{-/-}$ and *Atm*$^{+/+}$. "New" hotspots are those called only in *Atm*$^{-/-}$, although corresponding SPO11-oligo clusters are typically present in the *Atm*$^{+/+}$ map[9]. Hotspot pairs in *Atm*$^{-/-}$ are separated into classes according to whether each hotspot is shared or new.
(B) Distribution of distances separating hotspots in *Atm*$^{-/-}$ hotspot pairs, shown up to 250 kb. The proportion of each hotspot pair class from panel **A** is indicated. The increase in the number of hotspot pairs in *Atm*$^{-/-}$ separated by short distances (**Fig. 1A**) is primarily the result of the emergence of new hotspots close to shared hotspots.
(C) Strength of individual hotspots in hotspot pairs separated by 1–5 kb. Hotspot pair strength, the median of total RPM (reads per million) for the two hotspots in a pair, is indicated. Note that the RPM values do not account for the 11.3-fold increase in SPO11 oligos in *Atm*$^{-/-}$[4]. Outliers with >300 RPM (4 in *Atm*$^{-/-}$; 18 in *Atm*$^{+/+}$) are not shown.
(D) Hotspot strength for the individual hotspots in the Chr1 and Chr19 hotspot pairs. RPM values in *Atm*$^{-/-}$ are multiplied by 11.3, the average increase in whole-testis SPO11-oligo levels in *Atm*$^{-/-}$ mice. Hotspot rank is the rank of the individual hotspot within all hotspots for *Atm*$^{+/+}$ or *Atm*$^{-/-}$.

Figure S2. Deletions at the Chr19 hotspot pair.
(A) SPO11-oligo maps in wild type and $Atm^{-/-}$ at the Chr19 hotspot pair, with the scheme of the nested PCR assay below. The fold increase in the SPO11 oligos (RPM × 11.3) is indicated for each hotspot. Prior to amplification, genomic DNA is optionally digested with restriction enzymes to reduce amplification of parental products. The *Pip5k1b* exon located at the left hotspot is removed in many of the deletions.
(B) Representative gel of PCR products in $Atm^{-/-}$ from a 96-well plate. ~16,000 haploid genomes were seeded per PCR. Asterisks indicate deletions.
(C–F) Analysis of deletions. **C**, Deleted DNA sequences (black bars) from adult and juvenile testes. **D**, Arc diagram of deletions in $Atm^{-/-}$ (58 deletions total). See also **Fig. S3**. **E**, Distribution of individual breakpoints. **F**, Smoothed breakpoint and SPO11-oligo profiles (201-bp Hann filter for smoothing). Offsets between centers of hotspots and breakpoint distributions are indicated.

Chr1

SPO11 oligos (RPM)

All deletions

n=134

Atm⁻ᐟ⁻

Deletions with both breakpoints within 251 bp around hotspot centers

n=28 (21%)

Deletions with one breakpoint within 251 bp around hotspot centers

n=29 (22%)   n=25 (19%)

Deletions with both breakpoints outside 251 bp around hotspot centers

n=52 (39%)

-1500 -1000 -500 0 500 1000 1500 2000 2500 3000 3500 bp

251 bp around hotspot center   251 bp around hotspot center

89% of SPO11 oligos   72% of SPO11 oligos

Chr19

SPO11 oligos (RPM)

All deletions

n=58

Atm⁻ᐟ⁻

Deletions with both breakpoints within 251 bp around hotspot centers

n=11 (19%)

Deletions with one breakpoint within 251 bp around hotspot centers

n=10 (17%)   n=16 (28%)

Deletions with both breakpoints outside 251 bp around hotspot centers

n=21 (36%)

-1500 -1000 -500 0 500 1000 1500 2000 2500 3000 3500 bp

251 bp around hotspot center   251 bp around hotspot center

61% of SPO11 oligos   78% of SPO11 oligos

Figure S3. Distribution of deletions for the Chr1 and Chr19 hotspot pairs.
Deletion arcs from $Atm^{-/-}$ are segregated according to the positions of breakpoints relative to 251-bp zones around each hotspot center (yellow). The percentage of SPO11 oligos shown at bottom is the number of SPO11 oligos within each 251-bp zone divided by the total number of SPO11 oligos mapping between the left or right primer (arrows) and the midpoint between hotspot centers. Deletions often have one (red and blue arcs) or both breakpoints (violet arcs) within the 251-bp zones.

Figure S4. Distribution of deletions from experiments with or without prior genomic DNA digest.
Genomic DNA was optionally digested with restriction enzymes prior to PCR to suppress amplification of parental DNA. Nested primers are indicated with arrows. Restriction sites located in between the two hotspots of a pair are expected to be absent in deletion products. Consistent with this, the majority of deletions in $Atm^{-/-}$ at the Chr1 and Chr19 hotspot pairs had both breakpoints outside of the restriction site region (vertical blue lines). Similar results were obtained in experiments in which the digestion step was

omitted, or only the enzyme cutting uniquely inside this region was used (NcoI or KpnI, vertical black lines). A few deletion products contained restriction sites due to incomplete genomic DNA digestion ("Escapers", orange arcs) or were captured in experiments with no digest or when the unique central cutting enzyme was used ("Potential escapers", green arcs). Overall, deletion distributions were similar in all experiments, independent of the digest, with breakpoints clustered around hotspot centers. Note that some of these deletions had both breakpoints mapping to one hotspot of the hotspot pair (4 at the Chr1 pair; 1 at the Chr19 pair), likely arising from DSBs formed within one hotspot, as with microdeletions.

Figure S5. Deletions detected in the presence of ATM.
Deletions detected in *Atm*$^{+/+}$ (blue arcs) and *Atm*$^{+/-}$ (pink arcs) often had one breakpoint that was also found for a deletion in *Atm*$^{-/-}$ (red asterisks). The single deletion obtained from *Spo11*$^{-/-}$ at the Chr1 hotspot pair is also shown (grey arc).

A

B

Attempted double-strand gap repair

C

Precise insertion

Figure S6. Deletions in *Atm*$^{-/-}$ B6 × DBA F1 hybrid mice with models for haplotype switching.
(A) Arc diagram of deletions from the Chr1 hotspot pair (37 deletions total). Two interhomolog deletions are indicated (pink arcs).
(B, C). Models for haplotype switching in deletions. **B**, Attempted double-strand gap repair. Examples of one intrachromosomal (top) and one interhomolog (bottom) deletion with haplotype switches. In both cases, two DSBs produce a double-stranded gap between the two hotspots, releasing a double-cut fragment (lighter blue or red double line). One DSB end at the gap invades the homolog and is extended by DNA synthesis, but a DSB in the repair template leads to aborted synthesis, such that this end is ejected and joins the other end of the gap (intrachromosomal end joining) or the end of a DSB formed at the other hotspot in the homolog (interhomolog end joining). In the case of interhomolog deletions, it is also possible that only one DSB is formed on the chromosome initiating strand invasion. Thus, in both examples at least three DSBs occur, indicating extensive damage to multiple chromatids at adjacent hotspots when ATM is absent. The substantial template damage in this scenario may favor nonhomologous end-joining mechanisms. **C**, Precise insertion due to homologous annealing. An example of an intrachromosomal deletion with a haplotype switch. At a minimum four DSBs are formed, involving double cutting within one hotspot on one chromosome and at both hotspots on the other chromosome. A released double-cut fragment from one hotspot is precisely inserted at the double-stranded gap in the homolog through annealing of homologous sequences exposed by resection. Importantly this scenario depends on the relative position of the two DSBs in chromosomes. If the relative position were switched (bottom right), resection would not expose homologous sequences. Nonhomologous insertion is not compatible because it would produce a small deletion (bottom right) or a small duplication (bottom left). As we do not observe these types of repair products in our deletions, we consider the attempted double-strand gap repair scenario more likely.

A

B

Recurring deletions on Chr1

Microhomology	Microhomology length (bp)	No. deletions	Deletion (bp)	Strain
–	0	2	1416	B6
CT	2	2	1868	B6, B6/DBA
AGCC	4	2	1943	B6, B6/DBA
ACCCT	5	2	2372	B6
AGACA	5	2	2816	B6, B6/DBA
CACTGC	6	2	2371	B6
GGTTTAGA	8	8	2371	B6 (6), B6/DBA (2)
AGCCAGTCA	9	2	1624	B6, B6/DBA
	median 6 bp			

Inverted microhomology	Microhomology length (bp)	No. deletions	Deletion/Insertion (bp)	Strain
CACTGCCCTGGA	12*	2	2437/356	B6

24 of total 185 deletions

Figure S7. Microhomology at deletion breakpoints in *Atm*[−/−].
(A) Distribution of microhomology lengths for the Chr1 and Chr19 hotspot pairs deletion breakpoints. Only deletions without insertions are included (**Table S2**).
(B) Recurring deletions at the Chr1 hotspot pair in B6 inbred and B6 × DBA F1 hybrid mice. Recurring deletions often had longer microhomologies at breakpoints. Microhomologies within the larger imperfect repeat described in the text are marked in different colors as in **Fig. 3C**. A recurring deletion with inverted insertion (asterisk) with a 12-bp microhomology at one insert breakpoint is also shown (**Fig. 3C, S9B**).

Figure S8

Figure S8. Derivation of inserts mapping to DSB hotspots on other chromosomes.
SPO11 oligos at the genomic locations of inserts (blue bars) are shown for $Atm^{-/-}$ and $Atm^{+/+}$ and indicated as to whether they were called as hotspots in one or both genotypes or instead contain SPO11-oligo clusters. Inserts mapping to the non-PAR ChrY are shown in **Fig. S10**. One insert mapping to a microsatellite repeat is not shown. See also **Table S3**.

Figure S9. Characteristics of insertions in *Atm*⁻/⁻.

(A) Inverted insertions from the same hotspot pair have a broader distribution of lengths than inserts derived from other chromosomes. Median insert lengths are indicated. See also **Tables S3, S4**.

(B) Deletions with inverted insertions at the Chr1 hotspot pair in B6 and B6 × DBA mice. Asterisks denote deletions where one insert breakpoint had the same microhomology as the site with recurring deletions (**Fig. 3C**). Note that for the two deletions on the DBA chromosome, the inserts also derived from the DBA chromosome.

Figure S10. Inserts from the non-PAR ChrY map to multiple DSB hotspots.
Three inserts identified in $Atm^{-/-}$ deletions map to non-PAR ChrY repeats (**Table S3**), some of which have called hotspots. SPO11-oligo profiles are shown only for those repeats containing hotspots in $Atm^{-/-}$, $Atm^{+/+}$, or both, with the position of the inserts (blue bars) for each hotspot indicated. Inserts 1 and 2 are part of the same repeat family, and in one case they map to a single hotspot. Inserts 1 and 3 map to three and two hotspots, respectively. SPO11 oligos map uniquely to these hotspots ("Unique") or also map to other copies of the repeats ("Multi"). It remains possible that other repeat locations to which SPO11 oligos do not map uniquely also contain hotspots. Given that all insert-matching repeats (67 total) contain an intact PRDM9 motif, it is possible that DSBs are formed in many such repeats. The locations of multi-mapping SPO11 oligos were imputed, meaning they were distributed among mapped positions proportionally to local densities of uniquely mapping SPO11 oligos[9].

Figure S11. Evaluation of reads with microdeletions from deep sequencing of a single hotspot.
(A) Frequency of reads that had microdeletions of the indicated lengths. Four overlapping amplicons across the hotspot (1–4) were deep sequenced using MiSeq. For each amplicon, 576 separate PCRs were carried out with a total input of 9.6 million haploid genome equivalents, using a limited number of amplification cycles (see Methods). The PCR products for each amplicon were then pooled for sequencing. Four MiSeq runs were performed (I–IV) with the indicated amplicons and genotypes. For clarity, sequencing results are grouped by run (Top) or genotype (Bottom). At top, note that reads with small deletions (2–4 bp) were frequently observed in all amplicons for each genotype, including $Spo11^{-/-}$, and that their frequency varied between sequencing runs, indicating a significant batch effect. Short deletions (e.g., 2 bp) are known to arise from PCR and sequencing errors[63], and we saw that they became progressively less abundant in all genotypes up to 10 bp. Excluded from the 4-bp deletions is a product that was represented by >1000 reads in amplions 1–3. By contrast, reads with deletions >10 bp were more frequent in $Atm^{-/-}$ than in the other genotypes. At bottom, note that reads with 11–25-bp microdeletions were more frequent in $Atm^{-/-}$. Therefore, 11 bp was used as the lower cutoff to define a microdeletion for subsequent analyses. Our sequencing design (pooling of many low-cycle PCRs) is expected to minimize PCR jackpots and reduce instances where PCR duplicates from the same input DNA molecule are sequenced. However, two amplicons in $Atm^{+/+}$ showed multiple reads for one microdeletion each (amplicon 1: 26-bp microdeletion, 14 reads; amplicon 2: 31-bp microdeletion, 22 reads; **Table S5**), which therefore may be PCR duplicates. Thus, in some subsequent analyses we applied a stringent definition of independent events by including only one read for a given deletion product from a given amplicon.
(B) Recurring microdeletions within the hotspot. Top, smoothed (21-bp Hann filter) and unsmoothed SPO11-oligo maps at the Chr1 hotspot in $Atm^{-/-}$. Bottom, recurring microdeletions. These identical microdeletions (black bars) were found in independent PCRs for different amplicons. i) In $Atm^{-/-}$, amplicons that yielded the recurring microdeletions are indicated. For example, the microdeletion with the AG microhomology arose in amplicons 3 and 4. Note that one or both breakpoints for each of the recurring microdeletions occured at SPO11-oligo peaks (dashed vertical lines). ii) In $Atm^{+/-}$ and $Atm^{+/+}$, some microdeletions were identical to those found in $Atm^{-/-}$, in some cases in multiple amplicons. For example, the microdeletion with the GGAAGGAG microhomology was recovered in all three genotypes, but in different amplicons ($Atm^{-/-}$, 1; $Atm^{+/-}$, 2; $Atm^{+/+}$, 4). iii) Some of the microdeletions found in $Spo11^{-/-}$ recurred in other genotypes. Note that these map to the same region of the hotspot away from the center.
(C) Many of the duplicated reads in $Atm^{-/-}$ amplicons likely represent independent microdeletion events rather than PCR duplicates. For each amplicon, we randomly sampled with replacement from a set of reads representing all of the unique microdeletions observed. The sampled group was equal in number to the duplicated reads actually obtained for that amplicon. Then, the sampled reads with microdeletions mapping to the region where amplicons 1, 2, and 3 overlap were combined, and their breakpoint distribution was compared with the distribution of the observed duplicate reads. Breakpoint profiles of independently obtained reads, as in **Fig. 5C**, and of all reads are also shown. In the smoothed breakpoint profiles of the observed duplicates, the left breakpoints were particularly enriched in a peak that matched the central peak of SPO11 oligos, and the right breakpoints were enriched at a position that matched the secondary peak of SPO11 oligos. These patterns were not recapitulated by random sampling: both the left and right breakpoints were more widely dispersed without such clear enrichment coincident with the SPO11-oligo peaks. Because the observed duplicates were more likely than the random sample to have breakpoints aligned with the strongly favored locations of DSBs, these findings are consistent with the conclusion that most of the duplicated reads represent independent microdeletion events.

Figure S12. Extended analysis of microdeletions by genotype and microhomology.
(A) Arc diagrams of microdeletions stratified by length from *Atm*$^{-/-}$, as in **Fig. 5C**, with microdeletions in duplicated reads below. Microdeletions >160 bp are not shown, as their location may be more constrained by restricting analysis to the region where the amplicons overlap.

(B) Microdeletion length distribution for all amplicons by genotype for independent reads. Scatter plots and corresponding density plots are shown (>10 bp); densities in $Atm^{+/-}$, $Atm^{+/+}$, and $Spo11^{-/-}$ are normalized to $Atm^{-/-}$ to reflect read numbers in each genotype. As in the analysis of microdeletions in the overlap region of amplicons 1, 2, and 3 (**Fig. 5D**), distinct size classes were observed in the presence and absence of ATM, but not in $Spo11^{-/-}$. Larger microdeletion size classes (~140 bp, ~180 bp, ~230 bp) were well defined when all amplicons are considered. A smaller size class (~15 bp) was more prominent outside of the overlap region.

(C) Arc diagrams of the ~140-bp, ~180-bp, and ~230-bp microdeletions observed for $Atm^{-/-}$.

(D) Microhomology lengths at microdeletion breakpoints in the presence and absence of ATM. Analyses were performed for independent microdeletions in the overlapping region between amplicons 1, 2, and 3.

Table S1. Frequency of deletions in testis from individual mice.

Strain background	Hotspot pair	Mouse and genotype	No. haploid genomes	No. deletions	Frequency
B6	Chr1 adult	1794 $Atm^{-/-}$	5400000	67	12.41×10^{-6}
		*1793 $Atm^{+/+}$	5400000	0	$< 0.19 \times 10^{-6}$
		*2018 $Atm^{-/-}$	2066667	8	3.87×10^{-6}
		*2017 $Atm^{+/-}$	2066667	0	$< 0.48 \times 10^{-6}$
		*2066 $Atm^{+/-}$	1600000	0	$< 0.63 \times 10^{-6}$
		*2067 $Atm^{+/+}$	2266668	0	$< 0.44 \times 10^{-6}$
		*2656 $Atm^{-/-}$	1400000	6	4.29×10^{-6}
		*2657 $Atm^{+/-}$	3200000	0	$< 0.31 \times 10^{-6}$
		*2732 $Atm^{+/-}$	6400000	2	0.31×10^{-6}
		*2970 $Atm^{+/-}$	2000000	0	$< 0.50 \times 10^{-6}$
		*3162 $Atm^{+/+}$	6400000	0	$< 0.16 \times 10^{-6}$
		*3312 $Spo11^{-/-}$	6400000	0	$< 0.16 \times 10^{-6}$
		*3345 $Spo11^{-/-}$	6400000	0	$< 0.16 \times 10^{-6}$
		*3424 $Spo11^{-/-}$	1600000	1	0.63×10^{-6}
		Total $Atm^{-/-}$	8866667	81	9.14×10^{-6}
		Total $Atm^{+/-}$	15266667	2	0.13×10^{-6}
		Total $Atm^{+/+}$	14066668	0	$< 0.071 \times 10^{-6}$
		Total $Spo11^{-/-}$	14400000	1	0.07×10^{-6}
B6	Chr1 juvenile	*1 $Atm^{-/-}$	3200000	24	7.50×10^{-6}
		*2 $Atm^{+/-}$	3200000	0	$< 0.31 \times 10^{-6}$
		*3 $Atm^{+/+}$	3200000	0	$< 0.31 \times 10^{-6}$
F1 B6/DBA	Chr1 adult	2654 $Atm^{-/-}$	3600000	7	1.94×10^{-6}
		2675 $Atm^{-/-}$	5200000	30	5.77×10^{-6}
		2674 $Atm^{+/-}$	3200000	0	$< 0.31 \times 10^{-6}$
		2647 $Atm^{+/+}$	3200000	0	$< 0.31 \times 10^{-6}$
		Total $Atm^{-/-}$	8800000	37	4.20×10^{-6}
B6	Chr19 adult	*1793 $Atm^{+/+}$	5400000	2	0.37×10^{-6}
		*2018 $Atm^{-/-}$	266667	2	7.50×10^{-6}
		*2017 $Atm^{+/-}$	400000	0	$< 0.25 \times 10^{-5}$
		*2066 $Atm^{+/-}$	1600000	0	$< 0.63 \times 10^{-6}$
		*2067 $Atm^{+/+}$	1866667	0	$< 0.54 \times 10^{-6}$
		*2656 $Atm^{-/-}$	3200000	7	2.19×10^{-6}
		*2657 $Atm^{+/-}$	3200000	1	0.31×10^{-6}
		2733 $Atm^{-/-}$	4800000	29	6.04×10^{-6}
		*2732 $Atm^{+/-}$	4800000	3	0.63×10^{-6}
		2734 $Atm^{-/-}$	400000	2	5.00×10^{-6}
		2731 $Atm^{+/-}$	400000	0	$< 2.5 \times 10^{-6}$
		2780 $Atm^{-/-}$	400000	2	5.00×10^{-6}
		2779 $Atm^{+/-}$	400000	0	$< 2.5 \times 10^{-6}$
		2971 $Atm^{-/-}$	2000000	5	2.50×10^{-6}
		*2970 $Atm^{+/-}$	2000000	0	$< 0.50 \times 10^{-6}$
		*3162 $Atm^{+/+}$	6400000	0	$< 0.16 \times 10^{-6}$
		*3312 $Spo11^{-/-}$	6400000	0	$< 0.16 \times 10^{-6}$
		*3345 $Spo11^{-/-}$	6400000	0	$< 0.16 \times 10^{-6}$
		*3424 $Spo11^{-/-}$	1600000	0	$< 0.63 \times 10^{-6}$
		Total $Atm^{-/-}$	11066667	47	4.25×10^{-6}
		Total $Atm^{+/-}$	12800000	4	0.31×10^{-6}
		Total $Atm^{+/+}$	13666667	2	0.15×10^{-6}
		Total $Spo11^{-/-}$	14400000	0	$< 0.069 \times 10^{-6}$
B6	Chr19 juvenile	*1 $Atm^{-/-}$	1600000	11	6.88×10^{-6}
		*2 $Atm^{+/-}$	3200000	0	$< 0.31 \times 10^{-6}$
		*3 $Atm^{+/+}$	3200000	1	0.31×10^{-6}

Littermates are grouped. * The same mice used for both hotspot pairs.

Table S2. Deletions detected in *Atm-/-* with the analyses performed.

Hotspot pair	Strain background (primer set)[a]	No. PCR reactions	DNA (ng) seeded in PCR	No. haploid genomes	Total No. deletions	No. of deletions with insertions	Analyses performed
Chr1	B6 (b6)	168	25	1400000	16	0	frequency, breakpoint distribution, microhomology, insertions, recurring deletions
		640	50	10666667	89	12	frequency, breakpoint distribution, microhomology, insertions, recurring deletions
		72	100	2400000	29	3	breakpoint distribution, insertions, microhomology, recurring deletions
				Total Chr1 from B6	134	15	
	B6 (b6/dba)	48	50	640000	2	1	insertions, microhomology, recurring deletions
	B6/DBA (b6/dba)	528	50	8800000	37	2	frequency, breakpoint distribution, microhomology, insertions, recurring deletions
	B6/DBA (b6)	144	50	2400000	12	1	insertions, microhomology, recurring deletions
				Total Chr1	185	19	
Chr19	B6	8	10	266667	1	0	breakpoint distribution, microhomology, insertions, recurring deletions
		760	50	12666667	58[b]	8	frequency, breakpoint distribution, insertions, microhomology, recurring deletions
				Total Chr19	59[b]	8	

[a] For the Chr1 hotspot pair, two different primer sets were used as indicated.

[b] For one deletion event, one breakpoint could not be precisely mapped.

Table S3. Insertions derived from other hotspots.

Hotspot pair	Insert origin	Insert start	Insert end	SPO11 activity at insert origin *Atm*[-/-]	*Atm*[+/+]	Insert length (bp)	Deletion length (bp)	Microhomology at insert breakpoints (left/right)
Chr1	Chr3	116341240	116341350	cluster	no	111	2343	CAC/none
	Chr6	116019448	116019796	hotspot	cluster	349	2341	CACT/A
	Chr6	83540780	83540885	hotspot	cluster	106	1967	AG/none
	Chr7	41695870	41695945	hotspot	hotspot	76	1596	GTA/none
	Chr8	87936133	87936219	hotspot	hotspot	87	1513	(GATT insertion)/none
	Chr9	56053260	56053364	cluster	no	105	1671	C/TG
	Chr12	91711551	91711567	hotspot	hotspot	17	3326	CAAAGTGTCTG/TCT
	ChrX	69476686	69476920	hotspot	hotspot	235	2108	AT/none
Chr1 B/D	Chr9	55392465	55392589	no	cluster	125	2399	C/A
Chr19	Chr16	35066202	35066440	hotspot	hotspot	239	2005	none/none
	Chr17	78359789	78359826	hotspot	hotspot	38	2307	CA/none
	Chr17	80408805	80408819	hotspot pair	hotspot pair	15	1995	T/CAT

Hotspot pair	Insert from the non-PAR ChrY	Insert start	Insert end	SPO11 activity at insert origin *Atm*[-/-]	*Atm*[+/+]	Insert length (bp)	Deletion length (bp)	Microhomology at insert breakpoints (left/right)
Chr1	insert 1* (2 of 26 repeats)	28785715	28785921	hotspot	hotspot	207	2253	A/G
		58748224	58748430	cluster	hotspot			
		90475833	90476039	no	hotspot**			
	insert 2* (1 of 15 repeats)	90476619	90476713	no	hotspot**	95	1939	AAACA/G
	insert 3 (3 of 26 repeats)	5658645	5658703	hotspot	cluster	59	1500	none/none
		5944098	5944156	hotspot	cluster			

* Inserts 1 and 2 are part of the same repeat; both match 12 copies of the repeat.
** Same hotspot.

| Chr19 | Insert from Chr1 | 88533343 | 88533493 | microsatelite repeat (CTT)$_{27}$ | | 148 | 2643 | ACTGAGCTATCT/CCTCCTCCT |

Table S4. Insertions derived from the hotspot pair.

Hotspot pair	Hotspot left or right	Insert length (bp)	Deletion length (bp)	Distance between deletion and insertion breakpoints (bp)	Microhomology at insert breakpoints (left/right)
Chr1	left	709	2590	4	none/GAAGG
	left	22	2411	28	none/AGA
	right	343	2334	11	TCCAGGGCAGTGC/none
	right	356	2437	0	TCCAGGGCAGTG/none
	right	356	2437	0	TCCAGGGCAGTG/none
Chr1 B/D	left	493	2025	40	TTG/AAAG
	right	302	2465	56	TCCAGGGCAGT/ATTC
	right	216	2399	3	GT/T
Chr19	left	204	2591	0	none/C
	right	231		3	CTCTCTC/none
	left	130	1361	0	none/ATAGCT
	left	132	1949	0	none/TAGCT
	right	16	2063	44	GCTCT/A

Table S5. Frequency of microdeletions in individual amplicons.

Genotype	Amplicon	No. reads	No. reads with deletions (unique[a])	Frequency (adjusted[b])	P value[c] versus _Atm_[-/-] (adjusted[b])	P value[c] _Spo11_[-/-] (adjusted[b])
Atm[-/-]	1	3611491	60 (36)	1.66 x 10[-5]		
	2	3072566	65 (39)	2.12 x 10[-5]		
	3	3714378	50 (43)	1.35 x 10[-5]		
	4	3240515	41 (31)	1.27 x 10[-5]		
	Total	13638950	216 (149)	1.58 x 10[-5]		
Atm[+/-]	1	3615027	9 (7)	0.25 x 10[-5]	<0.0001	0.0161
	2	4036817	4 (4)	0.10 x 10[-5]	<0.0001	0.0825
	3	5334246	17 (16)	0.32 x 10[-5]	<0.0001	0.0592
	4	4704066	10 (10)	0.21 x 10[-5]	<0.0001	0.2799
	Total	17690156	40 (37)	0.23 x 10[-5]	<0.0001	0.4670
Atm[+/+]	1	5099332	23 (8)	0.45 (0.20) x 10[-5]	<0.0001 (<0.0001)	0.0004 (0.0387)
	2	4402709	53 (12)	1.20 (0.73) x 10[-5]	0.002 (<0.0001)	<0.0001 (0.0104)
	3	3360847	6 (4)	0.18 x 10[-5]	<0.0001	0.6158
	4	3253036	5 (4)	0.15 x 10[-5]	<0.0001	0.1313
	Total	16115924	87 (28)	0.54 (0.33) x 10[-5]	<0.0001 (<0.0001)	<0.0001 (0.0173)
Spo11[-/-]	1	3339211	1 (1)	0.03 x 10[-5]	<0.0001	
	2	2915459	8 (7)	0.27 x 10[-5]	<0.0001	
	3	4485773	6 (6)	0.13 x 10[-5]	<0.0001	
	4	3568282	12 (9)	0.34 x 10[-5]	<0.0001	
	Total	14308725	27 (23)	0.19 x 10[-5]	<0.0001	

[a]Total unique deletions obtained in that amplicon; note that the same deletion can be obtained independently in the overlapping amplicon.
[b]Adjusted frequencies and P values take into account potential jackpot events, in which 14 reads were obtained for the same microdeletion in amplicon 1, and 22 for another microdeletion in amplicon 2. Because these microdeletions were not obtained or were obtained just once in other amplicons they are considered to have arisen from PCR overamplification.
[c]Chi-square test.