Strengthening The Organization and Reporting of Microbiome Studies (STORMS): A Reporting Checklist for Human Microbiome Research

Chloe Mirzayi, Audrey Renson, Fatima Zohra, Shaimaa Elsafoury, Ludwig Geistlinger, Lora Kasselman, Kelly Eckenrode, Janneke van de Wijgert, Amy Loughman, Francine Z. Marques, STORMS Consortium, Genomic Standards Consortium, Massive Analysis and Quality Control Society, Nicola Segata, Curtis Huttenhower*, Jennifer B. Dowd*, Heidi E. Jones*, Levi Waldron*+

*Equal contribution
+Correspondence to: levi.waldron@sph.cuny.edu

Abstract

Background: Human microbiome research is a growing field with the potential for improving our understanding and treatment of diseases and other conditions. The field is interdisciplinary, making concise organization and reporting of results across different styles of epidemiology, biology, bioinformatics, translational medicine, and statistics a challenge. Commonly used reporting guidelines for observational or genetic epidemiology studies lack key features specific to microbiome studies.

Methods: A multidisciplinary group of microbiome epidemiology researchers reviewed elements of available reporting guidelines for observational and genetic studies and adapted these for application to culture-independent human microbiome studies. New reporting elements were developed for laboratory, bioinformatic, and statistical analyses tailored to microbiome studies, and other parts of these checklists were streamlined to keep reporting manageable.

Results: STORMS is a 17-item checklist for reporting on human microbiome studies, organized into six sections covering typical sections of a scientific publication, presented as a table with space for author-provided details and intended for inclusion in supplementary materials.

Conclusions: STORMS provides guidance for authors and standardization for interdisciplinary microbiome studies, facilitating complete and concise reporting and augments information extraction for downstream applications.

Availability: The STORMS checklist is available as a versioned spreadsheet from https://www.stormsmicrobiome.org/.

Introduction

Changes in the human microbiome have been associated with many disease and health states (1). However, reporting the results of human microbiome research is challenging as it often involves approaches from microbiology, genomics, biomedicine, bioinformatics, statistics, epidemiology, and other fields, resulting in a lack of consistent recommendations for reporting of methods and results. Inconsistent reporting can have consequences for the field by affecting the reproducibility of study results (2). While researchers have called for better reporting standards,(3) such as the Genomic Standards Consortium's MIxS checklist,(4) to provide a means for reporting sampling, processing and data generation; no comprehensive standardized guidelines spanning laboratory and epidemiological reporting have been proposed.

Standard reporting guidelines promote research consistency and, as a consequence, encourage reproducibility and improved study design. Editorial adoption of the CONsolidated Standards Of Reporting Trials (CONSORT) guidelines, for example, has been associated with an increase in quality of trial reporting.(5,6) Other epidemiological reporting guidelines have seen broad adoption, such as Strengthening the Reporting of OBservational studies in Epidemiology (STROBE)(7) and STrengthening the REporting of Genetic Association Studies (STREGA).(8) STROBE-metagenomics(9) proposes an extension to the STROBE checklist for metagenomics studies. Subsequent to the Minimum Information About a Microarray Experiment (MIAME) (9), Minimum Information about a MARKer gene Sequence (MIMARKS) and Minimum Information about any (x) Sequence (MIxS) checklists provide detailed guidance on reporting of sequencing studies in general. These are focused on the technical aspects of data generation, however, as are projects such as the Microbiome Quality Control Project (MBQC) (10) and International Human Microbiome Standards (IHMS).(11)(12) Together, these serve as useful foundations, but do not span the full range of reporting of human microbiome studies, include items intended for other types of studies, and provide limited guidance on manuscript preparation.

Studies of the human microbiome share many features with other types of molecular epidemiology, but also require unique considerations with their own methodological best practices and reporting standards. In addition to standard elements of epidemiological study design, culture-independent microbiome studies involve collection, handling and preservation of biological specimens, evolving approaches to laboratory processing with elevated potential for batch effects, bioinformatic processing, statistical analysis of sparse, unusually-distributed, high-dimensional data, and reporting of results on potentially thousands of microbial features.(13–15) Because there is no agreed-upon gold-standard method for microbiome research and the field has not reached consensus on many of these aspects, inconsistencies in reporting inhibit reproducibility and hamper efforts to draw conclusions across similar studies.

For these reasons, we convened a multi-disciplinary working group to develop guidelines tailored to microbiome study reporting. Members of this group include epidemiologists, biostatisticians, bioinformaticians, physician-scientists, genomicists, and microbiologists. The checklist is designed to balance completeness with burden of use, and is applicable to a broad

78 range of human microbiome study designs and analysis. The "Strengthening The Organization
79 and Reporting of Microbiome Studies (STORMS)" checklist draws relevant items from related
80 guidelines and adds new tailored guidelines to serve as a tool to organize study planning and
81 manuscript preparation, to improve the clarity of manuscripts, and to facilitate reviewers and
82 readers in assessing these studies.

Methods

Origin and development

The origins of these guidelines are rooted in a project to create a standardized database
of published literature reporting relationships between the microbiome and disease. The goal of
that project is to create a publicly available, standardized database of microbiome study findings
indexed by condition of interest (e.g. disease, health status, diet, environmental factor)
microbiome site (e.g. gut, mouth, skin), and microbial taxonomy to aid comparative analysis. As
of August 2020, 22 curators have extracted findings from 371 unique published studies.
Included studies must have examined the relationship between the microbiome and a condition
of interest and included findings on a taxonomic level (even if all findings were null).

This review revealed substantial reporting heterogeneity, particularly for epidemiology,
such as study design, confounding factors, and sources of bias. It also revealed microbiome-
specific issues, including statistical analysis of compositional relative abundance data and
handling 'batch' effects.(16) This heterogeneity highlighted the need for standardized reporting
guidelines, similar to those used in other fields of study. The curators determined that
standardized reporting guidelines would streamline the review process, but would more
importantly help researchers throughout the field of microbiome research communicate their
findings effectively.
101

The resulting multidisciplinary group of bioinformaticians, epidemiologists,
biostatisticians, and microbiologists was thus convened to discuss microbiome reporting
standards. The group began by reviewing existing reporting standards including STROBE,(17)
STROBE-ME,(18) STREGA,(8) MICRO,(19), MIMARKS,(4) and STROGAR.(20) The group also
reviewed existing articles containing recommendations for microbiome reporting.(21,22) The
STROBE and STREGA guidelines were used as a starting point for the STORMS checklist,
although aspects were incorporated from the other reporting standards.
109

Following the reporting standards development guidelines recommended by EQUATOR,
a comprehensive list of potential guideline items was created. From this list, group members
added, modified, and removed items based on their expertise. After the first round of edits, the
checklist was then applied to a recent microbiome study(23) by group members. Comments,
removals, and additions were harmonized after each round. Based on this process, additional
changes, simplifications, and clarifications were made. This process was repeated until there
was a group consensus that the checklist was ready for use.
117

In addition to the core working group, outside subject matter experts identified by
working group members were then invited to review the guidelines and provide feedback as
members of the STORMS Consortium. In total, 113 experts were invited to join the STORMS
Consortium and 61 accepted and provided feedback. Substantive feedback (i.e. not grammar,
spelling, or other small changes) was compiled into a feedback document (Supplement 1) and
the core working group responded to each piece of feedback individually. Additional revisions
were made based on their comments. After this round of revisions, consortium members were
once again invited to review the checklist prior to submission for publication.

Results

Checklist

The latest version of the checklist at time of publication is presented in Table 1 and a summary
of items is presented in Figure 1. Of the items in the latest version in the STORMS checklist,
nine items or sub-items were unchanged from STROBE, three were modified from STROBE,
one was modified from STREGA, and 57 new guidelines were developed. Nine Items with an
overlap with MIxS are specified. Rationale for new and modified items are presented below.
Documentation of items unmodified from STROBE and STREGA were presented in the
publications of those checklists.

Abstract (1.0-1.3)

Along with commonly included abstract materials such as a basic description of the participants
and results, authors should report the study design(24)--such as cross-sectional, case-control,
cohort, or randomized controlled trial--in the abstract of their article (item 1.1), as required by
other reporting guidelines. Communicating study design in the abstract allows readers to quickly
categorize the type of evidence provided. As part of this basic description, sequencing methods
should be mentioned (item 1.2). Body site(s) sampled should also be included (item 1.3).

Introduction (2.0-2.1)

The introduction should clearly describe the underlying background, evidence, or theory that
motivated the current study (item 2.0). Among other possibilities, this could include pilot study
data, previous findings from a similar study or topic, or a biologically plausible mechanism that
has been proposed. This clarifies for the reader the motivations for the present study. If the
study is exploratory in nature, it should be explained what motivated the current exploration and
the goals of the exploratory study. The hypothesis developed based on the background should
be included. If the study was exploratory and did not define a hypothesis, pre-specified study
objectives should be included (item 2.1).

Methods (3.0-8.5)

Participants (3.0-3.9)

The methods section should contain sufficient information for study replicability. Because study design is essential to understanding the study, it should be stated in the methods (item 3.0). When describing the participants in the study, the population of interest should be described and how participants were sampled from the source population (item 3.1). Because participant characteristics such as environment,(25) lifestyle behaviors, diet, biomedical interventions, demographics,(26) and geography(27) (item 3.2) can correspond with substantial differences in the microbiome, it is essential to include this description. Temporal context can be quite important as well so start and end dates for recruitment, follow-up, and data collection should be stated (item 3.3).

Specific criteria used to assess potential participants for eligibility in the study should also be reported, detailing both inclusion and exclusion criteria (item 3.4). Inclusion and exclusion criteria are pre-established characteristics used for selection of participants into the study, and describing these criteria is essential in understanding the study's target population.(28) This is expanded from STROBE, which requires eligibility criteria, but does not specify that both inclusion and exclusion criteria should be reported in detail. Any information collected about antibiotics or other treatments that could affect the microbiome should be described (item 3.5) as well as if any exclusion criteria included recent antibiotic or other medication use.

The final analytic sample sizes and read numbers should be stated as well as the reason for any exclusion of participants at any step of the recruitment, follow-up, or laboratory processes (item 3.6). STROBE suggests using a flow diagram to show when and why participants were removed from the study. Examples of such flow diagrams are presented in Figure 2. If participants were lost to follow-up or did not complete all assessments in a longitudinal study, details on how follow-ups were conducted should be stated and time-point specific sample sizes should also be reported (item 3.7). Additionally, studies that matched cases to controls should describe what variables were used in matching (item 3.4).

Laboratory methods (4.0-4.17)

Since STROBE does not cover laboratory methods, new items were developed for STORMS. Describe laboratory methods in sufficient detail to allow replication. The handling of lab samples should be described, including procedures for sample collection (items 4.1), shipping (item 4.2), and storage (item 4.3).

Because DNA extraction can be a major source of technical differences across studies,(10) DNA extraction methods should be described (item 4.4). Human DNA removal and microbial DNA enrichment, if performed, should also be included (item 4.5). Likewise, if positive controls (item 4.7), negative controls (item 4.8), or contaminant mitigation methods (item 4.9) were used, they should be identified and described.

192   Sequencing-related methods should be reported. This includes primer selection and DNA
193   amplification (including the variable region of the 16S rRNA gene if applicable) (item 4.6). Major
194   division of sequencing strategy, such as shotgun or amplicon sequencing, should be identified
195   (item 4.11). Finally, the methods used to determine relative abundances should be explained
196   (item 4.12) and the read numbers that serve as denominators recorded.
197
Batch effects should be discussed as a potential source of confounding, including steps taken to
ensure batch effects do not overlap with exposures or outcomes of interest (item 4.13).(29) If
conducting metatranscriptomics, metaproteomics, or metabolomics, provide details of those
methods (items 4.14 to 4.16).

Data sources/measurement (5.0)

For non-microbiome data (e.g. health outcomes, participant socioeconomic, behavioral, dietary,
biomedical characteristics, including disease location and activity, and environmental variables),
the measurement and definition of each variable should be described (item 5.0). For instance,
participant sex and age could be obtained from electronic medical records or from a
questionnaire distributed to participants; this data source should be described. Limitations of
measurement may also be discussed including potential bias due to misclassification or missing
data, as well as any attempts made to address these measurement issues.

Research design considerations for causal inference (6.0-6.1)

Observational data is often used to test associations that aim towards causal inference.
Methods include, for example, the use of multivariable analysis or matching to adjust for
confounding variables between a hypothesized exposure (such as abundance of a microbial
taxon) and the disease or condition under study.(30) If variables are adjusted for in the analysis,
theoretical justification for inclusion of these variables should be provided (item 6.0). As part of
this theoretical justification, consider including a directed acyclic graph showing the
hypothesized causal relationships of interest.(31,32) In addition to considering the theoretical
motivations for the present study, discuss the potential for selection or survival bias which can
distort the observed relationship between the microbiome and variable of interest (item 6.1). For
example, such bias may occur due to loss-to-follow-up (in longitudinal studies) or due to
participants not being included in the study due to the condition itself (e.g. participants who have
died of aggressive forms of colorectal cancer have not survived to be in a hypothetical study of
colorectal cancer microbiomes).(33)

Bioinformatics and Statistical Methods (7.0-7.9)

Adequate description of bioinformatic and statistical methods is essential to producing a
rigorous and reproducible research report. Data transformations (such as normalization,
rarefaction, and percentages) should be described (item 7.0). Quality control methods should be
fully disclosed so that readers understand criteria for filtering or removing reads or samples
(item 7.1). All statistical methods used to analyze the data should be stated, (item 7.3) including
how results of interest were selected (e.g. using a p-value, q-value, or other threshold) (item
7.8). Taxonomic, functional profiling, or other sequence analysis methods should be described
in detail (item 7.2) In the interest of reproducibility, all software, packages, databases, and

libraries used for the pre-processing and analysis of the data should be described and cited
including version numbers (item 7.9).

Reproducible Research (8.0-8.5)

Reproducible research practices serve as quality checks in the process of publication and
further transparency and knowledge sharing, as detailed in the rubric proposed by Schloss.(34)
Journals are increasingly implementing reproducible research standards that include the
publishing of data and code, and those guidelines should be followed when possible.(35,36)
STORMS itemizes the accessibility of data, methods, and code (items 8.0 through 8.5). If
possible, data should be deposited into a public repository (items 8.1 and 8.2). If data or code
are not or cannot be made publicly available, a description of how interested readers can
access the data should be provided. As stated in item 8.0, any protected information should be
described along with how such data can be accessed.

Results (9.0-10.4)

Descriptive Data (9.0)

Descriptive statistics about the study population should be reported (item 9.0). At a minimum,
age and sex of all participants should be reported, but other important participant characteristics
should be reported when possible including medication use or lifestyle factors such as diet.
Consider reporting these data in a descriptive statistics table. Packages such as the table1
package in R make creating such a table straightforward.(37)

Outcome Data (10.0-10.4)

The main outcomes of the study should be detailed including descriptive information, findings of
interest, and the results of any additional analyses. Descriptive microbiome analysis (for
instance, dimension reduction such as Principal Coordinates Analysis, measures of diversity,
gross taxonomic composition) should be reported for each group and each time point (item
10.0). This contextualizes the results of differential abundance analysis for readers. When
reporting differential abundance test results, the magnitude and direction of differential
abundance should be clearly stated (item 10.2) for each identifiable standardized taxonomic unit
(item 10.1). Results from other types of analyses such as metabolic function, functional
potential, MAG assembly, and RNAseq should be described in the results as well (items 10.3
and 10.4). Additional results (e.g. non-significant results or full differential abundance results)
can be included in supplements, but should not be excluded entirely. Although the problem has
been known for decades,(35) journals across many fields are recognizing the issue of
publication bias and therefore the issue of non-reporting of null results.(38) Including such
results in publications will help to reduce the severity of this bias and improve future systematic
reviews and meta-analyses.

Discussion (11.0-14.0)
Most recommendations for the Discussion section are similar to STROBE including a discussion
of the limitations of the present study and associated methods (item 13.0). One additional
recommendation is made: discuss the potential for biases and how they would influence the

study findings (item 13.1). Many forms of bias such as residual/unmeasured confounding, bias
related to compositional analysis,(39) measurement bias, or selection bias(40) could affect the
interpretation of the results of the study and it is important to acknowledge potential sources of
bias when discussing the results.(41) As described in STROBE, authors should also consider
the generalizability of their findings and if these findings could be applicable to the target
population or other populations (item 13.2). If different forms of bias were not assessed or
assumed to be negligible, this should be stated. Finally, authors should discuss potential future
or ongoing research based on findings of the present study (item 14.0).

Other Information (15.0-17.0)

In addition to a statement of funding (item 15.0), authors should also include acknowledgements
and conflicts of interest statements (items 15.1 and 15.2, respectively). Conflicts of interest
statements should be written according to the criteria established by the journal. Finally, the
paper should state where supplementary materials and data can be accessed (items 16.0 and
17.0).

Discussion

The STORMS checklist for reporting on human microbiome studies was developed with
the following priorities. The checklist should 1) be easy to understand and use by researchers
from various fields, through straightforward use of language and pruning of items rarely relevant
to the current literature, 2) be organized in the outline of a manuscript, so it can serve as a tool
for authors and for peer reviewers, particularly when included in manuscript submission as a
supplemental table with comments, 3) assist in the complete and organized reporting of a study,
not in enforcing any particular methods, 4) reuse or modify items from related checklists where
relevant, and 5) represent consensus across a broad cross-section of the human microbiome
research community. The checklist facilitates manuscript authors in providing a complete,
concise, and organized description of their study and its findings. Included as a supplemental
table to a manuscript, it also supports efficient peer review and post-publication interpretation.
298
While other efforts for extending STROBE for microbiome and metagenomic have been
proposed (9) and laboratory-focused reporting checklists have been released,(4,42) to our
knowledge STORMS is the first comprehensive reporting checklist for human microbiome
research. We aim for the STORMS guidelines to improve the quality and transparency of
microbiome epidemiology studies by introducing a shared grammar of study reporting in a
structured checklist format. Reporting checklists introduced in other disciplines have been
shown to improve the quality of journal articles.(5,6)
306
A major strength of STORMS is the rigor and transparency of its development by a
diverse, multidisciplinary consortium of subject matter experts. All feedback received from
consortium members, and responses, have been included as Supplement 1. The development
of STORMS is an ongoing process, and new versions of the checklist will be released to reflect
evolving standards and technological processes. A version control system with changelog has
been implemented, and annual reviews of the checklist are planned. Additionally, the working

group plans to evaluate STORMS impact on microbiome reporting by examining how many articles are fulfilling checklist items before and after its release. We invite interested readers to join the STORMS Consortium by contacting the corresponding author or by visiting the consortium website for more information (https://www.stormsmicrobiome.org/). We also encourage journals to include the STORMS checklist in their instructions to authors and advise peer reviewers to consult the checklist when reviewing submissions.

There are some limitations to the STORMS checklist. The checklist was not created to assess study or methodological rigor. It is meant to aid authors' organization and ease the process of reader assessment of how studies are conducted and analyzed. Conclusions about the quality of studies should not be made based on their adherence to STORMS guidelines, although we expect the reporting guidelines to help readers review studies critically. It does not encourage, discourage, or assume the use of null hypothesis significance testing(43) or methods of compositional data analysis(44), topics of some controversy in the field. In general the checklist avoids reference to or guidance on specific statistical methodological decisions.

Through the efforts of the STORMS Consortium working in an iterative, transparent, and collaborative process, the STORMS checklist provides a roadmap for researchers in reporting the results of a human microbiome study. The STORMS Consortium believes that the checklist is sufficiently flexible and user-friendly to support widespread adoption and contribution to microbiome study standards. Its adoption will ideally encourage thoughtful study design, reproducibility, collaboration, and open knowledge sharing between research groups as they explore the human microbiome.

Bibliography

1. Wirbel J, Pyl PT, Kartal E, et al. Meta-analysis of fecal metagenomes reveals global microbial signatures that are specific for colorectal cancer. *Nat. Med.* 2019;25(4):679–689.
2. Simoneau J, Dumontier S, Gosselin R, et al. Current RNA-seq methodology reporting limits reproducibility. *Brief. Bioinform.* [electronic article]. (https://academic.oup.com/bib/advance-article/doi/10.1093/bib/bbz124/5669860). (Accessed September 21, 2020)
3. Ten Hoopen P, Finn RD, Bongo LA, et al. The metagenomic data life-cycle: standards and best practices. *GigaScience.* 2017;6(8):1–11.
4. Yilmaz P, Kottmann R, Field D, et al. Minimum information about a marker gene sequence (MIMARKS) and minimum information about any (x) sequence (MIxS) specifications. *Nat. Biotechnol.* 2011;29(5):415–420.
5. Moher D, Jones A, Lepage L, et al. Use of the CONSORT Statement and Quality of Reports of Randomized Trials: A Comparative Before-and-After Evaluation. *JAMA.* 2001;285(15):1992–1995.
6. Plint AC, Moher D, Morrison A, et al. Does the CONSORT checklist improve the quality of reports of randomised controlled trials? A systematic review. 2006;185(5):5.
7. Vandenbroucke JP, Elm E von, Altman DG, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): Explanation and Elaboration. *PLOS Med.* 2007;4(10):e297.
8. STrengthening the REporting of Genetic Association Studies (STREGA)—an extension of the STROBE statement - Little - 2009 - Genetic Epidemiology - Wiley Online Library. (https://onlinelibrary.wiley.com/doi/abs/10.1002/gepi.20410). (Accessed January 29, 2020)

9.   Bharucha T, Oeser C, Balloux F, et al. STROBE-metagenomics: a STROBE extension
statement to guide the reporting of metagenomics studies. *Lancet Infect. Dis.*
2020;20(10):e251–e260.
10. Sinha R, Abu-Ali G, Vogtmann E, et al. Assessment of variation in microbial community
amplicon sequencing by the Microbiome Quality Control (MBQC) project consortium. *Nat.*
Biotechnol. 2017;35(11):1077–1086.
11. Costea PI, Zeller G, Sunagawa S, et al. Towards standards for human fecal sample
processing in metagenomic studies. *Nat. Biotechnol.* 2017;35(11):1069–1076.
12. Santiago A, Panda S, Mengels G, et al. Processing faecal samples: a step forward for
standards in microbial community analysis. *BMC Microbiol.* 2014;14(1):112.
13. Turner P, Fox-Lewis A, Shrestha P, et al. Microbiology Investigation Criteria for Reporting
Objectively (MICRO): a framework for the reporting and interpretation of clinical
microbiology data. *BMC Med.* 2019;17(1):70.
14. STROGAR – STrengthening the Reporting Of Genetic Association studies in
Radiogenomics - ScienceDirect.
(https://www.sciencedirect.com/science/article/pii/S0167814013003630). (Accessed
January 29, 2020)
15. Sinha R, Abnet CC, White O, et al. The microbiome quality control project: baseline study
design and future directions. *Genome Biol.* 2015;16(1):276.
16. Badal VD, Wright D, Katsis Y, et al. Challenges in the construction of knowledge bases for
human microbiome-disease associations. *Microbiome.* 2019;7(1):129.
17. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational
Studies in Epidemiology (STROBE) statement: guidelines for reporting observational
studies. *J. Clin. Epidemiol.* 2008;61(4):344–349.
18. Gallo V, Egger M, McCormack V, et al. STrengthening the Reporting of OBservational
studies in Epidemiology - Molecular Epidemiology (STROBE-ME): an extension of the
STROBE statement. *Eur. J. Clin. Invest.* 2012;42(1):1–16.
19. Microbiology Investigation Criteria for Reporting Objectively (MICRO): a framework for the
reporting and interpretation of clinical microbiology data | BMC Medicine | Full Text.
(https://bmcmedicine.biomedcentral.com/articles/10.1186/s12916-019-1301-1). (Accessed
January 29, 2020)
20. Kerns SL, Ruysscher D de, Andreassen CN, et al. STROGAR – STrengthening the
Reporting Of Genetic Association studies in Radiogenomics. *Radiother. Oncol.*
2014;110(1):182–188.
21. Sinha R, Ahsan H, Blaser M, et al. Next steps in studying the human microbiome and health
in prospective studies, Bethesda, MD, May 16–17, 2017. *Microbiome* [electronic article].
2018;6. (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6257978/). (Accessed September
20, 2019)
22. Hornung BVH, Zwittink RD, Kuijper EJ. Issues and current standards of controls in
microbiome research. *FEMS Microbiol. Ecol.* [electronic article]. 2019;95(5).
(https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6469980/). (Accessed March 16, 2020)
23. Min Y, Ma X, Sankaran K, et al. Sex-specific association between gut microbiome and fat
distribution. *Nat. Commun.* [electronic article]. 2019;10.
(https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6546740/). (Accessed September 24, 2020)
24. Pearce N. Classification of epidemiological study designs. *Int. J. Epidemiol.*
2012;41(2):393–397.
25. Jin Y, Wu S, Zeng Z, et al. Effects of environmental pollutants on gut microbiota. *Environ.*
Pollut. 2017;222:1–9.
26. Chen L, Zhang Y-H, Huang T, et al. Gene expression profiling gut microbiota in different
races of humans. *Sci. Rep.* 2016;6:23075.
27. Yatsunenko T, Rey FE, Manary MJ, et al. Human gut microbiome viewed across age and

geography. *Nature*. 2012;486(7402):222–227.

28. Patino CM, Ferreira JC, Patino CM, et al. Inclusion and exclusion criteria in research
studies: definitions and why they matter. *J. Bras. Pneumol.* 2018;44(2):84–84.

29. Yan L, Ma C, Wang D, et al. OSAT: a tool for sample-to-batch allocations in genomics
experiments. *BMC Genomics*. 2012;13:689.

30. VanderWeele TJ. Principles of confounder selection. *Eur. J. Epidemiol.* 2019;34(3):211–
219.

31. Textor J, van der Zander B, Gilthorpe MS, et al. Robust causal inference using directed
acyclic graphs: the R package 'dagitty.' *Int. J. Epidemiol.* 2017;dyw341.

32. Loughman A, Ponsonby A-L, O'Hely M, et al. Gut microbiota composition during infancy and
subsequent behavioural outcomes. *EBioMedicine*. 2020;52:102640.

33. Schooling CM. Selection bias in population-representative studies? A commentary on
Deaton and Cartwright. *Soc. Sci. Med. 1982*. 2018;210:70.

34. Schloss PD. Identifying and Overcoming Threats to Reproducibility, Replicability,
Robustness, and Generalizability in Microbiome Research. *mBio*. 2018;9(3).

35. Rosenthal R. The file drawer problem and tolerance for null results.
(http://pages.ucsd.edu/~cmckenzie/Rosenthal1979PsychBulletin.pdf). (Accessed March 23,
2020)

36. Munafò M, Neill J. Null is beautiful: On the importance of publishing null results. *J.*
Psychopharmacol. (Oxf.). 2016;30(7):585–585.

37. Rich B. benjaminrich/table1. (Accessed October 1,
2020).(https://github.com/benjaminrich/table1). (Accessed October 1, 2020)

38. Null is beautiful: On the importance of publishing null results - Marcus Munafò, Jo Neill,
2016. (https://journals.sagepub.com/doi/full/10.1177/0269881116638813). (Accessed March
23, 2020)

39. McLaren MR, Willis AD, Callahan BJ. Consistent and correctable bias in metagenomic
sequencing experiments. *eLife*. 2019;8.

40. Hernán M, Robins, JM. Causal Inference: What If. Boca Raton, FL: Chapman & Hall/CRC;
2020 (Accessed May 29, 2020).(https://www.hsph.harvard.edu/miguel-hernan/causal-
inference-book/). (Accessed May 29, 2020)

41. Writing a discussion section: how to integrate substantive and statistical expertise | BMC
Medical Research Methodology | Full Text.
(https://bmcmedresmethodol.biomedcentral.com/articles/10.1186/s12874-018-0490-1).
(Accessed March 23, 2020)

42. Raes J, Foerstner KU, Bork P. Get the most out of your metagenome: computational
analysis of environmental sequence data. *Curr. Opin. Microbiol.* 2007;10(5):490–498.

43. Szucs D, Ioannidis JPA. When Null Hypothesis Significance Testing Is Unsuitable for
Research: A Reassessment. *Front. Hum. Neurosci.* [electronic article]. 2017;11.
(https://www.frontiersin.org/articles/10.3389/fnhum.2017.00390/full). (Accessed June 17,
2020)

44. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, et al. Microbiome Datasets Are
Compositional: And This Is Not Optional. *Front. Microbiol.* [electronic article]. 2017;8.
(https://www.frontiersin.org/articles/10.3389/fmicb.2017.02224/full?report=reader).
(Accessed June 17, 2020)

Tables and Figures

Number	Item	Recommendation	Item Source	Additional Guidance	Yes/ No/ NA	Comments or location in manuscript
Abstract						
1.0	Structured or Unstructured Abstract	Abstract should include information on background, methods, results, and conclusions in structured or unstructured format.	STORMS			
1.1	Study Design	State study design in abstract.	STORMS	See 3.0 for additional information on study design.		
1.2	Sequencing methods	State the strategy used for metagenomic classification.	STORMS	For example, targeted 16S by qPCR or sequencing, shotgun metagenomics, metatranscriptomics, etc.		
1.3	Specimens	Describe body site(s) studied.	STORMS			
Introduction						
2.0	Background and Rationale	Summarize the underlying background, scientific evidence, or theory driving the current hypothesis as well as the study objectives.	STORMS			
2.1	Hypotheses	State the pre-specified hypothesis. If the study is exploratory, state any pre-specified study objectives.	STORMS			
Methods						
3.0	Study Design	Describe the study design.	STORMS	Observational (Case-Control, Cohort, Cross-sectional survey, etc.) or Experimental (Randomized controlled trial, Non-randomized controlled trial, etc.). For a brief description of common study designs see: DOI: 10.11613/BM.2014.022

If applicable, describe any blinding (e.g. single or double-blinding) used in the course of the study. | | |
| 3.1 | Participants | State what the population of interest is, and the method by which participants are | STORMS | Examples of the population of interest could be: adults with no chronic health | | |

			sampled from that population. Include relevant information on physiological state of the subjects or stage in the life history of disease under study when participants were sampled.		conditions, adults with type II diabetes, newborns, etc. This is the total population to whom the study is hoped to be generalizable to. The sampling method describes how potential participants were selected from that population. If the participants are from a substudy of a larger study, provide a brief description of that study and cite that study. Clearly state how cases and controls are defined. An example of relevant physiological state might be pre/post menopausal for a vaginal microbiome study; examples of stage in the life history of disease could be whether specimens were collected during active or dormant disease, or before or after treatment.	
3.2		Geographic location	State the geographic region(s) where participants were sampled from.	MIxS: geographic location (country and/or sea,region)	Geographic coordinates can be reported to prevent potential ambiguities if necessary.	
3.3		Relevant Dates	State the start and end dates for recruitment, follow-up, and data collection.	STORMS	Recruitment is the period in which participants are recruited for the study. In longitudinal studies, follow-up is the date range in which participants are asked to complete a specific assessment. Finally, data collection is the total period in which data is being collected from participants including during initial recruitment through all follow-ups.	
3.4		Eligibility criteria	List any criteria for inclusion and exclusion of recruited participants.	Modified STROBE	Among potential recruited participants, how were some chosen and others not? This could include criteria such as sex, diet, age, health status, or BMI.	

				If there is a primary and validation sample, describe inclusion/exclusion criteria for each.		
3.5	Antibiotics Usage	List what is known about antibiotics usage before or during sample collection.	STORMS	If participants were excluded due to current or recent antibiotics usage, state this here. Other factors (e.g. proton pump inhibitors, probiotics, etc.) that may influence the microbiome should also be described as well.		
3.6	Analytic sample size	Explain how the final analytic sample size was calculated, including the number of cases and controls if relevant, and reasons for dropout at each stage of the study. This should include the number of individuals in whom microbiome sequencing was attempted and the number in whom microbiome sequencing was successful.	STORMS	Consider use of a flow diagram. Also state sample size in abstract. If power analysis was used to calculate sample size, describe those calculations.		
3.7	Longitudinal Studies	For longitudinal studies, state how many follow-ups were conducted, describe sample size at follow-up by group or condition, and discuss any loss to follow-up.	STORMS	If there is loss to follow-up, discuss the likelihood that drop-out is associated with exposures, treatments, or outcomes of interest.		
3.8	Matching	For matched studies, give matching criteria.	Modified STROBE	"Matched" refers to matching between comparable study participants as cases and controls or exposed / unexposed. Indicate whether participants were individual or frequency matched and in what ratio were they matched (e.g. 1 case to 1 control).		
3.9	Ethics	State the name of the institutional review board that approved the study and protocols, protocol number and date of approval, and procedures for obtaining informed consent from participants.	STORMS			
4.0	Laboratory methods	State the laboratory/center where laboratory work was done.	STORMS	Provide a reference to complete lab protocols if previously published elsewhere such as on protocols.io. Note any modifications of lab protocols and the reason for protocol modifications.		
4.1	Specimen collection	State the body site(s) sampled from and how specimens were collected.	MIxS: sample	Use terms from the Uber-anatomy Ontology		

			collection device or method; host body site	(https://www.ebi.ac.uk/ols/ontologies/uberon) to describe body sites in a standardized format.		
4.2	Shipping	Describe how samples were stored and shipped to the laboratory.	STORMS	Include length of time from collection to receipt by the lab and if temperature control was used during shipping.		
4.3	Storage	Describe how the laboratory stored samples, including time between collection and storage and any preservation buffers or refrigeration used.	STORMS	State where each procedure or lot of samples was done if not all in the same place.

Include reagent/lot/catalogue #s for storage buffers. | | |
4.4	DNA extraction	Provide DNA extraction method, including kit and version if relevant.	MIxS: nucleic acid extraction	If any DNA quantification methods were used prior to DNA amplification or at the pooling step of library preparation, state so here.		
4.5	Human DNA sequence depletion or microbial DNA enrichment	Describe whether human DNA sequence depletion or enrichment of microbial or viral DNA was performed.	STORMS			
4.6	Primer selection	Provide primer selection and DNA amplification methods as well as variable region sequenced (if applicable).	MIxS: pcr primers			
4.7	Positive Controls	Describe any positive controls (mock communities) if used.	STORMS	If used, should be deposited under guidance provided in the 8.X items.		
4.8	Negative Controls	Describe any negative controls if used.	STORMS	If used, should be deposited under guidance provided in the 8.X items.		
4.9	Contaminant mitigation and identification	Provide any laboratory or computational methods used to control for or identify microbiome contamination from the environment, reagents, or laboratory.	STORMS	Includes filtering of reagents and other steps to minimize contamination. It is relevant to state whether the specimens of interest have low microbial load, which makes contamination especially relevant.		
4.10	Replication	Describe any biological or technical replicates included in the sequencing, including which steps were replicated between them.	STORMS	Replication may be biological (redundant biological specimens) or technical (aliquots taken at different stages of analysis) and used in extraction, sequencing, preprocessing, and/or data analysis.		
4.11	Sequencing strategy	Major divisions of strategy, such as shotgun or amplicon sequencing.	MIxS: sequencing	For amplicon sequencing (for example, 16S variable region), state the region		

			method	selected. State the model of sequencer used.		
4.12	Sequencing methods	State whether experimental quantification was used (QMP/cell count based, spike-in based) or whether relative abundance methods were applied.	STORMS	These include read length, sequencing depth per sample (average and minimum), whether reads are paired, and other parameters.		
4.13	Batch effects	Discuss any likely sources of batch effects, if known. Detail any blocking or randomization used in study design to avoid confounding of batches with exposures or outcomes.	STORMS	Sources of batch effects include sample collection, storage, library preparation, and sequencing and are commonly unavoidable in all but the smallest of studies.		
4.14	Metatranscriptomics	Detail whether any mRNA enrichment was performed and whether/how retrotranscription was performed prior to sequencing. Provide size range of isolated transcripts. Describe whether the sequencing library was stranded or not. Provide details on sequencing methods and platforms.	STORMS	Provide details on any internal standards which may have been used as well as parameters and versions of any software or databases used.		
4.15	Metaproteomics	Detail which protease was used for digestion. Provide details on proteomic methods and platforms (e.g. LC-MS/MS, instrument type, column type, mass range, resolution, scan speed, maximum injection time, isolation window, normalised collision energy, and resolution).	STORMS	Provide details on any internal standards which may have been used as well as parameters and versions of any software or databases used.		
4.16	Metabolomics	Detail which fractions were obtained (polar and/or non-polar) and how these were analysed. Provide details on metabolomics methods and platforms (e.g. derivatisation, instrument type, injection type, column type and instrument settings).	STORMS	Provide details on any internal standards which may have been used as well as parameters and versions of any software or databases used.		
5.0	Data sources/ measurement	For each non-microbiome variable, including the health condition, intervention, or other variable of interest, state how it was defined, how it was measured or collected, and any transformations applied to the variable prior to analysis.	MIxS: host disease status	State any sources of potential bias in measurements, for example multiple interviewers or measurement instruments, and whether these potential biases were assessed or accounted for in study design. Use terms from a standardized ontology such as the Experimental Factor Ontology (https://www.ebi.ac.uk/efo/) to describe variables of interest in a standardized format.		

6.0	Research design for causal inference	State which variables are controlled and state the rationale for controlling for them.	STORMS	Most observational research can be classified as predictive, descriptive, or causal. For causal inference, this item refers to describing the assumptions that would be required to draw causal inferences from observational data. See Hernán MA, Robins JM (2020). Causal Inference: What If. Boca Raton: Chapman & Hall/CRC for more details on causal inference. For example, hypothesized confounders may be controlled for by multivariable adjustment, but colliders or mediators should not be. Consider using a Directed Acyclic Graph (DAG) to sumarize hypothesized causal pathways.		
6.1	Selection bias	Discuss potential for selection or survival bias.	STORMS	Selection bias can occur when some members of the target study population are more likely to be included in the study/final analytic sample than others. Some examples include survival bias (where part of the target study population is more likely to die before they can be studied), convenience sampling (where members of the target study population are not selected at random), and loss to follow-up (when probability of dropping out is related to one of the things being studied).		
7.0	Bioinformatic and Statistical Methods	Describe any transformations to quantitative variables used in analyses (e.g. use of percentages instead of counts, normalization, rarefaction, categorization).	STORMS	If a variable is analyzed using different transformations, state rationale for the transformation and for each analyses which version of the variable is used. In case of any complex or multistep transformations, give enumerated instructions for reproducing those transformations.		
7.1	Quality Control	Describe any methods to identify or filter low quality reads or samples.	MIxS: sequence quality check	If samples were excluded based on quality or read depth, list the criteria used, the number of samples excluded, and the final sample size after quality		

				control.		
7.2	Sequence analysis	Describe any taxonomic, functional profiling, or other sequence analysis performed.	MIxS: feature prediction; similarity search method			
7.3	Statistical methods	Describe all statistical methods.	Modified STROBE	Describe any statistical tests used, exploratory data analysis performed, dimension reduction methods/unsupervised analysis, alpha/beta metrics, and/or methods for adjusting for measurement bias. If multiple statistical methods are possible, discuss why the methods used were selected. If a multiple hypothesis testing correction method was used, describe the type of correction used. State which taxonomic levels are analyzed.		
7.4	Longitudinal analysis	If the study is longitudinal, include a section that explicitly states what analysis methods were used (if any) to account for grouping of measurements by individual or patterns over time.	STORMS			
7.5	Subgroup analysis	Describe any methods used to examine subgroups and interactions.	STROBE			
7.6	Missing data	Explain how missing data were addressed.	STROBE	"Missing data" refers to participant measurements such as covariates, exposures, outcomes, or time points that should have been collected but were not, not to zeros in taxonomic abundance tables or data points not applicable to that observation.		
7.7	Sensitivity analyses	Describe any sensitivity analyses.	STROBE			
7.8	Findings	State criteria used to select findings for reporting.	STORMS	For example, false discovery rate with total number of tests, effect size threshold, significance threshold,		

				microbes of interest.		
7.9	Software	Cite all software (including read mapping software) and databases (including any used for taxonomic reference or annotating amplicons, if applicable) used. Include version numbers.	Modified STREGA	Installed packages, add-ons or libraries should be stated and cited in addition to the software used. All parameters employed that differ from the default of that software/version should be provided. This is in addition to, not a replacement for, publishing of code as outlined in the section Reproducible Research.		
8.0	Reproducible research	Make a statement about whether and how others can reproduce the reported analysis.	STORMS	Any protected information that has been excluded or provided under controlled access should be listed along with any relevant data access procedures. "On request from authors" is not sufficiently detailed; formal data access procedures and conditions should be defined. If data are unavailable, state so clearly. Consider using a specialized rubric for reproducible research (such as: https://mbio.asm.org/content/9/3/e00525-18.short). Consider preregistering the study protocol (such as on osf.io or https://plos.org/open-science/preregistration/).		
8.1	Raw data access	State where raw data may be accessed including demultiplexing information.	STORMS	Robust, long-term databases such as those hosted by NCBI and EBI are preferred. If using a private repository, provide rationale.		
8.2	Processed data access	State where processed data may be accessed.	STORMS	Unfiltered data should be provided. Robust, long-term databases such as those hosted by NCBI and EBI are preferred. If using a private repository, provide rationale.		
8.3	Participant data access	State where individual participant data such as demographics and other covariates may	STORMS	If re-categorized, transformed, or otherwise derived variables were used		

		be accessed, and how they can be matched to the microbiome data.		in the analysis, these variables or code for deriving them should be provided. Examples of how participant data can be matched to microbiome data are: using the same set of anonymized identifiers, or using different anonymized identifiers but providing a map. Provided data should be sufficient to independently replicate the current analysis.		
8.4	Source code access	State where code may be accessed.	STORMS	If a standard or formalized workflow was employed, reference it here.		
8.5	Full results	Provide full results of all analyses, in computer-readable format, in supplementary materials.	STORMS	For example, any fold-changes, p-values, or FDR values calculated, provided as a spreadsheet. Use a machine-readable, plain-text format such as csv or tsv.		
Results						
9.0	Descriptive data	Give characteristics of study participants (e.g. dietary, demographic, clinical, social) and information on exposures and potential confounders.	STROBE	Typically reported in a table included in the paper or as a supplementary table. Indicate number of participants with missing data for each variable of interest. This includes environmental and lifestyle factors that may affect the relationship between the microbiome and the condition of interest. Participant diet and medication use should be summarized, if known. At minimum, age and sex of all participants should be summarized.		
10.0	Microbiome data	Report descriptive findings for microbiome analyses with all applicable outcomes and covariates.	STORMS	This includes measures of diversity as well as relative abundances. These descriptive findings should be reported both for the sample overall and for individual groups.		
10.1	Taxonomy	Identify taxonomy using standardized taxon	STORMS	If not using full taxonomic hierarchy,		

		classifications that are sufficient to uniquely identify taxa.		make sure it is clear whether names stated are species, genera, family, etc. Italicize genus/species pairs. Consult journal guidelines or standardized references on taxonomic nomenclature. For instance, https://wwwnc.cdc.gov/eid/page/scientific-nomenclature		
10.2	Differential abundance	Report results of differential abundance analysis by the variable of interest and (if applicable) by time, clearly indicating the direction of change and total number of taxa tested.	STORMS	If there are more than two groups, include omnibus (multigroup) test results if applicable to the research question. If applicable, reported effect sizes should include a measure of uncertainty such as the confidence interval.		
10.3	Other data types	Report other data analyzed--e.g. metabolic function, functional potential, MAG assembly, and RNAseq.	STORMS			
10.4	Other statistical analysis	Report any statistical data analysis not covered above.	STORMS	This could include subgroup analysis, sensitivity analyses, and cluster analysis. Visualizations should be easily interpretable and colorblind-friendly. The caption and/or main text should provide a detailed description of visualizations for visually-impaired readers.		
Discussion						
11.0	Key results	Summarise key results with reference to study objectives	STROBE			
12.0	Interpretation	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence.	STROBE	Define or clarify any subjective terms such as "dominant," "dysbiosis," and similar words used in interpretation of results. When interpreting the findings, consider how the interpretation of the findings may be summarized or quoted for the general public such as in press releases or news articles.		

				If causal language is used in the interpretation (such as "alters," "affects," "results in," "causes," or "impacts"), assumptions made for causal inference should be explicitly stated as part of 6.0 and 13.0. Distinguish between function potential (ie inferred from metagenomics) and observed activity (ie metatranscriptomic, metabolomic, proteomic) if discussing microbial function.		
13.0	Limitations	Discuss limitations of the study, taking into account sources of potential bias or imprecision.	STROBE	Also consider limitations resulting from the methods (especially novel methods), the study design, and the sample size.		
13.1	Bias	Discuss any potential for bias to influence study findings.	STORMS	May include sampling method, representativeness of study participants, or potential confounding.		
13.2	Generalizability	Discuss the generalisability (external validity) of the study results	STROBE	To what populations or other settings do you expect the conclusions to generalize?		
14.0	Ongoing/future work	Describe potential future research or ongoing research based on the study's findings.	STORMS			
Other information						
15.0	Funding	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	STROBE			
15.1	Acknowledgements	Include acknowledgements of those who contributed to the research but did not meet critera for authorship.	STORMS	For general guidelines on authorship, see http://www.icmje.org and https://www.elsevier.com/authors/journal-authors/policies-and-ethics/credit-author-statement		
15.2	Conflicts of Interest	Include a conflicts of interest statement.	STORMS			
16.0	Supplements	Indicate where supplements may be accessed and what materials they contain.	STORMS			
17.0	Supplementary data	Provide supplementary data files of results with for all taxa and all outcome variables analyzed. Indicate the taxonomic level of all	STORMS	Depending on the analysis performed, examples of the supplemental results included could be mean relative		

		taxa.		abundance, differential abundance, raw p-value, multiple hypothesis testing-adjusted p-values, and standard error. All discussed taxa should include the taxonomic level (e.g. class, order, genus).		

454 Table 1. STORMS Checklist. For the latest version of the checklist, please visit https://www.stormsmicrobiome.org/.

Figure 1. STORMS checklist items by major manuscript heading. For detailed descriptions of each item and additional guidance, see the STORMS
checklist (Table 1).

460   Figure 2. Flowchart templates for item 3.6 (Analytic sample size).