Hidden heterogeneity and co-occurrence networks of soil prokaryotic

communities revealed at the scale of individual soil aggregates

Running title: Microbial diversity at soil aggregate-level

Márton Szoboszlay[1] and Christoph C. Tebbe[1]

[1]Thünen Institut für Biodiversität, Bundesforschungsinstitut für Ländliche Räume, Wald und

Fischerei, Braunschweig, Germany

Corresponding author: christoph.tebbe@thuenen.de

Address of corresponding author:

Prof. Dr. Christoph Tebbe, Thünen Institut für Biodiversität, Bundesallee 65, D-38116

Braunschweig, Germany

Abstract

Sequencing PCR-amplified gene fragments from metagenomic DNA is a widely applied method

for studying the diversity and dynamics of soil microbial communities. Typically DNA is extracted

from 0.25 to 1 g of soil. These amounts, however, neglect the heterogeneity of soil present at the

scale of soil aggregates; and thus, ignore a crucial scale for understanding the structure and

functionality of soil microbial communities. Here we show with a nitrogen-depleted agricultural

soil the impact of reducing the amount of soil used for DNA extraction from 250 mg to approx. 1

mg in order to access spatial information on the prokaryotic community structure as indicated by

16S rRNA-gene amplicon analyses. Furthermore, we demonstrate that individual aggregates from

the same soil differ in their prokaryotic communities. The analysis of 16S rRNA gene amplicon

sequences from individual soil aggregates allowed us, in contrast to 250 mg soil samples, to

construct a co-occurrence network that provides insight into the structure of microbial

associations in the studied soil. Two dense clusters were apparent in the network, one dominated

by *Thaumarchaeota*, known to be capable of ammonium oxidation at low N concentrations, and

the other by *Acidobacteria* subgroup 6 probably representing an oligotrophic lifestyle to obtain

energy from SOC. Overall this study demonstrates that DNA obtained from individual soil

aggregates provides new insights into how microbial communities are assembled.

Keywords

33   soil aggregates, soil microbial diversity, network analysis, spatial scales, soil DNA extraction,
34   microhabitats

Introduction

One of the major challenges in microbial ecology is to gain a predictive understanding of microbial diversity through elucidating the principles, patterns, and interactions which lead to the assembly of highly diverse microbial communities as found for example in soil (Green, Bohannan, & Whitaker, 2008). To achieve this, it is essential to consider temporal and spatial variation in micro-habitat conditions. In soil microbial ecology, the latter, however, is commonly ignored (Lombard, Prestat, van Elsas, & Simonet, 2011; Vos, Wolf, Jennings, & Kowalchuk, 2013). Instead, large composite samples are favored in order to obtain an overview of microbial diversity at the scale of a plot or a field. This neglects the fine scale heterogeneity of soil structure, and thus loses much information on patterns of community assembly (Thakur et al., 2020).

Soil structure develops as primary particles of different sizes and mineral composition, i.e., clay, silt, and sand, interact with each other, and with organic material to build microaggregates and macroaggregates, that are below or above 250 µm in diameter respectively (Six, Elliott, & Paustian, 2000). Most bacterial cells occur inside aggregates rather than on their surfaces (Ranjard, Poly, et al., 2000) and biogeochemical cycles, which are key ecosystem services driven by an interacting microbial community (Smith et al., 2015), are considered to mainly take place within aggregates (Wilpiszeski et al., 2019). Soil aggregates have been regarded as "massively concurrent evolutionary incubators" (Rillig, Müller, & Lehmann, 2017), or as "microbial villages" (Wilpiszeski et al., 2019), that represent small communities separated by distance and physical barriers and connected only periodically, e.g. during wetting events.

DNA-based methods to assess microbial diversity typically start with extracting 250 mg to 1 g of soil material (Young, Rawlence, Weyrich, & Cooper, 2014). This strategy has been useful to

investigate the overall microbial diversity of soils (Fierer & Jackson, 2006; Roesch et al., 2007), its

variation across geographical regions (Griffiths et al., 2011; Karimi et al., 2018), or its response to

land use change at a continental scale (Szoboszlay, Dohrmann, Poeplau, Don, & Tebbe, 2017).

However, in order to understand the processes and interactions taking place within soil microbial

communities, it would be rewarding to increase the spatial resolution of the community analysis

to individual aggregates and investigate microbial diversity in these spatial entities. Approaching

the "calling distance" of microbial interactions (Nunan, 2017), this would increase the likelihood

of detecting interacting microbial partners. Correlation networks have increasingly been applied

to reveal relationships between microbial community members as detected by PCR amplicon

sequence analyses (Banerjee et al., 2016; Barberan, Bates, Casamayor, & Fierer, 2012; Karimi et

al., 2020; F. Li, Chen, Zhang, Yin, & Huang, 2017). However, to interpret a positive correlation as

a mutualistic and a negative correlation as an antagonistic relationship is only possible for

community members sharing the same microhabitat (Weiss et al., 2016). Without distinction of

micro-habitats it is hard to assign the presence of taxa to niche exclusion (Faust & Raes, 2012).

Analyzing soil DNA extracted from 250 mg to 1 g represents mixed DNA from many microhabitats.

In contrast, working with individual soil aggregates, should strongly enhance the ecological

significance of soil microbial networks analyses.

The potential impact of the heterogeneous soil constituents on structuring microbial

communities at a microscale has already been demonstrated with pooled soil primary particles,

where the majority of abundant bacterial and fungal taxa exhibited particular preferences for

clay, silt, or sand-sized fractions (Hemkemeyer, Christensen, Tebbe, & Hartmann, 2019;

Hemkemeyer, Dohrmann, Christensen, & Tebbe, 2018). Furthermore, comparing pooled samples

of micro- and macroaggregates revealed that these two size classes also differ in microbial

community composition (Constancias et al., 2014; Davinic et al., 2012; Fox et al., 2018), diversity

(Bach, Williams, Hargreaves, Yang, & Hofmockel, 2018; Ivanova et al., 2015) and their response

to stress (Ranjard, Nazaret, et al., 2000). However, information on the heterogeneity of microbial

communities of individual aggregates within specific aggregate fractions is still lacking. A major

limitation of analyzing individual aggregates is the difficulty of obtaining a sufficient quantity of

nucleic acids from small amounts of soil for molecular methods. Attempts made so far, therefore,

either pooled several aggregates for DNA extraction (Bach et al., 2018; Bailey, Fansler, Stegen, &

McCue, 2013; Ivanova et al., 2015), sampled very large aggregates weighing 20 − 70 mg

(Kravchenko et al., 2014), or applied whole genome amplification (WGA) (Bailey, McCue, et al.,

2013). These solutions, however, do not deliver data on individual aggregates, provide coarse

spatial resolution, or generate substantial bias in the results (Direito, Zaura, Little, Ehrenfreund,

& Roling, 2014; Wang et al., 2016), respectively. To our knowledge, the only study that reported

the bacterial community composition in smaller, i.e. below 3 mm, individual aggregates without

applying WGA utilized taxonomic microarrays; a method of relatively low resolution, and focused

solely on linking enzyme activity profiles with community structure (Kim, Nunan, Dechesne,

Juarez, & Grundmann, 2015). Furthermore, applying molecular methods to small samples that

yield very low amounts of nucleic acids require validation to prove the consistent performance of

the methods and rule out the possibility of contamination and stochastic effects influencing the

results.

The tremendous scientific potential that individual soil aggregate-based microbial community

analysis should have for characterizing the heterogeneity of soil microbial communities at a

biologically and ecologically more meaningful scale motivated us testing the following hypotheses

in this study:

1. Metagenomic DNA of sufficient quantity and quality for PCR-based analyses can be

extracted from soil samples in the mg-range, thus representing the scale of macro-

aggregates

2. Increasing spatial resolution reveals heterogeneity in soil bacterial and archaeal

community structure and abundance

3. A higher heterogeneity seen among small soil samples is not a result of contamination or

sub-optimal performance of molecular methods

4. Comparing individual aggregates from the same soil unveils patterns of microbial co-

occurrence within the soil microbial community not seen with the commonly used

250 mg soil sample size.

Materials and methods

Overview of the experiments

Three experiments were conducted in this study. In the 1st experiment, samples decreasing in size

from 250 mg to 1 mg taken from the same soil were subjected to DNA extraction. To address the

first two hypotheses, qPCR and high-throughput amplicon sequencing targeting the 16S rRNA

gene were conducted to characterize the bacterial, archaeal, and fungal abundance and the

prokaryotic diversity in these samples. The 2nd experiment addressed the third hypothesis by

comparing 250 mg soil samples and aliquots of a homogenized soil slurry. The volumes of the

aliquots were chosen to contain the amount of DNA expected from 1 mg, 5 mg, and 25 mg soil

samples. Since all aliquots were taken from the same thoroughly homogenized soil slurry,

differences in prokaryotic community structure between these soil homogenate samples must be

results of contamination, stochastic effects, or sub-optimal performance of the DNA extraction

and PCR. In the 3rd experiment, DNA was extracted from individual aggregates and 250 mg soil

samples taken from the same soil to address the fourth hypothesis. All experiments included

several control samples to test for the presence of contamination.

Soil sampling and DNA extraction

The soil used in all experiments was a loam topsoil of a Haplic Chernozem (FAO classification)

from the Bad Lauchstädt experimental research station of the Helmholtz Centre for

Environmental Research in Germany (51°24'N 11°53'E) (Merbach & Schulz, 2013). It originated

form the Static Fertilization Experiment, initiated in the year 1902, and samples were collected in

December from a plot without any fertilization since 1903 (treatment NIL)(Ludwig et al., 2011)

which was under long term sugar beet – potato – winter wheat – barley rotation. Consequently,

the soil was compared to its fertilized variants depleted in nitrogen (Blair, Faulkner, Till,

Körschens, & Schulz, 2006). The soil samples had a pH value of 7.1 (in 0.01 M $CaCl_2$) and 17.7 mg

kg^{-1} organic C. It was sieved (2 mm mesh size) and stored at 4°C until use.

Before sampling, approximately 100 g soil was incubated at room T in the dark for 24 h. The soil

was then spread out in a sterile petri dish and samples were taken with sterilized spatulas directly

into the bead-beating tubes of the DNA extraction kit. Control samples were included in all

experiments. They received no soil but were handled together with and the same way as the soil

samples. DNA was extracted with the Quick-DNA Fecal/Soil Microbe Microprep Kit (Zymo

Research, Freiburg, Germany) including two 45 s bead beating cycles in an MP FastPrep-24 5G

Instrument (MP Biomedicals, Eschwege, Germany) at 6.5 m/s speed with a 300 s break in

between. The DNA extracts were eluted in 30 µl elution buffer. All work was done in a biosafety

cabinet decontaminated with UV light to minimize the chance of contamination. Measurement

of the DNA yield was attempted with Quant-iT PicoGreen dsDNA Assay Kit (Molecular Probes, Life

Technologies, Eugene, Or) but accurate quantification was not possible for many of the small

samples due to results close to the background fluorescence in the blank controls. In preparation

of this study, several DNA extractions methods were tested but were not found suitable. This

included the Fast DNA Spin kit for soil (MP Biomedicals, LLC, Illkrich, France) and variations of the

phenol-chloroform protocol (Miller, Bryant, Madsen, & Ghiorse, 1999).

In the 1st experiment, five different size-groups of soil samples were collected: 250 mg, 125 mg,

25 mg, 5 mg, and 1 mg, respectively. Eight samples were taken from each size-group along with

six control samples. Sample weights from all experiments are listed in Table 1. In the 2nd

experiment, ten samples of 250 mg soil and six control samples were taken. Eight of the soil

samples were processed normally in the DNA extraction, while for the other two, the DNA

extraction was interrupted after the centrifugation following the bead beating. By this point, the

samples had been turned into homogenized slurry by the bead beating, and the soil debris had

been separated from the supernatant that contained the metagenomic DNA. The supernatant

from the two samples were pooled and homogenized by vortexing. The mass of the resulting

suspension was 1 112 mg and it originated from 497 mg soil in total. Accordingly, a 55.9 mg aliquot

of this suspension contained the amount of DNA extractable from 25 mg soil, an 11.2 mg aliquot

the amount from 5 mg soil, and a 2.2 mg aliquot the amount from 1 mg soil. Eight aliquots from

each of these sizes, hereafter referred to as 25 mg, 5 mg, and 1 mg soil homogenate samples,

were taken and each mixed with 350 µl bashing bead buffer from the DNA extraction kit to

continue the DNA extraction. In the 3rd experiment, 37 individual soil aggregates, weighing 5.3

mg on average and similar in size (ca. 2 mm), were taken for DNA extraction along with 35 samples

of 250 mg soil and nine control samples.

Abundances of microbial groups assessed by qPCR

The abundance of *Bacteria*, *Archaea*, and *Fungi* were assessed by qPCR targeting the 16S rRNA

gene and the ITS region as previously described (Hemkemeyer, Christensen, Martens, & Tebbe,

2015). Archaeal and fungal abundance was only investigated in the 1st experiment. Reactions

were run in a Bio-Rad CFX96 real-time PCR system in duplicates from different dilutions of the

DNA extracts. In case of the 250 mg and 125 mg soil samples, 50- and 100-fold dilutions were

taken; from the 25 mg samples 10- and 20-fold dilutions; and from the 5 mg, 1 mg, individual

aggregate, and control samples undiluted DNA extracts and two-fold dilutions were used. PCR

efficiencies were 95.9 – 104.6% for the bacterial 16S rRNA gene, 94.2 – 96.2% for the archaeal

16S rRNA gene, and 83.3 – 84.3% for the fungal ITS with R^2 ≥ 0.995 in all cases. Results were

compared with Tukey's HSD tests, or Welch's t-test in case of the data from the 3rd experiment.

The analysis was carried out in R 3.4.4 (www.r-project.org). One of the 250 mg samples from the

1st experiment yielded a magnitude higher copy number in the fungal ITS qPCR assay than the

others. It was deemed to be an outlier and excluded from the analysis.

High-throughput sequencing of 16S rRNA gene amplicons and data processing

To characterize the bacterial and archaeal communities in the samples, DNA extracts were

subjected to high-throughput amplicon sequencing of the V4 region of the 16S rRNA gene

following the protocol of Kozich *et al.* (Kozich, Westcott, Baxter, Highlander, & Schloss, 2013) with

primers updated to match the modified 515f and 806r sequences according to Walters *et al.*

(Walters et al., 2016). In case of the small soil samples and control samples, due to the low DNA

yield, 10 µl DNA extract was used as template in the PCRs. Paired-end sequencing was done on

Illumina MiSeq instruments at StarSEQ (Mainz, Germany). Samples from the same experiment

were sequenced in the same run. All data is available at the European Nucleotide Archive under

the accession numbers PRJEB36881 (1st experiment), PRJEB36883 (2nd experiment), and

PRJEB36887 (3rd experiment).

The sequencing data from the three experiments were analyzed separately. Raw reads were

processed with the dada2 (version 1.6.0) pipeline (Callahan et al., 2016) in R 3.4.4. Forward and

reverse reads were truncated at positions 240 and 90, respectively. Reads with any ambiguous

bases were discarded as well as forward reads with over two and reverse reads with over one

expected error. The data from the 2nd experiment had higher quality allowing the reverse reads

to be truncated at position 130 and keeping those with two or less expected errors. Error models

were constructed from 10^6 randomly selected reads. Sequence variants (SVs) were inferred using

the pool option. Forward and reverse SVs were merged trimming overhangs, and the

removeBimeraDenovo function was employed to detect chimeras. The SVs were classified

according to the SILVA reference version 132 (Pruesse et al., 2007) accepting only results with

≥70% bootstrap support. SVs shorter than 220 nt or longer than 275 nt, or identified as chimeric,

mitochondrial, or chloroplast sequences, or not classified into *Bacteria* or *Archaea* were deleted.

Good's index was calculated to estimate the coverage of the SVs. SVs with ≥0.1% relative

abundance in any of the control samples of an experiment were regarded as potentially

contaminant and removed from the dataset of that experiment.

Analysis of the sequencing results

Principal component analysis (PCA) plots were created in R using the rda function of the vegan

package version 2.5-2 (Oksanen et al., 2018). To decrease the sparsity of the data, SVs not

reaching 0.1% relative abundance in any of the samples included in the PCA were removed.

Zeroes were replaced with the count zero multiplicative (CZM) method using the zCompositions

package version 1.1.1 (Palarea-Albaladejo & Martin-Fernandez, 2015) and centered log-ratio

(CLR) transformation was applied to the data to correct for compositional effects and differences

in sequencing depth (Gloor, Wu, Pawlowsky-Glahn, & Egozcue, 2016).

Aitchison distances between the samples were calculated as Euclidean distances in the CLR

transformed dataset (Gloor, Macklaim, Pawlowsky-Glahn, & Egozcue, 2017). SVs that didn't have

at least 0.1% relative abundance in any of the compared samples were removed and zeroes were

replaced with the CZM method to allow CLR transformation before calculating Euclidean

distances with the 'vegdist' function of the vegan package. Results were compared with Tukey's

HSD tests.

Plots illustrating the prevalence of abundant SVs among the soil samples were prepared in

Cytoscape 3.7.1 (www.cytoscape.org). CoNet 1.1.1 beta (K Faust & Raes, 2016) in Cytoscape was

used to construct co-occurrence networks from the data from the 3[rd] experiment. To limit the

number of parallel significance tests and the sparsity of the data, only SVs with ≥0.2% relative

abundance in at least one of the samples were included. Separate networks were constructed for

the 250 mg soil samples and the soil aggregates. However, the selection of SVs was done on the

joint data matrix to ensure that both networks include the same SVs. The data was relativized to

the total sequence count of each sample. Pearson and Spearman correlations, mutual

information (jsl setting), and Bray-Curtis and Kullback-Leibler (with a pseudo count of 10^{-8})

dissimilarities were calculated and the 1 000 highest and 1 000 lowest scoring edges from each of

the five metrics were kept. The ReBoot method (Faust et al., 2012), which mitigates compositional

effects, was used to assess the significance of the edges based on 1 000 permutations with

renormalization and 1 000 bootstrap iterations. In the network of the soil aggregates, edges with

scores below the 2.5^{th} and over the 97.5^{th} percentile of the bootstrap distribution or not

supported by at least three of the five metrics were considered unstable and removed. Brown's

method of p-value merging was applied followed by Benjamini-Hochberg correction. Only edges

with q ≤ 0.05 were included in the final network. In the network of the 250 mg samples, unstable

edges were not removed and the Benjamini-Hochberg correction was not applied as otherwise

no edges were retained. The networks were visualized in Cytoscape using the compound spring

embedder layout. Topological parameters were calculated with NetworkAnalyzer version 2.7

(Assenov, Ramirez, Schelhorn, Lengauer, & Albrecht, 2008).

Results

Microbial DNA can be extracted from soil samples in the mg-range

DNA could be extracted from soil samples as little as 0.87 mg as well as from intact soil aggregates. In all cases, the extracted DNA was sufficient for 16S rRNA gene amplicon sequencing and the qPCR assays. Accurate quantification of the DNA yield with PicoGreen assays was not possible because the fluorescence readings from many of the small samples were close to the background fluorescence in the blank controls.

To assess whether the DNA extraction could recover microbial DNA with similar efficiency from small quantities of soil as from 250 mg samples, estimates of the abundances of *Bacteria*, *Archaea*, and *Fungi* in one gram of soil were calculated from the qPCR results (Figure 1; Supplementary data file #1). Similar estimates were obtained from the 250 mg and 5 mg soil samples from the 1st experiment. Estimates from the 1 mg samples tended to be lower but were not significantly different. In contrast, the estimates of fungal abundance in a gram of soil were significantly lower from the 1 and 5 mg than from the 250 mg samples. Estimates of bacterial abundance obtained from the 25 mg and 125 mg samples and archaeal abundance from the 25 mg samples were significantly higher than from the 250 mg samples. Bacterial abundance estimates from the single aggregates and 250 mg soil samples of the 3rd experiment covered the same range (Figure S1), but the mean of the estimates from the single aggregates (2.27×10^9 copies/g soil) was lower (p < 0.001) than from the 250 mg samples (4.69×10^9 copies/g soil).

The control samples amplified in the qPCR assay targeting the bacterial 16S rRNA gene but yielded only 3 to 356 copies per µl DNA extract. In comparison, the 1 mg soil samples had 8 777 to 45 534

copies per µl DNA extract (Supplementary file 1). The control samples from the 1st experiment

had 0 to 13 fungal ITS copies per µl DNA extract and none of them showed amplification in the

archaeal 16S rRNA gene qPCR assays.

Removing potentially contaminant SVs from the sequencing results

It was possible to generate sequencing results from all control samples (the complete dataset

with the taxonomic classification of the SVs is in Supplementary file 2). Good's coverage index of

the SVs was over 0.993 in all of them indicating that almost their complete prokaryotic community

was captured by the sequencing (Table S1). They were very similar to each other in their

prokaryotic community structures but very different from the soil samples (Figure S2). Every SV

that reached 0.1% relative abundance in any of the control samples of an experiment was

considered as a potential contaminant. There were 450, 77, and 591 such SVs in the datasets of

the 1st, 2nd, and 3rd experiments respectively. In the data from the 1st experiment, these SVs

together covered 98.9 – 99.8% of the sequences obtained from the control samples and 2.5 –

6.6% of the sequences from the soil samples. In the 2nd and 3rd experiments, 92.5 – 99.5% and

98.6 – 99.6% of the sequences from the control samples, and 2.9 – 17.3% and 5.9 – 9.4% of the

sequences from the soil samples, respectively, were covered by the potentially contaminant SVs.

To mitigate the effect of contamination on the results, the potentially contaminant SVs were

deleted from the data matrices before further analysis.

Increasing spatial resolution reveals heterogeneity in soil bacterial and archaeal community structure but not in their abundance

The yield of high-quality 16S rRNA gene amplicon sequences was somewhat lower from the 1 mg samples than from the 5 mg and 25 mg samples in the 1st experiment (Figure 2A). Apart from this, however, the sequencing yield did not differ between the sample groups within any of the experiments. Thus, it is possible to compare the number of SVs detected in the samples without rarefying the data. The number of SVs in the 1st experiment was not significantly different between the 250 mg, 125 mg, and 25 mg samples, but decreased significantly in the 5 mg and even further among the 1 mg samples (Figure 2B). An opposite trend was apparent in the Good's coverage index (Table S1). Similarly, significantly lower numbers of SVs were detected in the single aggregates than in the 250 mg soil samples of the 3rd experiment.

Principal component analysis (PCA) from the sequencing results from the 1st experiment arranged all 250 mg and 125 mg samples, and most 25 mg samples into a single, tight group, indicating high similarity in their prokaryotic community structures (Figure 3A). In contrast, samples from the 5 mg and more so from the 1 mg categories, showed higher heterogeneity in community structure. Similarly, PCA indicated heterogeneity between individual aggregates that was not seen among the 250 mg samples in the 3rd experiment (Figure 3B).

The abundant SVs ($\geq$ 0.1% relative abundance in at least one of the samples) in the 250 mg, 125 mg, and 25 mg samples from the 1st experiment were almost all detectable in each individual sample, showing that the composition of the soil prokaryotic community appears very uniform when investigated at such a coarse spatial resolution (Figure 4). In contrast, 172 of the abundant SVs detected in the 1 mg samples were unique to just one or two of these samples. Among these

SVs, representatives of *Planctomycetes*, *Proteobacteria*, and *Acidobacteria* were especially

numerous, while *Thaumarchaeota* and *Actinobacteria* were dominant among the SVs present in

all samples. The 5 mg samples represented a level of spatial resolution at which some

heterogeneity in the prevalence of the abundant SVs was already apparent with 22 of them

detectable in two or only in a single sample.

In total, 5 620 SVs were detected in the eight 1 mg soil samples of the 1st experiment

(Supplementary file 2). Of these, 4 764 (85%) were also present in at least half of the 250 mg

samples. The remaining 856 SVs had low relative abundance in the 1 mg samples with only 59

reaching more than 0.1% relative abundance in any of them. The 5 mg samples together

contained 8 010 SVs, of which 6 443 (80%) were also detectable in at least half of the 250 mg

samples. Of the remaining 1 567 SVs, only 26 reached more than 0.1% relative abundance in any

of the 5 mg samples.

The qPCR results did not confirm our hypothesis that increasing spatial resolution would reveal

heterogeneity in microbial abundance. Bacterial and archaeal 16S rRNA gene, and fungal ITS copy

numbers did not show a larger variation among the 1 and 5 mg samples than between the 250

mg samples from the 1st experiment (Figure 1). Similarly, in the 3rd experiment, bacterial

abundance did not vary more in the single aggregates than in the 250 mg samples (Figure S1).

Impact of stochastic effects and inconsistent performance of the methods

The soil homogenate samples from the 2nd experiment served to test the influence of stochastic

effects and sub-optimal performance of the DNA extraction and PCR when extracting small

amounts of soil. The estimated bacterial abundance in a gram of soil based on the qPCR results

was in general higher in the samples from the 2nd experiment but followed the same pattern as

in the samples from the 1st experiment with no differences between the 1 mg, 5 mg, and 250 mg

samples but significantly higher values in the 25 mg samples (Figure S3). The small soil

homogenate samples did not show the degree of heterogeneity in prokaryotic community

structure we observed among the small soil samples of the 1st experiment. The abundant SVs ($\geq$

0.1% relative abundance in at least one sample) in the 25 mg soil homogenate samples were all

detectable in at least five of the eight replicates. Out of the 354 abundant SVs in the 5 mg soil

homogenate samples, one was present in only three of the samples but the others were

detectable in at least six. The 1 mg soil homogenate samples harbored 446 abundant SVs. None

of them was unique to a single sample and 442 were present in five or more of the eight samples.

The Aitchison distances of community structure were much higher among the 5 mg, and

especially among the 1 mg soil samples of the 1st experiment compared to the distances between

the 250 mg soil samples (Figure 5). In contrast, the distances between 5 mg or 25 mg soil

homogenate samples of the 2nd experiment were very similar to the distances among the 250 mg

soil samples indicating no difference in the heterogeneity of prokaryotic community structure

between these sample groups. The distances between the 1 mg soil homogenate samples were

only slightly increased.

Bacterial and archaeal co-occurrence patterns in 250 mg soil samples and aggregates

Networks of prokaryotic co-occurrence were constructed using the 272 SVs that reached $\geq$0.2%

relative abundance in at least one of the samples of the 3rd experiment. No network was obtained

from the 250 mg samples unless the removal of unstable edges and the correction of the p-values

for multiple testing were skipped. The resulting network has 78 edges between 35 nodes (Figure

6A). Thus, this spatial resolution revealed only a small number of putative associations many of which are likely false discoveries. In contrast, a network of 137 edges and 67 nodes (with the removal of unstable edges and control of the false discovery rate) was obtained from the individual soil aggregates (Figure 6B). A total of 54 of the nodes are part of a connected component in which there are three nodes with high betweenness centrality: SV15 (*Verrucomicrobia*, candidatus *Udaeobacter*), SV31 (*Actinobacteria*), and SV36 (*Acidobacteria* subgroup 6). Their relative abundance in the aggregates were 0.46 ± 0.16%, 0.33 ± 0.13%, and 0.30 ± 0.11%, respectively. These SVs potentially serve a keystone function by connecting two clusters in the network. One of the clusters contains several SVs of *Thaumarchaeota*, *Verrucomicrobia*, and *Actinobacteria*. The other is dominated by *Acidobacteria* subgroup 6. The hub of the latter cluster is SV58 (*Acidobacteria* subgroup 6) with a relative abundance of 0.23 ± 0.16% that has the highest degree in the network being connected to 19 nodes. SV399 (*Chloroflexi*) (0.06 ± 0.05%) and SV123 (*Acidobacteria* subgroup 6) (0.13 ± 0.06%) are linked with negative associations to several members of this cluster.

Discussion

The large disparity between the scale in which the soil microbiota is usually studied with molecular methods (0.25 to 1 g of soil) and the distance over which microbial interactions occur, impede the detection of interacting partners (Nunan, 2017). In order to gain information on the soil microbial diversity at an increased spatial resolution that considers soil structure, in this study we reduced the amount of soil used for DNA extraction from 250 to 1 mg and also extracted individual soil aggregates. Bacterial and archaeal DNA was recovered with not a significantly

different efficiency from the 250 mg and the 5 mg and 1 mg samples as shown by the qPCR results.

This was not true for fungal DNA. Either the DNA-extraction kit was not efficient in isolating fungal

DNA from samples below 25 mg, or fungi may preferentially colonize larger soil aggregates. Our

results show that the DNA extraction kit was most efficient in recovering bacterial and archaeal

DNA from 25 to 125 mg soil, although the variation in the yield of 16S rRNA gene copies was large

among these samples. Since this increased variation was apparent among the 25 mg soil

homogenate samples of the 2nd experiment as well, it is not an indication of an uneven

distribution of bacterial cells at the scale of 25 mg samples but must be due to this particular DNA

extraction method not working with consistent efficiency with this amount of soil.

As a consequence of sampling small amounts of soil, the DNA extracts had low template

concentrations for the subsequent PCR analyses. Thereby we had to anticipate a high risk of

contamination affecting the results (Weiss et al., 2014). In fact, quantifiable amounts of *Bacteria*

and *Fungi*, but not *Archaea*, were detected in the control samples without soil. However, they

reached no more than 4% of the number of bacterial rRNA genes copies in the smallest soil

samples, and thus, the influence of contamination on our results is negligible. The bacterial

community found in the control samples was clearly distinct from the soil communities suggesting

that the contamination originated from the reagents of the DNA extraction and sequencing library

preparation rather than cross-contamination between samples (Glassing, Dowd, Galandiuk,

Davis, & Chiodini, 2016; Salter et al., 2014). Another concern of working with small samples is that

molecular methods applied to such low amounts of template may perform inconsistently leading

to artificial variation in the results. The 2nd experiment showed that the DNA extraction, PCR, and

sequencing did not artificially generate more variation in the results from 5 mg samples than the

variation present among the 250 mg samples and the 1 mg samples showed only a slightly higher

variation. Therefore, the large heterogeneity in prokaryotic community composition and

structure among the 1 mg and 5 mg soil samples from the 1st experiment was not caused by

stochastic effects or PCR bias.

The samples of 25 mg up to 250 mg of soil were close to identical in prokaryotic community

composition, thus they provide a good representation of the overall prokaryotic diversity of our

soil. This is also indicated by the fact that increasing the amount of soil extracted up to 25 mg

increased the number of SVs detected in the samples, but larger soil samples did not yield more

SVs. Thus, the 25 mg samples already had a good coverage of the total prokaryotic community.

In contrast, the 1 mg and 5 mg samples and single aggregates were heterogeneous in community

structure. We found that while small soil samples could recover some SVs not necessarily

detected with the conventionally used 250 mg samples; these SVs were typically low in

abundance. Very few exceeded the relative abundance threshold we applied to control the

sparsity of the data in our analysis of community structure. Therefore, the large heterogeneity of

prokaryotic community structure we observed among the small samples was not because they

would have enabled the detection of more SVs. Instead, it appears that they contained different

subsets of the total community present in the 25 to 250 mg samples. This could be explained by

the fact that the smaller samples contain fewer micro-habitats, each of which harbors a local

community of fewer species (Leibold et al., 2004). Interestingly, the 1 mg and 5 mg samples did

not significantly differ in the abundance of *Bacteria, Archaea*, and *Fungi* compared to the 250 mg

samples. Similarly, the variation in bacterial abundance found with the individual aggregates was

not different from the 250 mg samples. Microbial abundance in soil has a patchy distribution at

the scale of a few micrometers (Nunan, Wu, Young, Crawford, & Ritz, 2003) but, for the soil of

this study, apparently not at the scale of macroaggregates or 1 to 5 mg samples.

Network analyses based on microbial co-occurrence have been applied to soil samples as large as

10 grams (Khan et al., 2019) and is typically used with 250 mg – 1 g samples (Barberan et al., 2012;

Karimi et al., 2020). In this study, however, we could not detect stable and significant associations

between SVs from 35 samples of 250 mg soil. These samples were taken from the same well-

mixed batch of soil and were very similar in their prokaryotic community composition. It is likely,

that each 250 mg sample gave a good representation of the overall prokaryotic diversity in our

soil, in which case, the variation of the relative abundance of SVs in these samples were mostly

random. Random patterns should not yield stable and significant associations in network analysis.

The value of utilizing much smaller samples is shown by our result that the 1 and 5 mg samples

contained subsets of the total soil microbial diversity captured by the 250 mg samples, and with

increasing spatial resolution the heterogeneity in the bacterial and archaeal community structure

increased among the samples. This results in detectable co-occurrence patterns. Furthermore,

the smaller spatial scale increases the likelihood that the observed co-occurrences actually

indicate interactions (Cordero & Datta, 2016).

From 37 soil aggregates, we obtained a complex network of bacterial and archaeal co-occurrence

that contained two clusters, one with several *Thaumarchaeota*, *Verrucomicrobia* and

Actinobacteria SVs, the other mainly with *Acidobacteria* subgroup 6 SVs. Three SVs, which could

represent keystone taxa, were found to connect these clusters. If these putative keystone SVs are

abundant in an aggregate, we can expect that members of both clusters are present there. The

two major clusters present in the network may provide complementary functions in the soil

ecosystem. There are indications that *Acidobacteria* subgroup 6, dominating one of the clusters,

prefer agricultural soils with low nitrogen input where it could be involved in the slower turnover

of soil organic carbon (SOC) originating from microbial necromass or plant material (Hester et al.,

2018; D. D. Li et al., 2018; Navarrete et al., 2013). The soil of this study originated from a long-

term nitrogen-depleted agricultural soil, thus supporting the preference to low nitrogen

concentrations and SOC turnover. The other cluster included several abundant SVs from phylum

Thaumarchaeota, which is known to be a strong contributor to ammonium oxidation in

agricultural soils (Leininger et al., 2006). Compared to ammonium-oxidizing bacteria,

Thaumarcheaota are thought to be adapted to lower nitrogen concentrations (Pester, Schleper,

& Wagner, 2011), thus the nitrogen-depleted soil of this study is likely a favorable environment

for them. The two clusters in our aggregate co-occurrence network could represent two distinct

types of metabolism adapted to a nitrogen depleted soil: a chemoorganotroph that oxidizes SOC,

and a chemolithotroph that oxidizes ammonia produced for example by ammonification from

crop residues. The presence of the less abundant *Verrucomicrobia* and *Actinobacteria* SVs within

the *Thaumarchaota* dominated cluster is possibly linked to an oligotrophic lifestyle (Bergmann et

al., 2011; Fierer, Bradford, & Jackson, 2007), but considering the limited information that 16S

rRNA gene analyses can provide for these phyla, this remains yet only a hypothesis. Shotgun

sequencing and metagenomic analysis of DNA extracted from individual soil aggregates could

shed more light on the nature of the associations we detected in the co-occurrence network. In

general, such aggregate-level analyses of the soil microbiota, which we call "aggregatomics",

could inspire new ways of linking structure to function in soil microbial communities.

While the spatial scale that we reached in this study is not yet fine enough to reveal most

microbial interactions as they may occur in microaggregates (Raynaud & Nunan, 2014), it should

be able to support the development of hypotheses and experiments to understand the patterns

and processes shaping the assembly of soil microbial communities and modelling their behavior

(Faust & Raes, 2012; Tecon & Or, 2017). Developing DNA extraction protocols from even smaller

soils samples, approaching the microaggregate-level, should be a way forward to fuel soil

aggregate-oriented   research   ("Aggregatomics";   https://www.thuenen.de/en/bd/fields-of-

activity/feld-und-laborstudien/microbiology-and-molecular-ecology/soil-aggregatomics/) for unveiling

hidden patterns of functions and ecological interactions.

Acknowledgements

We   thank   Ines   Merbach,   Department   Biozönoseforschung,   Helmholtz-Zentrum   für

Umweltforschung – UFZ, Bad Lauchstädt, Germany, for providing the soil of this study. We thank

Naomie Oßwald and Jerome Rischke who supported us at the early stages of this work as bachelor

students, and Karin Trescher for excellent technical assistance. The study was financially

supported by the German Research Foundation (DFG; Grant number TE 383/10-1) in context of

the Priority Programme "Rhizosphere spatiotemporal organization (SPP2089).

Conflict of interest

The authors declare no conflict of interest.

References

References

475 **References**

476 Assenov, Y., Ramirez, F., Schelhorn, S. E., Lengauer, T., & Albrecht, M. (2008). Computing
477     topological parameters of biological networks. *Bioinformatics, 24*(2), 282-284.
478     doi:10.1093/bioinformatics/btm554

479 Bach, E. M., Williams, R. J., Hargreaves, S. K., Yang, F., & Hofmockel, K. S. (2018). Greatest soil
480     microbial diversity found in micro-habitats. *Soil Biology & Biochemistry, 118*, 217-226.
481     doi:10.1016/j.soilbio.2017.12.018

482 Bailey, V. L., Fansler, S. J., Stegen, J. C., & McCue, L. A. (2013). Linking microbial community
483     structure to beta-glucosidic function in soil aggregates. *ISME Journal, 7*(10), 2044-2053.
484     doi:10.1038/ismej.2013.87

485 Bailey, V. L., McCue, L. A., Fansler, S. J., Boyanov, M. I., DeCarlo, F., Kemner, K. M., & Konopka, A.
486     (2013). Micrometer-scale physical structure and microbial composition of soil
487     macroaggregates. *Soil Biology & Biochemistry, 65*, 60-68.
488     doi:10.1016/j.soilbio.2013.02.005

489 Banerjee, S., Kirkby, C. A., Schmutter, D., Bissett, A., Kirkegaard, J. A., & Richardson, A. E. (2016).
490     Network analysis reveals functional redundancy and keystone taxa amongst bacterial and
491     fungal communities during organic matter decomposition in an arable soil. *Soil Biology &*
492     *Biochemistry, 97*, 188-198. doi:10.1016/j.soilbio.2016.03.017

493 Barberan, A., Bates, S. T., Casamayor, E. O., & Fierer, N. (2012). Using network analysis to explore
494     co-occurrence patterns in soil microbial communities. *ISME Journal, 6*(2), 343-351.
495     doi:10.1038/ismej.2011.119

496 Bergmann, G. T., Bates, S. T., Eilers, K. G., Lauber, C. L., Caporaso, J. G., Walters, W. A., . . . Fierer,
497     N. (2011). The under-recognized dominance of Verrucomicrobia in soil bacterial
498     communities. *Soil Biology & Biochemistry, 43*(7), 1450-1455.
499     doi:10.1016/j.soilbio.2011.03.012

500 Blair, N., Faulkner, R. D., Till, A. R., Körschens, M., & Schulz, E. (2006). Long-term management
501     impacts on soil C, N and physical fertility - Part II: Bad Lauchstädt static and extreme FYM
502     experiments. *Soil & Tillage Research, 91*(1-2), 39-47. doi:10.1016/j.still.2005.11.001

503 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016).
504     DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods,*
505     *13*(7), 581-+. doi:10.1038/nmeth.3869

506 Constancias, F., Prevost-Boure, N. C., Terrat, S., Aussems, S., Nowak, V., Guillemin, J. P., . . .
507     Ranjard, L. (2014). Microscale evidence for a high decrease of soil bacterial density and
508     diversity by cropping. *Agronomy for Sustainable Development, 34*(4), 831-840.
509     doi:10.1007/s13593-013-0204-3

510 Cordero, O. X., & Datta, M. S. (2016). Microbial interactions and community assembly at
511     microscales. *Current Opinion in Microbiology, 31*, 227-234. doi:10.1016/j.mib.2016.03.015

Davinic, M., Fultz, L. M., Acosta-Martinez, V., Calderon, F. J., Cox, S. B., Dowd, S. E., . . . Moore-Kucera, J. (2012). Pyrosequencing and mid-infrared spectroscopy reveal distinct aggregate stratification of soil bacterial communities and organic matter composition. *Soil Biology & Biochemistry, 46*, 63-72. doi:10.1016/j.soilbio.2011.11.012

Direito, S. O. L., Zaura, E., Little, M., Ehrenfreund, P., & Roling, W. F. M. (2014). Systematic evaluation of bias in microbial community profiles induced by whole genome amplification. *Environmental Microbiology, 16*(3), 643-657. doi:10.1111/1462-2920.12365

Faust, K., & Raes, J. (2012). Microbial interactions: from networks to models. *Nature Reviews Microbiology, 10*(8), 538-550. doi:10.1038/nrmicro2832

Faust, K., & Raes, J. (2016). CoNet app: inference of biological association networks using Cytoscape [version 2; peer review: 2 approved]. *5*(1519). doi:10.12688/f1000research.9050.2

Faust, K., Sathirapongsasuti, J. F., Izard, J., Segata, N., Gevers, D., Raes, J., & Huttenhower, C. (2012). Microbial co-occurrence relationships in the human microbiome. *PLoS Computational Biology, 8*(7), 17. doi:10.1371/journal.pcbi.1002606

Fierer, N., Bradford, M. A., & Jackson, R. B. (2007). Toward an ecological classification of soil bacteria. *Ecology, 88*(6), 1354-1364. doi:10.1890/05-1839

Fierer, N., & Jackson, R. B. (2006). The diversity and biogeography of soil bacterial communities. *Proceedings of the National Academy of Sciences of the United States of America, 103*(3), 626-631. doi:10.1073/pnas.0507535103

Fox, A., Ikoyi, I., Torres-Sallan, G., Lanigan, G., Schmalenberger, A., Wakelin, S., & Creamer, R. (2018). The influence of aggregate size fraction and horizon position on microbial community composition. *Applied Soil Ecology, 127*, 19-29. doi:10.1016/j.apsoil.2018.02.023

Glassing, A., Dowd, S. E., Galandiuk, S., Davis, B., & Chiodini, R. J. (2016). Inherent bacterial DNA contamination of extraction and sequencing reagents may affect interpretation of microbiota in low bacterial biomass samples. *Gut Pathogens, 8*. doi:10.1186/s13099-016-0103-7

Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V., & Egozcue, J. J. (2017). Microbiome datasets are compositional: And this is not optional. *Frontiers in Microbiology, 8*, 6. doi:10.3389/fmicb.2017.02224

Gloor, G. B., Wu, J. R., Pawlowsky-Glahn, V., & Egozcue, J. J. (2016). It's all relative: analyzing microbiome data as compositions. *Annals of Epidemiology, 26*(5), 322-329. doi:10.1016/j.annepidem.2016.03.003

Green, J. L., Bohannan, B. J. M., & Whitaker, R. J. (2008). Microbial biogeography: From taxonomy to traits. *Science, 320*(5879), 1039-1043. doi:10.1126/science.1153475

549   Griffiths, R. I., Thomson, B. C., James, P., Bell, T., Bailey, M., & Whiteley, A. S. (2011). The bacterial
550   biogeography of British soils. *Environmental Microbiology, 13*(6), 1642-1654.
551   doi:10.1111/j.1462-2920.2011.02480.x

552   Hemkemeyer, M., Christensen, B. T., Martens, R., & Tebbe, C. C. (2015). Soil particle size fractions
553   harbour distinct microbial communities and differ in potential for microbial mineralisation
554   of organic pollutants. *Soil Biology & Biochemistry, 90*, 255-265.
555   doi:10.1016/j.soilbio.2015.08.018

556   Hemkemeyer, M., Christensen, B. T., Tebbe, C. C., & Hartmann, M. (2019). Taxon-specific fungal
557   preference for distinct soil particle size fractions. *European Journal of Soil Biology, 94*.
558   doi:10.1016/j.ejsobi.2019.103103

559   Hemkemeyer, M., Dohrmann, A. B., Christensen, B. T., & Tebbe, C. C. (2018). Bacterial preferences
560   for specific soil particle size fractions revealed by community analyses. *Frontiers in*
561   *Microbiology, 9*. doi:10.3389/fmicb.2018.00149

562   Hester, E. R., Harpenslager, S. F., van Diggelen, J. M. H., Lamers, L. L., Jetten, M. S. M., Luke, C., . .
563   . Welte, C. U. (2018). Linking nitrogen load to the structure and function of wetland soil
564   and rhizosphere microbial communities. *Msystems, 3*(1). doi:10.1128/mSystems.00214-
565   17

566   Ivanova, E. A., Kutovaya, O. V., Tkhakakhova, A. K., Chernov, T. I., Pershina, E. V., Markina, L. G., .
567   . . Kogut, B. M. (2015). The structure of microbial community in aggregates of a typical
568   chernozem aggregates under contrasting variants of its agricultural use. *Eurasian Soil*
569   *Science, 48*(11), 1242-1256. doi:10.1134/s1064229315110083

570   Karimi, B., Terrat, S., Dequiedt, S., Saby, N. P. A., Horriguel, W., Lelievre, M., . . . Ranjard, L. (2018).
571   Biogeography of soil bacteria and archaea across France. *Science Advances, 4*(7).
572   doi:10.1126/sciadv.aat1808

573   Karimi, B., Villerd, J., Dequiedt, S., Terrat, S., Boure, N. C. P., Djemiel, C., . . . Ranjard, L. (2020).
574   Biogeography of soil microbial habitats across France. *Global Ecology and Biogeography,*
575   *29*(8), 1399-1411. doi:10.1111/geb.13118

576   Khan, M. A. W., Bohannan, B. J. M., Nusslein, K., Tiedje, J. M., Tringe, S. G., Parlade, E., . . .
577   Rodrigues, J. L. M. (2019). Deforestation impacts network co-occurrence patterns of
578   microbial communities in Amazon soils. *FEMS Microbiology Ecology, 95*(2).
579   doi:10.1093/femsec/fiy230

580   Kim, H., Nunan, N., Dechesne, A., Juarez, S., & Grundmann, G. (2015). The spatial distribution of
581   exoenzyme activities across the soil micro-landscape, as measured in micro- and macro-
582   aggregates, and ecosystem processes. *Soil Biology & Biochemistry, 91*, 258-267.
583   doi:10.1016/j.soilbio.2015.08.042

584   Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development
585   of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence
586   data on the MiSeq Illumina sequencing platform. *Applied and Environmental*
587   *Microbiology, 79*(17), 5112-5120. doi:10.1128/aem.01043-13

Kravchenko, A. N., Negassa, W. C., Guber, A. K., Hildebrandt, B., Marsh, T. L., & Rivers, M. L. (2014).
Intra-aggregate pore structure influences phylogenetic composition of bacterial
community in macroaggregates. *Soil Science Society of America Journal, 78*(6), 1924-1939.
doi:10.2136/sssaj2014.07.0308

Leibold, M. A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., . . .
Gonzalez, A. (2004). The metacommunity concept: a framework for multi-scale
community ecology. *Ecology Letters, 7*(7), 601-613. doi:10.1111/j.1461-
0248.2004.00608.x

Leininger, S., Urich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G. W., . . . Schleper, C. (2006). Archaea
predominate among ammonia-oxidizing prokaryotes in soils. *Nature, 442*(7104), 806-809.
doi:10.1038/nature04983

Li, D. D., Chen, L., Xu, J. S., Ma, L., Olk, D. C., Zhao, B. Z., . . . Xin, X. L. (2018). Chemical nature of
soil organic carbon under different long-term fertilization regimes is coupled with changes
in the bacterial community composition in a Calcaric Fluvisol. *Biology and Fertility of Soils,*
54(8), 999-1012. doi:10.1007/s00374-018-1319-0

Li, F., Chen, L., Zhang, J. B., Yin, J., & Huang, S. M. (2017). Bacterial community structure after
long-term organic and inorganic fertilization reveals important associations between soil
nutrients and specific taxa involved in nutrient transformations. *Frontiers in Microbiology,*
8. doi:10.3389/fmicb.2017.00187

Lombard, N., Prestat, E., van Elsas, J. D., & Simonet, P. (2011). Soil-specific limitations for access
and analysis of soil microbial communities by metagenomics. *FEMS Microbiology Ecology,*
78(1), 31-49. doi:10.1111/j.1574-6941.2011.01140.x

Ludwig, B., Geisseler, D., Michel, K., Joergensen, R. G., Schulz, E., Merbach, I., . . . Liu, X. (2011).
Effects of fertilization and soil management on crop yields and carbon stabilization in soils.
A review. *Agronomy for Sustainable Development, 31*(2), 361-372.
doi:10.1051/agro/2010030

Merbach, I., & Schulz, E. (2013). Long-term fertilization effects on crop yields, soil fertility and
sustainability in the Static Fertilization Experiment Bad Lauchstadt under climatic
conditions 2001-2010. *Archives of Agronomy and Soil Science, 59*(8), 1041-1058.
doi:10.1080/03650340.2012.702895

Miller, D. N., Bryant, J. E., Madsen, E. L., & Ghiorse, W. C. (1999). Evaluation and optimization of
DNA extraction and purification procedures for soil and sediment samples. *Applied and*
Environmental Microbiology, 65(11), 4715-4724.

Navarrete, A. A., Kuramae, E. E., de Hollander, M., Pijl, A. S., van Veen, J. A., & Tsai, S. M. (2013).
Acidobacterial community responses to agricultural management of soybean in Amazon
forest soils. *Fems Microbiology Ecology, 83*(3), 607-621. doi:10.1111/1574-6941.12018

Nunan, N. (2017). The microbial habitat in soil: Scale, heterogeneity and functional consequences.
Journal of Plant Nutrition and Soil Science, 180(4), 425-429. doi:10.1002/jpln.201700184

Nunan, N., Wu, K. J., Young, I. M., Crawford, J. W., & Ritz, K. (2003). Spatial distribution of bacterial
communities and their relationships with the micro-architecture of soil. *Fems*
Microbiology Ecology, 44(2), 203-215. doi:10.1016/s0168-6496(03)00027-8

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., . . . Wagner, H.
(2018). vegan: Community Ecology Package. R package version 2.5-2.

Palarea-Albaladejo, J., & Martin-Fernandez, J. A. (2015). zCompositions - R Package for
multivariate imputation of left-censored data under a compositional approach.
Chemometrics and Intelligent Laboratory Systems, 143, 85-96.
doi:10.1016/j.chemolab.2015.02.019

Pester, M., Schleper, C., & Wagner, M. (2011). The Thaumarchaeota: an emerging view of their
phylogeny and ecophysiology. *Current Opinion in Microbiology, 14*(3), 300-306.
doi:10.1016/j.mib.2011.04.007

Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W. G., Peplies, J., & Glockner, F. O. (2007).
SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA
sequence data compatible with ARB. *Nucleic Acids Research, 35*(21), 7188-7196.
doi:10.1093/nar/gkm864

Ranjard, L., Nazaret, S., Gourbiere, F., Thioulouse, J., Linet, P., & Richaume, A. (2000). A soil
microscale study to reveal the heterogeneity of Hg(II) impact on indigenous bacteria by
quantification of adapted phenotypes and analysis of community DNA fingerprints. *FEMS*
Microbiology Ecology, 31(2), 107-115. doi:10.1111/j.1574-6941.2000.tb00676.x

Ranjard, L., Poly, F., Combrisson, J., Richaume, A., Gourbiere, F., Thioulouse, J., & Nazaret, S.
(2000). Heterogeneous cell density and genetic structure of bacterial pools associated
with various soil microenvironments as determined by enumeration and DNA
fingerprinting approach (RISA). *Microbial Ecology, 39*(4), 263-272.

Raynaud, X., & Nunan, N. (2014). Spatial Ecology of Bacteria at the Microscale in Soil. *PLoS One,*
9(1). doi:10.1371/journal.pone.0087217

Rillig, M. C., Müller, L. A. H., & Lehmann, A. (2017). Soil aggregates as massively concurrent
evolutionary incubators. *ISEME Journal, 11*(9), 1943-1948. doi:10.1038/ismej.2017.56

Roesch, L. F., Fulthorpe, R. R., Riva, A., Casella, G., Hadwin, A. K. M., Kent, A. D., . . . Triplett, E. W.
(2007). Pyrosequencing enumerates and contrasts soil microbial diversity. *ISME Journal,*
1(4), 283-290. doi:10.1038/ismej.2007.53

Salter, S. J., Cox, M. J., Turek, E. M., Calus, S. T., Cookson, W. O., Moffatt, M. F., . . . Walker, A. W.
(2014). Reagent and laboratory contamination can critically impact sequence-based
microbiome analyses. *BMC Biology, 12*. doi:10.1186/s12915-014-0087-z

Six, J., Elliott, E. T., & Paustian, K. (2000). Soil macroaggregate turnover and microaggregate
formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biology &*
Biochemistry, 32(14), 2099-2103. doi:Doi 10.1016/S0038-0717(00)00179-6

Smith, P., Cotrufo, M. E., Rumpel, C., Paustian, K., Kuikman, P. J., Elliott, J. A., . . . Scholes, M. C.
(2015). Biogeochemical cycles and biodiversity as key drivers of ecosystem services
provided by soils. *Soil, 1*(2), 665-685. doi:10.5194/soil-1-665-2015

Szoboszlay, M., Dohrmann, A. B., Poeplau, C., Don, A., & Tebbe, C. C. (2017). Impact of land-use
change and soil organic carbon quality on microbial diversity in soils across Europe. *FEMS*
Microbiology Ecology, 93(12). doi:10.1093/femsec/fix146

Tecon, R., & Or, D. (2017). Biophysical processes supporting the diversity of microbial life in soil.
FEMS Microbiology Reviews, 41(5), 599-623. doi:10.1093/femsre/fux039

Thakur, M. P., Phillips, H. R. P., Brose, U., De Vries, F. T., Lavelle, P., Loreau, M., . . . Cameron, E. K.
(2020). Towards an integrative understanding of soil biodiversity. *Biological Reviews,*
95(2), 350-364. doi:10.1111/brv.12567

Vos, M., Wolf, A. B., Jennings, S. J., & Kowalchuk, G. A. (2013). Micro-scale determinants of
bacterial diversity in soil. *FEMS Microbiology Reviews, 37*(6), 936-954. doi:10.1111/1574-
6976.12023

Walters, W., Hyde, E. R., Berg-Lyons, D., Ackermann, G., Humphrey, G., Parada, A., . . . Knight, R.
(2016). Improved bacterial 16S rRNA gene (V4 and V4-5) and fungal internal transcribed
spacer marker gene primers for microbial community surveys. *Msystems, 1*(1), 10.
doi:10.1128/mSystems.00009-15

Wang, Y., Gao, Z. M., Xu, Y., Li, G. Y., He, L. S., & Qian, P. Y. (2016). An evaluation of multiple
annealing and looping based genome amplification using a synthetic bacterial community.
Acta Oceanologica Sinica, 35(2), 131-136. doi:10.1007/s13131-015-0781-x

Weiss, S., Amir, A., Hyde, E. R., Metcalf, J. L., Song, S. J., & Knight, R. (2014). Tracking down the
sources of experimental contamination in microbiome studies. *Genome Biology, 15*(12).
doi:10.1186/s13059-014-0564-2

Weiss, S., Van Treuren, W., Lozupone, C., Faust, K., Friedman, J., Deng, Y., . . . Knight, R. (2016).
Correlation detection strategies in microbial data sets vary widely in sensitivity and
precision. *ISME Journal, 10*(7), 1669-1681. doi:10.1038/ismej.2015.235

Wilpiszeski, R. L., Aufrecht, J. A., Retterer, S. T., Sullivan, M. B., Graham, D. E., Pierce, E. M., . . .
Elias, D. A. (2019). Soil aggregate microbial communities: Towards understanding
microbiome interactions at biologically relevant scales. *Applied and Environmental*
Microbiology, 85(14). doi:10.1128/aem.00324-19

Young, J. M., Rawlence, N. J., Weyrich, L. S., & Cooper, A. (2014). Limitations and
recommendations for successful DNA extraction from forensic soil samples: A review.
Science & Justice, 54(3), 238-244. doi:10.1016/j.scijus.2014.02.006

Data accessibility statement

Sequencing data is available at the European Nucleotide Archive under the accession numbers

PRJEB36881 (1st experiment), PRJEB36883 (2nd experiment), and PRJEB36887 (3rd experiment).

The qPCR data is in Supplementary file 1.

Author contributions

Both authors contributed to the design of the experiments, analysis of the results, and writing the

manuscript.

Tables and figures

Table 1 Number of samples and soil weights within each sample category

Soil weight class or sample type	Sample weight [mg] ± SD	Number of soil samples
1st Experiment		
250 mg	251 ± 1	8
125 mg	125 ± 1	8
25 mg	25.1 ± 0.4	8
5 mg	4.9 ± 0.2	8
1 mg	1.1 ± 0.2	8
Control, no soil		6
2nd Experiment		
250 mg	252 ± 3	8
25 mg soil homogenate		8
5 mg soil homogenate		8
1 mg soil homogenate		8

Control, no soil		6
3rd Experiment		
250 mg	251 ± 0	35
Soil aggregate	5.3 ± 1.6	37
Control		9

Figure 1 Estimates of (A) bacterial, (B) archaeal, and (C) fungal abundance in a gram of soil based

on qPCR from the samples from the 1st experiment (gene copy numbers per g of soil wet weight).

One of the 250 mg soil samples was an outlier in the fungal ITS qPCR results and is not included

in the plot. Sample groups not labelled with the same letter were significantly different in Tukey's

HSD tests. Thick lines indicate the median values, the upper and lower hinges the 75th and 25th

percentile, whiskers extend to the data extremes.

Figure 2 Number of (A) sequences and (B) sequence variants (SVs) in the samples after the removal of potentially contaminant SVs. Thick lines indicate the median values, the upper and lower hinges the 75th and 25th percentile, whiskers extend to the data extremes. Letters indicate significant differences between sample groups of the 1st experiment according to Tukey's HSD tests. * indicates significant difference based on Welch's t-test between the aggregate and the 250 mg soil samples of the 3rd experiment.

Figure 3 Principle component analyses (PCA) plots from the 16S rRNA gene sequencing data from

the 1st (A) and 3rd (B) experiments.

SVs detected in *N* samples

Figure 4 SVs arranged according to in how many of the 1 mg, 5 mg, 25 mg, 125 mg, or 250 mg samples from the 1ˢᵗ experiment they were detected. Only SVs that reached at least 0.1% relative abundance in any of the samples are included. Each node represents one SV colored based on its phylum-level classification and sized according to its average relative abundance across all samples excluding those in which it was not detected.

731   **Figure 5** Aitchison distances in the bacterial and archaeal community structure (16SrRNA gene

732   amplicons) within sample groups from the 1st and 2nd experiments. Thick lines indicate the median

733   values, the upper and lower hinges the 75th and 25th percentile, whiskers extend to the data

734   extremes. Sample groups from the same experiment not labelled with the same letter were

735   significantly different in Tukey's HSD tests.

A

B

Figure 6 Co-occurrence networks from the (A) 250 mg samples and (B) the single aggregates from

the 3rd experiment. The frames mark the two clusters discussed in the text. It should be noted

that in (A) unstable edges were *not* removed and the Benjamini-Hochberg correction for multiple

comparisons was not applied.