Hidden heterogeneity and co-occurrence networks of soil prokaryotic

communities revealed at the scale of individual soil aggregates

Running title: Microbiota at soil aggregate-level

Márton Szoboszlay and Christoph C. Tebbe[1]

Thünen Institut für Biodiversität, Bundesforschungsinstitut für Ländliche Räume, Wald und

Fischerei, Braunschweig, Germany

Corresponding author: christoph.tebbe@thuenen.de

Address of corresponding author:

Prof. Dr. Christoph Tebbe, Thünen Institut für Biodiversität, Bundesallee 65, D-38116

Braunschweig, Germany

<u>Declaration by the authors</u>:

Abstract

Sequencing PCR-amplified gene fragments from metagenomic DNA is a widely applied method for studying the diversity and dynamics of soil microbial communities. Typically DNA is extracted from 0.25 to 1 g of soil. These amounts, however, neglect the heterogeneity of soil present at the scale of soil aggregates; and thus, ignore a crucial scale for understanding the structure and functionality of soil microbial communities. Here we show with an arable soil the impact of reducing the amount of soil used for DNA extraction from 250 mg to approx. 1 mg in order to access spatial information on the prokaryotic community structure as indicated by 16S rRNA-gene amplicon analyses. Furthermore, we demonstrate that individual aggregates from the same soil differ in their prokaryotic communities. The analysis of 16S rRNA gene amplicon sequences from individual soil aggregates allowed us, in contrast to 250 mg soil samples, to construct a co-occurrence network that provides insight into the structure of microbial associations in the studied soil. Two dense clusters were apparent in the network, one dominated by *Thaumarchaeota*, *Verrucomicrobia*, and *Actinobacteria*, and the other by *Acidobacteria* subgroup 6, linked by three putative keystone taxa (candidatus *Udaeobacter*, Verrucomicrobia; a member of Actinobacteria and Acidobacteria, respectively).

Introduction

One of the major challenges in microbial ecology is to elucidate the principles, patterns, and interactions which lead to the formation of highly diverse microbial communities as found for example in soil [1, 2]. To gain a predictive understanding of soil microbial diversity, it is essential to consider spatial scale information linked to soil structure [3, 4].

Soil structure develops as primary particles of different sizes and mineral composition, i.e., clay, silt, and sand, interact with each other, and with organic material to build microaggregates and macroaggregates, both distinguished by their diameter, i.e., below or above 250 µm [5]. Most bacterial cells occur inside aggregates rather than on their surfaces [6]. Biogeochemical cycles, which are key ecosystem services driven by an interacting microbial community [7], are considered to mainly take place within aggregates [8]. Aggregates have been regarded as "massively concurrent evolutionary incubators" [9], or as "microbial villages" [8], that represent small communities separated by distance and physical barriers and connected only periodically, e.g. during wetting events.

DNA-based methods to assess microbial diversity typically start with extracting 250 mg to 1 g of soil [10]. This strategy has been useful to investigate the overall microbial diversity of soils [11, 12], its variation across geographical regions [13, 14], or its response to land use change at a continental scale [15]. However, in order to understand the processes and interactions taking place within soil microbial communities, it would be rewarding to increase the spatial resolution to individual aggregates and investigate microbial diversity in these spatial entities. This would make it possible to assess the micro-heterogeneity of soil microbial communities and increase the likelihood of detecting interacting microbial partners.

The potential impact of the heterogeneous soil constituents on structuring microbial

communities at a microscale has already been demonstrated with pooled soil primary particles,

where the majority of abundant bacterial and fungal taxa exhibited particular preferences for

clay, silt, or sand-sized fractions [16, 17]. Comparing pooled samples of micro- and

macroaggregates revealed that they also differ in microbial community composition [18-20],

diversity [21, 22] and their response to stress [23]. However, information on the heterogeneity

of microbial communities of individual aggregates within specific aggregate fractions is still

lacking. A major limitation of analyzing individual aggregates is the difficulty of obtaining a

sufficient quantity of nucleic acids from small amounts of soil for molecular methods. Attempts

made so far, therefore, either pooled several aggregates for DNA extraction [21, 22, 24],

sampled very large aggregates of 20 – 70 mg [25], or applied whole genome amplification [26].

These solutions, however, do not deliver data on individual aggregates, provide coarse spatial

resolution, or generate substantial bias in the results [27, 28], respectively. To our knowledge,

the only study that reported the bacterial community composition in smaller (2 mm) individual

aggregates without applying whole genome amplification utilized taxonomic microarrays; a

method of relatively low resolution, and focused solely on linking enzyme activity profiles with

community structure [29]. Furthermore, applying molecular methods to small samples that yield

very low amounts of nucleic acids require validation to prove the consistent performance of the

methods and rule out the possibility of contamination and stochastic effects influencing the

results.

The tremendous scientific potential that individual soil aggregate-based microbial community

analysis should have for characterizing the heterogeneity of soil microbial communities at a

biologically and ecologically more meaningful scale motivated us testing the following

hypotheses in this study:

1. Metagenomic DNA of sufficient quantity and quality for PCR-based analyses can be

extracted from soil samples as small as 1 mg

2. Increasing spatial resolution reveals heterogeneity in soil bacterial and archaeal

community structure and abundance

3. A higher heterogeneity seen among small soil samples is not a result of contamination or

sub-optimal performance of molecular methods.

4. Comparing individual aggregates from the same soil unveils patterns of microbial co-

occurrence within the soil microbial community not seen with the commonly used

250 mg soil sample size.

Materials and methods

Overview of the experiments

Three experiments were conducted in this study. In the 1st experiment, samples decreasing in

size from 250 mg to 1 mg taken from the same soil were subjected to DNA extraction. To

address the first two hypotheses, qPCR and high-throughput amplicon sequencing targeting the

16S rRNA gene were conducted to characterize the bacterial, archaeal, and fungal abundance

and the prokaryotic diversity in these samples. The 2nd experiment addressed the third

hypothesis by comparing 250 mg soil samples and aliquots of a homogenized soil slurry. The

volumes of the aliquots were chosen to contain the amount of DNA expected from 1 mg, 5 mg,

and 25 mg soil samples. Since all aliquots were taken from the same thoroughly homogenized

soil slurry, differences in prokaryotic community structure between these soil homogenate

samples must be results of contamination, stochastic effects, or sub-optimal performance of the

DNA extraction and PCR. In the 3rd experiment, DNA was extracted from individual aggregates

and 250 mg soil samples taken from the same soil to address the fourth hypothesis. All

experiments included several control samples to test for the presence of contamination.

Soil sampling and DNA extraction

The soil used in all experiments was a loam topsoil of a Haplic Chernozem (FAO classification)

from the Bad Lauchstädt experimental research station of the Helmholtz Centre for

Environmental Research in Germany (51°24'N 11°53'E). It was collected in December from a plot

under long term sugar beet – potato – winter wheat – barley rotation without fertilization. The

soil had pH 7.1 (in 0.01 M CaCl$_2$) and 17.7 g kg^{-1} organic C. It was sieved (2 mm) and stored at

4°C until use.

Before sampling, approximately 100 g soil was incubated at room T in the dark for 24 h. The soil

was then spread out in a sterile petri dish and samples were taken with sterilized spatulas

directly into the bead-beating tubes of the DNA extraction kit. Control samples were included in

all experiments. They received no soil but were handled together with and the same way as the

soil samples. DNA was extracted with the Quick-DNA Fecal/Soil Microbe Microprep Kit (Zymo

Research, Freiburg, Germany) including two 45 s bead beating cycles in an MP FastPrep-24 5G

Instrument (MP Biomedicals, Eschwege, Germany) at 6.5 m/s speed with a 300 s break in

between. The DNA extracts were eluted in 30 µl elution buffer. All work was done in a biosafety

cabinet decontaminated with UV light to minimize the chance of contamination. Measurement

of the DNA yield was attempted with Quant-iT PicoGreen dsDNA Assay Kit (Molecular Probes,

Life Technologies, Eugene, Or) but accurate quantification was not possible for many of the

small samples due to results close to the background fluorescence in the blank controls. In

preparation of this study, several DNA extractions methods were tested but were not found

suitable. This included the Fast DNA Spin kit for soil (MP Biomedicals, LLC, Illkrich, France) and

variations of the phenol-chloroform protocol [30].

In the 1^{st} experiment, five different size-groups of soil samples were collected: 250 mg, 125 mg,

25 mg, 5 mg, and 1 mg, respectively. Eight samples were taken from each size-group along with

six control samples. Sample weights from all experiments are listed in Table 1. In the 2^{nd}

experiment, ten samples of 250 mg soil and six control samples were taken. Eight of the soil

samples were processed normally in the DNA extraction, while for the other two, the DNA

extraction was interrupted after the centrifugation following the bead beating. By this point, the

samples had been turned into homogenized slurry by the bead beating, and the soil debris had

been separated from the supernatant that contained the metagenomic DNA. The supernatant

from the two samples were pooled and homogenized by vortexing. The mass of the resulting

suspension was 1 112 mg and it originated from 497 mg soil in total. Accordingly, a 55.9 mg

aliquot of this suspension contained the amount of DNA extractable from 25 mg soil, an 11.2 mg

aliquot the amount from 5 mg soil, and a 2.2 mg aliquot the amount from 1 mg soil. Eight

aliquots from each of these sizes, hereafter referred to as 25 mg, 5 mg, and 1 mg soil

homogenate samples, were taken and each mixed with 350 µl bashing bead buffer from the

DNA extraction kit to continue the DNA extraction. In the 3^{rd} experiment, 37 individual soil

aggregates, weighing 5.3 mg on average, were taken for DNA extraction along with 35 samples

of 250 mg soil and nine control samples.

Abundances of microbial groups assessed by qPCR

The abundance of *Bacteria*, *Archaea*, and *Fungi* were assessed by qPCR targeting the 16S rRNA

gene and the ITS region respectively following the protocol of Hemkemeyer *et al.* [31]. Archaeal

and fungal abundance was only investigated in the 1st experiment. Reactions were run in a Bio-

Rad CFX96 real-time PCR system in duplicates from different dilutions of the DNA extracts. In

case of the 250 mg and 125 mg soil samples, 50- and 100-fold dilutions were taken; from the 25

mg samples 10- and 20-fold dilutions; and from the 5 mg, 1 mg, individual aggregate, and

control samples undiluted DNA extracts and two-fold dilutions were used. PCR efficiencies were

95.9 − 104.6% for the bacterial 16S rRNA gene, 94.2 − 96.2% for the archaeal 16S rRNA gene,

and 83.3 − 84.3% for the fungal ITS with $R^2 \geq 0.995$ in all cases. Results were compared with

Tukey's HSD tests, or Welch's t-test in case of the data from the 3rd experiment. The analysis

was carried out in R 3.4.4 (www.r-project.org). One of the 250 mg samples from the 1st

experiment yielded a magnitude higher copy number in the fungal ITS qPCR assay than the

others. It was deemed to be an outlier and excluded from the analysis.

High-throughput sequencing of 16S rRNA gene amplicons and data processing

To characterize the bacterial and archaeal communities in the samples, DNA extracts were

subjected to high-throughput amplicon sequencing of the V4 region of the 16S rRNA gene

following the protocol of Kozich *et al.* [32] with primers updated to match the modified 515f

and 806r sequences according to Walters *et al.* [33]. In case of the small soil samples and

control samples, due to the low DNA yield, 10 µl DNA extract was used as template in the PCRs.

Paired-end sequencing was done on Illumina MiSeq instruments at StarSEQ (Mainz, Germany).

Samples from the same experiment were sequenced in the same run. All data is available at the

European Nucleotide Archive under the accession numbers PRJEB36881 (1st experiment),

PRJEB36883 (2nd experiment), and PRJEB36887 (3rd experiment).

The sequencing data from the three experiments were analyzed separately. Raw reads were

processed with the dada2 (version 1.6.0) pipeline [34] in R 3.4.4. Forward and reverse reads

were truncated at positions 240 and 90, respectively. Reads with any ambiguous bases were

discarded as well as forward reads with over two and reverse reads with over one expected

error. The data from the 2nd experiment had higher quality allowing the reverse reads to be

truncated at position 130 and keeping those with two or less expected errors. Error models

were constructed from 10^6 randomly selected reads. Sequence variants (SVs) were inferred

using the pool option. Forward and reverse SVs were merged trimming overhangs, and the

removeBimeraDenovo function was employed to detect chimeras. The SVs were classified

according to the SILVA reference version 132 [35] accepting only results with ≥70% bootstrap

support. SVs shorter than 220 nt or longer than 275 nt, or identified as chimeric, mitochondrial,

or chloroplast sequences, or not classified into *Bacteria* or *Archaea* were deleted. SVs with

≥0.1% relative abundance in any of the control samples of an experiment were regarded as

potentially contaminant and removed from the dataset of that experiment.

 Analysis of the sequencing results

Principal component analysis (PCA) plots were created in R using the rda function of the vegan

package version 2.5-2 [36]. To decrease the sparsity of the data, SVs not reaching 0.1% relative

abundance in any of the samples included in the PCA were removed. Zeroes were replaced with

the count zero multiplicative (CZM) method using the zCompositions package version 1.1.1 [37]

and centered log-ratio (CLR) transformation was applied to the data to correct for compositional

effects and differences in sequencing depth [38].

Aitchison distances between the samples were calculated as Euclidean distances in the CLR

transformed dataset [39]. SVs that didn't have at least 0.1% relative abundance in any of the

compared samples were removed and zeroes were replaced with the CZM method to allow CLR

transformation before calculating Euclidean distances with the 'vegdist' function of the vegan

package.

Plots illustrating the prevalence of abundant SVs among the soil samples were prepared in

Cytoscape 3.7.1 (www.cytoscape.org). CoNet 1.1.1 beta [40] in Cytoscape was used to construct

co-occurrence networks from the data from the 3rd experiment. To limit the number of parallel

significance tests and the sparsity of the data, only SVs with ≥0.2% relative abundance in at least

one of the samples were included. Separate networks were constructed for the 250 mg soil

samples and the soil aggregates. However, the selection of SVs was done on the joint data

matrix to ensure that both networks include the same SVs. The data was relativized to the total

sequence count of each sample. Pearson and Spearman correlations, mutual information (jsl

setting), and Bray-Curtis and Kullback-Leibler (with a pseudo count of 10^{-8}) dissimilarities were

calculated and the 1 000 highest and 1 000 lowest scoring edges from each of the five metrics

were kept. The ReBoot method [41], which mitigates compositional effects, was used to assess

the significance of the edges based on 1 000 permutations with renormalization and 1 000

bootstrap iterations. In the network of the soil aggregates, edges with scores below the 2.5[th]

and over the 97.5[th] percentile of the bootstrap distribution or not supported by at least three of

the five metrics were considered unstable and removed. Brown's method of p-value merging

was applied followed by Benjamini-Hochberg correction. Only edges with $q \leq 0.05$ were

included in the final network. In the network of the 250 mg samples, unstable edges were not

removed and the Benjamini-Hochberg correction was not applied as otherwise no edges were

retained. The networks were visualized in Cytoscape using the compound spring embedder

layout. Topological parameters were calculated with NetworkAnalyzer version 2.7 [42].

Results

Metagenomic DNA can be extracted from soil samples as small as 1 mg

DNA could be extracted from all soil samples including aggregates and amounts as little as 0.87

mg. In all cases, the extracted DNA was sufficient for 16S rRNA gene amplicon sequencing and

the qPCR assays. Accurate quantification of the DNA yield with PicoGreen assays was not

possible because the fluorescence readings from many of the small samples were close to the

background fluorescence in the blank controls.

To assess whether the DNA extraction could recover microbial DNA with similar efficiency from

small quantities of soil as from 250 mg samples, estimates of the abundances of *Bacteria*,

Archaea, and *Fungi* in one gram of soil were calculated from the qPCR results (Figure 1). Similar

estimates were obtained from the 250 mg and 5 mg soil samples from the 1[st] experiment.

Estimates from the 1 mg samples tended to be lower but were not significantly different. In

contrast, the estimates of fungal abundance in a gram of soil were significantly lower from the 1

and 5 mg than from the 250 mg samples. Estimates of bacterial abundance obtained from the

25 mg and 125 mg samples and archaeal abundance from the 25 mg samples were significantly

higher than from the 250 mg samples. Bacterial abundance estimates from the single

aggregates and 250 mg soil samples of the 3[rd] experiment covered the same range (Figure S1),

but the mean of the estimates from the single aggregates (2.27×10^9 copies/g soil) was lower (p

< 0.001) than from the 250 mg samples (4.69×10^9 copies/g soil).

Increasing spatial resolution reveals heterogeneity in soil bacterial and archaeal

community structure but not in their abundance

The yield of high-quality 16S rRNA gene amplicon sequences did not differ significantly between

small soil samples or single aggregates and the 250 mg samples in any of the three experiments

(Figure 2A). However, in the 1[st] experiment, the sequencing yield was somewhat lower from the

1 mg samples than from the 5 mg and 25 mg samples. The number of SVs detected in the

samples from the 1[st] experiment was not significantly different between the 250 mg, 125 mg,

and 25 mg samples, but decreased significantly in the 5 mg and even further among the 1 mg

samples (Figure 2B). Similarly, significantly lower numbers of SVs were detected in the single

aggregates than in the 250 mg soil samples of the 3[rd] experiment.

Principle component analysis (PCA) from the sequencing results from the 1[st] experiment

arranged all 250 mg and 125 mg samples, and most 25 mg samples into a single, tight group,

indicating high similarity in their prokaryotic community structures (Figure 3A). In contrast,

samples from the 5 mg and more so from the 1 mg categories, showed higher heterogeneity in

community structure. Similarly, PCA indicated heterogeneity between individual aggregates that

was not seen among the 250 mg samples in the 3rd experiment (Figure 3B).

The abundant SVs ($\geq$ 0.1% relative abundance in at least one of the samples) in the 250 mg, 125

mg, and 25 mg samples from the 1st experiment were almost all detectable in each individual

sample, showing that the composition of the soil prokaryotic community appears very uniform

when investigated at such a coarse spatial resolution (Figure 4). In contrast, 172 of the abundant

SVs detected in the 1 mg samples were unique to just one or two of these samples. Among

these SVs, representatives of *Planctomycetes*, *Proteobacteria*, and *Acidobacteria* were especially

numerous, while *Thaumarchaeota* and *Actinobacteria* were dominant among the SVs present in

all samples. The 5 mg samples represented a level of spatial resolution at which some

heterogeneity in the prevalence of the abundant SVs was already apparent with 22 of them

detectable in two or only in a single sample.

In total, 5 620 SVs were detected in the eight 1 mg soil samples of the 1st experiment. Of these,

4 764 (85%) were also present in at least half of the 250 mg samples. The remaining 856 SVs had

low relative abundance in the 1 mg samples with only 59 reaching more than 0.1% relative

abundance in any of them. The 5 mg samples together contained 8 010 SVs, of which 6 443

(80%) were also detectable in at least half of the 250 mg samples. Of the remaining 1 567 SVs,

only 26 reached more than 0.1% relative abundance in any of the 5 mg samples.

The qPCR results did not confirm our hypothesis that increasing spatial resolution would reveal

heterogeneity in microbial abundance. Bacterial and archaeal 16S rRNA gene, and fungal ITS

copy numbers did not show a larger variation among the 1 and 5 mg samples than between the

250 mg samples from the 1st experiment (Figure 1). Similarly, in the 3rd experiment, bacterial

abundance did not vary more in the single aggregates than in the 250 mg samples (Figure S1).

Impact of contamination, inconsistent performance of the methods, and stochastic

effects

The control samples amplified in the qPCR assay targeting the bacterial 16S rRNA gene but

yielded only 3 to 356 copies per µl DNA extract. In comparison, the 1 mg soil samples had 8 777

to 45 534 copies per µl DNA extract. The control samples from the 1st experiment had 0 to 13

fungal ITS copies per µl DNA extract and none of them showed amplification in the archaeal 16S

rRNA gene qPCR assays. It was possible to generate sequencing results from all control samples.

They were very similar to each other in their prokaryotic community structures but very

different from the soil samples (Figure S2). Every SV that reached 0.1% relative abundance in

any of the control samples of an experiment was considered as a potential contaminant and

removed from the datasets. There were 450, 77, and 591 such SVs in the datasets of the 1st, 2nd,

and 3rd experiments respectively. In the data from the 1st experiment, these SVs together

covered 98.9 − 99.8% of the sequences obtained from the control samples and 2.5 − 6.6% of the

sequences from the soil samples. In the 2nd and 3rd experiments, 92.5 − 99.5% and 98.6 − 99.6%

of the sequences from the control samples, and 2.9 − 17.3% and 5.9 − 9.4% of the sequences

from the soil samples, respectively, were covered by the potentially contaminant SVs.

The estimated bacterial abundance in a gram of soil based on the qPCR results was in general

higher in the samples from the 2nd experiment but followed the same pattern as in the samples

from the 1st experiment with no differences between the 1 mg, 5 mg, and 250 mg samples but

significantly higher values in the 25 mg samples (Figure S3). The small soil homogenate samples

did not show the degree of heterogeneity in prokaryotic community structure we observed

among the small soil samples of the 1st experiment. Abundant SVs ($\geq$ 0.1% relative abundance in

at least one sample) in the 25 mg soil homogenate samples were all detectable in at least five of

the eight replicates. Out of the 354 abundant SVs in the 5 mg soil homogenate samples, one

was present in only three of the samples but the others were detectable in at least six. The 1 mg

soil homogenate samples harbored 446 abundant SVs. None of them was unique to a single

sample and 442 were present in five or more of the eight samples. The Aitchison distances of

community structure were much higher among the 5 mg, and especially among the 1 mg soil

samples of the 1st experiment compared to the distances between the 250 mg soil samples

(Figure 5). In contrast, the distances between the 1 mg, 5 mg, or 25 mg soil homogenate

samples of the 2nd experiment were very similar to the distances among the 250 mg soil

samples indicating no difference in the heterogeneity of prokaryotic community structure

between these sample groups.

Bacterial and archaeal co-occurrence patterns in 250 mg-soil samples and aggregates

Networks of prokaryotic co-occurrence were constructed using the 272 SVs that reached ≥0.2%

relative abundance in at least one of the samples of the 3rd experiment. No network was

obtained from the 250 mg samples unless the removal of unstable edges and the correction of

the p-values for multiple testing were skipped. The resulting network has 78 edges between 35

nodes (Figure 6A). Thus, this spatial resolution revealed only a small number of putative

associations many of which are likely false discoveries. In contrast, a network of 137 edges and

67 nodes (with the removal of unstable edges and control of the false discovery rate) was

obtained from the individual soil aggregates (Figure 6B). A total of 54 of the nodes are part of a

connected component in which there are three nodes with high betweenness centrality: SV15

(*Verrucomicrobia*, candidatus *Udaeobacter*), SV31 (*Actinobacteria*), and SV36 (*Acidobacteria*

subgroup 6). These SVs potentially serve a keystone function by connecting two clusters in the

network. One of the clusters contains several SVs of *Thaumarchaeota*, *Verrucomicrobia*, and

Actinobacteria. The other is dominated by *Acidobacteria* subgroup 6. The hub of the latter

cluster is SV58 (*Acidobacteria* subgroup 6) that has the highest degree in the network being

connected to 19 nodes. SV399 (*Chloroflexi*) and SV123 (*Acidobacteria* subgroup 6) are linked

with negative associations to several members of this cluster.

Discussion

The large disparity between the scale in which the soil microbiota is usually studied with

molecular methods (0.25 to 1 g of soil) and the distance over which microbial interactions occur,

impede the detection of interacting partners [43]. In order to gain information on the soil

microbial diversity at an increased spatial resolution that considers soil structure, in this study

we reduced the amount of soil used for DNA extraction from 250 to 1 mg and also extracted

individual soil aggregates. Bacterial and archaeal DNA was recovered with similar efficiency

from the 250 mg and the 5 mg and 1 mg samples as shown by the qPCR results. This was not

true for fungal DNA. Either the DNA-extraction kit was not efficient in isolating fungal DNA from

samples below 25 mg, or fungi may preferentially colonize larger soil aggregates. Our results

show that the DNA extraction kit was most efficient in recovering bacterial and archaeal DNA

from 25 to 125 mg soil, although the variation in the yield of 16S rRNA gene copies was large

among these samples. Since this increased variation was apparent among the 25 mg soil

homogenate samples of the 2nd experiment as well, it must be a property of the DNA extraction

kit and not an indication of uneven distribution of bacterial cells at the scale of 25 mg samples.

As a consequence of sampling small amounts of soil, the DNA extracts had low template

concentrations for the subsequent PCR analyses. Thereby we had to anticipate a high risk of

contamination affecting the results [44]. In fact, quantifiable amounts of *Bacteria* and *Fungi*, but

not *Archaea*, were detected in the control samples without soil. However, they reached no

more than 4% of the number of bacterial rRNA genes copies in the smallest soil samples, and

thus, the influence of contamination on our results is negligible. The bacterial community found

in the control samples was clearly distinct from the soil communities suggesting that the

contamination originated from the reagents of the DNA extraction and sequencing library

preparation rather than cross-contamination between samples [45, 46]. Another concern of

working with small samples is that molecular methods applied to such low amounts of template

may perform inconsistently leading to artificial variation in the results. The 2nd experiment

showed that the DNA extraction, PCR, and sequencing did not artificially generate more

variation in the results from small samples than the variation present among the 250 mg

samples. Therefore, the heterogeneity in prokaryotic community composition and structure

among the 1 mg and 5 mg soil samples from the 1st experiment was not caused by stochastic

effects or PCR bias.

Samples of 25 mg up to 250 mg of soil were close to identical in prokaryotic community

composition, thus they provide a good representation of the overall prokaryotic diversity of our

soil. In contrast, the 1 mg and 5 mg samples and single aggregates were heterogeneous in

community structure. We found that while small soil samples could recover some SVs not

necessarily detected with the conventionally used 250 mg samples; these SVs were typically low

in abundance. Very few exceeded the relative abundance threshold we applied to control the

sparsity of the data in our analysis of community structure. Therefore, the large heterogeneity

of prokaryotic community structure we observed among the small samples was not because

they would have enabled the detection of more SVs. Instead, it appears that they contained

different subsets of the total community present in the 250 mg samples. Interestingly, the small

samples didn't have more variation in the abundance of *Bacteria, Archaea*, and *Fungi* than the

250 mg samples. Microbial cells in soil have a patchy distribution at the scale of a few

micrometers [47] but, for the soil of this study apparently not at the scale of macroaggregates

or 1 to 5 mg samples.

Network analyses based on microbial co-occurrence have been applied to soil samples as large

as 10 grams [48]. The scientific value of utilizing much smaller samples is shown by our result

that 1 and 5 mg samples contained subsets of the total microbial diversity. The smaller spatial

scale increases the likelihood that the observed co-occurrences actually indicate interactions

[49]. From 37 soil aggregates, we obtained a complex network of bacterial and archaeal co-

occurrence that contained two clusters, one with several *Thaumarchaeota, Verrucomicrobia*

and *Actinobacteria* SVs, the other mainly with *Acidobacteria* subgroup 6 SVs. Three SVs, which

could represent keystone taxa, were found to connect these clusters. Keystone taxa are

considered to be the best predictors for whole-community compositional changes [50]. If these

putative keystone SVs are abundant in an aggregate, we can expect that members of both

clusters are present there.

While the spatial scale that we reached in this study is not yet fine enough to reveal most

microbial interactions as they may occur in microaggregates [51], it can support the

development of hypotheses and experiments to understand the patterns and processes shaping

the soil microbiota and modelling its behavior [52-54]. Developing DNA extraction protocols

from even small soils samples, as typical for microaggregates, should be a way forward to unveil

interactions within soil microbial communities, thereby further enhancing our understanding of

how soil microbial communities are organized at structural and functional levels.

Acknowledgements

We thank Ines Merbach, Department Biozönoseforschung, Helmholtz-Zentrum für

Umweltforschung – UFZ, Bad Lauchstädt, Germany, for providing the soil of this study. We

thank Naomie Oßwald and Jerome Rischke who supported us at early stages of this work as

bachelor students, and Karin Trescher for excellent technical assistance. The study was

financially supported by the German Research Foundation (DFG; Grant number TE 383/10-1) in

context of the Priority Programme "Rhizosphere spatiotemporal organisation (SPP2089).

Conflict of interest

The authors declare no conflict of interest.

References

1. Green JL, Bohannan BJM, Whitaker RJ. Microbial biogeography: from taxonomy to traits.

Science. 2008. 320;5879:1039-43.

2. Widder S, Allen RJ, Pfeiffer T, Curtis TP, Wiuf C, Sloan WT, et al. Challenges in microbial

ecology: building predictive understanding of community function and dynamics. ISME J.

2016. 10;11:2557-68.

3. Vos M, Wolf AB, Jennings SJ, Kowalchuk GA. Micro-scale determinants of bacterial diversity in

soil. FEMS Microbiol Rev. 2013. 37;6:936-54.

4. Thakur MP, Phillips HRP, Brose U, De Vries FT, Lavelle P, Loreau M, et al. Towards an

integrative understanding of soil biodiversity. Biol Rev. 2020. 95;2:350-64.

5. Six J, Elliott ET, Paustian K. Soil macroaggregate turnover and microaggregate formation: a

mechanism for C sequestration under no-tillage agriculture. Soil Biol Biochem. 2000.

32;14:2099-103.

6. Ranjard L, Poly F, Combrisson J, Richaume A, Gourbiere F, Thioulouse J, et al. Heterogeneous

cell density and genetic structure of bacterial pools associated with various soil

microenvironments as determined by enumeration and DNA fingerprinting approach (RISA).

Microbial Ecol. 2000. 39;4:263-72.

7. Smith P, Cotrufo ME, Rumpel C, Paustian K, Kuikman PJ, Elliott JA, et al. Biogeochemical cycles

and biodiversity as key drivers of ecosystem services provided by soils. Soil. 2015. 1;2:665-85.

8. Wilpiszeski RL, Aufrecht JA, Retterer ST, Sullivan MB, Graham DE, Pierce EM, et al. Soil

aggregate microbial communities: towards understanding microbiome interactions at

biologically relevant scales. Appl Environ Microbiol. 2019. 85;14:UNSP e00324-19

9. Rillig MC, Müller LAH, Lehmann A. Soil aggregates as massively concurrent evolutionary

incubators. ISME J. 2017. 11;9:1943-8.

10. Young JM, Rawlence NJ, Weyrich LS, Cooper A. Limitations and recommendations for

successful DNA extraction from forensic soil samples: A review. Sci Justice. 2014. 54;3:238-

44.

11. Fierer N, Jackson RB. The diversity and biogeography of soil bacterial communities. Proc Nat

Acad Sci USA. 2006. 103;3:626-31.

12. Roesch LF, Fulthorpe RR, Riva A, Casella G, Hadwin AKM, Kent AD, et al. Pyrosequencing

enumerates and contrasts soil microbial diversity. ISME J. 2007. 1;4:283-90.

13. Griffiths RI, Thomson BC, James P, Bell T, Bailey M, Whiteley AS. The bacterial biogeography

of British soils. Environ Microbiol. 2011. 13;6:1642-54.

14. Karimi B, Terrat S, Dequiedt S, Saby NPA, Horriguel W, Lelievre M, et al. Biogeography of soil

bacteria and archaea across France. Sci Adv. 2018. 4;7:aat1808.

15. Szoboszlay M, Dohrmann AB, Poeplau C, Don A, Tebbe CC. Impact of land-use change and

soil organic carbon quality on microbial diversity in soils across Europe. FEMS Microbiol

Ecol. 2017. 93;12:fix146.

16. Hemkemeyer M, Dohrmann AB, Christensen BT, Tebbe CC. Bacterial preferences for specific

soil particle size fractions revealed by community analyses. Front Microbiol. 2018. 9:00149.

17. Hemkemeyer M, Christensen BT, Tebbe CC, Hartmann M. Taxon-specific fungal preference

for distinct soil particle size fractions. Eur J Soil Biol. 2019. 94:103103.

18. Constancias F, Prevost-Boure NC, Terrat S, Aussems S, Nowak V, Guillemin JP, et al.

Microscale evidence for a high decrease of soil bacterial density and diversity by cropping.

Agron Sustain Dev. 2014. 34;4:831-40.

19. Davinic M, Fultz LM, Acosta-Martinez V, Calderon FJ, Cox SB, Dowd SE, et al. Pyrosequencing

and mid-infrared spectroscopy reveal distinct aggregate stratification of soil bacterial

communities and organic matter composition. Soil Biol Biochem. 2012. 46:63-72.

20. Fox A, Ikoyi I, Torres-Sallan G, Lanigan G, Schmalenberger A, Wakelin S, et al. The influence

of aggregate size fraction and horizon position on microbial community composition. Appl

Soil Ecol 2018. 127;19-29.

21. Ivanova EA, Kutovaya OV, Tkhakakhova AK, Chernov TI, Pershina EV, Markina LG, et al. The

structure of microbial community in aggregates of a typical chernozem aggregates under

contrasting variants of its agricultural use. Eurasian Soil Sci. 2015. 48;11:1242-56.

22. Bach EM, Williams RJ, Hargreaves SK, Yang F, Hofmockel KS. Greatest soil microbial diversity

found in micro-habitats. Soil Biol Biochem. 2018. 118:217-26.

23. Ranjard L, Nazaret S, Gourbiere F, Thioulouse J, Linet P, Richaume A. A soil microscale study

to reveal the heterogeneity of Hg(II) impact on indigenous bacteria by quantification of

adapted phenotypes and analysis of community DNA fingerprints. FEMS Microbiol Ecol.

2000. 31;2:107-15.

24. Bailey VL, Fansler SJ, Stegen JC, McCue LA. Linking microbial community structure to beta-

glucosidic function in soil aggregates. ISME J. 2013. 7;10:2044-53.

25. Kravchenko AN, Negassa WC, Guber AK, Hildebrandt B, Marsh TL, Rivers ML. Intra-aggregate

pore   structure   influences   phylogenetic   composition   of   bacterial   community   in

macroaggregates. Soil Sci Soc Am J. 2014. 78;6:1924-39.

26. Bailey VL, McCue LA, Fansler SJ, Boyanov MI, DeCarlo F, Kemner KM, et al. Micrometer-scale

physical structure and microbial composition of soil macroaggregates. Soil Biol Biochem.

2013. 65;60-8.

27. Direito SOL, Zaura E, Little M, Ehrenfreund P, Roling WFM. Systematic evaluation of bias in

microbial community profiles induced by whole genome amplification. Environ Microbiol.

2014. 16;3:643-57.

28. Wang Y, Gao ZM, Xu Y, Li GY, He LS, Qian PY. An evaluation of multiple annealing and

looping based genome amplification using a synthetic bacterial community. Acta Oceanol

Sin. 2016. 35;2:131-6.

29. Kim H, Nunan N, Dechesne A, Juarez S, Grundmann G. The spatial distribution of exoenzyme

activities across the soil micro-landscape, as measured in micro- and macro-aggregates, and

ecosystem processes. Soil Biol Biochem. 2015. 91;258-67.

30. Miller DN, Bryant JE, Madsen EL, Ghiorse WC. Evaluation and optimization of DNA extraction

and purification procedures for soil and sediment samples. Appl Environ Microbiol. 1999.

65;11:4715-24.

31. Hemkemeyer M, Christensen BT, Martens R, Tebbe CC. Soil particle size fractions harbour

distinct microbial communities and differ in potential for microbial mineralisation of organic

pollutants. Soil Biol Biochem. 2015. 90;255-65.

32. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a dual-index

sequencing strategy and curation pipeline for analyzing amplicon sequence data on the

MiSeq Illumina sequencing platform. Appl Environ Microbiol. 2013. 79;17:5112-20.

33. Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, et al. Improved

bacterial 16S rRNA gene (V4 and V4-5) and fungal internal transcribed spacer marker gene

primers for microbial community surveys. mSystems. 2016. 1;1:e00009-15.

34. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-

resolution sample inference from Illumina amplicon data. Nat Methods. 2016. 13;7:581-3.

35. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig WG, Peplies J, et al. SILVA: a comprehensive

online resource for quality checked and aligned ribosomal RNA sequence data compatible

with ARB. Nucleic Acids Res. 2007. 35;21:7188-96.

36. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan: community

ecology   package.   R   package   version   2.5-2.   2018.   https://CRAN.R-

project.org/package=vegan.

37. Palarea-Albaladejo J, Martin-Fernandez JA. zCompositions - R Package for multivariate

imputation of left-censored data under a compositional approach. Chemometrics Intell Lab

Syst. 2015. 143;85-96.

38. Gloor GB, Wu JR, Pawlowsky-Glahn V, Egozcue JJ. It's all relative: analyzing microbiome data as compositions. Ann Epidemiol. 2016. 26;5:322-9.

39. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome datasets are compositional: and this is not optional. Front Microbiol. 2017. 8;6:02224.

40. Faust K, Raes J. CoNet app: inference of biological association networks using Cytoscape [version 2; peer review: 2 approved]. 2016. 5:1519. f1000research.9050.2.

41. Faust K, Sathirapongsasuti JF, Izard J, Segata N, Gevers D, Raes J, et al. Microbial Co-occurrence relationships in the human microbiome. PLoS Comput Biol. 2012. 8;7: e1002606.

42. Assenov Y, Ramirez F, Schelhorn SE, Lengauer T, Albrecht M. Computing topological parameters of biological networks. Bioinformatics. 2008. 24;2:282-4.

43. Nunan N. The microbial habitat in soil: Scale, heterogeneity and functional consequences. J Plant Nutr Soil Sci. 2017. 180;4:425-9.

44. Weiss S, Amir A, Hyde ER, Metcalf JL, Song SJ, Knight R. Tracking down the sources of experimental contamination in microbiome studies. Genome Biol. 2014. 15;12:564.

45. Glassing A, Dowd SE, Galandiuk S, Davis B, Chiodini RJ. Inherent bacterial DNA contamination of extraction and sequencing reagents may affect interpretation of microbiota in low bacterial biomass samples. Gut Pathog. 2016. 8:24.

46. Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, et al. Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. BMC Biol. 2014. 12:87.

47. Nunan N, Wu KJ, Young IM, Crawford JW, Ritz K. Spatial distribution of bacterial

communities and their relationships with the micro-architecture of soil. FEMS Microbiol

Ecol. 2003. 44;2:203-15.

48. Khan MAW, Bohannan BJM, Nüsslein K, Tiedje JM, Tringe SG, Parlade E, et al. Deforestation

impacts network co-occurrence patterns of microbial communities in Amazon soils. FEMS

Microbiol Ecol. 2019. 95;2:fiy230.

49. Cordero OX, Datta MS. Microbial interactions and community assembly at microscales. Curr

Opin Microbiol. 2016. 31:227-34.

50. Herren CM, McMahon KD. Keystone taxa predict compositional change in microbial

communities. Environ Microbiol. 2018. 20;6:2207-17.

51. Raynaud X, Nunan N. Spatial ecology of bacteria at the microscale in soil. PLOS ONE. 2014.

9;1:10.1371.

52. Faust K, Raes J. Microbial interactions: from networks to models. Nat Rev Microbiol. 2012.

10;8:538-50.

53. Rillig MC, Ryo M, Lehmann A, Aguilar-Trigueros CA, Buchert S, Wulf A, et al. The role of

multiple global change factors in driving soil functions and microbial biodiversity. Science.

2019. 366;6467:886-90.

54. Tecon R, Or D. Biophysical processes supporting the diversity of microbial life in soil. FEMS

Microbiol Rev. 2017. 41;5:599-623.

Figure legends

Figure 1 Estimates of (A) bacterial, (B) archaeal, and (C) fungal abundance in a gram of soil based on qPCR from the samples from the 1st experiment (gene copy numbers per g of soil wet weight). One of the 250 mg soil samples was an outlier in the fungal ITS qPCR results and is not included in the plot. Sample groups not labelled with the same letter were significantly different in Tukey's HSD tests.

Figure 2 Number of (A) sequences and (B) sequence variants (SVs) in the samples after the removal of potentially contaminant SVs. Letters indicate significant differences between sample groups of the 1st experiment according to Tukey's HSD tests. * indicates significant difference based on Welch's t-test between the aggregate and the 250 mg soil samples of the 3rd experiment.

Figure 3 Principle component analyses (PCA) plots from the 16S rRNA gene sequencing data from the 1st (A) and 3rd (B) experiments.

Figure 4 SVs arranged according to in how many of the 1 mg, 5 mg, 25 mg, 125 mg, or 250 mg samples from the 1st experiment they were detected. Only SVs that reached at least 0.1% relative abundance in any of the samples are included. Each node represents one SV colored based on its phylum-level classification and sized according to its average relative abundance across all samples excluding those in which it was not detected.

Figure 5 Aitchison distances in the bacterial and archaeal community structure (16SrRNA gene amplicons) within sample groups from the 1st and 2nd experiments.

549   **Figure 6** Co-occurrence networks from the (A) 250 mg samples and (B) the single aggregates

550   from the 3rd experiment. The frames mark the two clusters discussed in the text. It should be

551   noted that in (A) unstable edges were *not* removed and the Benjamini-Hochberg correction for

552   multiple comparisons was not applied.

553   **Table 1** Number of samples and soil weights within each sample category

Soil weight class or sample type	Sample weight [mg] ± SD	Number of soil samples
1^{st} Experiment		
250-mg	251 ± 1	8
125-mg	125 ± 1	8
25-mg	25.1 ± 0.4	8
5-mg	4.9 ± 0.2	8
1-mg	1.1 ± 0.2	8
Control, no soil	-	6
2^{nd} Experiment		
250-mg	252 ± 3	8
25-mg soil homogenate	-	8
5-mg soil homogenate	-	8
1-mg soil homogenate	-	8
Control, no soil	-	6
3^{rd} Experiment		
250-mg	251 ± 0	35
Soil aggregate	5.3 ± 1.6	37
Control	-	9

     Figure 1

560   Figure 2

563   Figure 3

Figure 4

569   Figure 5

572   Figure 6

A

B