1 **Consistent marine biogeographic boundaries across the tree of life despite centuries of**

2 **human impacts**

3
4

Luke E. Holman[1,*] (ORCID: 0000-0002-8139-3760)

Mark de Bruyn[2,3] (ORCID:0000-0003-1528-9604)

Simon Creer[2]

Gary Carvalho[2]

Julie Robidart[4] (ORCID: 0000-0001-9805-3570)

Marc Rius[1,5] (ORCID: 0000-0002-2195-6605)

[1] School of Ocean and Earth Science, National Oceanography Centre Southampton, University

of Southampton, United Kingdom

[2] Molecular Ecology and Fisheries Genetics Laboratory, School of Natural Sciences, Bangor

University, United Kingdom

[3] The University of Sydney, School of Life and Environmental Sciences, Australia

[4] Ocean Technology and Engineering Group, National Oceanography Centre Southampton,

United Kingdom

[5] Department of Zoology, Centre for Ecological Genomics and Wildlife Conservation,

University of Johannesburg, South Africa

* Corresponding author: lukeearlholman@gmail.com

Abstract

Over millennia, ecological and evolutionary mechanisms have shaped macroecological

distributions across the tree of life. Research describing patterns of regional and global

biogeography has traditionally focussed on the study of conspicuous species. Consequently,

there is limited understanding of cross-phyla biogeographic structuring, and an escalating need to

understand the macroecology of both microscopic and macroscopic organisms. Here we used

environmental DNA (eDNA) metabarcoding to explore the biodiversity of marine metazoans,

micro-eukaryotes and prokaryotes along an extensive and highly heterogeneous coastline. Our

results showed remarkably consistent biogeographic structure across the kingdoms of life, which

were underpinned by environmental and anthropogenic influence. Additionally, metazoan

communities displayed biographic patterns that suggest regional biotic homogenisation of

conspicuous species. Against the backdrop of global pervasive anthropogenic environmental

change, our work highlights the importance of considering multiple domains of life to understand

the maintenance and drivers of marine biodiversity across broad taxonomic, ecological and

geographical scales.

Introduction

Researchers have long recognised the importance of grouping global biota into distinct, geographically separated regions. Delineating these biogeographic areas is important to understand the factors shaping the range limits of species[1], to designate key geographic areas for biodiversity conservation[2] and in formulating predictive responses to environmental change[3,4]. One of the first efforts to define geographic regions of terrestrial biota were Alfred Russel Wallace's so-called 'Zoological Regions'[5], which included six major regions (hereafter realms) that are still recognised in subsequent classifications today[6]. The drivers responsible for these geographic classifications are frequently environmental conditions or physical barriers. Biogeographic studies have shown that deep divergence in the geographic arrangement of terrestrial biota arose as a result of plate tectonics, while shallow divergence has been most frequently attributed to climatic conditions[7]. In aquatic ecosystems, the relative importance of macroecological drivers is less understood, although both climatic (e.g. temperature)[8] and tectonic forces[9] have been identified as key determinants of marine biogeographic patterns. Recent studies have successfully partitioned the oceans into distinct ecoregions (i.e. a geographically defined area, smaller than realm, that contains characteristic assemblages of species)[1,10], but the description of marine ecoregions has mostly considered conspicuous or well-described species. Similarly, most macroecological marine research has focussed on readily identifiable eukaryotic species, principally metazoans[11], although demonstrable progress has been made in understanding global patterns of marine microbes[12]. In line with recent studies demonstrating strong cross-phylum interdependence[13], there is an increasing need to include prokaryotic species in our assessment of biogeographic patterns. The language of macroecology and microbial ecology is similar, both examining the incidence of species across different spatial scales, but these fields have long progressed independently. As a result, relatively few studies have explored biogeographic patterns simultaneously for both microscopic and macroscopic life, with examples of consistent and inconsistent patterns across different taxa[13-15]. Work is thus needed to explore the consistency of biogeographic breaks across different kingdoms of life.

Human driven habitat destruction, pollution and the introduction of non-native species are the main drivers of recent global biodiversity change[3] and therefore have the potential to alter

geographic patterns of biota at multiple spatial scales. Cumulatively, anthropogenic stressors not

only threaten vulnerable native species but whole-community structure and function[3,16,17]. The

magnitude and direction of human impacts are complex, with evidence for both gains and losses

in local species richness across biomes[18-20]. However, a consistent global pattern is emerging,

with a recent and rapid increase in species turnover[19] and an associated increase in community

similarity (β diversity) between two or more geographically separated sites[20]. Incidences of

increased community similarity are known as biotic homogenisation[21] and are driven by human

activities that promote extinctions and the introductions of non-native species. In light of

growing evidence that taxonomic, phylogenetic and functional diversity are strongly correlated[22],

the homogenisation of biological communities has the potential to negatively affect ecosystem

function. Furthermore, even uncommon species within an ecological community can contribute

significantly to ecosystem function[23], demonstrating the importance of studying inconspicuous

species to preserve ecosystem health. Studies have shown evidence for biotic homogenisation

around the globe, with examples from plants[18], vertebrates[24] and invertebrates[25] demonstrating

alteration of terrestrial biogeographic patterns. However, many studies are of limited taxonomic

scope, focussing on highly conspicuous species for which reliable data can be relatively easily

produced[24,25]. Thus, most work overlooks inconspicuous species (e.g. microbes and microscopic

eukaryotes), which show vastly different replication, demographic and dispersal patterns

compared to metazoans[13], but are known to be key actors shaping the assembly of ecological

communities and ultimately underpin ecosystem functioning[26]. Taken together, a more

comprehensive characterisation of ecological communities is clearly needed when testing the

role of biotic homogenisation on biogeographic patterns.

The advent of high-throughput sequencing has revolutionised our understanding of microbial

life, with studies examining global patterns of prokaryotic life now increasingly common[12].

Moreover, the recent and rapid development of methods to infer the incidence of larger

organisms using genetic material isolated from environmental samples (known as environmental

DNA or eDNA) has provided an unparalleled ability to identify species across the entire tree of

life[14,15,27]. Together, these methods can rapidly generate standardised biodiversity data for entire

communities at an unprecedented resolution, thereby minimising regional and taxonomic biases.

Such studies provide datasets that can be analysed without taxonomic assignment and DNA

samples can be repurposed to test novel hypotheses. A common technique is to amplify DNA

barcodes from eDNA and use high-throughput sequencing to produce high-resolution

biodiversity data. This method (eDNA metabarcoding) has been shown to reliably detect

organisms across many different ecosystems[27], but has infrequently been applied to understand

spatial patterns of biodiversity across different kingdoms of life[12,15].

A unique geographic setting for testing biogeographic hypotheses is the South African coastline,

where two large water masses (the Atlantic and Indian Oceans) meet, and a wide variety of

abiotic and biotic conditions are found in a relatively small region. This coastline has three well-

defined coastal ecoregions bounded by the cold western boundary Benguela Current and the

warm oligotrophic eastern boundary Agulhas Current. These ecoregions have been established

on the basis of studies over several decades involving a number of conspicuous metazoan

taxa[2,28]. Additionally, there is evidence for human exploitation of marine resources in the region

spanning thousands of years[29,30] and some areas of the coastline have been subject to heavy

maritime activity for centuries[31]. Other human activities also prevail such as the establishment of

aquaculture facilities or the construction of harbours and breakwaters[29,30]. Thus, the South

African coastline is an ideal study system to explore the mechanisms shaping biogeographic

boundaries.

Here we used eDNA metabarcoding to compare the biogeography of multiple marine kingdoms

along the diverse coastline of South Africa and to test the effects of life-history strategies and

anthropogenic and natural pressures on biogeographic patterns. We first investigated the

consistency of biogeographic boundaries across metazoans, protists and bacteria, and to what

extent these patterns could be explained by environmental conditions. We then evaluated if there

was evidence for homogenisation of ecological communities along a coastline that has been

affected by centuries of human impacts, and to what extent these patterns persisted across

taxonomic groups.

Methods

Field sampling

135
We sampled a range of sites along 2,000 km of coastline (Fig. 1) between October and

November of 2017 (see details in Supplementary Table 1), covering the three major marine

ecoregions of South Africa. In order to assess the effects of anthropogenic impacts, we compared

human altered 'artificial' sites (e.g. recreational marinas, harbours) and relatively unaltered rocky

shore and natural harbour sites, hereafter 'natural' (see Supplementary Table 1). The artificial

sites were previously sampled in Rius, et al. [32], and six adjacent natural sites were selected. The

natural sites were the nearest non-developed sites with matching aspect and exposure (Fig. 1) to

each of the artificial sampling sites. Three 400 ml seawater samples were filtered with 0.22 μm

polyethersulfone membrane Sterivex filters (Merck Millipore, MA, USA) following the

sampling scheme of Holman, et al. [33] at each sampling site. Consequently, we sampled a total of

1,200 ml of seawater per site, a volume that has been shown to differentiate fine scale (<1 km^2)

community structure in marine systems[33,34]. Filters were immediately preserved at ambient

temperature with the addition of 1.5 ml of Longmire's Solution for preservation until DNA

extraction. Field control filters and equipment cleaning blanks were taken, transported, stored

and sequenced as the rest of the field samples.

Environmental DNA extraction

152
We used a PCR-free laboratory separated from the main molecular biology laboratory facilities.

No post-PCR or high concentration DNA samples were permitted in the laboratory. All surfaces

and lab equipment were cleaned thoroughly before use with 1.25% sodium hypochlorite solution

(3:1 dilution of household bleach). DNA extraction followed the SX^{CAPSULE} method from Spens,

et al. [35]. Briefly, filters were first externally cleaned with sterile water and Longmire's Solution

was removed from the filter outlet using a sterile syringe, 720 μl Buffer ATL (Qiagen, Hilden,

Germany) and 80 μl Proteinase K (20mg/ml) was added and filters were incubated overnight at

56°C. The lysate was then removed from the filter inlet and subjected to DNA extraction using

the Qiagen DNeasy Blood and Tissue Kit under the manufacturers recommended protocol. DNA

was eluted using 200 μl Qiagen Buffer AE and re-eluted once to increase DNA yield. All DNA

samples were checked for PCR inhibition using the Primer Design Internal Positive Control

qPCR Kit (Primer Design, Southampton, UK) with 10 μl reactions under the manufacturer

recommended protocol. Inhibition was detected by an increase of >1.0 Ct in reactions containing

eDNA compared to reactions with extraction controls. As inhibition was detected in a minority

of samples, all samples were treated using the Zymo OneStep PCR Inhibition Removal Kit

(Zymo Research, California, USA) following the manufacturer recommended protocol. Inhibited

samples showed no evidence for inhibition post cleaning.

High throughput eDNA amplicon sequencing

Different sets of primers were used to generate three separate eDNA metabarcoding libraries for

all samples. Two gene regions were selected to target broad metazoan/eukaryotic diversity, a

section of the V4 region of the nuclear small subunit ribosomal DNA (hereafter 18S)[36] and a

section of the standard DNA barcoding region of cytochrome c oxidase subunit I (hereafter

COI)[37]. One gene region was used to target prokaryotes, the V3-V4 hypervariable region of

prokaryotic small subunit ribosomal DNA (hereafter 16S)[38]. Illumina unique double-indexed

metabarcoding amplicon libraries were constructed with a two-step PCR protocol as detailed in

Holman, et al. [33]. The first PCR setup was performed in a PCR-free laboratory. The three eDNA

samples per site were pooled and three independent technical replicates were sequenced per pool.

The process per sequenced pool was as follows. The first PCR reaction was conducted in

triplicate in a total reaction volume of 20 µl. Each reaction contained 10 µl Amplitaq GOLD 360

2X Mastermix (Applied Biosystems, California, USA), 0.8 µl (5 nmol ml^{-1}) of each forward and

reverse primers and 2 µl of undiluted environmental DNA template. The reaction conditions for

PCR were an initial denaturation step at 95°C for 10 minutes followed by 20 cycles of 95°C for

30 seconds, variable annealing temp (46°C for COI, 50°C for 18S and 55°C for 16S) for 30

seconds, and extension at 72°C for 1 minute. A final extension at 72°C was performed for 10

minutes. The triplicate first PCR replicates were then pooled and cleaned using AMPure XP

beads (Beckman Coulter, California, USA) at 0.8 beads:sample volume ratio following

manufacturer's instructions. The second PCR reaction was conducted in a total volume of 20 µl

containing 10 µl Amplitaq GOLD 360 2X Mastermix, 0.5 µl (10 nmol ml^{-1}) of both forward and

reverse primers and 5 µl of undiluted cleaned PCR product from the first reaction. PCR

conditions were an initial denaturation step at 95°C for 10 minutes followed by 15 cycles of

95°C for 30 seconds, annealing at 55°C for 30 seconds, and extension at 72°C for 1 minute. A

final extension at 72°C was performed for 10 minutes. PCR 2 products were cleaned using

AMPure XP beads as above. Negative control samples for the filters, extraction kit, PCR1 and 2

were included in library building and sequenced alongside experimental samples. Products were

quantified following the manufacturer's instructions using the NEBNext Library Quant qPCR kit

(New England Biolabs, Massachusetts, USA) and then normalised and pooled at an equimolar

concentration for each marker. Each gene region was sequenced independently using a V3

paired-end 300bp reagent kit on the Illumina MiSeq Instrument with 5% PhiX genomic library

added to increase sequence diversity.

Bioinformatics

Raw sequences were de-multiplexed using the GenerateFastQ (v2.0.0.9) module on the MiSeq

control software (v3.0.0.105). Cutadapt (v2.3)[39] was used to filter sequences to include only

those that contained both the forward and reverse primer sequence across both read pairs for

each gene fragment, remaining sequences then had the primer region removed for each gene

fragment using the default settings. Sequences were denoised using the DADA2 pipeline

(v1.12)[40] in R (v3.6.1)[41] with the default parameters unless noted as follows. Sequences were

filtered to retain only pairs of reads with an expected error of 1 or fewer per read. Read trimming

was performed after manual examination of the read quality profile, the forward reads were

trimmed to 250bp (COI), 240bp (18S) and 240 bp (16S) and the reverse reads were trimmed to

230bp (COI), 220bp (18S) and 220 bp (16S). As each marker was sequenced separately, the

differences in read trimming length reflect typical variation in sequencing runs rather than any

biological difference. The error rates per run were estimated and used to perform the denoising

using the DADA2 algorithm. The denoised sequence pairs were then merged and resulting

sequences were truncated if they were outside of the expected gene fragment range (303-323bp

for COI, 400-450bp for 18S and 390-450bp for 16S). Chimeras were identified and removed

before assembling a sample by ASV (amplicon sequence variant) table for analysis. The

denoised ASVs were then curated using the default settings of the LULU algorithm[42] which

merges sequences based on sequence similarity and co-occurrence. Assigning taxonomy to a set

of unknown sequences is a difficult task, particularly considering many marine species lack

DNA barcodes, are undescribed, or have erroneous barcodes in online public databases. We

therefore focused our analysis at a higher taxonomic level than species, assigning taxonomy to

sequences from the COI and 18S data as follows. The RDP classifier (v2.13)[43] was used to

assign taxonomy for COI using a previously published COI database[44] (v4.0) and a modified

version of the SILVA database[45] (v3.2 from https://github.com/terrimporter/18SClassifier). As

species level assignments have been shown to be accurate for COI data[33] an unconstrained (no

limits on sequence similarity or match length) BLAST search (v2.6.0+) was performed for each

sequence against the entire National Centre for Biotechnology Information nt database

(downloaded on 16th May 2019), 200 hits per sequence were retained (*-num_alignments*). These

sequences were then parsed using an R script to exclude hits below 65% coverage, remaining

assignments with percent identity above 97% for COI were used to collapse reads for ASVs

assigned to the same species. Recent analyses have suggested that only exact (100% identity)

matching of sequences to reference data is appropriate for species assignment for the prokaryotic

16S region[46]. The 16S sequences were matched to the SILVA database (release 132)[45] using the

default settings of the *assignTaxonomy* function from the DADA2 package to assign taxonomy

at genus level or above. The incidence of NUMTs (nuclear mitochondrial DNA) and chimeras in

the final ASV list was evaluated following Supplementary Information 1.

The following quality control filters were applied to the ASV by sample table produced by

DADA2. First, the minimum number of reads per observation was set at three. Any ASVs not

represented in at least one other sample were discarded. ASVs were then filtered to retain only

those found in all three technical replicates. For any ASV found in the negative control samples,

the largest value among the control samples was used as the zero value for all other samples (i.e.

any smaller values found in non-control samples were set to zero). The COI and 18S datasets

were then subset by the RDP classifier taxonomic assignments to produce datasets for the

protists and metazoans as follows. Phylum level assignments above a threshold of 30, a value

well above that shown to accurately assign phylum level taxonomy [44], were parsed to include

phyla that contained only metazoan or protist members for each group respectively, other

assignments or unknown assignments were discarded. This resulted in a protist and metazoan

dataset for each marker, which were subsequently used for separate analyses using these

groupings. The 16S data was parsed to include only bacterial ASVs. Within each taxonomic

dataset samples were then rarefied to the smallest number of reads (see Supplementary Table 2).

Technical replicates were then collapsed to produce a dataset containing the mean value of

rarefied reads per ASV. Finally, ASVs assigned using BLAST (with no cases of multiple

matches of equal quality) to the same species in the COI dataset were combined by summing

reads per site. The taxonomic assignment method used for the 16S data assigns to genus, and

species level assignments are not possible using the selected 18S region, so no ASVs from these

datasets were collapsed. In order to explore broad scale patterns of taxonomic diversity, the

number of ASVs per phyla and number of rarefied reads per phyla were collapsed to produce per

site assessments of taxonomic composition. As phylum level phylogeny is not resolved for all

protist species, the protist dataset was grouped by supergroup designations according to Burki, et

al. [47]. For plots, phyla represented by less than 2% of ASV counts were concatenated in an

'other' category.

Abiotic, human impact and geographic data

In situ temperature data reflects a snapshot of the total conditions experienced across the lifetime

of the species that make up marine communities. Therefore, abiotic variables for the sites

covering an ecologically relevant timescale were sourced as follows. High resolution ($1km^2$)

remote sensing average daily sea surface temperature (SST) data derived from multiple satellite

missions, combined with *in situ* data[48] was parsed in R to find the nearest datapoint to each site.

For each point, a mean from two years of data from November 2017 was calculated. Interpolated

average (2005-2017) sea surface salinity (SSS) data (0.25° grid resolution) generated using

gliders, oceanographic casts etc. from the 2018 World Ocean Atlas[49] was parsed to include only

surface data for the sites. Monthly global ocean colour data ($4km^2$) derived from multiple

satellite missions[50] was parsed to calculate an average value for chlorophyll *a* density per site

across two years from November 2017. Finally, a previously described[51] $1km^2$ global resolution

cumulative index for anthropogenic impact on marine ecosystems, comprising fishing pressure,

climate change, shipping and land-based pollution, was parsed to produce a value for each site

cumulatively across the entire period for which data were available (2003-2013). These global

datasets have excellent temporal resolution, but are only appropriate for testing large-scale

patterns as they have limited ability to discriminate highly localised observations.

Ecological statistics

Differences in the mean number of ASVs per coastline were assessed using an analysis of

variance (ANOVA) after testing for normally distributed residuals using a Shapiro-Wilk test and

equal variance between coasts using a Bartlett test. A Tukey's Honest Significant difference test

was used to evaluate significant ANOVA results. Differences in community similarity were

assessed using a Permutational Multivariate Analysis of Variance (PERMANOVA)[52]

implemented in R with the function *adonis* from the package *vegan* (v2.5-6)[53] to assess

differences in multivariate centroids and dispersion between coastlines. The PERMANOVA was

conducted on a matrix of Jaccard dissimilarities as this ecological index has been shown to be

appropriate for biogeographical studies[54]. Significant pairwise differences were assessed using

the R function *adonis.pair* from the *EcolUtils* package (v0.1)[55]. To analyse if groups of samples

have a difference in intra-group community variation, also known as heterogeneity of

multivariate dispersion, the PERMDISP2 procedure[56] was used, implemented in the R function

betadisper from the *vegan* package. The pairwise group differences in heterogeneity of

multivariate dispersion in the case of a global significant result from *betadisper* were analysed

using a Tukey's Honest Significant difference test. Non-metric multidimensional scaling

ordinations (nMDS) were calculated using Jaccard dissimilarities and the R function *metaMDS*

from the *vegan* package.

The influence of the abiotic and human impact data on the observed patterns of beta diversity

were evaluated as follows. It has previously been common to use a partial Mantel test to evaluate

the effect of a distance matrix (frequently environmental variables) on a second distance matrix

(species composition) while 'cancelling out' the effect of a third matrix (geographic distance).

However, this approach has been shown to be sensitive to spatial autocorrelation common in

ecological datasets[57]. A recently developed method[57], which corrects spurious inflations of the

parameter estimate for Mantel tests, was implemented in R. Across each taxonomic group

Mantel tests were conducted comparing Jaccard dissimilarity against Euclidean distance for each

environmental variable. For each test Moran spectral randomisation was performed including the

geographic distance data with 10,000 permutations to assess statistical significance using the *msr*

function in the *adespatial* package (v0.3-8).

Explanatory variables which had some correlation with the community dissimilarity after

adjusting for geographic distance were then evaluated as follows. First, a distance-based

redundancy analysis (dbRDA)[58], regressing site Jaccard dissimilarities against all remaining

variables, was performed using the R function *dbrda* from the *vegan* package. The significance

of terms was assessed with 10,000 permutations. The dbRDA ordination allows us to examine

linear changes in the beta diversity in response to a number of predictor variables in tandem, and

also to explore their relative impact. The function *varpart* from the *vegan* package was then used

to partition the variance in the community dissimilarity by the environmental variables. We then

used a generalised additive model to visualise the variation of each significant variable across the

nMDS space via a restricted maximum likelihood 2D smoother, implemented in the function

ordisurf in the R package *vegan*.

Distance-decay relationships were explored by first measuring compositional similarity (1-

Jaccard index) for each pair of sites, and then calculating distances between pairs of sites by

drawing a continuous transect 1km offshore parallel to the high-water mark using Google Earth

Pro (v 7.3.2.5776), taking the distance along the transect to measure distance between sites. We

then used these data in least-square regression models using the R function *lm* with an interaction

function between distance and site type (artificial or natural) terms against compositional

similarity as a response term. The compositional similarity values were log10 transformed to

linearise the response, untransformed values of zero (no overlap of species) were omitted to

avoid infinite response variable values.

Results

Sequencing

A total of 66.25 million sequences were produced across the three sequencing runs. The number

of unfiltered raw reads per experimental sample ranged from 61,958 to 859,580, with an average

per sample across all three markers of $347,536 \pm 109,665$ (*s.d.*) (see Supplementary Table 2 for

further details). Negative control samples exhibited very low levels of cross-contamination

(Supplementary Information 2).

Taxonomic assignments & alpha diversity

Analyses for each taxonomic group are presented for the marker which had the largest number of

observations (COI for metazoans, 18S for protists), analyses for the remaining subsets are

presented in Supplementary Information and were consistent with the main results. After

taxonomic assignment to phyla 1,054, 1,433 and 2,826 ASVs were retained for the metazoans,

protist and bacteria datasets respectively. Across all taxonomically grouped datasets the majority

of detected ASVs came from a small number (<5) of phyla or supergroups (Figure 1b). This

pattern was consistent across sites and no major changes in the identity of ASVs at phyla or

supergroup level across the study region were detected (Figure 1b).

Across all markers the greatest mean ASV richness was found along the southern coast (Fig. 1).

However, a one-way ANOVA showed a significant difference ($F_{2,15}$=7.18, p=0.007) between

coastlines only in the bacterial dataset with no difference found in the metazoan ($F_{2,15}$=1.941,

p=0.178) or protist data ($F_{2,15}$=1.416, p=0.273). A post-hoc Tukey test of the bacterial data

(Supplementary Table 3) showed that the east and west coasts had significantly fewer ASVs

compared to the south coast, but that they were not significantly different to one another in

overall ASV richness.

Figure 1. a Map of South Africa indicating the sampling sites and the site types (red crosses are artificial
sites and blue circles natural sites). Site codes as in Supplementary Table 1; **b** Proportion of ASVs per
phyla across each site for **i** metazoans, **ii** protists and **iii** bacteria. Each bar represents a site indicated by
the site code as in Supplementary Table 1; **c** Amplicon sequence variant (ASV) richness per site separated
by coast and taxonomic group, black line indicates mean ASV richness.

Beta diversity

Across all three taxonomic groups, non-metric multidimensional ordinations showed clustering

of sites consistent with ecoregions previously described in conspicuous metazoan species (Fig.

2). Furthermore, PERMANOVA models showed a significant ($p<0.001$) effect of coastline in all

cases (see Supplementary Table 4 for model output), with pairwise significant differences

($p<0.01$) between all pairs of coastlines in all taxa (Supplementary Table 5). There was evidence

(ANOVA on *betadispr* $F_{2,15}=4.09$, $p=0.038$) for heterogeneity of multivariate dispersion in the

bacterial dataset (Supplementary Table 6). A Tukey test revealed a significant ($p=0.031$)

pairwise difference between the east and west coast only, in line with the observations of Fig. 2.,

indicating that for bacteria, sites on the west coast were more variable in community composition

than the more homogenous communities found at sites on the east coast. When the datasets were

divided by phylum, they demonstrated consistent evidence for the same significant difference

between ecoregions across all tested phylum ($p<0.05$ in all cases, see Supplementary Information

3 for full model output and visualisation). A power analysis (Supplementary Information 4)

indicated that the number of ASVs allocated to each phylum was sufficient to detect a significant

difference given the study design.

The corrected Mantel tests indicated that in the metazoan and bacterial datasets, SST and the

human impact index were significantly correlated with the observed ASV dissimilarities after

geographic distance between sites was accounted for (SST $p<0.05$ in all cases; human impact

$p<0.01$ in all cases; full model outputs shown in Supplementary Table 7). In the protist dataset

chlorophyll *a* concentration and the impact index remained significant ($p>0.05$ in both cases). In

contrast, SSS showed no correlation ($p>0.05$ in all cases) with observed ASV dissimilarities in

any marker. These results indicate that both geographic and environmental distance have some

effect on the observed community structure and confirmed that the appropriate variables were

retained for analysis in each dataset. In all cases, partial Mantel tests gave similar R statistics and

p values (Supplementary Table 7).

The dbRDA showed a significant effect of both environmental variables ($p<0.001$ for SST and

chlorophyll *a* in all cases) and human impact ($p<0.05$ in all cases) on the site similarity in

metazoans, protists and bacteria, full model outputs are presented in Supplementary Table 8.

Variance partitioning of the dbRDA models showed that human impact had a relatively smaller

contribution to the observed dissimilarities compared to the chlorophyll *a* concentration or sea

surface temperature as shown in Figure 2. Across taxonomic groups there was negligible overlap

in the variance explained by impact and chlorophyll or temperature. Generalised additive models

with a 2D smoothed function showed significant terms ($p > 0.001$) across all models (individual

full model outputs shown in Supplementary Table 9), indicating that each variable separately

explains variation in the eDNA data between communities. SST and chlorophyll *a* concentration

showed surfaces across nMDS plots for all markers (Fig. 2) that were simple, with gradients that

were consistent with ecoregions. In contrast, human impact scores showed more complex

surfaces with multiple peaks across the ecoregions.

418
Figure 2. Observed patterns of β-diversity from environmental DNA metabarcoding of: **a** metazoans, **b**
protists and **c** bacteria; based on Jaccard dissimilarities between amplicon sequence variants along the
coast of South Africa. The first column of plots shows non-metric multidimensional scaling (nMDS)
ordinations. Coloured hulls show the spread of the data and lines indicate the spread around the centroid
grouped by coast with the east, south and west coasts denoted by orange, green and blue respectively. Site
abbreviations correspond with Figure 1, natural sites are denoted with triangles and artificial sites with
filled circles. The second column of plots shows the same nMDS ordinations as the first column including
the output of a generalised additive model with a 2D smoothed function for each of the significant
environmental / impact variables overlaid; temperature – mean sea surface temperature (°C); Chlorophyll
a – chlorophyll *a* concentration (mg m^{-3}); impact – human marine impact score (unitless measurement,
see details in text) against the two nMDS axes. The Venn diagrams indicate the percentage total of the
variance in the community dissimilarity explained by each significant variable, derived using variance
partitioning of a distance-based redundancy analysis.

Distance-decay

Distance-decay slopes for all observations showed an exponential decrease in compositional

similarity as the distance between sites increased (Fig. 3a). Regression models of log10

transformed compositional similarity indicated that this slope was statistically significant in all

cases (P<0.001 for all taxonomic groups, full model output in Supplementary Information 5). In

the metazoan dataset the model showed a significant difference in the slope between artificial

and natural sites ($F_{3,76}$=47.73, p < 0.001). No statistically significant difference was found

between site types in the protist or bacteria data, and the same pattern was observed within

taxonomic groups for the metazoan and protistan subsets across both the COI and 18S data

(Supplementary Information 5).

Figure 3. Plots showing distance between sites and community similarity measured using environmental
DNA metabarcoding across South Africa. Logarithmically (base 10) transformed compositional similarity
against distance is shown in **a**, which includes all datasets. Comparisons between artificial (coloured in
red) and natural (coloured blue) sites are shown for **b** metazoans, **c** COI protists and **d** bacteria. 95%
confidence intervals from the regression models are shown as light shaded areas around each regression
slope.
450

Discussion

Here, we showed that metazoans, protists and bacteria can have similar biogeographic patterns along an extensive and heterogenous coastline. We found that these remarkably consistent patterns could be partially explained by measured environmental conditions (chlorophyll *a* and temperature), and to a lesser extent, cumulative human impact. Additionally, we found evidence for anthropogenically driven homogenisation of communities exclusively in metazoans. Collectively, we provide evidence of congruent biogeographic boundaries across vastly different forms of life, and demonstrate that underlying processes, such as anthropogenic alterations, affect biogeographic patterns differentially across taxa.

Prokaryotes and eukaryotes diverged billions of years ago and have since evolved to inhabit a vast range of ecological niches. Previous studies have shown both similar[59,60] and dissimilar[61] patterns of β diversity between macro- and microscopic species across environmental and geographic gradients. Recent work has explored biogeographic regionalisation in marine plankton across kingdoms[15], showing that smaller planktonic organisms such as bacteria may have greater biogeographic structuring compared to larger metazoans or protists[15]. Together this evidence suggests that different ecological processes drive a number of taxon-specific responses to produce patterns that are not universal across systems at different spatial scales[13]. Here we observed similar biogeographic patterns across kingdoms (Fig. 2), providing clear evidence of cross-phyla biogeographical congruence.

Our analyses suggest that environmental variables such as temperature or chlorophyll *a* concentration influenced the structure of marine communities across the study region (Fig. 2). Global studies of biogeographic patterns have shown a central role of temperature in the structuring of both microbial[12] and larger planktonic life[15,62] across the ocean. There is growing evidence that the range boundaries of marine organisms closely track their thermal limits[4]. Therefore, a general expectation was that species would remain within their thermal niche resulting in temperature-structured communities as observed here. In contrast to temperature, salinity had a minor role in structuring the studied communities (Fig. 2), an observation previously reported for the global ocean[12], with exceptions found in microbial[63] and meiofaunal[64] life in regions with unusually strong salinity gradients (e.g. Baltic Sea). The SSS range across

our study system was very narrow (35.04 - 35.38ppt) and so the negligible observed effect was

expected. In the case of protists, biogeographic patterns showed a stronger association with

primary productivity (measured here as chlorophyll *a* concentration). Previous research has

shown little or no role of productivity in driving coastal and oceanic scale biodiversity

patterns[15,62]. However, these studies explored the global role of various environmental variables;

the significance of more localised oceanographic systems such as upwelling (as along the

western coast of southern Africa) might not be as apparent in global analyses.

Anthropogenic activities are known to alter both the physico-chemical properties of the marine

environment and the trophic and ecological properties of ecosystems[65]. In our study system,

human impact provided some explanatory power to understand the observed community

structure, but to a much lesser extent compared to environmental variables (Fig. 2). The human

impact index used here[51] covered a large number of different types of impact (e.g. pollution,

shipping intensity) but even this granular approach adequately explained a small proportion of

the total variation in ASVs observed among sampling sites. Previous work on marine metazoans

has shown a strong effect of proximate urbanisation[34] and the ecological drivers produced

through anthropogenic activities are well documented[65]. Interestingly, the pervasive and

conspicuous urbanisation of the marine environment in the study area showed a much weaker

effect on biogeographic patterns than other explanatory variables (Fig. 2). Anthropogenic

pressures have become a major ecological driver only relatively recently in evolutionary time,

with the most dramatic changes in biodiversity occurring within the 21st century[3]. It is clear that

human activities are altering evolutionary trajectories[65], either through extinction, range

expansions or contractions. However, our data suggests that centuries of human impacts in our

study system have not yet demonstrably altered the main observed biogeographic boundaries

across taxa.

Previous work on biotic homogenisation has shown a dramatic effect on whole communities at

both regional[18,66] and global scales[24,25]. Here, we found evidence for biotic homogenisation along

the South African coastline only in metazoan species, with a difference in the slope of a distance-

decay relationship between artificial and natural sites. This pattern was consistent for metazoans

across the gene regions considered (Supplementary Information 5), with more homogenous

community composition between artificial sites. Pervasive vessel activity in the region[31], along

with evidence that artificial environments are hotspots for metazoan invasions[33], suggest that

introduced metazoans are contributing to homogenisation of coastal communities. Further work

should incorporate time series data to explore biotic homogenisation, given the significant but

minor role of human impact in structuring ecological communities across the region.

Both environmental parameters and species interactions have a clear-cut effect on marine

community structure across kingdoms of life[67], but the comparative role of deterministic

(environmental filtering, niche processes, etc.) and stochastic processes (ecological drift, random

extinction, etc.) in explaining the observed patterns remains uncertain. The classical

deterministic theory (Baas Becking hypothesis[68], often summarised as 'everything is

everywhere') of microbial biogeography postulates that due to vast population sizes and

dispersal, microbes are found in all environments and the variation in abiotic conditions selects

for those that make up the vast majority of species in each region. This theory ignores neutral

processes which have been shown to have a critical role in structuring microbial biogeography

across biomes[69]. In line with previous efforts studying deterministic and stochastic processes

across taxonomic kingdoms[70,71], we found that the majority of the observed variation could not

be fully explained for both prokaryotic and eukaryotic species (Fig. 2). Indeed, recent

biogeographic research in the oceans has provided both theoretical[72] and empirical[15] evidence of

strong biogeographic patterns driven by both stochastic and deterministic forces, but much of the

observed variation between communities remains unexplained. Understanding the comparative

roles of different community structuring processes requires a more comprehensive examination

of observed variance between communities, the interactions between taxa, and the broader role

of the environmental conditions where they live.

Several recent innovations will provide valuable data to help uncover the unexplained variation

in community structure. For example, the extraction and analysis of sedimentary ancient DNA

allows the reconstruction of high-resolution biodiversity change over time[73]. By rewinding the

ecological clock, the approach provides evidence to evaluate the role of deterministic processes

in reconstructions of community composition through time relative to documented changes in

environmental proxies. In conjunction, the analysis of networks from molecular data, in

conjunction with life history parameters, could facilitate the hypothesis testing of species

interactions that may further elucidate the role of ecological interactions (e.g. Djurhuus, et al. [14])

in structuring biogeographic patterns. Finally, very high-resolution multi-spectral remote sensing

data (e.g. WorldView-3, <100m) will provide unparalleled insights into the role of

environmental forces on the distribution of ecological communities[74].

Acknowledgments

LH acknowledges the assistance of Molly Czachur and Thomas Grevesse during field surveys, Sophie von der Heyden for lab consumables and assistance with fieldwork logistics, and both Jamie Hudson and Ivan Haigh for assistance with remote sensing data. LH acknowledges Shirley Parker-Nance and the Elwandle Node of the South African Environmental Observation Network for assistance and in-country logistics. All marina owners and operators are acknowledged for field site access. We acknowledge the Environmental Sequencing Facility from the National Oceanography Centre, Southampton for access and sequencing assistance. We thank the IRIDIS High Performance Computing Facility, and associated support services at the University of Southampton. LH was supported by the Natural Environmental Research Council (grant number NE/L002531/1). The UK Research and Innovation Newton Fund (grant: ES/N013913/1) supported LH's research stay in South Africa.

Data Availability

Raw Illumina sequencing data are available from the European Nucleotide Archive under study accession number PRJEB38452, sample specific accessions are provided in Supplementary Table 10. Associated metadata, R scripts and intermediate files are available at the following DOI:10.5281/zenodo.4384644.

Code Availability

All code used in the current study can be found at the following DOI:10.5281/zenodo.4384644.

References

Spalding, M. D. *et al.* Marine ecoregions of the world: A bioregionalization of coastal and shelf areas. *BioScience* **57**, 573-583, doi:10.1641/B570707 (2007).

Awad, A. A., Griffiths, C. L. & Turpie, J. K. Distribution of South African marine benthic invertebrates applied to the selection of priority conservation areas. *Diversity and Distributions* **8**, 129-145, doi:10.1046/j.1472-4642.2002.00132.x (2002).

Pecl, G. T. *et al.* Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science* **355**, doi:10.1126/science.aai9214 (2017).

Sunday, J. M., Bates, A. E. & Dulvy, N. K. Thermal tolerance and the global redistribution of animals. *Nature Climate Change* **2**, 686-690, doi:10.1038/nclimate1539 (2012).

Wallace, A. R. *The geographical distribution of animals, with a study of the relations of living and extinct faunas as elucidating the past changes of the earth's surface.* (Macmillan and Co., 1876).

Holt, B. G. *et al.* An update of Wallace's zoogeographic regions of the world. *Science* **339**, 74-78, doi:10.1126/science.1228282 (2013).

Ficetola, G. F., Mazel, F. & Thuiller, W. Global determinants of zoogeographical boundaries. *Nature Ecology and Evolution* **1**, 89, doi:10.1038/s41559-017-0089 (2017).

Kocsis, A. T., Reddin, C. J. & Kiessling, W. The stability of coastal benthic biogeography over the last 10 million years. *Global Ecology and Biogeography* **27**, 1106-1120, doi:10.1111/geb.12771 (2018).

Zaffosa, A., Finnegan, S. & Peters, S. E. Plate tectonic regulation of global marine animal diversity. *Proceedings of the National Academy of Sciences of the United States of America* **114**, 5653-5658, doi:10.1073/pnas.1702297114 (2017).

Costello, M. J. *et al.* Marine biogeographic realms and species endemicity. *Nature Communications* **8**, doi:10.1038/s41467-017-01121-2 (2017).

Beck, J. *et al.* What's on the horizon for macroecology? *Ecography* **35**, 673-683, doi:10.1111/j.1600-0587.2012.07364.x (2012).

Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean microbiome. *Science* **348**, 1261359, doi:10.1126/science.1261359 (2015).

Shade, A. *et al.* Macroecology to Unite All Life, Large and Small. *Trends in Ecology and Evolution* **33**, 731-744, doi:10.1016/j.tree.2018.08.005 (2018).

Djurhuus, A. *et al.* Environmental DNA reveals seasonal shifts and potential interactions in a marine community. *Nature Communications* **11**, 254, doi:10.1038/s41467-019-14105-1 (2020).

Richter, D. J. *et al.* Genomic evidence for global ocean plankton biogeography shaped by large-scale current systems. *bioRxiv*, 867739, doi:10.1101/867739 (2019).

Naeem, S., Duffy, J. E. & Zavaleta, E. The functions of biological diversity in an age of extinction. *Science* **336**, 1401-1406, doi:10.1126/science.1215855 (2012).

Tilman, D., Isbell, F. & Cowles, J. M. Biodiversity and Ecosystem Functioning. *Annual Review of Ecology, Evolution, and Systematics* **45**, 471-493, doi:10.1146/annurev-ecolsys-120213-091917 (2014).

Finderup Nielsen, T., Sand-Jensen, K., Dornelas, M. & Bruun, H. H. More is less: net gain in species richness, but biotic homogenization over 140 years. *Ecology Letters* **22**, 1650-1657, doi:10.1111/ele.13361 (2019).

617   19   Blowes, S. A. *et al.* The geography of biodiversity change in marine and terrestrial
618   assemblages. *Science* **366**, 339-345, doi:10.1126/science.aaw1620 (2019).
619   20   Dornelas, M. *et al.* Assemblage time series reveal biodiversity change but not systematic
620   loss. *Science* **344**, 296-299, doi:10.1126/science.1248484 (2014).
621   21   Olden, J. D. & Rooney, T. P. On defining and quantifying biotic homogenization. *Global*
622   *Ecology and Biogeography* **15**, 113-120, doi:10.1111/j.1466-822X.2006.00214.x (2006).
623   22   Stuart-Smith, R. D. *et al.* Integrating abundance and functional traits reveals new global
624   hotspots of fish diversity. *Nature* **501**, 539-542, doi:10.1038/nature12529 (2013).
625   23   Mouillot, D. *et al.* Rare species support vulnerable functions in high-diversity
626   ecosystems. *PLoS Biology* **11**, e1001569, doi:10.1371/journal.pbio.1001569 (2013).
627   24   Bernardo-Madrid, R. *et al.* Human activity is altering the world's zoogeographical
628   regions. *Ecology Letters* **22**, 1297-1305, doi:10.1111/ele.13321 (2019).
629   25   Capinha, C., Essl, F., Seebens, H., Moser, D. & Pereira, H. M. The dispersal of alien
630   species redefines biogeography in the Anthropocene. *Science* **348**, 1248-1251,
631   doi:10.1126/science.aaa8913 (2015).
632   26   Azam, F. & Malfatti, F. Microbial structuring of marine ecosystems. *Nature Reviews*
633   *Microbiology* **5**, 782-791, doi:10.1038/nrmicro1747 (2007).
634   27   Deiner, K. *et al.* Environmental DNA metabarcoding: Transforming how we survey
635   animal and plant communities. *Molecular Ecology* **26**, 5872-5895,
636   doi:10.1111/mec.14350 (2017).
637   28   Emanuel, B. P., Bustamante, R. H., Branch, G. M., Eekhout, S. & Odendaal, F. J. A
638   zoogeographic and functional approach to the selection of marine reserves on the west
639   coast of South Africa. *South African Journal of Marine Science* **12**, 341-354,
640   doi:10.2989/02577619209504710 (1992).
641   29   Griffiths, C. L., Robinson, T. B., Lange, L. & Mead, A. Marine biodiversity in South
642   Africa: an evaluation of current states of knowledge. *PLoS One* **5**, e12008,
643   doi:10.1371/journal.pone.0012008 (2010).
644   30   Griffiths, C. L. *et al.* Impacts of human activities on marine animal life in the Benguela:
645   A historical overview. *Oceanography and Marine Biology: an Annual Review* **42**, 303-
646   392, doi:10.1201/9780203507810.ch8 (2004).
647   31   Kaluza, P., Kolzsch, A., Gastner, M. T. & Blasius, B. The complex network of global
648   cargo ship movements. *Journal of The Royal Society Interface* **7**, 1093-1103,
649   doi:10.1098/rsif.2009.0495 (2010).
650   32   Rius, M. *et al.* Range expansions across ecoregions: interactions of climate change,
651   physiology and genetic diversity. *Global Ecology and Biogeography* **23**, 76-88,
652   doi:10.1111/geb.12105 (2014).
653   33   Holman, L. E. *et al.* Detection of introduced and resident marine species using
654   environmental DNA metabarcoding of sediment and water. *Scientific Reports* **9**, 11559,
655   doi:10.1038/s41598-019-47899-7 (2019).
656   34   Kelly, R. P. *et al.* Genetic signatures of ecological diversity along an urbanization
657   gradient. *PeerJ* **4**, e2444, doi:10.7717/peerj.2444 (2016).
658   35   Spens, J. *et al.* Comparison of capture and storage methods for aqueous macrobial eDNA
659   using an optimized extraction protocol: advantage of enclosed filter. *Methods in Ecology*
660   *and Evolution* **8**, 635-645, doi:10.1111/2041-210x.12683 (2017).

36 Zhan, A. *et al.* High sensitivity of 454 pyrosequencing for detection of rare species in
aquatic communities. *Methods in Ecology and Evolution* **4**, 558-565, doi:10.1111/2041-
210x.12037 (2013).
37 Leray, M. *et al.* A new versatile primer set targeting a short fragment of the
mitochondrial COI region for metabarcoding metazoan diversity: application for
characterizing coral reef fish gut contents. *Frontiers in Zoology* **10**, 34,
doi:10.1186/1742-9994-10-34 (2013).
38 Takahashi, S., Tomita, J., Nishioka, K., Hisada, T. & Nishijima, M. Development of a
prokaryotic universal primer for simultaneous analysis of Bacteria and Archaea using
next-generation sequencing. *PLoS One* **9**, e105592, doi:10.1371/journal.pone.0105592
(2014).
39 Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet Journal **17**, 10-12 (2011).
40 Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon
data. *Nature Methods* **13**, 581-583, doi:10.1038/nmeth.3869 (2016).
41 R_Core_Team. R: a language and environment for statistical computing. *ISBN 3-900051-
07-0* (2019).
42 Frøslev, T. G. *et al.* Algorithm for post-clustering curation of DNA amplicon data yields
reliable biodiversity estimates. *Nature communications* **8**, 1188, doi:10.1038/s41467-017-
01312-x (2017).
43 Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy. *Applied and
environmental microbiology* **73**, 5261-5267 (2007).
44 Porter, T. M. & Hajibabaei, M. Automated high throughput animal CO1 metabarcode
classification. *Scientific Reports* **8**, 4226, doi:10.1038/s41598-018-22505-4 (2018).
45 Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Research* **41**, D590-596,
doi:10.1093/nar/gks1219 (2013).
46 Edgar, R. C. Updating the 97% identity threshold for 16S ribosomal RNA OTUs.
Bioinformatics **34**, 2371-2375, doi:10.1093/bioinformatics/bty113 (2018).
47 Burki, F., Roger, A. J., Brown, M. W. & Simpson, A. G. The new tree of eukaryotes.
Trends in Ecology & Evolution **35**, 43-55 (2020).
48 JPL_OurOceanProject. GHRSST Level 4 G1SST Global Foundation Sea Surface
Temperature Analysis. doi:10.5067/GHG1S-4FP01 (2010).
49 Zweng, M. M., J. R. Reagan, D. Seidov, T. P. Boyer, R. A. Locarnini, H. E. Garcia, A. V.
Mishonov, O. K. Baranova, K. Weathers, C. R. Paver, and I. Smolyar,. World Ocean
Atlas 2018, Volume 2: Salinity. 50 (2019).
50 European_Space_Agency. Ocean Colour Climate Change Initiative Dataset, Version 4.2.
(2020).
51 Halpern, B. S. *et al.* Recent pace of change in human impact on the world's ocean.
Scientific Reports **9**, 11609, doi:10.1038/s41598-019-47201-9 (2019).
52 Anderson, M. J. in *Wiley Stats Ref: Statistics Reference Online* (eds N. Balakrishnan *et
al.*) 1-15 (John Wiley & Sons, Ltd, 2014).
53 Oksanen, J. *et al.* Vegan: community ecology package. *R package, Version 2.5-6* (2011).

| 705 | 54 | Kreft, H. & Jetz, W. A framework for delineating biogeographical regions based on species distributions. *Journal of Biogeography* **37**, 2029-2053, doi:10.1111/j.1365-2699.2010.02375.x (2010). |

| 708 | 55 | Salazar, G. EcolUtils: Utilities for community ecology analysis. *R package version 0.1* (2018). |

| 710 | 56 | Anderson, M. J. Distance-Based Tests for Homogeneity of Multivariate Dispersions. *Biometrics* **62**, 245-253, doi:10.1111/j.1541-0420.2005.00440.x (2006). |

| 712 | 57 | Crabot, J., Clappe, S., Dray, S. & Datry, T. Testing the Mantel statistic with a spatially-constrained permutation procedure. *Methods in Ecology and Evolution* **10**, 532-540, doi:10.1111/2041-210x.13141 (2019). |

| 715 | 58 | McArdle, B. H. & Anderson, M. J. Fitting Multivariate Models To Community Data: A Comment On Distance-Based Redundancy Analysis. *Ecology* **82**, 290-297, doi:10.1890/0012-9658(2001)082[0290:Fmmtcd]2.0.Co;2 (2001). |

| 718 | 59 | Rapacciuolo, G., Beman, J. M., Schiebelhut, L. M. & Dawson, M. N. Microbes and macro-invertebrates show parallel β-diversity but contrasting α-diversity patterns in a marine natural experiment. *Proceedings of the Royal Society B: Biological Sciences* **286**, 20190999, doi:doi:10.1098/rspb.2019.0999 (2019). |

| 722 | 60 | Astorga, A. *et al.* Distance decay of similarity in freshwater communities: do macro- and microorganisms follow the same rules? *Global Ecology and Biogeography* **21**, 365-375, doi:10.1111/j.1466-8238.2011.00681.x (2012). |

| 725 | 61 | Wang, J. *et al.* Patterns of elevational beta diversity in micro- and macroorganisms. *Global Ecology and Biogeography* **21**, 743-750, doi:10.1111/j.1466-8238.2011.00718.x (2012). |

| 728 | 62 | Tittensor, D. P. *et al.* Global patterns and predictors of marine biodiversity across taxa. *Nature* **466**, 1098-U1107, doi:10.1038/nature09329 (2010). |

| 730 | 63 | Herlemann, D. P. *et al.* Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME Journal* **5**, 1571-1579, doi:10.1038/ismej.2011.41 (2011). |

| 733 | 64 | Broman, E. *et al.* Salinity drives meiofaunal community structure dynamics across the Baltic ecosystem. *Molecular Ecology* **28**, 3813-3829, doi:10.1111/mec.15179 (2019). |

| 735 | 65 | Shochat, E., Warren, P. S., Faeth, S. H., McIntyre, N. E. & Hope, D. From patterns to emerging processes in mechanistic urban ecology. *Trends in Ecology and Evolution* **21**, 186-191, doi:10.1016/j.tree.2005.11.019 (2006). |

| 738 | 66 | Blouin, D., Pellerin, S. & Poulin, M. Increase in non-native species richness leads to biotic homogenization in vacant lots of a highly urbanized landscape. *Urban Ecosystems* **22**, 879-892, doi:10.1007/s11252-019-00863-9 (2019). |

| 741 | 67 | Lima-Mendez, G. *et al.* Determinants of community structure in the global plankton interactome. *Science* **348**, 1262073, doi:10.1126/science.1262073 (2015). |

| 743 | 68 | Baas-Becking, L. G. M. *Geobiologie; of inleiding tot de milieukunde.* (WP Van Stockum & Zoon NV, 1934). |

| 745 | 69 | Hanson, C. A., Fuhrman, J. A., Horner-Devine, M. C. & Martiny, J. B. Beyond biogeographic patterns: processes shaping the microbial landscape. *Nature Reviews Microbiology* **10**, 497-506, doi:10.1038/nrmicro2795 (2012). |

| 748 | 70 | Farjalla, V. F. *et al.* Ecological determinism increases with organism size. *Ecology* **93**, 1752-1759, doi:10.1890/11-1144.1 (2012). |

750 71   Wu, W. X. *et al.* Contrasting the relative importance of species sorting and dispersal
751   limitation in shaping marine bacterial versus protist communities. *ISME Journal* **12**, 485-
752   494, doi:10.1038/ismej.2017.183 (2018).
753 72   Hellweger, F. L., van Sebille, E. & Fredrick, N. D. Biogeographic patterns in ocean
754   microbes emerge in a neutral agent-based model. *Science* **345**, 1346-1349,
755   doi:10.1126/science.1254421 (2014).
756 73   Balint, M. *et al.* Environmental DNA time series in ecology. *Trends in Ecology and*
757   *Evolution* **33**, 945-957, doi:10.1016/j.tree.2018.09.003 (2018).
758 74   He, K. S. *et al.* Will remote sensing shape the next generation of species distribution
759   models? *Remote Sensing in Ecology and Conservation* **1**, 4-18, doi:10.1002/rse2.7
760   (2015).
761
762