Consistent marine biogeographic boundaries across the tree of life despite centuries of human impacts

Luke E. Holman[1,*] (ORCID: 0000-0002-8139-3760)

Mark de Bruyn[2,3] (ORCID:0000-0003-1528-9604)

Simon Creer[2]

Gary Carvalho[2]

Julie Robidart[4] (ORCID: 0000-0001-9805-3570)

Marc Rius[1,5] (ORCID: 0000-0002-2195-6605)

[1] School of Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton, United Kingdom

[2] Molecular Ecology and Fisheries Genetics Laboratory, School of Natural Sciences, Bangor University, United Kingdom

[3] The University of Sydney, School of Life and Environmental Sciences, Australia

[4] Ocean Technology and Engineering Group, National Oceanography Centre Southampton, United Kingdom

[5] Department of Zoology, Centre for Ecological Genomics and Wildlife Conservation, University of Johannesburg, South Africa

* Corresponding author: Lukeearlholman@gmail.com

Abstract

Over millennia ecological and evolutionary mechanisms have shaped macroecological distributions across the tree of life. Research describing patterns of regional and global biogeography has traditionally focussed on the study of conspicuous species. Consequently, there is limited understanding of cross-phyla biogeographical structuring, and an escalating need to understand the macroecology of macrobial and microbial life. Here we used environmental DNA metabarcoding to explore the biodiversity of marine metazoans, micro-eukaryotes and prokaryotes along an extensive and highly heterogeneous coastline. We expected environmental forces to be responsible for creating similar biogeographical assemblages across kingdoms of life, while life-history strategies and pervasive anthropogenic impacts were predicted to disturb such patterns. We found that despite centuries of significant human activities, biogeographic patterns were consistent across kingdoms, underpinned primarily by abiotic factors. Our work highlights the importance of studying multiple domains of life to understand the maintenance of biodiversity and potential drivers of future change.

Introduction

Researchers have long recognised the importance of grouping global biota into distinct, geographically separated regions. Delineating these biogeographic areas is important to understand the factors shaping the range limits of species[1], to designate key areas for biodiversity conservation[2] and in formulating predictive responses to environmental change[3,4]. One of the first efforts to define geographic regions of terrestrial biota were Alfred Russel Wallace's so-called 'Zoological Regions'[5], which included six major regions (hereafter realms) that are still recognised in subsequent classifications today[6]. The drivers responsible for these geographical classifications are frequently environmental conditions or physical barriers. For example, broad tectonic dynamics can produce waterbodies or mountains, and dramatically different climatic conditions are known to define biogeographic boundaries and shape species distributions. Studies have shown that deep divergence in the geographic arrangement of terrestrial biota arose as a result of plate tectonics, while shallow divergence has been most frequently attributed to climate[7]. In aquatic ecosystems, the relative importance of macroecological drivers is less understood, although both climatic (e.g. temperature)[8] and tectonic forces[9] have been identified as key determinants of biogeographic patterns. Recent studies have successfully partitioned the oceans into distinct ecoregions (i.e. a geographically defined area, smaller than realm, that contains characteristic assemblages of species)[1,10], but the description of marine ecoregions has mostly considered conspicuous or well-described species. Similarly, most macroecological marine research has focussed on easily identifiable eukaryotic species, principally metazoans[11], although some progress has been made in understanding global patterns of marine microbes[12]. In line with recent studies demonstrating strong cross-phylum interdependence[13], there is an increasing need to include prokaryotic species in our assessment of biogeographic patterns. The language of macroecology and microbial ecology is similar, both examining the incidence of species across different spatial scales, but these fields have long progressed independently. As a result, only a few studies have simultaneously explored biogeographic patterns of both microscopic and macroscopic life to date[13,14], and to our knowledge no study has compared broad-scale biogeographic patterns across different kingdoms of life.

Human activities such as habitat destruction, pollution and the introduction of non-native species
are the main drivers of more recent global biodiversity change[3] and therefore have the potential
to alter geographic patterns of biota at multiple spatial scales. Cumulatively anthropogenic
stressors not only threaten vulnerable native species but whole-community structure and
function[3,15,16]. The magnitude and direction of human impacts is complex, with evidence for both
gains and losses in local species richness across biomes[17-19]. However, a consistent global pattern
is emerging, with a recent and rapid increase in species turnover[18] and an associated increase in
community similarity (β diversity) between two or more geographically separated sites[19]. These
incidences of increased community similarity are known as biotic homogenisation[20] and are
driven by human activities that promote extinctions and the introductions of non-native species.
In light of growing evidence that taxonomic, phylogenetic and functional diversity are strongly
correlated[21], the homogenisation of biological communities has the potential to negatively affect
ecosystem function. Furthermore, even uncommon species within an ecological community can
contribute significantly to ecosystem function[22], demonstrating the importance of studying
inconspicuous species to preserve ecosystems. Cumulatively, biotic homogenisation directly
disrupts biogeographic patterns, threatens endemism at both local and regional scales, and has
the potential to affect ecosystem function. Studies have shown evidence for biotic
homogenisation around the globe, with examples from plants[17], vertebrates[23] and invertebrates[24]
demonstrating alteration of terrestrial biogeographic patterns. While the overall magnitude and
direction of this effect is becoming clear, many studies are of limited taxonomic scope, focussing
on highly conspicuous species for which reliable data can be relatively easily produced[23,24].
Thus, most studies overlook inconspicuous species (e.g. microbes and microscopic metazoans),
which show vastly different reproductive, demographic and dispersal patterns compared to
eukaryotes[13], but are known to be key actors shaping the assembly of ecological communities
and ultimately underpin ecosystem functioning[25]. Taken together, a more comprehensive
characterisation of ecological communities is clearly needed when testing the role of biotic
homogenisation on biogeographic patterns.
98
The advent of high-throughput sequencing has revolutionised our understanding of microbial
life, with studies examining global patterns of prokaryotic life now increasingly common[12].
Moreover, the recent and rapid development of methods to infer the incidence of larger

organisms using genetic material isolated from environmental samples (known as environmental
DNA or eDNA) has provided an unparalleled ability to identify species across the entire tree of
life[26,27]. Together, these methods can rapidly generate standardised biodiversity data for entire
communities at an unprecedented resolution, thereby minimising regional and taxonomic biases.
Such studies provide datasets that can be analysed without taxonomic assignment and DNA
samples can be repurposed to test novel hypotheses. A common technique is to amplify DNA
barcodes from eDNA and use high-throughput sequencing to produce high-resolution
biodiversity data. This method (eDNA metabarcoding) has been shown to reliably detect
organisms across many different ecosystems[26], but has infrequently been applied to understand
spatial patterns of biodiversity across different kingdoms of life[12,28].

The South African coastline has three well-defined coastal ecoregions bounded by the cold
western boundary Benguela Current and the warm oligotrophic eastern boundary Agulhas
Current. These ecoregions have been delineated by the presence of a number of conspicuous
metazoan taxa[2,29] established over decades of study. Additionally, there is evidence for human
exploitation of marine resources in the region spanning thousands of years[30,31] and the coastline
has been an area of high maritime activity for centuries[32]. Other human activities have also been
prevalent such as the establishment of aquaculture facilities or the construction of harbours and
breakwaters[30,31]. Thus, this coastline is an ideal study system to explore the mechanisms shaping
biogeographic boundaries.

Here we used eDNA metabarcoding to examine the biogeography of multiple marine domains
along the ecologically diverse coastline of South Africa. We hypothesised that environmental
conditions would be responsible for the maintenance of clear and congruent cross-phyla
biogeographic boundaries. In turn, life-history strategies were expected to reshape the
geographic arrangement of different phyla. More specifically, we expected that microbial
organisms would show a more homogeneous distribution along the studied coastline compared to
eukaryotes, mainly due to the greater dispersal capacity of the former. Finally, in line with
previous observations, we expected that long-term anthropogenic impacts would disrupt well-
defined ecoregion boundaries and lead to an overall homogenisation of biogeographic patterns
across the kingdoms of life.

Methods

Field sampling

We sampled a range of sites along 2,000 km of coastline (Fig. 1) between October and November of 2017 (see details in Supplementary Table 1), representing the three major marine ecoregions. In order to assess the effects of anthropogenic impacts, we compared human altered 'artificial' sites (e.g. recreational marinas, harbours) and relatively unaltered rocky shore sites, hereafter 'natural' (see Supplementary Table 1). The artificial sites were previously sampled in Rius, et al. [33], and six adjacent natural rocky shore sites were selected, choosing the nearest site with matching aspect and exposure (Fig. 1) to an artificial site wherever possible. For each site, three 400 ml seawater samples were filtered with 0.22 μm polyethersulfone membrane Sterivex filters (Merck Millipore, MA, USA) following the sampling scheme of Holman, et al. [34]. Consequently, we sampled a total of 1,200 ml of seawater per site, a volume that has been shown to differentiate fine scale (<1 km^2) community structure in marine systems[34,35]. Filters were immediately preserved at ambient temperature with the addition of 1.5 ml of Longmire's Solution for preservation until DNA extraction. Field control filters and equipment cleaning blanks were taken, transported, stored and sequenced as the rest of the field samples.

Environmental DNA extraction

We used a PCR-free laboratory separated from the main molecular biology laboratory facilities. No post-PCR or high concentration DNA samples were permitted in the laboratory. All surfaces and lab equipment were cleaned thoroughly before use with 1.25% sodium hypochlorite solution (3:1 dilution of household bleach). DNA extraction followed the SX^{CAPSULE} method from Spens, et al. [36]. Briefly, filters were first externally cleaned with sterile water and Longmire's Solution was removed from the filter outlet using a sterile syringe, 720 μl Buffer ATL (Qiagen, Hilden, Germany) and 80 μl Proteinase K (20mg/ml) was added and filters were incubated overnight at 56°C. The lysate was then removed from the filter inlet and subjected to DNA extraction using the Qiagen DNeasy Blood and Tissue Kit under the manufacturers recommended protocol. DNA was eluted using 200 μl Qiagen Buffer AE and re-eluted once to increase DNA yield. All DNA samples were checked for PCR inhibition using the Primer Design Internal Positive Control qPCR Kit (Primer Design, Southampton, UK) with 10 μl reactions under the manufacturer

recommended protocol. Inhibition was detected by an increase of >1.0 Ct in reactions spiked

with eDNA compared to unspiked reactions with extraction controls. As inhibition was detected

in a minority of samples, all samples were treated using the Zymo OneStep PCR Inhibition

Removal Kit (Zymo Research, California, USA) following the manufacturer recommended

protocol. Inhibited samples showed no evidence for inhibition post cleaning.

High throughput eDNA amplicon sequencing

Three sets of primers were used to generate three separate eDNA metabarcoding libraries for all

samples. Two gene regions were selected to target broad metazoan/eukaryotic diversity, a section

of the V4 region of the nuclear small subunit ribosomal DNA (hereafter 18S)[37] and a section of

the standard DNA barcoding region of cytochrome c oxidase subunit I (hereafter COI)[38]. One

gene region was used to target Prokaryotes, the V3-V4 hypervariable region of prokaryotic small

subunit ribosomal DNA (hereafter 16S)[39]. Illumina double-indexed metabarcoding amplicon

libraries were constructed with a two-step PCR protocol as detailed in Holman, et al. [34]. The first

PCR was performed in a PCR-free laboratory. The three eDNA samples per site were pooled and

three independent technical replicates were sequenced per pool. The process per sequenced pool

was as follows. The first PCR reaction was conducted in triplicate in a total reaction volume of

20 μl. Each reaction contained 10 μl Amplitaq GOLD 360 2X Mastermix (Applied Biosystems,

California, USA), 0.8 μl (5 nmol ml^{-1}) of each forward and reverse primers and 2 μl of undiluted

environmental DNA template. The reaction conditions for PCR were an initial denaturation step

at 95°C for 10 minutes followed by 20 cycles of 95°C for 30 seconds, variable annealing temp

(46°C for COI, 50°C for 18S and 55°C for 16S) for 30 seconds, and extension at 72°C for 1

minute. A final extension at 72°C was performed for 10 minutes. The triplicate first PCR

replicates were pooled and cleaned using AMPure XP beads (Beckman Coulter, California,

USA) at 0.8 beads:sample volume ratio following manufacturer's instructions. The second PCR

reaction was conducted in a total volume of 20 μl containing 10 μl Amplitaq GOLD 360 2X

Mastermix, 0.5 μl (10 nmol ml^{-1}) of both forward and reverse primers and 5 μl of undiluted

cleaned PCR product from the first reaction. PCR conditions were an initial denaturation step at

95°C for 10 minutes followed by 15 cycles of 95°C for 30 seconds, annealing at 55°C for 30

seconds, and extension at 72°C for 1 minute. A final extension at 72°C was performed for 10

minutes. PCR 2 products were cleaned using AMpure XP beads as above. Negative control

samples for the filters, extraction kit, PCR1 and 2 were included in library building and

sequenced alongside experimental samples. Products were quantified following the

manufacturer's instructions using the NEBNext Library Quant qPCR kit (New England Biolabs,

Massachusetts, USA) and then normalised and pooled at an equimolar concentration for each

marker. Each gene region was sequenced independently using a V3 paired-end 300bp reagent kit

on the Illumina MiSeq Instrument with 5% PhiX genomic library added to increase sequence

diversity.

Bioinformatics

Raw sequences were de-multiplexed using the GenerateFastQ (v 2.0.0.9) module on the MiSeq

control software (v 3.0.0.105). Cutadapt (v 2.3)[40] was used to filter sequences to include only

those that contained both the forward and reverse primer sequence across both read pairs for

each gene fragment, remaining sequences then had the primer region removed for each gene

fragment using the default settings. Sequences were denoised using the DADA2 pipeline (v

1.12)[41] in R (v 3.6.1)[42] with the default parameters unless noted as follows. Sequences were

filtered to retain only pairs of reads with an expected error of 1 or fewer per read. Read trimming

was performed after manual examination of the read quality profile, the forward reads were

trimmed to 250bp (COI), 240bp (18S) and 240 bp (16S) and the reverse reads were trimmed to

230bp (COI), 220bp (18S) and 220 bp (16S). As each marker was sequenced separately, the

differences in read trimming length reflect typical variation in sequencing runs rather than any

biological difference. The error rates per run were estimated and used to perform the denoising

using the DADA2 algorithm. The denoised sequence pairs were then merged and resulting

sequences were truncated if they were outside of the expected gene fragment range (303-323bp

for COI, 400-450bp for 18S and 390-450bp for 16S). Chimeras were identified and removed

before assembling a sample by ASV (amplicon sequence variant) table for analysis. The

denoised ASVs were then curated using the default settings of the LULU algorithm[43] which

merges sequences based on sequence similarity and co-occurrence. Assigning taxonomy to a set

of unknown sequences is a difficult task, particularly considering many marine species lack

DNA barcodes, are undescribed, or have erroneous barcodes in online public databases. We

therefore focused our analysis at a higher taxonomic level than species, assigning taxonomy to

sequences from the COI and 18S runs as follows: an unconstrained (no limits on sequence

similar or match length) BLAST search (v 2.6.0+) was performed for each sequence against the entire nt database (downloaded on 16th May 2019), 200 hits per sequence were retained (-*num_alignments*). These sequences were then parsed using an R script as follows. All hits below 65% coverage and 85% percentage identity were removed, remaining sequences with percent identity above 97% for COI and 99% for 18S were marked as 'high confidence'. Remaining sequences were marked 'low confidence'. For all remaining sequences with conflicting (non-identical) taxonomic assignments an R script selected the lowest common ancestor (broadest taxonomic lineage) that covered all assignments and assigned this taxa to the sequence. The NCBI taxonomy (downloaded 16th May 2019) was used for the lowest common ancestor and the *NCBItax2lin* utility from Mahmoudabadi and Phillips [44] was used to convert NCBI taxonomic IDs into lineages. Recent analyses have suggested that only exact (100% identity) matching of sequences to reference data is appropriate for species assignment for the prokaryotic 16S region[45]. The 16S sequences were matched to the SILVA database (release 132)[46] using the default settings of the *assignTaxonomy* function from the DADA2 package to assign taxonomy at genus level or above. The incidence of NUMTs (nuclear mitochondrial DNA) and chimeras in the final ASV list was evaluated following Supplementary Information 1.

Abiotic, human impact and geographic data

In situ temperature data reflects a snapshot of the total conditions experienced across the lifetime of the species that make up marine communities. Therefore, abiotic variables for the sites covering an ecologically relevant timescale were sourced as follows. High resolution (1 km^2) remote sensing average daily sea surface temperature (SST) data derived from multiple satellite missions combined with *in situ* data[47] was parsed in R to find the nearest datapoint to each site. For each point a mean from two years of data from November 2017 was calculated. Interpolated average (2005-2017) sea surface salinity (SSS) data (0.25° grid resolution) generated using gliders, oceanographic casts etc. from the 2018 World Ocean Atlas[48] was parsed to include only surface data for the sites. Monthly global ocean colour data (4km^2) derived from multiple satellite missions[49] was parsed to calculate an average value for chlorophyll a density per site across two years from November 2017. Finally, a previously described[50] 1km^2 global resolution cumulative index for anthropogenic impact on marine ecosystems, comprising fishing pressure, climate change, shipping and land based pollution, was parsed to produce a value for each site

cumulatively across the entire period for which data were available (2003-2013). These global

datasets have excellent temporal resolution, but are only appropriate for testing large-scale

patterns as they have a limited ability to discriminate highly localised observations. Distances

between pairs of sites was estimated by drawing a continuous transect 1km offshore parallel to

the high water mark using Google Earth Pro (v 7.3.2.5776), and taking the distance along the

transect to measure distance between sites.

Ecological statistics

The following quality control filters were applied to the ASV by sample table produced by

DADA2. First, the minimum number of reads per observation was set at 3. Any ASVs not

represented in at least one other sample were discarded. ASVs were then filtered to retain only

those found in all three technical replicates. For any ASV found in the negative control samples,

the largest value among the control samples was used as the zero value for all other samples (i.e.

any smaller values found in non-control samples were set to zero). Samples were then rarefied to

the smallest number of reads found within each gene fragment (see Supplementary Table 2).

Technical replicates were then collapsed to produce a dataset containing the mean value of

rarefied reads per ASV. Finally, ASVs assigned with high confidence (and no cases of multiple

matches of equal quality) to the same species in both the COI and the 18S datasets were

combined. The taxonomic assignment method used for the 16S data assigns to genus, so no

ASVs from the 16S data were collapsed. In order to explore broad scale patterns and avoid

difficulties with taxonomic assignment, the number of ASVs per phyla and number of rarefied

reads per phyla were collapsed to produce per site assessments of taxonomic composition. For

plots phyla represented by less than 2% of ASV counts were concatenated in an 'other' category.

Differences in the mean number of ASVs per coastline were assessed using an analysis of

variance (ANOVA) after testing for normally distributed residuals using a Shapiro-Wilk test and

equal variance between coasts using a Bartlett test. A Tukey's Honest Significant difference test

was used to evaluate significant ANOVA results. Differences in community similarity were

assessed using a Permutational Multivariate Analysis of Variance (PERMANOVA)[51]

implemented in R with the function *adonis* from the package *vegan* (v 2.5-6)[52] to assess

differences in multivariate centroids and dispersion between coastlines. The PERMANOVA was

conducted on a matrix of Jaccard dissimilarities as this ecological index has been shown to be

appropriate for biogeographical studies[53]. Significant pairwise differences were assessed using

the R function *pairwise.adonis* from the *pairwiseAdonis* package (v.0.3)[54]. To analyse if groups

of samples have a difference in intra-group community variation, also known as heterogeneity of

multivariate dispersion, the PERMDISP2 procedure[55] was used, implemented in the R function

betadisper from the *vegan* package. The pairwise group differences in heterogeneity of

multivariate dispersion in the case of a global significant result from *betadisper* were analysed

using a Tukey's Honest Significant difference test. Non-metric multidimensional scaling

ordinations (nMDS) were calculated using Jaccard dissimilarities and the R function *metaMDS*

from the *vegan* package.

Two approaches were used to explore the influence of the abiotic and human impact data on the

observed patterns of beta diversity. First, a distance-based redundancy analysis (dbRDA)[56],

regressing site Jaccard dissimilarities against SST, SSS, chlorophyll a and human impact, was

performed using the R function *dbrda* from the *vegan* package. The significance of terms was

assessed with 10,000 permutations. Subsequently dbRDA models were constructed with every

combination of independent variables for models with three, two or a single variable for a total

of 15 models (including the full model). The adjusted R^2 values for each model were extracted to

show the unique information provided by each variable. The dbRDA ordination allows us to

examine linear changes in the beta diversity in response to a number of predictor variables in

tandem, and also to explore their relative impact. In addition, we used a generalised additive

model to assess the significance of each independent variable in explaining the variation in the

two non-metric multidimensional axes from above using a restricted maximum likelihood 2D

smoother, implemented in the function *ordisurf* in the R package *vegan*.

It has previously been common to use a partial Mantel test to evaluate the effect of a distance

matrix (frequently environmental variables) on a second distance matrix (species composition)

while 'cancelling out' the effect of a third matrix (geographic distance). However, this approach

has been shown to be sensitive to spatial autocorrelation common in ecological datasets[57]. A

recently developed method[57], which corrects spurious inflations of the parameter estimate for

Mantel tests, was implemented in R. Across each marker Mantel tests were conducted comparing

Jaccard dissimilarity against Euclidean distance for each environmental variable. For each test

Moran spectral randomisation was performed including the geographic distance data with 10,000

permutations to assess statistical significance using the *msr* function in the *adespatial* package

(v.0.3-8).

324
325   **Results**
326

Sequencing

A total of 66.25 million sequences were produced across the three sequencing runs. The number

of unfiltered raw reads per experimental sample ranged from 61,958 to 859,580, with an average

per sample across all three markers of $347,536 \pm 109,665$ (*s.d.*) (see Supplementary Table 2 for

further details). Negative control samples exhibited very low levels of cross-contamination

(Supplementary Information 2).

Alpha diversity and taxonomy

Across all markers the greatest mean ASV richness was found along the southern coast (Fig. 1).

However, a one-way ANOVA showed a significant difference ($F_{2,15}=7.18$, $p=0.007$) between

coastlines only in the 16S data with no difference found in the COI ($F_{2,15}=3.64$, $p=0.051$) or 18S

data ($F_{2,15}=1.82$, $p=0.195$). A post hoc Tukey test of the 16S data (Supplementary Table 3)

showed that the east and west coasts had significantly less ASVs compared to the south coast,

but that they were not significantly different to one another in terms of overall richness. The

taxonomic composition, measured by ASV richness per phyla, was consistent across sites and

markers (Fig. 1c). In contrast, the proportion of reads assigned to each phylum exhibited more

variation across sites (Supplementary Fig. 1) in the COI and 18S data, although the 16S data

were more even.

346
Figure 1. a Amplicon sequence variant (ASV) richness per site partitioned by coast (west = blue, south =
green, east = orange) from environmental DNA metabarcoding of COI, 18S and 16S markers, black line
indicates mean ASV richness; **b** Map of South Africa indicating the sampling sites and the site types (red
crosses are artificial sites and blue circles natural sites). Site codes as in Supplementary Table 1; **c**
Proportion of ASVs per phyla across each site for **i** COI, **ii** 18S and **iii** 16S. Each bar represents a site
indicated by the site code as in Supplementary Table 1.

Beta diversity

Across all three markers, the non-metric multidimensional ordinations showed clustering of sites consistent with ecoregions previously described in conspicuous metazoan species (Fig. 2). Furthermore, PERMANOVA modelling showed a significant (p<0.001) effect of coastline in all cases (see Supplementary Table 4 for model output), with pairwise significant differences (p<0.05) between all pairs of coastlines in all markers (Supplementary Table 5). There was evidence (ANOVA on *betadispr* $F_{2,15}$=3.94, p=0.042) for heterogeneity of multivariate dispersion in the 16S dataset (Supplementary Table 6). A Tukey test revealed a significant (p=0.036) pairwise difference between the east and west coast only, in line with the observations of Figure 2 with a large observed difference in multivariate spread between these coasts in the 16S data.

The dbRDA showed a significant global effect of the abiotic and human impact variables on the site similarity in COI ($F_{4,16}$=2.277, p<0.001), 18S ($F_{4,16}$=1.987, p<0.001) and 16S ($F_{4,16}$=2.456, p<0.001). Testing individual variables for statistical significance by permutation revealed an (p<0.05) effect in all variables across all markers except for human impact in the 16S dataset (p=0.052), full model outputs are presented in Supplementary Table 7. Decreasing the number of variables in the model decreased the proportion of explained variance in the ASV dissimilarity as shown in Figure 2. Across all markers, SST explained the most variance in the models (as measured by R^2 values in single term models) followed by SSS then chlorophyll a concentration, with the human impact index providing the lowest amount of explanatory information in models. Across all three markers, the model with all variables explained more variation than any model with three variables, indicating that all variables provided unique information. Generalised additive models with a 2D smoothed function showed significant terms (p>0.001) across all models (individual full model outputs shown in Supplementary Table 8), indicating that each variable separately explains variation in the eDNA data between communities. SST, SSS and chlorophyll a concentration showed surfaces across nMDS plots for all markers (Fig. 2) that were simple, with gradients across the plot consistent with ecoregions. In contrast, human impact scores showed more complex surfaces with multiple peaks across the ecoregions.

The corrected Mantel tests indicated that across all markers, SST and the human impact index
were significantly correlated with the observed ASV dissimilarities after geographic distance
between sites was accounted for (SST $p<0.05$ in all cases; human impact $p<0.01$ in all cases; full
model outputs shown in Supplementary Table 9). In contrast, SSS and chlorophyll a
concentration showed no correlation ($p>0.05$ in all cases) with observed ASV dissimilarities in
any marker. These results indicate that both geographic and environmental distance have some
effect on the observed community structure, but that out of the tested variables only temperature
and the human impact index have a substantial effect after geographic distance between sites has
been accounted for.
393

394
Figure 2. Observed patterns of β-diversity from environmental DNA metabarcoding of: **a** COI, **b** 18S and
c 16S; based on Jaccard dissimilarities between amplicon sequence variants along the coast of South
Africa. The first column of plots show non-metric multidimensional scaling (nMDS) ordinations.
Coloured ellipses indicate the 95% confidence interval around the centroid and are grouped by coast with
the east, south and west coasts denoted by orange, green and blue respectively. Site abbreviations
correspond with Figure 1, natural sites are denoted with triangles and artificial sites with filled circles.
The second column of plots show the same nMDS ordination as the first column including the output of a
generalised additive model with a 2D smoothed function for each of the environmental / impact variables
overlaid; SST – mean sea surface temperature (°C); SSS – mean sea surface salinity (parts per thousand);
Chl a – chlorophyll a concentration (mg m^{-3}); impact – human marine impact score (unitless
measurement, see details in text) against the two nMDS axes. The last column of plots depicts upset plots
showing the variance (%) in Jaccard dissimilarities among sites, explained by different terms using a
distance-based redundancy analysis. Each bar represents a model with the terms in the model indicated by
filled dots with the most complex model on the left and the least on the right.

Composition-distance-decay relationship

Distance-decay slopes for all observations showed an exponential decrease in compositional similarity as the distance between sites increased (Fig. 3). Regression models of log10 transformed compositional similarity indicated that this slope was statistically significant in all cases (P<0.005 for all markers, full model output in Supplementary Table 10). In the COI data the model showed a significant difference in both the slope and intercept between artificial and natural sites ($F_{3,77}$=77.92, additive p=0.018, interactive p=0.004, Supplementary Table 10 for full model). No statistically significant difference was found between site types in the 18S or 16S data. As shown in Figure 3b-d, there is much greater variance in the residuals for the 18S and 16S data compared to the COI data and the R^2 values from the models support these observations (COI R^2=0.730, 18S R^2=0.224, 16S R^2=0.473).

Figure 3. Plots showing distance between sites and community dissimilarity measured by environmental DNA metabarcoding across South Africa. a) Untransformed data for all gene regions. Logarithmically (base 10) transformed compositional similarity against distance is shown for b) COI, c) 18S and d) 16S data with comparisons between artificial sites coloured red and natural sites coloured blue. 95% confidence intervals from the least-squares linear regression models are shown as light shaded areas around each regression slope.

Phylum level analyses

Removing phyla with fewer than 20 ASVs resulted in 9, 11 and 6 Phyla remaining for the COI,

18S and 16S data respectively with a mean of 152 ASVs per phyla. All but two phyla showed a

significant difference between coastlines in PERMANOVA models, with *Chordata* and

Cryptophyta in the 18S data having no global statistical difference between coasts. In contrast

only three phyla showed a difference in the intercept or slope between artificial and natural sites.

Regression models for the *Bacillariophyta* from the COI data and *Epsilonbacteraeota* from the

16S data showed a significant ($p < 0.05$) difference in intercept and the *Proteobacteria* from the

16S data showed a difference in both the intercept and slope. Full model outputs for both

PERMANOVA and regression models are shown in Supplementary Table 11.

Discussion

Here, we showed that ecoregions are consistently delimited across the tree of life, with both

eukaryotes and prokaryotes showing similar biogeographic patterns along an extensive and

heterogenous coastline. On one hand, these observations were in line with our initial

expectations that abiotic factors would lead to the maintenance of the heterogeneity of large-

scale biodiversity patterns, and our findings clearly showed that seawater temperature was a

particularly important factor. On the other hand, our expectation of a more homogeneous

distribution along the studied coastline for microbial species was not met; suggesting that abiotic

forces are more important drivers than dispersal ability and other life-history characteristics of

niche occupancy. Finally, we expected a disrupting effect of anthropogenic impact on species

distributions, and although we found that it played a role in shaping the geographic arrangement

of some taxa reflected in the COI dataset, our results suggested limited effect of biotic

homogenisation (see comparisons between natural and artificial sites in Fig. 3). These findings

contradicted our expectations but highlighted the inherent difficulties of predicting the effects of

human impact. In addition, they showed that although anthropogenic impacts in ecosystems can

be pervasive, they may not be the central force structuring biogeographic patterns along

extensive coastlines.

Prokaryotes and eukaryotes diverged billions of years ago and have since evolved to inhabit a

vast range of ecological niches. Previous work has shown both similar[58,59] and dissimilar[60]

patterns of β diversity between macro- and microscopic species across environmental and

geographic gradients. This evidence suggests that different ecological processes drive a number

of taxon-specific responses to produce patterns that are not universal across systems at different

spatial scales[13]. Indeed evidence from metazoan species demonstrates that biogeography can

differ among different classes[6], which is a relatively shallow taxonomic level. Therefore, lack of

any universal biogeographic structure across taxa should be considered the null hypothesis for

the study of any system, particularly in the ocean where there are comparatively fewer barriers to

dispersal and migration. In our study, we observed remarkably similar biogeographic patterns

across kingdoms (Fig. 2), rejecting the above null hypothesis, and providing an example of

cross-phyla biogeographic congruence.

Our analyses suggested that temperature was the primary environmental variable structuring

marine communities across the study region (Fig. 2). In line with this, global studies of

biogeographic patterns have shown a central role of temperature in the structuring of both

microbial[12] and larger planktonic life[28,61] across the ocean. There is growing evidence that the

range boundaries of marine organisms closely track their thermal limits[4]. Therefore, a general

expectation was that regardless of taxonomic kingdom, species would remain within their

thermal niche resulting in temperature-structured communities as observed here. In contrast to

temperature, salinity had a minor role in structuring the studied communities (Fig. 2), an

observation previously reported for the global ocean[12], with exceptions found in microbial[62] and

meiofaunal[63] life in regions with unusually strong salinity gradients (e.g. Baltic Sea). The SSS

range across our study system was very narrow (35.04 - 35.38ppt) and so the negligible observed

effect was expected. Similarly, primary productivity (measured here as chlorophyll a

concentration) showed little effect in structuring biogeographic patterns (Fig. 1), in line with

previous research showing little or no role of productivity in driving coastal and oceanic scale

biodiversity patterns[28,61].

Anthropogenic activities are known to alter both the physico-chemical properties of the marine

environment and the trophic and ecological properties of ecosystems[64]. In our study system,

human impact provided some explanatory power to understand the observed community
structure, but to a much lesser extent compared to temperature (Fig. 2). The human impact index
used here[50] covered a large number of different types of impact (e.g. pollution, shipping
intensity) but even this granular approach explained the variation in ASVs observed among
sampling sites. Previous work on marine metazoans has shown a strong effect of proximate
urbanisation[35] and the ecological drivers produced through anthropogenic activities are well
documented[64]. Interestingly, the pervasive and conspicuous urbanisation of the marine
environment in the study area showed a much weaker effect on biogeographic patterns than
abiotic factors (Fig. 2). Anthropogenic pressures have become a major ecological driver only
relatively recently in evolutionary time, with the most dramatic changes in biodiversity occurring
within the 21st century[3]. It is clear human activities are altering evolutionary trajectories[64], either
through extinction, speciation or range expansion. However, centuries of human impacts in our
study system have not yet demonstrably altered the main biogeographic patterns across
kingdoms of life.
505
Previous work on biotic homogenisation has shown a dramatic effect on whole communities at
both regional[17,65] and global scales[23,24]. Here, we found limited evidence for biotic
homogenisation along the South African coastline. More specifically, we found no major
differences in the distance-decay relationship between artificial and natural sites. This pattern
was consistent for the vast majority of gene regions considered (Fig. 3) and subsets of the genes
when individual phyla were analysed (see Supplementary Table 11). However, we did find a
significant difference in the distance-decay relationship between site types in the COI dataset as
a whole (Fig. 3), with more homogenous community composition between artificial sites. As the
majority of COI phyla did not individually provide evidence of biotic homogenisation, the
difference in the distance decay relationship observed across the whole COI dataset must be
driven by the portion of the community not assigned to taxonomy, making interpretation
difficult. The remaining markers and phyla showed no evidence of biotic homogenisation when
sites were compared, likely a result of consistent species introductions and extirpations in both
natural and artificial sites, despite the higher rate of species introductions in anthropogenic
habitats. Further work should incorporate time series data to explore biotic homogenisation,
given the significant but minor role of human impact in structuring ecological communities

across the region. Additionally, seasonal effects may reveal some variation that might explain

biotic homogenisation patterns along highly dynamic coastlines.

Environmental parameters have a clear-cut effect on the community structure across kingdoms of

life, but the proportion of the total observed variance explained in the best RDA model was less

than half. Therefore, the comparative role of deterministic (environmental filtering, niche

processes, etc.) and stochastic processes (ecological drift, random extinction, etc.) in explaining

the observed patterns remains uncertain. The classical deterministic theory (Baas Becking

hypothesis[66]) of microbial biogeography (often summarised as 'everything is everywhere')

postulates that due to vast population sizes and dispersal microbes are found in all environments

and the variation in abiotic conditions selects for those that make up the vast majority of species

in each region. This theory ignores neutral processes which have been shown to have a critical

role in structuring microbial biogeography across biomes[67]. In line with previous efforts studying

deterministic and stochastic processes across taxonomic kingdoms[68,69], we found that the

majority of the observed variation could not be explained for both prokaryotic and eukaryotic

species (Fig 2.). Indeed recent biogeographic research in the oceans has provided both

theoretical[70] and empirical[28] evidence of strong biogeographic patterns driven by both stochastic

and deterministic forces, but much of the observed variation between communities remains

unexplained. Understanding the comparative roles of different community structuring processes

requires a more comprehensive view of the observed variance between communities, the

interactions between members of these communities, and the broader role of the environmental

conditions where they live.

Several recent innovations will provide valuable data to help uncover the unexplained variation

in community structure. For example, the extraction and analysis of sedimentary ancient DNA

allows the reconstruction of high-resolution biodiversity change over time[71]. By rewinding the

ecological clock, the approach provides evidence to evaluate the role of deterministic processes

in reconstructions of community composition through time relative to changes in environmental

proxies. In conjunction, the analysis of networks from molecular data facilitate the testing of

species interaction hypotheses that may further elucidate the role of ecological interactions (e.g.

Djurhuus, et al. [14]) in structuring biogeographic patterns. Finally, very high-resolution multi-

spectral remote sensing data (e.g. WorldView-3, <100m) will provide unparalleled insights into

the role of environmental forces on the distribution of ecological communities[72]. Taken together,

recently developed technological and analytical techniques will further advance our

understanding of the mechanisms underpinning the biogeography of ecological communities

across kingdoms of life.

Acknowledgments

We acknowledge the Environmental Sequencing Facility from the National Oceanography Centre, Southampton for access and sequencing assistance. We thank IRIDIS High Performance Computing Facility, and associated support services at the University of Southampton, for access. LH was supported by the Natural Environmental Research Council (grant number NE/L002531/1). The UK Research and Innovation Newton Fund (grant: ES/N013913/1) supported LH's research stay in South Africa. LH acknowledges Shirley Parker-Nance and the Elwandle Node of the South African Environmental Observation Network for assistance and in-country logistics. LH acknowledges the assistance of Molly Czachur and Thomas Grevesse during field surveys, Sophie von der Heyden for lab consumables and both Jamie Hudson and Ivan Haigh for assistance with remote sensing data. All marina owners and operators are acknowledged for field site access.

Data Availability

Raw Illumina sequencing data are available from the European Nucleotide Archive under study

accession number PRJEB38452. Associated metadata, R scripts and intermediate files are

available at the following DOI:10.5281/zenodo.3906304 .

Code Availability

All code used in the current study can be found at the following DOI:10.5281/zenodo.3906304 .

References

Spalding, M. D. *et al.* Marine ecoregions of the world: A bioregionalization of coastal and shelf areas. *BioScience* **57**, 573-583, doi:10.1641/B570707 (2007).

Awad, A. A., Griffiths, C. L. & Turpie, J. K. Distribution of South African marine benthic invertebrates applied to the selection of priority conservation areas. *Diversity and Distributions* **8**, 129-145, doi:10.1046/j.1472-4642.2002.00132.x (2002).

Pecl, G. T. *et al.* Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science* **355**, doi:10.1126/science.aai9214 (2017).

Sunday, J. M., Bates, A. E. & Dulvy, N. K. Thermal tolerance and the global redistribution of animals. *Nature Climate Change* **2**, 686-690, doi:10.1038/nclimate1539 (2012).

Wallace, A. R. *The geographical distribution of animals, with a study of the relations of living and extinct faunas as elucidating the past changes of the earth's surface.* (Macmillan and Co., 1876).

Holt, B. G. *et al.* An update of Wallace's zoogeographic regions of the world. *Science* **339**, 74-78, doi:10.1126/science.1228282 (2013).

Ficetola, G. F., Mazel, F. & Thuiller, W. Global determinants of zoogeographical boundaries. *Nature Ecology and Evolution* **1**, 89, doi:10.1038/s41559-017-0089 (2017).

Kocsis, A. T., Reddin, C. J. & Kiessling, W. The stability of coastal benthic biogeography over the last 10 million years. *Global Ecology and Biogeography* **27**, 1106-1120, doi:10.1111/geb.12771 (2018).

Zaffosa, A., Finnegan, S. & Peters, S. E. Plate tectonic regulation of global marine animal diversity. *Proceedings of the National Academy of Sciences of the United States of America* **114**, 5653-5658, doi:10.1073/pnas.1702297114 (2017).

Costello, M. J. *et al.* Marine biogeographic realms and species endemicity. *Nature Communications* **8**, doi:10.1038/s41467-017-01121-2 (2017).

Beck, J. *et al.* What's on the horizon for macroecology? *Ecography* **35**, 673-683, doi:10.1111/j.1600-0587.2012.07364.x (2012).

Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean microbiome. *Science* **348**, 1261359, doi:10.1126/science.1261359 (2015).

Shade, A. *et al.* Macroecology to Unite All Life, Large and Small. *Trends in Ecology and Evolution* **33**, 731-744, doi:10.1016/j.tree.2018.08.005 (2018).

Djurhuus, A. *et al.* Environmental DNA reveals seasonal shifts and potential interactions in a marine community. *Nature Communications* **11**, 254, doi:10.1038/s41467-019-14105-1 (2020).

Naeem, S., Duffy, J. E. & Zavaleta, E. The functions of biological diversity in an age of extinction. *Science* **336**, 1401-1406, doi:10.1126/science.1215855 (2012).

Tilman, D., Isbell, F. & Cowles, J. M. Biodiversity and Ecosystem Functioning. *Annual Review of Ecology, Evolution, and Systematics* **45**, 471-493, doi:10.1146/annurev-ecolsys-120213-091917 (2014).

Finderup Nielsen, T., Sand-Jensen, K., Dornelas, M. & Bruun, H. H. More is less: net gain in species richness, but biotic homogenization over 140 years. *Ecology Letters* **22**, 1650-1657, doi:10.1111/ele.13361 (2019).

Blowes, S. A. *et al.* The geography of biodiversity change in marine and terrestrial assemblages. *Science* **366**, 339-345, doi:10.1126/science.aaw1620 (2019).

19   Dornelas, M. *et al.* Assemblage time series reveal biodiversity change but not systematic
loss. *Science* **344**, 296-299, doi:10.1126/science.1248484 (2014).

20   Olden, J. D. & Rooney, T. P. On defining and quantifying biotic homogenization. *Global*
Ecology and Biogeography **15**, 113-120, doi:10.1111/j.1466-822X.2006.00214.x (2006).

21   Stuart-Smith, R. D. *et al.* Integrating abundance and functional traits reveals new global
hotspots of fish diversity. *Nature* **501**, 539-542, doi:10.1038/nature12529 (2013).

22   Mouillot, D. *et al.* Rare species support vulnerable functions in high-diversity
ecosystems. *PLoS Biology* **11**, e1001569, doi:10.1371/journal.pbio.1001569 (2013).

23   Bernardo-Madrid, R. *et al.* Human activity is altering the world's zoogeographical
regions. *Ecology Letters* **22**, 1297-1305, doi:10.1111/ele.13321 (2019).

24   Capinha, C., Essl, F., Seebens, H., Moser, D. & Pereira, H. M. The dispersal of alien
species redefines biogeography in the Anthropocene. *Science* **348**, 1248-1251,
doi:10.1126/science.aaa8913 (2015).

25   Azam, F. & Malfatti, F. Microbial structuring of marine ecosystems. *Nature Reviews*
Microbiology **5**, 782-791, doi:10.1038/nrmicro1747 (2007).

26   Deiner, K. *et al.* Environmental DNA metabarcoding: Transforming how we survey
animal and plant communities. *Molecular Ecology* **26**, 5872-5895,
doi:10.1111/mec.14350 (2017).

27   Pedersen, M. W. *et al.* Ancient and modern environmental DNA. *Philosophical*
Transactions of the Royal Society B: Biological Sciences **370**, 20130383,
doi:10.1098/rstb.2013.0383 (2015).

28   Richter, D. J. *et al.* Genomic evidence for global ocean plankton biogeography shaped by
large-scale current systems. *bioRxiv*, 867739, doi:10.1101/867739 (2019).

29   Emanuel, B. P., Bustamante, R. H., Branch, G. M., Eekhout, S. & Odendaal, F. J. A
zoogeographic and functional approach to the selection of marine reserves on the west
coast of South Africa. *South African Journal of Marine Science* **12**, 341-354,
doi:10.2989/02577619209504710 (1992).

30   Griffiths, C. L., Robinson, T. B., Lange, L. & Mead, A. Marine biodiversity in South
Africa: an evaluation of current states of knowledge. *PLoS One* **5**, e12008,
doi:10.1371/journal.pone.0012008 (2010).

31   Griffiths, C. L. *et al.* Impacts of human activities on marine animal life in the Benguela:
A historical overview. *Oceanography and Marine Biology: an Annual Review* **42**, 303-
392, doi:10.1201/9780203507810.ch8 (2004).

32   Kaluza, P., Kolzsch, A., Gastner, M. T. & Blasius, B. The complex network of global
cargo ship movements. *Journal of The Royal Society Interface* **7**, 1093-1103,
doi:10.1098/rsif.2009.0495 (2010).

33   Rius, M. *et al.* Range expansions across ecoregions: interactions of climate change,
physiology and genetic diversity. *Global Ecology and Biogeography* **23**, 76-88,
doi:10.1111/geb.12105 (2014).

34   Holman, L. E. *et al.* Detection of introduced and resident marine species using
environmental DNA metabarcoding of sediment and water. *Scientific Reports* **9**, 11559,
doi:10.1038/s41598-019-47899-7 (2019).

35   Kelly, R. P. *et al.* Genetic signatures of ecological diversity along an urbanization
gradient. *PeerJ* **4**, e2444, doi:10.7717/peerj.2444 (2016).

36 Spens, J. *et al.* Comparison of capture and storage methods for aqueous macrobial eDNA
using an optimized extraction protocol: advantage of enclosed filter. *Methods in Ecology*
and Evolution **8**, 635-645, doi:10.1111/2041-210x.12683 (2017).
37 Zhan, A. *et al.* High sensitivity of 454 pyrosequencing for detection of rare species in
aquatic communities. *Methods in Ecology and Evolution* **4**, 558-565, doi:10.1111/2041-
210x.12037 (2013).
38 Leray, M. *et al.* A new versatile primer set targeting a short fragment of the
mitochondrial COI region for metabarcoding metazoan diversity: application for
characterizing coral reef fish gut contents. *Frontiers in Zoology* **10**, 34,
doi:10.1186/1742-9994-10-34 (2013).
39 Takahashi, S., Tomita, J., Nishioka, K., Hisada, T. & Nishijima, M. Development of a
prokaryotic universal primer for simultaneous analysis of Bacteria and Archaea using
next-generation sequencing. *PLoS One* **9**, e105592, doi:10.1371/journal.pone.0105592
(2014).
40 Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet Journal **17**, 10-12, doi:10.14806/ej.17.1.200 (2011).
41 Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon
data. *Nature Methods* **13**, 581-583, doi:10.1038/nmeth.3869 (2016).
42 R_Core_Team. R: a language and environment for statistical computing. *ISBN 3-900051-*
07-0 (2019).
43 Frøslev, T. G. *et al.* Algorithm for post-clustering curation of DNA amplicon data yields
reliable biodiversity estimates. *Nature Communications* **8**, 1188, doi:10.1038/s41467-
017-01312-x (2017).
44 Mahmoudabadi, G. & Phillips, R. A comprehensive and quantitative exploration of
thousands of viral genomes. *Elife* **7**, doi:10.7554/eLife.31955 (2018).
45 Edgar, R. C. Updating the 97% identity threshold for 16S ribosomal RNA OTUs.
Bioinformatics **34**, 2371-2375, doi:10.1093/bioinformatics/bty113 (2018).
46 Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Research* **41**, D590-596,
doi:10.1093/nar/gks1219 (2013).
47 JPL_OurOceanProject. GHRSST Level 4 G1SST Global Foundation Sea Surface
Temperature Analysis. doi:10.5067/GHG1S-4FP01 (2010).
48 Zweng, M. M., J. R. Reagan, D. Seidov, T. P. Boyer, R. A. Locarnini, H. E. Garcia, A. V.
Mishonov, O. K. Baranova, K. Weathers, C. R. Paver, and I. Smolyar,. World Ocean
Atlas 2018, Volume 2: Salinity. 50 (2019).
49 European_Space_Agency. Ocean Colour Climate Change Initiative Dataset, Version 4.2.
(2020).
50 Halpern, B. S. *et al.* Recent pace of change in human impact on the world's ocean.
Scientific Reports **9**, 11609, doi:10.1038/s41598-019-47201-9 (2019).
51 Anderson, M. J. in *Wiley Stats Ref: Statistics Reference Online*   (eds N. Balakrishnan *et*
al.) Ch. 1 Permutational multivariate analysis of variance (PERMANOVA), 1-15 (John
Wiley & Sons, Ltd, 2014).
52 Oksanen, J. *et al.* Vegan: community ecology package. *R package, Version 2.5-6* (2011).
53 Kreft, H. & Jetz, W. A framework for delineating biogeographical regions based on
species distributions. *Journal of Biogeography* **37**, 2029-2053, doi:10.1111/j.1365-
2699.2010.02375.x (2010).

54   Martinez Arbizu, P. PairwiseAdonis: Pairwise multilevel comparison using adonis. *R*
package, Version 0.3 (2020).
55   Anderson, M. J. Distance-Based Tests for Homogeneity of Multivariate Dispersions.
Biometrics **62**, 245-253, doi:10.1111/j.1541-0420.2005.00440.x (2006).
56   McArdle, B. H. & Anderson, M. J. Fitting Multivariate Models To Community Data: A
Comment On Distance-Based Redundancy Analysis. *Ecology* **82**, 290-297,
doi:10.1890/0012-9658(2001)082[0290:Fmmtcd]2.0.Co;2 (2001).
57   Crabot, J., Clappe, S., Dray, S. & Datry, T. Testing the Mantel statistic with a spatially-
constrained permutation procedure. *Methods in Ecology and Evolution* **10**, 532-540,
doi:10.1111/2041-210x.13141 (2019).
58   Rapacciuolo, G., Beman, J. M., Schiebelhut, L. M. & Dawson, M. N. Microbes and
macro-invertebrates show parallel β-diversity but contrasting α-diversity patterns in a
marine natural experiment. *Proceedings of the Royal Society B: Biological Sciences* **286**,
20190999, doi:doi:10.1098/rspb.2019.0999 (2019).
59   Astorga, A. *et al.* Distance decay of similarity in freshwater communities: do macro- and
microorganisms follow the same rules? *Global Ecology and Biogeography* **21**, 365-375,
doi:10.1111/j.1466-8238.2011.00681.x (2012).
60   Wang, J. *et al.* Patterns of elevational beta diversity in micro- and macroorganisms.
Global Ecology and Biogeography **21**, 743-750, doi:10.1111/j.1466-8238.2011.00718.x
(2012).
61   Tittensor, D. P. *et al.* Global patterns and predictors of marine biodiversity across taxa.
Nature **466**, 1098-U1107, doi:10.1038/nature09329 (2010).
62   Herlemann, D. P. *et al.* Transitions in bacterial communities along the 2000 km salinity
gradient of the Baltic Sea. *ISME Journal* **5**, 1571-1579, doi:10.1038/ismej.2011.41
(2011).
63   Broman, E. *et al.* Salinity drives meiofaunal community structure dynamics across the
Baltic ecosystem. *Molecular Ecology* **28**, 3813-3829, doi:10.1111/mec.15179 (2019).
64   Shochat, E., Warren, P. S., Faeth, S. H., McIntyre, N. E. & Hope, D. From patterns to
emerging processes in mechanistic urban ecology. *Trends in Ecology and Evolution* **21**,
186-191, doi:10.1016/j.tree.2005.11.019 (2006).
65   Blouin, D., Pellerin, S. & Poulin, M. Increase in non-native species richness leads to
biotic homogenization in vacant lots of a highly urbanized landscape. *Urban Ecosystems*
22, 879-892, doi:10.1007/s11252-019-00863-9 (2019).
66   Baas-Becking, L. G. M. *Geobiologie; of inleiding tot de milieukunde.* (WP Van Stockum
& Zoon NV, 1934).
67   Hanson, C. A., Fuhrman, J. A., Horner-Devine, M. C. & Martiny, J. B. Beyond
biogeographic patterns: processes shaping the microbial landscape. *Nature Reviews*
Microbiology **10**, 497-506, doi:10.1038/nrmicro2795 (2012).
68   Farjalla, V. F. *et al.* Ecological determinism increases with organism size. *Ecology* **93**,
1752-1759, doi:10.1890/11-1144.1 (2012).
69   Wu, W. X. *et al.* Contrasting the relative importance of species sorting and dispersal
limitation in shaping marine bacterial versus protist communities. *ISME Journal* **12**, 485-
494, doi:10.1038/ismej.2017.183 (2018).
70   Hellweger, F. L., van Sebille, E. & Fredrick, N. D. Biogeographic patterns in ocean
microbes emerge in a neutral agent-based model. *Science* **345**, 1346-1349,
doi:10.1126/science.1254421 (2014).

71   Balint, M. *et al.* Environmental DNA time series in ecology. *Trends in Ecology and*
Evolution **33**, 945-957, doi:10.1016/j.tree.2018.09.003 (2018).
72   He, K. S. *et al.* Will remote sensing shape the next generation of species distribution
models? *Remote Sensing in Ecology and Conservation* **1**, 4-18, doi:10.1002/rse2.7
(2015).