Title

Wolbachia-conferred antiviral protection is determined by

developmental temperature

Ewa Chrostek[1,*, †], Nelson Martins[2,3,†], Marta S Marialva[4], Luis Teixeira[2,5, *]

1. Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, United
Kingdom
2. Instituto Gulbenkian de Ciência, Oeiras, Portugal.
3. Institut de Biologie Moléculaire et Cellulaire, Université de Strasbourg, Strasbourg,
France
4. Department for Biomedical Research, University of Bern, Switzerland
5. Faculdade de Medicina da Universidade de Lisboa, Lisboa, Portugal.

* **Corresponding authors:** Ewa Chrostek and Luis Teixeira

Email: e.chrostek@liv.ac.uk and lteixeira@igc.gulbenkian.pt

† These authors contributed equally to the work.

ORCID IDs:
Ewa Chrostek: 0000-0003-1825-372X
Nelson Martins: 0000-0002-3923-2998
Marta S Marialva: 0000-0002-5310-9224
Luis Teixeira: 0000-0001-8326-6645

Classification
Biological Sciences >> Microbiology

Keywords
Wolbachia, Drosophila, virus, temperature, symbiosis, development

Author Contributions
EC, MSM and LT designed research; EC and MSM performed research; EC, MSM, NM and LT
analyzed data; EC and LT edited figures; and EC and LT wrote the paper.

37
38

Abstract

Wolbachia is a maternally transmitted bacterium widespread in arthropods and filarial nematodes, and confers strong antiviral protection in *Drosophila melanogaster* and other insects. *Wolbachia*-transinfected *Aedes aegypti* are currently being deployed to fight transmission of dengue and Zika viruses. However, the mechanism of antiviral protection and factors influencing it are still not fully understood. Here we show that temperature modulates *Wolbachia*-conferred protection in *Drosophila melanogaster*. Temperature after infection directly impacts *Drosophila* C virus replication and modulates *Wolbachia* protection. At higher temperatures virus proliferates more and is more lethal, while *Wolbachia* confers lower protection. Strikingly, host developmental temperature is a determinant of *Wolbachia*-conferred antiviral protection. While there is a strong protection when flies are raised from egg to adult at 25°C, the protection is highly reduced or completely abolished when flies develop at 18°C. However, *Wolbachia*-induced changes during development are not sufficient to limit virus-induced mortality, as *Wolbachia* is still required to be present in adults at the time of infection. This developmental effect is general, since it was present in different host genotypes, *Wolbachia* variants and upon infection with different viruses. Overall, we show that *Wolbachia*-conferred antiviral protection is temperature dependent, being present or absent depending on the environmental conditions. This interaction likely impacts *Wolbachia*-host interactions in nature and, as a result, frequencies of host and symbionts in different climates. Dependence of *Wolbachia*-mediated pathogen blocking on developmental temperature could be used to dissect the mechanistic bases of protection and should be considered by programmes deploying *Wolbachia* as an antiviral agent in the field.

Significance Statement

Insects are often infected with beneficial intracellular bacteria. The bacterium *Wolbachia* can protect insects from pathogenic viruses. This effect can be used to prevent transmission of dengue and Zika viruses by *Wolbachia*-infected mosquitoes. To deploy *Wolbachia* in the field successfully and understand the biology of insects in the wild we need to discover which factors affect *Wolbachia*-conferred antiviral protection. Here we show that the temperature in which insects develop from eggs to adults can determine presence or absence of antiviral protection. The environment, therefore, influences this insect-bacterium interaction. Our work may help to provide insights into the mechanism of viral blocking by *Wolbachia* and inform programs using *Wolbachia* in mosquito-borne disease control.

Main Text

Introduction

Wolbachia is a maternally inherited intracellular bacterium infecting a wide range of arthropods and

some nematodes. This endosymbiont induces strong phenotypes in many of its hosts, which may

contribute to its success in invading and being maintained in host populations. *w*Mel, the *Wolbachia*

strain present in *Drosophila melanogaster,* confers a strong protection to a wide range of RNA

viruses, which can be a fitness benefit in nature (1, 2). This protection extends to non-native

Wolbachia-hosts associations, including mosquito vectors of human disease (3). Recently,

Wolbachia has become one of the most promising approaches to control dengue and Zika viruses

in the wild. It has been shown that the releases of *Wolbachia*-infected *Aedes aegypti* can reduce

the number of dengue cases in both, endemic and non-endemic areas (4–7). Although molecular

mechanisms of *Wolbachia*-conferred antiviral protection are not yet known, identification of factors

influencing protection can contribute to understanding the association of *Wolbachia* and hosts in

natural population, as well as guide *Wolbachia*-based field interventions.

Wolbachia-conferred antiviral protection is influenced by host and bacterial genetics. Different

Wolbachia strains in the same host genetic background vary in protection (8–12). In general,

differences in *Wolbachia* titres are correlated with differences in protection, with higher titres

conferring higher antiviral protection (8–12). However, some *Wolbachia* strains do not provide

protection despite high titres (13). Host genetic variation can also contribute to the strength of

Wolbachia-conferred protection, as seen in *Aedes aegypti* (14).

Environmental factors also affect this symbiont-mediated protection, either through modulation of

Wolbachia titre or by titre-independent mechanisms. Host diet rich in cholesterol reduces antiviral

protection in *Drosophila melanogaster* (15). Similarly, antibiotic treatment reducing *Wolbachia* titre

reduces the antiviral protection (16). Finally, temperature was shown to modulate *Wolbachia*-

conferred protection to parasites in an engineered *Wolbachia*-host association; in *Anopheles*

stephensi, temperature and somatic *Wolbachia* infection determined *Plasmodium* titre in

mosquitoes (17). On the other hand, in *Aedes aegypti* stably transinfected with *w*Me*l,* the

comparison of a constant and two fluctuating temperature regimes concluded, that the antiviral

protection is robust across conditions and unlikely to be compromised in the antiviral field trials

(18). However, different temperatures regimes affect dengue transmission independently of

Wolbachia (19).

Despite two studies tackling temperature effect on *Wolbachia*-conferred protection in artificial

associations, the impact of temperature on antiviral protection in natural *Wolbachia*-host symbiosis

remains unknown. Yet, it is known that temperature can affect the interaction between *Wolbachia*

and *D. melanogaster.* For instance, higher temperature leads to higher proliferation and cost of the

pathogenic variant *w*MelPop (20–22). Also, flies with different *w*Mel variants prefer different

temperatures, probably related with cost of these variants to the host (23). Geographic distribution

of *Wolbachia* in *D. melanogaster* populations could be explained by their relative fitness effects at

varying thermal conditions (24, 25). Antiviral protection could be a temperature-dependent fitness

benefit, which would be balanced against the potential cost of highly protective, high titre symbionts.

Here we tested how temperature affects *Wolbachia* protection to viruses in natural *Wolbachia-*

Drosophila associations. First, we asked how different virus doses and post-infection temperatures

affect *Wolbachia* densities and antiviral protection. Then, we dissected the effect of developmental

temperature on *Wolbachia*-carrying *Drosophila melanogaster*. We show strong interaction between

this environmental variable and *Wolbachia*-conferred protection against viruses.

Results

To test how different infection temperatures affect *Wolbachia*-conferred protection to *Drosophila* C

virus (DCV), we used the *Drosophila melanogaster* Drosdel w^{1118} isogenic line (*iso*) carrying a

natural *Wolbachia* variant, *w*MelCS_b (*Wolb+*), and a matching *Wolbachia*-free control (*Wolb-*) (1,

8). Flies were raised at 25°C and 3-6-day-old flies were challenged with serial dilutions of DCV and

subsequently placed at either 18°C or at 25°C (Fig. 1A, B). Virus-induced mortality was higher at

25°C than 18°C at all doses except the lowest one, where there is almost no mortality associated

with the infection (Temperature effect, $p \leq 0.001$ for all comparisons, except with infection dose of

10^5 TCID$_{50}$/ml). Importantly, *Wolbachia* protection against DCV varies with temperature

(*Wolbachia**Temperature interaction effect, $p < 0.001$), and it is stronger when temperature of

infection is 18°C (Cox hazard ratio (CHR) = -1.84, $p < 0.001$) than 25°C (CHR = -0.75 $p < 0.001$,

at 18°C). Overall, the temperature of infection affects the survival of the flies and *Wolbachia*

protects more at lower infection temperature (Fig. 1A,B).

Figure 1. Post-infection temperature modulates the strength of *Wolbachia*-conferred antiviral protection. (A,B) Survival curves of flies infected at different temperatures, with the schemes of the experimental designs shown above. *Wolb+* and *Wolb-* flies, 50 per *Wolbachia* status and condition, were raised at 25ºC, infected with different doses of DCV and placed at either 18°C (A) or 25°C (B) after DCV infection. Mortality was recorded daily. C) DCV titres in DCV-infected flies 3 dpi measured by RT-qPCR. Flies were kept at 25°C before infection and either at 18°C or 25°C after infection. (D) *Wolbachia* levels measured by qPCR at the day of infection (3-6 days old), or 3 days after in unchallenged (control), buffer- or DCV-challenged (10^7 TCID$_{50}$/ml) flies. Flies were kept at 25°C before and placed at either 18°C or 25°C after DCV infection. In (C) and (D) each dot is a sample of ten flies and horizontal lines are medians.

To test if the interaction of *Wolbachia* protection with temperature is also reflected in viral loads, we measured viral titres in flies three days post infection (dpi) by quantitative reverse transcription PCR (RT-qPCR) (Fig. 1C). We detected a strong interaction between *Wolbachia* presence,

temperature of infection, and dose of the virus ($p < 0.001$). At the lower temperature *Wolbachia*
confers more resistance at higher viral doses, as viral titres stay very low at lower doses in both
Wolb+ and *Wolb-* flies. At the higher temperature, it confers more resistance at the lower doses, as
virus titres are very high in *Wolb+* and *Wolb-* flies exposed to high virus doses. The mean viral load
was higher at 25ºC than at 18ºC ($p < 0.001$) and was lower in the presence of *Wolbachia* ($p <$
0.001). On average, *Wolbachia* induces higher resistance at 18°C than at 25°C. There was a 550-
fold reduction in viral load at 18°C ($p < 0.001$) and 50-fold reduction at 25°C ($p < 0.001$), producing
approximately 11-fold difference between these two conditions ($p = 0.003$). These results show
that the strength of *Wolbachia*-conferred protection to DCV, in terms of both, survival and viral
loads, depends on the temperature of infection. Protection is higher at a lower temperature. Since
post-infection temperature affects viral infection independently of *Wolbachia*, the lower capacity of
Wolbachia to protect at higher post-infection temperatures may be related with a higher replication
of the virus.

Antiviral protection is also usually positively correlated with *Wolbachia* levels (8, 9, 11, 16, 26, 27),
however, so we tested *Wolbachia* levels at the day of infection and after three days in DCV-infected,
buffer-pricked (control), or unmanipulated flies, at both temperatures (Fig. 1D). *Wolbachia* levels
were not affected by virus, buffer or temperature of infection ($p > 0.425$ for effect of temperature or
treatment). Thus, the difference in protection is likely independent of *Wolbachia* levels, as these
are not significantly affected within three days of the viral challenge.

We extended our analysis by testing how pre-infection temperature affects *Wolbachia* protection
(Fig. 2). We followed survival of infected flies in all four possible combinations of pre- and post-
infection temperatures of 18°C and 25°C (Fig. 2A, B). There is a strong interaction between pre-
infection temperature and *Wolbachia* ($p = 0.009$). Remarkably, in flies raised at 18°C *Wolbachia*
does not protect against DCV infection ($p = 0.207$), while in flies raised at 25°C it does (CHR = -
1.38, $p < 0.001$), irrespectively of the post-infection temperature. Also, in the absence of *Wolbachia*
the pre-infection temperature has no effect on the survival after DCV infection ($p = 0.276$). This
strong interaction between *Wolbachia*-conferred protection to DCV and pre-infection temperature
is also reflected in DCV loads (Fig. 2C, S1, $p < 0.001$). On average, *Wolbachia* reduced viral titres
14-fold at pre-infection temperature of 18°C and 1900-fold at 25°C (difference $p < 0.001$). In these
assays, pre-infection temperature also did not affect viral loads in the absence of *Wolbachia* ($p =$
0.534). Therefore, in contrast to post-infection temperature, pre-infection temperature does not
directly impact virus performance and modulates only *Wolbachia* protection. In summary, in flies
raised at lower temperature *Wolbachia* confers a reduced resistance to DCV in terms of viral titres
and no protection in terms of survival.

Figure 2. Pre-infection temperature determines *Wolbachia*-conferred antiviral protection. (A,B) *Wolb+*
and *Wolb-* flies, fifty per *Wolbachia* status per condition, were infected with DCV (10^8 TCID$_{50}$/ml) and checked
for survival every day. Flies were raised and kept at 25°C (A) or 18°C (B) before DCV infection and placed at
either 18°C or 25°C after infection. (C) DCV titres in flies 3 dpi measured by RT-qPCR. Flies were raised and
kept at either 18°C or 25°C before infection and at 18°C after infection. Replicate of this experiment is shown
in Fig. S1. (D) *Wolbachia* levels measured by qPCR in 3-6 days old flies raised at either 18°C or 25°C. For
(C) and (D) each dot is a sample of ten flies. Horizontal lines are medians of the samples.

Next, we measured *Wolbachia* levels in flies raised at 18°C and 25°C (Fig 2D). Flies raised at 18°C
had approximately 33% less *Wolbachia* than the flies raised at 25°C ($p < 0.001$). Although this
difference may contribute to the difference in protection, a larger, 50% difference in *Wolbachia* titres
between *w*Mel- and *w*MelCS-harbouring flies did not abolish the protection (8). Thus, differences
in *Wolbachia* titres do not seem to fully explain the difference in protection between flies raised at
different temperatures.

To test the robustness of pre-infection temperature effect on *Wolbachia*-conferred protection, we
tested flies with distinct host genetic backgrounds, Aljezur1 and Oregon-R W-20 (1), each
harbouring its original *w*Mel-like *Wolbachia* strain (Fig. 3A,B and S2A,B). In agreement with the
previous results, there was a significant effect of pre-infection temperature on *Wolbachia*-conferred
antiviral protection against DCV in both lines ($p<0.001$ for both). *Wolbachia* in Aljezur1 line
conferred protection only when the pre-infection temperature was 25°C ($p = 0.763$ for 18°C, and
CHR = -1.19, $p < 0.001$ for 25°C). In the Oregon-R W-20 line, *Wolbachia* still conferred some
protection when pre-infection temperature was 18°C, although it was significantly less than at 25°C
(CHR = -1.21, $p < 0.001$ for 18°C, and CHR = -3.02, $p < 0.001$ for 25°C, CHR difference = 1.8, $p <$
0.001). Next, we investigated if the pre-infection temperature effect was specific to DCV or if it was
also present upon challenge with another RNA virus, Flock house virus (FHV) (Fig. 3C and S2C),
There was an interaction between *Wolbachia*, pre-infection temperature, and dose in this assay (p
$= 0.013$). *Wolbachia* protection was significantly higher at flies raised at 25°C for all doses of virus
except for the lowest one (contrasts of CHR < -2.84, $p < 0.001$ for $10^7 - 10^9$ TCID$_{50}$/ml). At the
lowest dose, 10^6 TCID$_{50}$/ml, there was an overall low mortality of flies and the difference in survival
was not statistically significant (contrast of CHR = -1.00 but $p = 0.421$). The pre-infection
temperature had no effect on survival to FHV in the absence of *Wolbachia* ($p = 1$, in all
comparisons), as observed for DCV. In conclusion, the strong influence of pre-infection
temperature on *Wolbachia*-conferred protection is a general effect observed in different host
genetic backgrounds, *Wolbachia* variants and with the different viruses used.

Since there is a strong interaction between protection and constant pre-infection temperature, we
asked if protection exists when temperature is cycling daily between 18ºC and 25ºC (Fig. S3).
Wolbachia protects against DCV infection at the cycling temperature (CHR = - 0.67, $p = 0.014$).
This protection seems to be intermediate between the protection seen in flies developed at 18ºC
or 25ºC. This result shows that in daily cycling temperatures, which occur in natural conditions,
Wolbachia can protect against viruses.

Figure 3. Pre-infection temperature determines *Wolbachia*-conferred antiviral protection in different *Wolbachia* and *Drosophila* genotypes and against different viruses. (A) Aljezur 1 and (B) Oregon-R (W-20) *Wolb+* and *Wolb-* flies, fifty per *Wolbachia* status per condition, were pricked with three dilutions of DCV. (C) w^{1118} *iso* flies, fifty per *Wolbachia* status per condition, were infected with three doses of FHV. Flies were kept at 25°C or 18°C before the infection, and at 18°C after the infection. Mortality was recorded daily. A replicate of this experiment is shown in Fig. S2.

In the above protocols, flies developed from egg to adult at a given temperature and after collection they were aged for three more days at the same temperature, before infection with viruses. To dissect at which of these stages, development or aging, temperature influences *Wolbachia* protection, we tested the four possible combinations of the two different temperatures for these two stages (Fig. S4). Developmental temperature strongly influenced *Wolbachia* protection, which was higher at 25°C (CHR = -1.16, *p* < 0.001). Aging the flies at 25°C also interacted with *Wolbachia* protection, but increased it only slightly compared to aging at 18°C (CHR = -0.431, *p* = 0.011). Therefore, *Wolbachia* protection to viruses is mainly dependent on the temperature of development form egg to adult.

Since pre-infection events are crucial for *Wolbachia*-conferred resistance we asked how fast the

difference in virus titres arise between flies with and without *Wolbachia* (Fig. 4A). We observed a

Wolbachia-induced resistance of approximately 120-fold as soon as we can detect viral RNA, at

12h after infection, ($p < 0.001$ at 12h, and $p < 0.002$ for all posterior time points in Fig 4A and S5A).

Thus, *Wolbachia* lowers viral titres very early in the course of DCV infection. We observed a similar

pattern of early FHV blocking by *Wolbachia*, from day one post infection onwards, but with smaller

viral titre differences (*Wolbachia*-infected flies had 2.3 to 5.5 times less FVH from the first day

onwards, $p > 0.028$ for all these comparisons, Fig. S5B).

260
Figure 4. *Wolbachia* presence in adults is required for antiviral protection. (A) Time course analysis of
DCV titres in *Wolb+* and *Wolb-* flies raised at 25°C and kept at 18°C after DCV infection. Relative DCV levels
were determined by RT-qPCR. Time 0 corresponds to the time of infection. Each dot is a sample consisting
of 10 flies, lines are medians. (B) *Wolbachia* levels in single flies raised at 25°C, at eclosion (day 0), after 10
days of different antibiotic and control treatments at 25°C (day 10), and after additional 20 days of treatment
at 18°C (day 30), measured by qPCR. A replicate of this experiment is shown in Fig. S5C. (C) Survival of
Wolb+ and *Wolb-* flies, fifty per *Wolbachia* status per treatment, developed at 25°C, collected at eclosion, fed
antibiotics-containing food for 10 days at 25°C, infected with DCV (10^8 TCID$_{50}$/ml), and placed at 18°C on
antibiotic-containing food. Mortality was recorded daily. A replicate of this experiment is shown in Fig. S5E.

Developmental temperature determines protection and the protection can be detected as early as

we detect viral replication in flies. Thus, we asked if changes in the fly caused by *Wolbachia* during

development are sufficient to block viral infection in adults. To answer this, adult flies developed at

25°C were subsequently treated with antibiotics for ten days to remove *Wolbachia* before viral

infection (Fig. 4B, S5C). *Wolbachia* levels were reduced approximately 10-fold in ten days of

treatment with tetracycline and rifampicin ($p < 0.001$ for both) and stayed low until the end of the

experiment at 30 days ($p < 0.001$ for both). These treatments eliminated antiviral protection from

the flies ($p = 1.00$ for both, Fig. 4C, S5E). The control treatments with the antibiotics ampicillin and

streptomycin did not affect *Wolbachia* levels ($p > 0.254$ for both, Fig. 4B, S5C) and did not affect

endosymbiont-mediated protection (in both treatments *Wolbachia* protection is significant, $p <$

0.001 and not different from *Wolbachia* protection in controls, $p > 0.301$, Fig. 4C, S5E). As bacteria

in the fly gut were efficiently cleared by all antibiotic treatments (Fig. S5D), we conclude that it is

Wolbachia loss that leads to the loss of protection and not differential effect of the antibiotics on

gut-associated bacteria. These data show that *Wolbachia* presence in adults is required for the

Wolbachia-conferred antiviral protection, even though the presence or absence of protection is

determined by the temperature during development, before the challenge occurs.

Discussion

Here we show that temperature is a strong modulator of *Wolbachia*-conferred antiviral protection

in the natural host *D. melanogaster*. The temperature at which the infection progresses influences

this interaction, with *Wolbachia* giving more resistance and increasing survival at lower

temperatures. However, the most striking phenotype we report is the effect of host developmental

temperature on this interaction. While development at 25ºC leads to a strong antiviral protection in

terms of survival and resistance to DCV, development at 18ºC abrogates, or strongly reduces

protection. This is observed with different genotypes of *D. melanogaster,* different variants of

Wolbachia (*w*Mel and *w*MelCS), and different viruses, and is, therefore, likely to be a general

phenomenon.

This complex interaction between temperature and *Wolbachia* protection to viruses may play a role

in the natural environment. The outcome of *Wolbachia*-host interactions, in terms of antiviral

protection, may vary in regions with different climates or across seasons in the same region.

Therefore, *Wolbachia* may behave as a parasite or a mutualist, depending on the conditions.

Wolbachia's placement along the parasitism-mutualism continuum at any given time will depend

on the cost of harbouring *Wolbachia* and the benefit of antiviral protection to the host. Our results

lead to a prediction that *Wolbachia* would protect more in warmer conditions, given the strong

reduction in antiviral protection with a low developmental temperature. We have suggested before

that the antiviral protection is a fitness advantage that may explain *Wolbachia* prevalence in *D.*

melanogaster populations (1). Selection for hosts carrying *Wolbachia* would, therefore, be stronger

at higher temperatures. Consequently, geography and seasonality could impact the frequency of

Wolbachia in a population. Interestingly, there is a high variation in the frequency of *Wolbachia* in

D. melanogaster natural populations and a clinal distribution of this frequency. At lower latitudes,
and therefore warmer climates, the frequency *of Wolbachia* is higher (28). Although, the
relationship between *Wolbachia* frequency, temperature and other environmental parameters in *D.*
melanogaster is more complex at a smaller geographic scale, *Wolbachia* frequency is the highest
in regions with mean annual temperature of 22-26ºC (29). In insects, in general, there is also a
positive correlation between temperature and *Wolbachia* frequency but only in temperate climates
(25).

These results are also relevant for the deployment of *Wolbachia*-carrying mosquitoes to block
dengue, Zika, chikungunya and other arboviruses transmission (3, 30–32). It is already known that
heat stress impacts *w*Mel in transinfected *Aedes aegypti* , reducing its titres (33), heatwaves can
impact titres and frequency of *Wolbachia* in these mosquito populations (34) and at low cycling
temperatures *Wolbachia* titers decrease (35). On the other hand, temperatures varying between
25 and 28ºC do not seem to affect protection to viruses in this system (18). It would be important,
however, to assess how lower temperatures influence *Wolbachia*-induced pathogen blocking in
mosquitoes. Determining the temperature range of effectiveness of *Wolbachia*-deploying antiviral
field interventions will be crucial to plan where to use it and when to combine it with other
complementary approaches (e.g. insecticides).

Pre-infection temperature has a drastic and enigmatic effect on *Wolbachia*-conferred antiviral
protection. Lower temperatures during host development determine the level of protection in adult
life. Understanding the molecular mechanism of this effect will elucidate how environment interacts
with host-microbe symbioses and may be key to understanding *Wolbachia* antiviral protection. The
effect of development temperature acts solely on the interaction between virus and *Wolbachia,* and
does affect viral infection by itself. The protection conferred by *Wolbachia* seems onset immediately
or very soon after viral infection. We observe a difference in viral titres as soon as we detected viral
replication in *Wolbachia*-free flies, at 12h after infection. Thus, *Wolbachia* likely inhibits the viral
entry or first replication cycle *in vivo*. *Wolbachia* interference in early stages of viral infection is in
agreement with cell culture data (36–38). However, this is not a simple pre-set antiviral state of the
host because *Wolbachia* is still required at the time of infection. The lower *Wolbachia* titres in flies
developed at 18ºC, when compared with flies developed at 25ºC, could partly explain a reduction
in protection. However, this difference should not be enough for the complete lack of protection
when flies develop at 18ºC. We have previously observed significant protection to DCV in flies
carrying even lower titres of other *w*Mel variants (8). Nonetheless, tissue-specific differences in
titres, set during development, may underlie the phenotypic differences. It would, therefore, be
interesting to characterize in detail the spatial characterisation of *Wolbachia* tissue tropism and viral

infection, in flies developed at the different temperatures. Comparative transcriptomic and
metabolomic analysis of *D. melanogaster* and *Wolbachia* in these two conditions could not only
elucidate how temperature affects protection, but also the mechanism of *Wolbachia* antiviral
protection itself.

Temperature affects many insect-symbiont interactions and their phenotypes (39), including
protective symbiosis (40–42). Therefore, this environmental factor may play a general critical role
determining the outcome of complex host-endosymbiont-pathogen interactions and shape the
geographic distribution of insects and their symbionts. The phenotypic variation we report here,
ranging from no protection to strong protection to viruses, indicates that temperature could be a
crucial determinant of cost/benefit ratio of carrying *Wolbachia*. Therefore, temperature could
deeply impact the *D. melanogaster – Wolbachia* interaction in natural populations. Moreover, these
results may have important implications for the deployment of *Wolbachia*-carrying mosquitoes to
fight arboviruses transmission and may lead to new approaches to dissect the mechanism of
Wolbachia antiviral protection.

Materials and Methods

Fly strains and husbandry
DrosDel *w*[1118] *isogenic D. melanogaster* (*iso*) with *w*MelCS_b *Wolbachia* (*Wolb*+) and the matching
control without *Wolbachia* (*Wolb*-) were described elsewhere (1, 8, 43). Aljezur 1 and W-20 *D.*
melanogaster lines (*Wolb*+ and *Wolb*-) were also described before (1). We determined that the
Wolbachia variants in both these lines lack an IS5 transposon insertion in gene WD1310, based
on the primers described in (44). This insertion is present in all *w*MelCS-like variants, but not in
wMel variants (8, 44), therefore Aljezur-1 and W-20 are both *w*Mel-like *Wolbachia* variants. Stocks
were maintained at a constant temperature of 25°C on a diet consisting of: 45 g molasses, 75 g
sugar, 70 g cornmeal, 20 g yeast extract, 10 g agar, 1100 ml water and 25 ml of 10% Nipagin, with
addition of live yeast (Sigma).

Virus infection experiments
DCV and FHV were produced and titrated in cell culture as described before (1, 8). Flies for
experiments were produced by placing 12 females and 6 males per bottle for 4 days to produce
offspring at either 25°C, 18°C or at fluctuating temperature (18°C to 25°C gradual increase during
12 h, and 25°C to 18°C decrease during the subsequent 12 h). After 10 days (25°C), 15 days
(fluctuating temperature) or 20 days (18°C) the flies started to eclode. Unless specified otherwise,
0-3 days old flies were collected from the bottles and placed in the vials, ten males per vial, on food

without live yeast. Flies were aged for 3 more days at the developmental or otherwise indicated
temperature. These 3-6 days old flies were pricked intrathoracically with virus diluted in 50 mM
Tris-HCl, pH 7.5. After infection, flies were placed at the indicated temperatures. Survival was
monitored daily and vials were changed every 5 days.
388
Nucleic acids extractions and real-time qPCR
DNA for the quantification of *Wolbachia* was extracted from pools of 10 flies using DrosDel protocol
(http://www.drosdel.org.uk/molecular_methods.php) (43) or from single flies with the protocol
described before (45). RNA for assessment of viral titres was extracted using Trizol (Invitrogen)
and cDNA was prepared using M-MLV Reverse Transcriptase (Promega), as described previously
(8). Real-time qPCR reactions were carried out in 7900HT Fast Real-Time PCR System (Applied
Biosystems) with the iQ™ SYBR® Green supermix (Bio Rad) or in QuantStudio™ 7 Flex Real-
Time PCR System (Applied Biosystems) with iTaq™ universal SYBR® Green supermix (Bio Rad).
Wolbachia was quantified using *wsp* as the target gene and *Drosophila Rpl32* as the reference
gene. DCV was quantified with primers for DCV as the target gene and *Drosophila rpl32* as the
reference gene. Primers used were: *Wolbachia wsp,* 5'-CATTGGTGTTGGTGTTGGTG-3' and 5'-
ACCGAAATAACGAGCTCCAG-3';   DCV,   5'-TCATCGGTATGCACATTGCT-3'   and   5'-
CGCATAACCATGCTCTTCTG-3';   FHV,   5'-ACCTCGATGGCAGGGTTT-3'   and   5'-
CTTGAACCATGGCCTTTTG-3'; and *Drosophila rpl32*, 5'-CCGCTTCAAGGGACAGTATC-3' and
5'-CAATCTCCTTGCGCTTCTTG-3'. Thermal cycling protocol for *Wolbachia* amplification was:
initial 50°C for 2 min, denaturation for 10 min at 95°C followed by 40 cycles of 30 s at 95°C, 1 min
at 59°C and 30 s at 72°C. For DCV and FHV the same conditions were used, except for an
annealing temperature of 56°C. Relative levels of *Wolbachia* or DCV were calculated by the Pfaffl
method (46).
408

Antibiotic treatment of flies

Wolb+ and *Wolb-* flies were raised at 25°C, collected as 0-1 days old adults and placed on fly food with 100 mg/ml of tetracycline hydrochloride, rifampicin, ampicillin sodium or streptomycin sulphate (all Sigma) or control food with antibiotics solvent (water or ethanol). At day 10 of treatment, flies were challenged with DCV, placed at 18°C and survival was followed for additional 20 days. Food vials were changed every three days. At day 0, day 10 and day 30 flies were collected to assay *Wolbachia* levels by qPCR. At day 31, guts from three *Wolb+* flies per condition were dissected to assess the effect of antibiotics on the gut-associated bacteria. Guts were homogenized in 250 µl of sterile LB and 30 µl was plated on mannitol agar plates. This medium sustains growth of the main gut-associated bacteria in lab *D. melanogaster*, *Acetobacter* and *Lactobacillus* species (47). CFUs were counted after five days of incubation at 26°C.

Statistical analysis

All the statistical analysis was performed in R (48). The script of the statistical analysis and output of this analysis are presented in supplementary texts 1 and 2, respectively. All data for statistical analysis and presented in figures are available as supplementary files.

Analysis of survival data was performed with the Cox proportional hazard mixed effect models. Fixed effects, depending on the experiment, included temperature, dose of DCV, presence/absence of *Wolbachia,* and antibiotic treatment, while replicate vials within the same experiment or full experimental replicates were considered as random effects. Model fitting was performed using the coxme package in R version 2.2-16 (49).

Wolbachia and DCV titres were analysed with log-transformed qPCR data and linear models or general linear models. Model fitting was performed using lm or the lme4 package in R (50).

The effect of interaction between factors in the models was determined by ANOVA. Post hoc analysis of marginal (least square) means were compared between the conditions of interest using the lsmeans package in R (51).

Figures were produced using the ggplot2 package (52).

Acknowledgments

We thank Dr. Rita Valente for providing assistance with plating bacteria for CFU analyses. This work was funded by FEBS Long Term Fellowship and Marie Skłodowska-Curie Individual Fellowship MSCA-IF-2017-794507 to EC, FCT grant PTDC/BEX-GMG/3128/2014 to NM, and FCT grants PTDC/BIA-MIC/108327/2008, IF/00839/2015, Wellcome Trust grant 094664/Z/10/Z, and ERC grant 773260 to LT.

445
446
447   **References**
448
449   1.   Teixeira L, Ferreira A, Ashburner M (2008) The bacterial symbiont Wolbachia induces
450   resistance to RNA viral infections in Drosophila melanogaster. *PLoS Biol* 6(12):e1000002.

451   2.   Hedges LM, Brownlie JC, O'Neill SL, Johnson KN (2008) Wolbachia and virus protection in
452   insects. *Science* 322(5902):702.

453   3.   Moreira LA, et al. (2009) A Wolbachia symbiont in Aedes aegypti limits infection with
454   dengue, Chikungunya, and Plasmodium. *Cell* 139(7):1268–78.

455   4.   Nazni WA, et al. (2019) Establishment of Wolbachia strain wAlbB in Malaysian populations
456   of Aedes aegypti for dengue control. *bioRxiv*:775965.

457   5.   O'Neill SL (2018) The Use of Wolbachia by the World Mosquito Program to Interrupt
458   Transmission of Aedes aegypti Transmitted Viruses. *Dengue and Zika: Control and Antiviral*
459   *Treatment Strategies* (Springer Link), pp 355–360.

460   6.   O'Neill SL, et al. (2019) Scaled deployment of Wolbachia to protect the community from
461   dengue and other Aedes transmitted arboviruses. *Gates Open Res* 2:36.

462   7.   Chrostek E, Hurst GDD, McGraw EA (2020) Infectious Diseases: Antiviral Wolbachia Limits
463   Dengue in Malaysia. *Curr Biol* 30(1):R30–R32.

464   8.   Chrostek E, et al. (2013) Wolbachia variants induce differential protection to viruses in
465   Drosophila melanogaster: a phenotypic and phylogenomic analysis. *PLoS Genet*
466   9(12):e1003896.

467   9.   Chrostek E, Marialva MSP, Yamada R, O'Neill SL, Teixeira L (2014) High anti-viral
468   protection without immune upregulation after interspecies Wolbachia transfer. *PLoS One*
469   9(6):e99025.

470   10.   Martinez J, et al. (2017) Symbiont strain is the main determinant of variation in Wolbachia -
471   mediated protection against viruses across Drosophila species. *Mol Ecol*.
472   doi:10.1111/mec.14164.

473   11.   Martinez J, et al. (2014) Symbionts commonly provide broad spectrum resistance to viruses
474   in insects: a comparative analysis of Wolbachia strains. *PLoS Pathog* 10(9):e1004369.

475   12.   Osborne SE, Leong YS, O'Neill SL, Johnson KN (2009) Variation in antiviral protection
476   mediated by different Wolbachia strains in Drosophila simulans. *PLoS Pathog*
477   5(11):e1000656.

478   13.   Fraser JE, et al. (2020) Novel phenotype of Wolbachia strain wPip in Aedes aegypti
479   challenges assumptions on mechanisms of Wolbachia-mediated dengue virus inhibition.
480   *bioRxiv*:2020.02.20.957423.

481   14.   Ford SA, et al. (2019) Selection on Aedes aegypti alters Wolbachia-mediated dengue virus
482   blocking and fitness. *Nat Microbiol* 4(11):1832–1839.

483   15.   Caragata EP, et al. (2013) Dietary Cholesterol Modulates Pathogen Blocking by Wolbachia.
484   *PLoS Pathog* 9(6):e1003459.

16.   Osborne SE, Iturbe-Ormaetxe I, Brownlie JC, O'Neill SL, Johnson KN (2012) Antiviral

protection and the importance of Wolbachia density and tissue tropism in Drosophila
simulans. *Appl Environ Microbiol* 78(19):6922–9.

17.   Murdock CC, Blanford S, Hughes GL, Rasgon JL, Thomas MB (2014) Temperature alters
Plasmodium blocking by Wolbachia. *Sci Rep* 4:3932.

18.   Ye YH, Carrasco AM, Dong Y, Sgro CM, McGraw EA (2016) The effect of temperature on
Wolbachia-mediated dengue virus blocking in Aedes aegypti. *Am J Trop Med Hyg*
94(4):812–819.

19.   Lambrechts L, et al. (2011) Impact of daily temperature fluctuations on dengue virus
transmission by Aedes aegypti. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.1101377108.

20.   Min KT, Benzer S (1997) Wolbachia, normally a symbiont of Drosophila, can be virulent,
causing degeneration and early death. *Proc Natl Acad Sci U S A* 94(20):10792–6.

21.   Reynolds KT, Thomson LJ, Hoffmann AA (2003) The effects of host age, host nuclear
background and temperature on phenotypic effects of the virulent Wolbachia strain popcorn
in Drosophila melanogaster. *Genetics* 164(3):1027–34.

22.   Strunov AA, Ilinskii YY, Zakharov IK, Kiseleva E V. (2013) Effect of high temperature on
survival of Drosophila melanogaster infected with pathogenic strain of Wolbachia bacteria.
Russ J Genet Appl Res. doi:10.1134/S2079059713060099.

23.   Truitt AM, Kapun M, Kaur R, Miller WJ (2019) Wolbachia modifies thermal preference in
Drosophila melanogaster. *Environ Microbiol*. doi:10.1111/1462-2920.14347.

24.   Versace E, Nolte V, Pandey RV, Tobler R, Schlötterer C (2014) Experimental evolution
reveals habitat-specific fitness dynamics among Wolbachia clades in Drosophila
melanogaster. *Mol Ecol* 23(4):802–14.

25.   Charlesworth J, Weinert LA, Araujo E V., Welch JJ (2019) Wolbachia, Cardinium and
climate: An analysis of global data. *Biol Lett*. doi:10.1098/rsbl.2019.0273.

26.   Frentiu FD, Robinson J, Young PR, McGraw EA, O'Neill SL (2010) Wolbachia-mediated
resistance to dengue virus infection and death at the cellular level. *PLoS One* 5(10):e13398.

27.   Lu P, Bian G, Pan X, Xi Z (2012) Wolbachia Induces Density-Dependent Inhibition to
Dengue Virus in Mosquito Cells. *PLoS Negl Trop Dis* 6(7):e1754.

28.   Kriesner P, Conner WR, Weeks AR, Turelli M, Hoffmann AA (2016) Persistence of a
Wolbachia infection frequency cline in Drosophila melanogaster and the possible role of
reproductive dormancy. *Evolution (N Y)* 70(5):979–997.

29.   Gora N V., et al. (2020) Climate Factors and Wolbachia Infection Frequencies in Natural
Populations of Drosophila melanogaster. *Cytol Genet*. doi:10.3103/S0095452720030044.

30.   van den Hurk AF, et al. (2012) Impact of Wolbachia on infection with chikungunya and
yellow fever viruses in the mosquito vector Aedes aegypti. *PLoS Negl Trop Dis* 6(11):e1892.

31.   Aliota MT, et al. (2016) The wMel strain of Wolbachia Reduces Transmission of Zika virus
by Aedes aegypti. *Sci Rep* 6:28792.

32.   Dutra HLC, et al. (2016) Wolbachia Blocks Currently Circulating Zika Virus Isolates in
Brazilian Aedes aegypti Mosquitoes. *Cell Host Microbe* 19(6):771–774.

33. Ulrich JN, Beier JC, Devine GJ, Hugo LE (2016) Heat Sensitivity of wMel Wolbachia during
Aedes aegypti Development. *PLoS Negl Trop Dis* 10(7):e0004873.

34. Ross PA, et al. (2020) Heatwaves cause fluctuations in wMel Wolbachia densities and
frequencies in Aedes aegypti. *PLoS Negl Trop Dis*. doi:10.1371/journal.pntd.0007958.

35. Lau M-J, Ross PA, Endersby-Harshman NM, Hoffmann AA (2020) Impacts of Low
Temperatures on Wolbachia (Rickettsiales: Rickettsiaceae)-Infected Aedes aegypti
(Diptera: Culicidae). *J Med Entomol*. doi:10.1093/jme/tjaa074.

36. Rainey SM, et al. (2016) Wolbachia Blocks Viral Genome Replication Early in Infection
without a Transcriptional Response by the Endosymbiont or Host Small RNA Pathways.
PLoS Pathog 12(4). doi:10.1371/journal.ppat.1005536.

37. Schultz MJ, et al. (2018) Wolbachia wStri blocks Zika virus growth at two independent
stages of viral replication. *MBio*. doi:10.1128/mBio.00738-18.

38. Ekwudu O, Devine GJ, Aaskov JG, Frentiu FD (2020) Wolbachia strain wAlbB blocks
replication of flaviviruses and alphaviruses in mosquito cell culture. *Parasites and Vectors*.
doi:10.1186/s13071-020-3936-3.

39. Corbin C, Heyworth ER, Ferrari J, Hurst GDD (2017) Heritable symbionts in a world of
varying temperature. *Heredity (Edinb)*. doi:10.1038/hdy.2016.71.

40. Doremus MR, et al. (2018) Breakdown of a defensive symbiosis, but not endogenous
defences, at elevated temperatures. *Mol Ecol*. doi:10.1111/mec.14399.

41. Corbin C, Jones JE, Chrostek E, Fenton A, Hurst GDD (2020) Thermal sensitivity of the
Spiroplasma-Drosophila hydei protective symbiosis: The best of
climes, the worst of climes. *bioRxiv*:2020.04.30.070938.

42. Heyworth ER, Ferrari J (2016) Heat stress affects facultative symbiont-mediated protection
from a parasitoid wasp. *PLoS One*. doi:10.1371/journal.pone.0167180.

43. Ryder E, et al. (2004) The DrosDel collection: a set of P-element insertions for generating
custom chromosomal aberrations in Drosophila melanogaster. *Genetics* 167(2):797–813.

44. Riegler M, Sidhu M, Miller WJ, O'Neill SL (2005) Evidence for a global Wolbachia
replacement in Drosophila melanogaster. *Curr Biol* 15(15):1428–33.

45. Chrostek E, Teixeira L (2015) Mutualism breakdown by amplification of Wolbachia genes.
PLoS Biol 13(2):e1002065.

46. Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR.
Nucleic Acids Res 29(9):e45.

47. Pais IS, Valente RS, Sporniak M, Teixeira L (2018) Drosophila melanogaster establishes a
species-specific mutualistic interaction with stable gut-colonizing bacteria. *PLoS Biol*.
doi:10.1371/journal.pbio.2005710.

48. Team RC (2019) R: A language and environment for statistical computing. *R Found Stat
Comput Vienna, Austria*.

49. Therneau T (2012) coxme: mixed effects Cox models. R package version 2.2-3.
www.cran.R-project.org/package=coxme. *Oikos*. doi:10.1111/oik.01149.

50. Bates D, Mächler M, Bolker BM, Walker SC (2015) Fitting linear mixed-effects models using
lme4. *J Stat Softw*. doi:10.18637/jss.v067.i01.

51. Lenth R V. (2016) Least-squares means: The R package lsmeans. *J Stat Softw*.
doi:10.18637/jss.v069.i01.

52. Wickham H (2016) *ggplot2 Elegant Graphics for Data Analysis (Use R!)* doi:10.1007/978-
0-387-98141-3.

570
571
572

Relative DCV levels
(log10)

Wolb–Wolb+ *Wolb–Wolb+*
18°C–18°C 25°C–18°C

Fig. S1. Pre-infection temperature determines viral titres. DCV titres measured by RT-qPCR 3 dpi in *Wolb+* and *Wolb-* flies infected with 10^8 TCID$_{50}$/ml DCV. Flies were kept at 18°C or 25°C before the infection and at 18°C after the infection. Each dot is a sample, each sample consists of ten flies. Horizontal lines are medians of the replicates. This is a replicate of experiment shown in Fig. 2C.

584
Figure S2. Pre-infection temperature determines *Wolbachia*-conferred antiviral protection in
different *Wolbachia* and *Drosophila* genotypes and against different viruses. (A) Aljezur 1
and (B) Oregon-R (W-20) *Wolb+* and *Wolb-* flies, fifty per *Wolbachia* status per condition, were
pricked with three dilutions of DCV. (C) w^{1118} *iso* flies, fifty per *Wolbachia* status per condition, were
infected with three doses of FHV. Flies were kept at 25°C or 18°C before the infection, and at 18°C
after the infection. Mortality was recorded daily. This is a replicate of experiment shown in Fig. 3.
591

593
594
595 **Fig. S3. Wolbachia protects flies kept at a temperature cycling daily between 18 and 25ºC.**
596 Fifty *Wolb+* and *Wolb-* flies were raised with the temperature cycling daily between 25ºC (midday)
597 and 18ºC (midnight), infected with DCV (10^8 TCID$_{50}$/ml) and placed at the same cycling temperature
598 after infection. Mortality was recorded daily.
599

600
601

Figure S4. Developmental temperature determines *Wolbachia*-conferred antiviral protection.
Wolb+ and *Wolb-* flies, fifty per *Wolbachia* status per condition per replicate, were raised from egg
to adult at 18 or 25ºC (Dev), collected at the age of 0-1 days, aged for 3 days at an 18 or 25ºC
(Aging), pricked with DCV (10^8 TCID$_{50}$/ml), and placed at 18ºC. Mortality was recorded daily. Three
replicates of the same experiment are shown.

607
608
Figure S5. *Wolbachia* presence in adults is required for antiviral protection. (A) DCV titres in
Wolb+ and *Wolb-* flies raised at 25°C and kept at 18°C after DCV infection with two doses, 10^8 and
10^9 TCID$_{50}$/ml, and at two time points post infection, 12 and 24h. Relative DCV levels were
determined by RT-qPCR. Each dot is a sample consisting of 10 flies, lines are medians. (B) FHV
titres in *Wolb+* and *Wolb-* flies raised at 25°C and kept at 25°C after infection. Each dot is a sample
consisting of 10 flies, lines are medians, and time 0 corresponds to the time of infection. (C)
Wolbachia levels in single flies raised at 25°C, at eclosion (day 0), after 10 days of different
antibiotic/ control treatments at 25°C (day 10), and after additional 20 days of treatment at 18°C
(day 30) measured by qPCR. This is the replicate of the experiment shown in Fig. 4B. (D) Colony
forming units (CFUs) in dissected *Wolb+* fly guts at the day 30 of each antibiotic or control
treatment. Each dot is a pool of three guts. (E) Survival of *Wolb+* and *Wolb-* flies, fifty per *Wolbachia*
status per treatment, developed at 25°C, collected at eclosion, fed antibiotics-containing food for
10 days at 25°C, infected with DCV (10^8 TCID$_{50}$/ml), and placed at 18°C on antibiotic-containing
food. Mortality was recorded daily. This is the replicate of the experiment shown in Fig. 4C.
623

624 **Supplementary Information**
625
626 **SI Text S1 (text_s1.txt)**
627   Rmd script for statistical analysis
628
629 **SI Text S2 (text_s2.pdf)**
630   Output of statistical analysis
631
632 **SI Dataset S1 (dataset_s1.txt)**
633   Data for figures 1A, B
634
635
636 **SI Dataset S2 (dataset_s2.txt)**
637   Data for figure 1C
638
639 **SI Dataset S3 (dataset_s3.txt)**
640   Data for figure 1D
641
642 **SI Dataset S4 (dataset_s4.txt)**
643   Data for figures 2A, B, and S3
644
645 **SI Dataset S5 (dataset_s5.txt)**
646   Data for figures 2C and S1
647
648 **SI Dataset S6 (dataset_s6.txt)**
649   Data for figure 2D
650
651 **SI Dataset S7 (dataset_s7.txt)**
652   Data for figures 3A, B, and S2A, B
653
654 **SI Dataset S8 (dataset_s8.txt)**
655   Data for figures 3C and S2C
656
657 **SI Dataset S9 (dataset_s9.txt)**
658   Data for figure S4
659
660 **SI Dataset S10 (dataset_s10.txt)**
661   Data for figure 4A
662
663 **SI Dataset S11 (dataset_s11.txt)**
664   Data for figure S5A
665
666 **SI Dataset S12 (dataset_s12.txt)**
667   Data for figure S5B
668
669 **SI Dataset S13 (dataset_s13.txt)**
670   Data for figures 4B and S5C
671
672 **SI Dataset S14 (dataset_s14.txt)**
673   Data for figures 4C and S5E
674
675 **SI Dataset S15 (dataset_s15.txt)**
676   Data for figure S5D
677