3D Visualization of Macromolecule Synthesis

Timothy J. Duerr[1], Ester Comellas[2,3], Eun Kyung Jeon[1], Johanna E. Farkas[1], Marylou Joetzjer[4], Julien Garnier[4], Sandra J. Shefelbine[3,5], James R. Monaghan[1,*]

[1]Department of Biology, Northeastern University, Boston, MA 02115, USA

[2]Department of Mathematics, Laboratori de Càlcul Numeric (LaCàN), Universitat Politècnica de Catalunya (UPC), 08034 Barcelona, Spain

[3]Mechanical and Industrial Engineering, Northeastern University, Boston, MA 02115, USA

[4]University of Technology of Compiègne, Rue Roger Couttolenc, 60200 Compiègne, France

[5]Department of Bioengineering, Northeastern University, Boston, MA, USA

* Author for correspondence (j.monaghan@northeastern.edu)

Abstract

Measuring nascent macromolecular synthesis *in vivo* is key to understanding how cells and tissues progress through development and respond to external cues. Here, we perform *in vivo* injection of alkyne- or azide-modified analogs of thymidine, uridine, methionine, and glucosamine to label nascent synthesis of DNA, RNA, protein, and glycosylation. Three-dimensional volumetric imaging of nascent macromolecule synthesis was performed in axolotl salamander tissue using whole mount click chemistry-based fluorescent staining followed by light sheet fluorescent microscopy. We also developed an image processing pipeline for segmentation and classification of morphological regions of interest and individual cells, and we apply this pipeline to the regenerating humerus. We demonstrate our approach is sensitive to biological perturbations by measuring changes in DNA synthesis after limb denervation. This method provides a powerful means to quantitatively interrogate macromolecule synthesis in heterogenous tissues at the organ, cellular, and molecular levels of organization.

Introduction

The measurement of nascent DNA, RNA, and protein synthesis in animals provides critical information on the state of cells (dividing, growing, dying) in relation to their surrounding cells. Traditionally, radiolabeled or brominated nucleosides or amino acids are introduced to live animals, where they are incorporated into macromolecules during DNA synthesis[1,2], trancription[3,4], and translation[5]. Performing such approaches have allowed the characterization of cells actively undergoing macromolecular synthesis as well as quantification of synthesis rates, which facilitates the study of cell behavior during tissue remodeling, proliferation, stress, or disease [6-9].

In the past decade, bio-orthogonal macromolecule precursor analogs including 5-ethynyl-2′-deoxyuridine (EdU), 5-ethynyl-uridine (5-EU), and L-azidohomoalanine (AHA) have become commercially available. After injection of macromolecule precursor analogs into animals, precursors can later be detected in nascent DNA, RNA, and protein macromolecules with fluorescently labeled azides or alkynes through highly selective copper catalyzed azide-alkyne cycloaddition ("click") chemistry[10-12]. These powerful new analogs provide an alternative to the use of dangerous isotopes and the challenges associated with brominated precursors such as the

requirement of large secondary antibodies (~150kd) and harsh tissue retrieval methods that limit

their use in whole tissue samples.

Advantages of the click-chemistry labeling approach include the inert nature of macromolecule

precursor analogs that have minimal impact on the animal, the small size of fluorescently labeled

alkynes and azides, and the high selectivity of click-chemistry. These advantages have enabled

whole mount fluorescent labeling of DNA synthesis[11], RNA transcription[13], protein translation[14],

and glycans[15,16] in animals. These pioneering proof of principle experiments have demonstrated

that imaging macromolecular synthesis is possible, but the fact that most model organisms are

large, optically opaque, and consist of heterogenous tissues has made it a challenge to image

biological phenomena in deep tissues. Challenges such as photon penetration, differences in

refractive indices among different cellular components, light-induced photodamage, and

background fluorescence have limited the use of whole mount imaging of macromolecular

synthesis.

Advances in light sheet fluorescence microscopes (LSFMs) have recently enabled the imaging of

large biological specimens from millimeters to several centimeters in size[17]. LSFMs have been

utilized for volumetric imaging of many varieties including visualization of mRNA in whole

mount fluorescence *in situ* hybridization experiments[18] and *in vivo* interrogation of deep tissue

dynamics in transgenic reporter animals[19] among others[17]. Clearing methods like CLARITY[20],

CUBIC[21], and 3DISCO[22] have further enabled volumetric imaging by decreasing refractive index

mismatches and tissue clearing to advance large specimen imaging even further. Together, the

rise of three-dimensional imaging, new staining techniques, and tissue clearing has demanded

new means for cell counting, segmentation, and fluorescence quantification.

Here, we present a click-chemistry based method to visualize DNA synthesis, transcription,

translation, and protein glycosylation in whole mount samples using LSFM (Figure 1A). We

demonstrate the utility of this technique by imaging macromolecular synthesis in the

regenerating axolotl salamander limb. Following limb amputation, the axolotl regenerates its

limb by generating a mass of proliferating cells at the limb stump called a blastema[23]. The

blastema is an ideal environment to test our method because it is an accessible, heterogenous

tissue that increases DNA synthesis, transcriptional output, and translation rates compared to

uninjured tissue. Furthermore, DNA, RNA, and protein synthesis decrease after denervation of

the regenerating limb[24,25].

We use the regenerating limb to demonstrate that two click-it ready precursors can be

administered and subsequently visualized simultaneously in a single sample. We also show that

optical resolution of images can be improved with the clearing agent 2,2'-thiodiethanol (TDE),

and that this method works in a number of tissues. We outline an image analysis pipeline for

three dimensional (3D) morphology segmentation, cell counting, and fluorescence quantification

of stained tissues. We apply this pipeline to the regenerating humerus to demonstrate the

multiscale quantitative analysis capabilities of our method. Finally, we show that our method is

sensitive enough to detect and quantify changes in DNA synthesis rates in whole mount

innervated and denervated regenerating limbs. Taken together, our method provides a unique

approach to simultaneously interrogate cell state at the organ, cellular and molecular levels of

organization.

Materials and methods

Animal procedures

Axolotls (*Ambystoma mexicanum*: d/d RRID Catalog #101L) were either bred in captivity at

Northeastern University or purchased from the Ambystoma Genetic Stock Center at the

University of Kentucky. Experiments were performed in accordance with Northeastern

University Institutional Animal Care and Use Committee. Animals were grown to 4-6cm (Mean

5.3cm, SD 0.36) and 1-1.5g (Mean 1.3g, SD 0.19g) for use in all studies. For all experiments,

animals were anesthetized by treatment of 0.01% benzocaine until visually immobilized. Limbs

were amputated either at the distal end of the zeugopod or midway through the stylopod, and

bones were trimmed below the amputation plane to allow for uniform growth. At the date of

collection, animals were reanaesthetized and injected with either EdU (8.0ng/g animal), 5-EU

(270.0µg/g animal), AHA (180.59µg/g animal), or N-azidoacetylglucosamine-tetraacylated

(GlcNAz) (4.3µg/g animal) alone or simultaneously with the following combinations:

EdU/AHA, EdU/GlcNAz, 5-EU/AHA, 5-EU/GlcNAz. All monomer analogs were purchased

from www.clickchemistrytools.com and resuspended in DMSO at the following concentrations:

EdU- 300mM, 5-EU- 100mM, AHA- 100mM, GlcNAz- 100mM. Stocks were further diluted in

1X phosphate buffered saline (PBS) for injection. Three hours after injection, limbs were

collected from the upper stylopod and fixed in 4% paraformaldehyde (PFA) (diluted in 1X PBS)

at 4°C overnight. If limbs were denervated, the nerve supply was severed at the brachial plexus

24 hours prior to tissue collection.

Whole mount click-it protocol

Following fixation in 4% PFA, samples were washed three times with 1X PBS at room

temperature (~23°C) for 5 minutes. Samples were dehydrated in an increasing methanol series at

room temperature starting with 25% methanol (diluted in 1X PBS), 50% methanol, 75%

methanol, and 100% methanol for 5 minutes at each step. Samples could then be stored in 100%

methanol indefinitely at -20°C. For staining, samples were rehydrated in a decreasing methanol

series starting with 75% methanol (diluted in 1X PBS), 50% methanol, 25% methanol, and

finally placed in 100% 1X PBS for 5 minutes at each step. Samples were then washed 3 times

with 1X PBST (1X PBS with 1% Triton) for 5 minutes at room temperature. To aid in clearing,

samples were washed in 0.5% trypsin (diluted in 1X PBST) for 30-90 minutes on a rocker at

room temperature, or until sample appeared translucent. Samples were washed three times at

room temperature for 5 minutes with deionized water, then washed in 100% acetone for 20

minutes at -20°C and washed with deionized water again for 10 minutes. Samples were washed

in 1X PBST three times at room temperature for 5 minutes prior to applying click-it cocktail for

30 minutes at room temperature. The click-it cocktail was made in 1X TRIS buffered saline as

follows: 50µL 100mM Sodium Ascorbate, 20µL 4mM $CuSO_4$, and 2µL 500µM azide- or

alkyne-modified Alexa Flour, combined in order as listed. After the first round of staining,

samples were washed at room temperature 6 times for 30 minutes with rocking. For double-

labelling, samples were again placed in the click-it cocktail with a different fluorescent dye to

stain for the second analog at room temperature overnight. Both rounds of staining were

conducted in the dark to prevent photodegradation of fluorescent molecules. For staining with

DAPI, samples were washed 3 times for 5 minutes with 1X PBS, then placed in 277mM DAPI

for 4 days at room temperature. Samples were washed 3 times for 20 minutes with 1X PBS and

left in 1X PBS at 4°C for short term storage prior to imaging with LSFM. Whole mount samples

were cleared with 67% TDE (diluted in deionized water) overnight at room temperature in the

dark.

Tissue section click-it protocol

Following fixation in 4% PFA, samples were washed three times in 1X PBS each for 5 minutes,

and cryoprotected in 30% sucrose on a rocker for 1 hour or until the tissue fully sinks. Samples

were removed form sucrose and briefly washed in optimal cutting temperature (OCT) compound

before mounting in OCT compound and frozen at -80°C. 10µm sections were obtained using a

cryostat and baked at 65°C for 15 minutes. Slides were then washed with water for 30 minutes at

room temperature to remove residual OCT. Slides were washed once with 1X PBS for 5 minutes

at room temperature. The click-it cocktail (same as above) was applied to the slides and

incubated at room temperature in the dark for 30 minutes. If staining for a second

macromolecule, slides were washed five times for 5 minutes with 1X PBS at room temperature.

The samples were then stained for 30 minutes at room temperature in the dark using the above

click-it cocktail with a different a fluorophore dye. Following the final click-it reaction, slides

were washed once with 1X PBS at room temperature for 5 minutes, then stained with 277mM

DAPI for 5 minutes at room temperature. Slides were washed again with 1X PBS for 5 minutes

at room temperature and water for 5 minutes at room temperature and mounted with SlowFade™

Gold Antifade Mountant (Thermo S36936). Slides were imaged using a Zeiss LSM800 confocal

microscope.

Light sheet microscopy

All 3D images were acquired using a Zeiss light sheet Z.1 microscope paired

with Zen software. Unless otherwise indicated, samples were cleared and imaged in 67% TDE.

Post-processing for visualization purposes was performed with Arivis Vision4D v3.1.4 on

a workstation with a 64-bit Windows Embedded Standard operating system, and an Intel(R)

Xeon(R) CPU E5-2620 v3 @ 2.40GhZ (2 processors), 128GB RAM, and NVIDIA Quadro

K2200 GPU. Sub-volumes were stitched together with the Tile Sorter in Arivis, using the manual

projection option. Volume fusion was performed through automatic landmark registration of

manually selected points for alignment. Finally, constant background intensity was corrected

using the automatic functionality.

Data analysis

All data were processed on desktop computers with the aid of Fiji[26] and Matlab[27]. The custom

Fiji scripts and Matlab codes used are available in the supplementary material. We performed all

analyses on unprocessed .czi files acquired directly from the light sheet microscope.

<u>Organ level</u>: The goal of the organ level analysis was to determine gross morphology of the

organ, such as area, principal axes, and maximum diameters in said axes for each cross section.

Critical to this step is alignment of the images to a standard defined axis and reslicing in the

transverse direction (perpendicular to the long axis of the humerus). The image stack of a

regenerating axolotl elbow was imported into Fiji and aligned along the proximodistal axis of the

humerus (Figure 3A). Morphological segmentation of the humerus was performed semi-

automatically with the Segmentation Editor plugin (Figure 3B). The segmented surface was then

exported from Fiji as a mesh via an .stl file using 3D Viewer[28]. Volumetric analysis was

performed in Fiji by reslicing the aligned mask (Figure 3B) to obtain a stack of cross-sections

perpendicular to the proximodistal axis, and then quantified the shape and volume with the Slice

Geometry option in the BoneJ[29] Fiji plugin. Slice Geometry calculates cross-sectional geometric

properties of shapes, including area, second moment of area around the major and minor axes,

and maximum chord length from these major and minor axes.

<u>Cellular level</u>: The goal of the cellular level analysis was to determine the number of

proliferating cells (EdU$^+$) as a function of proximodistal position along the humerus, as well as

average size, shape, and orientation. The aligned image stack (Figure 3A) was cropped using the

morphological segmentation (Figure 3B) as a mask with the aid of the Image Calculator in Fiji.

Based on the EdU staining (Figure 3C), we segmented the nuclei of the proliferating cells with

Trainable Weka Segmentation 3D[30]. We used a combination of filters available in Fiji before and

after training the algorithm to improve segmentation results. Filters applied include the 3D Edge

and Symmetry, Background Subtraction, 3D Fill Holes, Gaussian Blur 3D, and 3D Watershed

Split. A detailed description of the sequence and parameters used is available in the

supplementary materials. The 3D Objects Counter then provided a list of identified cells as well

as their volume and position, among other information. In addition, the surfaces of the

segmented cell nuclei were exported and processed in MeshLab, similarly to the process

followed with the organ segmentation, to be visualized in Matlab (Figure 3E). The centroid

coordinates and corresponding object volumes identified in Fiji were imported into Matlab for

further processing and plotting. We computed the number of cell centroids within 50μm thick

slices perpendicular to the proximodistal axis to obtain a density-like measure of proliferating

cells. Slice thickness was selected as slightly larger than the average cell diameter to ensure cells

were not sampled across more than two slices.

Molecular level: The goal of the molecular level analysis was to determine molecular activity,

characterized by fluorescent signal, which in our study represented DNA replication (EdU). The

aligned and cropped image stack used as a starting point of the cellular-level analysis (Figure

3C) was also analyzed at the molecular level. We resliced the image stack along the

proximodistal axis in the organ under study (Figure 3F). The histogram and pixel intensity

statistics were listed for each slice via the getRawstatistics function. These allow for further

quantification of the fluorescent signal in Matlab, e.g. we calculated the mean intensity of each

plane along the proximodistal axis and the histogram of the whole organ. For the example

showing nerve-dependent regeneration in axolotl blastemas (Figure 6A), we compared overall

pixel intensity of the left (denervated) and right (control, innervated) forelimbs of the same

animal. To ensure full repeatability of our data analysis, we selected the same cubic volume in all

blastemas processed: a cube with 175μm sides, centered along the proximal-distal axis and at a

distance of 250μm from the distal tip (Figure 6A). The cube size was adjusted to maximize the

cube volume for all blastemas processed, while ensuring that the entire cube was contained

within the blastema. Histograms of each cubic volume and the ratio between the innervated and

denervated limb of the same animal were computed and plotted with Matlab (Figure 6B) to

quantify the changes in proliferation in innervated versus denervated limbs at different stages of

regeneration. All scripts for blastema cube quantification are provided in the supplementary

materials.

Results and discussion

Whole mount, click-it based visualization of macromolecule synthesis

To visualize macromolecule synthesis, we injected click-it compatible monomer analogs (Table

1) intraperitonially three hours prior to sample collection. During this time, analogs

metabolically incorporated into nascently synthesized macromolecules, resulting in *in vivo*

labeling of macromolecule synthesis. These labeled macromolecules contain either azide- or

alkyne-modified monomers that can be detected with click-it compatible fluorescent molecules,

enabling imaging of nascent macromolecules in whole mount tissues with LSFM (Figure 1). To

demonstrate the multiplexing capabilities of the approach, modified monomer analogs with

disparate functional groups (EdU/AHA, 5-EU/AHA, EdU/GlcNAz, 5-EU/GlcNAz) were co-

injected and visualized with LSFM (Figure 2A-D, Supplementary videos 1-2). Whole mount

samples were comparable to 2D longitudinal tissue sections of the same stains (Figure 2I-L),

showing that our method generates similar results in both whole mount and tissue sections

(Supplementary Figure 1). We demonstrate the specificity of N-Acetylglucosamine (GlcNAc)

incorporation by showing that a GlcNAz-specific antibody pretreated on tissue sections collected

from GlcNAz injected animals prevented the subsequent click-it reaction (Supplementary Figure

2).

An advantage of our method is that staining whole mount tissues eliminates the need for

sectioning, reducing the potential inconsistences that arise as a result of the sectioning process

(uneven tissue, different cutting planes, etc.). Additionally, traditional methods of obtaining 3D

images of thick tissues with confocal microscopy are impractically slow when imaging hundreds

of images in a single stack. LSFM allows for more rapid imaging of whole samples, requiring

only minutes to image each sample. However, several considerations exist when imaging whole

tissue in 3D with LSFM. Stain penetrates slower in whole 3D tissues compared to 10-20µm thick

tissue sections, requiring longer staining times. An advantage of our method is that the click

reaction requires molecularly diminutive reagents that readily pass through cell and nuclear

membranes, ensuring stain penetration in the center of dense tissues. Different refractive indices

between disparate cellular components and the imaging media cause light to scatter, which can

reduce the resolution and brightness of 3D images[31]. To improve image resolution and light

penetration, overnight refractive index matching with 67% 2,2'-thiodiethanol (TDE) can

sufficiently clear axolotl limbs for imaging with LSFM. This clearing method rapidly and

effectively improves the signal to noise ratio of stained samples compared to imaging in PBS

(Supplementary Figure 3). With careful attention to these challenges, our method provides a

means to obtain high quality 3D images from tissues 1mm in depth in less than 10 minutes per

sample (Figure 2A-H) with clear, consistent staining.

3D, multiscale quantitative analysis of the regenerating humerus

To demonstrate that our whole mount click-it method can obtain quantifiable data on the organ,

cellular, and molecular levels of organization, we applied our technique to the regenerating

humerus. After regenerating for 35 days, axolotls with mid-humeral amputations were injected

with EdU/AHA to identify cells within the humerus undergoing DNA synthesis (EdU) and

protein translation (AHA). We observed EdU staining in chondrocytes distal to the amputation

plane and AHA staining in the humerus perichondrium (Figure 3A). We outline a multiscale,

quantitative pipeline that leverages the staining patterns of these macromolecules for analysis of

3D humerus morphology and 3D macromolecule synthesis. This workflow combines available

plugins in Fiji[26] and scripts developed in Fiji and Matlab[27] for the data analysis process (see

supplementary information for a detailed description).

For 3D organ level analysis of macromolecule synthesis, AHA staining provided an adequate

outline of the humerus, allowing us to segment its 3D morphology (Figure 3A-B). We quantified

organ shape by assessing cross-sectional area (Figure 4A') and circularity (Figure 4A") of the

segmented humerus along the proximodistal axis with BoneJ[29] (Figure 4A). These measurements

of 3D organ size can be obtained with other methods such as microCT and focused ion beam

scanning electron microscopy (FIB-SEM). However, microCT is unable to image every tissue

due to stain limitations and uses hazardous radiation, while FIB-SEM can capture 3D surface

topography but not the entire organ morphology. Our method has the capability to fully image

the 3D structure of entire organs (size permitting) without the need for radiation while

simultaneously capturing information from both the cellular and molecular levels of

organization.

On the cellular level, we segmented the 3D morphology of proliferating chondrocytes based on

EdU staining using the Trainable Weka Segmentation 3D plugin[30] (Figure 4B"). With this

segmentation, we identify highly condensed regions of cells undergoing rapid rates of DNA

synthesis (Figure 4B). From these data, we observe cells synthesizing DNA most abundantly

distal to the plane of amputation, as expected for dividing chondrocytes. Traditionally, cell

quantification as such is conducted on 2D tissue sections. In heterogenous tissues, however, cells

are distributed non-uniformly; 2D sampling may not accurately capture the cellular distribution

and cannot be used to determine cell volume or shape. Our whole mount staining method allows

quantification of cells within an entire 3D tissue, resulting in a more accurate assessment of cell

density within heterogenous tissues.

To demonstrate quantitative molecular analysis of 3D macromolecule synthesis, we assessed

staining intensity of EdU in slices along the proximodistal axis of the regenerating humerus (Fig

4C). EdU intensity represents the rate at which a cell undergoes DNA synthesis, which is one of

the first steps of cell division and proliferation. Proliferating cells rapidly synthesize DNA,

enabling cells to integrate EdU into nascent DNA strands, providing ample opportunities for

covalent linkage of fluorescent molecules to the DNA strand. Thus, higher pixel values within

EdU$^+$ cells are representative of faster DNA synthesis rates. These data show that EdU intensity

is strongest in the regions distal to the amputation plane of the humerus (Figure 4C'), providing a

quantitative measure to assess macromolecule synthesis on the molecular level. Within a tissue,

cells synthesize macromolecules heterogeneously, reflected by different fluorescence values

between cells. Thus, quantifying macromolecule synthesis based on presence/absence of signal

instead of fluorescence does not account for variability in macromolecule synthesis rates among

cells. To compound this issue, quantifying fluorescence in tissue sections only represents the rate

of macromolecule synthesis from a fraction of cells in larger, heterogenous tissues. Our method

provides a means to capture this molecular heterogeneity in 3D samples, allowing us to observe

whole 3D regions in the regenerating humerus that synthesize DNA more rapidly than others,

which further demonstrates the utility of our whole mount click staining method.

Taken together, these results demonstrate that our method can provide quantifiable data on the

organ, cellular, and molecular levels of organization. This highlights the novelty of our method,

as we have not found previous examples of multiscale analysis as outlined here. We foresee this

multiscale, quantitative analysis having broad applications in the examination of dynamic cell

processes in 3D, such as in cancer metabolism and mammalian neurogenesis or other fields

where macromolecule synthesis is traditionally studied in tissue sections.

3D, molecular analysis of biological perturbations

To demonstrate that our method is sensitive enough to detect subtle changes in macromolecule

synthesis in vivo, we quantified the difference in EdU intensity between limbs regenerating with

and without a nerve supply. Cell proliferation in regenerating limbs is sustained by factors

secreted from nerves that innervate the limb[32,33]. Therefore, we amputated both forelimbs at the

mid-humerus, and the left limb was denervated at the brachial plexus one day prior to collection.

At 6, 9, 12, 15, 18, and 25 days post amputation (dpa) animals were pulsed with EdU for 3 hours

before collection to label proliferating blastema cells (Figure 5A). LSFM was used to image

samples (Figure 5B), ensuring pixel resolution was consistent between samples. We quantified

DNA synthesis in denervated limbs compared to innervated limbs by creating a 175x175x175µm

cube 250µm from the distal most tip of the blastema (Figure 6A). From these results, we

observed a marked decrease in blastema EdU incorporation due to denervation at 9, 12, and
15dpa (Figure 6B), demonstrating that our whole mount staining approach is capable of detecting
changes in macromolecule synthesis after biological perturbations. One potential limitation of
our method in the blastema is the inability to preform single cell segmentation. We predict that
imaging with higher magnification may overcome this limitation but will significantly increase
imaging time and file size.

Conclusions

The work presented here provides a fast, simple pipeline for visualizing macromolecule turnover
in the 3D space of whole tissues. While tissue sections were previously the standard for studying
these cellular processes, a new standard in the field must be expected where dynamic processes
like macromolecule synthesis are visualized in 3D to obtain a more complete understanding of
how these processes occur in a larger tissue context. Few modalities of imaging exist to provide
this level of analysis. Here, we outline a method to study macromolecule synthesis at the organ,
cellular, and molecular levels of organization, which is important in understanding cell state and
how cell state affects neighboring cells and tissues. To this end, we show that DNA synthesis,
transcription, translation, and protein glycosylation increase in the entire 3D space of the
regenerating blastema after limb amputation and that these processes can be visualized
concurrently. We outline a multiscale pipeline for analysis and quantification of heterogeneous
tissues at the organ, cellular, and molecular levels of organization. Additionally, we demonstrate
that our method is sensitive to detect biological perturbations by showing a decrease in DNA
synthesis in the blastema following limb denervation. We foresee our method being used to
similarly readdress other classical questions with modern techniques for a more exhaustive
understanding of biological processes; traditional questions within the fields of cancer biology
and neurobiology may especially benefit from a technology as such. Additionally, as more click-
it ready macromolecules monomer analogs are generated, our method will provide a means to
study more biological processes in whole tissues.

Acknowledgements

The authors would like to acknowledge Tyler Jensen for his early assistance in developing the
whole mount staining protocol, and both Alex Lovely and Gouxin Rong for their microscopy and
image analysis assistance. Images obtained from the Harvard University Center for Biological

Imaging and the Northeastern University Chemical Imaging of Living Systems core. We
acknowledge animal support from the Ambystoma Genetic Stock Center funded by NIH grant
P40-OD019794.

Author Contributions

334 SJS and JRM conceived and supervised the study. TJD, EKJ, and JEF preformed experiments
335 and imaging. EC, MJ, and JG designed and implemented codes for multiscale data analysis. TJD
336 and EC performed the analysis. TJD, EC, SJS, and JRM wrote the paper. All authors approved
337 the final version of the manuscript.

Funding

339 This project has received funding from the National Science Foundation under grant #1727518
340 to SJS, #1656429 and #1558017 to JRM. E.C acknowledges funding from the European Union's
341 Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant
342 agreement No. 841047. EKJ acknowledges support from a Northeastern University Matz
343 Scholarship and Undergraduate Research Fellowship.

Competing interests

345 The authors declare no competing or finical interests.

References

1   Sidman, R. L., Miale, I. L. & Feder, N. Cell proliferation and migration in the primitive
ependymal zone: an autoradiographic study of histogenesis in the nervous system. *Exp*
Neurol **1**, 322-333, doi:10.1016/0014-4886(59)90024-x (1959).

2   Gratzner, H. G. Monoclonal antibody to 5-bromo- and 5-iododeoxyuridine: A new
reagent for detection of DNA replication. *Science* **218**, 474-475,
doi:10.1126/science.7123245 (1982).

3   Uddin, M., Altmann, G. G. & Leblond, C. P. Radioautographic visualization of
differences in the pattern of [3H]uridine and [3H]orotic acid incorporation into the RNA
of migrating columnar cells in the rat small intestine. *J Cell Biol* **98**, 1619-1629,
doi:10.1083/jcb.98.5.1619 (1984).

4   Wansink, D. G. *et al.* Fluorescent labeling of nascent RNA reveals transcription by RNA
polymerase II in domains scattered throughout the nucleus. *J Cell Biol* **122**, 283-293,
doi:10.1083/jcb.122.2.283 (1993).

5   Garlick, P. J., McNurlan, M. A. & Preedy, V. R. A rapid and convenient technique for
measuring the rate of protein synthesis in tissues by injection of [3H]phenylalanine.
Biochem J **192**, 719-723, doi:10.1042/bj1920719 (1980).

6   Yoshizawa, F., Nagasawa, T., Nishizawa, N. & Funabiki, R. Protein synthesis and
degradation change rapidly in response to food intake in muscle of food-deprived mice. *J*
Nutr **127**, 1156-1159, doi:10.1093/jn/127.6.1156 (1997).

7   Rombouts, Y. *et al.* Acute phase inflammation is characterized by rapid changes in
plasma/peritoneal fluid N-glycosylation in mice. *Glycoconj J* **33**, 457-470,
doi:10.1007/s10719-015-9648-9 (2016).

8   Brien, T. & Lis, J. T. Rapid changes in Drosophila transcription after an instantaneous
heat shock. *Mol Cell Biol* **13**, 3456, doi:10.1128/MCB.13.6.3456 (1993).

9   Rose, R. J., Cran, D. G. & Possingham, J. V. Changes in DNA synthesis during cell
growth and chloroplast replication in greening spinach leaf disks. *J Cell Sci* **17**, 27-41
(1975).

10   Kolb, H. C., Finn, M. G. & Sharpless, K. B. Click Chemistry: Diverse Chemical Function
from a Few Good Reactions. *Angew Chem Int Ed Engl* **40**, 2004-2021, doi:10.1002/1521-
3773(20010601)40:11<2004::aid-anie2004>3.3.co;2-x (2001).

11   Salic, A. & Mitchison, T. J. A chemical method for fast and sensitive detection of DNA
synthesis in vivo. *Proc Natl Acad Sci U S A* **105**, 2415-2420,
doi:10.1073/pnas.0712168105 (2008).

12   Best, M. D. Click chemistry and bioorthogonal reactions: unprecedented selectivity in the
labeling of biological molecules. *Biochemistry* **48**, 6571-6584, doi:10.1021/bi9007726
(2009).

13   Jao, C. Y. & Salic, A. Exploring RNA transcription and turnover in vivo by using click
chemistry. *Proc Natl Acad Sci U S A* **105**, 15779-15784, doi:10.1073/pnas.0808480105
(2008).

14   Hinz, F. I., Dieterich, D. C., Tirrell, D. A. & Schuman, E. M. Non-canonical amino acid
labeling in vivo to visualize and affinity purify newly synthesized proteins in larval
zebrafish. *ACS Chem Neurosci* **3**, 40-49, doi:10.1021/cn2000876 (2012).

15   Sawa, M. *et al.* Glycoproteomic probes for fluorescent imaging of fucosylated glycans in
vivo. *Proc Natl Acad Sci U S A* **103**, 12371-12376, doi:10.1073/pnas.0605418103 (2006).

16   Laughlin, S. T. & Bertozzi, C. R. Imaging the glycome. *Proc Natl Acad Sci U S A* **106**,
12-17, doi:10.1073/pnas.0811481106 (2009).
17   Power, R. M. & Huisken, J. A guide to light-sheet fluorescence microscopy for
multiscale imaging. *Nat Methods* **14**, 360-373, doi:10.1038/nmeth.4224 (2017).
18   Mano, T. *et al.* Whole-Brain Analysis of Cells and Circuits by Tissue Clearing and Light-
Sheet Microscopy. *J Neurosci* **38**, 9330-9337, doi:10.1523/jneurosci.1677-18.2018
(2018).
19   Tomer, R., Khairy, K., Amat, F. & Keller, P. J. Quantitative high-speed imaging of entire
developing embryos with simultaneous multiview light-sheet microscopy. *Nature*
Methods **9**, 755-763, doi:10.1038/nmeth.2062 (2012).
20   Chung, K. *et al.* Structural and molecular interrogation of intact biological systems.
Nature **497**, 332-337, doi:10.1038/nature12107 (2013).
21   Susaki, E. A. *et al.* Whole-brain imaging with single-cell resolution using chemical
cocktails and computational analysis. *Cell* **157**, 726-739, doi:10.1016/j.cell.2014.03.042
(2014).
22   Erturk, A. *et al.* Three-dimensional imaging of solvent-cleared organs using 3DISCO.
Nat Protoc **7**, 1983-1995, doi:10.1038/nprot.2012.119 (2012).
23   Stocum, D. L. Mechanisms of urodele limb regeneration. *Regeneration* **4**, 159-200,
doi:10.1002/reg2.92 (2017).
24   Singer, M. & Caston, J. D. Neurotrophic dependence of macromolecular synthesis in the
early limb regenerate of the newt, Triturus. *Journal of*
Embryology and Experimental Morphology **28**, 1 (1972).
25   Dresden, M. H. Denervation effects on newt limb regeneration: DNA, RNA, and protein
synthesis. *Dev Biol* **19**, 311-320, doi:10.1016/0012-1606(69)90044-x (1969).
26   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature*
Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).
27   MATLAB version 9.7.0.813654 (R2019b). Natick, Massachusetts: The MathWorks Inc.;
2019
28   Pietzsch, T., Saalfeld, S., Preibisch, S. & Tomancak, P. BigDataViewer: visualization and
processing for large image data sets. *Nature Methods* **12**, 481-483,
doi:10.1038/nmeth.3392 (2015).
29   Doube, M. *et al.* BoneJ: Free and extensible bone image analysis in ImageJ. *Bone* **47**,
1076-1079, doi:10.1016/j.bone.2010.08.023 (2010).
30   Arganda-Carreras, I. *et al.* Trainable Weka Segmentation: a machine learning tool for
microscopy pixel classification. *Bioinformatics* **33**, 2424-2426,
doi:10.1093/bioinformatics/btx180 (2017).
31   Fadero, T. C. & Maddox, P. S. Live imaging looks deeper. *Elife* **6**, e30515,
doi:10.7554/eLife.30515 (2017).
32   Singer, M. The influence of the nerve in regeneration of the amphibian extremity. *Q Rev*
Biol **27**, 169-200, doi:10.1086/398873 (1952).
33   Farkas, J. E. & Monaghan, J. R. A brief history of the study of nerve dependent
regeneration. *Neurogenesis (Austin)* **4**, e1302216, doi:10.1080/23262133.2017.1302216
(2017).

Table 1. Monomer analogs used to demonstrate whole mount visualization method

Name	Macromolecule analog	Biological process	Click-it modification	Reference
5-ethynyl-2′-deoxyuridine (EdU)	Thymidine	DNA synthesis	Alkyne	Salic & Mitchison 2008
5-Ethynyl Uridine (5-EU)	Uracil	Transcription	Alkyne	Jao & Salic 2008
L-Azidohomoalanine (AHA)	Methionine	Translation	Azide	Wang et al. 2017
Azide-modified glucosamine (GlcNAz)	Glucoseamine	Protein glycosylation	Azide	Laughlin et al. 2006

Figure 1. Outline of staining/analysis pipeline and exemplary images.

(A) Overview of entire sample preparation, imaging, and data analysis pipeline. (B-D) Once macromolecules are labeled *in vivo*, synthesis can be visualized throughout the injected animal. Here, we show DNA synthesis (EdU) in the torso (B) and both DNA synthesis and protein glycosylation (GlcNAz) in the hand (C) and finger (D). Images from panels B and C were uncleared and imaged at 5X magnification. Image from panel D was also uncleared and imaged at 10X magnification. Scale bars for panels B&C= 600µm for each axis. Scale bars for panel D= 200µm for each axis.

Figure 2. Dual staining of macromolecule synthesis in whole mount.

(A-D) Stitched and fused 3D reconstruction of 13 dpa blastemas stained for multiple macromolecules obtained by LSFM. (B-H) Single Z-plane from A-D that represents the entirety of the blastema. (I-L) Tissue section from identically treated limbs as A-H showing similar macromolecule staining patterns, indicating that the whole mount staining method does not alter macromolecule synthesis staining patterns. Scale bars for panels A-D= 600µm for each axis. Scale bars for panels E-L= 200µm.

Figure 3. Workflow for 3D, multiscale analysis of the regenerating axolotl humerus.

Multiscale analysis of a 35 dpa regenerating axolotl humerus, stained for AHA (red) and EdU

(green). The humerus in the image stack was (A) aligned along the proximodistal (P-D) axis and

(B) its morphology was segmented. The resulting mask was used to analyze the organ volume

and shape by (D) reslicing it along the P-D axis and studying the cross-sections obtained. (C)

The segmented morphology in B was used to mask the green channel for cellular- and

molecular-level analyses. (E) Cells in the humerus were segmented and their spatial distribution

was analyzed to obtain cellular number and density. (F) The masked image stack in C was

resliced along the P-D axis and the pixel maps of the cross-sections were used to characterize the

molecular intensity and distribution within the humerus. The vertical yellow line in A and C

indicates the plane of amputation.

Figure 4. 3D quantification across scales of a regenerating axolotl humerus.

(A) The cross-sections of the humerus in Figure 3D were analyzed with the Fiji plugin BoneJ to quantify humerus shape and volume. (A') The cross-sectional area along the proximodistal (P-D) axis provides a measure of volume distribution along the humerus. (A") The ratio of the maximum chord length from the minor axis ($2R_1$) with respect to the maximum chord length from the major axis ($2R_2$) provides a measure of cross-sectional circularity in the humerus. Values closer to 1.0 in the proximal side indicate a more circular cross-section in this zone. (B) The Fiji plugin Trainable Weka Segmentation 3D and 3D Objects Counter were used in the cellular analysis of proliferating chondrocytes illustrated in Figure 3E. (B') The number of EdU$^+$ cells within a 50µm slice along the P-D axis was divided by the slice volume to obtain a density-like measure. (B") The center of mass of each cell was plotted in 3D, with point size and color proportional to the segmented cell volume. (C) The molecular intensity and distribution were analyzed based on the resliced pixel intensity maps of the masked green channel in Figure 3F. (C') Mean intensity of each slice perpendicular to the P-D axis. (C") The histogram of the EdU staining in the humerus can provide a measure of DNA synthesis rate. The vertical yellow line in the top row images indicates the plane of amputation.

478 **Figure 5. 3D visualization of DNA synthesis in innervated/denervated regenerating limbs.**

479 (A) Schematic of experimental design used to obtain samples from B-O. (B-O) Time course of

480 regeneration in innervated and 24 hour denervated limbs at 0, 6, 9, 12, 15, 18, and 25 dpa. Scale

481 bars for panels B-O=600µm for each axis.

Figure 6. 3D quantification of DNA synthesis in innervated/denervated regenerating limbs.

(A) A cube with sides of 175µm was cropped along the proximodistal axis 250µm from the distal tip of each blastema. (B) The overall histogram of the cubes confirms that innervated blastemas have faster DNA synthesis rates than their denervated counterparts. The difference is quantified by computing the ratio between the mean pixel intensity of the innervated versus denervated blastema cube of a same animal (inset plot).

Supplementary figure 1. Single staining of macromolecule synthesis in whole mount.

(A-C) Stitched and fused 3D reconstruction of 13 dpa blastemas stained for one macromolecule

obtained by LSFM. (D-F) Single Z-plane from A-C that represents the entirety of the blastema.

(G-I) Tissue section from identically treated limbs as A-F. Scale bars for panels A-C= 600µm in

each axis. Scale bars for panels D-I= 200µm.

Supplementary figure 2. Specificity of GlcNAz staining.

(A) Tissue section of a regenerating axolotl limb where the click-it cocktail (green) for GlcNAz

was applied prior to staining with GlcNAc antibodies (Ab) (red). (B) Tissue section of a

regenerating axolotl limb where GlcNAc antibodies (green) were applied prior to treatment with

the click-it cocktail (red) for GlcNAz, demonstrating the specificity of the *in vivo* GlcNAz

labeling. Scale bars=100µm.

Supplementary figure 3. Comparison of imaging in PBS and 67%TDE

(A-B) Single Z-plane of 13 dpa blastema imaged in PBS (A) or cleared and imaged in 67% TDE (B). Red indicates EdU staining whereas blue represents DAPI staining. (C-D) Pixel intensity map of PBS imaged blastema (C) and 67% TDE imaged blastema (D). Scale bars are in units of microns.

Supplementary video 1

Rotating axolotl hand stained for EdU (green) and GlcNAz (red).

Supplementary video 2

Rotating axolotl limb stained for EdU (green), AHA (red), and DAPI (blue).

Supplementary video 3

Scroll through of Z-stack from supplementary video 2.