Oxytocin selectively excites interneurons and inhibits output neurons of the bed nucleus of the

stria terminalis (BNST)

Running title: Oxytocin regulates intrinsic inhibitory network of the BNST

Walter Francesconi [1,2], Fulvia Berton [1,2], Valentina Olivera-Pasilio [1,2,3], and Joanna Dabrowska
[1,2,3]

[1] Center for the Neurobiology of Stress Resilience and Psychiatric Disorders, Rosalind Franklin

University of Medicine and Science, North Chicago, IL 60064, USA

[2] Discipline of Cellular and Molecular Pharmacology, Chicago Medical School, Rosalind Franklin

University of Medicine and Science, North Chicago, IL 60064, USA

[3] School of Graduate and Postdoctoral Studies, Rosalind Franklin University of Medicine and

Science, North Chicago, IL 60064, USA

* To whom correspondence should be addressed:

Joanna Dabrowska, Center for the Neurobiology of Stress Resilience and Psychiatric Disorders,

Rosalind Franklin University of Medicine and Science, North Chicago, IL 60064, USA

joanna.dabrowska@rosalindfranklin.edu

Phone: (847) 578-8664

Fax: (847) 578-3268

Number of figures (7), number of tables (1)

Abstract

The dorsolateral bed nucleus of the stria terminalis ($BNST_{DL}$) has high expression of oxytocin receptors, but their role in the modulation of $BNST_{DL}$ activity remains elusive. $BNST_{DL}$ contains GABA-ergic neurons classified based on intrinsic membrane properties into three types. Using *in vitro* patch-clamp and cell-attached recordings in male rats, we demonstrate that oxytocin excites and increases spontaneous firing of Type I, putative $BNST_{DL}$ interneurons. As a consequence, oxytocin increases the frequency of spontaneous inhibitory post-synaptic currents (sIPSCs) (tetrodotoxin-sensitive) and reduces spontaneous firing of Type II neurons. In contrast, in Type III neurons, oxytocin reduces the amplitude of both sIPSCs and evoked IPSCs, suggesting a direct postsynaptic inhibitory effect. As Type II and Type III are the $BNST_{DL}$ projection neurons, we present a model of fine-tuned modulation by oxytocin, which selectively excites Type I $BNST_{DL}$ interneurons and inhibits Type II and Type III output neurons, via an indirect and direct mechanism, respectively.

1. Introduction

Oxytocin (OT) is a nine amino acid peptide serving as a hypothalamic hormone, which regulates uterine contractions during labor, milk ejection reflex during lactation (Caldeyro-Barcia and Poseiro, 1959; Nickerson et al., 1954), as well as water-electrolyte balance (Han et al., 1993; Verbalis et al., 1993). As a neuromodulator in the central nervous system, OT facilitates social behaviors (Bosch and Young, 2018; Pedersen et al., 1992; Tabbaa et al., 2016), regulates stress response, stress-coping behaviors, and fear and anxiety-like behaviors, for review see (Janeček and Dabrowska, 2019; Neumann and Slattery, 2016; Olivera-Pasilio and Dabrowska, 2020). Although OT has been shown to modulate fear responses primarily in the central amygdala (CeA) (Knobloch et al., 2012; Terburg et al., 2018; Viviani et al., 2011), we recently showed that oxytocin receptor (OTR) transmission in the dorsolateral bed nucleus of the stria terminalis (BNST$_{DL}$) facilitates the formation of cued fear and improves fear discrimination to signaled vs. un-signaled threats (Martinon et al., 2019; Moaddab and Dabrowska, 2017), for review see (Olivera-Pasilio and Dabrowska, 2020).

On a cellular level, OT has shown robust modulatory influence on the neurocircuitry of the hippocampus (Harden and Frazier, 2016; Maniezzi et al., 2019; Owen et al., 2013; Tirko et al., 2018), auditory cortex (Marlin et al., 2015), ventral tegmental area, VTA, (Hung et al., 2017; Xiao et al., 2017), and the CeA (Huber et al., 2005; Knobloch et al., 2012; Terburg et al., 2018; Viviani et al., 2011). Notably, although the BNST has one of the highest expression levels of OTR in the rodent brain (Dabrowska et al., 2011; Dumais et al., 2013; Tribollet et al., 1992; Veinante and Freund-Mercier, 1997) and receives OT inputs from hypothalamic nuclei (Dabrowska et al., 2011; Knobloch et al., 2012; Martinon et al., 2019), the effects of OT on excitability and synaptic transmission of BNST neurons remain unknown. Studies in the 90's from the Ingram group using electrophysiological approaches have shown that OT increases the spontaneous firing frequency in a subpopulation of BNST neurons in lactating female rats

(Ingram et al., 1990; Ingram and Moos, 1992; Wilson et al., 2005), but their phenotype is not
known.

$BNST_{DL}$ neurons are GABA-ergic (Cullinan et al., 1993; Dabrowska et al., 2013a; Sun
and Cassell, 1993), and functionally interconnected via a dense intrinsic inhibitory network (Xu
et al., 2016). As originally described by Rainnie and colleagues, $BNST_{DL}$ neurons can be
classified into three major neuron types (I-III) based on their electrophysiological phenotype
defined as a unique pattern of voltage deflection in response to transient depolarizing and
hyperpolarizing current injections (Daniel et al., 2017; Hammack et al., 2007) and/or based on a
single-cell mRNA expression of ion channel subunits (Hazra et al., 2011). Recently, it was
shown that both Type II and Type III are output neurons of the $BNST_{DL}$, with major projections to
the CeA (mainly Type II neurons), lateral hypothalamus (LH), and the VTA (mainly Type III
neurons) (Yamauchi et al., 2018).

In the current study, using *in vitro* slice patch-clamp recordings in a whole-cell and cell-
attached mode, we investigated the effects of OT on intrinsic excitability, inhibitory synaptic
transmission and spontaneous firing in Type I-III $BNST_{DL}$ neurons in male Sprague-Dawley rats.
We show that OT is a powerful modulator of the intrinsic inhibitory network such as it selectively
increases the intrinsic excitability and the frequency of spontaneous firing of regular spiking,
Type I neurons, which leads to increased inhibitory synaptic transmission (spontaneous
Inhibitory Postsynaptic Currents, sIPSCs) and reduced spontaneous firing rate of Type II
$BNST_{DL}$ neurons. We also combined injections of a fluorescent retrograde tracer into the CeA
with electrophysiological recordings from fluorescent $BNST_{DL}$ neurons and we confirm that all
recorded CeA-projecting (*BNST➔CeA*) neurons are Type II neurons. Finally, we demonstrate
that OT reduces the amplitude of both spontaneous and evoked IPSCs selectively in Type III
neurons via a direct, OTR-dependent, postsynaptic mechanism. These results show that OT
directly excites Type I $BNST_{DL}$ interneurons and inhibits two classes of the $BNST_{DL}$ output
neurons via an indirect (Type II neurons) and a direct (Type III neurons) mechanism.

2. Results

2.1. The effects of OT on intrinsic membrane properties and spontaneous firing in Type I-III BNST$_{DL}$ neurons.

Whole-cell patch-clamp recordings were performed from BNST$_{DL}$ neurons, primarily in the oval nucleus of the BNST (BNST$_{OV}$). At the beginning of the recording sessions in current clamp mode, neurons were electrophysiologically characterized into three types (Type I-III) using current pulses (450 msec in duration) from -350 pA to 80 pA in 5 or 10 pA increments. Based on their characteristic voltage responses (Hammack et al., 2007), three types of neurons were identified in the BNST$_{DL}$ (see Methods for details). We investigated the effects of OT (0.2 μM based on (Crane et al., 2020; Xiao et al., 2017) on intrinsic membrane properties: resting membrane potential (RMP), input resistance (Rin), rheobase (Rh), threshold of the first action potential (Th) (calculated as the voltage at which the depolarization rate exceeded 5 mV/ms), and latency of the first spike (Lat) evoked at the Rh current. Intrinsic membrane properties were investigated in a presence of synaptic transmission's blockers. In Type I neurons, OT induced a significant depolarization of RMP from -62.39 ± 0.84 mV to -57.94 ± 1.39 mV in comparison to baseline (pre, P = 0.0013) and washout (post, P = 0.0304, **Fig. 1A-B**). This was associated with a significant increase of Rin from 170.98 ± 8.6 MΩ to 233.64 ± 18.6 MΩ (P = 0.0085, **Fig. 1C**). As a consequence, OT induced a significant reduction of Rh from 21.96 ± 2.29 pA to 17.32 ± 2.26 pA (P = 0.0023, **Fig. 1D**). The Th of the first spike was also significantly reduced from -38.38 ± 0.68 mV to -39.75 ± 0.88 mV after OT (P = 0.0346, **Fig. 1E**). When the selective and potent OTR antagonist, d(CH2)5(1), D-Tyr(2), Thr(4), Orn(8), des-Gly-NH2(9)]-Vasotocin trifluoroacetate salt (Manning et al., 2012) (0.4 μM, OTA) was applied for 10 min before and during OT application, OT did not significantly affect RMP (P = 0.0710), Rh (P = 0.1231), Th (P = 0.3532) or Lat (P = 0.6252, **not shown**). In contrast to Type I neurons, OT did not affect the intrinsic membrane properties of Type II and III neurons (**Table 1**).

Figure 1. **Oxytocin affects intrinsic membrane properties of Type I BNST$_{DL}$ neurons. A**: A representative trace showing that oxytocin (OT, 0.2 µM) shortens the latency to the 1st spike at rheobase current (Rh) and depolarizes resting membrane potential (RMP, from -64 mV to -60 mV). This is associated with increased input resistance (Rin), measured with a depolarizing current step below the Rh. OT also reduces Rh from 30 pA (pre) to 20 pA (OT) **B-D**: Line graphs with individual neurons' responses (thin gray) and average responses for all neurons recorded (thick black) showing that OT significantly increases the RMP (**B**, $P = 0.0002$, F (1.593, 19.12) = 15.76, n = 14, mixed-effect analysis), increases Rin (**C**, $P = 0.0085$, n = 9, paired t-test), reduces the Rh (**D**, $P = 0.0457$, F (1.341, 16.09) = 4.226, n = 14, mixed-effect analysis), reduces the 1st spike threshold (**E**, P = 0.0164, F (1.472, 20.61) = 5.749, and reduced the 1st spike latency (**F**, $P = 0.0185$, F (1.971, 23.66) = 4.775, n = 14, mixed-effect analysis). Thick black lines show average responses for all Type I neurons recorded, **$P < 0.01$, *$P < 0.05$ in comparison to pre or post-OT.

Table 1. Intrinsic membrane properties and firing activity of Type I-III BNST$_{DL}$ neurons after OT (0.2 µM) application respective to baseline (pre-OT). ****P < 0.0001, **P < 0.01, *P < 0.05. In Type II neurons OT did not affect RMP (P = 0.0985), Rin (P = 0.1377), Rh (P = 0.0964), or Th (P = 0.5027). Similarly, in Type III neurons OT did not significantly change RMP (P = 0.1066), Rin (P = 0.0788), Rh (P = 0.1664), or Th (P = 0.5150).

Type I neurons	Pre	OT
RMP (mV)	-62.39 ± 0.84	**57.94 ± 1.39 ****
Rin (MOhms)	170.98 ± 8.6	**233.64 ± 18.6 ***
Rh (pA)	21.96 ± 2.29	**17.32 ± 2.26 ****
Th (mV)	-38.38 ± 0.68	**39.75 ± 0.88 ***
Lat (ms)	326.44 ± 28.63	**224.58 ± 32.51 ****
fAHP (mV)	8.63 ± 0.89	**7.47 ± 0.77 ****
mAHP (mV)	10.69 ± 0.61	**8.45 ± 0.72 *****
lAHP (mV)	0.49 ± 0.12	0.41 ± 0.1
Type II neurons		
RMP (mV)	-57.04 ± 0.86	-56.74 ± 0.78
Rin (MOhms)	221.8 ± 29.27	276.03 ± 46.66
Rh (pA)	16.92 ± 2.52	15 ± 2.66
Th (mV)	-39.87 ± 0.95	-40.32 ± 1.05
Type III neurons		
RMP (mV)	-65.73 ± 1.42	-63.3 ± 1.83
Rin (MOhms)	120.66 ± 23.12	156.93 ± 29.7
Rh (pA)	56.46 ± 8.92	48.33 ± 7.36
Th (mV)	-37.28 ± 0.85	-37.08 ± 0.93

We also investigated spiking ability by measuring the input-output (I-O) relationship

before and after OT application. The steady state frequency (SSF) was measured by applying

depolarizing pulses (1 sec in duration) of different amplitudes (0-80 pA in 10 pA increments).

The SSF was determined as the average of the inverse of the inter-spike intervals (ISI) from all

action potentials starting from the second action potential. The I/O curve was built before (pre),

during, and after OT application (post). OT caused a leftward shift of input/output (I-O)

relationship in Type I neurons, demonstrating a significant increment effect on SSF at 40 pA, 50

pA, and 60 pA. Post-hoc test showed a significant increase of the SSF after OT from 6.81 ±

0.68 Hz to 8.96 ± 0.63 Hz in comparison to baseline (pre, P = 0.0032) and washout (post, P =

0.0095) at 40 pA; from 8.73 ± 0.73 Hz to 11.20 ± 0.70 Hz (P = 0.0163, pre and P = 0.0226, post)

at 50 pA; and from 9.22 ± 0.57 Hz to 11.89 ± 0.99 Hz (P = 0.0271, pre and P = 0.0054, post) at

60 pA. The OT effect reached a trend at 30 pA (from 5.64 ± 0.5 Hz to 7.31 ± 0.53 Hz, P =

0.0820) and at 70 pA (from 10.79 ± 0.6 Hz to 13.62 ± 0.99 Hz, P = 0.0794). The effect was not

significant at 20 pA (P = 0.1268) and 80 pA (P = 0.1320, **Fig. 2A-B**). Importantly, in the

presence of OTA, OT did not affect the SSF of Type I neurons at 80 pA (P = 0.7960), 70 pA (P

= 0.4237), 60 pA (P = 0.6312), 50 pA (P = 0.4299), 40 pA (P = 0.3365), or 30 pA (P = 0.4817,

Fig. 2C). OT did not affect SSF in Type II and Type III neurons (**Fig. 2D-E**).

This increase in the spiking ability of Type I neurons after OT was associated with a

significant shortening of the first spike's latency from 326.44 ± 28.63 msec to 224.58 ± 32.51

msec (P = 0.0083, **Fig. 1F**).

A variety of afterhyperpolarizations (AHPs) following a single or a train of action

potentials have been described to regulate the neuron excitability and spike frequency

adaptation (Gu et al., 2008; Lopez de Armentia and Sah, 2004). They come in a temporal

sequence of three AHPs: the fast AHP (fAHP) lasting 1-5 msec, the medium AHP (mAHP)

lasting 50-200 msec, which is responsible for early spike frequency adaptation during a train of

action potentials, and the late AHP (lAHP) lasting several seconds after a train of action

potentials (Gu et al., 2005a). We investigated the effect of the OT on the amplitude of fAHP,

Figure 2

Figure 2. **Oxytocin increases intrinsic excitability of Type I, but not Type II or Type III, BNST$_{DL}$ neurons**. **A**: A representative trace showing voltage deflections in response to hyperpolarizing and depolarizing current injections characteristic of Type I neuron (left) and the increase in firing frequency of Type I neuron in the presence of OT respective to baseline (pre) (right). **B**: OT causes a leftward shift of input/output (I-O) relationship in Type I neurons. The steady state frequency (SSF) was determined as an average of an inverse of the inter-spike intervals (ISI) from all action potentials starting from the second action potential. In Type I neurons, OT shows a significant increment effect on SSF at 40 pA (P = 0.0024, F (1.667, 15.01) = 10.11), at 50 pA (P = 0.0271, F (1.863, 16.76) = 4.628), and at 60 pA (P = 0.0271, F (1.688, 11.82) = 5.267, mixed-effects analysis). **C**: Application of OTR antagonist, OTA (0.4 µM), prevents the leftward shift of the I-O relationship induced by OT application (40 pA, P = 0.3365, 50 pA, P = 0.4299, 60 pA, P = 0.6312, n = 5, RM ANOVA). **D-E**: OT application does not modify the I-O relationship in Type II (**D**) or Type III BNST$_{DL}$ neurons (**E**). In Type II neurons, OT did not affect SSF at 80 pA (P = 0.3116), 70 pA (P = 0.8236), 60 pA (P = 0.9519), 50 pA (P = 0.8993), 40 pA (P = 0.7636), 30 pA (P = 0.8673), or at 20 pA (P = 0.6990, **D**). Similarly, in Type III neurons, there was no effect of OT on SSF at 80 pA (P = 0.1151), 70 pA (P = 0.1138), 60 pA (P = 0.180), 50 pA (P = 0.0847), 40 pA (P = 0.1041), 30 pA (P = 0.0790), or at 20 pA (P = 0.0681, **E**). Each data point represents the mean ± SEM, **P < 0.01, *P < 0.05 in comparison to pre-OT, ## P < 0.01, # P < 0.05 in comparison to post-OT.

mAHP and lAHP in all types of the $BNST_{DL}$ neurons and show that OT significantly reduced the amplitude of fAHP (from 8.63 ± 0.89 mV to 7.47 ± 0.77 mV, P = 0.0062, **Fig. 3A-B**) and mAHP (from 10.69 ± 0.61 mV to 8.45 ± 0.72 mV, P = 0.0001 vs. pre, and P = 0.0003 vs. post), **Fig. 3A, D**). Notably, OT did not affect the amplitudes of fAHP (P = 0.2220, **Fig. 3C**) and mAHP (P = 0.3968, **Fig. 3E**) in the presence of OTA. Lastly, OT did not significantly affect the amplitude of lAHP (P = 0.2401, **not shown**).

We also investigated if these changes in AHP amplitude could modify the spike frequency adaptation. We measured the instantaneous frequency for every action potential interval. The ISI number is the inter-stimulus interval between two consecutive action potentials. We found that OT significantly increased the instantaneous frequency of ISIs and post-hoc test showed a significant OT effect on frequency (in Hz) of ISI number 2 (ISI-2) from 16.77 ± 1.55 to 19.57 ± 1.2 (P = 0.0006), ISI-3 from 12.48 ± 0.88 to 16.19 ± 1.09 (P < 0.0001), ISI-4 from 11.92 ± 0.88 to 14.87 ± 0.76 (P = 0.0006), ISI-5 from 11.36 ± 0.68 to 14.58 ± 0.77 (P = 0.0001), ISI-6 from 11.26 ± 0.72 to 13.93 ± 0.76 (P = 0.0079), and ISI-7 from 10.58 ± 0.64 o 13.44 ± 0.88 (P = 0.0051, **Fig. 3F**).

The spontaneous firing activity of Type I and II $BNST_{DL}$ neurons was recorded in a cell-attached voltage-clamp mode without synaptic transmission's blockers. During cell-attached recordings, the spontaneous firing was present in 7 out of 9 Type I neurons and OT significantly increased the firing rate in the spontaneously firing neurons from 0.88 ± 0.66 Hz to 2.19 ± 0.93 Hz (P = 0.0242, paired t-test, **Fig. 4A-B**). In addition, although only 6 out of 18 Type II neurons showed spontaneous firing in a cell-attached mode, in all spontaneously firing Type II neurons, OT reduced the firing rate from 1.85 ± 0.74 Hz to 1.15 ± 0.69 Hz, which nearly reached statistical significance (P = 0.0525, paired t-test, **Fig. 4C-D**).

Overall, these results suggest that OT increases the intrinsic excitability and facilitates the spontaneous firing selectively in Type I $BNST_{DL}$ neurons, whereas it reduces spontaneous firing rate in Type II $BNST_{DL}$ neurons.

Figure 3. Oxytocin reduces fast and medium afterhyperpolarizations and modifies early and late spike frequency adaptation in Type I BNST$_{DL}$ neurons. **A**: A representative trace from Type I neuron showing a reduction of fast AHP (fAHP) and medium AHP (mAHP) after OT application respective to baseline (pre). The amplitudes of the fAHP and mAHP following the first action potential evoked at Rh were calculated as the difference from the action potential threshold to the membrane voltage measured after 2-5 msec and 50-100 msec, respectively. The lAHP was measured at 2000 msec after the end of the depolarizing pulse, following generation of five action potentials. **B**: Line graphs showing individual neurons' responses (gray) and reduction of the fAHP during OT application (P = 0.0124, F (1.473, 17.68 = 6.493, mixed-effects analysis, n = 14). **C**: OTR antagonist, OTA (0.4 µM), prevents the OT-induced reduction of fAHP (P = 0.2220, n = 7, RM one-way ANOVA). **D**: Line graph showing reduction of mAHP during OT application (P <0.0001, F (1.844, 22.13) = 23.25, mixed effect analysis, n = 14). **E**: OT-induced reduction of mAHP is blocked by OTR antagonist, OTA (P = 0.3968, n = 7, RM one-way ANOVA). Thick black lines show average responses for all Type I neurons recorded. **F**: OT modifies early and late spike frequency adaptation. The early and late spike frequency adaptation was investigated using a depolarizing pulse of 1 sec and measuring the instantaneous frequency for every action potential interval. The ISI number is the inter-stimulus interval between two consecutive action potentials. OT significantly increased the instantaneous frequency of ISI (P < 0.0001, F (1, 13) = 82.69, mixed effects model). Each data point represents the mean ± SEM, ****P < 0.0001, ***P < 0.001, **P < 0.01, *P < 0.05.

Figure 4. Oxytocin increases spontaneous firing rate of Type I neurons and reduces spontaneous firing rate of Type II BNST$_{DL}$ neurons in cell-attached recordings. **A**: A representative trace of Type I neuron showing an increase in spontaneous firing rate after OT application. **B**: Line graph showing that OT increased spontaneous firing rate in all spontaneously firing Type I neurons (*P* = 0.0242, paired *t*-test, n = 7), gray lines represent individual neurons' responses, thick black line represents average response. **C**: A representative trace of Type II neuron showing a reduction in spontaneous firing rate after OT application. **D**: Line graph showing that OT reduced spontaneous firing rate in spontaneously firing Type II neurons in cell-attached mode (*P* = 0.0525, paired *t*-test, n = 6).

2.2. The effects of OT on inhibitory synaptic transmission in Type I-III BNST$_{DL}$ neurons.

To study the effect of OT on spontaneous inhibitory postsynaptic currents (sIPSCs), the

events were recorded from Type I-III neurons in a voltage clamp mode at holding potential of -

70 mV, in a presence of glutamatergic transmission's blockers. In Type II BNST$_{DL}$ neurons,

application of OT caused a significant increase in the frequency of sIPSCs in 12 out of 14

recorded neurons from 1.03 ± 0.20 Hz to 2.64 ± 0.47 Hz vs. baseline (pre, P = 0.0033), and vs.

washout (post, P = 0.0018, **Fig. 5A-B**). Notably, the effect of OT on the sIPSCs frequency was

not associated with a significant change in the amplitude (P = 0.9205, **Fig. 5C**) or in the total

charge (P = 0.5744, **not shown**). OT did not increase the sIPSCs frequency in the presence of

OTA (P = 0.3053, **Fig. 5D**). In addition, when the spike-driven sIPSCs were blocked (P =

0.0387) with Tetrodotoxin (TTX, 1 μM), OT no longer affected the frequency of miniature IPSCs

(mIPSCs) of Type II neurons (P = 0.9975, **Fig. 5E**). Finally, after electrical stimulation of the

Stria Terminalis, OT did not affect the evoked IPSCs (eIPSCs) amplitude (at the Vh of +10 mV)

in Type II neurons (from 266.76 ± 55.43 pA (pre) to 316.17 ± 82.49 pA, post, P = 0.2459, paired

t-test, **Fig. 6A-B**). These results demonstrate that OT indirectly increases the frequency of

spike-driven sIPSCs in Type II neurons.

In Type I neurons, OT did not affect the sIPSCs frequency (P = 0.3018), amplitude (P =

0.3347), or total charge (P = 0.6087, **not shown**). Moreover, OT did not change the amplitude

of the eIPSC in Type I neurons (P = 0.2449, **not shown**). In Type III neurons, OT did not affect

the frequency (P = 0.9440, **Fig. 5F**), but it reduced amplitude (P = 0.0314, **Fig. 5G**) and total

charge of sIPSCs (P = 0.0372, **not shown**). In Type III neurons, OT significantly reduced the

amplitude of the eIPSCs from 286.76 ± 97.41 (pre) to 210.69 ± 78.57 pA (P = 0.0105, paired t-

test, **Fig. 6C-D**). This reduction of the eIPSC amplitude was not associated with a significant

change in the paired pulse ratio (PPR) (P = 0.5672, paired t-test, **Fig. 6E**), suggesting that in

Type III neurons OT affects the synaptic response through a postsynaptic mechanism. Notably,

Figure 5. Oxytocin modulates inhibitory synaptic transmission in BNST_DL neurons. **A**: Representative recordings showing an increase in the frequency of spontaneous inhibitory postsynaptic currents (sIPSCs) after OT application respective to baseline (pre). **B-C**: Line graphs (gray) showing that 12 out of 14 recorded Type II neurons showed an increase in sIPSCs frequency (**B**, $P = 0.0022$, $F (1.058, 12.17) = 14.41$, n = 14, mixed-effect analysis) but not amplitude (**C**, $P = 0.5855$, $F (1.563, 17.19) = 0.4718$, n = 14, mixed-effect analysis) after OT application. **D**: OT does not affect sIPSC frequency in a presence of OTR antagonist, OTA (0.4 µM) ($P = 0.1788$, $F (1.469, 8.814) = 2.135$, n = 6, RM ANOVA). **E**: When spike-driven IPSCs were blocked with TTX ($P = 0.0151$, $F (1.030, 6.179) = 10.98$, n = 6, RM ANOVA), OT does not affect the sIPSCs frequency. **F-G**: OT does not affect frequency (**F**, $P = 0.9362$) but reduces amplitude of sIPCS in Type III neurons (**G**, $P = 0.0160$, paired *t*-test). Thick black lines show averaged responses for all neurons recorded. Each data point represents the mean ± SEM, **$P < 0.01$, *$P < 0.05$ in comparison to pre- or post-OT.

Figure 6. Oxytocin increases amplitude of evoked IPSCs in Type III, but not Type II BNST$_{DL}$ neurons. **A**: Representative traces of eIPSC in response to the electrical stimulation of the stria terminalis in Type II neurons. **B**: Line graph showing that OT does not significantly change the mean amplitude of eIPSC in Type II neurons (P = 0.2459, paired *t*-test). Thick black lines show averaged responses for all neurons recorded. **C**: Representative traces of eIPSCs in Type III neurons showing reduction of the eIPSCs amplitude after OT application. **D-E**: Line graphs showing individual Type III neurons and the reduction of the eIPSC amplitude after OT (**D**, P = 0.0105) without any significant change in the PPR (**E**, P = 0.5672).

Figure 7. Type II neurons are output neurons of the BNST$_{DL}$ to the central amygdala. A-B: Microphotographs showing neurons expressing green fluorescent microbeads in the BNST$_{DL}$ after the retrobeads were injected into the central amygdala (CeA). **A**: BNST neurons projecting to the CeA (BNST⇒CeA neurons) were primarily found in the dorsolateral BNST, primarily the oval nucleus, BNST$_{OV}$, and to a lesser extent in the anteromedial BNST, BNST$_{AM}$ (20x, LV – Lateral ventricle, IC – Internal capsulae, scale bar 100 μm). **B**: High magnification (60x) microphotographs showing green fluorescent beads present in membranes and processes of the BNST⇒CeA neurons (scale bar 10 μm). **C-D**: Electrophysiological characterization of the BNST⇒CeA neurons using visually guided patching (**C**, 40x, scale bar 20 μm). Representative recording of the BNST⇒CeA fluorescent neuron showing the electrophysiological properties of a Type II neuron such as the presence of a sag during the hyperpolarized current pulses and the spike rebound at the end of the 450 ms current pulse injection (**D**). **E**: Diagram showing that OT regulates the intrinsic inhibitory network and inhibits output neurons of the BNST$_{DL}$. By directly increasing frequency of spontaneous firing of Type I regular spiking neurons, OT inhibits firing of Type II BNST$_{DL}$ neurons. Type II neurons are output neurons of the BNST$_{DL}$ sending projections to the CeA (BNST⇒CeA neurons). OT also inhibits another class of output neurons, Type III, via a direct postsynaptic mechanism. Therefore, OT excites BNST$_{DL}$ interneurons (Type I) and inhibits two distinct classes of the BNST$_{DL}$ projection neurons via indirect (Type II) and direct (Type III) mechanisms, respectively.

in a presence of OTR antagonist, OT did not reduce the amplitude of eIPSCs in Type III

neurons (P = 0.8286, **not shown**).

2.3. CeA-projecting BNST$_{DL}$ neurons

To determine an electrophysiological phenotype of central amygdala-projecting BNST$_{DL}$

(*BNST➔CeA)* neurons, rats were first bilaterally injected into the CeA with a retrograde tracer,

green microbeads. Retro-labeled fluorescent neurons were found in subdivisions of the BNST$_{DL}$,

primarily in the oval nucleus of the BNST$_{DL}$ (BNST$_{OV}$) and to a lesser extent in anteromedial

BNST (BNST$_{AM}$, **Fig. 7A-B**). We found healthy fluorescent neurons that were suitable for

recordings in 6 out of 17 rats. In the slices from those 6 rats, we recorded 8 fluorescent neurons

in the BNST$_{OV}$ (**Fig. 7C**), and all recorded neurons were classified as Type II neurons (**Fig. 7D**).

3. Discussion

Here we demonstrate that OT regulates the intrinsic inhibitory network of the BNST$_{DL}$, where it

increases intrinsic excitability and spontaneous firing of Type I, regular spiking interneurons,

which leads to a potentiation of GABA-ergic inhibition and a reduction of spontaneous firing of

Type II, burst firing neurons. Using visually guided patching of retro-labeled BNST$_{DL}$ neurons,

we confirm that Type II are output neurons of the BNST$_{DL}$, which project to the CeA

(*BNST➔CeA* neurons). Furthermore, we show that OT inhibits another class of BNST$_{DL}$

projection neurons, Type III, via a direct postsynaptic mechanism. Therefore, we demonstrate

that there are three groups of OT-responsive neurons in the BNST$_{DL}$; a population of

interneurons (Type I) that is directly excited by OTR activation, and two distinct groups of

projection neurons, which are indirectly (Type II) and directly (Type III) inhibited by OT (**Fig. 7E).**

The OT-mediated inhibition of the BNST$_{DL}$ output has powerful implications for the modulation of

fear and anxiety-like behaviors.

Neurons of the BNST$_{DL}$ are GABA-ergic (Cullinan et al., 1993; Dabrowska et al., 2013a;

Sun and Cassell, 1993), and can be categorized into three major types (I-III) (Daniel et al.,

2017; Hammack et al., 2007). Type I neurons, characterized by regular firing and lack of burst-

firing activity (Hammack et al., 2007; Hazra et al., 2011), were classified as interneurons by

Paré and colleagues (Rodríguez-Sierra et al., 2013). This has been recently supported by

retrograde labeling showing a very small fraction of Type I neurons (in comparison to Type II

and Type III) projecting outside of the $BNST_{DL}$ (Yamauchi et al., 2018). We show that Type I

neurons are the primary OT-responsive neurons of the $BNST_{DL}$ as demonstrated by their

increased excitability and spontaneous firing rate after OT application. Previous studies using

single unit recordings showed that OT increased the spontaneous firing rate of 50% of BNST

neurons in lactating female rats (Ingram et al., 1990), and the magnitude of the response was

significantly potentiated during lactation (Ingram and Moos, 1992). Notably, in our study, the

excitatory effects of OT in Type I neurons are paralleled by increased inhibitory transmission

observed in Type II $BNST_{DL}$ neurons. The specific effect of OT on the sIPSCs frequency, but not

amplitude, suggests a presynaptic mechanism in Type II neurons. In addition, since the effect

was no longer observed when the spike-driven IPSCs were suppressed with TTX, this indicates

an indirect influence driven by an increased spontaneous firing of $BNST_{DL}$ interneurons. This

has been further supported by the lack of the OT effect on eIPSCs in Type II neurons,

suggesting that the inhibitory effect of OT in Type II neurons is not a result of the stimulation of

the extrinsic inhibitory projections.

In contrast, in Type III neurons, OT reduced the amplitude and the total charge, but not

the frequency of sIPSCs, as well as reduced the amplitude of eIPSC without changing the PPR,

overall indicating a postsynaptic effect. OT was shown previously to depress GABA-ergic

synapses in the supraoptic nucleus of the hypothalamus via postsynaptic increase of Ca^{2+} due

to the activation of G-protein coupled receptors (Alger and Nicoll, 1980; Brussaard et al., 1996).

However, since in our hands the effect on the amplitude was accompanied by a reduction in the

total charge, this could potentially indicate shortening of GABA channel opening (Martina et al.,

1994) via OTR-mediated increases in intracellular Ca^{2+}, as demonstrated after OTR activation

before (Brussaard et al., 1996). Therefore, in Type III neurons OT might reduce the amplitude of

sIPSCs and eIPSC through a postsynaptic mechanism involving signal transduction-mediated

regulation of intracellular Ca^{2+} levels. In addition, direct inhibitory effect of OT mediated via OTR

linked to inhibitory Gi-subunit was demonstrated in sensory neurons of spinal cord before

(Eliava et al., 2016).

To further investigate the OT effects on intrinsic excitability and firing, we demonstrated

that OT reduced fAHP and mAHP selectively in Type I neurons. AHPs generated after a single

action potential or a train of action potentials limit the frequency of neuronal firing (Alger and

Nicoll, 1980), as demonstrated by a significant increase in action potential firing after

pharmacological inhibition of AHPs (Pedarzani and Storm, 1993). The fAHP, mediated by BK

channels, is responsible for early spike frequency adaptation in many neuronal types (Gu et al.,

2007). The mAHP is mediated by the non-inactivating M-current, which blockade with selective

antagonist reduces late spike adaptation (Gu et al., 2008, 2005b). An OTR-mediated increase of

neuronal excitability was recently shown in the hippocampus, where a selective OTR agonist

induced membrane depolarization and increased membrane resistance in CA2 pyramidal cells

leading to a leftward shift in their firing frequency-current relationship (Tirko et al., 2018). Here

we show that OT-mediated excitation of Type I neurons in the BNST$_{DL}$ was associated with a

significant increase in instantaneous frequency of action potentials due to the reduction in the

early and late adaptation. Therefore, by acting on potassium conductances OT might increase

the intrinsic excitability and the spontaneous firing of Type I neurons.

According to our knowledge, this is the first study utilizing cell-attached mode in a cell-

type specific manner to show that in the BNST$_{DL}$, the majority of Type I neurons fire

spontaneously, and this effect is further potentiated by OT. Notably, although we showed that

the majority of Type II neurons remain silent even during cell-attached configuration, in all

spontaneously firing Type II neurons, OT application reduced the rate of firing. Notably, the

indirect inhibitory effect on spontaneous firing of Type II neurons was further confirmed by the

lack of the OT effect on current-evoked firing in these neurons, which was investigated in a

presence of synaptic transmission blockers. The latter further suggests that OT does not alter

intrinsic properties of Type II neurons and instead, reduces the spontaneous firing rate and so

their inhibitory synaptic transmission via the direct excitatory effect on Type I neurons.

The BNST was not entirely isolated during recordings, such as the slice also potentially

contained GABA-ergic neurons in the medial preoptic area (Okamura et al., 1990), which

express OTR (Sharma et al., 2019), and project to the BNST (Simerly and Swanson, 1988).

However, this projection primarily targets the principal encapsulated nucleus in the posterior

BNST (Maejima et al., 2015), where OTR are involved in reproductive behavior (Kremarik et

al., 1991). Notably, the posterior BNST was not a part of our slice preparation. In addition,

although we cannot entirely rule out a possibility that some inhibitory inputs to the $BNST_{DL}$ might

be originating from anteromedial ($BNST_{AM}$) or ventral BNST ($BNST_{VEN}$), which were included in

our slice, the intrinsic inhibitory projections in the BNST have been primarily shown to originate

in the $BNST_{DL}$, mainly the $BNST_{OV}$, and target $BNST_{AM}$ and $BNST_{VEN}$, whereas opposite

projections are rare (Gungor and Paré, 2016). Therefore, our results strongly suggest that Type

I neurons are $BNST_{DL}$ interneurons providing inhibitory synaptic input to Type II neurons, an

effect potentiated by OT.

It is important to note that the BNST also contains neurons producing arginine-

vasopressin (AVP) (al-Shamma and De Vries, 1996) and has a high expression of AVP

receptors, V1AR (Grundwald et al., 2016). As OT shows a relatively high affinity toward AV1R,

this raises a possibility of non-specific effects via V1AR. However, all the significant OT effects

in our study were blocked by OTA, which is highly selective toward OTR with a negligible affinity

toward V1AR, see (Manning et al., 2012).

Similar modulation of the intrinsic inhibitory network by OT was shown in a variety of

limbic brain regions, including the CeA. The CeA and the BNST share many similarities,

including predominantly GABA-ergic nature, analogous peptidergic cell types, as well as similar

intrinsic and extrinsic connectivity, and together they form an extended amygdala complex

(Alheid, 2003; Sun and Cassell, 1993). Interestingly, in the CeA, there are two groups of OT-

responsive neurons. A group of interneurons located in the lateral nucleus of the CeA (CeL)

(Gozzi et al., 2010) is excited by OTR activation (Huber et al., 2005; Terburg et al., 2018; Viviani

et al., 2011). Once excited by OT, these interneurons provide inhibitory input to medial nucleus

of the CeA (CeM) output neurons, which were shown to selectively project to lateral

periaqueductal gray and reduce contextual fear (Viviani et al., 2011), for review see (Beyeler

and Dabrowska, 2020). Similarly to the effects we observed in the BNST$_{DL}$, OTR-mediated

indirect inhibition of the CeM output neurons was demonstrated as increased frequency of

sIPSC during whole-cell recordings and reduced spontaneous firing rate in a cell-attached

configuration. Similarly, in the hippocampus (Harden and Frazier, 2016; Maniezzi et al., 2019;

Owen et al., 2013; Tirko et al., 2018; Zaninetti and Raggenbass, 2000), lateral amygdala (Crane

et al., 2020), and the prefrontal cortex (Marlin et al., 2015; Nakajima et al., 2014), OT was

shown to excite a population of interneurons, including fast-spiking parvalbumin interneurons,

which were shown to inhibit glutamatergic projection neurons.

We show that in the BNST$_{DL}$, OT inhibits two classes of projection neurons via indirect

(Type II) and direct (Type III) mechanisms. Recently, both Type II and Type III were

demonstrated output neurons of the BNST$_{DL}$, with major projections to the CeA (62% of Type II

and 33% of Type III), LH (83% of Type III), and the VTA (17% of Type II, 79% of Type III

neurons). Although at this time we cannot unequivocally state that OT specifically inhibits the

BNST➔CeA neurons, OT increased the sIPSCs frequency and reduced the spontaneous firing

in 86% and 100% of recorded Type II neurons, respectively. In addition, all the retrogradely

labeled *BNST➔CeA* neurons we recorded from the BNST$_{OV}$ were identified as Type II.

Therefore, there is a high probability that OT inhibits the *BNST➔CeA* neurons. It would still

need to be determined whether OT inhibits specific Type III projection neurons. Although

neuropeptidergic phenotypes corresponding to Type I-III neurons remain elusive, Type III are

putative corticotropin-releasing factor (CRF) neurons of the BNST$_{OV}$ (Dabrowska et al., 2013b)

and they were shown projecting to the LH, VTA, and the CeA, among many other brain regions

(Dabrowska et al., 2016).

Notably, Type II neurons are the most abundant and heterogeneous neuronal population

in rat $BNST_{DL}$ (Hazra et al., 2011; Yamauchi et al., 2018). Although the projection from the

$BNST_{DL}$ to the CeA has been shown before (McDonald et al., 1999), a functional inhibitory

projection was recently demonstrated using optogenetics (Gungor et al., 2015; Yamauchi et al.,

2018). The BNST is known to mediate stress-induced anxiety-like responses (Dabrowska et al.,

2013b; Daniel and Rainnie, 2016; Davis et al., 2010; Sparta et al., 2013), contextual fear (Davis

et al., 2010; Sullivan et al., 2004; Waddell et al., 2006), as well as fear responses to un-

signaled, diffuse, or unpredictable threats (Duvarci et al., 2009; Goode et al., 2020, 2019). The

recent optogenetic study showed that activation of the *BNST➔CeA* projection increases

anxiety-like behavior (Yamauchi et al., 2018), suggesting that the *BNST➔CeA* neurons are

important output neurons recruited during heightened states of anxiety. In addition, the BNST

has been shown to inhibit cued fear responses to discrete threats (Meloni et al., 2006). Notably,

our recent behavioral studies have demonstrated that OTR transmission in the $BNST_{DL}$

facilitates acquisition of cued fear and reduces fear responses to un-signaled/diffuse threats

(Janeček and Dabrowska, 2019; Martinon et al., 2019; Moaddab and Dabrowska, 2017),

showing opposite effects to the effects of the BNST activation. Our current electrophysiology

results suggest that OTR activation in the $BNST_{DL}$ facilitates cued fear and reduces anxiety by

inhibiting the *BNST➔CeA* output neurons.

4. Material and methods

4.1. Animals

Male Sprague-Dawley rats (Envigo, Chicago, IL; 240-300 g) were housed in groups of three on

a 12 h light/dark cycle (light 7 a.m. to 7 p.m.) with free access to water and food. Rats were

habituated to this environment for one week before the experiments began. Experiments were

performed in accordance with the guidelines of the NIH and approved by the Animal Care and

Use Committee at RFUMS. A total of 144 rats were used for the experiments. The effects of OT

on membrane properties were tested in 14 Type I neurons (from 10 rats), 13 Type II (12 rats),

and 9 Type III neurons (8 rats). The effect of OT on evoked firing was tested on 12 Type I (11

rats), 8 Type II (7 rats), and 9 Type III neurons (8 rats). The effect of OT on the spontaneous

firing was tested in 9 Type I (8 rats) and 18 Type II (15 rats). To study the effect of OT on

sIPSCs, we recorded 14 Type II (12 rats), 6 Type I (5 rats), and 8 Type III neurons (8 rats). We

recorded 6 Type II neurons (4 rats) in experiment with OTR antagonist, and 6 Type II neurons (5

rats) in TTX experiment. The effect of OT on the evoked IPSCs was tested in 5 Type I (5 rats), 8

Type II (8 rats), and 6 Type III (5 rats) neurons. We recorded 5 Type III (4 rats) neurons in

experiments with OTR antagonist. We recorded 8 retrogradely labeled Type II neurons (17 rats).

4.2. Slice preparation for electrophysiology

Rats were anaesthetized with Isoflurane USP (Patterson Veterinary, Greeley, CO, USA) before

decapitation. Brains were quickly removed and coronal 300 μm slices containing the $BNST_{DL}$

were prepared in ice-cold cutting solution (saturated with 95% O_2/ 5% CO_2) containing in mM:

NaCl 130, KCl 3.5, $NaHCO_3$ 30, KH_2PO_4 1.1, $CaCl_2 \cdot 2H_2O$ 1, D-glucose 10, $MgCl_2$ $6H_2O$ 6,

Kynurenate 2, pH 7.4, 290-300 mOsm. The slices were prepared using a Leica vibratome

(VT1200; Leica, Wetzlar, Germany), then incubated for 1 hr at 34°C in the cutting solution and

subsequently transferred to room temperature in artificial cerebrospinal fluid (aCSF) containing

in mM: NaCl 130, KCl 3.5, $NaHCO_3$ 30, KH_2PO_4 1.1, $CaCl_2 \cdot 2H_2O$ 2.5, D-glucose 10, and

$MgCl_2 \cdot 6H_2O$ 1.3, Ascorbate 0.4, Thiourea 0.8, Na Pyruvate 2, pH 7.4, with osmolarity of 290-300

mOsm, continuously oxygenated with 95% O_2/ 5% CO_2. Next, the slices were transferred to a

submerged recording chamber and perfused with aCSF at 33°C ± 1°C saturated with 95% O_2/

5% CO_2 at a flow rate of 2-3 ml/min. $BNST_{DL}$ neurons were visualized using infrared differential

interference contrast (IR-DIC) optic with an Upright microscope (Scientifica SliceScope Pro

1000, Clarksburg, NJ, USA).

4.3. Patch-clamp recordings

Whole-cell patch-clamp recordings were performed from $BNST_{DL}$ neurons, primarily in the oval

nucleus of the BNST ($BNST_{OV}$), using glass pipette (4-8 MΩ) pulled from thick-walled

borosilicate glass capillaries with a micropipette puller (Model P-97; Sutter Instrument, Novato,

CA, USA) as before (Dabrowska et al., 2013b; Francesconi et al., 2009). To study the effects of

OT (0.2 μM concentration based on (Crane et al., 2020; Xiao et al., 2017), oxytocin acetate salt,

H-2510, Bachem, Switzerland) on intrinsic membrane properties and firing activity the patch

pipettes were filled with potassium gluconate internal solution containing in mM: K-gluconate

135, KCl 2, $MgCl_2·6H_2O$ 3, HEPES 10, Na-phosphocreatine 5, ATP-K 2, GTP-Na 0.2, pH 7.3

adjusted with KOH, 300-305 mOsm. Intrinsic membrane properties experiments were performed

in a presence of synaptic transmission blockers: the AMPA receptor antagonist CNQX (10 μM),

the NMDA receptor antagonist AP-5 (50 μM), and GABA-A receptor antagonist, picrotoxin (25

μM). To study the effects of OT on the spontaneous inhibitory postsynaptic currents (sIPSCs),

the patch pipettes were filled with high chloride internal solution in mM: KOH 110, $MgCl_2·6H_2O$

2, KCl 50, EGTA 0.2, ATP-Na 2, GTP-Na 0.3, HEPES 10, and Spermine 0.1, the pH was

adjusted to 7.3 with D-gluconic acid and osmolarity was 300-305 mOsm. The evoked IPSCs

(eIPCSs) were recorded with pipettes filled with K-gluconate solution as described above. In

order to isolate GABAergic eIPSCs, the eIPSCs were recorded at the holding voltage (Vh) of

+10 mV, which is the reversal potential for the excitatory postsynaptic current. A stimulating

pipette filled with aCSF was placed in the Stria Terminalis. For each neuron recorded the

stimulus intensity was adjusted to give an eIPSCs of about 1/3 of the maximal amplitude and

maintained throughout the experiment. We applied 10 paired pulse stimuli (ISI 50 msec) at the

frequency of 0.2 Hz. The peak amplitude of the eIPSC was measured from the averaged

responses. The responses were acquired each minute for 10 min before and 12 min after OT

application. The paired pulse stimulus was used to calculate the PPR between the mean

amplitude of the second eIPSC divided by the amplitude of the first eIPSC. Here we used the

PPR to evaluate a possible presynaptic mechanism of the OT effect on the BNST neurons

inhibitory synaptic responses.

To verify that the effects of OT were mediated by OTRs, selective and potent OTR

antagonist d(CH2)5(1), D-Tyr(2), Thr(4), Orn(8), des-Gly-NH2(9)]-Vasotocin trifluoroacetate salt

(Manning et al., 2012) (0.4 μM, OTA, V-905 NIMH Chemical Repository) was applied 10 min

before OT in all whole-cell experiments. Recordings were performed with a Multiclamp 700B

amplifier (Molecular Device, Sunnyvale, CA, USA). Current-clamp and voltage-clamp signals

were acquired at 10 kHz with a 16-bit input-output board NI USB-6251 (National Instruments,

Austin, TX) using custom MatLab scripts (Math-Work, Natick, MA) written by Dr. Niraj Desai

(Desai et al., 2017). All neurons included in the analysis had a seal resistance > 1GΩ and an

initial access resistance < 20 MΩ maintained for the entire recording period. Any cells showing

a 10% change in the serial resistance were not included in the analysis.

4.3.1. Electrophysiological characterization of BNST$_{DL}$ neurons

Type I neurons are regular firing and display spike frequency adaptation. These neurons also

display moderate voltage sag that indicates the presence of the hyperpolarization-activated

cation current (I_h). As a distinguishing feature of Type II neurons, post-inhibitory spikes are

produced in response to preceding negative current steps that are related to the action of the

low-threshold Ca^{2+} current. Additionally, Type II neurons display strong voltage sags under

hyperpolarizing current pulses that indicate a high level of I_h current. Type III neurons differ from

the previous two types in several aspects: they exhibit high rheobase and no voltage sag under

negative current steps; they display prominent inward rectification that is caused by the

activation of the inward rectifying K^+ current (I_{KIR}) at membrane potentials more negative than

approximately −50 mV; and start firing after a characteristic slow voltage ramp mediated by the

K^+ delayed current (I_D).

4.3.2. Cell-attached recordings

The spontaneous firing activity of Type I and II BNST$_{DL}$ neurons was recorded in a cell-attached

voltage-clamp mode, using patch pipettes filled with K-gluconate solution as described above,

without blockers of the synaptic transmission. To avoid a possible effect of amplifier-generated

currents on the spontaneous firing activity during the recording the command potential was

carefully adjusted at the value at which no current was flowing from the amplifier through the cell

membrane as described by (Perkins, 2006). At the end of the spontaneous firing recording

period, a gentle negative pressure was applied to break through the membrane into whole cell

configuration and determine the neuron type in a current-clamp mode as described above.

4.3.3. The effect of OT on intrinsic membrane properties and spontaneous firing of Type

I-III BNST$_{DL}$ neurons

The amplitudes of the fAHP and mAHP following the first action potential evoked at Rh

were calculated as the difference from the action potential threshold to the membrane voltage

measured after 2-5 msec and 50-100 msec, respectively. The lAHP was measured at 2000

msec after the end of the depolarizing pulse, following generation of five action potentials. As

the K$^+$ currents underlying these AHPs are considered to regulate spike frequency adaptation

(Gu et al., 2008; Lopez de Armentia and Sah, 2004), we measured early and late spike

frequency adaptation before and after OT application using a depolarizing pulse of 1 sec. We

measured the spike frequency adaptation rather than the spike frequency, because the firing of

the BNST neurons, as described by Paré and colleagues (Rodríguez-Sierra et al., 2013), is

adapting and not regular. This is a characteristic of all types of BNST$_{DL}$ neurons, including Type

I (described as regular firing), which also adapt on a long current pulse. We also performed

recordings in cell-attached mode in Type I-II neurons and measured the spontaneous firing

frequency before and after OT. The spontaneous firing frequency (Hz) was determined as the

number of spontaneous spikes for every minute of recording.

4.3.4. The effect of OT on inhibitory synaptic activity of Type I-III BNST$_{DL}$ neurons

To study the effect of OT on spontaneous inhibitory postsynaptic currents (sIPSCs), the events

were recorded from Type I-III neurons in a voltage clamp mode at holding potential of -70 mV.

To isolate the sIPSCs, the recordings were made in the presence of the AMPA receptor

antagonist CNQX (10 μM) and the NMDA receptor antagonist AP-5 (50 μM). The sIPSCs were

characterized by their mean frequency, amplitude, and charge (calculated as the area under the

curve) using custom scripts in MatLab. To reach an adequate number of events, the sIPSCs

were acquired continuously in 2-minute sweeps during the entire recording time. The sIPSCs

were recorded for 10 min in aCSF, followed by 12 min bath application of OT and followed by

the drug washout (at least 10 more min after the OT application was completed). As sIPSCs are

a combination of spike-independent IPSCs and spike-driven IPSCs, in some experiments the

action potential generation was blocked with Tetrodotoxin (TTX, 1 μM, #1078; Tocris

Bioscience, UK), leaving only the spike-independent IPSCs (miniature IPSCs, mIPSCs). $GABA_A$

receptor antagonist, picrotoxin (25 μM), was used at the end of the recordings to verify the

GABA-ergic origin of the sIPSCs. Finally, to study the effect of OT on the evoked inhibitory

synaptic transmission in Type I-III neurons, we recorded the eIPSCs by electrical stimulation of

the Stria Terminalis.

4.3.5. CeA-projecting BNST$_{DL}$ neurons

To determine an electrophysiological phenotype of _BNST➔CeA_ neurons, rats were first

bilaterally injected into the CeA with a retrograde tracer, green microbeads (Lumafluor Inc., 100

nl; AP: -2.4 mm, ML: +4.2 mm, DV: -8.2 mm), and were allowed to recover for 4-7 days before

electrophysiological recordings. Stereotaxic surgeries were performed as before (Martinon et

al., 2019; Moaddab and Dabrowska, 2017). The slices were illuminated with CoolLED pE-

300lite (Scientifica Ltd., Uckfield, UK) and ET - EGFP (FITC/Cy2) filter (Chroma Technology

Corp, Bellows Falls, VT, US) to enable visually guided patching of fluorescent neurons. Slices

were stored in 10% stabilized formalin post-recordings, mounted with Krystalon Mounting

medium (Millipore Sigma, #64969-71), and analyzed with Nikon Eclipse E600 microscope

(Melville, NY, US) and Olympus Fv10i confocal microscope for high magnification imaging

(Center Valley, PA, US).

4.4. Statistical analysis

All data are presented as MEAN ± standard error of mean (SEM). Distribution of data sets was

first evaluated with Kolmogorov-Smirnov test. Based on the normal distribution, the effects of

OT on the sIPSCs, excitability, and firing frequency were analyzed by a repeated measures

(RM) one-way analysis of variance (ANOVA) or mixed effects model, or paired *t*-tests, when

applicable. Where the F-ratio was significant, all-pairwise post hoc comparisons were made

using Tukey's or Sidak's tests. The statistical analyses were completed using GraphPad Prism

version 8.4.1 (GraphPad Software Inc., San Diego, CA). $P < 0.05$ was considered significant.

5. Acknowledgements

We would like to thank Dr. Attilio Caravelli for his help with Matlab. We also thank Rachel

Chudoba for proofreading the manuscript. This work was supported by grant from the National

Institute of Mental Health R01MH113007 and R01MH113007-04S1 to JD and start-up funds

from the Chicago Medical School, Rosalind Franklin University of Medicine and Science to JD.

6. Contributions

JD conceived the study. JD and WF designed the study. WF, FB, and VOP acquired data. WF,

FB, VOP, and JD analyzed data. WF, FB, VOP and JD interpreted data. JD and WF wrote the

manuscript. JD obtained funding.

7. Financial Disclosures

The authors declare no conflict of interest and have no financial interests to disclose. Dr.

Joanna Dabrowska reports submission of a provisional patent application entitled: Method and

System for Testing for Stress-related Mental Disorders, Susceptibility, Disease Progression,

Disease Modulation and Treatment Efficacy (# 62/673447).

8. Figure legends

Figure 1. Oxytocin affects intrinsic membrane properties of Type I BNST$_{DL}$ neurons. A: A

representative trace showing that oxytocin (OT, 0.2 µM) shortens the latency to the 1st spike at

rheobase current (Rh) and depolarizes resting membrane potential (RMP, from -64 mV to -60

mV). This is associated with increased input resistance (Rin), measured with a depolarizing

current step below the Rh. OT also reduces Rh from 30 pA (pre) to 20 pA (OT) **B-D**: Line

graphs with individual neurons' responses (thin gray) and average responses for all neurons

recorded (thick black) showing that OT significantly increases the RMP (**B**, P = 0.0002, F

(1.593, 19.12) = 15.76, n = 14, mixed-effect analysis), increases Rin (**C**, P = 0.0085, n = 9,

paired t-test), reduces the Rh (**D**, P = 0.0457, F (1.341, 16.09) = 4.226, n = 14, mixed-effect

analysis), reduces the 1st spike threshold (**E**, P = 0.0164, F (1.472, 20.61) = 5.749, and reduced

the 1st spike latency (**F**, P = 0.0185, F (1.971, 23.66) = 4.775, n = 14, mixed-effect analysis).

Thick black lines show average responses for all Type I neurons recorded. Each data point

represents the mean ± SEM, **P < 0.01, *P < 0.05 in comparison to pre or post-OT.

Figure 2. Oxytocin increases intrinsic excitability of Type I, but not Type II or Type III,

BNST$_{DL}$ neurons. **A**: A representative trace showing voltage deflections in response to

hyperpolarizing and depolarizing current injections characteristic of Type I neuron (left) and the

increase in firing frequency of Type I neuron in the presence of OT respective to baseline (pre)

(right). **B**: OT causes a leftward shift of input/output (I-O) relationship in Type I neurons. The

steady state frequency (SSF) was determined as an average of an inverse of the inter-spike

intervals (ISI) from all action potentials starting from the second action potential. In Type I

neurons**,** OT shows a significant increment effect on SSF at 40 pA (P = 0.0024, F (1.667, 15.01)

= 10.11), at 50 pA (P = 0.0271, F (1.863, 16.76) = 4.628), and at 60 pA (P = 0.0271, F (1.688,

11.82) = 5.267, mixed-effects analysis). **C**: Application of OTR antagonist, OTA (0.4 µM),

prevents the leftward shift of the I-O relationship induced by OT application (40 pA, P = 0.3365,

50 pA, P = 0.4299, 60 pA, P = 0.6312, n = 5, RM ANOVA). **D-E:** OT application does not modify

the I-O relationship in Type II (**D**) or Type III BNST$_{DL}$ neurons (**E**). In Type II neurons, OT did not

affect SSF at 80 pA (P = 0.3116), 70 pA (P = 0.8236), 60 pA (P = 0.9519), 50 pA (P = 0.8993),

40 pA (P = 0.7636), 30 pA (P = 0.8673), or at 20 pA (P = 0.6990, **D**). Similarly, in Type III

neurons, there was no effect of OT on SSF at 80 pA (P = 0.1151), 70 pA (P = 0.1138), 60 pA (P

= 0.180), 50 pA (P = 0.0847), 40 pA (P = 0.1041), 30 pA (P = 0.0790), or at 20 pA (P = 0.0681,

E). Each data point represents the mean ± SEM, **P < 0.01, *P < 0.05 in comparison to pre-OT,

P < 0.01, # P < 0.05 in comparison to post-OT.

Figure 3. Oxytocin reduces fast and medium afterhyperpolarizations and modifies early

and late spike frequency adaptation in Type I BNST$_{DL}$ neurons. **A**: A representative trace

from Type I neuron showing a reduction of fast AHP (fAHP) and medium AHP (mAHP) after OT

application respective to baseline (pre). The amplitudes of the fAHP and mAHP following the

first action potential evoked at Rh were calculated as the difference from the action potential

threshold to the membrane voltage measured after 2-5 msec and 50-100 msec, respectively.

The lAHP was measured at 2000 msec after the end of the depolarizing pulse, following

generation of five action potentials. **B**: Line graphs showing individual neurons' responses (gray)

and reduction of the fAHP during OT application (P = 0.0124, F (1.473, 17.68 = 6.493, mixed-

effects analysis, n = 14). **C**: OTR antagonist, OTA (0.4 µM), prevents the OT-induced reduction

of fAHP (P = 0.2220, n = 7, RM one-way ANOVA). **D**: Line graph showing reduction of mAHP

during OT application (P <0.0001, F (1.844, 22.13) = 23.25, mixed effect analysis, n = 14). **E***:

OT-induced reduction of mAHP is blocked by OTR antagonist, OTA (P = 0.3968, n = 7, RM one-

way ANOVA). Thick black lines show average responses for all Type I neurons recorded. **F**: OT

modifies early and late spike frequency adaptation. The early and late spike frequency

adaptation was investigated using a depolarizing pulse of 1 sec and measuring the

instantaneous frequency for every action potential interval. The ISI number is the inter-stimulus

interval between two consecutive action potentials. OT significantly increased the instantaneous

frequency of ISI (P < 0.0001, F (1, 13) = 82.69, mixed effects model). Each data point

represents the mean ± SEM, ****P < 0.0001, ***P < 0.001, **P < 0.01, *P < 0.05.

Figure 4. Oxytocin increases spontaneous firing rate of Type I neurons and reduces spontaneous firing rate of Type II BNST$_{DL}$ neurons in cell-attached recordings. **A**: A representative trace of Type I neuron showing an increase in spontaneous firing rate after OT application. **B**: Line graph showing that OT increased spontaneous firing rate in all spontaneously firing Type I neurons (P = 0.0242, paired t-test, n = 7), gray lines represent individual neurons' responses, thick black line represents average response. **C**: A representative trace of Type II neuron showing a reduction in spontaneous firing rate after OT application. **D**: Line graph showing that OT reduced spontaneous firing rate in spontaneously firing Type II neurons in cell-attached mode (P = 0.0525, paired t-test, n = 6).

Figure 5. Oxytocin modulates inhibitory synaptic transmission in BNST$_{DL}$ neurons. A: Representative recordings showing an increase in the frequency of spontaneous inhibitory postsynaptic currents (sIPSCs) after OT application respective to baseline (pre). **B-C**: Line graphs (gray) showing that 12 out of 14 recorded Type II neurons showed an increase in sIPSCs frequency (**B**, P = 0.0022, F $(1.058, 12.17)$ = 14.41, n = 14, mixed-effect analysis) but not amplitude (**C**, P = 0.5855, F $(1.563, 17.19)$ = 0.4718, n = 14, mixed-effect analysis) after OT application. **D**: OT does not affect sIPSC frequency in a presence of OTR antagonist, OTA (0.4 µM) (P = 0.1788, F $(1.469, 8.814)$ = 2.135, n = 6, RM ANOVA). **E**: When spike-driven IPSCs were blocked with TTX (P = 0.0151, F $(1.030, 6.179)$ = 10.98, n = 6, RM ANOVA), OT does not affect the sIPSCs frequency. **F-G**: OT does not affect frequency (**F**, P = 0.9362) but reduces amplitude of sIPCS in Type III neurons (**G**, P = 0.0160, paired t-test). Thick black lines show averaged responses for all neurons recorded. Each data point represents the mean ± SEM, **P < 0.01, *P < 0.05 in comparison to pre- or post-OT.

Figure 6. Oxytocin increases amplitude of evoked IPSCs in Type III, but not Type II neurons. **A:** Representative traces of eIPSC in response to the electrical stimulation of the stria terminalis in Type II neurons. **B**: Line graph showing that OT does not significantly change the mean amplitude of eIPSC in Type II neurons (P = 0.2459, paired t-test). Thick black lines show

averaged responses for all neurons recorded. **C:** Representative traces of eIPSCs in Type III

neurons showing reduction of the eIPSCs amplitude after OT application. **D-E:** Line graphs

showing individual Type III neurons and the reduction of the eIPSC amplitude after OT (**D**, P =

0.0105) without any significant change in the PPR (**E**, P = 0.5672).

Figure 7. Type II neurons are output neurons of the $BNST_{DL}$ to the central amygdala. A-B:

Microphotographs showing neurons expressing green fluorescent microbeads in the $BNST_{DL}$

after the retrobeads were injected into the central amygdala (CeA). **A**: BNST neurons projecting

to the CeA (*BNST➔CeA* neurons) were primarily found in the dorsolateral BNST, primarily the

oval nucleus, and to a lesser extent in the anteromedial BNST (20x, LV – Lateral ventricle, IC –

Internal capsulae, $BNST_{AM}$ - anteriomedial BNST, $BNST_{OV}$ - oval nucleus of the BNST, scale

bar 100 μm). **B**: High magnification (60x) microphotographs showing green fluorescent beads

present in membranes and processes of the *BNST➔CeA* neurons (scale bar 10 μm). **C-D**:

Electrophysiological characterization of the *BNST➔CeA* neurons using visually guided patching

(**C**, 40x, scale bar 20 μm). Representative recording of the BNST➔CeA fluorescent neuron

showing the electrophysiological properties of a Type II neuron such as the presence of a sag

during the hyperpolarized current pulses and the spike rebound at the end of the 450 ms current

pulse injection (**D**). **E**: Diagram showing that OT regulates the intrinsic inhibitory network and

inhibits output neurons of the $BNST_{DL}$. By directly increasing frequency of spontaneous firing of

Type I regular spiking neurons, OT inhibits firing of Type II $BNST_{DL}$ neurons. Type II neurons

are output neurons of the $BNST_{DL}$ sending projections to the central amygdala (*BNST➔CeA*

neurons*). OT also inhibits another class of output neurons, Type III, via a direct postsynaptic

mechanism. Therefore, OT excites $BNST_{DL}$ interneurons (Type I) and inhibits two classes of the

$BNST_{DL}$ projection neurons via distinct indirect (Type II) and direct (Type III) mechanisms,

respectively.

9. References

al-Shamma HA, De Vries GJ. 1996. Neurogenesis of the sexually dimorphic vasopressin cells of

the bed nucleus of the stria terminalis and amygdala of rats. *J Neurobiol* **29**:91–98.

doi:10.1002/(SICI)1097-4695(199601)29:1<91::AID-NEU7>3.0.CO;2-2

Alger BE, Nicoll RA. 1980. Epileptiform burst afterhyperolarization: calcium-dependent

potassium potential in hippocampal CA1 pyramidal cells. *Science* **210**:1122–1124.

doi:10.1126/science.7444438

Alheid GF. 2003. Extended amygdala and basal forebrain. *Ann N Y Acad Sci* **985**:185–205.

Beyeler A, Dabrowska J. 2020. Neuronal diversity of the amygdala and the bed nucleus of the

stria terminalis. *Handb Behav Neurosci* **26**:63–100. doi:10.1016/b978-0-12-815134-

1.00003-9

Bosch OJ, Young LJ. 2018. Oxytocin and Social Relationships: From Attachment to Bond

Disruption. *Curr Top Behav Neurosci* **35**:97–117. doi:10.1007/7854_2017_10

Brussaard AB, Kits KS, de Vlieger TA. 1996. Postsynaptic mechanism of depression of

GABAergic synapses by oxytocin in the supraoptic nucleus of immature rat. *J Physiol*

497 (Pt 2):495–507. doi:10.1113/jphysiol.1996.sp021783

Caldeyro-Barcia R, Poseiro JJ. 1959. Oxytocin and contractility of the pregnant human uterus.

Ann N Y Acad Sci **75**:813–830.

Crane JW, Holmes NM, Fam J, Westbrook RF, Delaney AJ. 2020. Oxytocin increases inhibitory

synaptic transmission and blocks development of long-term potentiation in the lateral

amygdala. *J Neurophysiol* **123**:587–599. doi:10.1152/jn.00571.2019

Cullinan WE, Herman JP, Watson SJ. 1993. Ventral subicular interaction with the hypothalamic

paraventricular nucleus: evidence for a relay in the bed nucleus of the stria terminalis. *J*

Comp Neurol **332**:1–20. doi:10.1002/cne.903320102

Dabrowska J, Hazra R, Ahern TH, Guo J-D, McDonald AJ, Mascagni F, Muller JF, Young LJ,

Rainnie DG. 2011. Neuroanatomical evidence for reciprocal regulation of the

corticotrophin-releasing factor and oxytocin systems in the hypothalamus and the bed

nucleus of the stria terminalis of the rat: Implications for balancing stress and affect.

Psychoneuroendocrinology **36**:1312–1326. doi:10.1016/j.psyneuen.2011.03.003

Dabrowska J, Hazra R, Guo J-D, Dewitt S, Rainnie DG. 2013a. Central CRF neurons are not

created equal: phenotypic differences in CRF-containing neurons of the rat

paraventricular hypothalamus and the bed nucleus of the stria terminalis. *Front Neurosci*

7:156. doi:10.3389/fnins.2013.00156

Dabrowska J, Hazra R, Guo J-D, Li C, Dewitt S, Xu J, Lombroso PJ, Rainnie DG. 2013b.

Striatal-enriched protein tyrosine phosphatase-STEPs toward understanding chronic

stress-induced activation of corticotrophin releasing factor neurons in the rat bed nucleus

of the stria terminalis. *Biol Psychiatry* **74**:817–826. doi:10.1016/j.biopsych.2013.07.032

Dabrowska J, Martinon D, Moaddab M, Rainnie DG. 2016. Targeting Corticotropin-Releasing

Factor Projections from the Oval Nucleus of the Bed Nucleus of the Stria Terminalis

Using Cell-Type Specific Neuronal Tracing Studies in Mouse and Rat Brain. *J*

Neuroendocrinol **28**. doi:10.1111/jne.12442

Daniel SE, Guo J, Rainnie DG. 2017. A comparative analysis of the physiological properties of

neurons in the anterolateral bed nucleus of the stria terminalis in the Mus musculus,

Rattus norvegicus, and Macaca mulatta. *J Comp Neurol* **525**:2235–2248.

doi:10.1002/cne.24202

Daniel SE, Rainnie DG. 2016. Stress Modulation of Opposing Circuits in the Bed Nucleus of the

Stria Terminalis. *Neuropsychopharmacology* **41**:103–125. doi:10.1038/npp.2015.178

Davis M, Walker DL, Miles L, Grillon C. 2010. Phasic vs sustained fear in rats and humans: role

of the extended amygdala in fear vs anxiety. *Neuropsychopharmacology* **35**:105–135.

doi:10.1038/npp.2009.109

Dumais KM, Bredewold R, Mayer TE, Veenema AH. 2013. Sex differences in oxytocin receptor

binding in forebrain regions: correlations with social interest in brain region- and sex-

specific ways. *Horm Behav* **64**:693–701. doi:10.1016/j.yhbeh.2013.08.012

Duvarci S, Bauer EP, Paré D. 2009. The bed nucleus of the stria terminalis mediates inter-

individual variations in anxiety and fear. *J Neurosci* **29**:10357–10361.

doi:10.1523/JNEUROSCI.2119-09.2009

Eliava M, Melchior M, Knobloch-Bollmann HS, Wahis J, da Silva Gouveia M, Tang Y, Ciobanu

AC, Triana Del Rio R, Roth LC, Althammer F, Chavant V, Goumon Y, Gruber T, Petit-

Demoulière N, Busnelli M, Chini B, Tan LL, Mitre M, Froemke RC, Chao MV, Giese G,

Sprengel R, Kuner R, Poisbeau P, Seeburg PH, Stoop R, Charlet A, Grinevich V. 2016.

A New Population of Parvocellular Oxytocin Neurons Controlling Magnocellular Neuron

Activity and Inflammatory Pain Processing. *Neuron* **89**:1291–1304.

doi:10.1016/j.neuron.2016.01.041

Francesconi W, Berton F, Repunte-Canonigo V, Hagihara K, Thurbon D, Lekic D, Specio SE,

Greenwell TN, Chen SA, Rice KC, Richardson HN, O'Dell LE, Zorrilla EP, Morales M,

Koob GF, Sanna PP. 2009. Protracted withdrawal from alcohol and drugs of abuse

impairs long-term potentiation of intrinsic excitability in the juxtacapsular bed nucleus of

the stria terminalis. *J Neurosci* **29**:5389–5401. doi:10.1523/JNEUROSCI.5129-08.2009

Goode TD, Acca GM, Maren S. 2020. Threat imminence dictates the role of the bed nucleus of

the stria terminalis in contextual fear. *Neurobiol Learn Mem* **167**:107116.

doi:10.1016/j.nlm.2019.107116

Goode TD, Ressler RL, Acca GM, Miles OW, Maren S. 2019. Bed nucleus of the stria terminalis

regulates fear to unpredictable threat signals. *Elife* **8**. doi:10.7554/eLife.46525

Gozzi A, Jain A, Giovannelli A, Giovanelli A, Bertollini C, Crestan V, Schwarz AJ, Tsetsenis T,

Ragozzino D, Gross CT, Bifone A. 2010. A neural switch for active and passive fear.

Neuron **67**:656–666. doi:10.1016/j.neuron.2010.07.008

Grundwald NJ, Benítez DP, Brunton PJ. 2016. Sex-Dependent Effects of Prenatal Stress on

Social Memory in Rats: A Role for Differential Expression of Central Vasopressin-1a

Receptors. *J Neuroendocrinol* **28**. doi:10.1111/jne.12343

Gu N, Hu H, Vervaeke K, Storm JF. 2008. SK (KCa2) channels do not control somatic

excitability in CA1 pyramidal neurons but can be activated by dendritic excitatory

synapses and regulate their impact. *J Neurophysiol* **100**:2589–2604.

doi:10.1152/jn.90433.2008

Gu N, Vervaeke K, Hu H, Storm JF. 2005a. Kv7/KCNQ/M and HCN/h, but not KCa2/SK

channels, contribute to the somatic medium after-hyperpolarization and excitability

control in CA1 hippocampal pyramidal cells. *J Physiol (Lond)* **566**:689–715.

doi:10.1113/jphysiol.2005.086835

Gu N, Vervaeke K, Hu H, Storm JF. 2005b. Kv7/KCNQ/M and HCN/h, but not KCa2/SK

channels, contribute to the somatic medium after-hyperpolarization and excitability

control in CA1 hippocampal pyramidal cells. *J Physiol (Lond)* **566**:689–715.

doi:10.1113/jphysiol.2005.086835

Gu N, Vervaeke K, Storm JF. 2007. BK potassium channels facilitate high-frequency firing and

cause early spike frequency adaptation in rat CA1 hippocampal pyramidal cells. *J*

Physiol (Lond) **580**:859–882. doi:10.1113/jphysiol.2006.126367

Gungor NZ, Paré D. 2016. Functional Heterogeneity in the Bed Nucleus of the Stria Terminalis.

J Neurosci **36**:8038–8049. doi:10.1523/JNEUROSCI.0856-16.2016

Gungor NZ, Yamamoto R, Paré D. 2015. Optogenetic study of the projections from the bed

nucleus of the stria terminalis to the central amygdala. *J Neurophysiol* **114**:2903–2911.

doi:10.1152/jn.00677.2015

Hammack SE, Mania I, Rainnie DG. 2007. Differential expression of intrinsic membrane

currents in defined cell types of the anterolateral bed nucleus of the stria terminalis. *J*

Neurophysiol **98**:638–656. doi:10.1152/jn.00382.2007

Han JS, Maeda Y, Knepper MA. 1993. Dual actions of vasopressin and oxytocin in regulation of

water permeability in terminal collecting duct. *Am J Physiol* **265**:F26-34.

doi:10.1152/ajprenal.1993.265.1.F26

Harden SW, Frazier CJ. 2016. Oxytocin depolarizes fast-spiking hilar interneurons and induces

GABA release onto mossy cells of the rat dentate gyrus. *Hippocampus* **26**:1124–1139.

doi:10.1002/hipo.22595

Huber D, Veinante P, Stoop R. 2005. Vasopressin and oxytocin excite distinct neuronal

populations in the central amygdala. *Science* **308**:245–248.

doi:10.1126/science.1105636

Hung LW, Neuner S, Polepalli JS, Beier KT, Wright M, Walsh JJ, Lewis EM, Luo L, Deisseroth

K, Dölen G, Malenka RC. 2017. Gating of social reward by oxytocin in the ventral

tegmental area. *Science* **357**:1406–1411. doi:10.1126/science.aan4994

Ingram CD, Cutler KL, Wakerley JB. 1990. Oxytocin excites neurones in the bed nucleus of the

stria terminalis of the lactating rat in vitro. *Brain Res* **527**:167–170.

Ingram CD, Moos F. 1992. Oxytocin-containing pathway to the bed nuclei of the stria terminalis

of the lactating rat brain: immunocytochemical and in vitro electrophysiological evidence.

Neuroscience **47**:439–452.

Janeček M, Dabrowska J. 2019. Oxytocin facilitates adaptive fear and attenuates anxiety

responses in animal models and human studies-potential interaction with the

corticotropin-releasing factor (CRF) system in the bed nucleus of the stria terminalis

(BNST). *Cell Tissue Res* **375**:143–172. doi:10.1007/s00441-018-2889-8

Knobloch HS, Charlet A, Hoffmann LC, Eliava M, Khrulev S, Cetin AH, Osten P, Schwarz MK,

Seeburg PH, Stoop R, Grinevich V. 2012. Evoked axonal oxytocin release in the central

amygdala attenuates fear response. *Neuron* **73**:553–566.

doi:10.1016/j.neuron.2011.11.030

Kremarik P, Freund-Mercier MJ, Stoeckel ME. 1991. Autoradiographic detection of oxytocin-

and vasopressin-binding sites in various subnuclei of the bed nucleus of the stria

terminalis in the rat. Effects of functional and experimental sexual steroid variations. *J*

Neuroendocrinol **3**:689–698. doi:10.1111/j.1365-2826.1991.tb00335.x

Lopez de Armentia M, Sah P. 2004. Firing properties and connectivity of neurons in the rat

lateral central nucleus of the amygdala. *J Neurophysiol* **92**:1285–1294.

doi:10.1152/jn.00211.2004

Maejima S, Ohishi N, Yamaguchi S, Tsukahara S. 2015. A neural connection between the

central part of the medial preoptic nucleus and the bed nucleus of the stria terminalis to

regulate sexual behavior in male rats. *Neurosci Lett* **606**:66–71.

doi:10.1016/j.neulet.2015.08.047

Maniezzi C, Talpo F, Spaiardi P, Toselli M, Biella G. 2019. Oxytocin Increases Phasic and Tonic

GABAergic Transmission in CA1 Region of Mouse Hippocampus. *Front Cell Neurosci*

13:178. doi:10.3389/fncel.2019.00178

Manning M, Misicka A, Olma A, Bankowski K, Stoev S, Chini B, Durroux T, Mouillac B, Corbani

M, Guillon G. 2012. Oxytocin and vasopressin agonists and antagonists as research

tools and potential therapeutics. *J Neuroendocrinol* **24**:609–628. doi:10.1111/j.1365-

2826.2012.02303.x

Marlin BJ, Mitre M, D'amour JA, Chao MV, Froemke RC. 2015. Oxytocin enables maternal

behaviour by balancing cortical inhibition. *Nature* **520**:499–504.

doi:10.1038/nature14402

Martina M, Kilić G, Cherubini E. 1994. The effect of intracellular Ca2+ on GABA-activated

currents in cerebellar granule cells in culture. *J Membr Biol* **142**:209–216.

doi:10.1007/BF00234942

Martinon D, Lis P, Roman AN, Tornesi P, Applebey SV, Buechner G, Olivera V, Dabrowska J.

2019. Oxytocin receptors in the dorsolateral bed nucleus of the stria terminalis (BNST)

bias fear learning toward temporally predictable cued fear. *Transl Psychiatry* **9**:140.

doi:10.1038/s41398-019-0474-x

McDonald AJ, Shammah-Lagnado SJ, Shi C, Davis M. 1999. Cortical afferents to the extended

amygdala. *Ann N Y Acad Sci* **877**:309–338.

Meloni EG, Jackson A, Gerety LP, Cohen BM, Carlezon WA. 2006. Role of the bed nucleus of

the stria terminalis (BST) in the expression of conditioned fear. *Ann N Y Acad Sci*

1071:538–541. doi:10.1196/annals.1364.059

Moaddab M, Dabrowska J. 2017. Oxytocin receptor neurotransmission in the dorsolateral bed

nucleus of the stria terminalis facilitates the acquisition of cued fear in the fear-

potentiated startle paradigm in rats. *Neuropharmacology* **121**:130–139.

doi:10.1016/j.neuropharm.2017.04.039

Nakajima M, Görlich A, Heintz N. 2014. Oxytocin modulates female sociosexual behavior

through a specific class of prefrontal cortical interneurons. *Cell* **159**:295–305.

doi:10.1016/j.cell.2014.09.020

Neumann ID, Slattery DA. 2016. Oxytocin in General Anxiety and Social Fear: A Translational

Approach. *Biol Psychiatry* **79**:213–221. doi:10.1016/j.biopsych.2015.06.004

Nickerson K, Bonsness RW, Douglas RG, Condliffe P, Du Vigneaud V. 1954. Oxytocin and milk

ejection. *Am J Obstet Gynecol* **67**:1028–1034.

Okamura H, Abitbol M, Julien JF, Dumas S, Bérod A, Geffard M, Kitahama K, Bobillier P, Mallet

J, Wiklund L. 1990. Neurons containing messenger RNA encoding glutamate

decarboxylase in rat hypothalamus demonstrated by in situ hybridization, with special

emphasis on cell groups in medial preoptic area, anterior hypothalamic area and

dorsomedial hypothalamic nucleus. *Neuroscience* **39**:675–699. doi:10.1016/0306-

4522(90)90252-y

Olivera-Pasilio V, Dabrowska J. 2020. Oxytocin Promotes Accurate Fear Discrimination and

Adaptive Defensive Behaviors. *Front Neurosci* **14**:583878.

doi:10.3389/fnins.2020.583878

Owen SF, Tuncdemir SN, Bader PL, Tirko NN, Fishell G, Tsien RW. 2013. Oxytocin enhances

hippocampal spike transmission by modulating fast-spiking interneurons. *Nature*

500:458–462. doi:10.1038/nature12330

Pedarzani P, Storm JF. 1993. PKA mediates the effects of monoamine transmitters on the K+

current underlying the slow spike frequency adaptation in hippocampal neurons. *Neuron*

11:1023–1035. doi:10.1016/0896-6273(93)90216-e

Pedersen CA, Caldwell JD, Peterson G, Walker CH, Mason GA. 1992. Oxytocin activation of

maternal behavior in the rat. *Ann N Y Acad Sci* **652**:58–69.

Perkins KL. 2006. Cell-attached voltage-clamp and current-clamp recording and stimulation

techniques in brain slices. *J Neurosci Methods* **154**:1–18.

doi:10.1016/j.jneumeth.2006.02.010

Rodríguez-Sierra OE, Turesson HK, Pare D. 2013. Contrasting distribution of physiological cell

types in different regions of the bed nucleus of the stria terminalis. *J Neurophysiol*

110:2037–2049. doi:10.1152/jn.00408.2013

Sharma K, LeBlanc R, Haque M, Nishimori K, Reid MM, Teruyama R. 2019. Sexually dimorphic

oxytocin receptor-expressing neurons in the preoptic area of the mouse brain. *PLoS*

ONE **14**:e0219784. doi:10.1371/journal.pone.0219784

Simerly RB, Swanson LW. 1988. Projections of the medial preoptic nucleus: a Phaseolus

vulgaris leucoagglutinin anterograde tract-tracing study in the rat. *J Comp Neurol*

270:209–242. doi:10.1002/cne.902700205

Sparta DR, Jennings JH, Ung RL, Stuber GD. 2013. Optogenetic strategies to investigate neural

circuitry engaged by stress. *Behav Brain Res* **255**:19–25. doi:10.1016/j.bbr.2013.05.007

Sullivan GM, Apergis J, Bush DEA, Johnson LR, Hou M, Ledoux JE. 2004. Lesions in the bed

nucleus of the stria terminalis disrupt corticosterone and freezing responses elicited by a

contextual but not by a specific cue-conditioned fear stimulus. *Neuroscience* **128**:7–14.

doi:10.1016/j.neuroscience.2004.06.015

Sun N, Cassell MD. 1993. Intrinsic GABAergic neurons in the rat central extended amygdala. *J*

Comp Neurol **330**:381–404. doi:10.1002/cne.903300308

Tabbaa M, Paedae B, Liu Y, Wang Z. 2016. Neuropeptide Regulation of Social Attachment: The

Prairie Vole Model. *Compr Physiol* **7**:81–104. doi:10.1002/cphy.c150055

Terburg D, Scheggia D, Triana Del Rio R, Klumpers F, Ciobanu AC, Morgan B, Montoya ER,

Bos PA, Giobellina G, van den Burg EH, de Gelder B, Stein DJ, Stoop R, van Honk J.

2018. The Basolateral Amygdala Is Essential for Rapid Escape: A Human and Rodent

Study. *Cell* **175**:723-735.e16. doi:10.1016/j.cell.2018.09.028

Tirko NN, Eyring KW, Carcea I, Mitre M, Chao MV, Froemke RC, Tsien RW. 2018. Oxytocin

Transforms Firing Mode of CA2 Hippocampal Neurons. *Neuron* **100**:593-608.e3.

doi:10.1016/j.neuron.2018.09.008

Tribollet E, Dubois-Dauphin M, Dreifuss JJ, Barberis C, Jard S. 1992. Oxytocin receptors in the

central nervous system. Distribution, development, and species differences. *Ann N Y*

Acad Sci **652**:29–38.

Veinante P, Freund-Mercier MJ. 1997. Distribution of oxytocin- and vasopressin-binding sites in

the rat extended amygdala: a histoautoradiographic study. *J Comp Neurol* **383**:305–325.

Verbalis JG, Blackburn RE, Olson BR, Stricker EM. 1993. Central oxytocin inhibition of food and

salt ingestion: a mechanism for intake regulation of solute homeostasis. *Regul Pept*

45:149–154.

Viviani D, Charlet A, van den Burg E, Robinet C, Hurni N, Abatis M, Magara F, Stoop R. 2011.

Oxytocin selectively gates fear responses through distinct outputs from the central

amygdala. *Science* **333**:104–107. doi:10.1126/science.1201043

Waddell J, Morris RW, Bouton ME. 2006. Effects of bed nucleus of the stria terminalis lesions

on conditioned anxiety: aversive conditioning with long-duration conditional stimuli and

reinstatement of extinguished fear. *Behav Neurosci* **120**:324–336. doi:10.1037/0735-

7044.120.2.324

Wilson BC, Terenzi MG, Ingram CD. 2005. Differential excitatory responses to oxytocin in sub-

divisions of the bed nuclei of the stria terminalis. *Neuropeptides* **39**:403–407.

doi:10.1016/j.npep.2005.04.001

Xiao L, Priest MF, Nasenbeny J, Lu T, Kozorovitskiy Y. 2017. Biased Oxytocinergic Modulation

of   Midbrain   Dopamine   Systems.   *Neuron*   **95**:368-384.e5.

doi:10.1016/j.neuron.2017.06.003

Xu X, Ikrar T, Sun Y, Santos R, Holmes TC, Francesconi W, Berton F. 2016. High-resolution

and cell-type-specific photostimulation mapping shows weak excitatory vs. strong

inhibitory inputs in the bed nucleus of the stria terminalis. *J Neurophysiol* **115**:3204–

3216. doi:10.1152/jn.01148.2015

Zaninetti M, Raggenbass M. 2000. Oxytocin receptor agonists enhance inhibitory synaptic

transmission in the rat hippocampus by activating interneurons in stratum pyramidale.

Eur J Neurosci **12**:3975–3984. doi:10.1046/j.1460-9568.2000.00290.x