Systematic evaluation of parameters for genome-scale metabolic models of cultured mammalian cells

Song-Min Schinn[1], Carly Morrison[2], Wei Wei[2], Lin Zhang[2], Nathan E. Lewis[1,3,4]

1. Department of Pediatrics, University of California, San Diego

2. Pfizer, Biotherapeutics Pharmaceutical Sciences, Andover, MA

3. Department of Bioengineering, University of California, San Diego

4. Novo Nordisk Foundation Center for Biosustainability at UC San Diego

Abstract

Genome-scale metabolic models describe cellular metabolism with mechanistic detail. Given their high complexity, such models need to be parameterized correctly to yield accurate predictions and avoid overfitting. Effective parameterization has been well-studied for microbial models, but it remains unclear for higher eukaryotes, including mammalian cells. To address this, we enumerated model parameters that describe key features of cultured mammalian cells – including cellular composition, bioprocess performance metrics, mammalian-specific pathways, and biological assumptions behind model formulation approaches. We tested these parameters by building thousands of metabolic models and evaluating their ability to predict the growth rates of a panel of phenotypically diverse Chinese Hamster Ovary cell clones. We found the following considerations to be most critical for accurate parameterization: (1) cells limit metabolic activity to maintain homeostasis, (2) cell morphology and viability change dynamically during a growth curve, and (3) cellular biomass has a particular macromolecular composition. Depending on parameterization, models predicted different metabolic phenotypes, including contrasting mechanisms of nutrient utilization and energy generation, leading to varying accuracies of growth rate predictions. Notably, accurate parameter values broadly agreed with experimental measurements. These insights will guide future investigations of mammalian metabolism.

1 *Introduction*

Cultured mammalian cells are prominent expression systems for large-scale biotherapeutics

manufacturing. However, conventional bioprocess engineering has largely been empirical due to

lacking knowledge of cell biology. Recent advances in high dimensional data and computational

methods are improving and accelerating industrial cell line and bioprocess development[1–3]. In

particular, metabolism has been a key target in cell line development[4–8], given its role in meeting

the anabolic and bioenergetic needs of cell growth and protein production[9,10]. Correspondingly,

there has been increased interest to investigate the metabolism of mammalian cells, such as

Chinese hamster ovary (CHO) cells, using genome-scale metabolic network models[11–18]. These

models contextualize large-scale biological data with curated biochemical knowledge, and have

been used with a wide array of *in silico* methods[19,20] to probe the molecular basis of

metabolism[21,22], disease[23–25], and human-microbiome interactions[26,27]. More recently, metabolic

network models are being applied to industrial bioprocess development[28–30] – e.g. predict

metabolic phenotype[31,13,32], identify metabolic bottlenecks[16], and optimize media formulation[15,33].

A general challenge of genome-scale metabolic network modeling is to identify relevant

insights from an underdetermined solution space[34], all the while avoiding overfitting. That is,

these models typically consist of thousands of metabolites, and reactions that far outnumber

experimental measurements available as boundary conditions, often leading to an

underdetermined system that is vulnerable to overfitting. To address this challenge, the following

established steps can be taken: (1) create a context-specific sub-model consisting of the most

relevant metabolic reactions for a given context by interpreting gene expression data[17,35,36], (2)

input experimental data as boundary conditions, and (3) hypothesize likely metabolic flux values

by applying optimality principles to the model reaction network[37–39]. Each of these steps invite

the modeler to make assumptions about cellular features – e.g. cellular makeup[40,41], the relevance

of specific pathways[42,43] and 'cellular objectives'[38,39,44–46] – that improve the model's solution

space reflect biology. Here, we define the model implementation of these assumptions as 'model

parameters.' In the last two decades, such model parameters have been extensively explored and

refined for model microbes such as *Escherichia coli*[47,48], *Saccharomyces cerevisae*[49,50] and

others[51,52]. However, parameters for higher eukaryotes remain under-characterized, despite

notable recent advances[53–55].

Mammalian cells are distinct from microbes in their metabolic pathways, auxotrophic

requirements, sub-compartmentalization and regulation[56], and therefore require parameterization

that capture such differences. A few studies have investigated mammalian-specific parameters

describing cell biomass[57], novel metabolic pathways[12] and cellular objectives[58,14]. These

parameters, however, have not been rigorously evaluated against wide-ranging conditions;

meanwhile, other features of mammalian metabolism and growth remain unexplored altogether.

To address this, we systematically investigate a panel of parameters for describing diverse

cultured mammalian cell metabolic phenotypes, with a focus on CHO cells. Our work identifies

key parameters for predicting metabolic phenotypes, challenges conventional parameterization

approaches, and guides future modeling efforts for mammalian metabolism.

Results

A cell line-specific model was constructed for phenotypically diverse CHO clones

Cultured mammalian cells, such as CHO cells, consume media nutrients – e.g. glucose,

amino acids and cofactors – to produce biomass, recombinant proteins and byproducts.

Depending on genotype and environmental conditions, CHO cells vary in their metabolic rates

and pathway usage, resulting in high metabolic heterogeneity. Here, we have examined a variety

of CHO metabolic phenotypes to identify general modeling principles and avoid overfitting.

Specifically, we investigated 10 CHO clones that differed in recombinant antibody, bioreactor

conditions and gene knockout treatment (see Methods). Consequently, the cells exhibited

varying patterns of nutrient consumption and byproduct secretion (Fig. 1A), ultimately leading to

divergent growth and productivity phenotypes (Fig. 1B; Fig. S1). We also observed the 10 clones

between culture days 4 to 11 (Fig. 1C) for temporal variation in metabolism[59,60].

Broadly, we saw several distinct types of metabolisms and nutrient utilization efficiencies.

Some cells displayed high glucose consumption and high proliferation, suggesting an efficient

and fast-moving metabolism (e.g. clones Z2, Z3). Other cells showed low glucose consumption

and low proliferation but high protein production, suggesting an efficient but attenuated

metabolism (e.g. clones A1, A2). Still others displayed high glucose consumption but low

proliferation and low protein production, suggesting a severely inefficient metabolism under

cellular stress.

We hypothesized that these diverse CHO metabolisms could be described by a well-

parameterized metabolic network model. To test this, we constructed a cell line-specific

metabolic network model by modifying a previously published genome-scale model of CHO

metabolism[11]. Briefly, this was done by the following steps: (1) transcriptomics data were

analyzed to quantify the relative activity of 210 metabolic tasks to hypothesize a list of active

metabolic genes[36,61,62], (2) these active genes and their associated model reactions were refined

into a fully functional model via the mCADRE algorithm[63], and (3) the subsequent draft model

was manually curated.

Mammalian-specific parameters embedded biological assumptions onto model

To enable the curated model of describing diverse CHO metabolisms, we identified salient

features of mammalian cell cultures and formulated them as model parameters (Table 1; see

Supplementary Document for detailed and technical discussions on their computational

implementation, including annotated code). First, we considered cellular biomass'

macromolecular makeup and dry weight (Table 1, parameters 1-2). These parameters are

necessary for the model to describe proliferation via the biomass production reaction[37], and to

convert experimental measurements (*per cell* units) into model-compatible values (*per cellular*

dry weight units). In other words, these parameters are critical to bounding the model solutions to

experimental data. While the importance of these parameters are well-established

previously[57,41,64,40], here we compare them with other novel parameters in the context of

mammalian cell biology.

Second, we considered that cellular dry weight change during fed-batch culture, indicating

internal shifts in metabolism and productivity[65]. Indeed, in the presented experiments, cellular

mass increased by as much as 40-70% throughout the culture period (Fig. S2). Up to now,

modeling efforts have treated this feature as static, even for multi-week time-course studies. A

dynamic cellular dry weight could facilitate more accurate experiment-to-model unit conversions.

It could also refine calculations between proliferation and biomass production. For example, it

would enable the model to consider that CHO cells may still produce significant amounts of

biomass during stationary phase when they proliferate more slowly but grow in size (see also Fig.

S2C). The impact of these considerations on model results is unclear.

We also observed cell loss despite high viability (>99%) during the transition between

exponential growth phase and stationary phase (Fig. S4), presumably due to shear stress-induced

apoptosis[66,67]. The cell loss was primarily observed on clones grown in Fed Batch #1 (Fig. 1, Fig

S4A). Accounting for these lost cells could improve calculations of total biomass produced

during later stages of cell culture. We explore these several bioprocess-related phenomena (Table

1, parameters 3-5) as novel model parameters.

Third, we improved descriptions of mammalian metabolic pathways which are not found in

microbes (Table 1, parameters 6-8). For example, we included in the model an intricate multi-

organelle secretory pathway[68] to improved estimations of metabolic expenditures towards

producing complex recombinant proteins (Table S3). Similarly, we also considered metabolic

pathways for producing various byproducts from amino acid catabolism[69] (Table S4), and

cellular death and turnover[70].

Fourth, we reevaluated the biological premises behind applying the optimality principle to

the metabolic network model (Table 1, parameter 9). This mathematical operation assumes that

cellular metabolism operates at some optimal condition and is crucial to constraining the

metabolic network model. As an example, proliferative bacteria have often been described to

maximize biomass production for rapid growth, which is computationally formulated as a

biomass objective function[37]. This objective function presumes that biomass production is

principally limited by nutrient availability. Meanwhile, a variety of alternative objective

functions have also been explored[38,39] – e.g. optimal energy generation per substrate[71],

minimized redox potential[58,72,73], efficient use of enzyme capacity[74,75] and streamlined amino

acid transport[14]. Compared to microbes, mammalian cells are larger, more complex and wired for

multicellularity, all of which increase the costs of homeostatic maintenance and limit

proliferation. Therefore, we compare the conventional *maximize biomass production* function

against alternative objective functions describing mammalian metabolism as chiefly limited by

various homeostasis requirements – e.g. thermal[76], proteomic[77,78], reactive oxygen species or

redox homeostasis[79]. These alternative objective functions were realized by imposing 'penalties'

on model reactions (see Methods; Table S5, 6).

Experimental measurements help produce robust model parameters

We investigated how the above-described model parameters impacted model prediction of

metabolism. The nine parameters were randomly permuted in value to within-literature-reported

ranges thousands of times. A resulting set of specified parameter values is called here as a

'parameter setting'. In total, 4000 distinct parameter settings were generated and used to bound

the solution space. We tested these parameter settings against experimental data for 10 clones

between culture days 4 to 11, resulting in 320,000 model predictions (Fig. 2a; Table S7). Each of

these predictions yielded a model-predicted growth rate that could be compared to experimental

measured growth rates (Fig. 2b). For ease of interpretation, prediction errors were preprocessed

to yield an 'accuracy' metric (Fig. 2c). Specifically, the prediction errors were transformed by a

negative-log function and normalized so that the average error metric for all clones and

timepoints would lie between 0 and 1. For reference, when growth rates were predicted to within

5% of experimental measurements for all 80 points, the accuracy metric equaled 0.8.

Overall, the various parameter settings resulted in a wide range of prediction accuracies

(Fig. S4). About one-eighth of the parameter settings robustly described at least eight of the ten

clones with high accuracy (accuracy > 0.8). Half of the parameter settings predicted only two or

fewer clones with high accuracy, indicating that the models were highly overfit or poorly

parameterized. The remaining one-third of the parameter settings predicted a moderate range of

clones with high accuracy, suggesting various degrees of overfitting. Overfit parameters of one

clone tended to also describe biologically similar clones better, recapitulating known differences

in cell line lineage and bioprocess conditions (Fig. S5).

Experimental data helped produce robust models. A model parameterized according to

experimental measurements (Table 1, parenthesized) yielded an average accuracy of 0.82 across

all clones and timepoints (Fig. 2b; 2c, blue line). Likewise, parameter settings with values close

to experimental measurements (within ±15%) resulted in comparable accuracies (0.83 ± 0.11).

Only a minority of parameter settings (2%) significantly deviated from experimental values yet

managed to perform comparably. The value of experimental measurements was especially

apparent for the *cell death rate* parameter describing late-stage cell death observed only in some

clones. The *cell death rate* parameter helped the model recapitulate observed cross-clonal

variation (Fig. S6).

Selection of objective function has greatest influence on model prediction

We estimated the importance of the parameters for prediction accuracy using a linear

regression model (Fig. 3). Of the nine parameters, *objective function formulation* had an outsized

influence on prediction accuracy. Specifically, the *maximize biomass production* function was

correlated strongly with poor predictions. In contrast, the *minimize cytosolic NADPH*

regeneration function was well correlated with accurate predictions. The following parameters

also improved model predictions: *cell death rate*, *biomass composition* and *consideration of*

dynamic dry weights to calculate growth rate (Fig. S8). Interestingly, the time-course of biomass

weight was more important than the baseline weight value itself. The remaining parameters had

negligible impact on model accuracy (effect size < 0.01; see Fig. S7).

The *objective function formulations* provided detailed and mechanistic hypotheses of the

intracellular metabolic activities underlying the growth rate predictions (Fig. 4), as revealed by

flux sampling analysis (see Methods). The *maximize biomass production* objective function

predicted an efficient and highly active metabolism, which utilized almost all consumed

substrates towards generating energy and biomass. As a result, the model consistently

overestimated growth rates across clones and timepoints (Fig. S9). Overestimations were

particularly pronounced for inefficient metabolisms, such as clones B1 and B2. In contrast,

several homeostasis-related objective functions predicted less efficient metabolisms that partially

discard consumed nutrients via various shunting mechanisms. This assumption, supported by

carbon balance estimations (Fig. S10), was key to predicting CHO metabolisms with differing

nutrient utilization efficiencies in a well-rounded manner.

Specifically, the homeostasis-related objective functions hypothesized two different carbon

shunting mechanisms: (1) secretion of glycerol, which is converted from the glycolytic

intermediate dihydroxyacetone phosphate via glycerol-3-phosphate dehydrogenase and glycerol-

3-phosphate phosphatase (Fig. 3, *B*), and (2) secretion of malate, an intermediate of the citric acid cycle and malate-aspartate shuttle (Fig. 3, *D*). The objective functions *minimize enzyme costs*, *minimize Gibbs energy dissipation* and *minimize cytosolic NADPH regeneration* hypothesized shunting via glycerol, with varied fluxes through glycolysis, glycerol synthesis, mitochondrial respiration and pyruvate carboxylate. The objective function *minimize mitochondrial NADH regeneration* hypothesized shunting via malate. This loss of flux in the citric acid cycle was then counteracted by converting carbon dioxide from respiration to bicarbonate, to eventually oxaloacetate via pyruvate carboxylate[80] (Fig. 3, *F*). The secretions of both glycerol and malate have been reported previously in CHO fed-batch systems[81,82,10], although only glycerol was observed in the present study to accumulate in the spent medium (Fig. S11).

Lastly, the *minimize ROS synthesis* objective function hypothesized negligible activities in glycolysis and citric acid cycle, deviating from known CHO biology[83]. Instead, most consumed glucose was diverted to the folate cycle via phosphoserine transamination to produce energy and byproduct formic acid (Fig. 3, *C*), resulting in an improbable metabolic phenotype. Additional comparison of objective functions and their influence on metabolic configurations are also provided (Fig. S12).

Discussion

Parameters help constrain metabolic networks sufficiently and accurately by embedding knowledge and data onto models. We compared thousands of distinct parameter settings for their ability to describe CHO clones with wide-ranging phenotypes. The resulting analyses lead to important insights for parameterizing metabolic network models of cultured mammalian cells. Specifically, this study (1) confirms that experimental data improve model parameterization, (2) identifies relevant parameters for mammalian metabolic models, most prominently the *objective function formulation*, and (3) challenges the popular use of the *maximize biomass production* objective function for modeling exponentially growing mammalian cells and explores promising alternatives.

Our analysis of model parameters agreed with experimental observations in several notable ways. First, experimentally measured parameter values resulted in robust and accurate model predictions, confirming the value of experimental measurements in parameterization. Second, parameter settings that described one clone well also tended to describe similar clones well, recapitulating differences in cell line lineages and glucose-feeding strategies. Lastly, parameter values of *cell death rate* recapitulated experimental observations at the clone level. These broad agreements demonstrate that well-parameterized metabolic network models can describe diverse metabolic states.

Our analysis identified parameters that strongly affected model precision. Of all the investigated parameters, the *objective function formulation* had a predominant influence on model-predicted growth rates and their underlying metabolic activities. *Biomass composition* was also reconfirmed as a key parameter, agreeing with previous findings in microbial models[40,64,41]. Lastly, novel parameters accounting for time-course changes in cell size and viability improved model predictions substantially. This emphasizes the relevance of dynamic and bioprocess-specific features of mammalian cell cultures for parameterization.

Notably, the widely-used *maximize biomass production* objective function performed poorly in describing mammalian metabolism. For microbial metabolism, this formulation is supported by theoretical and experimental evidence[84], and has been widely predictive for many experimental conditions and different models[39,73]. Accordingly, several recent studies using mammalian metabolic models (e.g., CHO cells, cancer cells, stem cells, immune cells) adopted this assumption. In the present study, however, *maximize biomass production* consistently overestimated growth rates and intracellular metabolic activity. Alternatively, five other presented formulations assumed that mammalian cells limit biomass production to maintain various homeostatic conditions. These homeostasis-limited assumptions performed markedly better than *maximize biomass production* by hypothesizing a restrained CHO metabolism. Objective functions considering cytosolic redox homeostasis and enzyme capacity predicted metabolic configurations that agreed especially well with observed growth rates and exometabolomic measurements. Future studies can rigorously refine and validate these formulations by comparing predicted intracellular flux distributions with experimental flux measurements, following the well-established footsteps of *Escherichia coli* models[38,39].

Conclusion

Proper parameter selection is essential for metabolic network models to provide accurate descriptions of cellular metabolism. Above all, the objective function parameter is highly influential in predicting metabolic and bioprocess performance phenotype. For this purpose, mammalian cells may be described as limiting their metabolic activities to maintain homeostasis. Various objective functions predicted different growth rates and metabolic pathway usage, leaving room for further validation and refinements. In addition, parameters describing biomass and time-course metabolic shifts also improved model predictions, especially when set to experimentally measured values. These results will guide future efforts to develop and improve models for mammalian cell metabolism, with applications ranging from biotherapeutic production[18,85] to unraveling the metabolic basis of diverse diseases[86–88].

Methods

Cell culture experiments

Two production fed batch processes were used, Fed batch 1 and Fed batch 2. Both fed batch processes used chemically defined media and feeds over the 12-day cell culture. Fed batch 1 used a glucose restricted fed batch process called HiPDOG[65]. Glucose concentration is kept low during the initial phase of the process, Day 2-7, through intermittent addition of feed medium containing glucose at the high end of pH dead-band and then glucose was maintained above 1.5 g/L thereafter. These conditions help restrict lactate production in fed batch cultures without compromising the proliferative capability of cells. In Fed batch 2 a conventional cell culture process was used where glucose was maintained above 1.5 g/L throughout the process.

For both process conditions, bioreactor vessels were inoculated at 2×10^6 viable cells/mL. The following bioprocess characteristics were quantified daily using a NOVA Flex BioProfile Analyzer (Nova Biomedical, Waltham, MA): viable cell density, average live cell diameter and concentrations of glucose, lactate, glutamate, and glutamine. Viable cell density (VCD) data was converted to growth rates by following equation to be compared to model-predicted growth rates.

$$Growth\ rate = \frac{1}{vcd} \cdot \frac{\Delta vcd}{\Delta time} = \frac{1}{vcd_0} \cdot \frac{vcd_{+1} - vcd_{-1}}{time_{+1} - time_{-1}}$$ (1)

In addition, cell cultures were sampled on specific days (Day 0, 3, 5, 7, 10 and 12) for cell

pellets and supernatant for transcriptomic (RNA-Seq), metabolomics analyses, and titer

measurements. Titers were analyzed using a protein A HPLC (model 1100 HPLC, Agilent

Technologies, Inc., Santa Clara, CA, protein A column model 2-1001-00, Applied Biosystems,

Foster City, CA). The RNA libraries were mapped to the CHO genome[89,90] using STAR aligner[91]

and processed to quantify gene expression counts with HTSeq-count[92].

Flash-frozen cell pellets (10E6 cells) and supernatant (1 mL) were collected from

bioreactor runs for clones A1 and A2 for each sampling day. Similarly, cell pellet and

supernatant samples were collected from bioreactor runs for clones B1 and B2 for days 7, 10 and

12. Collected samples were sent to Metabolon (Metabolon Inc, Morrisville, NC) for

metabolomics analyses. Proteins were removed by methanol precipitation and the metabolites

were recovered by vigorous shaking and centrifugation. The extracted samples were run for

reverse-phase Ultrahigh Performance Liquid Chromatography-Tandem Mass Spectroscopy with

negative ion mode ESI. Raw data was extracted, peak-identified and processed for quality

control using Metabolon's hardware and software. The raw ion count data was normalized

against the extracted proteins quantified using a Bradford assay.

Biomass measurement experiments

During cell culturing of clones Z1-Z4, 20 x 10^6 cells were sampled during days 0, 3, and 7.

At day 7, an additional 1 mL of culture was sampled. The samples were sub-divided and stored

at -80 °C for the following measurements.

Cellular dry weights were measured by weighing dehydrated cells, as previously

described[93]. Briefly, for each sample, an aluminum weigh boat was dried at 70 °C for 48 hours

and pre-weighed after being cooled down to room temperature. Stored samples (5×10^6 cells)

were thawed in ice, centrifuged, washed with PBS and centrifuged again. The cell pellets were

resuspended in deionized water and transferred to the pre-weighed aluminum boats and dried for

48 hours at 70° C. Then, the samples were cooled back to room temperature and weighed in

triplicates. The weighed mass was divided by cell count to calculate cellular dry weights.

Cellular protein contents were quantified by the Bicinchoninic Acid assay using a

commercial kit (Pierce™ BCA Protein Assay Kit; Thermo Fisher, Waltham, MA). Stored

samples (1×10^6 cells) were thawed in ice, centrifuged, and washed with PBS. Assay standards

were prepared as instructed for the range of 0 – 2000 µg/mL protein. The cells were disrupted by

a mixture of cell lysis agent (CelLytic™ M; Sigma Aldrich, St. Louis, MO) and protease

inhibitor cocktail (Sigma Aldrich, St. Louis, MO). A negative control was prepared from the

mixture without cell samples. The standards, samples and control were treated with assay

reagents, incubated and measured following kit instructions. The resulting standard curve was

confirmed to be linear ($R^2 = 0.988$). Measured absorbance values were converted to protein

concentrations following this standard curve and considering background absorbance described

by the negative control. Protein content in biomass was calculated to be 64.9% (±6.8) for days 3

and 7, agreeing well with previously published measurements[11,57].

The lipid contents of the cells were extracted using the Blight and Dyer protocol[94]. Briefly,

for each bioreactor run, 1 mL stored culture samples were centrifuged and re-suspended in water.

Chloroform and methanol were added sequentially in 1:2 ratio and mixed well by vortex. Then,

chloroform and water are sequentially added in 1:1 ratio and mixed well in between by vortex.

The resulting mixture was then centrifuged at 4000 RPM for 15 minutes at 20 °C, resulting in

phase separation. The upper phase and cell debris were carefully discarded. The lower

chloroform phase containing the lipids was carefully transferred to pre-weighed glass beakers.

Chloroform evaporated in a fume hood for 24 hours, after which the lipid samples were weighed.

The measured weights were divided by cell count and measured cellular dry weight, yielding a

cellular lipid content of 21.62% (±2.46), agreeing well with previously published measurements.

Constraint-based modeling and analysis

Flux balance analyses were conducted using the COBRA Toolbox 2.0[95] and the Gurobi

solver version 8.0.0 (Gurobi Optimization; Beaverton, Oregon) in MATLAB R2018b

(MathWorks; Natick, Massachusetts, USA).

Constructing a context-specific model

A community-consensus genome-scale model[11] was modified to produce a single cell line-

specific model for the cell lines from Pfizer by qualitatively interpreting transcriptomics data for

clone A1, following benchmarking results[36,61,62]. Specifically, transcript abundance data for each

gene was pre-processed by equation 2 to produce a binary gene score metric (ON: >5, OFF: 0).

The equation's threshold values were calculated from the mean transcript abundance across all

samples. Genes with transcript abundances of top and bottom 25 percentile were binarized as ON

and OFF, respectively. The binary gene scores were converted to reaction scores by considering

multimeric or isozyme relationships as described by the model's gene-protein-relationship matrix.

These binary reaction scores were used as input data for model extraction by the mCADRE

algorithm[63].

(2) $Gene\ Score = 5 * \log_2(1 + \frac{transcript\ abundance}{threshold})$

(3) $Reaction\ Score = \text{GPR}\ (\ Gene\ Score_1 \dots Gene\ Score_n)$

The resulting draft model was then manually curated to be consistent with experimental

data and established cell biology. Given the defined media of Pfizer's bioprocess, we allowed

extracellular transport only for metabolites with experimental basis – e.g. glucose, lactate,

proteinogenic amino acids, oxygen, carbon dioxide etc. We confirmed amino acid essentialities

and CHO-specific auxotrophies for arginine, cysteine and proline. We also ensured the inclusion

of synthesis pathways of non-essential amino acids and degradation pathways for all amino acids.

We confirmed the model utilized glucose via central carbon metabolism reactions, and removed

unlikely reactions – e.g. the methylglyoxal pathway or cross-membrane transport of metabolites

not found in media formulation. We ensured cytosolic and mitochondrial compartmentalization

of redox cofactors such as NAD/H and NADP/H. The final model consisted of 2375 irreversible

reactions, 587 metabolites and 1043 genes (Table S1, S2).

Model implementation of parameters

Here we detail how the nine parameters were implemented in the model. The workflow and

decision-making described here are also visualized as a flowchart (Fig. S13). The parameter

value of *biomass weight* (i.e. cellular dry weight) was used to convert the units of experimental measurements from [pg·cell^{-1}·day^{-1}] to [mmol·g$_{DW}$$^{-1}$·hr^{-1}]. When *considering dynamic dry weight (I)* for data input, the dry weight values were estimated from cell diameter measurements by assuming a constant cellular density and geometry[96]. When *considering dynamic dry weight (II)* for growth rate calculations, the following equation was used.

$$(4) \qquad \text{Adjusted Growth Rate} = \text{Growth Rate} * \frac{Future\ dry\ weight}{Current\ dry\ weight}$$

The *biomass composition* parameter was adjusted by altering the stoichiometric formulation of the biomass production reaction. Specifically, macromolecular protein content and lipid content were inversely varied, while assuming content of nucleotides and carbohydrates were stable. This was because protein and lipid contents were observed to vary the most[57]. Molecular makeup of amino acids and specific lipid molecules was unchanged from the genome-scale model, which was based on approximations from hybridoma cells. This was because variations in molecular composition did not affect model predictions. The *cell death rate* was implemented by the following equation for timepoints between days 7-11, in accordance to experimental observations.

$$(5) \qquad \text{Adjusted Growth Rate} = \text{Growth Rate} - Death\ Rate$$

Consideration of secretion costs was implemented by joining a model of the CHO secretory pathway[68] onto our metabolic network model. The secretory pathway model included about 100 reactions for folding, transport, post-translational modification and related activities in the endoplasmic reticulum and Golgi body (Table S3). We approximated the structure, folding requirements and glycosylation needs of Pfizer's monoclonal antibodies by that of Rituximab. *Biomass turnover rate* was implemented a single ATP hydrolysis reaction.

Consider byproduct synthesis was implemented for the following byproduct molecules: glycerol, formic acid, 2-hydroxybutyurate, isovalerate, acetic acid, hydroxyphenylpyruvate, hydroxyphenyl-lactate, phenyl-lactate, homocysteine, indole 3-lactate, citrate, and malate. These notably include products of amino acid catabolism as well as citric acid cycle intermediates.

Glycerol was assumed to be synthesized from dihydroxyacetone phosphate via glycerol-3-phosphate dehydrogenase and glycerol-3-phosphate phosphatase. Formic acid was assumed to be

synthesized in the folate cycle during tetrahydrofolate synthesis from 10-formyltetrahydrofolate.

The synthesis of isovalerate, 2-hydroxybutyurate, acetic acid, hydroxyphenylpyruvate,

hydroxyphenyl-lactate, phenyl-lactate, homocysteine, and indole 3-lactate were formulated as

catabolic byproducts, as previously published[69]. The model already included synthesis reactions

for homocysteine, acetic acid, citrate and malate. Synthesis and transport reactions were enabled

for all byproducts (Table S4).

Formulation of the objective function

The most widely used objective function is the maximization of biomass production, which

formulates the model as a linear programming optimization problem around the biomass

production reaction[97]. Alternatively, we formulated the objective function as a two-step linear

programming optimization to reflect cellular limitations to biomass production. First, we

enumerated possible limitations to biomass production, as discussed above. Second, we

annotated model reactions for 'penalty' metrics describing such limitations (Table S5, S6). We

derived penalty values for Gibbs dissipation[76] and enzyme costs[98] from previous works. Third,

the model was bounded by experimentally measured nutrient consumption rates (Table S8).

Fourth, for a penalty of choice, we calculated the minimal amount of penalty to process the

consumed nutrients. This calculated value was then used as an additional boundary condition.

Fifth, given all these bounds, we calculated the maximum possible amount of biomass

production. This predicted value, then, describes the maximum anabolic activity given nutrient

consumption behavior and theoretical cellular limitations.

This presented workflow builds upon previous work that minimizes network flux[99] and

parsimonious flux balance analysis[78] and the 'max biomass per unit flux' objective[38] with

important differences. Previous methods implicitly assumed that biomass production was gated

by substrate availability and therefore first maximized biomass production and then minimized

total unit flux as a penalty metric. Here, we assume cellular limitations restrict growth rather than

substrate availability, and therefore inverse the order of optimization. Also, while previous

methods penalized reactions uniformly, we apply weighted penalties following biological

assumptions. For example, reactions that are catalyzed by complex multimers are penalized

correspondingly more severely. Like previous methods, this workflow requires the model to be
irreversible.

Estimation of parameter impact on model accuracy

4000 parameter settings were generated by randomly varying parameters to within a range
based on literature or experimental measurements (Table 1, Table S7). For each setting,
parameter values were implemented to yield a flux balance analysis prediction. The prediction
used as input data the following experimental measurements: specific productivity and the
consumption rates of glucose, lactate and 20 proteinogenic amino acids (Table S8). The
prediction was repeated 80 times by using experimental data from 10 clones across 8 days. The
resulting 80 model-predicted growth rates were compared to their respective experimental values
to yield prediction errors (Table S7). To facilitate human interpretation, the prediction errors
were first transformed by the negative log function. Then, the transformed values were
normalized so that mean metric values for all 80 predictions laid between 0 and 1. The resulting
metric was called 'accuracy'.

A regression analysis between accuracy and parameter values were performed, using
Python 3.7.3 in the Jupyter Notebook environment. For this analysis, parameter values were
normalized to be between 0 and 1 by referencing minimum and maximum possible values (Table
1). A linear regression analysis was performed according to equation 6, where B_i and X_i
represent respectively the effect size and normalized parameter values of parameter *i*. The
resulting effect size of parameters and their p-values were used to identify parameter relevance.

(6) $\text{Accuracy} = B_0 + \sum_{i=1}^{9} B_i \cdot X_i$

Flux sampling analysis

We computed distributions of likely fluxes for each reaction in the model by stochastically
sampling 5000 points within the solution space via the Markov chain Monte Carlo sampling via
artificially centered hit-and-run algorithm, as described previously[100]. First, the metabolic
network model was parameterized according to experimentally observed values. Then,

experimental measurements for all clones on day 4 were used to constrain model reactions for

biomass production, monoclonal antibody secretion and consumption of glucose, lactate and

proteinogenic amino acids. A set of non-uniform 'points' or flux values was generated within the

feasible flux space. Each point was subsequently moved randomly, while remaining within the

feasible flux space. To do this, a random direction was first chosen. Second, the limit for how far

the point can travel in the randomly-chosen direction was calculated. Lastly, a new random point

on this line was selected. This process was iterated until the set of points approached a uniform

sample of the solution space. Thereafter, the sampled fluxes were normalized by total model

flux[101]. The normalized values were analyzed to explore in detail the impact of objective

functions on model predictions. These analyses were done using Python 3.7.3 in the Jupyter

Notebook environment.

References

1. Smiatek J, Jung A, Bluhmki E. Towards a Digital Bioprocess Replica: Computational Approaches in Biopharmaceutical Development and Manufacturing. *Trends Biotechnol*. Published online June 6, 2020. doi:10.1016/j.tibtech.2020.05.008

2. Abt V, Barz T, Cruz-Bournazou MN, et al. Model-based tools for optimal experiments in bioprocess engineering. *Curr Opin Chem Eng*. 2018;22:244-252. doi:10.1016/j.coche.2018.11.007

3. Lewis AM, Abu☐Absi NR, Borys MC, Li ZJ. The use of 'Omics technology to rationally improve industrial mammalian cell line performance. *Biotechnol Bioeng*. 2016;113(1):26-38. doi:10.1002/bit.25673

4. Richelle A, Lewis NE. Improvements in protein production in mammalian cells from targeted metabolic engineering. *Curr Opin Syst Biol*. 2017;6:1-6. doi:10.1016/j.coisb.2017.05.019

5. Mulukutla BC, Mitchell J, Geoffroy P, et al. Metabolic engineering of Chinese hamster ovary cells towards reduced biosynthesis and accumulation of novel growth inhibitors in fed-batch cultures. *Metab Eng*. 2019;54:54-68. doi:10.1016/j.ymben.2019.03.001

6. Wang Q, Yin B, Chung C-Y, Betenbaugh MJ. Glycoengineering of CHO Cells to Improve Product Quality. *Methods Mol Biol Clifton NJ*. 2017;1603:25-44. doi:10.1007/978-1-4939-6972-2_2

7. Baik JY, Gasimli L, Yang B, et al. Metabolic engineering of Chinese hamster ovary cells: towards a bioengineered heparin. *Metab Eng*. 2012;14(2):81-90. doi:10.1016/j.ymben.2012.01.008

8. Gupta SK, Srivastava SK, Sharma A, et al. Metabolic engineering of CHO cells for the development of a robust protein production platform. *PLOS ONE*. 2017;12(8):e0181455. doi:10.1371/journal.pone.0181455

9. Golabgir A, Gutierrez JM, Hefzi H, et al. Quantitative feature extraction from the Chinese hamster ovary bioprocess bibliome using a novel meta-analysis workflow. *Biotechnol Adv*. 2016;34(5):621-633. doi:10.1016/j.biotechadv.2016.02.011

10. Sumit M, Dolatshahi S, Chu A-HA, et al. Dissecting N-Glycosylation Dynamics in Chinese Hamster Ovary Cells Fed-batch Cultures using Time Course Omics Analyses. *iScience*. 2019;12:102-120. doi:10.1016/j.isci.2019.01.006

11. Hefzi H, Ang KS, Hanscho M, et al. A Consensus Genome-scale Reconstruction of Chinese Hamster Ovary Cell Metabolism. *Cell Syst*. 2016;3(5):434-443.e8. doi:10.1016/j.cels.2016.10.020

12. Calmels C, McCann A, Malphettes L, Andersen MR. Application of a curated genome-scale metabolic model of CHO DG44 to an industrial fed-batch process. *Metab Eng*. 2019;51:9-19. doi:10.1016/j.ymben.2018.09.009

13. Popp O, Müller D, Didzus K, et al. A hybrid approach identifies metabolic signatures of high-producers for chinese hamster ovary clone selection and process optimization. *Biotechnol Bioeng*. 2016;113(9):2005-2019. doi:10.1002/bit.25958

14. Chen Y, McConnell BO, Gayatri Dhara V, et al. An unconventional uptake rate objective function approach enhances applicability of genome-scale models for mammalian cells. *NPJ Syst Biol Appl*. 2019;5. doi:10.1038/s41540-019-0103-6

15. Fouladiha H, Marashi S-A, Torkashvand F, Mahboudi F, Lewis NE, Vaziri B. A metabolic network-based approach for developing feeding strategies for CHO cells to increase monoclonal antibody production. *Bioprocess Biosyst Eng*. Published online March 24, 2020. doi:10.1007/s00449-020-02332-6

16. Zhuangrong H, Seongkyu Y. Identifying metabolic features and engineering targets for productivity improvement in CHO cells by integrated transcriptomics and genome-scale metabolic model. *Biochem Eng J*. Published online May 5, 2020:107624. doi:10.1016/j.bej.2020.107624

17. Huang Z, Yoon S. Integration of Time-Series Transcriptomic Data with Genome-Scale CHO Metabolic Models for mAb Engineering. *Processes*. 2020;8(3):331. doi:10.3390/pr8030331

18. Huang Z, Lee D-Y, Yoon S. Quantitative intracellular flux modeling and applications in biotherapeutic development and production using CHO cell cultures. *Biotechnol Bioeng*. 2017;114(12):2717-2728. doi:10.1002/bit.26384

19. Lewis NE, Nagarajan H, Palsson BO. Constraining the metabolic genotype-phenotype relationship using a phylogeny of in silico methods. *Nat Rev Microbiol*. 2012;10(4):291-305. doi:10.1038/nrmicro2737

20. Heirendt L, Arreckx S, Pfau T, et al. Creation and analysis of biochemical constraint-based models using the COBRA Toolbox v.3.0. *Nat Protoc*. 2019;14(3):639-702. doi:10.1038/s41596-018-0098-2

21. Bordbar A, Monk JM, King ZA, Palsson BO. Constraint-based models predict metabolic and associated cellular functions. *Nat Rev Genet*. 2014;15(2):107-120. doi:10.1038/nrg3643

22. Sastry AV, Gao Y, Szubin R, et al. The Escherichia coli transcriptome mostly consists of independently regulated modules. *Nat Commun*. 2019;10(1):5536. doi:10.1038/s41467-019-13483-w

23. Yizhak K, Chaneton B, Gottlieb E, Ruppin E. Modeling cancer metabolism on a genome scale. *Mol Syst Biol*. 2015;11(6):817. doi:10.15252/msb.20145307

24. Mardinoglu A, Agren R, Kampf C, Asplund A, Uhlen M, Nielsen J. Genome-scale metabolic modelling of hepatocytes reveals serine deficiency in patients with non-alcoholic fatty liver disease. *Nat Commun*. 2014;5:3083. doi:10.1038/ncomms4083

25. Lewis NE, Abdel-Haleem AM. The evolution of genome-scale models of cancer metabolism. *Front Physiol*. 2013;4:237. doi:10.3389/fphys.2013.00237

26. Magnúsdóttir S, Thiele I. Modeling metabolism of the human gut microbiome. *Curr Opin Biotechnol*. 2018;51:90-96. doi:10.1016/j.copbio.2017.12.005

27. Sen P, Orešič M. Metabolic Modeling of Human Gut Microbiota on a Genome Scale: An Overview. *Metabolites*. 2019;9(2). doi:10.3390/metabo9020022

28. Zhang C, Hua Q. Applications of Genome-Scale Metabolic Models in Biotechnology and Systems Medicine. *Front Physiol*. 2016;6. doi:10.3389/fphys.2015.00413

29. Kim HU, Lee SY. Applications of genome-scale metabolic network models in the biopharmaceutical industry. *Pharm Bioprocess*. 2013;1(4):337-339. doi:10.4155/pbp.13.37

30. Sommeregger W, Sissolak B, Kandra K, von Stosch M, Mayer M, Striedner G. Quality by control: Towards model predictive control of mammalian cell culture bioprocesses. *Biotechnol J*. 2017;12(7). doi:10.1002/biot.201600546

31. Varma A, Palsson BO. Stoichiometric flux balance models quantitatively predict growth and metabolic by-product secretion in wild-type Escherichia coli W3110. *Appl Environ Microbiol*. 1994;60(10):3724-3731.

32. Aurich MK, Fleming RMT, Thiele I. A systems approach reveals distinct metabolic strategies among the NCI-60 cancer cell lines. *PLOS Comput Biol*. 2017;13(8):e1005698. doi:10.1371/journal.pcbi.1005698

33. Traustason B. Amino Acid Requirements of the Chinese Hamster Ovary Cell Metabolism during Recombinant Protein Production. *bioRxiv*. Published online October 7, 2019:796490. doi:10.1101/796490

34. Bonarius HPJ, Schmid G, Tramper J. Flux analysis of underdetermined metabolic networks: the quest for the missing constraints. *Trends Biotechnol*. 1997;15(8):308-314. doi:10.1016/S0167-7799(97)01067-6

35. Blazier AS, Papin JA. Integration of expression data in genome-scale metabolic network reconstructions. *Front Physiol*. 2012;3. doi:10.3389/fphys.2012.00299

36. Opdam S, Richelle A, Kellman B, Li S, Zielinski DC, Lewis NE. A Systematic Evaluation of Methods for Tailoring Genome-Scale Metabolic Models. *Cell Syst*. 2017;4(3):318-329.e6. doi:10.1016/j.cels.2017.01.010

37. Feist AM, Palsson BO. The biomass objective function. *Curr Opin Microbiol*. 2010;13(3):344-349. doi:10.1016/j.mib.2010.03.003

38. Schuetz R, Kuepfer L, Sauer U. Systematic evaluation of objective functions for predicting intracellular fluxes in Escherichia coli. *Mol Syst Biol*. 2007;3:119. doi:10.1038/msb4100162

39. Costa RS, Nguyen S, Hartmann A, Vinga S. Exploring the Cellular Objective in Flux Balance Constraint-Based Models. In: Mendes P, Dada JO, Smallbone K, eds. *Computational Methods in Systems Biology*. Lecture Notes in Computer Science. Springer International Publishing; 2014:211-224. doi:10.1007/978-3-319-12982-2_15

40. Dikicioglu D, Kırdar B, Oliver SG. Biomass composition: the "elephant in the room" of metabolic modelling. *Metabolomics*. 2015;11(6):1690-1701. doi:10.1007/s11306-015-0819-2

41. Lakshmanan M, Long S, Ang KS, Lewis N, Lee D-Y. On the impact of biomass composition in constraint-based flux analysis. *bioRxiv*. Published online May 28, 2019:652040. doi:10.1101/652040

42. Dikicioglu D, Oliver SG. Extension of the yeast metabolic model to include iron metabolism and its use to estimate global levels of iron-recruiting enzyme abundance from cofactor requirements. *Biotechnol Bioeng*. 2019;116(3):610-621. doi:10.1002/bit.26905

43. Monk JM, Lloyd CJ, Brunk E, et al. iML1515, a knowledgebase that computes Escherichia coli traits. *Nat Biotechnol*. 2017;35(10):904-908. doi:10.1038/nbt.3956

44. García Sánchez CE, Torres Sáez RG. Comparison and analysis of objective functions in flux balance analysis. *Biotechnol Prog*. 2014;30(5):985-991. doi:10.1002/btpr.1949

45. Zhao Q, Stettner AI, Reznik E, Paschalidis ICh, Segrè D. Mapping the landscape of metabolic goals of a cell. *Genome Biol*. 2016;17(1):109. doi:10.1186/s13059-016-0968-2

46. Sánchez BJ, Zhang C, Nilsson A, Lahtvee P-J, Kerkhoven EJ, Nielsen J. Improving the phenotype predictions of a yeast genome-scale metabolic model by incorporating enzymatic constraints. *Mol Syst Biol*. 2017;13(8):935. doi:10.15252/msb.20167411

47. Edwards JS, Palsson BO. The Escherichia coli MG1655 in silico metabolic genotype: its definition, characteristics, and capabilities. *Proc Natl Acad Sci U S A*. 2000;97(10):5528-5533. doi:10.1073/pnas.97.10.5528

48. Orth JD, Conrad TM, Na J, et al. A comprehensive genome-scale reconstruction of Escherichia coli metabolism--2011. *Mol Syst Biol*. 2011;7:535. doi:10.1038/msb.2011.65

49. Förster J, Famili I, Fu P, Palsson BØ, Nielsen J. Genome-scale reconstruction of the Saccharomyces cerevisiae metabolic network. *Genome Res*. 2003;13(2):244-253. doi:10.1101/gr.234503

50. Heavner BD, Smallbone K, Price ND, Walker LP. Version 6 of the consensus yeast metabolic network refines biochemical coverage and improves model performance. *Database*. 2013;2013. doi:10.1093/database/bat059

51. Gu C, Kim GB, Kim WJ, Kim HU, Lee SY. Current status and applications of genome-scale metabolic models. *Genome Biol*. 2019;20(1):121. doi:10.1186/s13059-019-1730-3

52. Yilmaz S, Walhout A. Metabolic network modeling with model organisms. *Curr Opin Chem Biol*. 2017;36:32-39.

53. Sigurdsson MI, Jamshidi N, Steingrimsson E, Thiele I, Palsson BØ. A detailed genome-wide reconstruction of mouse metabolism based on human Recon 1. *BMC Syst Biol*. 2010;4:140. doi:10.1186/1752-0509-4-140

54. Witting M, Hastings J, Rodriguez N, et al. Modeling Meets Metabolomics-The WormJam Consensus Model as Basis for Metabolic Studies in the Model Organism Caenorhabditis elegans. *Front Mol Biosci*. 2018;5:96. doi:10.3389/fmolb.2018.00096

55. Brunk E, Sahoo S, Zielinski DC, et al. Recon3D enables a three-dimensional view of gene variation in human metabolism. *Nat Biotechnol*. 2018;36(3):272-281. doi:10.1038/nbt.4072

56. Gerdtzen ZP. Modeling Metabolic Networks for Mammalian Cell Systems: General Considerations, Modeling Strategies, and Available Tools. *Genomics Syst Biol Mamm Cell Cult*. Published online 2011:71-108. doi:10.1007/10_2011_120

57. Szeliova D, Ruckerbauer D, Galleguillos S, et al. What CHO is made of: Variations in the biomass composition of Chinese hamster ovary cell lines. *Metab Eng*. Published online 2020.

58. Savinell JM, Palsson BO. Network analysis of intermediary metabolism using linear optimization. I. Development of mathematical formalism. *J Theor Biol*. 1992;154(4):421-454. doi:10.1016/s0022-5193(05)80161-4

59. Ahn WS, Antoniewicz MR. Metabolic flux analysis of CHO cells at growth and non-growth phases using isotopic tracers and mass spectrometry. *Metab Eng*. 2011;13(5):598-609. doi:10.1016/j.ymben.2011.07.002

60. Dean J, Reddy P. Metabolic analysis of antibody producing CHO cells in fed-batch production. *Biotechnol Bioeng*. 2013;110(6):1735-1747. doi:10.1002/bit.24826

61. Richelle A, Chiang AWT, Kuo C-C, Lewis NE. Increasing consensus of context-specific metabolic models by integrating data-inferred cell functions. *PLOS Comput Biol*. 2019;15(4):e1006867. doi:10.1371/journal.pcbi.1006867

62. Richelle A, Joshi C, Lewis NE. Assessing key decisions for transcriptomic data integration in biochemical networks. *PLOS Comput Biol*. 2019;15(7):e1007185. doi:10.1371/journal.pcbi.1007185

63. Wang Y, Eddy JA, Price ND. Reconstruction of genome-scale metabolic models for 126 human tissues using mCADRE. *BMC Syst Biol*. 2012;6(1):153. doi:10.1186/1752-0509-6-153

64. Chan SHJ, Cai J, Wang L, Simons-Senftle MN, Maranas CD. Standardizing biomass reactions and ensuring complete mass balance in genome-scale metabolic models. *Bioinformatics*. 2017;33(22):3603-3609. doi:10.1093/bioinformatics/btx453

65. Pan X, Dalm C, Wijffels RH, Martens DE. Metabolic characterization of a CHO cell size increase phase in fed-batch cultures. *Appl Microbiol Biotechnol*. 2017;101(22):8101-8113. doi:10.1007/s00253-017-8531-y

66. Neunstoecklin B, Stettler M, Solacroup T, Broly H, Morbidelli M, Soos M. Determination of the maximum operating range of hydrodynamic stress in mammalian cell culture. *J Biotechnol*. 2015;194:100-109. doi:10.1016/j.jbiotec.2014.12.003

67. Tanzeglock T, Soos M, Stephanopoulos G, Morbidelli M. Induction of mammalian cell death by simple shear and extensional flows. *Biotechnol Bioeng*. 2009;104(2):360-370. doi:10.1002/bit.22405

68. Gutierrez JM, Feizi A, Li S, et al. Genome-scale reconstructions of the mammalian secretory pathway predict metabolic costs and limitations of protein secretion. *Nat Commun*. 2020;11(1):1-10. doi:10.1038/s41467-019-13867-y

69. Mulukutla BC, Kale J, Kalomeris T, Jacobs M, Hiller GW. Identification and control of novel growth inhibitors in fed-batch cultures of Chinese hamster ovary cells. *Biotechnol Bioeng*. 2017;114(8):1779-1790. doi:10.1002/bit.26313

70. Gunn JM, Brancheau MR. Protein turnover, growth and proliferation in CHO cells. Variation within and between mutant classes for salvage pathway enzymes. *Biochem J*. 1992;282 (Pt 1):49-57. doi:10.1042/bj2820049

71. Ramakrishna R, Edwards JS, McCulloch A, Palsson BO. Flux-balance analysis of mitochondrial energy metabolism: consequences of systemic stoichiometric constraints. *Am J Physiol Regul Integr Comp Physiol*. 2001;280(3):R695-704. doi:10.1152/ajpregu.2001.280.3.R695

72. Knorr AL, Jain R, Srivastava R. Bayesian-based selection of metabolic objective functions. *Bioinforma Oxf Engl*. 2007;23(3):351-357. doi:10.1093/bioinformatics/btl619

73. Sánchez CEG, García CAV, Sáez RGT. Predictive Potential of Flux Balance Analysis of Saccharomyces cerevisiae Using as Optimization Function Combinations of Cell Compartmental Objectives. *PLOS ONE*. 2012;7(8):e43006. doi:10.1371/journal.pone.0043006

74. Noor E, Flamholz A, Bar-Even A, Davidi D, Milo R, Liebermeister W. The Protein Cost of Metabolic Fluxes: Prediction from Enzymatic Rate Laws and Cost Minimization. *PLOS Comput Biol*. 2016;12(11):e1005167. doi:10.1371/journal.pcbi.1005167

75. Yeo HC, Hong J, Lakshmanan M, Lee D-Y. Enzyme capacity-based genome scale modelling of CHO cells. *Metab Eng*. 2020;60:138-147. doi:10.1016/j.ymben.2020.04.005

76. Niebel B, Leupold S, Heinemann M. An upper limit on Gibbs energy dissipation governs cellular metabolism. *Nat Metab*. 2019;1(1):125-132. doi:10.1038/s42255-018-0006-7

77. Klumpp S, Scott M, Pedersen S, Hwa T. Molecular crowding limits translation and cell growth. *Proc Natl Acad Sci*. 2013;110(42):16754-16759. doi:10.1073/pnas.1310377110

78. Lewis NE, Hixson KK, Conrad TM, et al. Omic data from evolved E. coli are consistent with computed optimal growth from genome-scale models. *Mol Syst Biol*. 2010;6:390. doi:10.1038/msb.2010.47

79. Ray PD, Huang B-W, Tsuji Y. Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cell Signal*. 2012;24(5):981-990. doi:10.1016/j.cellsig.2012.01.008

80. Nicolae A, Wahrheit J, Bahnemann J, Zeng A-P, Heinzle E. Non-stationary 13C metabolic flux analysis of Chinese hamster ovary cells in batch culture using extracellular labeling highlights metabolic reversibility and compartmentation. *BMC Syst Biol*. 2014;8(1):50. doi:10.1186/1752-0509-8-50

81. Sellick CA, Croxford AS, Maqsood AR, et al. Metabolite profiling of recombinant CHO cells: designing tailored feeding regimes that enhance recombinant antibody production. *Biotechnol Bioeng*. 2011;108(12):3025-3031. doi:10.1002/bit.23269

82. Chong WPK, Reddy SG, Yusufi FNK, et al. Metabolomics-driven approach for the improvement of Chinese hamster ovary cell growth: Overexpression of malate dehydrogenase II. *J Biotechnol*. 2010;147(2):116-121. doi:10.1016/j.jbiotec.2010.03.018

83. O'Brien CM, Mulukutla BC, Mashek DG, Hu W-S. Regulation of Metabolic Homeostasis in Cell Culture Bioprocesses. *Trends Biotechnol*. 2020;0(0). doi:10.1016/j.tibtech.2020.02.005

84. Towbin BD, Korem Y, Bren A, Doron S, Sorek R, Alon U. Optimality and sub-optimality in a bacterial growth law. *Nat Commun*. 2017;8. doi:10.1038/ncomms14123

85. Gutierrez JM, Lewis NE. Optimizing eukaryotic cell hosts for protein production through systems biotechnology and genome-scale modeling. *Biotechnol J*. 2015;10(7):939-949. doi:10.1002/biot.201400647

86. Bordbar A, Palsson BO. Using the reconstructed genome-scale human metabolic network to study physiology and pathology. *J Intern Med*. 2012;271(2):131-141. doi:10.1111/j.1365-2796.2011.02494.x

87. Chen Y, Li G, Nielsen J. Genome-Scale Metabolic Modeling from Yeast to Human Cell Models of Complex Diseases: Latest Advances and Challenges. *Methods Mol Biol Clifton NJ*. 2019;2049:329-345. doi:10.1007/978-1-4939-9736-7_19

88. Geng J, Nielsen J. In silico analysis of human metabolism: Reconstruction, contextualization and application of genome-scale models. *Curr Opin Syst Biol*. 2017;2:29-38. doi:10.1016/j.coisb.2017.01.001

89. Lewis NE, Liu X, Li Y, et al. Genomic landscapes of Chinese hamster ovary cell lines as revealed by the Cricetulus griseus draft genome. *Nat Biotechnol*. 2013;31(8):759-765. doi:10.1038/nbt.2624

90. Li S, Cha SW, Heffner K, et al. Proteogenomic Annotation of Chinese Hamsters Reveals Extensive Novel Translation Events and Endogenous Retroviral Elements. *J Proteome Res*. 2019;18(6):2433-2445. doi:10.1021/acs.jproteome.8b00935

91. Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 2013;29(1):15-21. doi:10.1093/bioinformatics/bts635

92. Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics*. 2015;31(2):166-169. doi:10.1093/bioinformatics/btu638

93. Edros RZ, McDonnell S, Al-Rubeai M. Using molecular markers to characterize productivity in Chinese hamster ovary cell lines. *PloS One*. 2013;8(10):e75935. doi:10.1371/journal.pone.0075935

94. Beck AE, Hunt KA, Carlson RP. Measuring Cellular Biomass Composition for Computational Biology Applications. *Processes*. 2018;6(5):38. doi:10.3390/pr6050038

95. Schellenberger J, Que R, Fleming RMT, et al. Quantitative prediction of cellular metabolism with constraint-based models: the COBRA Toolbox v2.0. *Nat Protoc*. 2011;6(9):1290-1307. doi:10.1038/nprot.2011.308

96. Milo R, Jorgensen P, Moran U, Weber G, Springer M. BioNumbers--the database of key numbers in molecular and cell biology. *Nucleic Acids Res*. 2010;38(Database issue):D750-753. doi:10.1093/nar/gkp889

97. Orth JD, Thiele I, Palsson BØ. What is flux balance analysis? *Nat Biotechnol*. 2010;28(3):245-248. doi:10.1038/nbt.1614

98. Consortium TU. UniProt: a hub for protein information. *Nucleic Acids Res*. 2015;43(D1):D204-D212. doi:10.1093/nar/gku989

99. Klitgord N, Segrè D. The importance of compartmentalization in metabolic flux models: yeast as an ecosystem of organelles. *Genome Inform Int Conf Genome Inform*. 2010;22:41-55.

100. Nam H, Lewis NE, Lerman JA, et al. Network Context and Selection in the Evolution to Enzyme Specificity. *Science*. 2012;337(6098):1101-1104. doi:10.1126/science.1216861

101. Lewis NE, Schramm G, Bordbar A, et al. Large-scale in silico modeling of metabolic interactions between cell types in the human brain. *Nat Biotechnol*. 2010;28(12):1279-1285. doi:10.1038/nbt.1711

1 *Figures*

Figure 1: 10 CHO clones with diverse metabolic phenotypes were studied – The examined clones expressed different monoclonal antibodies (1: ○, 2: △, 3: □) and were subjected to different bioprocess conditions (Fed batch #1: empty; Fed batch #2: filled). Due to these differences, the cells exhibited diverse metabolic phenotypes: (a) the cells consumed key nutrients such as glucose at varying amounts, and variously consumed or secreted lactate, (b) resulting in distinct growth and productivity performances. (c) These clones were observed between culture days 4 and 11, during which the cells traversed exponential and stationary phases (highlighted).

Figure 2: Analysis of model parameters, workflow scheme & results – (a) 4000 parameter settings were evaluated against 80
datapoints to produce thousands of model predictions. (b) Here, we provide an example model prediction of growth rate for 80
points. These particular predictions were made by assuming parameter values equal to experimental measurements – that is, a
baseline biomass dry weight of 280 picograms/cell at day 3, and 65% protein composition of biomass; biomass weight was also
varied according to time-course cell diameter measurements. Biomass production was assumed to be limited by enzyme capacity.
(c) Analysis results were examined by their 'accuracy', which was calculated and normalized from the negative-log of mean
prediction residuals. Predictions assuming experimental values, featured in panel (b), resulted in an average accuracy of 0.82
(blue line).

Objective function formulation

Figure 3: Key model parameters impact model accuracy – A regression model estimated the importance of parameters by their effect size. *Objective function formulation* was treated as six independent binary parameters representing the various formulations (italicized and highlighted in grey). Several of these formulations were highly correlated to prediction accuracy. Other important parameters were *cell death rate*, *biomass composition* and *consideration of dynamic biomass weight* (BW)*for growth rate calculations*. The remaining parameters had negligible effects, which are detailed elsewhere (Fig. S6)

Figure 4: Objective functions vary in hypothesized metabolic activities underlying the predicted growth rates – Objective functions helped determine the metabolic activities of key reactions in the central carbon metabolism. The heatmap displays the reactions' fractional contribution to total metabolic flux. These reactions include the first step of glycolysis catalyzed by hexokinase 1 (*A*); carbon shunting variously towards glycerol (*B*), folate cycle intermediates (*C*), malate secretion (*D*); mitochondrial respiration (*E*); anaplerosis of respiratory carbon dioxide via bicarbonate and pyruvate carboxylase (*F*). The flowchart on the right contextualizes the reactions within the central carbon metabolism.

Table 1: Parameters varied in this study – *the table provides names and descriptions of the parameters, along with range of values explored in our analysis; mean experimental values of the presented clones are given in parentheses when applicable.*

#	Parameter name & description	Parameter values
1	**Biomass weight:** a baseline biomass dry weight value at day 4.	210 – 400 pg/cell (280)
2	**Biomass composition:** as a percentage of protein in cellular biomass.	45 – 75% (65)
3	**Consider dynamic biomass weight (I)** when inputting experimental data as boundary conditions.	TRUE / FALSE
4	**Consider dynamic biomass weight (II)** to refine growth rate calculations (Fig. S1).	TRUE / FALSE
5	**Cell death rate:** a constant cell death rate presumably due to shear stress during decreased viability (Fig. S3).	0 – 0.005 hr^{-1}
6	**Consider secretion costs:** to estimated energy and material costs for monoclonal antibody secretion.	TRUE / FALSE
7	**Consider amino acid catabolism byproduct synthesis** as recently described[69].	TRUE / FALSE
8	**Biomass turnover rate:** a constant rate of energy expenditure for biomass maintenance.	0 – 0.01 mmol g^{-1} hr^{-1}
9	**Objective function formulation:** the assumption of a 'cellular objective', which renders the metabolic network as a linear programming optimization problem.	• max. biomass production • min. enzyme costs • min. Gibbs energy dissipation • min. ROS synthesis • min. cytosolic NADPH regeneration • min. mitochondrial NADH regeneration