**Coastal winds and larval fish abundance indicate a recruitment mechanism for southeast**

**Australian estuarine fisheries**

Hayden T. Schilling[1,2]*, Charles Hinchliffe[2], Jonathan P. Gillson[3], Anthony Miskiewicz[2,4], Iain M.

Suthers[1,2]

1. Sydney Institute of Marine Science, Chowder Bay Road, Mosman, 2088, Australia

2. Centre for Marine Science & Innovation, UNSW Australia, 2052 Australia

3. The Centre for Environment, Fisheries and Aquaculture Science, Pakefield Road, Lowestoft,

NR33 0HT, United Kingdom

4. Australian Museum, College St., Sydney 2000, Australia

*Corresponding author: h.schilling@unsw.edu.au

Running title: Coastal wind driven recruitment

**Abstract**

Coastal winds transport water masses and larval fish onshore or offshore which may influence

estuarine recruitment, yet our understanding of the mechanism underlying this relationship is

limited. Here, we combine datasets from a historical database of larval fish off southeast Australia

with a high-resolution atmospheric reanalysis model to show that normalised abundance of coastally

spawned larvae increased with weak to moderate upwelling favourable winds 14 days prior to

sampling. The increase in abundance may reflect increased nutrient and plankton availability for

larval fish. Normalised larval abundance decreased following strong upwelling favourable winds but

increased after onshore (downwelling favourable) winds, due to wind-driven transport. By

combining a commercial estuarine fisheries catch-rate dataset (4 species, 8 estuaries, 10 years) and

the high-resolution atmospheric reanalysis model, we show that negative effects of upwelling

favourable winds during the spawning period can be detected in lagged estuarine commercial

fisheries catch rates (lagged by 2 – 8 years depending on species' growth rates), potentially

representing the same mechanism proposed for larval fish. Upwelling favourable winds in the

southeast Australian region have increased since 1850 while onshore winds have decreased, which

may reduce larval recruitment to estuaries. Coastal winds are likely an important factor for estuarine

recruitment in the southeast Australian region and future research on the estuarine recruitment of

fish should incorporate coastal winds.

**Keywords:** larval fish, retention, wind driven upwelling, fisheries production, Harald Dannevig,

downwelling, estuarine recruitment

**Introduction**

Strong year classes of marine fish have long interested fisheries oceanographers, with many studies attempting to better understand the drivers of variability in population dynamics (Houde, 2008). Previous studies have linked strong year classes with abiotic and biotic factors during the larval period, with hypotheses either focusing on increased growth or reduced mortality resulting in a higher recruitment (Garvine *et al.*, 1997; Helbig and Pepin, 1998; Pepin, 2016). Some of the key factors influencing recruitment include the biomass of the spawning stock, location and timing of spawning, the vertical distribution of eggs and larvae and upwelling generated productivity. The spawning stock biomass (SSB) is the basis for recruitment success (Cury *et al.*, 2014) and in combination with fecundity, egg viability and sex ratio, determines the number of eggs and larvae which subsequent stochastic processes act upon to shape recruitment (Kell *et al.*, 2016).

Stochastic processes are often more influential than SSB due to the high variability in mortality and subsequent recruitment (Szuwalski *et al.*, 2015). The location and timing of spawning events is also important for overall recruitment, with seasonal and interannual variations in oceanography often shaping dispersal and potential recruitment (Ciannelli *et al.*, 2015; Schilling *et al.*, 2020). As larvae have limited swimming potential in their early stages (hence oceanographic driven dispersal), larvae which can control their vertical positioning often exploit vertical variation in cross-shelf flow on continental shelves to remain near favourable areas for juvenile habitat, potentially increasing recruitment (Hare and Govoni, 2005; Ospina-Alvarez *et al.*, 2018).

Productivity, which is often generated by upwelling (Everett *et al.*, 2014), is often suggested as a driver of faster growth in fish larvae through the intermediate production of phytoplankton and zooplankton thereby reducing mortality from predation as the larvae increase in swimming ability (Anderson, 1988; McFarlane and Beamish, 1992), and increasing recruitment (Bailey and Houde, 1989). On the other hand, upwelling can also have negative effects in some circumstances due to offshore transport away from suitable recruitment habitat. While the influence of upwelling has been clearly demonstrated in some regions including the southern Benguela upwelling region (Wilhelm *et al.*, 2005) and the Northeast Atlantic (Santos *et al.*, 2004), this process is likely region specific and remains uncertain in many regions, including the southwest Pacific.

The coastal waters of eastern Australia are dominated by the East Australian Current (EAC; Oke *et al.*, 2019). The EAC generally transports larvae poleward along a narrow continental shelf (<30km) to coastal and estuarine nursery habitats (Roughan *et al.*, 2011; Schilling *et al.*, 2020). Together with the EAC, onshore winds are an important driver of upwelling and downwelling

through Ekman transport mechanisms (Roughan and Middleton, 2002; Schaeffer *et al.*, 2013, 2014).

Wind is an important driver of cross-shelf flows in the area (McClean-Padman and Padman, 1991;

Middleton *et al.*, 1996). Winds from the northeast (NE) blow along the coastline, promoting offshore

Ekman transport and upwelling of cold nutrient-rich water along the coast, while winds from the

southeast (SE) blow onto the coast, promoting onshore transport and downwelling (Griffin and

Middleton, 1992; Middleton *et al.*, 1996).

It was first hypothesised that onshore winds may be affecting nearshore retention and

potentially driving fluctuations in commercial catches in southeastern Australian estuaries in the

early 20$^{th}$ century (Dannevig, 1907; Suthers *et al.*, 2020). Dannevig (1907) hypothesised that onshore

winds reduced unfavourable advection of fish eggs and larvae, retaining them nearer the coast and

therefore increasing recruitment to estuarine fisheries. When testing this hypothesis, Dannevig

(1907) found a positive correlation between estuarine commercial catch rates and onshore winds

lagged by three or four years (Suthers *et al.*, 2020). Since the inception of this hypothesis, many

studies have shown relationships between wind and juvenile recruitment with the effects largely

being system specific (Nelson *et al.*, 1977; Caputi *et al.*, 2001; Takeshige *et al.*, 2013; Wilson and

Laman, 2021), likely due to complex estuary specific recruitment processes (Boehlert and Mundy,

1988).

The winds in many of the world's coastal upwelling systems are being altered due to climate

change, resulting in both increases (California, Benguela and Humboldt systems) and decreases (e.g.

Iberian system) in upwelling favourable winds (Sydeman *et al.*, 2014). Changing winds are likely to

have a range of effects including enrichment of waters, regional changes in stratification and basin-

scale changes in thermocline structure, all of which may influence the productivity of local fisheries

(Bakun *et al.*, 2010). Concerns have been expressed with regards to changing coastal winds altering

upwelling regimes which provide important nutrients to ecosystems (Bakun and Weeks, 2008; Bakun

*et al.*, 2010; Sorte, 2013). In addition to changes in nutrient supply, variation in upwelling patterns

may also affect the advection and recruitment of larval fish to estuaries.

Estuary specific characteristics also contribute to variation in commercial catch rates

through both short term catchability effects and longer term productivity differences among

estuaries. Within estuarine environments, fish and fisher behaviour can change in response to

external effects such as weather or freshwater flow (Gillson, 2011). Key estuarine fisheries families

such as Sparidae have been shown to reverse their diel behaviour patterns following heavy rain

(Payne *et al.*, 2013), with the catch rates of many species differing between drought and non-

drought conditions (Gillson *et al.*, 2009). Within eastern Australia, it has been demonstrated that

there are consistent differences in hydrology and fisheries production between types of estuaries

defined by their geomorphology (Pease, 1999; Roy *et al.*, 2001). When investigating variation in

estuarine fisheries production, potentially due to larval recruitment effects, it is therefore necessary

to control for differences originating from differences in freshwater flow or estuarine type.

In the present study, we test the original hypothesis of Dannevig (1907) using both larval fish

abundance and commercial fisheries data. To evaluate offshore and onshore winds (upwelling

favourable and downwelling favourable respectively) as a driver of estuarine fisheries production in

southeast Australia, model-based estimates of surface wind were first compared to normalised

abundances of 132 taxa of coastally spawned fish larvae (1990 - 2016), and then to estuarine fishery

catches (4 species, 8 estuaries, July 1997 – June 2007). Our present study extends the idea of

Dannevig (1907) by investigating both larval fish and commercial catch data to test three specific

aims. Firstly, we examine whether coastal winds influence the abundance of coastally spawned

larval fish near the coast. Secondly, we evaluate whether the same winds during the spawning

period influence commercial fisheries harvest when lagged by an appropriate growth period for each

species. Finally, we explore changes in the upwelling and downwelling (onshore) favourable winds

since 1850. We expect that coastally spawned larval abundance will be greater during periods of

onshore winds. If the onshore transport of coastal fish larvae does occur during downwelling

favourable winds, then the effects of this retention should result in increased larval supply to

estuaries, and assuming juvenile fish are resident in estuaries until size of capture, may result in a

detectable effect on commercial fisheries catch rates.

**Method**

*Dataset Descriptions*

*Wind Data*

To provide a consistent estimate of winds, we used the wind speed and direction data from the

Australian Bureau of Meteorology Atmospheric high-resolution Regional Reanalysis for Australia

(BARRA; Su *et al.*, 2019, 2020). This re-analysis product provides hourly wind speeds at a 12 km

resolution over the Australian domain with downscaled 1.5 km resolution within several sub-

domains between 1990 and 2019. For all analyses in the present study, we used near-surface (10 m)

wind data from the 1.5 km resolution data from within the Eastern New South Wales subdomain (Su

*et al.*, 2020). This 1.5 km reanalysis model demonstrates high agreement with observations, with

higher skill than the 12 km, particularly in coastal areas (Su *et al.*, 2020).

To provide a long-term context, we also used wind speed and directions from the 20[th] Century

Reanalysis V2c data collated by the NOAA/OAR/ESRL PSD ([www.esrl.noaa.gov/psd/](www.esrl.noaa.gov/psd/)), which provides

three-hourly wind speed and direction data from 1850 to 2014 at a resolution of ≈200 km (Compo *et*

*al.*, 2015). As both wind products are reanalysis datasets, the models are constrained by observed

values.

*Larval Fish Data*

To investigate the response of larval fish abundance to wind, we used data from the Australian

Integrated Marine Observing System (IMOS) Larval Fish Database (Smith *et al.*, 2018). Fish larvae

were all collected from the top 150m of water using horizontal and oblique tows from variety of

plankton nets with a 300 – 500 μm mesh size (see Smith *et al.* (2018) for full dataset description and

Table S1 for a summary of the samples included in the current analysis). Each sample refers to a

single plankton tow and the abundance of each taxa collected. The database contains abundance

information for 218 taxa collected over 12 research projects (1983 - 2016; Smith *et al.*, 2018).

*Commercial Fisheries Data*

To assess whether the effects of onshore winds during the spawning period, on the abundance of

coastally spawned fish larvae could be detected at the commercial fisheries scale, we used an

estuarine catch-per-unit-effort (CPUE) dataset from eastern Australia (Gillson *et al.*, 2009). While

CPUE is often not considered a robust measure of fish abundance (Richards and Schnute, 1986;

Harley *et al.*, 2001; Haggarty and King, 2006), for our purpose this passive gear provided a measure of fisheries catch (≥80mm mesh size) from which to assess the impacts of variation in recruitment on CPUE and presented the most consistent method available from which an index of abundance could be inferred (Gillson *et al.*, 2009). This dataset consisted of ten years of monthly CPUE data (combined to annual values; July 1997 – June 2007) from gillnet fisheries in located inside eight estuaries along the east Australian coast (Roy *et al.*, 2001; Table S1). An extensive description of the dataset is given in Gillson *et al.* (2009). Briefly, these CPUE data were derived from monthly catch (kg of harvest) and effort (fishing days) for four fish species; yellowfin bream (*Acanthopagrus australis*; Sparidae), sea mullet (*Mugil cephalus*; Mugilidae), dusky flathead (*Platycephalus fuscus*; Platycephalidae) and sand whiting (*Sillago ciliata*; Sillaginidae). These fish species represent the dominant contribution to both commercial and recreational estuarine fisheries harvest. All four of these species are known to spawn in nearshore coastal waters before juveniles recruit to estuaries (spawning seasons and references detailed in Table 1). Sand whiting and dusky flathead larvae are usually found at deeper depths (15-50 m) while sea mullet larvae are found in surface waters (Collins and Stender, 1989). Yellowfin bream larvae are found equally throughout the top 30 m of the water column, with some larvae found up to 70 m deep (Gray, 1993; Gray *et al.*, 2019). Luderick (*Girella tricuspidata*; Girellidae) was in the original data (Gillson *et al.*, 2009), but was not included in the present study due to the variable spawning periods between populations in this region (Gray *et al.*, 2012). Due to reporting requirements at the time, effort represents total gillnet effort on monthly timescale, undifferentiated by species. There is some seasonality to the fishing effort which is generally higher in winter months, corresponding to increased CPUE for flathead and bream, suggesting fishers may target bream and flathead in winter months. As we analysed our data on an annual time scale and there are no changes in annual effort (Figure S3), the seasonality is consistent between years.

*Data Preparation*

All data preparation and analysis was conducted using R v4.0.2 (R Core Team, 2020).

*Wind Data*

Wind speed and direction data were extracted based upon the specific dates and locations of samples (larval fish or estuary locations) using the R packages 'tidyverse' (Wickham, 2017), 'raster' (Hijmans, 2019), 'ncdf4' (Pierce, 2019) and 'RedaS' (Hatzinger *et al.*, 2015). For each sample, the wind direction and speed were taken as the mean of 10 pixels (15 km x 15 km) centred over the sample latitude/longitude.

To investigate the interaction between offshore advection/upwelling favourable (NE) and

retention/downwelling favourable (SE) winds, we separated the onshore winds in eastern Australia

into SE and NE components resulting in two variables. The SE and NE components of wind are most

relevant to the transport of larvae. To calculate the magnitude of wind in a particular direction (SE or

NE winds) from the known direction and speed, we converted the wind direction to radians and

applied a sine function, taking the absolute value to create an effect size for wind displacement. This

can be interpreted as the net movement of air moving along a specified axis (NE or SE). If the wind

was directly from 45° (for NE winds; 135° for SE), then it was a full effect (1); but if it was at a slight

angle, then it was reduced (<1) according to the sine function.

195
$$E = |\sin(r(d + a))|$$

Where $E$ = effect size for the wind displacement, $r$ = a function to convert direction wind is going

towards to radians, $d$ = direction wind is going towards (°), $a$ = 45° for NE calculation or 135° for SE

calculation.

The effect size was then multiplied by the wind speed to get the displacement in each direction per

hour and -1 to account for the data being direction towards and ensure offshore winds were

negative.

202
$$D = E \times s \times -1$$

Where $D$ = displacement (km h$^{-1}$) and $s$ = wind speed (km h$^{-1}$).

Displacement values were then summed to generate a total net displacement (km) over the time

span of interest. A positive net displacement means that a greater amount of air moved onto the

coast than off the coast during the time period. The use of net displacement rather than other

methods aligns with the original work of Dannevig (1907) and Suthers *et al.* (2020) while presenting

a similar metric to Schlaefer *et al.* (2018). Henceforth, references to upwelling favourable winds (NE

winds) or downwelling favourable winds (SE winds) refer to net displacement of air in either

upwelling or downwelling favourable directions. While we are calculating wind displacement, rather

than direct upwelling or downwelling (Ekman transport), wind is a key driver of upwelling in this

region (Schaeffer *et al.*, 2013).

*Larval Fish Data*

We subset the larval fish abundance data to include samples taken on the continental shelf (≤ 1000

m bathymetry; Figure 1) between 30–36° S for the period 1990 to 2016 (to match wind data

availability). In order to investigate recruitment mechanisms affecting all coastally spawned fish,

within each larval fish sample we focused only on taxa known to spawn coastally in this region (Table

S2; Neira *et al.*, 1998, Miskiewicz Unpublished Data). While this may miss some species-specific

effects, we are assuming that coastally spawned taxa will respond in similar ways prior to their

swimming ability improving with larval development, which facilitates movement from coastal

waters into estuarine nursery habitats. To avoid any one taxon dominating the abundance of coastal

larvae, the abundance of each taxon was normalised (i.e. the abundance of each family summed to

1) by transposing the dataset and using the '*normalize.rows()*' function from the 'vegetarian' R

package (Charney and Record, 2012). This created a relative abundance measure and ensured that

each taxon had equal weighting during subsequent analysis, removing species specific effects. The

normalised abundance of each taxon was summed together to create a single metric for analysis.

Following normalisation, the normalised relative coastal abundance across all samples had a mean

of 0.07 (SD = 0.11), with a maximum of 1.48 and the mean abundance was stable between sampling

projects (Figure S1). By using the relative abundance of a suite of species, the possibility exists that

some individual species-level effects may be missed, but during the larval stage (and without size

information) we are assuming that swimming ability is poor and that all species will respond similarly

to coastal winds. The larval fish sampling in Smith *et al.* (2018) was spatially inconsistent and in this

region larval abundance is also generally higher in shallower areas (Hinchliffe *et al.*, 2021).

Therefore, to control potential bias, we calculated the distance (km) to the nearest point of the

Australian mainland for each sample to use as a covariate during modelling.

To separate the effects of advection from upwelling favourable (NE) winds and onshore transport

from downwelling favourable (SE) winds on larval fish abundance, the net air displacement for

upwelling favourable and downwelling favourable winds for each larval fish sample was calculated

by summing the hourly displacement measurements of the previous 14 days. This time period was

selected to quantify both potential upwelling and retention against the coast as there are often

lagged effects on larval abundance through physical transport and nutrient enrichment (Dalley *et al.*,

2002; Ings *et al.*, 2008). While larval transport can happen over short periods of time, 14 days is the

approximate period it takes for upwelling to generate secondary production (increased

phytoplankton and zooplankton) in this region and therefore provide a potentially favourable

environment for fish larvae to develop (Baird *et al.*, 2006).

*Commercial Catch*

For our investigation, we aggregated all monthly data to an annual scale (July – June). Over the ten-year period, there were no major regulation or fisher behaviour changes within this fishery and both total catch and CPUE fluctuated despite relatively stable fishing effort within each estuary (Figures S2 – S4). We therefore proceeded under the assumption that fluctuations in annual CPUE may be a reliable proxy for fish abundance (Gillson *et al.*, 2009).

To investigate the effects of upwelling and downwelling favourable winds during the spawning periods of the commercially important fish species and the effects on lagged annual CPUE, onshore winds were calculated for each estuary (located 0.15°E of the estuary mouth; Figure 1), centred, and scaled according to the above method with upwelling (NE) and downwelling (SE) favourable components. The net air displacement during the spawning period was determined by identifying the spawning periods of each species from published literature (Table 1). Rather than exploring multiple lags, we identified lags *a priori* by using the modal age of these species caught by gillnets in these estuaries (Gray *et al.*, 2015). The modal age is the age at which the species are fully recruited to the fishery, removing any effect of gear selectivity. This age was then used to lag the spawning period winds to correspond to the most abundant (modal) age class, which are therefore most likely to show an effect of the onshore winds if they were influencing larval recruitment. As drought has previously been shown to be an important driver for this CPUE dataset (Gillson *et al.*, 2009), we included drought as the number of months each estuary was 'drought declared' during each 12-month CPUE period based on the New South Wales Department of Primary Industries drought situation maps. Our dataset also contained three types of estuaries (barrier river, drowned river valley & barrier lagoon) which might have differing responses to drought due to their different flushing schemes (Roy *et al.*, 2001).

*Statistical Analysis*

*Effects of Wind on Larval Fish Abundance*

The larval fish data covered a period of 27 years (1990 – 2016) and included 1,489 larval fish samples that were used to calculate the normalised coastal abundance metric for each sample described above. A total of 60 larval fish samples (4%) contained no (zero) coastally spawned larvae. Rather than remove these samples we used a two-stage gamma hurdle model to test the effects of

upwelling favourable (NE) and downwelling favourable (SE) winds on larval fish abundance. This

gamma hurdle model first analyses all the data in a presence-absence method using a Bayesian

binomial model with a logit link. This is followed by a Bayesian generalised linear mixed model with a

gamma error distribution using a log link for the presence only data because the abundance data

had a continuous, positively skewed, distribution. The model included the fixed effects of

downwelling favourable winds, upwelling favourable winds and distance from the coast (km) as well

as interaction terms and separate quadratic terms for both downwelling and upwelling favourable

winds with interactions with distance to coast. Both linear and quadratic terms for the downwelling

and upwelling favourable winds were included in the models because it was hypothesised that the

winds would have a disproportionate effect on larval fish abundance at low or high speeds, hence a

non-linear fit may be appropriate. Distance to coast was included as a covariate to control for the

natural increase in larval abundances closer to the Australian mainland (Hinchliffe *et al.*, 2021).

Distance to coast, upwelling favourable (NE) and downwelling favourable (SE) winds were centred

and standardised to assist the model fitting process and increase interpretability of the model

coefficients (Schielzeth, 2010). This means a wind of 0 is interpreted as moderate (i.e. the mean)

while positive values are stronger winds and negative values are weaker than the mean.

Since the Australian IMOS Larval Fish database includes data from a variety of projects which used

slightly different sampling methodologies (Smith *et al.*, 2018), the model also included a random

intercept effect for Project. As each Project occurred in a discrete time period, this random effect

also controls for any temporal inconsistencies such as a traditional year effect. The use of normalised

larval abundances also creates a more consistent dataset despite the differences in sampling over

time. The fitted model thus had the form:

$$Abund_{it} = \begin{cases} 0 & with\ probability\ p\ and \\ Gamma(\mu_{i,t}, \varphi) & with\ probability\ 1-p \end{cases}$$

$$\eta_{i,t} = \alpha + Up\_Winds_{i,t} + Up\_Winds^2_{1,t} + Down\_Winds_{i,t} + Down\_Winds^2_{1,t} + Up\_Winds_{i,t}$$

$$* Down\_Winds_{i,t} + Distance\_to\_Coast_{i,t} + v_t$$

$$v_t \sim Normal(0, \sigma_t)$$

Where the probability that the relative abundance of coastally spawned larvae (*Abund*) is 0 was

modelled as logit(*p*); $\alpha$ is a constant intercept; Variables relate wind and distance to coast variables

to the abundance of coastally spawned larvae in sample *i* from project *t*. * represents an interaction

term in the model. *Abund* was modelled as Gamma distributed with mean $\mu_{i,t}$ fitted via a log link

with linear predictor $\eta_{i,t}$, and shape parameter $\varphi$. $v_t$ is a random intercept by project with mean 0

and variance $\sigma_t^2$.

Model parameters were estimated by MCMC using Stan (Carpenter *et al.*, 2017) using the function

brm() of R package 'brms' (Bürkner, 2018). Four parallel MCMC chains were run for 10,000 iterations

(5,000 warm up and 5,000 samples each chain) and all parameter estimates were presented with

their 95% Bayesian credible intervals (CI). Convergence was assessed by visually examining MCMC

trace plots and assessment of the Gelman–Rubin statistic (Brooks and Gelman, 1998). The overall

model fit was assessed by extracting residuals from the model and generating standard Pearson's

residual vs fitted plots and normal quantile plots, following the model checking procedure of Gillson

*et al*. (2020). The model was considered stable as the chains were well mixed (Figure S5) the

Gelman–Rubin test statistic < 1.01 for all parameters. The shape and hurdle parameter estimates did

not overlap zero (shape 95% CI: 0.54 – 0.62, hu 95% CI: 0.13 – 0.17), thereby justifying the added

model complexity in this case. There was some deviance from model fit (Figure S6), particularly in

the tails of the dataset but due to the large number of samples and appropriateness of the gamma

hurdle structure we proceeded with the analysis.

Default weakly informative priors were used for all parameters: improper flat priors over the reals

for covariate effects, Student-$t(\mu = -3.1, \sigma = 2.5, \nu = 3)$ for the intercept, Student-$t(\mu = 0, \sigma = 2.5, \nu = 3)$

3) for the standard deviation terms, Gamma(0.01, 0.01) for $\varphi$ and logistic(0, 1) for the zero $Abund$

probability parameter. To visualise the effects of the model predictors, marginal effects were

calculated using the '*ggeffects()*' function in the 'ggeffects' R package (Lüdecke, 2018).

*Effect of wind on estuarine fisheries catch rates*

A Bayesian linear mixed model with gaussian error distribution was used to assess the effects of

coastal winds on annual CPUE. The model included fixed effects of downwelling favourable winds,

upwelling favourable winds (NE), drought months and estuary type, interactions between drought

months and estuary type, upwelling and downwelling favourable winds as well as quadratic terms

for both upwelling and downwelling favourable winds. To investigate the overall effect of onshore

winds on annual CPUE and incorporate the dependency structure among observations from the

same estuary or species, we used estuary as a random intercept and species as a random slope

effect as part of a Bayesian linear mixed model. This controls for our data coming from the same

eight estuaries over a ten-year period with species crossed with estuary. The random intercept lets

each estuary have different overall level of production while the random slope effect lets each

species respond differently to the environmental variables. Using the annual CPUE as the response

variable, the linear mixed model was fit and assessed using the same method described above for

the larval fish analysis. The fitted model thus had the form:

$$CPUE_{jns} = UpWind_{jns} + DownWind_{jnsit} + UpWind_{jns} \times DownWind_{jns} + UpWind_{jns}^2$$
$$+ DownWind_{jns}^2 + EstuaryType_{jns} + DroughtMonths_{jns}$$
$$+ EstuaryType_{jns} \times DroughtMonths_{jns} + Estuary_n \times Species_s$$

$$Estuary_n \sim N(0, \sigma_n)$$

$$Species_s \sim N(0, \sigma_s)$$

Where $CPUE_{jns}$ is the $j$th CPUE observation in $Estuary\ n$ for $Species\ s$. $Estuary_n$ is the random

intercept and $Species_s$ is the random slope effect. $UpWind$ is the standardised upwelling

favourable winds, $DownWind$ is the standardised downwelling favourable winds, $EstuaryType$ is

the type of estuary an observation was from, $DroughtMonths$ is the number of months of

'declared drought' during the year. Uninformative flat priors were used for all variables except the

overall intercept and random intercepts for $Estuary$ and $Species$. The overall intercept used a

Student-$t$ ($\mu$ = 30.8, $\sigma$ = 40.3, $v$ = 3) and the random intercepts for $Estuary$ and $Species$ both used

Student-$t$ ($\mu$ = 0, $\sigma$ = 40.3, $v$ = 3). The CPUE model showed good convergence of chains with

Gelman–Rubin test statistic < 1.01 for all parameters but there was some divergence in model fit

towards the tails (Figures S9 & S10).

*Historical Wind Changes*

To investigate whether or not upwelling or downwelling favourable winds have changed over time in

southeast Australia, we used the 20[th] Century Reanalysis V2c dataset (1850–2014; Compo *et al.*,

2015) and calculated upwelling favourable (NE) and downwelling favourable (SE) winds as described

above, except net annual displacement was derived by summing all winds during a Gregorian

calendar year (centred on Sydney 33.839° S 151.309° E; Figure 1). To ensure consistency with the

high-resolution BARRA model used for the prior analyses, we initially tested the correlation between

the BARRA model and 20[th] Century Reanalysis V2c model using both annual displacement for

upwelling favourable (NE) and downwelling favourable (SE) winds for the overlapping years (1990–

2014) centred over Sydney. There was a moderate to high correlation for both upwelling favourable

($r$ = 0.599, $t_{23}$ = 3.589, $P$ = 0.002) and downwelling favourable ($r$ = 0.697, $t_{23}$ = 4.658, $P$ < 0.001)

winds. We then applied two separate Bayesian linear models with gaussian error distributions for

upwelling favourable (NE) and downwelling favourable (SE) winds with year as a fixed effect (flat

uninformative prior). Initial exploration revealed temporal autocorrelation in the first and second

year of the wind time-series data as measured with the '*acf*' function from the 'stats' R package (R

Core Team, 2020). Therefore, wind data from every third year were used in the final analysis to

remove the presence of temporal autocorrelation. Bayesian linear models were fit and assessed

using the same method described above for the larval fish analysis, with good mixing of chains and

model fit (Figures S11 – S14).

*Interpretation and Sensitivity Analysis*

For interpretation of all model outputs in this study we used the median posterior estimate for

parameters and the 95% Bayesian CI. If the 95% CI did not overlap zero, we deemed a parameter

important (van der Linden and Chryst, 2017). Conditional (full model) and marginal (fixed factors

only) $R^2$ values were calculated using the method of Gelman *et al.* (2019) and implemented in the

function '*r2_bayes()*' of R package 'performance' (Lüdecke *et al.*, 2020).

For both the larval and CPUE models, sensitivity analyses were performed on the lag times used to

assess the robustness of our findings. This involved running the models multiple times while altering

the duration of the lag period. For the larval models, we varied the lead up time for winds by 9 – 20

days prior to sampling and for the CPUE models we simulated altering the lag by ± 2 years from the

identified modal age. This is important for the CPUE models as there are potential influences in the

data due to the harvesting of multiple age classes. As we choose the modal age from gill net

sampling, our original lag should have the strongest effect, and if multiple age classes are present in

a sample, then the effects of wind should be similar but potentially weaker if an age class is less

abundant.

**Results**

*Coastally Spawned Larval Fish*

A total of 175,112 larval fish from 132 coastal spawning taxa were present in 1,489 observations on the continental shelf between 30 and 36° S. Examining the winds 14 days prior to sampling revealed evidence of an interaction between distance to the coast, upwelling favourable (NE) wind and downwelling favourable winds (Estimate = -0.48, 95% CI: -0.85 – -0.13; Figure 2). This interaction can be interpreted as upwelling and downwelling favourable winds interact with each other in different ways depending on the distance a sample is taken from the coast. An overall decline in abundance was observed with increasing distance from the coast (Figure 3). Upwelling favourable winds had a nonlinear effect on abundance where when upwelling favourable winds increased from below average to moderate amounts (0 on standardised axis), larval abundance increased but then as the winds strengthened further, abundance decreased (Figure 3). This resulted in an optimum threshold where larvae were most abundant following moderate amounts of upwelling favourable wind. Downwelling favourable winds showed a linear positive effect on abundance (Figure 3). Both the upwelling and downwelling favourable wind relationships were strongest near the coast and the relationships became less defined with increasing distance from the coast (Figure S15).

Our 14 day wind model explained a small amount of variance in larval fish abundance (conditional $R^2$ = 0.059, marginal $R^2$ = 0.044), and the sensitivity analysis of the wind lead up times in the larval fish model showed that estimates of most effect sizes did not clearly change between 9 – 21 days lead up (overlapping error bars; Figure S16). Two changes were evident in the sensitivity analysis: 1) the interactions between upwelling favourable winds and downwelling favourable winds became inconsequential when a 9 – 12 day lag was tested (error bars overlaps 0); and 2) the effect size of interaction between upwelling favourable winds, downwelling favourable winds and distance to coast became smaller at a 9 day lag, although still remained important and negative.

*Commercial Estuarine Catch Rates and Historical Winds*

The multi-species model performed well (conditional $R^2$ = 0.822, marginal $R^2$ = 0.506) and showed strong evidence for a negative effect of upwelling favourable winds during the spawning period on CPUE (Estimate: -230.48, 95% CI: -374.58 – -85.05; Figures 4 & 5). There was no evidence of any effect from downwelling favourable winds on CPUE (Estimate: -88.62, 95% CI: -227.90 – 50.63; Figure 4).

Aside from the wind effects, there was evidence of a positive effect of drought on CPUE (Estimate =
8.16, 95% CI: 4.56 – 11.70) although this effect was variable depending on estuary type. Barrier
Lagoon estuaries had a more positive response to drought compared with Barrier Rivers and
Drowned River Valleys (Figure S17).

The sensitivity analysis of the wind lag times in the CPUE model showed that altering the lag times
only influenced the wind effects with the standard error of all other estimates overlapping the lag
change 0 estimates for all the lags considered (Figure S18). When the lag was shortened by one year
or increased by two years, there was an interaction evident between upwelling favourable winds
and downwelling favourable winds (Figure S18). The effect size of upwelling favourable winds varied
with lag time. The effect became inconsequential when the lag was shortened by two years while a 1
year longer lag had a stronger effect on CPUE compared to a 1 year shorter lag. The effect of altering
the lag time of downwelling favourable winds was more variable with a non-linear pattern occurring
if the lag was increased by two years. If the lag was reduced by 1 year, the effect of downwelling
favourable winds became inconsequential.

Between 1850 and 2014, there was clear evidence of an increase in upwelling favourable winds
(Estimate of annual change: 52.86, 95% CI: 19.00 – 85.59; Figure 6) and a decline in onshore
transport causing downwelling favourable winds (Estimate of annual change: -40.93, 95% CI: -78.18
– -3.71; Figure 6). The upwelling favourable winds shifted from a negative net displacement to a
positive net displacement, while the downwelling favourable winds remained net positive but
declined to approximately half of their initial levels (Figure 6).

**Discussion**

By combining three datasets from a larval fish database, a high-resolution wind re-analysis model and commercial CPUE time-series data, we have demonstrated that coastal winds may influence the abundance of coastally spawned taxa (Figures 2 & 3), potentially resulting in detectable effects on commercial fisheries catch rates (Figures 5 & 6). While we initially expected samples collected following strong onshore winds to have the highest abundance of coastal larvae, we found both upwelling and downwelling favourable winds are important for fish larvae. Moderate amounts of upwelling favourable winds potentially increased planktonic production (Armbrecht *et al.*, 2014), although above average upwelling favourable winds negatively affected coastal larvae abundance, perhaps due to offshore advection. Strong downwelling favourable winds had a positive influence on the abundance of coastally spawned larvae, likely driving onshore transport which may facilitate estuarine recruitment (Agostini and Bakun, 2002). Using lagged winds from each species' spawning season, we showed a detectable negative effect of strong upwelling favourable winds on CPUE, possibly due to the advection of larvae away from favourable habitat. This was a similar effect to that observed in the larval fish with strong upwelling favourable wind being correlated with low relative abundance of coastal larvae. We found no evidence of the hypothesised positive effect of downwelling favourable winds in the CPUE analysis. The high variance explained in the CPUE models (conditional $R^2$ = 0.82) in the present study indicates that incorporating wind from the spawning period into future recruitment models might improve forecasts of estuarine fisheries CPUE. This is important as we have demonstrated that coastal winds have changed since 1850 and are expected to continue changing in the future (Bakun *et al.*, 2010; Sydeman *et al.*, 2014).

*A proposed recruitment mechanism*

We showed a negative effect of upwelling favourable wind on larval fish abundance when it was above average strength. This is likely caused by offshore transport as strong upwelling drives advection away from the coast and juvenile estuarine and nearshore habitat. It is also recognised that over an approximate 14 day cycle, upwelling can generate increased nutrients and chlorophyll at the surface which may in turn flow into the lower trophic levels (Gasol *et al.*, 1997; Buesa, 2019). This upwelling may increase prey availability for larval fish, thereby increasing growth and survival rates (Zenitani *et al.*, 2007). Onshore transport from downwelling favourable winds is important as it retains larvae near the coast. Based upon previous research, the positive effects of this retention are

most visible after upwelling preconditions the ecosystem with nutrients, thereby creating a
favourable environment near juvenile habitat (Rykaczewski and Checkley, 2008).

Our sensitivity analysis showed a consistently positive effect of downwelling favourable winds
regardless of lag time. We believe this supports our hypothesis that onshore transport may increase
recruitment into estuaries by larval fishes as they are geographically closer and stochastic dispersal
will be reduced (Bruno *et al.*, 2018).

*Commercial Estuarine Fisheries*

Consistent with Dannevig's (1907) observation, we found coastal winds during the spawning period
can influence commercial fisheries catch rates. This is consistent with the relationships we found
between coastal winds and the abundance of coastally spawned larval fish. However, likely due to
the coarse temporal resolution of the CPUE data, there was no evidence of a positive effect of
downwelling favourable winds which represent more immediate transport effects. Based on the
demonstrated effects of these winds on coastal larval fish abundance, the events most likely to
contribute to successful recruitment would be short periods of upwelling (to precondition the area
and generate larval prey), followed by periods of onshore transport (downwelling). We propose our
results are showing a negative effect of upwelling favourable winds on CPUE because offshore
transport is increased during strong upwelling, which negatively affects recruitment. As our analysis
is correlative rather than a manipulative experiment, the exact mechanisms underlying the
relationship between upwelling favourable winds and CPUE cannot be pinpointed. Temperature
mediated changes in growth and survival are likely to be a key driver of this relationship. Upwelling
favourable winds cause upwelling which draws colder water up from depth (Schaeffer *et al.*, 2013). If
larvae are not advected offshore with the displaced surface water they would remain in a much
cooler habitat which would likely result in slower growth, potentially increasing risk of predation
(Pepin, 1991; Buckley *et al.*, 2008). To understand possible temperature dynamics, future work
should potentially consider incorporating water temperature into studies of both larval abundance
and growth as well as commercial catch data.

The generally weak positive effect of drought on CPUE (with different effects in some types of
estuaries) agrees with previous research using this dataset that found monthly bream CPUE
increased during periods of drought (Gillson *et al.*, 2009). Despite this, previous research also
showed that the monthly CPUE of other species generally declined during periods of drought
suggesting that the effect of drought on CPUE needs further investigation (Gillson *et al.*, 2009). We

believe the contrasting results in our study compared to Gillson *et al.* (2009) are the result of using

different temporal scales, with shorter timescales likely more representative of changes in both fish

and fisher behaviour (Gillson *et al.*, 2009).

The sensitivity analysis conducted on the lag times between the winds and year of capture identified

that the modal ages show the strongest effects of coastal winds, as would be expected with ages

(represented by lags) which are less common in the harvest having smaller effect sizes. In the future,

it may be possible to conduct more detailed analyses of single species by identifying particular

bottlenecks in the lifecycle including recruitment, which may be affected by coastal winds.

*Historical changes in onshore winds*

Since 1850, upwelling favourable winds have increased while downwelling favourable winds have

decreased in the southeast Australian region. While the direct cause of these changes is uncertain, it

is possible this has been a response to global climate change. For example, intensification of surface

winds have been attributed to a decline in ozone around Antarctica, which resulted in large scale

changes in southern hemisphere winds including over Australia (Cai, 2006). The demonstrated

changes in wind in southeast Australia would likely have reduced the onshore transport of larval fish,

impacting recruitment to estuaries because increased upwelling and decreased downwelling would

increase prey availability but reduce retention near the coast, potentially transporting larvae further

away from estuaries.

Despite this, very few fish species spawn all year round, and it is possible that seasonal changes in

wind may be more important than annual changes. The temporal analysis presented here does not

test alternating upwelling and downwelling favourable winds, and it would be useful to calculate a

metric that encompasses the alternating nature of upwelling and downwelling favouring winds. A

possible approach could be to document the number and duration of upwelling and downwelling

events.

Previous research has indicated that the intensity of wind driven upwelling is likely to increase in

most regions, with the exception of southwestern Africa (Bakun *et al.*, 2010). This agrees with our

finding that upwelling favourable winds have historically increased in southeast Australia. While this

may result in increased fisheries production by increasing nutrient availability, it is a complex system

in which changing winds could have a multitude of effects including increasing productivity and

nutrient concentrations of source waters, regional changes in stratification and basin-scale changes

in thermocline structure (Bakun *et al.*, 2010).

*Limitations of this study*

While our study successfully combined three datasets to assess the effects of wind on larval fish

retention and recruitment, it does have limitations. Our study used correlative analyses and several

potential explanatory variables were not included in our models, which may have captured

additional variance in either larval fish abundance or commercial fisheries catch rates. These include

varying population spawning biomasses, water temperature, oceanographic currents, improved

larval swimming ability with development or the effects of density dependence, all of which

influence spawning and/or recruitment (Ottersen and Sundby, 1995; Schilling *et al.*, 2020).

An important limitation of the present study was that by using a historical larval dataset, which was

collected opportunistically, we could not directly quantify the movement or survival of larval fish

during wind events and instead we could only look at abundances at the time of sampling while

controlling for variation such as distance from coast and the individual project which collected the

data. A more robust analysis would be to structure a survey that repeatedly samples in a grid pattern

during wind events and directly quantifies the movement of larvae. As no larval fish size data were

available, swimming ability was not considered, which is known to be an important factor

influencing larval movement (Leis, 2007; Drake *et al.*, 2018). While this could potentially bias

analyses, we considered the swimming ability of each sample to be representative of a normal

distribution (modelled as part of the random project effect) and any strong swimmers would have

likely avoided capture by the slow plankton net.

The CPUE analysis has several key limitations which may obscure the effects of wind during the larval

period. The most important of these is the duration between recruitment and commercial harvest,

which can be up to 7 years in our study. This is a large time interval during which there may be many

sources of mortality or population limitation. A potential limiting factor in estuaries is habitat

limitation, if there is insufficient habitat to support recruiting larvae as juveniles, then the total

population of the species will be limited by habitat rather than larval recruitment. While no

information is available on potential habitat limitation in the estuaries we studied, in nearby

estuaries closed to commercial fishing there is some evidence of habitat limitation (Folpp *et al.*,

2020). The other limitation is that multiple age classes are harvested at a time and by choosing a

modal age (based upon 2008-2010 harvest; Gray *et al*. 2015), any effects with the influence of other

ages are blurred. While not possible for this study, a more rigorous approach would have been to

only use the CPUE of the modal age class being harvested rather than the total CPUE. This blurring

effect may be a contributor to the variation observed in the CPUE sensitivity analysis (Figure S18).

The present study also did not investigate the abundance of larval fish entering estuaries (Boehlert

and Mundy, 1988), which would be an important metric to confirm increased recruitment following

favourable wind conditions. By expanding investigations beyond the larval period, fish recruitment

mechanisms in this region might be better understood.

*Conclusions*

Various studies have demonstrated positive effects (Nelson *et al.*, 1977) and negative effects (Parrish

*et al.*, 1981; Takeshige *et al.*, 2013) of wind-driven Ekman transport of larval fish for estuarine

recruitment. Despite the low variance explained in the larval fish analysis, potentially caused using a

large database containing samples taken over large temporal and spatial scales, the present study is

the first to show a correlation between coastal winds and fish larvae and detect a similar

corresponding effect on the commercial fishery. We suggest a recruitment mechanism involving

wind driven coastal enrichment and larval retention during the spawning season, which is evident in

commercial fisheries catch rates. This mechanism involves increased productivity from wind-driven

upwelling and increased retention near the coastline, which work in tandem to increase overall

recruitment of larvae into estuaries. These recruitment dynamics were potentially detected in

commercial estuarine fisheries data when appropriate lags are applied and by incorporating

spawning period winds into recruitment models, may improve predictions of commercial fisheries

catch rates. As climate change is altering wind patterns, it is likely that wind driven ocean dynamics

will continue to vary and it is recommended that scientists and management consider potential

changes in recruitment.

**Acknowledgements**

This study was prompted by Dennis Reid (Australian Museum) who brought to our attention Harald

Dannevig's original insight on the effect of coastal winds on estuarine fisheries. Support for the 20th

Century Reanalysis Project version 2c dataset is provided by the U.S. Department of Energy, Office of

Science Biological and Environmental Research (BER), and by the National Oceanic and Atmospheric

Administration Climate Program Office. HTS was supported by the UNSW Network Laboratory for

Ocean Collaboration and NSW Research, Acceleration and Attraction Program. CH is supported by an

Australian Government Research Training Program Scholarship. Larval fish data were sourced from

the Integrated Marine Observing System (IMOS), an initiative of the Australian Government being

conducted as part of the National Collaborative Research Infrastructure Strategy and the Super

Science Initiative. This research includes computations using the computational cluster Katana

supported by Research Technology Services at UNSW Sydney.

**Author contributions**

HTS, IMS, AM & JPG collected the data, IMS conceived the ideas, HTS, CH, JPG & AM designed the

analysis, HTS performed the analysis and led the writing of the manuscript. All authors contributed

critically to the drafts and gave final approval for publication.

**Data Availability Statement**

The larval fish data is freely available from the Australian Ocean Data Network

https://portal.aodn.org.au/ (Smith *et al.*, 2018). The BARRA wind model is freely available upon

request to the Australian Bureau of Meteorology (Su *et al.*, 2019). 20th Century Reanalysis V2c data

provided by the NOAA/OAR/ESRL PSL, Boulder, Colorado, USA, from their Web site at

https://psl.noaa.gov/ (Compo *et al.*, 2015). NSW Commercial Fisheries Catch Data is available upon

request to the NSW Department of Primary Industries Fisheries. Extracted data and code used in this

analysis is available at https://github.com/HaydenSchilling/Wind-and-Fisheries.

**Conflict of Interest Statement**

The authors declare no conflicts of interest.

**References**

629 Agostini, V. N., and Bakun, A. 2002. `Ocean triads' in the Mediterranean Sea: physical mechanisms
630   potentially structuring reproductive habitat suitability (with example application to
631   European anchovy, *Engraulis encrasicolus*). Fisheries Oceanography, 11: 129–142.
632 Anderson, J. 1988. A review of size dependent survival during pre-recruit stages of fishes in relation
633   to recruitment. Journal of Northwest Atlantic Fishery Science, 8: 55–66.
634 Armbrecht, L. H., Roughan, M., Rossi, V., Schaeffer, A., Davies, P. L., Waite, A. M., and Armand, L. K.
635   2014. Phytoplankton composition under contrasting oceanographic conditions: Upwelling
636   and downwelling (Eastern Australia). Continental Shelf Research, 75: 54–67.
637 Bailey, K. M., and Houde, E. D. 1989. Predation on Eggs and Larvae of Marine Fishes and the
638   Recruitment Problem. *In* Advances in Marine Biology, pp. 1–83. Ed. by J. H. S. Blaxter and A.
639   J. Southward. Academic Press.
640 Baird, M. E., Timko, P. G., Suthers, I. M., and Middleton, J. H. 2006. Coupled physical–biological
641   modelling study of the East Australian Current with idealised wind forcing: Part II. Biological
642   dynamical analysis. Journal of Marine Systems, 59: 271–291.
643 Bakun, A., and Weeks, S. J. 2008. The marine ecosystem off Peru: What are the secrets of its fishery
644   productivity and what might its future hold? Progress in Oceanography, 79: 290–299.
645 Bakun, A., Field, D. B., Redondo-Rodriguez, A., and Weeks, S. J. 2010. Greenhouse gas, upwelling-
646   favorable winds, and the future of coastal ocean upwelling ecosystems. Global Change
647   Biology, 16: 1213–1228.
648 Boehlert, G., and Mundy, B. 1988. Roles of behavioral and physical factors in larval and juvenile fish
649   recruitment to estuarine nursery areas. *In* American Fisheries Society Symposium, pp. 51–
650   67.
651 Brooks, S. P., and Gelman, A. 1998. General methods for monitoring convergence of iterative
652   simulations. Journal of Computational and Graphical Statistics, 7: 434–455. Taylor & Francis.
653 Bruno, D. O., Delpiani, S. M., and Acha, E. M. 2018. Diel variation of ichthyoplankton recruitment in a
654   wind-dominated temperate coastal lagoon (Argentina). Estuarine Coastal and Shelf Science,
655   205: 91–99.
656 Buckley, L. J., Caldarone, E. M., and Clemmesen, C. 2008. Multi-species larval fish growth model
657   based on temperature and fluorometrically derived RNA/DNA ratios: results from a meta-
658   analysis. Marine Ecology Progress Series, 371: 221–232.
659 Buesa, R. J. 2019. Plankton based energy transfer rates in four Cuban coastal lagoons. Estuarine,
660   Coastal and Shelf Science, 216: 118–127.
661 Burchmore, J. J., Pollard, D. A., Middleton, M. J., Bell, J. D., and Pease, B. C. 1988. Biology of four
662   species of whiting (Pisces: Sillaginidae) in Botany Bay, NSW. Marine and Freshwater
663   Research, 39: 709–727.
664 Bürkner, P.-C. 2017. brms: An R package for bayesian multilevel models using Stan. Journal of
665   Statistical Software, 80: 1–28.
666 Bürkner, P.-C. 2018. Advanced bayesian multilevel modeling with the R package brms. The R Journal,
667   10: 395–411.
668 Cai, W. 2006. Antarctic ozone depletion causes an intensification of the Southern Ocean super-gyre
669   circulation. Geophysical Research Letters, 33: L03712.
670 Caputi, N., Chubb, C., and Pearce, A. 2001. Environmental effects on recruitment of the western rock
671   lobster, *Panulirus cygnus*. Marine and Freshwater Research, 52: 1167–1174.
672 Carpenter, B., Gelman, A., Hoffman, M. D., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M., *et al.*
673   2017. Stan: A probabilistic programming language. Journal of Statistical Software, 76: 1–32.
674 Charney, N., and Record, S. 2012. vegetarian: Jost diversity measures for community data, R package
675   version 1.3.
676 Ciannelli, L., Bailey, K., and Olsen, E. M. 2015. Evolutionary and ecological constraints of fish
677   spawning habitats. ICES Journal of Marine Science, 72: 285–296.

Collins, M. R., and Stender, B. W. 1989. Larval striped mullet (*Mugil Cephalus*) and white mullet
(*Mugil Curema*) off the southeastern United States. Bulletin of Marine Science, 45: 580–589.
Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Allan, R. J., McColl, C., Yin, X., Giese, B. S., *et al.*
2015. NOAA/CIRES Twentieth Century Global Reanalysis Version 2c.
Cury, P. M., Fromentin, J.-M., Figuet, S., and Bonhommeau, S. 2014. Resolving Hjort's dilemma: How
is recruitment related to spawning stock biomass in marine fish? Oceanography, 27: 42–47.
Oceanography Society.
Dalley, E. L., Anderson, J. T., and deYoung, B. 2002. Atmospheric forcing, larval drift, and recruitment
of capelin (*Mallotus villosus*). ICES Journal of Marine Science, 59: 929–941.
Dannevig, H. C. 1907. On some peculiarities in our coastal winds and their influence upon the
abundance of fish in inshore waters. Journal of the Royal Society of NSW, 41: 27–45.
Drake, P. T., Edwards, C. A., Morgan, S. G., and Satterthwaite, E. V. 2018. Shoreward swimming
boosts modeled nearshore larval supply and pelagic connectivity in a coastal upwelling
region. Journal of Marine Systems, 187: 96–110.
Everett, J. D., Baird, M. E., Roughan, M., Suthers, I. M., and Doblin, M. A. 2014. Relative impact of
seasonal and oceanographic drivers on surface chlorophyll a along a Western Boundary
Current. Progress in Oceanography, 120: 340–351.
Folpp, H. R., Schilling, H. T., Clark, G. F., Lowry, M. B., Maslen, B., Gregson, M., and Suthers, I. M.
2020. Artificial reefs increase fish abundance in habitat-limited estuaries. Journal of Applied
Ecology, 57: 1752–1761.
Garvine, R. W., Epifanio, C. E., Epifanio, C. C., and Wong, K. C. 1997. Transport and recruitment of
blue crab larvae: a model with advection and mortality. Estuarine Coastal and Shelf Science,
45: 99–111.
Gasol, J. M., del Giorgio, P. A., and Duarte, C. M. 1997. Biomass distribution in marine planktonic
communities. Limnology and Oceanography, 42: 1353–1363.
Gelman, A., Goodrich, B., Gabry, J., and Vehtari, A. 2019. R-squared for bayesian regression models.
The American Statistician, 73: 307–309. Taylor & Francis.
Gillson, J., Scandol, J., and Suthers, I. 2009. Estuarine gillnet fishery catch rates decline during
drought in eastern Australia. Fisheries Research, 99: 26–37.
Gillson, J. 2011. Freshwater flow and fisheries production in estuarine and coastal systems: Where a
drop of rain is not lost. Reviews in Fisheries Science, 19: 168–186. Taylor & Francis.
Gillson, J. P., Maxwell, D. L., Gregory, S. D., Posen, P. E., Riley, W. D., Picken, J. L., and Assunção, M.
G. L. 2020. Can aspects of the discharge regime associated with juvenile Atlantic salmon
(*Salmo salar* L.) and trout (*S. trutta* L.) densities be identified using historical monitoring data
from five UK rivers? Fisheries Management and Ecology, 27: 567–579.
Gray, C. A. 1993. Horizontal and vertical trends in the distributions of larval fishes in coastal waters
off central New South Wales, Australia. Marine Biology, 116: 649–666.
Gray, C. A., Haddy, J. A., Fearman, J., Barnes, L. M., Macbeth, W. G., and Kendall, B. W. 2012.
Reproduction, growth and connectivity among populations of *Girella tricuspidata* (Pisces:
Girellidae). Aquatic Biology, 16: 53–68.
Gray, C. A., Young, C. L., Johnson, D. D., and Rotherham, D. 2015. Integrating fishery-independent
and -dependent data for improved sustainability of fisheries resources and other aspects of
biodiversity. Sydney, Australia.
Gray, C. A., and Barnes, L. M. 2015. Spawning, maturity, growth and movement of *Platycephalus*
*fuscus* (Cuvier, 1829) (Platycephalidae): fishery management considerations. Journal of
Applied Ichthyology, 31: 442–450.
Gray, C. A., Miskiewicz, A. G., Otway, N. M., and Kingsford, M. J. 2019. Historical daytime vertical
structure of larval fish assemblages in southeast Australian coastal waters: A benchmark for
examining regional ecosystem change. Regional Studies in Marine Science, 29: 100634.
Griffin, D. A., and Middleton, J. H. 1992. Upwelling and internal tides over the inner New South
Wales continental shelf. Journal of Geophysical Research: Oceans, 97: 14389–14405.

Haggarty, D. R., and King, J. R. 2006. CPUE as an index of relative abundance for nearshore reef
fishes. Fisheries Research, 81: 89–93.
Hare, J., and Govoni, J. 2005. Comparison of average larval fish vertical distributions among species
exhibiting different transport pathways on the southeast United States continental shelf.
Fishery Bulletin, 103: 728–736.
Harley, S. J., Myers, R. A., and Dunn, A. 2001. Is catch-per-unit-effort proportional to abundance?
Canadian Journal of Fisheries and Aquatic Sciences. NRC Research Press Ottawa, Canada.
https://cdnsciencepub.com/doi/abs/10.1139/f01-112 (Accessed 18 March 2021).
Hatzinger, R., Hornik, K., Nagel, H., and Maier, M. J. 2015. Companion to: *In* R: Einführung durch
angewandte Statistik, 2nd Edition. Pearson Studium, München.
Helbig, J. A., and Pepin, P. 1998. Partitioning the influence of physical processes on the estimation of
ichthyoplankton mortality rates. I. Theory. Canadian Journal of Fisheries and Aquatic
Sciences, 55: 2189–2205.
Hijmans, R. J. 2019. raster: Geographic data analysis and modeling, R package version 3.0.2.
Hinchliffe, C., Smith, J. A., Everett, J. D., Falster, D. S., Lara-Lopez, A., Miskiewicz, A. G., Richardson, A.
J., *et al.* 2021. Modelling the distribution of larval fish in a western boundary current using a
multi-voyage database. Reviews in Fish Biology and Fisheries, 31: 399–415.
Houde, E. D. 2008. Emerging from Hjort's shadow. Journal of Northwest Atlantic Fishery Science, 41:
53–70.
Ings, D. W., Gregory, R. S., and Schneider, D. C. 2008. Episodic downwelling predicts recruitment of
Atlantic cod, Greenland cod and white hake to Newfoundland coastal waters. Journal of
Marine Research, 66: 529–561.
Kell, L. T., Nash, R. D. M., Dickey-Collas, M., Mosqueira, I., and Szuwalski, C. 2016. Is spawning stock
biomass a robust proxy for reproductive potential? Fish and Fisheries, 17: 596–616.
Leis, J. M. 2007. Behaviour as input for modelling dispersal of fish larvae: behaviour, biogeography,
hydrodynamics, ontogeny, physiology and phylogeny meet hydrography. Marine Ecology
Progress Series, 347: 185–193.
Lüdecke, D. 2018. ggeffects: Tidy data frames of marginal effects from regression models. Journal of
Open Source Software, 3: 772.
Lüdecke, D., Makowski, D., Waggoner, P., and Patil, I. 2020. performance: Assessment of regression
models performance, R package version 0.5.0. https://easystats.github.io/performance/.
McClean-Padman, J., and Padman, L. 1991. Summer upwelling on the Sydney inner continental shelf:
The relative roles of local wind forcing and mesoscale eddy encroachment. Continental Shelf
Research, 11: 321–345.
McFarlane, G. A., and Beamish, R. J. 1992. Climatic Influence Linking Copepod Production with
Strong Year-Classes in Sablefish, *Anoplopoma fimbria*. Canadian Journal of Fisheries and
Aquatic Sciences, 49: 743–753.
Middleton, J. H., Cox, D., and Tate, P. 1996. The oceanography of the Sydney region. Marine
Pollution Bulletin, 33: 124–131.
Neira, F. J., Miskiewicz, A. G., and Trnski, T. 1998. Larvae of temperate Australian fishes: laboratory
guide for larval fish identification. UWA Publishing.
Nelson, W. R., Ingham, M. C., and Schaaf, W. E. 1977. Larval transport and year-class strength of
Atlantic menhaden, *Brevoortia tyrannus*. Fishery Bulletin, 75: 23–41.
Ochwada-Doyle, F., Roberts, D., Gray, C., Barnes, L., Haddy, J., and Fearman, J. 2012. Characterizing
the biological traits and life history of *Acanthopagrus* (Sparidae) hybrid complexes:
implications for conservation and management. Journal of Fish Biology, 81: 1540–1558.
Oke, P. R., Roughan, M., Cetina-Heredia, P., Pilo, G. S., Ridgway, K. R., Rykova, T., Archer, M. R., *et al.*
2019. Revisiting the circulation of the East Australian Current: Its path, separation, and eddy
field. Progress in Oceanography, 176: 102139.

Ospina-Alvarez, A., Weidberg, N., Aiken, C. M., and Navarrete, S. A. 2018. Larval transport in the
upwelling ecosystem of central Chile: The effects of vertical migration, developmental time
and coastal topography on recruitment. Progress in Oceanography, 168: 82–99.

Ottersen, G., and Sundby, S. 1995. Effects of temperature, wind and spawning stock biomass on
recruitment of Arcto-Norwegian cod. Fisheries Oceanography, 4: 278–292.

Parrish, R. H., Nelson, C. S., and Bakun, A. 1981. Transport mechanisms and reproductive success of
fishes in the California Current. Biological Oceanography, 1: 175–203.

Payne, N. L., van der Meulen, D. E., Gannon, R., Semmens, J. M., Suthers, I. M., Gray, C. A., and
Taylor, M. D. 2013. Rain reverses diel activity rhythms in an estuarine teleost. Proceedings of
the Royal Society B: Biological Sciences, 280: 20122363. Royal Society.

Pease, B. C. 1999. A spatially oriented analysis of estuaries and their associated commercial fisheries
in New South Wales, Australia. Fisheries Research, 42: 67–86.

Pepin, P. 1991. Effect of temperature and size on development, mortality, and survival rates of the
pelagic early life history stages of marine fish. Canadian Journal of Fisheries and Aquatic
Sciences, 48: 503–518.

Pepin, P. 2016. Reconsidering the impossible — linking environmental drivers to growth, mortality,
and recruitment of fish. Canadian Journal of Fisheries and Aquatic Sciences, 73: 205–215.

Pierce, D. 2019. ncdf4: Interface to unidata netCDF (Version 4 or earlier) format data files, R package
version 1.17.

R Core Team. 2020. R: A language and environment for statistical computing v4.0.2. R Foundation for
Statistical Computing, Vienna, Austria.

Richards, L. J., and Schnute, J. T. 1986. An experimental and statistical approach to the question: Is
CPUE an index of abundance? Canadian Journal of Fisheries and Aquatic Sciences. NRC
Research Press Ottawa, Canada. https://cdnsciencepub.com/doi/abs/10.1139/f86-151
(Accessed 19 March 2021).

Roughan, M., and Middleton, J. H. 2002. A comparison of observed upwelling mechanisms off the
east coast of Australia. Continental Shelf Research, 22: 2551–2572.

Roughan, M., Macdonald, H. S., Baird, M. E., and Glasby, T. M. 2011. Modelling coastal connectivity
in a Western Boundary Current: Seasonal and inter-annual variability. Deep Sea Research
Part II: Topical Studies in Oceanography, 58: 628–644.

Roy, P. S., Williams, R. J., Jones, A. R., Yassini, I., Gibbs, P. J., Coates, B., West, R. J., *et al.* 2001.
Structure and Function of South-east Australian Estuaries. Estuarine, Coastal and Shelf
Science, 53: 351–384.

Rykaczewski, R. R., and Checkley, D. M. 2008. Influence of ocean winds on the pelagic ecosystem in
upwelling regions. Proceedings of the National Academy of Sciences of the United States of
America, 105: 1965–1970.

Santos, A. M. P., Peliz, A., Dubert, J., Oliveira, P. B., Angélico, M. M., and Ré, P. 2004. Impact of a
winter upwelling event on the distribution and transport of sardine (*Sardina pilchardus*) eggs
and larvae off western Iberia: a retention mechanism. Continental Shelf Research, 24: 149–
165.

Schaeffer, A., Roughan, M., and Morris, B. D. 2013. Cross-shelf dynamics in a western boundary
current regime: Implications for upwelling. Journal of Physical Oceanography, 44: 2812–
2813.

Schaeffer, A., Roughan, M., and Wood, J. E. 2014. Observed bottom boundary layer transport and
uplift on the continental shelf adjacent to a western boundary current. Journal of
Geophysical Research-Oceans, 119: 4922–4939.

Schielzeth, H. 2010. Simple means to improve the interpretability of regression coefficients.
Methods in Ecology and Evolution, 1: 103–113.

Schilling, H. T., Everett, J. D., Smith, J. A., Stewart, J., Hughes, J. M., Roughan, M., Kerry, C., *et al.*
2020. Multiple spawning events promote increased larval dispersal of a predatory fish in a
western boundary current. Fisheries Oceanography, 29: 309–323.

Schlaefer, J. A., Wolanski, E., Lambrechts, J., and Kingsford, M. J. 2018. Wind conditions on the Great
Barrier Reef influenced the recruitment of snapper (*Lutjanus carponotatus*). Frontiers in
Marine Science, 5. Frontiers.
Smith, J. A., Miskiewicz, A. G., Beckley, L. E., Everett, J. D., Garcia, V., Gray, C. A., Holliday, D., *et al.*
2018. A database of marine larval fish assemblages in Australian temperate and subtropical
waters. Scientific Data, 5: 180207.
Sorte, C. J. B. 2013. Predicting persistence in a changing climate: flow direction and limitations to
redistribution. Oikos, 122: 161–170.
Stewart, J., Hegarty, A.-M., Young, C., and Fowler, A. M. 2018. Sex-specific differences in growth,
mortality and migration support population resilience in the heavily exploited migratory
marine teleost *Mugil cephalus* (Linnaeus 1758). Marine and Freshwater Research, 69: 385–
394.
Su, C.-H., Eizenberg, N., Steinle, P., Jakob, D., Fox-Hughes, P., White, C. J., Rennie, S., *et al.* 2019.
BARRA v1.0: the Bureau of Meteorology Atmospheric high-resolution Regional Reanalysis for
Australia. Geoscientific Model Development, 12: 2049–2068.
Su, C.-H., Eizenberg, N., Jakob, D., Fox-Hughes, P., Steinle, P., White, C. J., and Franklin, C. 2020.
BARRA v1.0: Kilometre-scale downscaling of an Australian regional atmospheric reanalysis
over four midlatitude domains. Geoscientific Model Development Discussions [Preprint]: 1–
34. Copernicus GmbH.
Suthers, I. M., Reid, D. D., Moksness, E., and Schilling, H. T. 2020. Novel fisheries investigations by
Harald Dannevig: some parallels with Johan Hjort on the other side of the world. ICES
Journal of Marine Science.
Sydeman, W. J., García-Reyes, M., Schoeman, D. S., Rykaczewski, R. R., Thompson, S. A., Black, B. A.,
and Bograd, S. J. 2014. Climate change and wind intensification in coastal upwelling
ecosystems. Science, 345: 77–80.
Szuwalski, C. S., Vert-Pre, K. A., Punt, A. E., Branch, T. A., and Hilborn, R. 2015. Examining common
assumptions about recruitment: a meta-analysis of recruitment dynamics for worldwide
marine fisheries. Fish and Fisheries, 16: 633–648.
Takeshige, A., Miyake, Y., Nakata, H., Kitagawa, T., and Kimura, S. 2013. Effect of wind stress on the
catch of Japanese anchovy *Engraulis japonicus* off northwestern Kyushu, Japan. Fisheries
Science, 79: 989–998.
van der Linden, S., and Chryst, B. 2017. No Need for Bayes Factors: A Fully Bayesian Evidence
Synthesis. Frontiers in Applied Mathematics and Statistics, 3. Frontiers.
https://www.frontiersin.org/articles/10.3389/fams.2017.00012/full (Accessed 9 November
2020).
Wickham, H. 2017. tidyverse: Easily install and load the 'Tidyverse'.
Wilhelm, M. R., Painting, S. J., Field, J. G., Kerstan, M., Durholtz, M. D., Wilhelm, M. R., Painting, S. J.,
*et al.* 2005. Impact of environmental factors on survival of larval and juvenile Cape anchovy
*Engraulis encrasicolus* (G.) in the southern Benguela upwelling region, determined from
hatchdate distributions: implications for recruitment. Marine and Freshwater Research, 56:
561–572. CSIRO PUBLISHING.
Wilson, M. T., and Laman, N. 2021. Interannual variation in the coastal distribution of a juvenile
gadid in the northeast Pacific Ocean: The relevance of wind and effect on recruitment.
Fisheries Oceanography, 30: 3–22.
Zenitani, H., Kono, N., and Tsukamoto, Y. 2007. Relationship between daily survival rates of larval
Japanese anchovy (*Engraulis japonicus*) and concentrations of copepod nauplii in the Seto
Inland Sea, Japan. Fisheries Oceanography, 16: 473–478.
876

878    **Table 1** Details of the spawning season and modal age of capture in gillnet fisheries for four coastally

879    spawning fish species, used to define the choice of lags for winds during the spawning season with

880    an example of the wind intervals used for the annual catch-per-unit-effort (CPUE) between July 1997

881    and June 1998. The ranges of harvested ages (> 5 % of the catch) is shown in brackets with the

882    modal harvested age. Full age distributions can be seen in Gray *et al.* (2015).

| Species | Spawning season | Modal harvested age (Range; Gray *et al.*, 2015) | Example wind intervals used for annual CPUE for July 1997 – June 1998 |
|---|---|---|---|
| bream (*Acanthopagrus australis*) | October – February (Ochwada-Doyle *et al.*, 2012) | 7 (4 – 11) | October 1989 – February 1990 |
| dusky flathead (*Platycephalus fuscus*) | December – March (Gray and Barnes, 2015) | 3 (2-5) | December 1993 – March 1994 |
| sand whiting (*Sillago ciliata*) | December – February (Burchmore *et al.*, 1988) | 6 (3 -10) | December 1991 – February 1992 |
| sea mullet (*Mugil cephalus*) | April – July (Stewart *et al.*, 2018) | 2 (1-5) | April – July 1995 |

886    **Figures**

888    **Figure 1** Location of southeast Australia showing the position of the larval fish samples included in

889    the study (empty grey circles), the estuaries used in the catch-per-unit-effort analysis (filled red

890    circles) and Sydney (filled blue circle). The black line represents the 1000 m isobath.

**Figure 2** Caterpillar plots showing the Bayesian parameter estimates for the coastally spawned larval

fish abundance models based upon winds in the 14 days prior to sampling. Points show the estimate

of the parameter with the horizontal bar representing the 66% (thick bar) and 95% (thin bar)

credible interval of the estimate. Model terms include distance to coast (Dist), upwelling favourable

winds (Up) and downwelling favourable winds (Down). Parameter estimates are deemed important

if the 95% credible interval does not cross the dashed red line which marks an estimate of zero. For

exact estimate values see Table S3.

**Figure 3** Visualisation of the predicted normalised coastal larval abundance based on the generalised

linear mixed model for winds 14 days prior to sampling: a) the effect of distance from the coast,

based upon mean (0 on the standardised scale) upwelling and downwelling favourable winds with

shading showing the 95% credible interval; b) the effect of upwelling favourable winds; and c) the

effect of downwelling favourable winds. Predictions for each variable were made while holding all

other variables within the model to mean values (0 for wind variables and 10km for distance from

the coast). For the winds, 0 represents mean winds, with 1 and −1 representing 1 and −1 SD from the

mean, respectively.

**Figure 4** Caterpillar plot showing the Bayesian parameter estimates for the multi-species catch-per-unit-effort (CPUE) generalised linear mixed model. Model terms include upwelling favourable winds (Up), downwelling favourable winds (Down) and estuary type (Est Type). Points show the estimate of the parameter with the horizontal bar representing the 66% (thick bar) and 95% (thin bar) credible intervals (CI) of the estimate. The dashed red line marks an estimate of zero. The 95% CI does not cross zero for Up Winds, Species: Whiting, Species: Mullet, Drought Months, Estuary Type: Drowned River Valley * Drought Months and Estuary Type: Barrier River * Drought Months. For exact estimate values see Table S4.

**Figure 5** Visualisation of predicted catch-per-unit-effort (CPUE) based upon upwelling favourable

winds during the spawning period as predicted by the results of the CPUE Bayesian linear mixed

model. The grey area represents the 95% credible interval. There were no detectable effects of

downwelling favourable winds and the effects of drought are shown in the supplementary material

(Figure S17). Predictions were made while holding all other variables within the model to mean

values. Note due to the random slope and intercept effects in the model, the actual y scale is

relative.

**Figure 6** Temporal change in annual net displacement in upwelling (from northeast; Estimate of annual change: 52.86, 95% credible interval: 19.00 – 85.59) and downwelling favourable winds (from southeast; Estimate of annual change: -40.93, 95% credible interval: -78.18 – -3.71) between 1850 and 2014. The grey lines show 20,000 posterior sample estimates with the median trend line in blue. Only every third year of annual net displacement data was included in the models to account for temporal autocorrelation in the wind time-series data.