1

Saliva sampling is an excellent option to increase the number of SARS CoV2

diagnostic tests in settings with supplies shortages

Joaquín Moreno-Contreras[1], Marco A. Espinoza[1], Carlos Sandoval-Jaime[1], Marco A. Cantú-Cuevas[2], Héctor Barón-Olivares[3], Oscar D. Ortiz-Orozco[3], Virginia A. Muñoz-Rangel[3], Manuel Hernández-de Jesús[3], César M. Eroza-Osorio[3], Carlos F. Arias[1], Susana López[1]*

[1]Departamento de Genética del Desarrollo y Fisiología Molecular, Instituto de Biotecnología UNAM, Av. Universidad 2001, Col. Chamilpa. Cuernavaca, Morelos; [2]Secretaría de Salud del Edo. de Morelos, Ajusco #2 Col. Buena Vista, Cuernavaca Morelos; [3]Servicios de Salud del Edo. de Morelos. Callejón Borda 3 Col. Centro, Cuernavaca Morelos.

*Corresponding author e-mail susana@ibt.unam.mx

Running title: A simpler and faster method to diagnose SARS CoV-2

Abstract

As part of any plan to lift or ease the confinement restrictions that are in place in many

different countries, there is an urgent need to increase the capacity of laboratory testing for SARS

CoV-2. Detection of the viral genome through RT-qPCR is the golden standard for this test,

however, the high demand of the materials and reagents needed to sample individuals, purify

the viral RNA, and perform the RT-qPCR test has resulted in a worldwide shortage of several of

these supplies. Here, we show that saliva samples treated with a lysis buffer can serve as a

suitable source for viral RNA detection, that is cheaper and can be as efficient as the classical

protocol that involves column purification of the viral RNA. In addition, it skips the need for swab

sampling, decreases the risk of the healthcare personnel involved in this process, and accelerates

the diagnostic procedure.

INTRODUCTION

With the worldwide COVID-19 health emergency, there is an urgent need for rapid and

reliable methods of diagnostic for SARS-CoV-2. The accepted golden standard for detection of

this virus is the amplification of regions of the viral genome by RT-qPCR in nasopharyngeal and

oropharyngeal swabs[1][2]. Unfortunately, given the enormous demand of the reagents needed

to take the biological samples, and to purify the viral RNA, there have been shortages of many of

the reagents needed for the diagnostic tests. Swabs, viral transport media, and kits for viral RNA

extraction are amongst the most common consumables that have become scarce, compromising

the number of tests that can be done in many parts of the world.

Recently, several reports have demonstrated the possibility of using saliva instead of oral

and nasal swabs to detect the genome of SARS-COV-2 [3][4][5]. Saliva collection also has many

collateral benefits, including self-collection, what decreases the risk of healthcare workers in

charge of taking the swabs, and does not require the use of PPE (personal protecting equipment)

that has also become a scarce item in this pandemia [6][7]. In addition, the methods to extract

the RNA from biological samples require the use of purification kits whose availability has also

become limited due to the heavy worldwide demand.

In this report, we compared the RT-qPCR results from 253 paired samples obtained from

saliva and swabs of ambulatory patients; the RNA in the swab samples was extracted using a

commercial RNA purification kit, and the saliva samples were directly mixed with a lysis buffer,

boiled, and used for the RT-qPCR protocol. We found a very good correlation of results between

both types of samples, and propose that this method, which simplifies the sampling of patients,

and accelerates the preparation of the RNA for the RT-qPCR test represents an excellent

alternative that will facilitate to sample and diagnose a larger number of persons at a reduced

cost.

MATERIALS AND METHODS

Sample collection. 253 paired samples from oropharyngeal and/or nasopharyngeal swabs, and

saliva were collected during a span of 30 days (from May 2nd to 31st) by healthcare workers from

the epidemiology department of the health ministry of the state of Morelos (Secretaría de Salud

Morelos, SSM). All but 3 samples, were from ambulatory patients, the 3 exceptions were

collected from hospitalized patients. After their collection, swabs were placed in 2.5 ml of viral

transport medium. Saliva were self-collected by patients that were asked to spit several occasions

into sterile urine cup containers until completing roughly 2-3 ml, both samples were maintained

at 4ºC and transported to the Institute of Biotechnology, UNAM for their analysis, within 24 hours

after sample collection.

Nucleic acid extraction and SARS-CoV-2 detection by RT-qPCR. Total RNA was extracted from

swab samples using the QIAamp viral RNA mini kit (QIAGEN) following the manufacturer´s

protocol, using 140 μl of viral transport medium from each swab and the RNA was eluted in 60

μl of elution buffer.

Saliva samples were treated with the Quick Extract TM DNA Extraction Solution (QE,

Lucigen) by mixing 50 μl of saliva with 50 μl of QE and heating for 5 minutes at 95ºC, the mixture

was then cooled on ice and kept at 4ºC until their use (within 1 hour of QE treatment). In saliva

samples that had high viscosity, 1 volume of sterile phosphate-buffered saline (PBS) was added

and mixed by repeated pippeting, the diluted saliva sample was the extracted as mentioned

above.

SARS-CoV-2 detection was performed using the Berlin protocol, using the reported oligos

and probes for viral gene E and for human RNase P [19]. The RT-qPCR reactions were performed

using the StarQ One-Step RT-qPCR (Genes 2 Life) kit, using 5 μl of the column extracted total RNA

in a 20 μl of reaction mix, or 2.5 μl of the QE treated saliva into 22.5 μl of reaction mix. Samples

were analyzed in an ABI Prism 7500 Sequence Detector System (Applied Biosystems) with the

following thermal protocol: 50ºC for 15 min, 95ºC for 2 min and then 45 cycles of 95ºC for 15 s

and 60ºC for 30 s. All samples with a Ct value equal or less than 38 were classified as positive.

Determination of viral copy number. To determine the viral copy number in a sample, a standard

curve was generated using a 10-fold serial dilution of an *in vitro* T7 RNA transcript that encodes

the sequence recognized by oligonucleotides and probe for gene E. Briefly, the logarithm of

concentration of each dilution was plotted against the Ct and the viral copy number from

unknown samples was determined by extrapolating the Ct value onto the corresponding

standard curve.

Statistical analysis. Statistical analysis was performed using GraphPad Prism 6.0 (GraphPad

Software Inc.) as described in the results section.

RESULTS

Viral titer determination in swabs and saliva. To evaluate if saliva is a good source of viral RNA

for the RT-qPCR tests we determined the presence of the SARS-CoV-2 genome in paired saliva

and swab samples from 253 ambulatory patients. All patients had two or more symptoms related

to COVID-19. All patients (100%) had two or more symptoms related to COVID-19, 115 (45.4%)

were male and 137 (54.1%) female, with a median age of 41 (+/- 14.4) years (Table 1)[8][9].

Samples were taken from ambulatory patients in the respiratory triage of the Tlaltenango health

Center, in Cuernavaca, Morelos. The RT-qPCR Berlin protocol was used to detect SARS-CoV-2,

using only the primers and probe for gene E, since previous studies have shown a weak detection

of viral RNA when the RdRp gene is probed [10][11]. As an internal control of RNA content in the

samples, the RNase P gene was detected. Total RNA was purified from swabs using the QIAamp

viral RNA mini kit, the RNA in saliva was directly obtained using the QE lysis buffer (Lucigen) and

boiled for 5 minutes, as reported [12].

From the 253 samples analyzed, 139 (54.9 %) were negative for the presence of SARS-

CoV-2 both in swab and saliva samples, and 114 (45.1%) were positive for SARS-CoV-2 in at least

one type of sample. Out of the 114 positive samples, 60 (52.6%) were positive both in swab and

saliva, while 34 (29.8%) were positive only in saliva, and 20 (17.5%) were detected only in swabs

(Figure 1). According to these results, saliva samples allowed the detection of 82.4% of total

positive samples, while through swabs only 70.1% positives were detected.

Quantitation of viral RNA. When the number of viral copies in the positive paired samples were

compared, a significant difference between the geometric mean of the viral copy number in saliva

and swab samples was detected, with saliva samples having a 1.2 $\log_{10}$ higher titer than that

observed in swabs ($p<0.005$, Figure 2A). This can be better appreciated when the viral copy

number in paired swabs and saliva from the same patient, is plotted and represented as

connecting lines (Figure 2 B); in 42 of the paired samples the number of viral copies was higher

in saliva samples than in swabs. Human RNase P was used as an internal control of sampling

quality; the comparison between the mean of Ct values obtained from swabs and saliva showed

a difference of at least 7.6 Ct's between both types of samples (Figure 2 C), indicating that there

is more cellular material in saliva, as reported in other studies[13].

Taken together these results suggest that that saliva is a good source of SARS-CoV-2, and

can be used for diagnostic tests using QE extraction buffer in place of the column-based method.

The higher virus titer as well as the higher level of RNase P detected in saliva as compared to

swabs, support this conclusion.

DISCUSSION

In this study we analyzed 253 paired samples from pharyngeal and/or nasopharyngeal

swabs and saliva. RNA purified from swabs using commercial column kits was compared with

saliva treated by direct lysis using QE buffer and surpassing the RNA extraction protocol. More

positive samples (82.4%) and a higher viral copy number (1.2 $\log_{10}$) were detected in saliva as

compared to the paired nasopharyngeal swab (70.1%) samples. In this regard, the number of

false negatives in swabs was almost twice (29.8%) as those in saliva samples (17%).

The failure of identification of SARS-CoV-2 in swab samples when the saliva samples were

positive for the virus could be due to bad swab sampling, what can be corroborated by the high

Ct values of RNase P detected in these samples (Figure 2 C), with the consequent low viral copy

number. This is a major concern, since for natural reasons the medical personnel in charge of

taking the samples frequently do not do it correctly for the risk associated with this process. Of

note, most (28/34) of the samples that were positive for saliva and negative for swabs

corresponded to swab samples that were taken only from the oropharinx (single swab). It has

been reported that nasopharyngeal swabs have a higher viral titer compared with oropharyngeal

swabs [2]; thus, this could contribute to the discrepancies observed. In addition, it has also been

previously reported that nasopharyngeal swabs have a lower viral titer than saliva samples [13],

what can also contribute to explain our findings. On the other hand, the false negatives in saliva

could be due to problems during the collection, transport and or storage of the sample before its

arrival to the laboratory.

SARS-CoV-2 has been detected in saliva at higher titers during the first days after the

onset of symptoms, with the viral titer declining over time. It is not clear how long after the

symptom onset the viral RNA can be detected in saliva, although some reports suggest a short

period of detection (~13 days) as compared with nasopharyngeal swabs (~19 days) [14].

However, other reports have recently demonstrated the detection of viral RNA in saliva for longer

periods of time (~20 days or longer) [3][15]. The patients included in this study were ambulatory,

and according to their clinical interview had between 1 and 7 days, (with a median of 4 days) of

the onset of symptoms. We did not find a significative difference between the onset of symptoms

and the results obtained from samples that were only positive in saliva versus those that were

only detected in swabs.

Direct lysis of nasopharyngeal or oropharyngeal swab samples in viral transport medium

using the QE buffer has been reported as a suitable method for direct qRT-PCR for SARS-COV-2

detection, with rates similar to methods based on column purification [16][12]. However, in our

experience we have found a great variability in the results obtained using the QE lysis protocol

when applied to swab samples, most likely due to variations in the material of the swabs used

and to variations in the preparation of the viral transport medium employed (data not shown).

In this regard, it has recently been reported that the composition of viral transport media can

affect the detection of viral RNA from SARS-CoV-2 and other viruses [17] and, due to the scarcity

of it, several laboratories have started to prepare their own transport media introducing an

additional confusion factor. A similar situation occurs with the swabs, since in view of the scarce

suitable materials, other materials are being employed, despite the fact that some materials are

known to inhibit the RT-PCR reactions [18].

The use of saliva samples offers the advantage that, in general, saliva has a constant

composition and no additives or transport media need to be used for their preservation if stored

in cold and analyzed up to 36 h after their collection. Our results indicate that a rapid processing

of saliva using direct lysis with QE buffer, and avoiding an RNA extraction protocol, offers an

excellent alternative to the current swab analysis that uses RNA column purification, since it is a

sensitive, fast and cheap method that can be used for massive screening, in particular in those

settings where common supplies needed for the classical methods are in shortage.

ACKNOWLEDGMENTS

We are grateful to the healthcare workers of Servicios Estatales de Salud de Morelos for their

invaluable help in collecting the samples, and to the personnel of the Laboratorio Estatal de Salud

Pública del Edo de Morelos, for their support in the preparation and transporting of the samples.

This work was supported by grant 314343 from CONACyT.

FIGURE LEGENDS

Figure 1. Comparation of SARS-CoV-2 positive samples in swab and saliva. Percent number(n)

of positive samples (+) and negative samples (-) detected in swabs and saliva as indicated.

Figure 2. SARS-CoV-2 is detected at higher titer in total positive saliva samples. A) Viral titer

(viral copies/ml) from all positive samples detected in swabs (n=94) and in saliva (n=80); B) Viral

titer (viral copies/ml) in positive swab and saliva samples (n=60) were represented by lines

connected between both samples. Data were compared by a Wilcoxon test (p<0.005); C) RT-PCR

cycle threshold Ct values for RNase P. Data were compared by Mann-Whitney test (p< 0.005 and

p<0.0001). Bars represent the geometric median and 95% CI.

196
197

REFERENCES

[1]   Wang Y, Kang H, Liu X, Tong Z. Combination of RT-qPCR testing and clinical features
        for diagnosis of COVID-19 facilitates management of SARS-CoV-2 outbreak. *J Med
        Virol*. 2020;92(6):538-539. doi:10.1002/jmv.25721.

[2]   Pan Y, Zhang D, Yang P, Poon LLM, Wang Q. Viral load of SARS-CoV-2 in clinical
        samples. *Lancet Infect Dis*. 2020;20(4):411-412. doi:10.1016/S1473-3099(20)30113-4.

[3]   To KK, Tsang OT, Chik-Yan Yip C, et al. Consistent detection of 2019 novel coronavirus
        in saliva. *Clin Infect Dis*. 2020;ciaa149. doi:10.1093/cid/ciaa149.

[4]   Williams E, Bond K, Zhang B, Putland M, Williamson DA. Saliva as a non-invasive
        specimen for detection of SARS-CoV-2. *J Clin Microbiol*. 2020;JCM.00776-20.
        doi:10.1128/JCM.00776-20.

[5]   Kojima N, Turner F, Slepnev V, et al. Self-Collected oral fluid and nasal swabs demonstrate
comparable sensitivity to clinician collected nasopharyngeal swabs for Covid-19 detection.
medRxiv 2020.04.11.20062372; doi: https://doi.org/10.1101.

[6]   Ng K, Poon BH, Kiat Puar TH, et al. COVID-19 and the Risk to Health Care Workers: A
Case Report. *Ann Intern Med*. 2020;172(11):766-767. doi:10.7326/L20-0175.

[7]   Khurshid Z, Asiri FYI, Al Wadaani H. Human Saliva: Non-Invasive Fluid for Detecting
Novel Coronavirus (2019-nCoV). *Int J Environ Res Public Health*. 2020;17(7):2225. Published
2020 Mar 26. doi:10.3390/ijerph17072225.

[8]   Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel
        coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506. doi:10.1016/S0140-
        6736(20)30183-5.

[9]   Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With
        2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China.
        JAMA.2020;323(11):1061-1069. doi:10.1001/jama.2020.1585.

[10]   Nalla AK, Casto AM, Huang MW, et al. Comparative Performance of SARS-CoV-2
        Detection Assays Using Seven Different Primer-Probe Sets and One Assay Kit. *J Clin
        Microbiol*. 2020;58(6):e00557-20. doi:10.1128/JCM.00557-20.

[11]   Ramírez JD, Muñoz M, Hernández C, et al. Genetic diversity among SARS-CoV2 strains in
        South America may impact performance of Molecular detection. medRxiv
        2020.06.18.20134759; doi: 10.1101/2020.06.18.20134759.

[12]   Ladha A, Joung J, Abudayyeh OO, et al. A 5-min RNA preparation method for COVID-19
        detection with RT-qPCR. medRxiv 2020.05.07.20055947; doi:
        10.1101/2020.05.07.20055947.

[13]   Wyllie AL, Fournier J, Casanovas-Massana A, et al. Saliva is more sensitive for SARS-CoV-2
        detection in COVID-19 patients than nasopharyngeal swabs. MedRxiv
        2020.04.16.20067835; doi:10.1101/2020.04.16.20067835.

[14]   Iwasaki S, Fujisawa S, Nakakubo S, et al. Comparison of SARS-CoV-2 detection in
        nasopharyngeal swab and saliva. *J Infect*. 2020;S0163-4453(20)30349-2.
        doi:10.1016/j.jinf.2020.05.071.

[15]   To KK, Tsang OT, Leung WS, et al. Temporal profiles of viral load in posterior
        oropharyngeal saliva samples and serum antibody responses during infection by SARS-
        CoV-2: an observational cohort study. *Lancet Infect Dis*. 2020;20(5):565-574.
        doi:10.1016/S1473-3099(20)30196-1.

[16] Sentmanat M, Kouranova E, Cui X. One-step RNA extraction for RT-qPCR detection of
2019-nCoV. bioRxiv 2020.04.02.022384; doi:10.1101/2020.04.02.022384.
[17] Kirkland PD, Frost MJ. The impact of viral transport media on PCR assay results for the
detection of nucleic acid from SARS-CoV-2 and other viruses. bioRxiv 2020.06.09.142323;
doi:10.1101/2020.06.09.142323.
[18] Hedman J, Rådström P. Overcoming inhibition in real-time diagnostic PCR. *Methods Mol*
Biol. 2013;943:17-48. doi:10.1007/978-1-60327-353-4_2.
[19] V. M. Corman *et al.*, "Detection of 2019 -nCoV by RT-PCR," *Euro Surveill*, vol. 25, no. 3,
pp. 1–8, 2020.
252
253
254

Figure 1

318 Figure 2

TABLE 1. Patient data

	Total paired samples (253)*
Male	115 (45.4%)
Female	137 (54.1%)
Age mean	41 +/- 14.4 years
Hospitalized	3 (1.1%)
Ambulatory	229 (90.5%)

*Data from some patients were not obtained