HUMAN PAPILLOMAVIRUS TYPE 16 SUB-LINEAGES AND INTEGRATION IN CANCER

Robert Jackson[1,2,*,#], **Alejandro Ortigas-Vásquez**[2,3], **Ingeborg Zehbe**[2,4,5]

[1]Biotechnology Program, Lakehead University; [2]Probe Development & Biomarker Exploration,

Thunder Bay Regional Health Research Institute; [3]Bioinformatics Program; [4]Department of

Biology; [5]Northern Ontario School of Medicine, Lakehead University, Thunder Bay, Canada

[*]**Corresponding author:** Robert Jackson, PhD (e-mail: rjackso1@lakeheadu.ca)

[#]**Affiliation history:** Robert Jackson was affiliated with Lakehead University and the Thunder Bay

Regional Health Research Institute at the time the study was performed and is currently affiliated

with the University of Arizona.

KEYWORDS

The Cancer Genome Atlas (TCGA), Human papillomavirus type 16 (HPV16), Sub-lineage,

Variants, Integration, Cervical cancer, Head and neck cancer

SHORT TITLE

HPV16 sub-lineage integration

ABSTRACT

Our lab has been intrigued by the fact that viral genomes often take on the role of mobile elements to perpetuate their existence in a complex organism's genome. Multiple DNA viruses such as Epstein-Barr virus, hepatitis B virus, and human papillomavirus (HPV) can invade their host genome, as "genomic parasites". We have been investigating HPV type 16 (HPV16), which is a prominent human tumour virus. In our recent *in vitro* work using 3D organoids, a common variant of HPV16's coding region elicited early integration into the host genome compared to the HPV16 prototype sequence. Next-generation sequencing (NGS) data confirmed a transcriptomic profile of increased proliferation and chromosomal instability—both hallmarks of cancer. Epidemiologically, this variant is associated with a high cervical cancer incidence. To take inquiries a step further, we investigated variant-specific integration across HPV16-related cancers using NGS data from population-derived clinical samples in The Cancer Genome Atlas (TCGA)-curated database. Data were analyzed for HPV16 positivity, sub-lineage, and viral-host integration using a bioinformatic pipeline of open-source tools, including HPVDetector. Here, we report the analysis of 120 cervical cancer cases comprising HPV16 positive and negative samples as well as their different sub-lineage and integration status. The integration signature between variant and prototype did not differ quantitatively but qualitatively: that of the variant being related to hypoxia/energetics (Warburg effect) and that of the prototype being much more varied to include host immune abrogation and cancer pathways activation. We conclude by discussing challenges and future directions for expanding these analyses.

INTRODUCTION

The co-evolutionary interplay between pathogens and their hosts involves a plethora of intriguing biological phenomena, including genomic integration of pathogen DNA into host cells. While cellular integration of viral genomic material is a means of propagation for families such as

Retroviridae (*e.g.*, human immunodeficiency virus, HIV), it can also occur during persistent

infections with DNA viruses and is often associated with virally induced cancers. One such family

of common DNA tumour viruses, *Papillomaviridae*, is responsible for causing ~5% of all human

cancers worldwide [Ghittoni *et al.*, 2015]. There are more than 600 types of papillomaviruses

(PVs) discovered as of July 2021 based on the PaVE reference genome database [PaVE:

https://pave.niaid.nih.gov, Van Doorslaer *et al.*, 2013; 2017a]. The number of PV types, including

the non-human PVs (currently 226) is expected to continue rising due to high-throughput

genomics and sampling [Van Doorslaer and Dillner, 2019]. The majority of PVs currently identified

are the human papillomaviruses (HPVs) [Van Doorslaer *et al.*, 2017b] and these types (448 to

date) differ in their propensity for inducing cancer as well as their epithelial tropism, with "high-

risk" types being the primary cause of malignancies: predominantly via sexually-transmitted

persistent HPV type 16 (HPV16) infection of anogenital and oropharyngeal mucosa [Ndiaye *et*

al., 2014]. Although this and other HPV genomes are considered "small", only ~7.9 kb containing

nine open reading frames (ORFs) and a variety of alternatively spliced and multi-cistronic variant

transcripts [Graham & Faizo, 2017], it encodes potent oncoproteins such as E6 and E7 which

drive host cells to acquire the hallmarks of cancer [Hanahan and Weinberg, 2000; 2011; Mesri *et*

al., 2014]. The combined E6 and E7 actions disable protective apoptotic mechanisms and

promote host cell proliferation as outlined by Hoppe-Seyler and colleagues [2018]—a

phenomenon that could potentially be exploited therapeutically at an RNA [Togtema *et al.*, 2018]

or protein-level [Togtema *et al.*, 2019]. Of particular relevance for this study, the HPV-augmented

environment can permit chromosomal instability (an enabling characteristic among the hallmarks

of cancer) and eventual integration of HPV16 sequences into human DNA via double-strand break

(DSB) repair events [Winder *et al.*, 2007], yielding unique integration signatures found in next-

generation sequencing (NGS) data from human carcinomas [Holmes *et al.*, 2016].

HPV16 sub-lineages (variants) via epidemiological and lab-based studies have been reported to infer differing cancer risk [reviewed by Mirabello *et al.*, 2018 and references within; Clifford *et al.*, 2019]. HPV16 variant designations were originally based on their geographical region of origin and have since been updated to lineage (1.0 to 10.0% whole viral genome sequence difference) and sub-lineage (0.5 to 1.0% difference) nomenclature. The European "prototype" sub-lineage (EP; now A1 according to new nomenclature) was the first HPV16 genome published [Seedorf *et al.*, 1985]. The remaining sub-lineages include two additional European groups (E; now A2 and A3), one Asian (As; now A4), eight African (Af-1a, -1b and -2; now B1-4 and C1-4), two North-American (NA; now D1 and D4) and two Asian-American (AA-1 and -2; now D3 and D2, respectively) [Burk *et al.*, 2013; Mirabello *et al.*, 2018]. We have put our efforts toward comparing the EP and AA sub-lineage variants, which differ by only three amino acid changes at residues 14 (Q to H), 78 (H to Y) and 83 (L to V) in the major transforming protein E6 [Richard *et al.*, 2010; Niccoli *et al.*, 2012; Jackson *et al.*, 2014; 2016; Togtema *et al.*, 2015; Cuninghame *et al.*, 2017]. Epidemiological studies revealed that the AA sub-lineage infers a higher risk for dysplasia and an earlier onset of invasive tumours than EP [Xi *et al.*, 1997; Berumen *et al.*, 2001]. Our functional assays showed that AAE6 has a greater transforming, migratory, and invasive potential than EPE6 when the respective E6 gene was retrovirally transduced into primary human keratinocytes in long-term *in vitro* immortalization studies [Richard *et al.*, 2010; Niccoli *et al.*, 2012; Togtema *et al.*, 2015]. These observations may be due, at least in part, to an AAE6-mediated altered metabolic phenotype reminiscent of the Warburg effect [Richard *et al.*, 2010; Cuninghame *et al.*, 2017]. Consistent with the hypothesis that AAE6 is more oncogenic than EPE6, our recent experimental work of early HPV16 carcinogenesis revealed that AAE6 in the context of the full-length viral genome is more prone to host genome integration than is EPE6 [Jackson *et al.*, 2014; 2016]. At that time, one study mentioned HPV16 sub-lineage variants and integration but did not find a difference (P = 0.28, two-tailed Fisher's exact test) between EPE6 (3

episomal and 20 integrated cases) and the E-T350G variant (6 episomal and 16 integrated cases,

responsible for one of the residue changes also found in AAE6: L83V) [Xu *et al.*, 2013]. Notably,

however, the lone tumour sample that contained the AA variant was in integrated form.

Based on our assumption that the propensity for HPV16 integration may vary between

sub-lineages, we set out to do a search of clinical samples with available genomic and

transcriptomic data reported in databases. Two landmark publications of The Cancer Genome

Atlas (TCGA) reported on cervical squamous cell carcinoma and endocervical adenocarcinoma

(CESC) [TCGA Research Network, 2017] and head and neck squamous cell carcinoma (HNSC)

[TCGA Research Network, 2015]. More recently, Cantalupo *et al.* [2018] reported further analysis

of additional TCGA data, including additional cancer sets that have become available in recent

years. They also used a clearly defined, rules-based approach to filtering and detection: first

extracting high-quality non-human reads followed by removal of non-relevant viral families

(bacteriophage, plant viruses, insect viruses) and artifacts, with 10 or more high-quality

alignments required for virus detection. Prior to the primary TCGA data analysis papers, there

was a publication about DNA viruses in TCGA RNA-Seq data across a variety of human cancers

[Khoury *et al.*, 2013]. An additional literature search found several more related papers: HPV and

HNSC comprehensive analysis [Castellsagué *et al.*, 2016] and an HPV16-focused follow-up

TCGA-HNSC analysis [Nulton *et al.*, 2017]. In the latter, three different groupings of integration

status were reported: episomal, integrated, and human-viral episomal hybrids, challenging

integration data by Parfenov *et al.* [2014]. The Parfenov *et al.* study disclosed HPV typing and

sub-lineage analyses of the 35 HPV+ HNSC cases with 29 containing HPV16: one of which was

an AA variant. Interestingly, integration with 16 breakpoints in the human genome (*i.e.*, ~ 4-fold

more compared to EP) was shown. In the Nulton *et al.* study, for the same case, a new

phenomenon of human-viral hybrids in the host genomic DNA subsequently found in episomal

form was reported [Nulton *et al.*, 2017]. Recently, Zapatka *et al.* [2020] further analyzed TCGA

data to determine viral associations with cancer, including the impact of integrations on host copy

number variations. Most recently, Lou *et al.* [2020] report that integration rates indeed differ

between sub-lineages in cervical cancers. In a Guatemalan population with a 24%, 31%, and 45%

prevalence of D2, D3, and A1 sub-lineages, respectively, they found that D2 had a higher rate of

integration than D3 (76% vs 44%), coupled with an earlier age of cancer diagnosis. Interestingly,

they also found that A1 had an equally high rate of integration (78%) as D2, which is in contrast

with our *in vitro* data (at least, within this specific population) [Jackson et al., 2016]. It is noteworthy

that A1 is phylogenetically distant from D2/D3, with ~2% genome difference (>150 single-

nucleotide polymorphisms, SNPs, and insertions/deletions, indels), whereas D2 and D3 differ

from each other by only ~0.5% (<40 SNPs and indels, and hence it remains open what the

biological meaning for these findings may be).

Our overall objective was to screen TCGA for all HPV16-containing samples, verify their

sub-lineage genotypes, then detect and characterize instances of viral integration along with a

complete picture of its impact on the host genome and identify potential contributing factors in the

surrounding molecular landscape. The most pertinent research question addressed here is

whether there are specific integration mechanisms associated with different HPV16 sub-lineage

genotypes: is the AA variant (D2 and D3) more frequently associated with integration than the EP

variant (A1) and/or are there unique integration patterns with potential functional relevance? Such

findings would resonate with our previous variant-specific integration work [Jackson *et al.*, 2014;

2016] and provide evidence for a novel mechanism in differing variant tumourigenesis. Our

approach of HPV16 sub-lineage genotyping and integration detection, along with an attempt to

reconcile and verify existing data, has not been previously addressed for TCGA data. We

analyzed TCGA cervical cancer cases (n = 120), including DNA- and RNA-level data for each (n

= 240 total samples). We did not find significantly more or less integration events between AA

and EP samples but differences in the integration pattern. AA—in a singular manner—seems to

adapt host cells to a hypoxia/energetics (Warburg effect)-associated phenotype whereas EP

alters host cells to various hallmarks of cancer including host immune surveillance and cancer

pathway signaling. A variety of bioinformatics pipelines exist for sub-lineage genotyping and

integration detection. These are however susceptible to biological (*e.g.*, low copies and

incomplete viral coverage) and technical (*e.g.*, inadequate read depth) hurdles, which are

important considerations for future analyses.

RESULTS & DISCUSSION

HPV16 D2/D3 integration data are limited for cervical and head and neck cancers

When seeking to test the relationship between HPV16 sub-lineage and integration

propensity we started by compiling a sample summary of previously published and available data

from TCGA [**Table 1**]. Next, we performed a preliminary sub-lineage and integration analysis for

available CESC sample data [**Figure 1**] as it was stratified into sub-lineage for the A lineage,

whereas HNSC data were not further stratified between A1, A2, and A3. Overall, while the

proportion of integrated cases was 1.23x higher in D2/D3-containing CESC samples (9/10, 90.0%

vs 51/70, 72.9% for A1), the sample size was very low rendering this initial analysis underpowered

and the finding as not statistically significant ($P = 0.242$, X^2 test).

Table 1. Sample summary breakdown for each TCGA case set. Sample genotypes for cancer

cases from the Cervical Squamous Cell Carcinoma and Endocervical Adenocarcinoma (CESC)

as well as Head and Neck Squamous Cell Carcinoma (HNSC) datasets. CESC HPV16 data are

based on TCGA Research Network [2017] (using the sub-lineages identified at the time, which

were only 10 of the current 16). *HNSC HPV16 data are based on Parfenov *et al.* [2014], where

the A sub-lineage was grouped as "EUR" genotypes without specifying A1, A2 or A3.

Sample genotype	CESC (*n* = 178)	HNSC (*n* = 279)
Total HPV16 cases	103 (57.9% of CESC cases)	29 (10.4% of HNSC cases)
A1	70 (68.0% of HPV16 cases)	
A2	12 (11.7%)	22 (75.9% of HPV16 cases)*
A3	4 (3.9%)	
A4	4 (3.9%)	2 (6.9%)
B1	2 (1.9%)	3 (10.3%)
B2	–	–
C	1 (1.0%)	1 (3.4%)
D1	–	–
D2	2 (1.9%)	–
D3	8 (7.8%)	1 (3.4%)

Figure 1. Preliminary analysis of CESC samples. Previously available HPV16 sub-lineage and integration data for the CESC dataset, published by the TCGA Research Network [2017], were analyzed to determine whether there was a higher proportion of integrated samples

corresponding to D2/D3 (which have viral genomes that differ from each other by ~0.5%, but have

identical E6 amino-acid sequences) vs the A1 sub-lineage (which differs ~2% from D2/D3 viral

genomes). While these sub-lineage classifications were based on 10 reference genomes, the set

has since been expanded to 16 sub-lineages, as described in Mirabello *et al.* [2018].

The untapped potential of TCGA

To expand upon these preliminary results, we were using our own HPV16 detection and

sub-lineage analyses to verify sub-lineage calls by prior studies, resolve discrepancies, and

further analyze the "unknown" samples. From the National Cancer Institute's Genomic Data

Commons (GDC) Data Portal, there are currently 835 relevant CESC (n = 307) and HNSC cases

(n = 528) available for analysis, including 2,568 binary alignment map (BAM) files (27.84 TB total,

an average of 33.34 GB/case, including all whole-exome sequencing (WXS) and RNA-Seq

sequencing read files and excluding miRNA-Seq files as they are not primarily relevant for HPV16

genotyping). There are 307 CESC cases (924 BAM files, 10.44 TB) and 528 HNSC cases (1,644

BAM files, 17.41 TB), with both sets including samples from primary tumours, metastases, normal

blood, and normal solid tissues. It appears the TCGA legacy data portal also contains whole-

genome sequencing (WGS) files for these cancer sets. Based on the TCGA landmark studies

[TCGA Research Network, 2015; 2017], we expect the majority of CESC samples are HPV

positive, whereas a smaller proportion of HNSC samples are HPV positive. While HPV16 is

expected to be the predominant type, multi-type infections and co-infections with other non-

commensal microorganisms are also possible. As with most nucleic-acid based techniques,

cross-contamination between samples is a potential source of confusion, especially for adjacent

cases where sequence libraries could be contaminated [Cantalupo *et al.*, 2015].

The number of CESC and HNSC samples that were previously marked as being positive

for HPV16 differ between studies, likely due to differences in analysis pipelines (*e.g.*, data types

analyzed, thresholding, and filtering for contamination) [TCGA Research Network, 2017;

Cantalupo *et al.*, 2018]. While both RNA-Seq and WXS were used for integration analysis in the

landmark CESC study, it seems that the authors mostly if not totally relied on RNA-Seq [TCGA

Research Network, 2017]. Moreover, for integration status analysis, an in-house tool was used,

which cannot be directly compared with other published and available tools such as ViFi [Nguyen

et al., 2018], HPVDetector [Chandrani *et al.*, 2015] or VirTect [Xia *et al.*, 2019] (and others detailed

in the **METHODS** section). This poses major challenges to a current, valid analysis of sub-lineage

association with integration based on these previous publications.

Figure 2. Flowchart summary of all 240 samples analyzed. The 240 samples analyzed were

broken down based on positivity (based on RPM), lineage and sub-lineage (based on reference

genome with most mapped reads). Samples mapping equally to more than one reference genome

were grouped under "Undefined".

HPV16 positivity quantification reveals discrepancy between RNA-Seq and WXS samples

We analyzed both RNA-Seq and WXS CESC samples independently to detect the

presence of HPV16 in a subset of 120 cases (240 samples, **Figure 2**) representing all 103

available HPV16 positive cases identified in the landmark study as well as 17 negatives to

determine how robust our methods are [**Figure 3A**]. As expected, control sample SiHa was a low-

positive (5.51 VRPM) and CaSki was a high-positive (502.66 VRPM). TCGA RNA-Seq samples

had a range of 0.00 to 1,182.78 VRPM, whereas WXS samples ranged from 0.00 to 109,994.41

VRPM. In total, 85/120 (70.8%) of the cases we analyzed were HPV16 positive based on RNA-Seq and used for subsequent sub-lineage and integration analysis. However, for the same cases, DNA and RNA level HPV16 quantification were often inconsistent, with only 67/120 (55.8%) cases yielding consistent HPV16 quantification (negative, low-positive, or high-positive).

To resolve these discrepancies, we compared our data for both RNA-Seq and WXS samples to data from the landmark CESC study [TCGA Research Network, 2017], summarized in [**Figure 4**]. We found that positivity results for RNA-Seq samples were the most consistent with previous findings, as all cases that had been reported as negative for HPV16 were also found negative in our analysis at the RNA level, but not necessarily at the DNA level, where we found 7 WXS samples above the low-positive threshold (2 of which are even above the high-positive threshold). Interestingly, 9/10 (90.0%) of the D lineage samples were positive at both the DNA and RNA level, whereas only 62/94 (66.0%) of the A lineage samples analyzed were positive at both DNA and RNA levels. These discrepant findings could be due to biological or technical phenomena, such as insufficient read depth, too few viral reads (small amount of viral DNA in a subset of cells, relative to host DNA), or no/weak transcription of integrated viral sequences (in the cases where RNA-Seq signal is lacking). These issues could be further compounded by computational artefacts and potential contaminating viral sequences, which makes it essential (yet challenging) to develop tools that successfully distinguish contaminating from authentic viral sequences [Cantalupo *et al.*, 2018].

Overall, for this set of 120 cases there was a weak positive correlation between DNA and RNA HPV16 levels (R^2 = 0.3205) [**Figure 3B**]. Lineage-specific HPV16 levels were compared for DNA and RNA, with both similar between A and D lineages [**Figure 3C**]. The majority of HPV16 positivity was attributable to the A lineage [**Figure 3D**].

Figure 3. HPV16 positivity in RNA-Seq and WXS TCGA samples. (A) A total of 240 samples

were analyzed to determine whether they were positive for HPV16, with 120 corresponding to

RNA-Seq and 120 corresponding to WXS samples. Samples with less than 0.99 viral reads per

million (RPM) (marked by the lowest dashed line) were classified as negative. Positive samples

were then subdivided based on strength of positivity, with RPM values below 99 (marked by the

highest dashed line) indicating low positivity and those above 99 indicating high positivity. TCGA

case IDs are colour-coded based on their CESC landmark study lineage determination (A = green,

B = purple, and D = blue). Cervical cancer control cell lines, SiHa and CaSki (shaded grey), were

included as low- and high-positive controls, respectively. Values are inverse hyperbolic sine (IHS)

transformed. **(B)** Correlation plot of RNA vs DNA-level HPV16 VRPM. **(C)** Sina plot [Sidiropoulos

et al., 2018] of VRPM for HPV16 lineage A vs D. **(D)** HPV16 lineage pie chart, for the 87 HPV16

positive (RNA-level) cases analyzed.

Figure 4. Flowchart summary of all 120 cases analyzed. The 120 cases analyzed were broken down based on positivity [depending on whether both samples were positive (+/+), one was positive and the other negative (+/-), or whether both were negative (-/-)], lineage [defined by both corresponding samples giving a matching lineage call] and integration [based on number of repeated integration reads].

HPV16 lineage, but not sub-lineage, was consistent between RNA-Seq and WXS

Our genotyping results were highly consistency among RNA-Seq and WXS samples at the lineage level, with all sample pairs with matching positivity calls also possessing matching lineage calls. However, of these cases, only 7 matched at the sub-lineage level (1 match for D2 and 6 matches for D3). No exact matches were found at the sub-lineage level for samples of the A or B lineages. This could be due to the high degree of sequence similarity between members of the A lineage as opposed to those belonging to the D lineage, making the sub-lineage determination dependent on a higher degree of difference between references. In the case of WXS sample TCGA-IR-A3LA, our genotyping algorithm was not capable of distinguishing between sub-lineages A2, A3 and A4, indicating equal identity for all three sequences based on mapping statistics. The final lineage distribution of the 72 cases positive for HPV16 can be seen in **Figure 4**. For our controls, SiHa and CaSki were both within A lineage, with SiHa identified as A1/A3 and CaSki as sub-lineage A2. Overall, our results are consistent with others reporting that sub-lineage identification is challenging, with difficulty in confidently assigning a single sub-lineage among highly similar members of the same lineage [Dawson *et al.*, 2019].

An important consideration is that accurate sub-lineage identification can be negatively

impacted by integration itself, due to truncation of viral genomic regions which may decrease the

amount of genomic space available for alignment. Also, it is possible that some integration events

may render the viral genome transcriptionally inactive, but still tumourigenic due to insertional

mutagenesis (*e.g.*, disrupting a tumour suppressor). In this theoretical example, DNA-level high-

throughput sequencing (*e.g.*, WGS and WXS) would be required to resolve the integration site,

but low sequencing depth and a low abundance of virus relative to host DNA could yield false-

negatives for viral positivity, sub-lineage identification, and integration. This is a limitation with

TCGA datasets. Viral sequence capture enrichment could address this concern, so long as

probes are designed to target across the entire viral genome and flanking host regions.

289   *The proportion of HPV16 integrated reads does not differ between lineages*

Initial raw output indicated the presence of viral integration sites in all samples that had

been found positive for HPV16 in earlier steps of the pipeline, suggesting a higher than anticipated

false-positive rate. While it is possible these could represent a low proportion of truly integrated

reads, we decided that by considering the sequencing depth of each sample we could obtain a

more reliable metric in the form of percentage of integrated reads [**Figure 5A**]. Samples were

then filtered based on this value and compared to findings from the landmark CESC study [TCGA

Research Network, 2017]. Once again, results from RNA-Seq samples were the most consistent

with previous reports, indicating presence of integrated HPV16 [**Figure 5A**]. In this case, the

proportion of integrated reads was higher in RNA-Seq vs WXS samples. The HPV16 positive

cervical carcinoma controls resulted in positive integration for SiHa, while CaSki was considered

negative with the current methodology. Clearly, CaSki's tandemly repeated full genomes pose a

problem for detection and represent a false-negative, highlighting the importance of

complimentary techniques and a careful biological interpretation and analysis. Integrated reads

 (%) in HPV16 lineages A vs D were compared and indicate no difference at the RNA or DNA level

[Figure 5B], yet a high amount of variability among the HPV16 positive samples.

Figure 5. Integrated HPV16 reads in RNA-Seq and WXS samples. (A) For each of the HPV16

positive samples, the total number of integrated reads reported by HPVDetector were converted

to a percentage of the total number of viral reads. Samples with 1.14% or more integrated reads

(marked by the dashed line) were classified as positive for HPV16 integration. Samples below

this threshold were classified as predominately containing the episomal form of HPV16. SiHa and

CaSki controls were also included, as they are both known to have integrated HPV16. **(B)**

Proportion of integrated reads for RNA-Seq and WXS samples comparing HPV16 lineages A and

D as a Sina plot [Sidiropoulos *et al.*, 2018].

Integrated genes and known associations with carcinogenesis

Qualitative analysis criteria are based on 63 A lineage and 9 D lineage cases with at least

10 reads in either DNA or RNA-Seq or both [**Figure 4, 6**]. For the A lineages, 28 cases showed

corresponding RNA and DNA samples, and for the D lineages, 3/4 cases corresponded, albeit in

the 4th case corresponding DNA was found but only with 3 hits. For the A lineages the 31 affected

genes were *TAOK3, SUSD1, MIR3134, IKZF3, PLCD3, MAP3K14, IGFBP6, PALLD, VPS13D,*

IL9R, NFIA, MPRIP, UBA6, UTP11L, DYRK1A, CEACAM5, RXRA, MIPOL1, FLNB, CTSE,

FAM220A, *ATXN10*, *RNF166*, *GRIK4*, *RAD51B*, *FHAD1*, *C6orf203*, *SIPA1L3*, *MACROD2*,

PGAP3, *EPHA3*. There are close to 143 genes with more than 10 hits in the DNA-Seq samples

with either no or less than 10 RNA-Seq hits, *e.g.*, *TP73*, *DLG*, *MYC* and *RAD51B*. TP73 is

particularly interesting as it comes up twice with very high (5342) and low reads (19), respectively.

There are 47 genes with more than 10 hits in the RNA-Seq samples. Generally, the remaining

DNA-Seq only samples contain less than 100 hits, and with the exception of *MYC*, which shows

a corresponding 3 hits in the RNA-Seq, show either no corresponding RNA-Seq data or just 1 hit.

It is noteworthy, that the other cases are negative even for such low amounts of hits. In contrast,

samples with 10 or more hits in the RNA-Seq often show a corresponding 1 hit of DNA-Seq not

only in that particular sample but in several other samples as well. We have not found any real

hotspots but earlier-reported *GRIK4* and *RNF166* come up multiple times. For the D lineages, the

affected genes were *CUL2*, *SLC31A1*, *MRPS28* and *LEPREL1*. While the signature for the A

lineages centres around several functions like immune system [*RNF166*, *CEACAM5*, *MPRIP*,

CTSE, *IL9R*, *FAM220A*, *PALLD*, *IKZF3*], proto-oncogenes [*PGAP3*], DNA-related processes

[*MACROD2*, *IGFBP6*, *FHAD1*, *RXRA*, *SIPA1L3*, *UTP11L*, *NFIA*], cellular processes [*UBA6*,

FLNB, *VPS13D*], signaling [*EPHA3*, *GRIK4*, *TAOK3*], and mitochondria [*DYRK1A*, *C6orf203*], the

signature of the D lineages is related to hypoxia/energetics (the Warburg effect). While this may

be related to the fewer samples compared to the A lineages, each affected gene was related to

this energy axis. This is particularly interesting because it closely relates to earlier findings in our

group [Richard *et al.*, 2010, Niccoli *et al.*, 2012, Cuninghame *et al.*, 2017, Dayer *et al.*, 2020].

HPV16 A1 sub-lineage: Our search with glutamate receptor, ionotropic, kainate 4 encoded

by *GRIK4* only retrieved a pharmacogenetic study [Yeon *et al.*, 2015] nevertheless associating

this neurotransmitter (altered in schizophrenia and depression) in a gene ontology pathway

analysis with Wnt/beta-catenin signaling implicated in cancer. Serine/threonine-protein kinase

encoded by *TAOK3* was found to be associated with Hippo signaling and a cancer stem cell

phenotype [Plouffe *et al.*, 2016, Bian *et al.,* 2019]. Exons can undergo non-linear reverse splicing

to form circ(ular)RNA. Such function has been elucidated in a breast cancer study for Post-GPI

attachment to proteins factor 3 encoded by *PGAP3* where overexpressed circPGAP3 inhibited

miR-330-3p and upregulated MYC leading to tumour progression and metastases [He *et al.,*

2020]. Other studies have also reported a deregulation of PGAP3 in pancreatic [Zhong *et al.*,

2020] and breast cancer [Pan *et al.*, 2019] without detailing biological consequences but

concluding to use PGAP3 respectively as tumour marker or to improve diagnostic accuracy.

Mitochondrial transcription rescue factor 1 encoded by *MTRES1* alias *C6orf203* binds highly

structured RNA *in vitro* and interacts with the mitochondrial ribosomal large subunit in human

cells. Because the loss of C6orf203 leads to reduced mitochondrial translation and consequently,

to respiratory inability it was reasoned that C6orf203 is required for efficient mitochondrial

translation [Gopalakrishna *et al.,* 2019]. Interestingly, C6orf203 showed the highest number of

hits in RNA-Seq, which together with DYRK1 could impact A1's adaption to energetics potentially

favouring oxidative phosphorylation (rather than aerobic glycolysis).

HPV16 D2/3 sub-lineages: With only 5 breakpoints in the DNA and 4 in the RNA, this sub-

lineage showed remarkably fewer hits than the A1 sub-lineage. Interestingly, both CUL2 and

MRPS28 proteins are associated with closely-related hallmarks of cancer: hypoxia and

deregulated energetics where the altered metabolism "prepares" malignant cells to the hypoxic

tumour environment. MRPS28, normally implicated in mitochondrial metabolism, was reported to

be modulated in thyroid tumours [Jacques *et al.*, 2013] connecting this finding to the Warburg

theory of cancer described in 1926 by Otto Heinrich Warburg: "Stoffwechsel der Tumoren" [Brand,

2010]. Warburg argued that rather than fully respiring in the presence of adequate oxygen cancer

cells instead resort to aerobic glycolysis fermenting sugar to lactate (rather than pyruvate) [Vander

Heiden, 2009]. A newly sparked interest in the *Warburg effect* in the past decade can be gleaned

from close to 3,000 publications on the PubMed server alone. Interestingly, MRPS28 is also

associated with adaption to hypoxia [Wang *et al.*, 2019]. Another finding in this respect was

HPV16 D2/D3 integration into the *CUL2* gene [Faull *et al.,* 2019, Fouad *et al.,* 2019], which

interestingly codes for a E3 ubiquitin ligase that targets pRb for degradation [Lee and Zhou, 2010].

While hypoxia-inducing factor 1-alpha (HIF1A) is also targeted by the CUL2 protein, this only

leads to its degradation during *normoxia* but not during *hypoxia* where this transcription factor

instead accumulates in the cell [Lee and Zhou, 2010 and references therein]. This underlines our

previous observations based on lab data that HIF1A is significantly increased in keratinocytes

transduced with D2/D3 E6 compared to A1 E6 in the hypoxic environment [Cuninghame *et al.,*

2017]. Notably, tumour hypoxia is one of the most challenging aspects of cancer treatment

[Cuninghame *et al.,* 2014 and references therein]. In the past 20 years, the importance of copper

in carcinogenesis (*e.g.*, solute carrier family 31 member 1, SLC31A1 alias CTR1) has been

gaining attention [Barresi *et al.,* 2016]. Copper homeostasis is tightly coupled to GSH redox

regulation in the mitochondria [Baker *et al.*, 2017]. Metal transporting solute carriers mediate

control over the availability of endogenous metal ions (*e.g.*, copper, iron and zinc) and may have

key roles in regulating tumour angiogenesis, cell proliferation, epithelial-to-mesenchymal

transition (EMT) and aberrant MAPK and STAT3 signal transduction in cancer [Jong *et al.*,

2014]. Indeed, copper was demonstrated to increase tumour growth in mice as an outcome of

elevated copper levels in drinking water and the switch to aerobic glycolysis (Warburg effect) may

be due in part to insufficient copper supply in the tumour microenvironement [Ishida *et al.*, 2013].

Similarly, disintegrin and metalloproteinase domain-containing protein 23 is a non-catalytic

protein encoded by *ADAM23* with potential roles in tumour progression related to EMT and

metastasis [Zmetakova *et al.*, 2019].

Overall, it is tempting to speculate that keratinocytes infected with the HPV16 D2/3 sub-

lineage adapt to a hypoxia- and Warburg effect-associated phenotype reminiscent of Warburg's

initial notion that cancer is a mitochondrial metabolic disease. Intriguingly, this underlines our

experimental findings using cell culture models [Richard *et al.*, 2010, Cuninghame *et al.*, 2017].

Moreover, patients with cervical cancer containing D2/D3 sub-lineage HPV16 are 7.7 years

younger than patients infected with the A1 sub-lineage [Berumen *et al.*, 2001] suggesting a more

aggressive tumour evolution for D2/D3. The A1 sub-lineage integration signature on the other

hand shows variable traits of carcinogenesis related to cancer pathway signaling, and most

importantly the potential to tap into oxidative phosphorylation rather than aerobic glycolysis for its

energy supply as evidenced by the high expression level of C6orf203. Notably, Warburg's notion

may not be complete since normal proliferating cells with intact mitochondria frequently use

aerobic glycolysis as energetics [Vander Heiden, 2009], and it is likely that mixed phenotypes,

e.g., with partially intact mitochondria exist in malignant tumours [Jia *et al.*, 2018].

Figure 6. Number of repeated integrated reads per human chromosomal location.

Manhattan plots for both A1 and D2/D3 variants based on HPVDetector analyzed RNA-Seq and

WXS data indicating the number of repeated integrated reads and the corresponding

chromosome number, chromosomal region, and name of the affected gene on the human

genome: **(A)** A1 RNA-Seq **(B)** A1 WXS **(C)** D2/D3 RNA-Seq **(D)** D2/D3 WXS. Horizontal lines

represent cut-offs: blue = 6 reads; red = 9 reads.

Present challenges and future directions

While we have addressed relevant literature, acquired access to TCGA sequencing data,

designed and setup a working algorithm, and in CESC samples carried out analyses for the

HPV16 genotype, sub-lineages, and integration, continued efforts are required to complete our

work. Even with TCGA datasets, a major limitation in answering these questions is the small

number of D2/D3 compared to A1 samples, and it may be necessary to extend these analyses

beyond the well-curated TCGA database, but more broadly into other databases (*e.g.*, the

Sequence Read Archive, SRA). While additional samples could possibly be found and

incorporated into our analyses, these data may lack sufficient metadata to be easily searchable

and screened (*e.g.*, experimental vs clinical data). Since the needs of a specific HPV-based

analysis may not have been considered *a priori* when data were collected for those samples,

there may be significant variability in data formats, read lengths, depths, and quality. Specifically,

whether sequence capture or enrichment for HPV sequences was performed would greatly affect

the probability of detecting HPV sequences as previously and independently reported by several

groups [Holmes *et al*., 2016; Jackson *et al*., 2016]. Sequencing errors could further confound

variant identification, requiring careful mitigation [Stewart *et al*., 2018]. Overall, while these

challenges exist, potentially relevant datasets deserve analysis and could be useful contributions

toward our research question.

Additional work is also required to determine the factors involved in a HPV16 sub-lineage-

specific genomic instability and integration risk, including the role of defective DNA damage repair

pathways [Seiwert *et al*., 2015; Ratnaparkhe *et al*., 2018]. It is worthwhile to explore relationships

between E6 splice variants and genomic stability [Olmedo-Nieva *et al*., 2018]. Researchers

continue to explore novel mechanisms surrounding HPV16 integration, such as "super-enhancer-

like elements" [Warburton *et al*., 2018] and interruption of tumour suppressor genes [Zhao *et al*.,

2016]. Future research could continue exploring these relationships, such as determining the role

of non-coding genome elements in genomic instability (*e.g.*, lncRNAs) [Munschauer *et al.*, 2018],

the similarities of viral-human integration with mobile elements and human fusion genes

[Imielinski and Ladanyi, 2018], as well as evolutionary perspectives [Petrie *et al.*, 2018]. Finally,

it remains to be seen how clinically-relevant HPV16 sub-lineage genotyping could be, and

whether this could lead to improved care and outcome for patients. One strategy could be to

compute individual risk scores based on the co-factors affecting the risk of disease progression

[Bastarache *et al.*, 2018]. New screening techniques could also be useful, such as urine sampling

[Van Keer *et al.*, 2018] followed by high-throughput sequencing and HPV genotyping, as well as

assessing methylation signatures [Cook *et al.*, 2018].

Full characterization of the genomic landscape surrounding detected HPV16 integration

sites within human chromosomes is required to comprehensively address our research question,

quantitatively and *qualitatively*. Having a more qualitative approach, by analyzing the affected

genes and potential functional consequences of the HPV integration process, makes us conclude

that an investigation beyond quantitative statistical analyses needs to be considered. Using both

approaches for biological interpretation are important for avoiding computational reproducibility

issues: the "alchemy problem" [Rahimi, 2018]. Further characterization of these data includes

analyzing viral and host nucleotide positions, sequence overlap/microhomologies, functional

annotations of viral and host features, such as structure and gene proximity (inside/outside,

exon/intron), promoters, enhancers, transcription factor binding sites, repeat elements, and

proximity to fragile sites. "Virtual ChIP-seq" [Karimzadeh and Hoffman, 2018] could be a useful

tool to predict nearby transcription factor binding sites (*e.g.*, CTCF [Doolittle-Hall *et al.*, 2015;

Paris *et al.*, 2015]). As described previously [Jackson *et al.*, 2016], the region surrounding a viral

integration site can be scanned for repeat elements, which may be prone to rearrangements,

using the UCSC human genome browser RepeatMasker track. Another important aspect for

qualitative assessment and interpretation will be to visualize and compare the A1 vs D2/D3 cases;

example visualizations exist in the literature [Akagi *et al.*, 2014; Tang *et al.*, 2013; Holmes *et al.*,

2016; Jackson *et al.*, 2016; Warburton *et al.*, 2018; Lagström *et al.*, 2019].

CONCLUDING REMARKS

From a basic science perspective, big data analysis may aid in unraveling the mechanisms

by which viruses can become integrated into their hosts, akin to mobile elements, and as a result

lead to persistence of those sequences with consequences for the host genome. In the case of

HPV16, a small number of changes in its DNA may be responsible for differential host genomic

instability and integration propensity, as evidenced by a difference in the rate of integration

between HPV16 D2 and D3 sub-lineages [Lou *et al.*, 2020]. Additional bioinformatics analysis of

this study and other *ex vivo* data, coupled with *in vitro* experimental work, is required to further

test the HPV integration hypothesis and the impact of other co-factors. Hypothesis-free

exploration of these data will also be important to clarify the gaps in our existing knowledge and

identify fascinating phenomena and molecular mechanisms to test in future experiments [Pipas,

2019]. From an applied science and clinical perspective, HPV16 sub-lineage genotyping can be

used for preventative screening and routine monitoring, and next-generation sequencing

specifically could be used on blood to detect integrated HPV sequences in circulating tumour DNA

[Holmes *et al.*, 2016] as a tumour-specific diagnostic biomarker and for monitoring residual

disease and recurrence after treatments.

METHODS

Data access and acquisition

Access to controlled sample data from TCGA, using the database of Genotypes and

Phenotypes (dbGaP), was acquired via application to the electronic Research Administration

(eRA) commons. We registered our host institution, Lakehead University, as a new organization

and received a Data Universal Numbering System (DUNS) identifier for access. Our proposal title

was "Characterization of human papillomavirus type 16 integration sites in cervical and head and

neck tumour biopsies", with approval allowing access to TCGA Cervical Squamous Cell

Carcinoma and Endocervical Adenocarcinoma (TCGA-CESC) as well as Head and Neck

Squamous Cell Carcinoma (TCGA-HNSC) files for analysis. After receiving access to the data (a

multi-step administrative process, outlined in **Figure 7A**) TCGA case files were transferred via

their universally unique identifier (UUID) using the command-line implementation of the Genomics

Data Commons Transfer Tool v1.4.0 and analyzed on the Compute Canada Cedar cluster, which

uses the Secure Shell (SSH) encryption protocol for secure data transmission.

High-throughput sequencing data from HPV16-positive cervical carcinoma-derived cell

lines SiHa [Friedl *et al.*, 1970] and CaSki [Pattillo *et al.*, 1977] were used as controls. For SiHa,

which contains low-copy number integrated HPV16, we used the same sample used in the

HPVDetector paper [Chandrani *et al.*, 2015], a whole-genome sequence from the DDBJ (DNA

Data Bank of Japan) Sequence Read Archive (DRA) run SRR1609142, from study SRP048769.

For CaSki, which contains a high-copy number of tandemly repeated HPV16 genome

integrations, we downloaded RNA-Seq data from the Sequence Read Archive (SRA) run

SRR9924750, from project PRJNA484129 [Sen *et al.*, 2020].

Figure 7. Access and analysis of TCGA data. (A) Access to controlled sample genomic and transcriptomic data from TCGA required a multi-step authorization process before being able to access relevant datasets. This administrative process, which requires annual renewal and project revision, takes approximately three months initially. **(B)** Our analytical workflow involved the following computational steps: TCGA high-throughput sequence read data (WXS or RNA-Seq) were first accessed and converted from pre-aligned BAM/SAM format to FASTQ read format. Data were then subjected to HPV16 sub-lineage determination and integration analysis.

HPV16 positivity and sub-lineage genotyping analysis

While HPV16-positive cervical cancer (TCGA-CESC) samples had associated sub-

lineages already identified (albeit, with 10 of the 16 current sub-lineage reference genomes

[TCGA Research Network, 2017]), sub-lineage genotyping was not immediately available for the

head and neck (TCGA-HNSC) cases [TCGA Research Network, 2015]. It was thus necessary to

develop a pipeline for ascertaining the sub-lineages of unknown identified samples (*i.e.*, those

that were identified as HPV16 positive by Cantalupo *et al.*, [2018] as well) in addition to confirming

their status as HPV16 positive through an analysis of our own. Our pipeline follows an analytical

workflow starting with the pre-aligned BAM files from TCGA [**Figure 7B**]. First, BAM files were

converted into FASTQ files using SAMtools v1.7 [Li *et al.*, 2009]. The reads were then mapped to

the reference genomes of each HPV16 sub-lineage [**Table 2**] using HISAT2 v2.1.0 [Kim *et al.*,

2015]. From the sixteen separate alignments performed, the one with the highest number of

mapped reads was used for quantifying the presence of HPV16 by calculating the number of viral

reads per million sequencing reads (RPM = 10^6 x number of mapped reads/total reads, which

accounts for sequence depth variation between sample libraries). RPM ranges for positivity

analysis were based on a prior study which sought reasonable cutoffs for HPV detection in cancer

tissues [Montgomery *et al.*, 2016]. If RPM < 0.99, the sample was marked as negative across all

categories and the analysis was terminated. Samples with RPM between 0.99 and 99 were

marked as having "low" positivity for HPV16, while samples of RPM > 99 were marked as having

"high" positivity. Given their large range, all RPM values were subjected to an inverse hyperbolic

sine (IHS) transformation [Burbidge *et al*, 1988] following IHS(x) = $\log_{10}(x + (x^2 + 1)^{1/2})$, an

alternative to traditional logarithmic transformation that allows values lower than 1 to have a

plottable log value and which has seen recent use when analyzing biological data [Haines *et al.*,

2020]. SAMtools v1.7 was then used to sort and prepare the files corresponding to these samples

for conversion to VCF format. BCFtools v1.9 [Li, 2011] was used for variant calling and generating

the consensus sequence. The new consensus sequences along with all reference sequences

were aligned against each other using MAFFT v7.310 [Katoh *et al.*, 2009] with default parameters.

Maximum likelihood trees were inferred using RAxML HPC v8.2.11 [Stamatakis, 2014], under the

general time-reversible nucleotide model with gamma-distributed rate heterogeneity and invariant

sites (GTRGAMMAI). Sub-lineages were determined from Newick tree files using ETE Toolkit

v3.0.0b34 [Huerta-Cepas *et al.*, 2010] based on topological distance. Each sample was

incorporated into the reference tree [**Figure 8**] and assigned a sub-lineage corresponding to the

nearest neighbour.

Table 2. HPV16 sub-lineage reference genomes used for genotyping. Derived from reference

data summarized in Burk *et al.* [2013], Chen *et al.* [2018], and Mirabello *et al.* [2018].

HPV16 sub-lineage	GenBank #	Other ID	Original nomenclature
A1	K02718	NC_001526.4 (latest)	European Prototype (EP)
A2	AF536179	w0122	European (E)
A3	HQ644236	AS411	European (E)
A4	AF534061	w0724	Asian (As)
B1	AF536180	w0236	African-1a (Af-1a)
B2	KU053915	IARC1100085NI	African-1 (Af-1)
B3	HQ644298	Z109	African-1b (Af-1b)
B4	KU053914	IARC907912AL	African-1 (Af-1)
C1	AF472509	R460	African-2 (Af-2)
C2	HQ644244	IARC240211GU	African-2 (Af-2)
C3	KU053920	IARC040105BE	African-2 (Af-2)
C4	KU053925	IARC304612MO	African-2 (Af-2)
D1	HQ644257	Qv00512	North-American-1 (NA1)
D2	AY686579	Qv15321	Asian-American (AA) 2
D3	AF402678	Qv00995	Asian-American (AA) 1
D4	KU053931	IARC903812AL	North-American-2 (NA2)

Figure 8. HPV16 sub-lineage reference tree for phylogenetic analyses. HPV16 phylogenetic

maximum likelihood tree used as a reference for sub-lineage determination. Branches correspond

to four lineages (A, B, C, and D), each containing four sub-lineage genomes (1 to 4). The tree

was built with FigTree v1.4.4 using standard settings.

Viral-human integration detection and characterization

Several open-source software tools for integration analysis are available for WGS, WXS, and whole-transcriptome sequencing (RNA-Seq): *e.g.*, VirusSeq [Chen *et al.*, 2013], ViralFusionSeq [Li *et al.*, 2013], VirusFinder [Wang *et al.*, 2013], HPVDetector v1.0 [Chandrani *et al.*, 2015], ViFi [Nguyen *et al.*, 2018] and VirTect [Xia *et al.*, 2019]. The latter tool name, VirTect, was also used by another group for viral genome sequencing and host genes related to carcinogenesis using RNA-Seq [Khan *et al.*, 2018]. HPVDetector v1.0 was chosen since it was specifically designed for detecting and annotating HPV integration events, as well its compatibility with the Compute Canada Cedar cluster environment. VirTect was also considered but is currently incompatible with available modules in Cedar, requiring an older version of Python (Python 2.7 vs 3.0). Integration analysis was performed using command-line execution of HPVDetector v1.0 [**Figure 7B**] set to "Integration" mode for all samples and "Transcriptome" or "Exome" mode depending if the sample contained WXS or RNA-Seq data, respectively. Raw output was obtained in the form of the number of integrated reads. This value was then converted to a percentage of the total number of viral reads to account for sequencing depth variation amongst selected samples and used as a metric to quantify viral-host integration. Samples with values at or above 1.14% were classified as positive for HPV16 integration, a threshold obtained through comparison to the landmark CESC study cases to maximize concordance. The remaining samples were considered as positive for the episomal form of HPV16.

DECLARATIONS

Acknowledgements

The results reported here are based upon data generated by The Cancer Genome Atlas Research Network (cancergenome.nih.gov) and we are grateful to the participants and Network members for this resource. This research was enabled in part by computational support provided

by Compute Ontario (www.computeontario.ca) and Compute Canada (www.computecanada.ca).

Thank you to Darryl Willick and Dr. Wely Floriano for their assistance in helping set up and

maintain required bioinformatics tools (such as the Galaxy platform) hosted at the Lakehead

University High Performance Computing Centre (LUHPCC) as well as Dr. Zigui Chen who shared

a HPV sub-lineage detection pipeline. We are also thankful for Bruce Bogacki's technical support

at the Thunder Bay Regional Health Research Institute (TBRHRI). Additional thanks to Kathleen

Roulston, Vanessa Masters, Anirudh Shahi, Dallas Nygard, and Christopher Gibb for assisting

with preliminary literature review, initial data access, and early analyses. Preliminary work related

to this study was presented as a poster in Heidelberg, Germany (11-14 Oct 2017) at The Mobile

Genome: Genetic and Physiological Impacts of Transposable Elements conference and talks in

Thunder Bay, Canada at the Health & Information Technology (HIT) Research Group Workshop

(12 Oct 2018) and the Health & Information Technology & Biotechnology & Allied Sciences

(HITBASS) Symposium (27 Oct 2019). This work was supported by the Natural Sciences and

Engineering Research Council of Canada (NSERC) grants to IZ (#RGPIN-2015-03855) and RJ

(CGS-D#454402-2014). The funding body had no role in study design, data collection, data

analysis and interpretation, or preparation of the manuscript. We declare no conflicts of interest.

600   *Authors' contributions*

601   The study was initially conceived of and designed by IZ and RJ. Data access was

602   managed by RJ, IZ, and AOV. Methods were developed, optimized, and analyses performed by

603   AOV with input from RJ and IZ. Biological interpretation, literature searches and manuscript

604   writing were led by RJ and IZ, with input from AOV, while all authors were involved in providing

605   critical revision and feedback.

REFERENCES

Akagi K, Li J, Broutian TR, Padilla-Nash H, Xiao W, Jiang B, *et al.* Genome-wide analysis of HPV integration in human cancers reveals recurrent, focal genomic instability. *Genome Res.* **2014**; 24:185-99. DOI: 10.1101/gr.164806.113

Baker ZN, Cobine PA, Leary SC. The mitochondrion: a central architect of copper homeostasis. *Metallomics* **2017**; 9:1501-1512. DOI: 10.1039/c7mt00221a

Barresi V, Trovato-Salinaro A, Spampinato G, Musso N, Castorina S, Rizzarelli E, *et al.* Transcriptome analysis of copper homeostasis genes reveals coordinated upregulation of SLC31A1,SCO1, and COX11 in colorectal cancer. *FEBS Open Bio.* **2016**; 6:794-806. DOI: 10.1002/2211-5463.12060

Bastarache L, Hughey JJ, Hebbring S, Marlo J, Zhao W, Ho WT, *et al.* Phenotype risk scores identify patients with unrecognized Mendelian disease patterns. *Science* **2018**; 359:1233-9. DOI: 10.1126/science.aal4043

Berumen J, Ordoñez RM, Lazcano E, Salmeron J, Galvan SC, Estrada RA, *et al.* Asian American variant of human papillomavirus 16 and risk for cervical cancer: a case–control study. *J Natl Cancer Inst.* **2001**; 93:1325-30. DOI: 10.1093/jnci/93.17.1325

Bian Y, Teper Y, Mathews Griner LA, Aiken TJ, Shukla V, Guha R, *et al.* Target Deconvolution of a Multikinase Inhibitor with Antimetastatic Properties Identifies TAOK3 as a Key Contributor to a Cancer Stem Cell-Like Phenotype. *Mol Cancer Ther.* **2019**; 18:2097-2110. DOI: 10.1158/1535-7163.MCT-18-1011

Brand RA. Biographical sketch: Otto Heinrich Warburg, PhD, MD. *Clin Orthop Relat Res.* **2010**; 468:2831-2. DOI: 10.1007/s11999-010-1533-z

Burbidge JB, Magee L, Robb AL. Alternative transformations to handle extreme values of the dependent variable. *J Am Stat Assoc.* **1988**; 83:123-7. DOI: 10.1080/01621459.1988.10478575

Burk RD, Harari A, Chen Z. Human papillomavirus genome variants. *Virology*. **2013**; 445:232-43.

DOI: 10.1016/j.virol.2013.07.018

Cantalupo PG, Katz JP, Pipas JM. HeLa nucleic acid contamination in The Cancer Genome Atlas

leads to the misidentification of HPV18. *J Virol*. **2015**; 89:4051-7. DOI: 10.1128/JVI.03365-

14

Cantalupo PG, Katz JP, Pipas JM. Viral sequences in human cancer. *Virology*. **2018**; 513:208-

16. DOI: 10.1016/j.virol.2017.10.017

Castellsagué X, Alemany L, Quer M, Halec G, Quirós B, Tous S, *et al*. HPV involvement in head

and neck cancers: comprehensive assessment of biomarkers in 3680 patients. *J Natl*

Cancer Inst. **2016**; 108:djv403. DOI: 10.1093/jnci/djv403

Chandrani P, Kulkarni V, Iyer P, Upadhyay P, Chaubal R, Das P, *et al*. NGS-based approach to

determine the presence of HPV and their sites of integration in human cancer genome. *Br*

J Cancer **2015**; 112:1958-65. DOI: 10.1038/bjc.2015.121

Chen Y, Yao H, Thompson EJ, Tannir NM, Weinstein JN, Su X. VirusSeq: software to identify

viruses and their integration sites using next-generation sequencing of human cancer

tissue. *Bioinformatics.* **2013**; 29:266-267. DOI: 10.1093/bioinformatics/bts665

Chen Z, DeSalle R, Schiffman M, Herrero R, Wood CE, Ruiz JC, *et al*. Niche adaptation and viral

transmission of human papillomaviruses from archaic hominins to modern humans. *PLoS*

Pathog. **2018**; 14:e1007352. DOI: 10.1371/journal.ppat.1007352

Clifford GM, Tenet V, Georges D, Alemany L, Pavón MA, Chen Z. Human Papillomavirus 16 sub-

lineage dispersal and cervical cancer risk worldwide: whole viral genome sequences from

7116 HPV16-positive women. *Papillomavirus Res.* **2019**; 7:67-74. DOI:

10.1016/j.pvr.2019.02.001

Cook DA, Krajden M, Brentnall AR, Gondara L, Chan T, Law JH, *et al.* Evaluation of a validated

methylation triage signature for human papillomavirus positive women in the HPV FOCAL

cervical cancer screening trial. *Int J Cancer* **2018**: 9 Nov 2018. DOI: 10.1002/ijc.31976

Cuninghame S, Jackson R, Lees SJ, Zehbe I. Two common variants of human papillomavirus

type 16 E6 differentially deregulate sugar metabolism and hypoxia signalling in permissive

human keratinocytes. *J Gen Virol.* **2017**; 98:2310-9. DOI: 10.1099/jgv.0.000905

Cuninghame S, Jackson R, Zehbe I. Hypoxia-inducible factor 1 and its role in viral carcinogenesis.

Virology **2014**; 456-457:370-83. DOI: 10.1016/j.virol.2014.02.027

Dawson ET, Wagner S, Roberson D, Yeager M, Boland J, Garrison E, *et al.* Viral coinfection

analysis using a MinHash toolkit. *BMC Bioinformatics* **2019**; 20:389. DOI: 10.1186/s12859-

019-2918-y

Dayer G, Masoom ML, Togtema M, Zehbe I. Virus–Host Protein–Protein Interactions between

Human Papillomavirus 16 E6 A1 and D2/D3 Sub-Lineages: Variances and Similarities. *Int.*

J. Mol. Sci. **2020**; 21:7980. DOI: 10.3390/ijms21217980

Doolittle-Hall J, Cunningham Glasspoole D, Seaman W, Webster-Cyriaque J. Meta-analysis of

DNA tumor-viral integration site selection indicates a role for repeats, gene expression and

epigenetics. *Cancers* **2015**; 7:2217-2235. DOI: 10.3390/cancers7040887

Faull SV, Lau AMC, Martens C, Ahdash Z, Hansen K, Yebenes H, *et al.* Structural basis of Cullin

2 RING E3 ligase regulation by the COP9 signalosome. *Nat Commun.* **2019**; 10:3814. DOI:

10.1038/s41467-019-11772-y

Fouad S, Wells OS, Hill MA, D'Angiolella V. Cullin Ring Ubiquitin Ligases (CRLs) in Cancer:

Responses to Ionizing Radiation (IR) Treatment. *Front Physiol.* **2019**; 10:1144. DOI:

10.3389/fphys.2019.01144

Friedl F, Kimura I, Osato T, Ito Y. Studies on a new human cell line (SiHa) derived from carcinoma

of uterus. I. Its establishment and morphology. *Proc Soc Exp Biol Med.* **1970**; 135:543-5.

DOI: 10.3181/00379727-135-35091a

Ghittoni R, Accardi R, Chiocca S, Tommasino M. Role of human papillomaviruses in

carcinogenesis. *Ecancermedicalscience.* **2015**; 9:526. DOI: 10.3332/ecancer.2015.526

Gopalakrishna S, Pearce SF, Dinan AM, Schober FA, Cipullo M, Spåhr H, *et al.* C6orf203 is an

RNA-binding protein involved in mitochondrial protein synthesis. *Nucleic Acids Res.* **2019**;

47:9386-99. DOI: 10.1093/nar/gkz684

Graham SV, Faizo AA. Control of human papillomavirus gene expression by alternative splicing.

Virus Res. **2017**; 231:83-95. DOI: 10.1016/j.virusres.2016.11.016

Haines SR, Siegel JA, Dannemiller KC. Modeling microbial growth in carpet dust exposed to

diurnal variations in relative humidity using the "Time-of-Wetness" framework. *Indoor Air.*

2020. DOI: 10.1111/ina.12686

Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell.* **2011**; 144:646-74. DOI:

10.1016/j.cell.2011.02.013

Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell.* **2000**; 100:57-70. DOI: 10.1016/S0092-

8674(00)81683-9

He D, Yang X, Kuang W, Huang G, Liu X, Zhang Y. The Novel Circular RNA Circ-PGAP3

Promotes the Proliferation and Invasion of Triple Negative Breast Cancer by Regulating the

miR-330-3p/Myc Axis. *Onco Targets Ther.* **2020**; 13:10149-59. DOI: 10.2147/OTT.S274574

Holmes A, Lameiras S, Jeannot E, Marie Y, Castera L, Sastre-Garau X, *et al.* Mechanistic

signatures of HPV insertions in cervical carcinomas. *Genome Med.* **2016**; 1:16004. DOI:

10.1038/npjgenmed.2016.4

Hoppe-Seyler K, Bossler F, Braun JA, Herrmann AL, Hoppe-Seyler F. The HPV E6/E7 oncogenes: key factors for viral carcinogenesis and therapeutic targets. *Trends Microbiol.* **2018**; 26:158-68. DOI: 10.1016/j.tim.2017.07.007

Huerta-Cepas J, Dopazo J, Gabaldón T. ETE: a python Environment for Tree Exploration. BMC Bioinformatics **2010**; 11:24. DOI: 10.1186/1471-2105-11-24

Imielinski M, Ladanyi M. Fusion oncogenes—genetic musical chairs. *Science* **2018**; 361:848-9. DOI: 10.1126/science.aau8231

Ishida S, Andreux P, Poitry-Yamate C, Auwerx J, Hanahan D. Bioavailable copper modulates oxidative phosphorylation and growth of tumors. *Proc Natl Acad Sci USA* **2013**; 110:19507-12. DOI: 10.1073/pnas.1318431110

Jackson R, Rosa BA, Lameiras S, Cuninghame S, Bernard J, Floriano WB, *et al.* Functional variants of human papillomavirus type 16 demonstrate host genome integration and transcriptional alterations corresponding to their unique cancer epidemiology. *BMC Genomics.* **2016**; 17:851. DOI: 10.1186/s12864-016-3203-3

Jackson R, Togtema M, Lambert PF, Zehbe I. Tumourigenesis driven by the human papillomavirus type 16 Asian-American E6 variant in a three-dimensional keratinocyte model. *PLoS One* **2014**; 9:e101540. DOI: 10.1371/journal.pone.0101540

Jacques C, Guillotin D, Fontaine JF, Franc B, Mirebeau-Prunier D, Fleury A, *et al.* DNA microarray and miRNA analyses reinforce the classification of follicular thyroid tumors. *J Clin Endocrinol Metab.* **2013**; 98:E981-9. DOI: 10.1210/jc.2012-4006

Jia D, Park JH, Jung KH, Levine H, Kaipparettu BA. Elucidating the Metabolic Plasticity of Cancer: Mitochondrial Reprogramming and Hybrid Metabolic States. *Cells* **2018**; 7:21. DOI: 10.3390/cells7030021

Jong NN, McKeage MJ. Emerging roles of metal solute carriers in cancer mechanisms and treatment. *Biopharm Drug Dispos.* **2014**; 35:450-62. DOI: 10.1002/bdd.1903

Karimzadeh M, Hoffman MM. Virtual ChIP-seq: Predicting transcription factor binding by learning

from the transcriptome. *bioRxiv* **2018**:168419. DOI: 10.1101/168419

Katoh K, Asimenos G, Toh H. Multiple alignment of DNA sequences with MAFFT. *Methods Mol*

Biol **2009**; 537:39-64. DOI: 10.1007/978-1-59745-251-9_3

Khan A, Liu Q, Chen X, Stucky A, Sedghizadeh PP, Adelpour D, *et al*. Detection of human

papillomavirus in cases of head and neck squamous cell carcinoma by RNA-seq and

VirTect. *Mol Oncol*. **2019**; 13:829-39. DOI: 10.1002/1878-0261.12435

Khoury JD, Tannir NM, Williams MD, Chen Y, Yao H, Zhang J, *et al*. Landscape of DNA virus

associations across human malignant cancers: analysis of 3,775 cases using RNA-Seq. *J*

Virol. **2013**; 87:8916-26. DOI: 10.1128/JVI.00340-13

Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements.

Nat Methods **2015**; 12:357-60. DOI: 10.1038/nmeth.3317

Lagström S, Umu SU, Lepistö M, Ellonen P, Meisal R, Christiansen IK, *et al*. TaME-seq: an

efficient sequencing approach for characterisation of HPV genomic variability and

chromosomal integration. *Sci Rep*. **2019**; 9:524. DOI: 10.1038/s41598-018-36669-6

Lee J, Zhou P. Cullins and cancer. *Genes Cancer* **2010**; 1:690-9. DOI:

10.1177/1947601910382899

Li H. A statistical framework for SNP calling, mutation discovery, association mapping and

population genetical parameter estimation from sequencing data. *Bioinformatics* **2011**;

27:2987-2993. DOI: 10.1093/bioinformatics/btr509

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, *et al*. The sequence alignment/map

format and SAMtools. *Bioinformatics* **2009**; 25:2078-9. DOI: 10.1093/bioinformatics/btp352

Li JW, Wan R, Yu CS, Co NN, Wong N, Chan TF. ViralFusionSeq: accurately discover viral

integration events and reconstruct fusion transcripts at single-base resolution.

Bioinformatics **2013**; 29:649-51. DOI: 10.1093/bioinformatics/btt011

Lou H, Boland JF, Torres-Gonzalez E, Albanez A, Zhou W, Steinberg M, *et al*. Human

papillomavirus 16 positive cervical cancer in Guatemala: The D2 and D3 sublineages differ

in integration rate and age of diagnosis. *bioRxiv* **2020**. DOI: 10.1101/2020.01.07.897546

Mesri EA, Feitelson MA, Münger K. Human viral oncogenesis: a cancer hallmarks analysis. *Cell*

Host Microbe. **2014**; 15:266-82. DOI: 10.1016/j.chom.2014.02.011

Mirabello L, Clarke MA, Nelson CW, Dean M, Wentzensen N, Yeager M, *et al*. The intersection

of HPV epidemiology, genomics and mechanistic studies of HPV-mediated carcinogenesis.

Viruses **2018**; 10:80. DOI: 10.3390/v10020080

Montgomery ND, Parker JS, Eberhard DA, Patel NM, Weck KE, Sharpless NE, *et al*. Identification

of Human Papillomavirus Infection in Cancer Tissue by Targeted Next Generation

Sequencing. *Appl Immunohistochem Mol Morphol*. **2016**; 24:490-5. DOI:

10.1097/PAI.0000000000000215

Munschauer M, Nguyen CT, Sirokman K, Hartigan CR, Hogstrom L, Engreitz JM, *et al*. The

NORAD lncRNA assembles a topoisomerase complex critical for genome stability. *Nature*

2018; 561:132-136. DOI: 10.1038/s41586-018-0453-z

Ndiaye C, Mena M, Alemany L, Arbyn M, Castellsagué X, Laporte L, *et al*. HPV DNA, E6/E7

mRNA, and p16^{INK4a} detection in head and neck cancers: a systematic review and meta-

analysis. *Lancet Oncol*. **2014**; 15:1319-1331. DOI: 10.1016/S1470-2045(14)70471-1

Nguyen N-PD, Deshpande V, Luebeck J, Mischel PS, Bafna V. ViFi: accurate detection of viral

integration and mRNA fusion reveals indiscriminate and unregulated transcription in

proximal genomic regions in cervical cancer. *Nucleic Acids Res*. **2018**; 46:3309-25. DOI:

10.1093/nar/gky180

Niccoli S, Abraham S, Richard C, Zehbe I. The Asian-American E6 variant protein of human

papillomavirus 16 alone is sufficient to promote immortalization, transformation, and

migration of primary human foreskin keratinocytes. *J Virol.* **2012**; 86:12384-96. DOI: 10.1128/JVI.01512-12

Nulton TJ, Olex AL, Dozmorov M, Morgan IM, Windle B. Analysis of The Cancer Genome Atlas sequencing data reveals novel properties of the human papillomavirus 16 genome in head and neck squamous cell carcinoma. *Oncotarget* **2017**; 8:17684-99. DOI: 10.18632/oncotarget.15179

Olmedo-Nieva L, Muñoz-Bello J, Contreras-Paredes A, Lizano M. The role of E6 spliced isoforms (E6*) in human papillomavirus-induced carcinogenesis. *Viruses* **2018**; 10:45. DOI: 10.3390/v10010045

Pan X, Hu X, Zhang YH, Chen L, Zhu L, Wan S, *et al.* Identification of the copy number variant biomarkers for breast cancer subtypes. *Mol Genet Genomics* **2019**; 294:95-110. DOI: 10.1007/s00438-018-1488-4

Parfenov M, Pedamallu CS, Gehlenborg N, Freeman SS, Danilova L, Bristow CA, *et al.* Characterization of HPV and host genome interactions in primary head and neck cancers. *Proc Natl Acad Sci USA.* **2014**; 111:15544-15549. DOI: 10.1073/pnas.1416074111

Paris C, Pentland I, Groves I, Roberts DC, Powis SJ, Coleman N, *et al.* CCCTC-binding factor recruitment to the early region of the human papillomavirus 18 genome regulates viral oncogene expression. *J Virol.* **2015**; 89:4770-4785. DOI: 10.1128/JVI.00097-15

Pattillo RA, Hussa RO, Story MT, Ruckert AC, Shalaby MR, Mattingly RF. Tumor antigen and human chorionic gonadotropin in CaSki cells: a new epidermoid cervical cancer cell line. *Science* **1977**; 196:1456-8. DOI: 10.1126/science.867042

Petrie KL, Palmer ND, Johnson DT, Medina SJ, Yan SJ, Li V, *et al.* Destabilizing mutations encode nongenetic variation that drives evolutionary innovation. *Science* **2018**; 359:1542-1545. DOI: 10.1126/science.aar1954

Pipas JM. DNA tumour viruses and their contributions to molecular biology. *J Virol.* **2019**;
JVI.01524-18. DOI: 10.1128/JVI.01524-18

Plouffe SW, Meng Z, Lin KC, Lin B, Hong AW, Chun JV, *et al.* Characterization of Hippo Pathway
Components by Gene Inactivation. *Mol Cell.* **2016**; 64:993-1008. DOI:
10.1016/j.molcel.2016.10.034

Rahimi A. Quotes within article by Hutson M. AI researchers allege that machine learning is
alchemy. *Science* **2018**. DOI: 10.1126/science.aau0577

Ratnaparkhe M, Wong J, Wei PC, Hlevnjak M, Kolb T, Haag D, *et al.* Defective DNA damage
repair leads to frequent catastrophic genomic events in murine and human tumors. *Nat*
Commun. **2018**; 9:4760. DOI: 10.1038/s41467-018-06925-4

Richard C, Lanner C, Naryzhny S, Sherman L, Lee H, Lambert PF, *et al.* The immortalizing and
transforming ability of two common human papillomavirus 16 E6 variants with different
prevalences in cervical cancer. *Oncogene* **2010**; 29:3435-3445. DOI: 10.1038/onc.2010.93

Seedorf K, Krämmer G, Dürst M, Suhai S, Röwekamp WG. Human papillomavirus type 16 DNA
sequence. *Virology.* **1985**; 145:181-185. DOI: 10.1016/0042-6822(85)90214-4

Seiwert TY, Zuo Z, Keck MK, Khattri A, Pedamallu CS, Stricker T, *et al.* Integrative and
comparative genomic analysis of HPV-positive and HPV-negative head and neck
squamous cell carcinomas. *Clin Cancer Res.* **2015**; 21:632-641. DOI: 10.1158/1078-
0432.CCR-13-3310

Sen P, Ghosal S, Hazra R, Arega S, Mohanty R, Kulkarni KK, *et al.* Transcriptomic analyses of
gene expression by CRISPR knockout of miR-214 in cervical cancer cells. *Genomics* **2020**;
112:1490-9. DOI: 10.1016/j.ygeno.2019.08.020

Sidiropoulos N, Sohi SH, Pedersen TL, Porse BT, Winther O, Rapin N, *et al.* SinaPlot: an
enhanced chart for simple and truthful representation of single observations over multiple
classes. *J Comput Graph Stat.* **2018**; 27:673-6. DOI: 10.1080/10618600.2017.1366914

Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **2014**; 30:1312-1313. DOI: 10.1093/bioinformatics/btu033

Stewart C, Leshchiner I, Hess J, Getz G. Comment on "DNA damage is a pervasive cause of

sequencing errors, directly confounding variant identification". *Science* **2018**;

361:eaas9824. DOI: 10.1126/science.aas9824

Tang KW, Alaei-Mahabadi B, Samuelsson T, Lindh M, Larsson E. The landscape of viral

expression and host gene fusion and adaptation in human cancer. *Nat Commun*. **2013**;

4:2513. DOI: 10.1038/ncomms3513

TCGA Research Network. Comprehensive genomic characterization of head and neck squamous

cell carcinomas. *Nature* **2015**; 517:576-582. DOI: 10.1038/nature14129

TCGA Research Network. Integrated genomic and molecular characterization of cervical cancer.

Nature **2017**; 543:378-384. DOI: 10.1038/nature21386

Togtema M, Hussack G, Dayer G, Teghtmeyer MR, Raphael S, Tanha J, *et al*. Single-domain

antibodies represent novel alternatives to monoclonal antibodies as targeting agents

against the human papillomavirus 16 E6 protein. *Int J Mol Sci*. **2019**; 20:2088. DOI:

10.3390/ijms20092088

Togtema M, Jackson R, Grochowski J, Villa PL, Mellerup M, Chattopadhyaya J, *et al*. Synthetic

siRNA targeting human papillomavirus 16 E6: a perspective on *in vitro* nanotherapeutic

approaches. *Nanomedicine* **2018**; 13:455-474. DOI: 10.2217/nnm-2017-0242

Togtema M, Jackson R, Richard C, Niccoli S, Zehbe I. The human papillomavirus 16 European-

T350G E6 variant can immortalize but not transform keratinocytes in the absence of E7.

Virology **2015**; 485:274-282. DOI: 10.1016/j.virol.2015.07.025

Vander Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg effect: the metabolic

requirements of cell proliferation. *Science* **2009**; 324:1029-33. DOI:

10.1126/science.1160809

Van Doorslaer K, Dillner J. The launch of an international animal papillomavirus reference center.

Viruses **2019**; 11:55. DOI: 10.3390/v11010055

Van Doorslaer K, Li Z, Xirasagar S, Maes P, Kaminsky D, Liou D, *et al*. The Papillomavirus

Episteme: a major update to the papillomavirus sequence database. *Nucleic Acids Res.*

2017a; 45:D499-506. DOI: 10.1093/nar/gkw879

Van Doorslaer K, Ruoppolo V, Schmidt A, Lescroël A, Jongsomjit D, Elrod M, *et al*. Unique

genome organization of non-mammalian papillomaviruses provides insights into the

evolution of viral early proteins. *Virus Evol.* **2017b**; 3:vex027. DOI: 10.1093/ve/vex027

Van Doorslaer K, Tan Q, Xirasagar S, Bandaru S, Gopalan V, Mohamoud Y, *et al*. The

Papillomavirus Episteme: a central resource for papillomavirus sequence data and analysis.

Nucleic Acids Res. **2013**; 41:D571-578. DOI: 10.1093/nar/gks984

Van Keer S, Tjalma WA, Pattyn J, Biesmans S, Pieters Z, Van Ostade X, *et al*. Human

papillomavirus genotype and viral load agreement between paired first-void urine and

clinician-collected cervical samples. *Eur J Clin Microbiol Infect Dis.* **2018**; 37:859-869. DOI:

10.1007/s10096-017-3179-1

Wang H, Chai Z, Hu D, Ji Q, Xin J, Zhang C, *et al*. A global analysis of CNVs in diverse yak

populations using whole-genome resequencing. *BMC Genomics* **2019**; 20:61. DOI:

10.1186/s12864-019-5451-5

Wang Q, Jia P, Zhao Z. VirusFinder: software for efficient and accurate detection of viruses and

their integration sites in host genomes through next generation sequencing data. *PLoS One.*

2013; 8:e64465. DOI: 10.1186/s12920-018-0461-8

Warburton A, Redmond CJ, Dooley KE, Fu H, Gillison ML, Akagi K, *et al*. HPV integration hijacks

and multimerizes a cellular enhancer to generate a viral-cellular super-enhancer that drives

high viral oncogene expression. *PLoS Genet.* **2018**; 14:e1007179. DOI:

10.1371/journal.pgen.1007179

Winder DM, Pett MR, Foster N, Shivji MK, Herdman MT, Stanley MA, *et al*. An increase in DNA

double-strand breaks, induced by Ku70 depletion, is associated with human papillomavirus

16 episome loss and de novo viral integration events. *J Pathol*. **2007**; 213:27-34. DOI:

10.1002/path.2206

Xi LF, Koutsky LA, Galloway DA, Kiviat NB, Kuypers J, Hughes JP, *et al*. Genomic variation of

human papillomavirus type 16 and risk for high grade cervical intraepithelial neoplasia. *J*

Natl Cancer Inst. **1997**; 89:796-802. DOI: 10.1093/jnci/89.11.796

Xia Y, Liu Y, Deng M, Xi R. Detecting virus integration sites based on multiple related sequencing

data by VirTect. *BMC Med Genomics* **2019**; 12:19. DOI: 10.1186/s12920-018-0461-8

Xu B, Chotewutmontri S, Wolf S, Klos U, Schmitz M, Dürst M, *et al*. Multiplex identification of

human papillomavirus 16 DNA integration sites in cervical carcinomas. *PLoS One* **2013**;

8:e66693. DOI: 10.1371/journal.pone.0066693

Yeon B, Ahn E, Kim KI, Kim IW, Oh JM, Park T. Analysis of pharmacogenomic variants associated

with population differentiation. PLoS One. 2015 Mar 25;10(3):e0119994. doi:

10.1371/journal.pone.0119994. PMID: 25807276; PMCID: PMC4373713.

Zapatka M, Borozan I, Brewer DS, Iskar M, Grundhoff A, Alawi M, *et al.* The landscape of viral

associations in human cancers. *Nat Genet.* **2020**; 320-330. DOI: 10.1038/s41588-019-

0558-9

Zhao JW, Fang F, Guo Y, Zhu TL, Yu YY, Kong FF, *et al*. HPV16 integration probably contributes

to cervical oncogenesis through interrupting tumor suppressor genes and inducing

chromosome instability. *J Exp Clin Cancer Res*. **2016**; 35:180. DOI: 10.1186/s13046-016-

0454-4

Zhong J, Jermusyk A, Wu L, Hoskins JW, Collins I, Mocci E, *et al*. A Transcriptome-Wide

Association Study Identifies Novel Candidate Susceptibility Genes for Pancreatic Cancer. *J*

Natl Cancer Inst. **2020**; 112:1003-12. DOI: 10.1093/jnci/djz246

Zmetakova I, Kalinkova L, Smolkova B, Horvathova Kajabova V, Cierna Z, Danihel L, *et al*. A

disintegrin and metalloprotease 23 hypermethylation predicts decreased disease-free

survival in low-risk breast cancer patients. *Cancer Sci*. **2019**; 110:1695-704. DOI:

10.1111/cas.13985