**Title:**

**More than noise: Context-dependant luminance contrast discrimination in a**

**coral reef fish (*Rhinecanthus aculeatus*)**

**Running title:**

**Luminance contrast discrimination in reef fish**

Authors: Cedric P. van den Berg[1,3]*, Michelle Hollenkamp[2], Laurie J. Mitchell[1,3], Erin J. Watson[1],

Naomi F. Green[1], N. Justin Marshall[3], Karen L. Cheney[1,3]

*[1] School of Biological Sciences, The University of Queensland, Brisbane, QLD 4072, Australia*

*[2] Department of Ecology and Evolutionary Biology, University of Colorado Boulder, Boulder, CO 80309, USA*

*[3] Queensland Brain Institute, The University of Queensland, Brisbane, QLD 4072, Australia*

*\*corresponding author: c.vandenberg@uq.edu.au*

## Abstract

Achromatic (luminance) vision is used by animals to perceive motion, pattern, space and texture. Luminance contrast sensitivity thresholds are often poorly characterised for individual species and are applied across a diverse range of perceptual contexts using over-simplified assumptions of an animal's visual system. Such thresholds are often estimated using the Receptor Noise Limited model (RNL) using quantum catch values and estimated noise levels of photoreceptors. However, the suitability of the RNL model to describe luminance contrast perception remains poorly tested.

Here, we investigated context-dependent luminance discrimination using triggerfish (*Rhinecanthus aculeatus)* presented with large achromatic stimuli (spots) against uniform achromatic backgrounds of varying absolute and relative contrasts. 'Dark' and 'bright' spots were presented against relatively dark and bright backgrounds. We found significant differences in luminance discrimination thresholds across treatments. When measured using Michelson contrast, thresholds for bright spots on a bright background were significantly higher than for other scenarios, and the lowest threshold was found when dark spots were presented on dark backgrounds. Thresholds expressed in Weber contrast revealed increased contrast sensitivity for stimuli darker than their backgrounds, which is consistent with the literature. The RNL model was unable to estimate threshold scaling across scenarios as predicted by the Weber-Fechner law, highlighting limitations in the current use of the RNL model to quantify luminance contrast perception. Our study confirms that luminance contrast discrimination thresholds are context-dependent and should therefore be interpreted with caution.

38    **Keywords**

39    Receptor Noise Limited Model, experimental psychophysics, perceptual thresholds, visual

40    ecology, visual modelling

## Introduction

The perception of chromatic (colour) and achromatic (luminance) information from the

surrounding environment enables animals to perform complex behaviours such as navigation,

mate choice, territorial defence, foraging and predator avoidance. Chromatic information is

largely used to assess the spectral composition and quality of objects or other organisms (Osorio

and Vorobyev, 2005), whereas achromatic information is predominantly used for object grouping,

pattern and texture detection, figure-ground segregation, and the perception of motion and

depth (Anderson, 2011; Brooks, 2014; Elder and Sachs, 2004; Elder and Velisavljevic, 2010;

Gilchrist, 2008; Gilchrist and Radonjic, 2009).

Behavioural experiments to examine colour and luminance discrimination thresholds

enable inferences on the perception of visual information by non-human observers (for discussion

see Olsson et al., 2018). Thresholds may be influenced by spatiotemporal and spatiochromatic

properties of a visual scene, as the perception of colour, pattern, luminance and motion interact

when low-level retinal information is processed along pathways in the visual cortex (Monnier and

Shevell, 2003; Shapley and Hawken, 2011; Shevell and Kingdom, 2008), or at even earlier stages

(Heath et al., 2020; Zhou et al., 2020). For example, the perception of luminance contrast in

animals is influenced by a range of factors, including perceived illumination and reflectance

(which in turn depend on illumination) in addition to various spatial and temporal properties,

such as depth perception, adaptation, stimulus geometry and viewer expectation of the position

and shape of a stimulus (Corney and Lotto, 2007; Craik, 1938; Gilchrist and Radonjic, 2009;

Heinemann and Chase, 1995; Kingdom, 2011; Lind et al., 2012; Pelli and Bex, 2013). The impact

of post-photoreceptor, and particularly post-retinal neuronal processing, on luminance

perception is often illustrated by visual displays targeting these effects, such as simultaneous

contrast illusions (Fig. 1). To investigate the design, function and evolution of animal visual signals,

the context sensitivity of visual threshold measurement is important to define.

**Figure 1:** The simultaneous contrast effect: despite having identical luminance, the left most internal square appears darker than the right one as a result of the background contrast against which they each square is viewed.

Luminance contrast of objects against their visual background or between objects can be

measured in a number of different ways, including Michelson contrast (MC), Weber contrast (WC)

and Root Mean Square (RMS) (Bex and Makous, 2002; Moulden et al., 1990; Vorobyev and Osorio,

1998). MC is commonly used to describe the contrast between two comparably sized objects or

sine gratings (Bex and Makous, 2002; Pelli and Bex, 2013). The WC, particularly popular in

psychophysics, is designed to describe the contrast of an object against a dominating background,

while accounting for the Weber-Fechner law that states that psychometric thresholds scale with

stimulus intensity at a constant ratio: the Weber fraction (Dzhafarov and Colonius, 1999; Norwich,

1987; Treisman, 1964). Luminance discrimination thresholds in animals have been obtained from

behavioural experiments and measured in MC, and most commonly in WC (e.g. Lind et al., 2013;

Scholtyssek et al., 2008). For example, human luminance discrimination thresholds are between

0.11 and 0.14 WC (Cornsweet and Pinsker, 1965), which is similar to seals (0.11-0.14 WC)

(Scholtyssek and Dehnhardt, 2013; Scholtyssek et al., 2008). Other animals have poorer

luminance discrimination thresholds, including birds (0.18-0.22 WC) (Lind et al., 2013), dogs (0.22-

0.27 WC) (Pretterer et al., 2004), manatees 0.35 WC (Griebel and Schmid, 1997) and horses (0.42-

0.45 WC) (Geisbauer et al., 2004).

Behavioural experiments measuring discrimination thresholds are often time-consuming

and unfeasible, especially when studying non-model organisms. Furthermore, focal species may

not be suitable for behavioural testing due to ethical, legal or logistical restrictions. Therefore, in

studies on visual ecology, the 'Receptor Noise Limited' (RNL) model (Vorobyev and Osorio, 1998)

has been adopted as a means of estimating whether both colour and luminance contrast within

and between animal colour patterns, or between animals and their backgrounds, are perceivable

to a species The model was initially designed for colour contrast modelling; however, the

achromatic interpretation of the RNL model (Siddiqi, 2004).has been used in a large number of

studies to quantify the perception of luminance contrast by non-human observers (e.g. Cheney

and Marshall, 2009; Marshall et al., 2016; Spottiswoode and Stevens, 2010; Stoddard and Stevens,

2010; Troscianko and Stevens, 2015). In contrast to using WC or MC, the RNL model allows the

prediction of contrast discriminability without the need of behavioural experimentation, but

instead can even be applied using conservatively chosen estimates of vision parameters (Olsson

et al., 2018).

The RNL model assumes that signal discrimination under 'ideal viewing conditions' is

limited by noise originating in the receptors and subsequent opponent processing (Vorobyev and

Osorio, 1998; Vorobyev et al., 2001). It was designed to estimate when a signal receiver could

discriminate between two colours that were spectrally similar, adjacent, of fixed size and

luminance. The point at which the contrast between two stimuli surpasses a behaviourally

determined threshold (e.g. 75% correct choice in a pairwise choice paradigm) is then expressed

as a 'Just Noticeable Difference' (JND) corresponding to a Euclidian distance ($\Delta S$) in an $n$-

dimensional space, where $n$ is the number of colour or luminance processing channels (Hempel

de Ibarra et al., 2001). The model predicts a JND is equal to 1 $\Delta S$ if all model assumptions (ideal

viewing conditions) are met (Vorobyev and Osorio, 1998; Vorobyev et al., 2001).

However, in many animals, the neuronal pathways leading to the perception of luminance

contrast vary significantly from those involved in the perception of colour contrast. In humans for

example, the magnocellular and parvocellular pathways segregate colour and luminance tasks

(Zeki, 1993) which can interact (to varying degrees) during subsequent neuronal processing (e.g.

Bruce et al., 2010; Gegenfurtner and Kiper, 1992; Shapley and Hawken, 2011; Simmons and

Kingdom, 2002; Webster and Wilson, 2000). The pronounced context-dependent sensitivity of

luminance contrast perception is partly due to the fact that achromatic vision in vertebrates lacks

a process as efficient as colour constancy (Kelber et al., 2003; Land, 1986; Osorio and Vorobyev,

2008; Wallach, 1948) which enables the perceived color of objects to remain relatively constant

under varying illumination conditions (but see Lotto and Purves, 2000; Simpson et al., 2016).

However, despite assuming receptor noise levels to be the limiting factor shaping both chromatic

and achromatic contrast perception, behavioural validations of perceptual distances calculated

using the RNL model are required in various visual contexts (as suggested by Olsson, Lind, &

Kelber, 2018 but see Skorupski & Chittka, 2011; Vasas, Brebner, & Chittka, 2018). Olsson et al.

(2018) have further suggested a conservative threshold of up to 1 JND = 3 $\Delta S$ for colour

discrimination, as both parameter choice and behavioural threshold validation are often difficult.

The use of such conservative chromatic discrimination thresholds in perceptually complex

contexts has recently been supported by empirical work (Escobar-Camacho et al., 2019; Sibeaux

et al., 2019). However, no empirical evidence exists for choosing conservative luminance

(achromatic) contrast thresholds using the RNL model.

In this study, we performed behavioural experiments with triggerfish, *Rhinecanthus*

*aculeatus*, to determine luminance discrimination thresholds in a foraging task using large stimuli

under well-illuminated (photopic) conditions. We refer to the task of discriminating a stimulus

from its background as a detection task, as this reflects a common use of the achromatic RNL

model in visual ecology, most prominently when quantifying the efficiency of animal camouflage

(e.g. Troscianko *et al.*, 2016). The ability to detect the presence of a potential prey item is the pre-

requisite for more complex cognitive processes and decision making by a predator (Endler, 1991)

and as such more likely to reflect low-level retinal and post-retinal properties of visual contrast

processing, such as the ones the RNL model has been developed to reflect.

Fish were trained to first locate a target spot that was randomly placed on an achromatic

background from which the spot differed in terms of luminance, and then peck it to receive a food

reward. Luminance discrimination thresholds were measured for both increasing and decreasing

luminance, on both a relatively bright and a dark background. We report thresholds in terms of

Michelson and Weber contrast, but then translate these thresholds into achromatic ΔS using the

log transformed RNL model, as per Siddiqi *et al.* (2004). To our knowledge, this is the first time

that achromatic discrimination thresholds have been quantified in a marine vertebrate, using a

'detection' task (as opposed to a pairwise choice paradigm as in Siebeck et al., (2014)) as well as

doing so using animals which have been trained to detect both randomly placed brighter and

darker stimuli simultaneously.

**Materials and methods**

**Study species**

We used triggerfish *Rhinecanthus aculeatus* (n = 15), which ranged in size from 6 to 16 cm

(standard length, SL). This species inhabits shallow tropical reefs and temperate habitats

throughout the Indo-Pacific and feeds on algae, detritus and invertebrates (Randall et al., 1997).

They are relatively easy to train for behavioural studies (e.g. Green *et al.*, 2018), and their visual

system has been well-studied (Champ et al., 2014; Champ et al., 2016; Cheney et al., 2013;

Pignatelli et al., 2010). They have trichromatic vision based on one single cone, containing short-

wavelength sensitive visual pigment (sw photoreceptor $\lambda_{max}$ = 413 nm); and a double cone, which

houses the medium-wavelength sensitive pigment (mw photoreceptor $\lambda_{max}$ = 480 nm) and long-

wavelength sensitive pigment (lw photoreceptor $\lambda_{max}$ = 528 nm) (Cheney et al., 2013). The double

cone members are used independently in colour vision (Pignatelli et al., 2010), but are also

thought to be used in luminance vision (Marshall et al., 2003; Siebeck et al., 2014), as per other

animals such as birds and lizards (Lythgoe, 1979).

However, it is not clear if both members of the double cone are used for luminance

perception via electrophysiological coupling (Marchiafava, 1985; Siebeck et al., 2014). We have

based our experiment on the assumption of both members contributing as per previous studies

modelling luminance perception in *R. aculeatus* (Mitchell et al., 2017; Newport et al., 2017). These

studies have used the added input of both double cone members (mw + lw), whereas our study

uses the averaged output of both members (mw + lw / 2) as suggested by Pignatelli & Marshall,

(2010) and Pignatelli et al., (2010). Additionally, Cheney et al., (2013) have used the lw receptor

response rather than both double cone members for luminance contrast modelling in *R.*

*aculeatus*, based on discussions in Marshall et al., (2003). However, MC/WC/$\Delta$S contrast values

are identical for $f_{t/b}$ = mw + lw and $f_{t/b}$ = mw + lw / 2 (eq. 2). Using the lw member of the double

cone only (as opposed to both members) causes less than 1% difference (well below

measurement error) in receptor stimulation due to the lack of chromaticity of the stimuli and the

strong overlap of spectral sensitivities of both double cone members (Cheney et al., 2013).

Fish were obtained from an aquarium supplier (Cairns Marine Pty Ltd, Cairns), shipped to

The University of Queensland, Brisbane and housed in individual tanks of 120L (W: 40cm; L: 80

cm, H: 40cm). They were acclimatised for at least one week before training commenced.

Experiments were conducted in September-November 2017. All experimental procedures for this

study were approved by the University of Queensland Animal Ethics Committee

(SBS/111/14/ARC).

**Stimulus creation and calibration**

We used a custom programme in Matlab (MathWorks, 2000) to create the stimuli

(available on [GitHub](GitHub)). This programme allowed us to specify the RGB values of the background

and target spot, and randomly allocate the target spot (1.6cm diam) to a position on the

background. The size of spot was chosen to be well within the spatial acuity of *R. aculeatus*

(Champ et al., 2014) and could be easily resolved by the fish from anywhere in their aquaria.

Stimuli, distractors and backgrounds were printed on TrendWhite ISO 80 A4 recycled paper using

a HP Laserjet Pro 400 color M451dn printer. Stimuli were then laminated using matte laminating

pouches. Throughout the experiment, any stimuli with detectable scratches or damage were

replaced immediately.

To ensure all stimuli were achromatic, reflectance measurements were plotted in colour

space as per Cheney et al. (2019). Target and background colours were < 1 $\Delta S$ from the achromatic

locus in the RNL colour space as per equations 1-4 in Hempel de Ibarra et al. (2001).

Photoreceptor stimulation was calculated using spectral sensitivities of triggerfish from Cheney

et al. (2013). Measures of photoreceptor noise are not available in this species, therefore we

assumed a cone ratio of 1:2:2 (SW:MW:LW) with a standard deviation of noise in a single cone of

0.05 as per (Champ et al., 2016; Cheney et al., 2019). The cone abundance was normalised relative

to the LW cone, which resulted in channel noise levels (univariant Weber fractions) of

0.07:0.05:0.05 (SW:MW:LW).

We quantified luminance contrast using calibrated digital photography (Stevens et al.,

2007) using an Olympus E-PL5 Penlight camera fitted with a 60mm macro lens to take pictures of

each stimulus combination (Suppl. material). Two EcoLight KR96 30W white LED lights (Eco-lamps

Inc. – Hong Kong) were used to provide even illumination between 400-700nm wavelength

(Suppl. Material). Pictures were analysed using the 'Multispectral Image Calibration and Analysis'

(MICA) Toolbox (Troscianko and Stevens, 2015) to calculate cone capture quanta of the double

cone. The double cone stimulation was calculated as the average stimulation of the medium-

wavelength (MW) and long-wavelength (LW) cone, as per Pignatelli *et al.* (2010). We used a

spatial acuity estimation of 2.75 cycles per degree (Champ et al., 2014) at 15cm viewing distance

using AcuityView (Caves and Johnsen, 2018) implemented in MICA's QCPA package (van den Berg

et al., 2020).

Stimulus contrast was measured as Michelson contrast using the MICA derived cone catch

values of the double cones. The stimuli contrasts were evenly spaced around an area of interest

in which the threshold was expected to lie, according to pilot trials. Weber contrast of the

thresholds was calculated as $\Delta I_t/I_s$; where $\Delta I_t$ is the stimulus contrast at threshold and $I_s$ is the

intensity of the distractor or background respectively as per Lind *et al.*, (2013). Achromatic ΔS

values were calculated according to equation 7 in Siddiqi *et. al* (2004) (Eq. 1).

$$\Delta S = |\Delta f_{dbl}/\omega|$$    **Equation 1**

Where $\Delta f_{dbl}$ describes the contrast in von Kries corrected double cone stimulation

between the stimulus $(f_t)$ and its background $(f_b)$, calculated as per equation 4 in Siddiqi *et. al*

(2004) (Eq. 2) in relation to the weber fraction $(\omega)$ of the double cone channel. When using the

natural logarithm of the quantum catches $\omega = e_i$

$$\Delta f_{dbl} = ln(f_t) - ln(f_b)$$     **Equation 2**

A total of 6 stimuli were created for each scenario (Fig. 2, Table 1).

**Experimental setup**

Aquaria were divided in two halves by a removable grey, opaque PVC partition. This

enabled the fish to be separated from the testing arena while the stimuli were set up. Stimuli

were displayed on vertical, grey, PVC boards and placed against one end of the aquaria. Tanks

were illuminated using the same white LED lights (EcoLight KR96 30W) used for stimulus

calibration. To ensure equal light levels in all tanks, sidewelling absolute irradiance was measured

using a calibrated OceanOptics USB2000 spectrophotometer, a 180° cosine corrector and a

400nm optic fibre cable fixed horizontally in the tank (Suppl. Material).

**Animal training**

Fish were trained to peck at the target dot using a classic conditioning approach. First, fish

were trained to pick a small piece of squid off a black or white (randomly chosen) spot (1.6 cm

diam) on the grey background corresponding to the treatment group ('bright' or 'dark', Table 1).

We trained the fish to detect target spots on both brighter and darker backgrounds to reduce

hypersensitivity through anticipation by applying the principle of 'constant stimuli' thresholds

(Colman, 2008; Laming and Laming, 1992; Pelli and Bex, 2013). We will be referring to stimuli with

greater luminance than their background as bright or brighter to facilitate reading. However, the

perception of luminance is complex, and the term brightness means specifically the perception

of surface luminance is often used wrongly and/or in confusion with lightness which refers to the

perception of surface reflectance (Kingdom, 2011). Training fish to react to stimuli being either

brighter or darker intended to produce thresholds more closely related to a natural context, as

prey items in the natural environments can be both brighter and darker than their natural

background. Second, once fish consistently removed the food reward from the black and white

target spots, a second food reward was presented from above using forceps. Once fish were

confident with this, the final stage of training was a food reward given from above once they had

tapped at the target stimulus (without food). Training consisted of up to two sessions per day,

with six to ten trials per session. Fish moved to the testing phase when fish were successful in

performing the task in > 80% trials over at least 6 consecutive sessions. A trial was considered

unsuccessful if the fish took longer than 90 seconds to make a choice or if it pecked at the

background more than twice. Testing was suspended for the day if the fish showed multiple

timeouts for obviously easy contrasts, assuming the fish was not motivated to perform the task.

However, this occurred rarely (<1% of trials) with smaller fish being more susceptible to having

been fed enough to lose appetite.

**Animal testing**

We randomly allocated fish into two groups: Group 1 (n = 7) had to find and peck at target

spots that were brighter ($T_{bd}$) or darker ($T_{dd}$) than a relatively dark background; Group 2 (n = 8)

had to find and peck target spots that were brighter ($T_{bb}$) or darker ($T_{db}$) than a relatively bright

background (Fig. 2, Table 1). As with the training of the animals, the target spots were presented

in a random position against an A4 sized achromatic background in two sessions per day

consisting of 6-10 trials per session depending on the appetite of the fish. The trials for each

session were chosen pseudo-randomly from all possible contrasts, thus fish were presented with

both darker and brighter spots compared to their background in each session. Each stimulus was

presented a minimum of 6 times (Table 1). We ensured that both easier and harder contrast

stimuli were presented in each session to maintain fish motivation. Motivation was considered

low when the animal did not engage in the trial immediately, and if this occurred, trials were

ceased for that fish until the next session. However, this rarely occurred and was further

minimised by carefully avoiding overfeeding the animals. As per training, trial was considered

unsuccessful if the fish took longer than 90 seconds to make a choice or if it pecked at the

background more than twice. Wrong pecks were recorded and time to detection was recorded as

the time between the moment the fish moved past the divider and the successful peck at the

target spot.

**Figure 2:** Schematic representation of detection scenarios. Figure proportions are not to scale. Group 1 (dark background) and group 2 (bright background) are shown with darker and brighter target spots with the maximum contrast used in the experiment. The top left of each scenario shows the corresponding abbreviation. $T_{bd}$ = bright spot on dark background, $T_{dd}$ = dark spot on dark background, $T_{bb}$ = bright spot on bright background, $T_{db}$ = dark spot on bright background. Background were A4 size and the spots 1.6cm in diameter, randomly placed of each trial.

| Group 1 (dark background, n = 7) [ΔS] / [Michelson Contrast] (median trials per fish ± sd) | | Group 2 (bright background, n = 8) [ΔS] / [Michelson Contrast] (median trials per fish ± sd) | |
|---|---|---|---|
| Bright Spot ($T_{bd}$) | Dark Spot ($T_{dd}$) | Bright Spot ($T_{bb}$) | Dark Spot ($T_{db}$) |
| 15.34 / 0.37* (6.0 ± 0.5) | 9.26 / 0.23* (7.0 ± 1.1) | 17.87 / 0.42* (8.5 ± 1.5) | 15.51 / 0.37* (9.0 ± 1.8) |
| 5.98 / 0.15 (8.0 ± 1.3) | 6.55 / 0.16 (6.0 ± 0.4) | 8.84 / 0.22 (8.5 ± 1.8) | 7.99 / 0.20 (8.0 ± 1.3) |
| 4.82 / 0.12 (6.0 ± 0.5) | 5.04 / 0.13 (8.0 ± 1.7) | 5.19 / 0.13 (7.5 ± 0.9) | 5.92 / 0.15 (8.5 ± 1.5) |
| 3.94 / 0.10 (8.0 ± 1.5) | 3.03 / 0.08 (9.0 ± 1.5) | 3.98 / 0.10 (9.0 ± 1.7) | 4.65 / 0.12 (8.5 ± 1.3) |
| 2.34 / 0.06 (8.0 ± 1.2) | 1.24 / 0.03 (9.0 ± 1.6) | 1.82 / 0.05 (8.0 ± 1.3) | 2.46 / 0.06 (6.0 ± 1.6) |
| 0.58 / 0.01 (7.0 ± 1.1) | 0.89 / 0.02 (9.0 ± 1.6) | 0.84 / 0.02 (6.5 ± 1.0) | 1.58 / 0.04 (7.0 ± 1.4) |

**Table 1:** Summary of all stimulus contrasts across both groups in ΔS and Michelson contrast. Number of trials per fish are indicated in brackets below each stimulus contrast.

**Statistical analysis**

Psychometric curves were fitted to the data with % correct choice per stimulus as the

response variable and stimulus contrast measured in Michelson contrast as the independent

variable, using the R package *quickpsy* (Linares and Lopez-Moliner, 2015; R Core Team, 2015). The

best model fit (cumulative normal or logistic) was determined using the lowest AIC as per Yssaad-

Fesselier & Knoblauch (2006) and Linares & Lopez-Moliner (2015) and is expressed both

individually for each scenario as well as the sum across all scenarios. We interpolated the 50%

correct choice thresholds with a 95% confidence interval from these curves. Thresholds between

the fitted curves for each scenario were compared as per Jörges *et al.* (2018) using the Bootstrap

(Boos, 2003) implemented in *quickpsy* (100 permutations). The Bonferroni method (Bland and

Altman, 1995) was used to adjust the significance level of the confidence intervals to 1-0.05/n,

with n corresponding to the number of comparisons.

**Results**

A total of 1365 trials were conducted across all animals and treatments (Table 1). The total success

rate was 68.5% across all 24 stimuli with a median (± sd) time to detection of 3.1 ± 12.6 s with the

fastest success at 0.3 seconds and the slowest at 89.9 s. The median time for successful detection

was similar across all scenarios (± sd): $T_{dd}$ = 2.9 ± 12.9 s, $T_{bd}$ = 2.8±10.8 seconds, $T_{db}$ = 3.1 ± 13.5 s,

$T_{bb}$ = 3.22 ± 12.58 s.

Detection thresholds (50% correct choice) for all scenarios are presented in Figure 3 and

Table 2. The sum of AIC across all four detection scenarios (fit = cumulative normal) was 162.4

($T_{dd}$ = 24.2, $T_{bd}$ = 50.8, $T_{bb}$ = 50.1, $T_{db}$ = 37.3). In group 1 (dark background), the detection thresholds

for the bright and dark spot were not significantly different from each other, with the threshold

for detecting a spot brighter than a dark background being slightly higher than a spot darker than

a dark background ($T_{bd}$ - $T_{dd}$ = 0.007 MC, $CI_{diff}$ [0.002 / 0.017]). However, the detection thresholds

in group 2 (bright background) were significantly different from each other, with the threshold

for detecting a dark spot against a bright background being significantly lower than the threshold

for detecting a bright spot against a bright background ($T_{db}$ - $T_{bb}$ = -0.028MC, $CI_{diff}$ [0.014 / 0.041]).

While the threshold for detecting a bright spot against a dark background was not

different from that for detecting a dark spot against a bright background (its 'inverse' scenario)

($T_{bd}$ -$T_{db}$ =-0.003, $CI_{diff}$ [-0.013 / -0.016]) all other detection thresholds varied significantly from

each other when compared across group 1 & 2 (Fig. 3 & 4, Table 2).

**Figure 3:** 50% probability of a fish successfully pecking a target spot. Estimated using a logistic regression fitted to the detection data. Thresholds for each scenario in Michelson contrast, error bars represent the 95% confidence intervals.

**Figure 4:** Detection thresholds for individual fish. Individually estimated discrimination thresholds in Michelson contrast for each scenario. Error bars represent the 95% confidence intervals.

| | | Scenario Significance (MC / ΔS) | Michelson Contrast (95% CI) | Weber Contrast (95% CI) | ΔS (95% CI) |
|---|---|---|---|---|---|
| $T_{bd}$ | **Group 1** | (abd / a)  | 0.063 (0.057-0.071) | 0.313 (0.282-0.349) | 2.543 (2.286-2.831) |
| $T_{dd}$ | | (b / a)  | 0.056 (0.051-0.063) | 0.278 (0.241-0.309) | 2.252 (1.955-2.510) |
| $T_{bb}$ | **Group 2** | (c / b)  | 0.095 (0.086-0.104) | 0.322 (0.287-0.354) | 3.799 (3.379-4.118) |
| $T_{db}$ | | (d / a)  | 0.066 (0.060-0.073) | 0.226 (0.197-0.253) | 2.662 (2.317-2.979) |

**Table 2:** Summary of results for the 50% correct choice threshold contrasts. Letters above scenario drawings indicate significant differences in MC thresholds as per bootstrap sampling (letters on the left) or a 1 ΔS RNL contrast (letters on the right).

## Discussion

Our study demonstrates that for triggerfish, *Rhinecanthus aculeatus*, the ability to discriminate a large, well-illuminated achromatic stimulus against a uniform achromatic background depends on both the relative luminance contrast between target and background ($f_t$ vs. $f_b$) as well as the absolute luminance level ($f_t + f_b$) at which the contrast is perceived (eq. 2). For example, discrimination thresholds, measured as Michelson contrast (MC), were significantly lower when fish were presented with a bright spot against a dark background, as opposed to a bright spot against a bright background (Table 2). However, when expressed in terms of Weber contrast (i.e. scaling the contrast with the luminance level at which the luminance contrast is perceived) these two thresholds were almost identical (Table 2). This finding supports the Weber-Fechner law that states the ability to discriminate a target stimulus against its background scales with the intensity at which the discrimination is made. The same holds true for the discrimination thresholds of dark spot against a dark background ($T_{dd}$) as opposed to a bright background ($T_{db}$), which have an almost identical Weber contrast (Table 2). Furthermore, the contrast sensitivity depends on the direction of the contrast ($f_t > f_b \neq f_t < f_b$), that is, the Weber contrast for detecting stimuli darker than their respective backgrounds is lower ( = more sensitive) from that for stimuli which are brighter than their backgrounds (WC 0.23 − 0.28 for dark spots and 0.31 − 0.32 for bright ones) (Table 2).

Our results agree with previous findings that humans (e.g. Bowen, Pokorny, & Smith, 1989; Emran et al., 2007; Lu & Sperling, 2012), non-human vertebrates (e.g. Baylor et al., 1974), and invertebrate visual systems (e.g. Smithers et al., 2019) are consistently better at detecting darker stimuli. Increasing and decreasing luminance changes are thought to be processed

differently: darker stimuli are detected by off-centre ganglion cells, while lighter ones are

detected by on-centre ganglion cells (Schiller et al., 1986). Dark stimuli cause depolarization of

photoreceptors, whereas light ones are detected as hyperpolarization (Baylor et al., 1974). For

example, investigation of turtle photoreceptors has shown that dark stimuli result in much

greater depolarization of photoreceptors, than the magnitude of hyperpolarization resulting from

light ones (Baylor et al., 1974).This asymmetry is thought to be a crucial contributor to object and

motion detection in post-retinal processing (e.g. Oluk et al., 2016; Vidyasagar and Eysel, 2015).

**Behavioural calibration of the RNL**

The relationship of absolute (background + stimulus) and relative luminance (background

vs. stimulus) contrast does not hold when expressing thresholds as achromatic ΔS (Table 2). The

exclusion of signal intensity is a fundamental assumption when calculating chromatic contrasts

using the RNL model (Vorobyev and Osorio, 1998), which was designed to quantify contrast

perception between two closely opposed chromatic stimuli viewed against an achromatic

background. As a result, the RNL equations used by Siddiqi et al. (2004) calculate a relative

comparison of two background adapted receptor responses without scaling the difference in

photoreceptor stimulation between stimulus and background in relation to the overall brightness

of a scene. Thus, the commonly used RNL equations in Siddiqi *et al.* (2004) fail to reflect the

Weber-Fechner law for the discrimination of a stimulus from its background. Olsson et al. (2018)

proposed the use of an adaptation where the Weber contrast at the behaviourally determined

discrimination threshold ($WC_t$) should be used in place of the receptor noise:

$$\Delta S = |\Delta f_{dbl}/WC_t|$$ **Equation 3**

This renders the following ΔS values at threshold: $T_{dd}$ = $T_{bd}$ = 0.41 ΔS ± 0.0001 and $T_{bb}$ = $T_{db}$

= 0.59 ΔS ± 0.001 using the WC determined in this experiment. This makes the RNL model, as

modified by Olsson et al. 2018, conform with the Weber-Fechner law while preserving the

difference in contrast sensitivity regarding increments and decrements. Furthermore, the

thresholds are well below 1 ΔS, making the assumption of a 'Just Noticeable Difference' (JND)

corresponding to a threshold of 1 ΔS a comfortably conservative (but not extreme) threshold. It

should be noted that the general conclusions of Siddiqi et al., (2004) remain most likely correct,

but we can now realise a closer description of the underlying mechanisms.

Olsson et al. (2018) propose the use of Michelson contrast (MC) in place of receptor noise

in order to estimate the channel specific noise ($e_i$). First, the contrast sensitivity (CS) is calculated

as the inverse of the behaviourally determined Michelson Contrast ($C_t$):

393
$$CS = \frac{1}{C_t}$$
**Equation 4**

394     Next, this CS (which is sensitive to the absolute level of luminance as our results confirm)

395     can be used to calculate the relative quantum catch of stimulus 2 ($q_{stim2}$):

396
$$q_{stim2} = \frac{1 - C_t}{1 + C_t}$$
**Equation 5**

397     We can then use $q_{stim2}$ as $f_t$ and our originally measured $f_b$ in eq. 2 to derive the channel

398     noise ($e_i$) (see Olsson et al. 2018 for further details). With the assumption of ΔS = 1 at threshold,

399     this produces:

400
$$e_i = \ln\frac{\Delta S}{q_{stim2}}$$
**Equation 6**

Thus, we obtain the following channel noise estimates ($e_i \pm$ 95% CI): $e_{Tdd}$ = 0.113 (0.098 –

0.125) & $e_{Tbd}$ = 0.127 (0.114 – 0.142) for group 1 (dark background) and $e_{Tdb}$ = 0.133 (0.116 –

0.149) & $e_{Tbb}$ = 0.190 (0.169 – 0.209) for group 2 (bright background). This is the same as setting

ω (which is equal to $e_i$) in eq. 1 so that the observed ΔS in our experiments (Table 2) would be

equal to 1 (as we determined $f_t$ at threshold by fitting a model to the observed MC). This is

interesting, as these noise values are up to almost 4 times as high as the 'conservative' (!) standard

deviation of noise estimate of 0.05, currently used for modelling vertebrate vision across the field

of visual ecology.

Given that WC is meant to be used for comparably small stimuli against large backgrounds

and MC to be used for contrasts between stimuli of comparable size, we recommend a

differentiated use of either equation 3 or 4-6 depending on the visual context in which a

discrimination threshold is used. For example, as the scenario in this study involved the

discrimination of a single spot against a much larger background, we would assume equation 3 to

be more relevant than equations 4-6 (e.g. equations 4-6 still produce a higher noise ratio for

Group 2 (bright background), especially $T_{bb}$). Thus, equations 4-6 would likely be more relevant

when discriminating between two objects of equal size. This further implies that one could plot

the discrimination curves as a function of WC rather than MC to obtain the discrimination

threshold. The thresholds would then only be distinguishable based on the relative direction of

the contrast (bright spot or dark spot) and not the background intensity (Table 2). It also implies

that thresholds obtained from experiments using a discrimination scenario more fitting to

equations 4-6 (e.g. Lind et al., 2013) should not be used to infer the detectability of most likely

relatively small prey items against their most likely large visual backgrounds.

**Future directions**

The specific mechanisms causing the observed difference in WC between the detection of

a dark spot and a bright spot (or mathematical approximations thereof), or an explanation as to

why $e_{achromatic}$ is much higher than the conservatively chosen receptor noise of 0.05, remain

speculative. Further investigations might seek advances in the understanding of

neurophysiological mechanisms underlying luminance contrast perception in *R. aculeatus*. These

include knowledge of the detailed anatomy and receptor noise of double cone photoreceptors,

the relative contribution of each double-cone member to luminance contrast sensitivity (Siebeck

et al., 2014) as well as the precise mechanism by which photoreceptor stimulation is integrated

in post-receptor structures such as edge detecting receptive fields. Behavioural experiments with

closely related species with different retinal morphologies would be of interest to further

investigate e.g. the role of retinal neuroanatomy on luminance contrast perception.

The adaptations to the RNL model in Olsson et al. (2018), while apparently effective, do

not account for the effects of spatial frequency on luminance contrast sensitivity when

discriminating objects against visual backgrounds. This is probably the most notable confounding

effect on low-level processing of luminance contrast as a result of post-receptor lateral-inhibition

(Veale et al., 2017). One possible approach would be the use of contrast sensitivity functions (CSF)

to scale Weber fractions as a function of spatial frequency in a visual scene. However, given that

these are determined using a perceptually different experimental setup (da Silva Souza et al.,

2011) this should be investigated using context specific behavioural experimentation.

Our results warrant caution in the use of uniform contrast sensitivity thresholds (be it

achromatic or chromatic) across widely diverse perceptual contexts, independently of which

models are used to describe them. Luminance discrimination, as expected, is not just limited by

photoreceptor noise and therefore cannot be adequately represented by the use of a singular

detection or discrimination threshold determined using the equations in Siddiqi et al. (2004) as

currently common in behavioural ecology studies. This realisation shares many parallels with

ongoing discussions regarding the use of the RNL model outside of model assumptions (Marshall,

2018; Olsson et al., 2018; Osorio et al., 2017; Sibeaux et al., 2019; Stuart-Fox, 2018; Vasas et al.,

2018). Our results suggest the use of conservative achromatic RNL threshold assumption of 3ΔS

(e.g. Spottiswoode & Stevens, 2010) without adaptations such as those proposed by Olsson et al.

(2018) might warrant caution.

We show that the noise in the achromatic channel of *R. aculeatus* can be substantially

higher than anticipated in previous studies modelling its luminance contrast sensitivity using

'conservative' receptor noise estimates. However, this increase in channel noise ($e_i$) can be

originating from many potential sources, including electrophysiological coupling of receptors in

the double cone of *R.aculeatus* (but also a generally higher noise level in receptor responsible for

luminance contrast detection) or downstream (post-receptor) processing of visual information.

As such it is wrong to conclude receptor noise from such behavioural calibration (Vasas et al.,

2018) and it would be more appropriate to refer to the noise of the entire pathway involved in

the performance of a task based on the animal's ability to perceive luminance contrast in a

specific visual context.

Despite having investigated luminance contrast sensitivity using two different levels of

background luminance, our study only considered discrimination of large, uniform and

achromatic circular target stimulus against a uniform grey background. In future studies, more

realistic backgrounds and illumination should be taken into account (e.g. Matchette et al., 2020),

as a variety of factors can fundamentally influence luminance contrast perception in most

circumstances (Gilchrist, 2014; Gilchrist and Radonjic, 2009; Gilchrist et al., 1999; Kingdom, 2011;

Maniatis, 2014). Unsurprisingly then, there is evidence that luminance contrast modulates the

salience of objects at stages well beyond the retina (Einhäuser and König, 2003).

**Summary**

Our findings provide insight into the processing of achromatic information as well as the

use of the RNL model to quantify achromatic discrimination by non-human observers. We show

that the current use of the RNL model for the quantification of luminance contrast sensitivity

thresholds warrants caution. More specifically, our study suggests the lack of adequate scaling of

thresholds by the RNL model to the average luminance of a scene and the need for context

specific behavioural experimentation whenever possible.

One of the main reasons why researchers use the RNL model is that, presumably, the

discriminability of visual contrasts can be reliably predicted by using a set of conservatively

estimated physiological parameters such as photoreceptor noise, abundance and spectral

sensitivity. While this seems to work satisfyingly well for colour contrast perception across a range

of animals, our study suggests quite the opposite to be the case for achromatic contrast. Despite

the possibility of calibrating the RNL using contextualised behavioural experiments (as suggested

by Olsson et al. 2018), the result remains unsatisfying. However, we recommend the use of

behaviourally determined discrimination thresholds suitable to the given visual context in which

they are to be applied as well as generous caution when predicting the discriminability of

luminance contrast.

Our study indeed suggests that one cannot reliably use the RNL to predict achromatic

contrast perception without context specific behavioural experimentation. This has direct

implications on the design of behavioural experiments where validated discrimination thresholds

are unavailable. For example, given the difficulty of predicting luminance discriminability,

luminance contrast should be thoroughly randomised (as opposed to attempting iso-luminance

between stimuli) in any behavioural experiment than can potentially be influenced by luminance

contrast perception.

## Contributions
497

498    Cedric P. van den Berg conceived and conducted the study and wrote the manuscript. Laurie J.

499    Mitchell, Michelle Hollenkamp and Erin Watson assisted with training and testing of the animals.

500    Karen L. Cheney, Naomi F. Green and N. Justin Marshall helped review the document.

## Acknowledgements
502

503    We would like to thank Natalie Meiklejohn for assisting with experiments and data entry, Ama

504    Wakwella for preliminary experimentation and many volunteers for assisting with animal

505    husbandry, Almut Kelber, Anna Hughes & Simon Laughlin for valuable discussions and William

506    Allen & Nick Scott-Samuel for valuable feedback on the manuscript.

## Funding
508

509    This study was funded by an ARC Discovery Grant DP150102710 awarded to Karen Cheney and

510    N. Justin Marshall.

**Conflict of interest**

We declare to have no conflicting interests.

**References**

**Anderson, B. L.** (2011). Visual perception of materials and surfaces. *Curr. Biol.* **21**, R978-83.

**Baylor, D. A., Hodgkin, A. L. and Lamb, T. D.** (1974). The electrical response of turtle cones to

flashes and steps of light. *J. Physiol.* **242**, 685–727.

**Bex, P. J. and Makous, W.** (2002). Spatial frequency, phase, and the contrast of natural images.

*J. Opt. Soc. Am. A* **19**, 1096.

**Bland, j. M. and Altman, D. G.** (1995). Statistics notes: Multiple significance tests: the

Bonferroni method. *BMJ* **310**, 170–170.

**Boos, D. D.** (2003). Introduction to the Bootstrap World. *Stat. Sci.* **18**, 168–174.

**Bowen, R. W., Pokorny, J. and Smith, V. C.** (1989). Sawtooth contrast sensitivity: decrements

have the edge. *Vision Res.* **29**, IN1-1509.

**Brooks, J. L.** (2014). *Traditional and New Principles of Perceptual Grouping*. (ed. Wagemans, J.)

Oxford University Press.

**Bruce, V., Green, P. R., Georgeson, M. A. and Dynan, L.** (2010). *Visual perception: physiology,*

*psychology, & ecology*. Fourth edi. Hove, England ; New York, New York: Psychology Press.

**Caves, E. M. and Johnsen, S.** (2018). AcuityView: An r package for portraying the effects of

visual acuity on scenes observed by an animal. *Methods Ecol. Evol.* **9**, 793–797.

**Champ, C., Wallis, G., Vorobyev, M., Siebeck, U. and Marshall, J.** (2014). Visual acuity in a

species of coral reef fish: Rhinecanthus aculeatus. *Brain. Behav. Evol.* **83**, 31–42.

**Champ, C. M., Vorobyev, M. and Marshall, N. J.** (2016). Colour thresholds in a coral reef fish. *R.*

*Soc. Open Sci.* **3**, 160399.

**Cheney, K. L. and Marshall, N. J.** (2009). Mimicry in coral reef fish: how accurate is this

deception in terms of color and luminance? *Behav. Ecol.* **20**, 459–468.

**Cheney, K. L., Newport, C., McClure, E. C. and Marshall, N. J.** (2013). Colour vision and response

bias in a coral reef fish. *J. Exp. Biol.* **216**, 2967–2973.

**Cheney, K. L., Green, N. F., Vibert, A. P., Vorobyev, M., Marshall, N. J., Osorio, D. C. and**

**Endler, J. A.** (2019). An Ishihara-style test of animal colour vision. *J. Exp. Biol.* **222**,

jeb189787.

**Colman, A. M.** (2008). *A Dictionary of Psychology*. Oxford University Press.

**Corney, D. and Lotto, R. B.** (2007). What are lightness illusions and why do we see them? *PLoS*

*Comput. Biol.* **3**, e180.

**Cornsweet, T. N. and Pinsker, H. M.** (1965). Luminance discrimination of brief flashes under

various conditions of adaptation. *J. Physiol.* **176**, 294–310.

**Craik, K. J. W.** (1938). The effect of adaptation on differential brightness discrimination. *J.*

*Physiol.* **92**, 406–421.

**da Silva Souza, G., Gomes, B. D. and Silveira, L. C. L.** (2011). Comparative neurophysiology of

spatial luminance contrast sensitivity. *Psychol. Neurosci.* **4**, 29–48.

**Dzhafarov, E. N. and Colonius, H.** (1999). Fechnerian metrics in unidimensional and

multidimensional stimulus spaces. *Psychon. Bull. Rev.* **6**, 239–268.

**Einhäuser, W. and König, P.** (2003). Does luminance-contrast contribute to a saliency map for

overt visual attention? *Eur. J. Neurosci.* **17**, 1089–1097.

**Elder, J. H. and Sachs, A. J.** (2004). Psychophysical receptive fields of edge detection

mechanisms. *Vision Res.* **44**, 795–813.

**Elder, J. H. and Velisavljevic, L.** (2010). Cue dynamics underlying rapid detection of animals in

natural scenes. *J. Vis.* **9**, 787–787.

**Emran, F., Rihel, J., Adolph, A. R., Wong, K. Y., Kraves, S. and Dowling, J. E.** (2007). OFF

ganglion cells cannot drive the optokinetic reflex in zebrafish. *Proc. Natl. Acad. Sci.* **104**,

19126–19131.

**Endler, J. A.** (1991). Interactions between predator and prey. In *Behavioural Ecology* (ed. JA, K.)

and NB, D.), pp. 169–196. Oxford: Blackwell Scientific.

**Escobar-Camacho, D., Taylor, M. A., Cheney, K. L., Green, N. F., Marshall, N. J. and Carleton, K.**

**L.** (2019). Color discrimination thresholds in a cichlid fish: Metriaclima benetos. *J. Exp. Biol.*

**222**, jeb201160.

**Gegenfurtner, K. R. and Kiper, D. C.** (1992). Contrast detection in luminance and chromatic

noise. *J Opt. Soc. Am. A.* **9**, 1880–1888.

**Geisbauer, G., Griebel, U., Schmid, A. and Timney, B.** (2004). Brightness discrimination and

neutral point testing in the horse. *Can. J. Zool.* **82**, 660–670.

**Gilchrist, A. L.** (2008). Perceptual organization in lightness. In *Oxford Handbook of Perceptual*

*Organisation*, pp. 1–25.

**Gilchrist, A.** (2014). Response to Maniatis critique of anchoring theory. *Vision Res.* **102**, 93–96.

**Gilchrist, A. L. and Radonjic, A.** (2009). Anchoring of lightness values by relative luminance and

relative area. *J. Vis.* **9**, 13–13.

**Gilchrist, A., Kossyfidis, C., Bonato, F., Agostini, T., Cataliotti, J., Li, X., Spehar, B., Annan, V.**

**and Economou, E.** (1999). An anchoring theory of lightness perception. *Psychol. Rev.* **106**,

795–834.

**Green, N. F., Urquhart, H. H., van den Berg, C. P., Marshall, N. J. and Cheney, K. L.** (2018).

Pattern edges improve predator learning of aposematic signals. *Behav. Ecol.* **29**, 1–6.

**Griebel, U. and Schmid, A.** (1997). Brightness discrimination ability in the West Indian manatee

(Trichechus manatus). *J. Exp. Biol.* **200**, 1587–92.

**Heath, S. L., Christenson, M. P., Oriol, E., Saavedra-Weisenhaus, M., Kohn, J. R. and Behnia, R.**

(2020). Circuit Mechanisms Underlying Chromatic Encoding in Drosophila Photoreceptors.

*Curr. Biol.* **30**, 264-275.e8.

**Heinemann, E. G. and Chase, S.** (1995). A quantitative model for simultaneous brightness

induction. *Vision Res.* **35**, 2007–2020.

**Hempel de Ibarra, N., Giurfa, M. and Vorobyev, M.** (2001). Detection of coloured patterns by

honeybees through chromatic and achromatic cues. *J. Comp. Physiol. A Sensory, Neural,*

*Behav. Physiol.* **187**, 215–224.

**Jörges, B., Hagenfeld, L. and López-Moliner, J.** (2018). The use of visual cues in gravity

judgements on parabolic motion. *Vision Res.* **149**, 47–58.

**Kelber, A., Vorobyev, M. and Osorio, D. C.** (2003). Animal colour vision – behavioural tests and

physiological concepts. *Biol. Rev. Camb. Philos. Soc.* **78**, S1464793102005985.

**Kingdom, F. A. A.** (2011). Lightness, brightness and transparency: A quarter century of new

ideas, captivating demonstrations and unrelenting controversy. *Vision Res.* **51**, 652–673.

**Laming, D. and Laming, J.** (1992). F. Hegelmaier: On memory for the length of a line. *Psychol.*

*Res.* **54**, 233–239.

**Land, E. H.** (1986). Recent advances in retinex theory. *Vision Res.* **26**, 7–21.

**Linares, D. and Lopez-Moliner, J.** (2015). quickpsy: An R package to fit psychometric functions

for multiple groups. 1–11.

**Lind, O., Sunesson, T., Mitkus, M. and Kelber, A.** (2012). Luminance-dependence of spatial

vision in budgerigars (Melopsittacus undulatus) and Bourke's parrots (Neopsephotus

bourkii). *J. Comp. Physiol. A* **198**, 69–77.

**Lind, O., Karlsson, S. and Kelber, A.** (2013). Brightness discrimination in budgerigars

(Melopsittacus undulatus). *PLoS One* **8**, e54650.

**Lotto, R. B. and Purves, D.** (2000). An empirical explanation of color contrast. *Proc. Natl. Acad.*

*Sci.* **97**, 12834–12839.

**Lu, Z.-L. and Sperling, G.** (2012). Black-white asymmetry in visual perception. *J. Vis.* **12**, 8–8.

**Lythgoe, J. N.** (1979). *The ecology of vision*. Oxford University Press.

**Maniatis, L. M.** (2014). A theory divided: Current representations of the anchoring theory of

lightness contradict the original's core claims. *Vision Res.* **102**, 89–92.

**Marchiafava, P. L.** (1985). Cell coupling in double cones of the fish retina. *Proc. R. Soc. London.*

*Ser. B. Biol. Sci.* **226**, 211–215.

**Marshall, N. J.** (2018). Do not be distracted by pretty colors: A comment on Olsson et al. *Behav.*

*Ecol.* **29**, 286–287.

**Marshall, N. J., Jennings, K., McFarland, W. N., Loew, E. R. and Losey, G. S.** (2003). Visual

biology of hawaiian coral reef fishes. III. Environmental light and an integrated approach to

the ecology of reef fish vision. *Copeia* **2003**, 467–480.

**Marshall, K. L. A., Philpot, K. E. and Stevens, M.** (2016). Microhabitat choice in island lizards

enhances camouflage against avian predators. *Sci. Rep.* **6**, 19815.

**Matchette, S. R., Cuthill, I. C., Cheney, K. L., Marshall, N. J. and Scott-Samuel, N. E.** (2020).

Underwater caustics disrupt prey detection by a reef fish. *Proceedings. Biol. Sci.* **287**,

20192453.

**MathWorks** (2000). Matlab.

**Mitchell, L., Cheney, K. L., Cortesi, F., Marshall, N. J. and Vorobyev, M.** (2017). Triggerfish uses

chromaticity and lightness for object segregation. *R. Soc. Open Sci.* **4**, 171440.

**Monnier, P. and Shevell, S. K.** (2003). Large shifts in color appearance from patterned chromatic

backgrounds. *Nat. Neurosci.* **6**, 801–802.

**Moulden, B., Kingdom, F. and Gatley, L. F.** (1990). The standard deviation of luminance as a

metric for contrast in random-dot images. *Perception* **19**, 79–101.

**Newport, C., Green, N. F., McClure, E. C., Osorio, D. C., Vorobyev, M., Marshall, N. J. and**

**Cheney, K. L.** (2017). Fish use colour to learn compound visual signals. *Anim. Behav.* **125**,

93–100.

**Norwich, K. H.** (1987). On the theory of Weber fractions. *Percept. Psychophys.* **42**, 286–298.

**Olsson, P., Lind, O. and Kelber, A.** (2018). Chromatic and achromatic vision: parameter choice

and limitations for reliable model predictions. *Behav. Ecol.* **29**, 273–282.

**Oluk, C., Pavan, A. and Kafaligonul, H.** (2016). Rapid motion adaptation reveals the temporal

dynamics of spatiotemporal correlation between on and off pathways. *Sci. Rep.* **6**, 1–10.

**Osorio, D. and Vorobyev, M.** (2005). Photoreceptor spectral sensitivities in terrestrial animals:

Adaptations for luminance and colour Vision. *R. Soc.* **272**, 1745–1752.

**Osorio, D. and Vorobyev, M.** (2008). A review of the evolution of animal colour vision and visual

communication signals. *Vision Res.* **48**, 2042–2051.

**Osorio, D., Vorobyev, M. and Simmons, L.** (2017). Principles and application of the receptor

noise model of color discrimination: A comment on Olsson et al. *Behav. Ecol.* **29**, 1–2.

**Pelli, D. G. and Bex, P.** (2013). Measuring contrast sensitivity. *Vision Res.* **90**, 10–14.

**Pignatelli, V. and Marshall, J.** (2010). Morphological characterization of retinal bipolar cells in

the marine teleost Rhinecanthus aculeatus. *J. Comp. Neurol.* **518**, 3117–3129.

**Pignatelli, V., Champ, C., Marshall, J. and Vorobyev, M.** (2010). Double cones are used for

colour discrimination in the reef fish, Rhinecanthus aculeatus. *Biol. Lett.* **6**, 537–539.

**Pretterer, G., Bubna-Littitz, H., Windischbauer, G., Gabler, C. and Griebel, U.** (2004). Brightness

discrimination in the dog. *J. Vis.* **4**, 10.

**R Core Team** (2015). R: A language and environment for statistical computing.

**Randall, J. E., Allen, G. R. and Steene, R. C.** (1997). *Fishes of the Great Barrier Reef and Coral*

*Sea*. Crawford House Press.

**Schiller, P. H., Sandell, J. H. and Maunsell, J. H. R.** (1986). Functions of the On and Off channels

of the visual system. *Nature* **322**, 824–825.

**Scholtyssek, C. and Dehnhardt, G.** (2013). Brightness discrimination in the South African fur seal

(Arctocephalus pusillus). *Vision Res.* **84**, 26–32.

**Scholtyssek, C., Kelber, A. and Dehnhardt, G.** (2008). Brightness discrimination in the harbor

seal (Phoca vitulina). *Vision Res.* **48**, 96–103.

**Shapley, R. and Hawken, M. J.** (2011). Color in the Cortex: single- and double-opponent cells.

*Vision Res.* **51**, 701–717.

**Shevell, S. K. and Kingdom, F. A. A.** (2008). Color in Complex Scenes. *Annu. Rev. Psychol.* **59**,

143–166.

**Sibeaux, A., Cole, G. L. and Endler, J. A.** (2019). Success of the receptor noise model in

predicting colour discrimination in guppies depends upon the colours tested. *Vision Res.*

**159**, 86–95.

**Siddiqi, A.** (2004). Interspecific and intraspecific views of color signals in the strawberry poison

frog Dendrobates pumilio. *J. Exp. Biol.* **207**, 2471–2485.

**Siebeck, U. E., Wallis, G. M., Litherland, L., Ganeshina, O. and Vorobyev, M.** (2014). Spectral

and spatial selectivity of luminance vision in reef fish. *Front. Neural Circuits* **8**, 118.

**Simmons, D. R. and Kingdom, F. A. A.** (2002). Interactions between chromatic- and luminance-

contrast-sensitive stereopsis mechanisms. *Vision Res.* **42**, 1535–1545.

**Simpson, E. E., Marshall, N. J. and Cheney, K. L.** (2016). Coral reef fish perceive lightness

illusions. *Sci. Rep.* **6**, 35335.

**Skorupski, P. and Chittka, L.** (2011). Is colour cognitive? *Opt. Laser Technol.* **43**, 251–260.

**Smithers, S. P., Roberts, N. W. and How, M. J.** (2019). Parallel processing of polarization and

intensity information in fiddler crab vision. *Sci. Adv.* **5**, eaax3572.

**Spottiswoode, C. N. and Stevens, M.** (2010). Visual modeling shows that avian host parents use

multiple visual cues in rejecting parasitic eggs. *Proc. Natl. Acad. Sci.* **107**, 8672–8676.

**Stevens, M., Parraga, C. A., Cuthill, I. C., Partridge, J. C. and Troscianko, T. S.** (2007). Using

digital photography to study animal coloration. *Biol. J. Linn. Soc.* **90**, 211–237.

**Stoddard, M. C. and Stevens, M.** (2010). Pattern mimicry of host eggs by the common cuckoo,

as seen through a bird's eye. *Proc. R. Soc. B Biol. Sci.* **277**, 1387–1393.

**Stuart-Fox, D.** (2018). Opening the "black box" of modeling animal color vision: A comment on

Olsson et al. *Behav. Ecol.* **29**, 284.

**Treisman, M.** (1964). Noise and Weber's law: The discrimination of brightness and other

dimensions. *Psychol. Rev.* **71**, 314–330.

**Troscianko, J. and Stevens, M.** (2015). Image calibration and analysis toolbox – a free software

suite for objectively measuring reflectance, colour and pattern. *Methods Ecol. Evol.* **6**,

1320–1331.

**Troscianko, J., Wilson-Aggarwal, J., Stevens, M. and Spottiswoode, C. N.** (2016). Camouflage

predicts survival in ground-nesting birds. *Sci. Rep.* **6**, 19966.

**van den Berg, C. P., Troscianko, J., Endler, J. A., Marshall, N. J. J. and Cheney, K. L.** (2020).

Quantitative Colour Pattern Analysis (QCPA): A comprehensive framework for the analysis

of colour patterns in nature. *Methods Ecol. Evol.* **11**, 316–332.

**Vasas, V., Brebner, J. S. and Chittka, L.** (2018). Color discrimination is not just limited by

photoreceptor noise: a comment on Olsson et al. *Behav. Ecol.* **29**, 285–286.

**Veale, R., Hafed, Z. M. and Yoshida, M.** (2017). How is visual salience computed in the brain?

Insights from behaviour, neurobiology and modelling. *Philos. Trans. R. Soc. B Biol. Sci.* **372**,

20160113.

**Vidyasagar, T. R. and Eysel, U. T.** (2015). Origins of feature selectivities and maps in the

mammalian primary visual cortex. *Trends Neurosci.* **38**, 475–485.

**Vorobyev, M. and Osorio, D.** (1998). Receptor noise as a determinant of colour thresholds.

*Proc. R. Soc. London. Ser. B Biol. Sci.* **265**, 351–358.

**Vorobyev, M., Brandt, R., Peitsch, D., Laughlin, S. B. and Menzel, R.** (2001). Colour thresholds

and receptor noise: behaviour and physiology compared. *Vision Res.* **41**, 639–653.

**Wallach, H.** (1948). Brightness constancy and the nature of achromatic colors. *J. Exp. Psychol.*

711            **38**, 310–324.

712      **Webster, M. A. and Wilson, J. A.** (2000). Interactions between chromatic adaptation and

713            contrast adaptation in color appearance. *Vision Res.* **40**, 3801–3816.

714      **Yssaad-Fesselier, R. and Knoblauch, K.** (2006). Modeling psychometric functions in R. *Behav.*

715            *Res. Methods* **38**, 28–41.

716      **Zeki, S.** (1993). *A vision of the brain*. Oxford, England: Blackwell Scientific.

717      **Zhou, M., Bear, J., Roberts, P., Janiak, F., Semmelhack, J., Yoshimatsu, T. and Baden, T.** (2020).

718            What the Zebrafish's eye tells the Zebrafish's brain: Retinal ganglion cells for prey capture

719            and colour vision. *bioRxiv* 2020.01.31.927087.

720