A warming Southern Ocean may compromise Antarctic blue whale foetus growth

Carl Smith[1,2,3*]

[1]School of Animal, Rural and Environmental Sciences, Nottingham Trent University,

Southwell, Nottinghamshire NG25 0QF, UK

[2]Institute of Vertebrate Biology, Academy of Sciences of the Czech Republic, Brno, Czech

Republic

[3]Department of Ecology and Vertebrate Zoology, University of Łódź, Łódź, Poland

[*] Corresponding author: Carl Smith, carl.smith01@ntu.ac.uk

Abstract

After declining in abundance due to commercial whaling during the 20[th] Century, populations

of the Antarctic blue whale (*Balaenoptera musculus intermedia*) have failed to recover to pre-

exploitation levels. Using historical whaling data from 1926-1954, in combination with

temperature data for the Southern Ocean, a gamma GLMM with temporal dependency was

fitted to 20,144 records of *B. m. intermedia* foetus size using Bayesian inference. There was a

negative relationship between antecedent winter sea surface temperature (SST) in the Southern

Ocean on foetus size. This relationship is proposed as being mediated by a positive effect of

the extent of winter sea ice on Antarctic krill (*Euphasia superba*) abundance on which *B. m.*

intermedia feed. There was also a positive density-dependent effect of a 'krill surplus' at low

whale population sizes. However, the positive effect of a 'krill surplus' at low *B. m. intermedia*

population size on foetus growth was reversed at elevated winter SST due to a negative impact

on *E. superba* recruitment. Projected increases in temperature in the Southern Ocean are

predicted to compromise the growth rates of *B. m. intermedia* foetuses, with implications for

the capacity of the subspecies to recover from overexploitation.

Keywords: Antarctica, cetacean, density dependence, IWC, population dynamics, recovery

Introduction

The large-scale harvest of baleen whales (Balaenopteridae) in the Southern Ocean, which began at the start of the 20th Century, saw rapid declines in the abundance of blue (*Balaenoptera musculus*), humpback (*Megaptera novaeangliae*), fin (*B. physalus*) and sei (*B. borealis*) whales (Baker & Clapham, 2004). After several decades of protection, populations of these whales have shown varying degrees of recovery (Best, 1993; Branch, Matsuoka, & Miyashita, 2004; Clapham, Young, & Brownell, 1999). *M. novaeangliae* have shown strong recovery, *B. physalus* populations have broadly shown increases in numbers, while the status of *B. borealis* populations is poorly known (Thomas, Reeves, & Brownell, 2016). *B. musculus* populations show wide variation in status and recovery; some populations have disappeared entirely or remain at extremely low levels (Thomas et al., 2016), while at least one population from the eastern North Pacific appears to have recovered to pre-exploitation levels (Monnahan, Branch, Punt, 2015). The Antarctic blue whale (*B. m. intermedia*) is classified as Critically Endangered (Reilly et al., 2008). Prior to the onset of commercial whaling the abundance of *B. m. intermedia* was estimated at approximately 239,000 individuals. Abundance declined to as few as 370 individuals, despite a moratorium on the hunting of this species in 1965 (Branch et al., 2004). The most recent conservative population estimate for this subspecies was 2,280 individuals, barely 1% of pre-exploitation levels (Branch, 2007).

The underlying reasons for the slow recovery of *B. m. intermedia* have not been identified but may, in part, arise from illegal hunting by the Soviet Union into the 1970s, as well as from on-going mortalities associated with boat strikes and entanglement in fishing gear (Clapham et al., 1999; Yablakov, 1994). An additional explanation for their slow recovery could relate to a decline in food availability, particularly of Antarctic krill (*Euphasia superba*) on which *B. m. intermedia* feed almost exclusively (Bannister, 2008). In the southern hemisphere, *B. m. intermedia* migrate to Antarctic waters to feed during the austral summer, before migrating back to lower latitudes for the remainder of the year, where feeding is typically more limited,

though there are departures from this general pattern (Hucke-Gaete et al., 2018). The period spent feeding is critical (Mackintosh, 1965; Hucke-Gaete et al., 2018), particularly as females arrive in the Southern Ocean pregnant, with foetal growth contingent on the accumulation of energy reserves by the mother on the feeding grounds (Christiansen, Víkingsson, Rasmussen, & Lusseau, 2014).

E. superba are an important species in the Southern Ocean (Fraser & Hofmann, 2003), feeding on blooms of phytoplankton at the edge of the Antarctic sea ice with their abundance closely linked to the extent of ice (Braithwaite, Meeuwig, Letessier, Jenner, & Brierley, 2015; Brierly et al., 2002). Declines in the extent and duration of Antarctic winter sea ice since the 1970s have resulted in a decline in the abundance of *E. superba* (Atkinson, Siegel, Pakhomov, & Rothery, 2004; Nicol, 2006), with potential consequences for the predators that rely on them, including baleen whales (Ainley et al., 2006, 2007; Alonzo & Mangal, 2001; Reid & Croxall, 2001). Braithwaite et al. (2015) linked the extent of sea ice with the quantity of oil extracted from the carcasses of humpback whales caught during commercial whaling between 1947-1963. They showed that the extent of winter sea ice correlated positively with oil yields, suggesting that *M. novaeangliae* body condition was directly linked to *E. superba* abundance, mediated by the extent of sea ice.

Balaenopterid whales display the highest foetal growth rates of all mammals (Frazer & Huggett, 1973), demanding substantial allocation of maternal somatic resources for embryonic and foetal growth (Lockyer, 1986). Rapid foetal growth in marine mammals permits the production of large neonates with high survival rates (Boltnev, York, & Antonelis, 1998; McMahon, Burton, & Bester, 2003) and in the case of balaenopterid whales is synchronised with their annual feeding migration from low to high latitudes (Chivers, 2008). Energy supplied to the developing foetus comes from reserves accumulated during the intense feeding phase of their annual migratory cycle. If female condition is compromised, the outcome is a decline in fertility

(Williams et al., 2013) and reduced investment in embryonic and foetal growth (Christiansen

et al., 2014), with potential negative consequences for calf survival (Lockyer, 2007).

Declines in balaenopterid population size due to whaling can release resources (a 'krill surplus')

to survivors and thereby facilitate recovery through density-dependent effects (Ainley et al.,

2007; Best, 1993; Fraser, Trivelpiece, Ainley, & Trivelpiece, 1992; Laws, 1977). Density-

dependent population growth has been demonstrated in *B. physalus* (Williams et al., 2013).

However, while there is some evidence for the modest recovery of *B. m. intermedia* (Branch,

2008), the continued low abundance of this species means that it remains critically endangered.

Here I indirectly test whether foetal growth of *B. m. intermedia* is determined by the availability

of *E. superba* by examining the effect of temperature on *B. m. intermedia* foetal body size. A

prediction is that a reduction in the extent of sea ice in response to elevated temperatures in the

Southern Ocean will suppress *E. superba* productivity thereby compromising *B. m. intermedia*

foetus development with negative consequences for their subsequent post-natal survival to

maturity. An additional prediction is that a decline in the abundance of *B. m. intermedia* will

result in a 'krill surplus', detectable as increased foetus growth rate. A further prediction is that

these processes are linked, with temperatures in the Southern Ocean interacting with the effect

of adult *B. m. intermedia* abundance. Thus, a combination of low winter temperatures combined

with low *B. m. intermedia* abundance are predicted to enhance foetal growth rates while the

converse, elevated winter temperatures and high adult abundances, would tend to supress foetal

growth as a consequence of reduced krill availability. I test these scenarios using historical

whaling records and temperature data to model *B. m. intermedia* foetus size in response to sea

surface temperature (SST) and *B. m. intermedia* abundance.

Methods

Data

Blue whale data came from International Whaling Statistics (IWS) database and comprised

measurements of female body length and foetus length, whether the foetus was single or a twin,

along with the date of capture (www.iwc.int). The complete dataset included 20,144 complete

records spanning 30 years from 1925-1954 with data grouped by Antarctic whaling season,

spanning the austral summer (October - March). Data for the whaling seasons 1940 - 1946 were

missing due to limited hunting during WWII and its immediate aftermath. SST anomaly data

for the Southern Ocean came from the HadISST1 data set, which comprises a globally complete

SST dataset from 1871 onwards at a spatial resolution of 1° latitude by 1° longitude, compiled

by the UK Met Office Hadley Centre (www.metoffice.gov.uk/hadobs/hadisst) (details in

Rayner et al., 2003). A nominal day of conception for *B. m. intermedia* of 1st August was used

to estimate the day of gestation on which a female was killed and its foetus collected. This date

was based on an analysis by Roston, Lickorish, & Buchholtz (2013), which used previous

published studies combined with IWS data and assuming a gestation period of 350 days. While

the evidence indicates this date represents a peak in conception frequency, there is a potentially

wide temporal window in the timing of conception in this species. While time-series data for

E. superba abundance in the Southern Ocean are available (Atkinson et al., 2017) and could

potentially be used to link food abundance with foetus size, for the period of this study these

data are incomplete and were not collected using consistent methodologies.

Density-dependent effects on foetus size were modelled using the total cumulative catch of *B.*

m. intermedia in the whaling season preceding the one in which a foetus was collected. *B. m.*

intermedia catch data came from Branch et al. (2008) and included a correction for Soviet

misreporting of catches. Catch data, rather than estimated population size, was used as an

inverse index of abundance since catch data were relatively reliably recorded, while estimates

of population size necessarily include a margin of error (Christensen, 2006).

Data analysis

Before applying statistical models, a data exploration was undertaken following the protocol

described in Ieno & Zuur (2015). The data were examined for outliers in the response and

explanatory variables, homogeneity and zero inflation in the response variable and the nature

of relationships between the response and explanatory variables. Collinearity between

explanatory variables was examined using co-plots and by estimating variance inflation (VIF).

Model covariates were not collinear and VIF of covariates in the model were all <2. Model

assumptions were verified by plotting model residuals against fitted values, each covariate in

the model and covariates not included in the model. Model residuals were additionally assessed

for evidence of non-linearity (Zuur & Ieno, 2016). Twin foetuses, of which there were 1,014 in

the complete dataset, were consistently smaller that singletons and were dropped from the

analysis. If twin foetuses were retained in the model the qualitative outcomes were identical.

Data were modelled using R (version 3.6.3; R Development Core Team, 2020) with models

fitted in a Bayesian framework using Integrated Nested Laplace Approximation (R-INLA; Rue

et al., 2017). The advantage of using Bayesian inference was that it provides probability

distributions for model parameters of interest, so that probability statements about the

magnitude of model parameters can be made with confidence. This approach avoids reliance

on hypothesis testing and P-values, which are increasingly recognised as unreliable statistical

tools for any but the simplest models (Burnham & Anderson, 2014; Nuzzo, 2014; Wasserstein

& Lazar, 2016).

To accommodate temporal dependency, foetus size was modelled with a gamma GLMM with

a second order random walk trend fitted for day of gestation. The model was formulated as:

$Foetus_{it} \sim Gamma(\mu_{it}, \phi)$

$E(Foetus_{it}) = \mu_i$ and $var(Foetus_{it}) = \frac{\mu_{it}^2}{\phi}$

$\log(\mu_{it}) = \eta_{it}$

$\eta_{it} = \beta_1 + \beta_2 \times Female_{it} + \beta_3 \times Catch_{it} \times \beta_4 \times SST_{it} + DoG_t$

$DoG_t \sim N(0, \sigma_{DoG}^2)$

Where *Foetus*$_{it}$ is the mean size (μ) of foetus *i* on day of gestation (*DoG*) *t*. The parameter ϕ is

a dispersion parameter (Zuur, Ieno & Saveliev, 2017). The model contained linear effects for

female length (*Female*$_{it}$), cumulative *B. m. intermedia* catch in the whaling season preceding

the one in which the foetus was collected (*Catch*$_{it}$), and mean SST of the Southern Ocean for

the preceding austral winter (April-September) (*SST*$_{it}$). Non-informative priors were put on δ

and τ, with Jeffreys prior ($a_1 = b_1 = 0$) on δ and a Gelman (2006) recommended prior ($a_2 = -$

$0.5, b_2 = 0$) on τ.

Because the window of conceptions for *B. m. intermedia* is potentially wide (Roston et al.,

2013), any consistent environmental effects on the time of conception could undermine

conclusions relating to effects of environmental temperature and foetus size. To accommodate

this possibility, a model was also fitted to the difference between observed foetus size and fitted

estimates of foetus size as a function of mean SST for the preceding austral winter using a

Gaussian GLM, with the prediction that any relationship between foetus size and SST (either

positive or negative) would indicate an association between conception date and antecedent

winter SST.

Results

There was a statistically important negative interaction between cumulative *B. m. intermedia*

catch and SST anomaly on *B. m. intermedia* foetus size (Table 1; Fig. 1). Female *B. m.*

intermedia body size was positively associated with foetus size, after controlling for variance

in day of gestation, SST anomaly and *B. m. intermedia* catch (Table 1). The smoother for the

random effect of day of gestation showed a strong positive temporal trend (Fig. 2).

Model estimates of foetus size over time showed a comparable trend to observed data (Fig. 3),

though with lower variance. There was no association between observed foetus size and fitted

estimates of foetus size as a function of mean SST (Fig. 4) indicating that, while there was

substantial variability around the nominal day of conception of 1st August, there was no

association between conception date and SST (Table 2). The body size of pregnant *B. m.*

intermedia over the period for which data are presented showed no trend, either upward or

downward (Fig. 5).

Discussion

A statistically important negative interaction between SST anomaly in the Southern Ocean and

abundance of adult *B. m. intermedia* on foetus growth was detected (Fig. 1, Table 1). Elevated

winter temperatures in the Southern Ocean are associated with weak *E. superba* recruitment

through limitation of the extent and duration of winter sea ice (Atkinson et al., 2004;

Braithwaite et al., 2015; Brierly et al., 2002). A low abundance of *E. superba*, the main

component of the diet of *B. m. intermedia* (Bannister, 2008), is predicted to have a negative

effect on foetus growth. At the same time, the cumulative catch of *B. m. intermedia* through

whaling was predicted to have a positive effect on foetus growth, through the greater abundance

of *E. superba* as the population declined under hunting pressure (Fraser et al., 1992; Laws,

1977) (Fig. 1). The interaction between SST and *B. m. intermedia* catch on foetus size

demonstrates that a positive effect of reduced blue whale density can be reversed at high SST

(Fig. 1). An implication of this result is that while low Antarctic blue whale population size is

predicted to result in increased foetal size-at-age through positive density dependence,

increased temperatures in the Southern Ocean may offset any positive effects of reduced adult

density through reduced recruitment of *E. superba*.

The fitness consequences of reduced foetus size in *B. m. intermedia* are not known, but

compromised feeding by pregnant females is associated with reproductive failure in other

marine vertebrates. In Adélie penguins (*Pygoscelis adeliae*), foraging and breeding

performance is closely linked to *E. superba* abundance and extent of sea ice (Fraser &

Hofmann, 2003). Similarly, Brierly et al. (1997) showed a direct link between fluctuation in *E.*

superba abundance and the reproductive performance of Gentoo (*Pygoscelis papua*) and

Macaroni penguins (*Eudyptes chrysolophus*). In marine mammals, pregnant females in poor

condition limit investment in foetus growth. In the minke whale (*Balaneoptera acutorostrata*)

investment in foetal growth by females is in direct proportion to body condition, which is a

function of feeding success by pregnant females (Christansen et al., 2014). Near-term foetal

body weight was reported to be lower in years with diminished female body condition and

feeding conditions in Icelandic populations of *B. physalus* (Lockyer, 1990, 2007). Feeding

conditions were also shown to be related to blubber thickness and pregnancy rates in *B.

physalus* (Lockyer, 1986; Williams et al., 2013). In southern right whales (*Eubalena australis*)

calf abundance between 1997-2013 off the coast of southern Brazil correlated positively with

E. superba abundance in the waters around South Georgia nine months earlier (Seyboth et al.,

2016). Calving success of the same population between 1971- 2000 was further shown to be

negatively correlated with SST at South Georgia the previous year (Leaper et al., 2006),

demonstrating a link between SST, *E. superba* abundance and reproductive success. The

impacts of nutritional stress on the development and early survival of offspring of pregnant

females are well described in a range of taxa (Lindström, 1999), including marine mammals in

which poorly provisioned foetuses experience slow growth and, following parturition, suffer

reduced survival rates through an elevated risk of starvation, greater susceptibility to predation

and reduced buoyancy (Boltnev et al., 1998; Lockyer, 2007; McMahon et al., 2000, 2003;

Reeves et al., 2001).

The data presented here span 30 years, from 1925-1954. In the interval following this period,

the extent of Antarctic sea ice and duration of the sea-ice season have declined sharply, in

association with elevated SST in the Southern Ocean (de la Mare, 1997; Jacobs & Cosimo,

1997; Smith & Stammerjohn, 2001), with well-characterised negative effects on *E. superba*

abundance (Atkinson et al., 2004; Brierly et al., 2002). The impact of these trends on the fitness

of *B. m. intermedia*, have hitherto not been demonstrated, though several authors have

considered the likely knock-on effects of declining *E. superba* stocks on baleen whale

populations (Braithwaite et al., 2015; Brierly et al., 1997, 2002). The implications of the present

analysis are that the fitness of *B. m. intermedia* will be compromised by reduced *E. superba*

stocks through a negative effect on foetal growth, undermining any positive density-dependent

effects through a 'krill surplus' at low *B. m. intermedia* abundance. This finding offers a

potential explanation for the failure of *B. m. intermedia* to show recovery to anything

approaching pre-exploitation levels, with reduced food availability in Antarctic waters

predicted to result in under-sized calves with low predicted survival rates. With the prospect of

a 95% decline in *E. superba* stocks in regions of the Southern Ocean over the coming century

in response to an increase of 1-2 °C in SST (Murphy et al., 2007), strong recovery of *B. m.*

intermedia populations, in consequence, appears doubtful.

Because foetuses were not aged, the results presented here could be explained through

environmental effects on the time of conception in *B. m. intermedia*, rather than effects on foetal

growth rates. Thus, if conception dates were later in years with high SST in the Southern Ocean

the same effect would be observed, with smaller foetuses, on average, on a given day of the

year, but unrelated to the nutritional state of the female. Model estimates of foetus size-at-age

closely matched observed data, though with lower variance (Fig. 3). Some of this difference in

variance could be attributed to inaccuracies in the measurement and recording of foetal lengths

and with variation in pregnant female condition and foetal growth rates (Frazer & Huggett,

1973). However, an additional explanation for this disparity in predicted and observed size-at-

age could be a wider window in which conceptions occurred than assumed. Despite this

disparity, there was no evidence for a directional effect of time of conception as a function of

environmental conditions in the data (Fig. 4), indicating that model predictions did not differ

systematically from observed as a function of temperature. There is also no obvious mechanism

by which conception dates might be influenced by SST in the Southern Ocean, since breeding

in *B. m. intermedia* takes place at low latitudes during the preceding winter (Roston et al., 2013),

though it may influence ovulation. In addition, the data showed clear evidence of density

dependence in foetus growth, with a positive relationship between foetus size and cumulative

B. m. intermedia catch, which indicates foetal sensitivity to female feeding conditions as

predicted.

Another potential caveat to this study stems from a temporal shift in whalers taking *B. m.*

intermedia to the capture of the smaller pygmy blue whale (*B. m. brevicauda*). However, this

switch came in the early 1960s, approximately a decade after the period for which data are

presented here (Branch, 2007; Branch et al., 2004). It is notable that pregnant female body size

showed no trend, either upward or downward, over the period for which data are presented (Fig.

5), suggesting that pregnant *B. m. brevicauda* did not make a significant contribution to the

catch, if at all. Similarly, while the impact of commercial fisheries for finfish and *E. superba*

on the trophic structure of the Southern Ocean have been substantial (Hill, Murphy, Reid,

Trathan, & Constable, 2006), the onset of commercial fishing in the Southern Ocean came over

a decade after the end of the period for which data are presented (Kock, 1994).

In conclusion, the analysis presented here provides circumstantial evidence for a negative

relationship between *B. m. intermedia* foetus size and antecedent winter SST in the Southern

Ocean. The influence of winter temperature on foetus size is most likely mediated by the

positive effect of the extent and duration of winter sea ice on *E. superba* abundance. The effect

of winter temperature was shown to interact with *B. m. intermedia* abundance, with a positive

density-dependent effect of a 'krill surplus' at low whale population sizes lost at elevated winter

SST. The effect of predicted SST rises of 1-2 °C in the Southern Ocean (Murphy et al., 2007)

are anticipated to compromise the growth of *B. m. intermedia* foetuses, with implications for

the capacity of the sub-species to recover from overexploitation.

Acknowledgements

I am grateful to Andy Brierley, Phil Hammond and Rowena Spence for comments and to John Smith

for meticulous data compilation.

Data Accessibility

Whaling data came from the IWC (www.iwc.int) and are open access. Temperature anomaly data came

from the HadISST1 data set compiled by the UK Met Office Hadley Centre

(www.metoffice.gov.uk/hadobs/hadisst).

References

Ainley, D., Toniolo, V., Ballard, G., Barton, K., Eastman, J., Karl, B., Focardi, S., Kooyman, G.,
Lyver, P., Olmastroni, S., Stewart, B., Testa, J.W., & Wilson, P. (2006). Managing ecosystem
uncertainty: Critical Habitat and dietary overlap of top-predators in the Ross Sea. CCAMLR
document EMM 06–07, Hobart, Tasmania, Australia.

Ainley, D., Ballard, G., Ackley, S., Blight, L.K., Eastman, J.T., Emslie, S.D., Lescroël, A.,
Olmastroni, S., Townsend, S.E., Tynan, C.T., & Wilson, P. (2007) Paradigm lost, or is top-down
forcing no longer significant in the Antarctic marine ecosystem? *Antarctic Science, 19*, 283-290.

Alonzo, S.H., & Mangel, M. (2001). Survival strategies and growth of krill: avoiding predators in
space and time. *Marine Ecology Progress Series, 209*, 203-217.

Atkinson, A., Hill, S.L., Pakhomov, E., Siegel, V., Anadon, R., Chiba, S., Daly, K.L., Downie, R.,
Fielding, S., Fretwell, P., Gerrish, L., Hosie, G.W., Jessopp, M.J., Kawaguchi, S., Krafft, B.A.,
Loeb, V., Nishikawa, J., Peat, H.J., Reiss, C.S., Ross, R.M., Langdon, B., Schmidt, K.,
Steinberg, D.K., Subramaniam, R.C., Tarling, G.A., & Ward, P. (2017). KRILLBASE: a
circumpolar database of Antarctic krill and salp numerical densities, 1926–2016. *Earth System
Science Data*, *9*, 193-2107.

Atkinson, A., Siegel, V., Pakhomov, E., & Rothery, P. (2004). Long-term decline in krill stock and
increase in salps within the Southern Ocean. *Nature*, *432*, 100-103.

Baker, C.S., & Clapham, P.J. (2004). Modelling the past and future of whales and whaling. *Trends in
Ecology & Evolution, 19*, 365-371.

Bannister, J.L. (2008). Baleen whales (mysticetes). In W.F. Perrin, B. Würsig & J.G.M. Thewissen
(Eds.), *Encyclopedia of marine mammals*, 2nd edn (pp. 80-89). New York, NY: Academic Press.

Best, P.B. (1993). Increase rates in severely depleted stocks of baleen whales. *ICES Journal of Marine
Science, 50*, 169-186.

Boltnev, A.I., York, A.E., & Antonelis, G.A. (1998). Northern fur seal young: interrelationships
among birth size, growth, and survival. *Canadian Journal of Zoology, 76*, 843–854.

Braithwaite, J.E., Meeuwig, J.J., Letessier, T.B., Jenner, K.C.S., & Brierley, A.S. (2015). From sea ice
to blubber: linking whale condition to krill abundance using historical whaling records. *Polar
Biology, 38*, 1195-1202.

Branch, T.A. (2007). Abundance of Antarctic blue whales south of 60 °S from three complete
circumpolar sets of surveys. *Journal of Cetacean Research and Management, 9*, 87–96.

Branch, T.A. (2008). Biological parameters for pygmy blue whales. Paper SC/60/SH6 presented to the

Scientific Committee of the International Whaling Committee, Santiago, Chile. 11 pp.

Branch, T.A., Matsuoka, K., & Miyashita, T. (2004). Evidence for increases in Antarctic blue whales

based on Bayesian modelling. *Marine Mammal Science, 20*, 726-54.

Brierley, A.S., Fernandes, P.G., Brandon, M.A., Armstrong, F., Millard, N.W., McPhail, S.D.,

Stevenson, P., Pebody, M., Perrett, J., Squires, M., & Bone, D.G. (2002). Antarctic krill under sea

ice: elevated abundance in a narrow band just south of ice edge. *Science, 295*, 1890-1892.

Brierley, A.S., Watkins, J.L., & Murray, A.W. (1997). Interannual variability in krill abundance at

South Georgia. *Marine Ecology Progress Series, 150*, 87-98.

Burnham, K.P., & Anderson, D.R. (2014). P values are only an index to evidence: 20th-vs. 21st-

Century statistical science. *Ecology, 95*, 627-630.

Clapham, P.J., Young, S.B., & Brownell, R.L. (1999). Baleen whales: conservation issues and the

status of the most endangered populations. *Mammal Review, 29*, 37-62.

Chivers, S.J. (2008) Cetacean life history. In W.F. Perrin, B. Würsig & J.G.M. Thewissen (Eds.),

Encyclopedia of marine mammals, 2nd edn (pp. 215-220). New York, NY: Academic Press.

Christensen, L.B. (2006). Marine mammal populations: reconstructing historical abundances at the

global scale. *Fisheries Centre Research Reports, 14*, 1-161.

Christiansen, F., Víkingsson, G.A., Rasmussen, M.H., & Lusseau, D. (2014). Female body condition

affects foetal growth in a capital breeding mysticete. *Functional Ecology, 28*, 579-588.

Analysis, 1, 515-533.

Fraser, W.R., Trivelpiece, W.Z., Ainley, D.G., & Trivelpiece, S.G. (1992). Increases in Antarctic

penguin populations: reduced competition with whales or a loss of sea ice due to global warming?

Polar Biology, 11, 525–531

Fraser, W.R., & Hofmann, E.E. (2003). A predator's perspective on causal links between climate

change, physical forcing and ecosystem response. *Marine Ecology Progress Series, 265*, 1-15.

Frazer, J.F.D., & Huggett, A.S.G. (1973). Specific foetal growth rates of cetaceans. *Journal of*

Zoology, 169, 111–126.

Gelman, A. (2006). Prior distributions for variance parameters in hierarchical models. *Bayesian*

Hill, S.L., Murphy, E.J., Reid, K., Trathan, P.N., & Constable, A.J. (2006). Modelling Southern Ocean

ecosystems: krill, the food-web, and the impacts of harvesting. *Biological Reviews, 81*, 581-608.

Hucke-Gaete, R., Bedriñana-Romano, L., Viddi, F.A., Ruiz, J.E., Torres-Florez, J.P., & Zerbini, A.N.

(2018). From Chilean Patagonia to Galapagos, Ecuador: novel insights on blue whale migratory

pathways along the Eastern South Pacific. *PeerJ, 6*, e4695.

Ieno, E.N., & Zuur, A.F. (2015). *A Beginner's guide to data exploration and visualisation with R.*

Newburgh: Highland Statistics Limited.

Jacobs, S.J., & Comiso, J.C. (1997). Climate variability in the Amundsen and Bellingshausen seas.

Journal of Climate, 10, 697–709.

Kock, K.H. (1994). Fishing and conservation in southern waters. *Polar Record, 30*, 3-22.

Laws, R.M. (1977). Seals and whales of the Southern Ocean. *Philosophical Transactions of the Royal*
Society B, 279, 81-96.

Leaper, R., Cooke, J., Trathan, P., Reid, K., Rowntree, V., & Payne, R. (2006). Global climate drives
southern right whale (*Eubalaena australis*) population dynamics. *Biology Letters, 2*, 289-292.

Lindström, J. (1999). Early development and fitness in birds and mammals. *Trends in Ecology &*
Evolution, 14, 343-348.

Lockyer, C. (1986). Body fat condition in northeast Atlantic fin whales, *Balaenoptera physalus*, and
its relationship with reproduction and food resource. *Canadian Journal of Fisheries and Aquatic*
Sciences, 43, 142–147.

Lockyer, C. (1990). The importance of biological parameters in population assessments with special
reference to fin whales from the NE Atlantic. *North Atlantic Studies*, *2*, 22-31.

Lockyer, C. (2007). All creatures great and smaller: a study in cetacean life history energetics. *Journal*
of the Marine Biological Association of the United Kingdom, 87, 1035-1045.

Mackintosh, N.A. (1965). *The stocks of whales*. London: Fishing News (Books) Ltd.

de la Mare, W.K. (1997). Abrupt mid-twentieth-century decline in Antarctic sea-ice extent from
whaling records. *Nature, 389*, 57-60.

McMahon, C., Burton, H.R., & Bester, M.N. (2000). Weaning mass and the future survival of juvenile
southern elephant seals, *Mirounga leonina*, at Macquarie Island. *Antarctic Science, 12*, 149–153.

McMahon, C.R., Burton, H.R., & Bester, M.N. (2003). A demographic comparison of two southern
elephant seal populations. *Journal of Animal Ecology, 72*, 61–74.

Monnahan, C.C., Branch, T.A., & Punt, A.E. (2015). Do ship strikes threaten the recovery of
endangered eastern North Pacific blue whales? *Marine Mammal Science, 31*, 279-297.

Murphy, E.J., Trathan, P.N., Watkins, J.L., Reid, K., Meredith, M.P., Forcada, J., Thorpe, S.E.,
Johnston, N.M., & Rothery, P. (2007). Climatically driven fluctuations in Southern Ocean
ecosystems. *Proceedings of the Royal Society of London B: Biological Sciences, 274*, 3057-3067.

Nicol, S. (2006). Krill, currents, and sea ice: *Euphausia superba* and its changing environment.
Bioscience, 56, 111-120.

Nuzzo, R. (2014). Scientific method: statistical errors. *Nature, 506*, 150-152.

R Development Core Team (2019). *R: A language and environment for statistical computing*. Vienna,
Austria: R Foundation for Statistical Computing.

Rayner, N.A., Parker, D.E., Horton, E.B., Folland, C.K., Alexander, L.V., Rowell, D.P., Kent, E.C., &
Kaplan, A. (2003). Global analyses of sea surface temperature, sea ice, and night marine air
temperature since the late nineteenth century. *Journal of Geophysical Research: Atmospheres,*
108, 4407.

Reeves, R.R., Rolland, R., & Clapham, P.J. (2001). Causes of reproductive failure in North Atlantic
right whales: new avenues and research. Northeast Fisheries Science Center Reference Document
01–16.

Reid, K., & Croxall, J.P. (2001). Environmental response of upper trophic-level predators reveals a
system change in an Antarctic marine ecosystem. *Proceedings of the Royal Society of London B:*
Biological Sciences, 268, 377-384.

Reilly, S.B., Bannister, J.L., Best, P.B., Brown, M., Brownell Jr., R.L., Butterworth, D.S., Clapham,
P.J., Cooke, J., Donovan, G.P., Urbán, J., & Zerbini, A.N. (2008). *Balaenoptera musculus* ssp.
intermedia. The IUCN Red List of Threatened Species 2008, eT41713A98837960.

Roston, R.A., Lickorish, D., & Buchholtz, E.A. (2013). Anatomy and age estimation of an early blue
whale (*Balaenoptera musculus*) foetus. *The Anatomical Record, 296*, 709-722.

Rue, H., Riebler, A., Sørbye, S.H., Illian, J.B., Simpson, D.P., & Lindgren, F.K. (2017). Bayesian
computing with INLA: a review. *Annual Review of Statistics and its Application, 4*, 395-421.

Sears, R., & Perrin, W.F. (2008). Blue whale: *Balaenoptera musculus*. In W.F. Perrin, B. Würsig &
J.G.M. Thewissen (Eds.), *Encyclopedia of marine mammals*, 2nd edn (pp. 120-124). New York,
NY: Academic Press.

Seyboth, E., Groch, K.R., Dalla, R.L., Reid, K., Flores, P.A., & Secchi, E.R. (2016). Southern right
whale (*Eubalaena australis*) reproductive success is influenced by krill (*Euphausia superba*)
density and climate. *Scientific Reports*, *6*, 28205.

Smith, R.C., & Stammerjohn, S.E. (2001). Variations of surface air temperature and sea ice extent in
the Western Antarctic Peninsula (WAP) region. *Annals of Glaciology, 33*, 493-500.

Thomas, P.O., Reeves, R.R., & Brownell, R.L. (2016). Status of the world's baleen whales. *Marine*
Mammal Science, 32, 682-734.

Wasserstein, R.L., & Lazar, N.A. (2016). The ASA's statement on p-values: context, process, and
purpose. *The American Statistician, 70*, 129-133.

Williams, R., Vikingsson, G.A., Gislason, A., Lockyer, C., New, L., Thomas, L., & Hammond, P.S.
(2013). Evidence for density-dependent changes in body condition and pregnancy rate of North
Atlantic fin whales over four decades of varying environmental conditions. *ICES Journal of*
Marine Science, 70, 1273-1280.

Yablokov, A.V. (1994). Validity of whaling data. *Nature, 367*, 108.

Zuur, A.F. and Ieno, E.N., 2016. A protocol for conducting and presenting results of regression-type
analyses. *Methods in Ecology and Evolution, 7*, 636-645.

Zuur, A.F., Ieno, E.N., & Saveliev, A.A. (2017). *Spatial, temporal and spatial-temporal ecological*
data analysis with R-INLA. Newburgh: Highland Statistics Limited.

Figure legends

Figure 1. Posterior mean (solid line) fitted Antarctic blue whale (*Balaenoptera musculus intermedia*) foetus length (m), with 95% credible intervals (shaded area), as a function of the interaction between cumulative *B. m. intermedia* catch and antecedent winter sea surface temperature (SST) anomaly in the Southern Ocean, showing a reversal of the positive relationship between foetus length and catch at low SST (-1.3 °C, -0.8 °C), to a negative relationship at elevated SST (0.3 °C).

Figure 2. Posterior mean (solid line) with 95% credible intervals (shaded area) of the random effects of the random walk gamma GLMM for Antarctic blue whale (*Balaenoptera musculus intermedia*) foetus lengths against day of gestation assuming a nominal day of conception of 1st August.

Figure 3. Plot of observed (red dots) and model fitted values (black dots) of Antarctic blue whale (*Balaenoptera musculus intermedia*) foetus lengths against day of year modelled with a random walk gamma GLMM assuming a nominal day of conception of 1st August.

Figure 4. Difference between posterior fitted estimates of Antarctic blue whale (*Balaenoptera musculus intermedia*) foetus length (m) and observed foetus length as a function of average Southern Ocean temperature anomaly (°C) during the preceding austral winter (April – September). Horizontal black dashed line indicates unity.

Figure 5. Boxplots of pregnant female Antarctic blue whale (*Balaenoptera musculus intermedia*) lengths (m) caught in the 1926 – 1954 whaling seasons. Horizontal red dashed line indicates mean female length. N is the number of females measured in each whaling season.

Cumulative Antarctic blue whale catch

Table 1. Posterior mean estimates of *B. m. intermedia* foetus size modelled using a random walk gamma GLMM fitted using INLA with non-informative priors. CrI is the 95% Bayesian credible interval. Credible intervals that do not contain zero in bold to indicate statistical importance. *Female size* is the length of female *B. m. intermedia* from which a foetus was taken. *SST* is the mean sea surface temperature anomaly of the Southern Ocean for the preceding austral winter (April-September). *Catch* is the cumulative catch of *B. m. intermedia* in the preceding whaling season.

Model parameter	Posterior mean	Lower CrI	Upper CrI
Intercept	**0.831**	**0.815**	**0.848**
Female size	**0.090**	**0.085**	**0.096**
Catch	-0.006	-0.017	0.004
SST	-0.003	-0.025	0.018
Catch x *SST*	**-0.033**	**-0.052**	**-0.014**

Table 2. Posterior mean estimates of the difference between observed foetus size and model estimates of foetus size as a function of SST modelled using a Gaussian GLM fitted using INLA with default priors. CrI is the 95% Bayesian credible interval. If credible intervals do not contain zero it indicates statistical importance. *SST* is the mean sea surface temperature anomaly of the Southern Ocean for the preceding austral winter (April-September).

Model parameter	Posterior mean	Lower CrI	Upper CrI
Intercept	0.002	-0.020	0.024
SST	0.006	-0.037	0.049