Patterns of genetic variation in a prairie wildflower, *Silphium*
integrifolium, suggest a non-prairie origin and untapped variation
available for improved breeding.

Andrew R. Raduski[1], Adam Herman[1], Cloe Pogoda[2],

Kevin M. Dorn[3] David L. Van Tassel[4], Nolan Kane[2], and Yaniv Brandvain[1]

Running head: Population genetics of *Silphium integriolium*
Keywords: population genetics, transcriptome, perennial agriculture, de novo domestication

[1]Department of Plant and Microbial Biology, University of Minnesota Twin Cities, 140 Gortner Laboratory, 1479 Gortner Ave, St. Paul, MN 551108 USA.

[2]Ecology and Evolutionary Biology Department, University of Colorado, Ramaley N122, 1900 Pleasant St, Boulder, CO 80309-0334 USA.

[3]USDA-ARS, 1701 Centre Ave, Fort Collins, CO 80526 USA.

[4]The Land Institute, 2440 E. Water Well Road, Salina, KS 67401 USA.

₉   ABSTRACT

₁₀ *Premise*

₁₁ Understanding the relationship between genetic structure and geography provides
₁₂ information about a species' evolutionary history and can be useful to breeders interested
₁₃ in de novo domestication. *Silphium integrifolium* is an iconic perennial American prairie
₁₄ wildflower that is targeted for domestication as an oilseed crop. Germplasm in the existing
₁₅ breeding program is derived from accessions collected in restricted geographic regions. We
₁₆ present the first application of population genetic data in this species to address the
₁₇ following goals (1) improve existing breeding programs by characterizing genetic structure
₁₈ and (2) identify the species geographic origin and potential targets and drivers of natural
₁₉ selection as the range has expanded.

₂₀ *Methods*

₂₁ We developed a reference transcriptome to use as a genotyping reference for dozens of
₂₂ samples from throughout the species range. Population genetic analyses were used to
₂₃ describe the distribution of genetic variation and demographic inference modeling was used
₂₄ to characterize potential processes that have shaped variation. Divergence outlier scans for
₂₅ selection and associations with environmental variables were used to identify loci linked to
₂₆ putative targets and drivers of natural selection.

₂₇ *Key results*

₂₈ Genetic variation partitions samples into three geographic clusters. Patterns of variation
₂₉ and demographic modeling suggests that the geographic origin of the species is in the
₃₀ American southeast. Breeding program accessions are from the region with lowest observed
₃₁ genetic variation.

₃₂ *Conclusions*

₃₃ This iconic prairie species did not originate within the modern prairie. Breeding programs
₃₄ can be improved by including accessions from outside of the germplasm founding region,
₃₅ which has relatively little genetic variation compared to the south and east. The observed
₃₆ geographic structuring of variation coupled with the identified targets and environmental
₃₇ drivers of adaptation during range expansion can guide future collecting efforts towards
₃₈ wild populations with beneficial agronomic traits.

Species of the genus *Silphium* (Asteraceae) are deep-rooted, perennial wildflowers found throughout the United States, east of the Rocky Mountains. Common *Silphium* species are best known for being charismatic flora of the North American prairie and their ability to tolerate extreme abiotic stresses (Weaver et al. 1935; Leopold 1968). Recently there have been efforts to better incorporate the ecological benefits of *Silphium* species into improving habitat restoration and agricultural practices (Van Tassel et al. 2017). Roots of perennial plants in the soil protect against soil erosion, promote a diverse soil microbiome, and act as a carbon sequestration resource. Because of these benefits, combined with high biomass accumulation requiring relatively few inputs, species of *Silphium* show promise for forage and biogas production (Ustak and Munoz 2018; Wever et al. 2019) or as a perennial oil seed crop (Van Tassel et al. 2017). Additionally, the incorporation of native wildflowers in prairie restoration projects or as buffer strips among agriculture fields are thought to improve pollinator abundance (Isaacs et al. 2009). *Silphium* species are known to attract beneficial insects (Fiedler and Landis 2007) and are currently being evaluated for such conservation practices because they are large plants, competitive with weeds after established, have potential for economic return (seed or bioenergy) and breeding programs may improve seedling vigor compared to other wildflowers (Butters et al. 2019). Despite increased interests across applied settings, relatively little is known about patterns in the distribution of genetic variation in long-lived perennial prairie species such as *Silphium*.

Knowledge of the origin of this iconic prairie species is important given the relatively recent assemblage of the current North American prairie ecosystem, within the past few million years (Fox and Koch 2003, 2004). The North American prairie habitat has been well-studied as a grassland ecosystem (Weaver 1954; Lauenroth 1979; Matthews 1988; Wilson and Hartnett 1998; Samson et al. 2004; Knapp et al. 2008). However, surprisingly little is known about the evolutionary history of common North American prairie forbs from population genetic data, and it is not currently known if there are general patterns across species of directional range expansion or geographic sources of ancestral variation. Changes in species composition have drastically altered the biome as C4 grasses became dominant on the landscape over the past few million years (Fox and Koch 2003, 2004). Data from fossil pollen deposits have suggested that now common prairie wildflower families, such as the Asteraceae, were likely not plentiful on the prairie landscape at the time that C4 grasses became dominant (Williams et al. 2004). Understanding the evolutionary and demographic history of how a species has come to inhabit its current range can allow us to better understand the present relationship between a species's genetic variation and geography.

As there has been a recent movement to domesticate *S. integrifolium* and *S. perfoliatum*, understanding the underlying extent and distribution of genomic variation within and among populations will support ongoing applied *Silphium* research. Breeding trials to develop *S. perfoliatum* as a potential cover crop began in the 1950's and continue today with additional effort to develop it as a biogas resource (Stanford 1990; Wever et al. 2019). Most recently, *S. integrifolium* has been targeted for development as an oilseed

crop, in part, because of the similar fatty acid composition profile in its seeds to sunflower seeds (Reinert et al. 2019). Accessions used in *S. perfoliatum* breeding have little genetic diversity from (often) unknown geographic origins (Wever et al. 2019). To date, most *S. integrifolium* breeding progress has been made from accessions that originated within a small geographic area (Kansas, USA) (Vilela et al. 2018), with a general lack of genomic resources (Van Tassel et al. 2017). The extent to which breeding populations are representative of species-wide diversity has not been genetically evaluated. However, if populations that produced *S. integrifolium* breeding accessions are not representative of species-wide diversity, greater improvements may be achieved by incorporating accessions with unique diversity. Additionally, the choice of locations for field trials may be informed by a better understanding of the relationship between natural genetic variation and geography.

Here, we characterize genetic variation among populations of *S. integrifolium* with the goal of informing and improving existing breeding programs, as well as understanding the history and evolution of the species. Presently, *Silphium* genomic resources are lacking, in part, because of the difficulty of assembling its large, repetitive genome. All *Silphium* species are diploid (2n = 14) with genome sizes that are nearly triple that of their close relative sunflower (*Helianthus annuus*) (Bai et al. 2012). We utilize transcriptome resequencing of *S. integrifolium* to characterize and quantify functional genetic variation throughout the species range. We assemble a reference transcriptome of a diverse set of tissues from a single individual and use the reference to identify variants from a panel of resequenced young leaf transcriptomes.

The goals of our research are twofold. First, we aim to investigate the distribution of neutral genetic variation within and among wild populations while maintaining an interest in identifying a possible region of geographic origin of the species. Additionally, we hypothesize that populations of this widespread species are locally adapted to their unique environments and thus, quantifying the relationship between genetic variation and geography is important. Therefore, we next sought to identify genes whose patterns of genetic variation have been shaped by natural selection. Such environmentally associated variants highlight traits experiencing geographically variable selection in this species, and could identify important variation to include in the breeding program germplasm.

Methods

De novo transcriptome sampling, sequencing, and assembly

Five tissues were collected from a single *S. integrifolium* plant from the breeding program of The Land Institute in Salina, Kansas. Root, crown, flower, leaf, and stem tissues were collected and immediately frozen in liquid nitrogen. Flash frozen samples were transported to the University of Minnesota on dry ice where total RNA was extracted using Qiagen RNeasy kits (Germantown, Maryland USA). Pooled RNAseq stranded libraries were constructed by the University of Minnesota Genome Center and sequenced on two lanes of an Illumina (San Diego, California USA) HiSeq instrument (2 x 125 paired-end reads per

121 lane).

122   Raw sequence reads were trimmed of low quality sites and adaptor sequences using
123 BBDuk (Bushnell 2018) and *de novo* assembled into a reference transcriptome with Trinity
124 (Haas et al. 2013) (version 2.1.1, non default settings: SS_lib_type_type RF
125 min_contig_length 300 min_kmer_cov 2 min_glue 5).

126   *Assembled transcriptome completeness and annotation*

The completeness of the initial Trinity *S. integrifolium* assembly was measured by assessing
the number of benchmark universal single-copy orthologs with the program BUSCO v3.0.2
with the *embryophyta_odb9* gene set (Simão et al. 2015). The longest isoform of each
transcript was used for BUSCO searches.

The number of nearly full length protein coding transcripts was counted by querying
the Swiss UniProt protein database (Boutet et al. 2016) with the full set of transcripts
using BLASTx with an e-value cutoff of 1e-20 (Madden 2013). For each unique result from
the protein database, the percent of matching sequence length of the best matching
transcript was recorded. Transcripts that matched a database entry over more than 80% of
its length were classified as nearly full-length transcripts.

Transcript abundance was estimated with the RSEM software (Li and Dewey 2011)
after mapping cleaned sequence reads to the Trinity assembly using Bowtie v1.1.2
(Langmead and Salzberg 2012). Gene abundance matrices were built after calculating the
number of gene transcripts per million sampled transcripts (TPM) using Trinity scripts.
Transcript abundance matrices per gene were built to determine the isoform prevalence
within each gene. The median contig length (N50) of the transcriptome assembly was
calculated using all transcripts (E100N50) and only transcripts with expression levels
above the 90th percentile (E90N50). Although the N50 statistic is widely reported without
regard for expression level, it may be an inaccurate representation of transcriptome
assembly because of a bias from low expressed transcripts.

Functional annotation of the longest isoform of each contig was performed using the
standard Trinotate pipeline (Bryant et al. 2017). Possible protein coding sequences were
identified using TransDecoder (Haas and Papanicolaou 2017). Putative proteins longer
than 100 amino acid residues were kept. BLASTx searches using transcripts and BLASTp
searches using predicted amino acids as queries were performed against the Uniprot protein
sequence database (Madden 2013; Boutet et al. 2016).

153   *Reference transcriptome filtering*

The Trinity assembly was further filtered to create a subset of contigs to serve as a
genotyping reference for downstream population genetic analyses. We initially retained
only contigs that had a significant BLAST result to a plant after using the longest isoform
of each contig as a query to the Swiss UniProt database and the NCBI non-redundant
nucleotide database with BLASTp and BLASTn searches, respectively. E-value cutoffs of
1e-20 (BLASTp) or 1e-30 (BLASTn) were used. Plant contigs were identified by keeping

queries with a best BLAST result belonging to a genera on the NCBI viridiplantae
taxonomy list (Sayers et al. 2009). The completeness of the filtered contig set was assessed
by again performing a BUSCO analysis of TransDecoder predicted amino acids with the
embryophyta_odb9 gene set. The plant filtered contig set was further reduced to include
only contigs that were inferred to be single copy orthologous genes in both *S. integrifolium*
and sunflower, *H. annuus* with OrthoFinder (Emms and Kelly 2015). The resulting set of
contigs served as a genotyping reference for downstream resequencing of samples.

Transcriptome resequencing

Fresh young leaf tissue was collected and flash frozen from 68 wild *S. integrifolium* plants
and one *S. lacinatum* plant to be used as an outgroup (Appendix S1; see Supplemental
Data with this article). All plants were grown from wild collected seed in a common garden
at The Land Institute in Salina, Kansas. Pooled RNAseq stranded libraries were
constructed by the University of Minnesota Genome Center and sequenced using an
Illumina HiSeq instrument (2 x 125 PE reads per lane).

Variant calling workflow

Reads from each resequenced leaf transcriptome were trimmed of adaptor sequences using
scythe v0.991 (`https://github.com/vsbuffalo/scythe`) and aligned to the genotyping
reference using STAR v2.5.2b (Dobin et al. 2013). Reads were aligned in a two-pass mode
to accommodate the possible mapping of isoforms across splice junctions. PCR duplicates
were removed from each alignment using Picard Tools
(`http://broadinstitute.github.io/picard/`). Sample genotypes at variant single
nucleotide polymorphisms (SNPs) and non-variant sites were called using Freebayes with all
samples assigned to a single population (Garrison and Marth 2012). A minimum site depth
of 10 reads was required within each sample for a site to be included in genotype calling.

Population structure and genetic diversity analyses

A principal components analysis (PCA) was performed on *S. integrifolium* genotypes to
assess genetic population structure across the species range. SNPs were filtered to remove
rare variants (minor allele frequency (MAF) $\leq$ 0.05) and sites in linkage disequilibrium
(LD) (LD threshold = 0.2, missingness = 0%) prior to performing PCA. We performed a
Procrustes transformation of sample principal components (PC1 and PC2) coordinates
onto their sampling latitude and longitude coordinates to better visualize the relationship
between genetic variation structuring and geographic space (Wang et al. 2010).
    The relative effect of isolation-by-distance on the observed population structure
patterns was estimated using `conStruct` (Bradburd et al. 2018). Similar to other
STRUCTURE-like programs (Pritchard et al. 2000), `conStruct` assigns a proportion of
ancestry from different genetic groups (K) to individuals. Additionally, `conStruct` can
incorporate a spatial element into its model to account for isolation-by-distance. We ran

₁₉₇ spatial and non-spatial models over a range of underlying genetic groups (K = 1 through K
₁₉₈ = 6), and found spatial models always had higher predictive accuracy. We ran multiple
₁₉₉ spatial models (K = 1 through K = 6,) to identify the best K value with a scree-plot of
₂₀₀ predictive accuracy. For each value of K, we ran four independent runs with a 1000
₂₀₁ iteration chain length. Five independent runs, each with four MCMC chains of 5000
₂₀₂ iterations were then performed with the inferred optimal K value.

₂₀₃ Genetic relationships among *S. integrifolium* samples were inferred using `SVD`
₂₀₄ `Quartets` (Chifman and Kubatko 2014). `SVD Quartets` estimates an unrooted phylogeny
₂₀₅ describing the relationships between sets of four taxa. The correct phylogeny topology
₂₀₆ under a coalescent process is inferred by singular value decomposition of the genetic
₂₀₇ distance matrix containing the four taxa. Multiple quartet phylogenies are then assembled
₂₀₈ into a single phylogeny that contains all taxa. A matrix of biallelic SNPs was used with
₂₀₉ exhaustive quartet sampling, 100 bootstrap replicates and the Quartet MaxCut tree
₂₁₀ assembly method to produce an unrooted phylogeny within `Paup` (Swofford 2000). A single
₂₁₁ individual from Illinois was removed from the SNP data matrix prior to running `SVD`
₂₁₂ `Quartets` because we observed evidence that it was of admixed ancestry.

₂₁₃ Results from PCA, `conStruct`, and `SVD Quartets` indicated that the major axes of
₂₁₄ genetic variation separate *S. integrifolium* samples into three clusters that broadly
₂₁₅ correspond to geographic regions. Hereafter we refer to these broad groupings as the
₂₁₆ eastern, southern, and western regions. We use 'population' to refer to the American state
₂₁₇ within each region.

₂₁₈ The level of support for region-level relationships among groups within *S. integrifolium*
₂₁₉ across markers was assessed using `TWISST` (Martin and Van Belleghem 2017). `TWISST`
₂₂₀ prunes a given phylogeny down to one tip from each user-defined taxon group and reports
₂₂₁ the topology of the pruned tree. The process is repeated and support for each topology is
₂₂₂ recorded as the number of observations. A maximum likelihood phylogeny was inferred for
₂₂₃ each contig using RAxML v8.0 (Stamatakis 2014) and each tip was assigned to a regional
₂₂₄ taxon inferred from the above clustering methods and population genetic analyses. The
₂₂₅ pruned topology most often observed for each contig's reconstructed phylogeny was
₂₂₆ recorded.

₂₂₇ Genetic differentiation among *S. integrifolium* populations was measured by
₂₂₈ calculating Wright's F_{ST} (Weir and Cockerham 1984) between all pairwise combinations of
₂₂₉ populations using a stringently filtered marker dataset with the R package `StAMPP`
₂₃₀ (Pembleton et al. 2013). Retained SNP markers were biallelic in *S. integrifolium*, had a
₂₃₁ Phred quality score $\geq$ 20, had an allele balance within heterozygotes between 0.25 and 0.75
₂₃₂ or within homozygotes less than 0.05, had average sequencing depth greater than 10 and
₂₃₃ less than 200, and MAF $\geq$ 0.05. One hundred bootstrap replicates were performed across
₂₃₄ loci to calculate 95% confidence intervals around F_{ST} estimates.

₂₃₅ Average numbers of nucleotide differences per base pair at four-fold degenerate sites
₂₃₆ were calculated between all samples to measure genetic diversity (Nei 1987). We organized
₂₃₇ the diversity measurements into within (π) and between (D_{xy}) populations. Variation was
₂₃₈ calculated across contig bootstrap replicates to calculate 95% confidence intervals around

mean values. Tajima's D was calculated within each population on every contig using vcftools (Danecek et al. 2011). The level of partitioning of genetic variance was measured using an AMOVA performed on genetic distances (Nei 1972) between individuals with populations and regions as hierarchical levels of stratification with the R packages StAMPP and pegas (Pembleton et al. 2013; Paradis 2010). Statistical significance of variance partitioning among stratification levels was assessed with 10000 random permutations.

We found that a single population, Illinois, exhibited patterns of nucleotide variation, Tajima's D, and inferred ancestry that were unexpected within the larger contextual patterns of each metric across the species landscape. We performed the three population f_3 admixture tests in TreeMix to test the hypothesis that the Illinois population is the result of past admixture between Wisconsin and a third population instead of the null expectation that the relationships among the sampled populations are the product of a tree-like branching process (Reich et al. 2009; Pickrell and Pritchard 2012). Each test took on the form $f_3\{IL; WI, X\}$, where evidence that IL is the result of admixture between WI and X populations is given by a significantly negative test statistic. For all identities of X, Z-scores and standard errors of estimates were reported from jackknifing 50 SNP blocks. We subsequently removed these recently and heavily admixed samples from our demographic inference so as to not confuse historical processes with anthropogenic movement.

Data from the three-dimensional joint site frequency spectra was used to infer general branching patterns of *S. integrifolium* regional taxa using Moments (Jouganous et al. 2017). For three-dimensional spectra, the tested branching scenarios were those that would result in the tree topologies (East, (South, West)), (South, (East, West)), and (West, (South, East)). For each branching pattern scenario, we fit demographic models with no migration, symmetric migration between each taxa pairs, and asymmetric migration rates. The fit of each model was assessed across 100 replicates using different starting parameters selected from bounded limits decided by trial and error. The likelihood of each model was optimized using Powell's method and the quality of each model's fit to the data was assessed with AIC scores. After the best-fitting model was selected for each scenario, uncertainty estimates around each parameter were calculated using the Godambe Information Matrix with 100 bootstrapped spectra (Jouganous et al. 2017).

Scans for selection

To identify loci possibly under divergent selection, we used BayeScan (Foll and Gaggiotti 2008; Foll 2012) to find statistical outlier SNPs that significantly deviate from neutral expectations. We calculated global F_{ST} scans across sites between regions using the same marker set used to perform an AMOVA. Default BayeScan model parameters were used (burn-in length = 50,000, thinning interval = 10, sample size = 5,000, number of pilot runs = 20, pilot run length = 5,000). Annotation of resulting outlier loci was done using BLASTx, BLASTp and eggNOG searches using Trinotate scripts. We then sought to identify environmental variables that may be associated with the selective pressures acting on each outlier protein. We identified environmental variables that had significant associations with genetic markers within each protein after accounting for the coancestry of

²⁸⁰ samples using BayEnv2 (Günther and Coop 2013). Environmental variables from the
²⁸¹ geographic midpoint of each population were extracted from the ClimateNA dataset (Wang
²⁸² et al. 2016). Brief descriptions of the climatic variables used are provided in Appendix S3.
²⁸³ For each BayeScan identified protein, the Bayes factor, absolute Spearman's rank, and
²⁸⁴ absolute Pearson correlation coefficients with the largest quantile were used to identify
²⁸⁵ putative environmental variables driving local adaptation.

Results

De novo transcriptome and assembly

²⁸⁸ Trinity assembled 541,120,911 paired-end sequencing reads into 280,573 transcripts and
²⁸⁹ 166,458 contigs, totaling over 246.9 megabases of assembled nucleotides. The median
²⁹⁰ contig length of all transcripts (E100N50) was 1,159bp. The median contig length of
²⁹¹ transcripts within the top 90th percentile of expression (E90N50) was 2,014bp. The filtered
²⁹² assembly with the longest isoform of each contig totaled 123.5 megabases. A BUSCO
²⁹³ analysis of the longest isoform of each assembled contig found that 80.7% of the 1,440
²⁹⁴ genes in the *embryophyta_odb9* dataset were found in complete form (71.3% as single copy
²⁹⁵ and 9.4% as duplicated). BLASTx searches of the Swiss UniProt database identified 13,143
²⁹⁶ unique protein results. Of those, 8,418 (64.0%) were found as full-length or nearly
²⁹⁷ full-length transcripts in the full Trinity transcriptome assembly.

Reference transcriptome filtering

²⁹⁹ 51,238 contigs remained after filtering the full Trinity for contigs with a BLASTp or
³⁰⁰ BLASTn result from genera on the NCBI viridiplantae taxonomy list. Results of a BUSCO
³⁰¹ analysis of this filtered contig set were similar to the full assembly analyses (80.7% of 1,440
³⁰² BUSCO genes found complete, of which, 71.4% were single copy and 9.3% were
³⁰³ duplicated). The plant-filtered contig set was further reduced to 10,575 contigs that
³⁰⁴ contained orthologous genes that were inferred to be single copy in *Silphium* and sunflower,
³⁰⁵ *H. annuus.* This final contig set was used as a reference to align resequencing reads and to
³⁰⁶ call variants.

Transcriptome resequencing and variant calling

³⁰⁸ Leaf transcriptome resequencing data mapped to 7,718 of the 10,575 contigs in the
³⁰⁹ genotyping reference set. Genotyping was restricted to contigs with mapped reads and
³¹⁰ covered 12,410,795 total sites (852,148 SNPs) across all samples. One *S. integrifolium*
³¹¹ sample from Missouri was removed from downstream analyses because of low average
³¹² sequencing depth (< 10) and a high rate of marker missingness ($> 60\%$).

The PCA of *S. integrifolium* genotypes resulted in the first two PC axes explaining 13.79%
of the variation in the genetic data. Genotypes formed three distinct groups on the first
two PC axes (Fig. 1). The groups broadly consisted of plants from the eastern (Arkansas,
Illinois, Mississippi, Missouri, and Wisconsin), southern (Oklahoma and Texas), and
western (Kansas and Nebraska) regions of the species range. Eastern and western regions
were separated along the first PC axis. The southern region was separated from eastern
and western regions along the second PC axis.

Our `conStruct` analyses that incorporated isolation-by-distance into population
structure (spatial models) always performed better than non-spatially aware models.
Results from spatially aware runs with different numbers of K indicated that relatively
little predictive power was gained after increasing the number of genetic layers past K = 3.
The patterns of assigned ancestry across all samples supported the PCA pattern of
subdivision. Within the eastern region, samples were overwhelmingly comprised of ancestry
from a single dominant genetic layer. Samples from the southern and western regions are
inferred to be admixed, with both regions being made up of a different dominant ancestry
layer (Figs. 2A, 2B). In western samples, the second largest ancestry layer is the dominant
layer in southern samples. The reciprocal is true in most southern samples.

Reconstruction of the phylogenetic relationships of samples using `SVD Quartets`
resulted in strong geographic clustering with 100% bootstrap support for all nodes in the
phylogeny (Fig. 2C). All but one sample from the eastern region formed a large
monophyletic clade, with most samples from within a population forming smaller
monophyletic groups. Mississippi samples exhibit a polyphyletic pattern throughout the
larger eastern clade, with various samples being found clustered with Missouri or Arkansas
samples. A single Mississippi sample was reconstructed to be outside of a larger clade
containing all other *S. integrifolium* samples. Southern samples are paraphyletic with most
Oklahoma samples forming a monophyletic clade that is sister to a single Texas sample.
Most Texas samples form a monophyletic clade that is sister to a larger clade that contains
a single Oklahoma sample and a larger monophyletic clade of all western samples (Fig. 2).

To investigate the level of support for the general `SVD Quartets` phylogeny across
markers, we performed a `TWISST` analysis. Each maximum likelihood phylogeny was rooted
with *S. laciniatum* as the outgroup and the region-level topology of *S. integrifolium* taxa
phylogeny was recorded. For each taxa in a tree the identity of its sister tip was recorded.
If the nearest sister was a node, a descendant of the sister node was randomly sampled and
given a weight proportionate to the number of descendants of the sister node. Similar to
the `SVD Quartets` phylogeny topology, southern and western regions were sister to each
more often than either was sister to the eastern region. The eastern region was found to be
sister to western and southern regions an approximate equal number of times across
markers (Fig. 2D).

Pairwise F_{ST} ranged from 0.0340 to 0.2204 across all population (i.e. state) pairs.
Differentiation patterns among populations supported the structure inferred from PCA. As
a general pattern, differentiation within regions was approximately $F_{ST} \leq 0.10$ regardless of

geographic distance separating populations (Fig. 3). Population differentiation between regions was generally higher than within region comparisons with increased differentiation observed between geographically distant populations (Fig. 3). The three highest within-region comparisons all involved the suspect IL population, likely reflecting the recent admixture in this sample. The highest levels of differentiation were observed between eastern and western populations, and between eastern and southern populations. The greatest allele frequency divergence observed was between Nebraska (western region) and Illinois (eastern region) populations (F_{ST} = 0.2204). Our AMOVA results showed that genetic variation is strongly partitioned both among regions and among populations within regions (Table 1).

The average numbers of pairwise nucleotide differences per base pair at four-fold degenerate sites (π) between samples within regions were calculated to estimate genetic diversity (Fig. 4B, Appendix S2). The eastern region had the highest mean genetic diversity of any group (π = 0.01013, 95% C.I. 0.00986 - 0.01041). The southern region had slightly lower genetic diversity (π = 0.00982, 95% C.I. 0.00952 - 0.01013). The western region had lowest genetic diversity (π = 0.00894, 95% C.I. 0.00865 - 0.00923).

Within the eastern and southern regions, populations with lower latitudes had higher π, consistent with those regions possibly serving as founder populations for northward expansion (Fig. 4A). Mississippi and Arkansas contained more genetic diversity than most more eastern region populations (π = 0.01018, 95% C.I. 0.00987 - 0.01051 and π = 0.00972, 95% C.I. 0.00935 - 0.01008, respectively). Within the southern region, genetic diversity in Texas was greater than Oklahoma (π = 0.0098, 95% C.I. 0.00944 - 0.01017 and π = 0.00949, 95% C.I. 0.00918 - 0.0098, respectively). The reduction in genetic diversity found in Kansas and Nebraska populations (π = 0.00846, 95% C.I. 0.00815 - 0.00878 and π 0.00948, 95% 0.00918 - 0.00978, respectively)) is suggestive of a past bottleneck of *S. integrifolium* into the western region.

Patterns of nucleotide divergence between populations (D_{xy}) support the regional structure grouping of plants found through PCA and `conStruct`. The largest D_{xy} observed, on a regional scale, was between eastern and southern regions (D_{xy} = 0.0105, 95% C.I. 0.01021 - 0.0108), only slightly exceeding π in the diverse eastern region. The largest levels of divergence were found between Mississippi and Texas (D_{xy} = 0.01069, 95% C.I. 0.01036 - 0.01102) followed by Texas and Wisconsin (D_{xy} = 0.01068, 95% C.I. 0.01035 - 0.01102), and Mississippi and Nebraska (D_{xy} = 0.01067, 95% C.I. 0.01036 - 0.01098). Plants from the western region had the smallest divergence from southern region plants, suggesting that the western populations are a result of a population bottleneck originating from southern populations (Figs. 3, 4C).

Tajima's D, calculated within each population across contigs, was consistently negative in eight of nine populations (Fig. 6), suggesting population growth across the range. We note that this runs counter to the known recent contraction of prairie species due to habitat loss, and likely reflects population growth since the glacial retreat. The western region populations (Kansas and Nebraska), Oklahoma, and Mississippi populations had the most negative Tajima's D values. Illinois was the only population that had an average

Tajima's D value greater than zero (D = 0.201), likely reflecting the excess of intermediate frequency variants induced by rampant recent admixture.

The relatively high genetic variation (π), elevated Tajima's D estimates, and amounts of mixed ancestry inferred by `construct` within the IL population led us to test for a history of admixture using the f_3 test for treeness (Reich et al. 2009). We assumed that the IL and WI populations have recently diverged from a common ancestor population and thus tested the null hypothesis that a tree-like branching pattern would best describe the process that formed any trio of populations containing IL and WI in the form $f_3\{$IL; WI, X$\}$. We found significantly negative Z-scores for the f_3 test when the third population in the trio was any western or southern population (KS, NE, OK, or TX), but not any other eastern population (AR, MO, or MS) (Table 2). Overall, this pattern suggests that IL is the product of admixture between the general eastern region lineage and the western or southern regions lineages or the common ancestor of the two.

Data from the joint site frequency spectra was used to model the demographic history of *S. integrifolium* under three general scenarios. The scenarios differed from each other by the identities of the two regional taxa assumed to be sister to each other *e.g.* (East,(West,South)), (South,(East,West)), and (West,(East,West)). Under each model, an ancestral population was forced to split into two lineages, with one lineage later experiencing a second split to produce three total taxa. The proportions of a lineage that went into a split at a branching event, timing of branching events, migration rates between all lineages, and population sizes were estimated for each model. Migration rates differed in each model to either be nonexistent, symmetric, or asymmetric.

Under all branching scenarios considered, models that included migration were preferred over no-migration models. Models with migration rates constrained to be symmetric were preferred over asymmetric migration models. The three different branching scenarios considered for three-dimensional spectra each resulted in similar maximum likelihoods and AIC scores (Fig. 5, Table 3). Overall, the best fitting model produced the (East,(South,West)) topology with substantial amounts of population size changes in the South and West with migration throughout all populations. This model estimates that approximately 40% of the ancestral population split $\approx 3.7 N_e$ generations ago to form the common ancestor of South and West regions. This lineage then approximately doubled in size before splitting again $\approx 1.6 N_e$ generations ago to create the South and West regions. At the most recent branching event, the majority of the population (83.7%) split to form the South region, which has since decreased by nearly half. The West region's population size has increased nearly fourfold since the most recent split. The population of the East region has been stable in size since its initial split from the common ancetor. Estimated migration rates are highest between East-South and South-West regions, however confidence intervals around migration rates are wide. Parameter estimates under all models are reported in Table 3.

 Scans for selection

We performed a transcriptome wide scan for divergence outliers on a set of 41,159 high
quality markers with BayeScan. We detected 47 SNPs that were F_{ST} outliers while using a
maximum false discovery rate of 0.001. The outlier SNPs were spread across 40 contigs, 34
of which we were able to gather annotation information from BLAST and eggNOG
database searches. The summarized annotation information for selected proteins are given
in Appendices S4, S5, and S6. For each protein identified by BayeScan as containing an
outlier SNP, we found the environmental variable that had the highest correlation with any
SNP based on Bayes factor, Spearman rank correlations, or Pearson correlation coefficient
quantiles. We speculated that the combined knowledge of the protein function from
BLAST and eggNOG searches and associated environmental variables would lead to better
understanding past selective pressures *S. integrifolium* experienced while expanding to its
current geographic range. Several genes with known functions associated with abiotic
stress tolerance show associations with climatic variables. Below, we highlight
environmental variables that are associated with genes to further understand biological
processes that may have enabled range expansion. For example, Delta(8)-fatty-acid
desaturase has a role in tolerating cold and chilling environments (Chen et al. 2012) and
was found to be associated with both continentality (temperature difference between
warmest and coldest months) and mean annual precipitation variables. Aquaporin TP1
encodes a channel protein involved in water transport that has been shown to influence cell
turgor and expansion (Maurel et al. 1993) and was found to be associated with annual heat
moisture index, a composite measure of heat and precipitation, and longitude. The E3
ubiquitin ligase SUD1 gene is involved in drought tolerance through its effects on cuticle
wax formation and transpiration rate (Lü et al. 2012) and was found to be associated with
the mean warmest month temperature and extreme warm temperatures.

461   DISCUSSION

462   *Reference transcriptome as a resource*

The development of *Silphium* genomic resources has been hindered because of difficulties
imposed by its large genome that is replete with repetitive elements. We sought
transcriptome resequencing as a reliable method to obtain a common set of reduced
representation genetic markers from this non-model organism. Towards this goal, we built
a reference transcriptome of a diverse set of tissues from a single plant and used it as a
genotyping reference for the resequencing of a species wide panel of *S. integrifolium*
samples. The initial transcriptome assembly contained roughly 250Mb of sequence data
assembled into more than 166,000 contigs, some of which are products of sequence
contamination and assembly error. An assembly of the longest isoform of each contig
contained a majority of nearly full-length transcripts and an almost complete set of
universal benchmark genes predicted to be present in all plants. Our final genotyping
reference was the product of filtering to include only contigs that contained genes that have

been sequenced in other plants and present as single copies in both *Silphium* and its close
relative, sunflower. Assuming approximate conservation of the gene number present across
plants, the reference transcriptome contains roughly one third of genes in *Silphium* and can
presumably serve as a helpful genotyping resource in place of a quality reference genome
(Sterck et al. 2007).

S. integrifolium comprises three genetic groups

We resequenced the young leaf transcriptomes from dozens of *S. integrifolium* plants from
localities that cover the species range. Multiple analyses to identify genetic population
structure across samples found that *S. integrifolium* comprises three clusters (Figs. 1, 2).
Although the genetic clusters are each found within particular geographic regions of the
United States, there are no obvious physical barriers that may help maintain cluster
integrity by restricting gene flow. Population pairs that are geographically near ($\approx$200 km)
but belong to different regional groupings show greater differentiation, as measured by F_{ST},
than the most distant ($\sim$1000 km) population pairs from the same regional grouping (Fig.
3). Although we found considerable levels of neutral genetic variation within each
population or region, both are lower than levels found in other self-incompatible wild
sunflower species (Fig. 4). Using a set of nuclear genes, Liu and Burke (2006) found
approximately three times the amount of synonymous variation in wild *H. annuus* (wild
populations $\pi = 0.0315$) than what we observe in *S. integrifolium.* The discrepancy
suggests that *S. integrifolium* experienced a much smaller historical effective population
size than sunflower, perhaps reflecting *S. integrifolium*'s longer generation time and/or
other features of its life history (Andreasen and Baldwin 2001).

Plants from the eastern region form the most genetically variable cluster (Fig. 4B).
Within the eastern region there is a general negative correlation between latitude and π
(Fig. 4A). The Illinois population was found to have appreciable amounts of admixed
ancestry (Figs. 2A, 2B). Results from f_3 tests for introgression having a role in the Illinois
population's past were significant (Table 2). Introgression from the west could potentially
have proceeded naturally through unsampled populations, *e.g.* Iowa, or from western *S.
integrifolium* seed introduced by humans in prairie restoration projects. Within the eastern
region, populations are separated by latitude in phylogenetic analyses (Fig. 2C).

The southern region maintains a breadth of genetic variation that is comparable to the
variation of the eastern region (Fig. 4B). There is a negative correlation between latitude
and genetic diversity across the two populations. All southern region samples were found
to be composed of admixed ancestry (Figure 2B). The southern region has a predominant
ancestry layer that is more rarely found in eastern and western regions. The two southern
populations differ in the makeup of their inferred minor ancestry layers. The Texas
population shows more admixture with the eastern region, and the Oklahoma population
shows more admixture with the western region.

The western region contains the least amount of nucleotide variation (Figure 4B). The
same negative correlation between latitude and genetic variation is observed in western
samples (Figure 4A). Almost all western samples were inferred to be overwhelmingly made

up of a single ancestry group (Figure 2B). The reduced levels of polymorphism in the west result in strong phylogenetic clustering that largely groups with geography. Kansas samples are paraphyletic with Nebraska samples forming a monophyletic clade that is embedded within the larger western clade. All western samples form a monophyletic clade that is sister to a sample from the Oklahoma population (Figure 2C).

Geographic origin of S. integrifolium

It has been claimed that *S. integrifolium* originated from within the historical range of the American tallgrass prairie, *i.e.* the "prairie origin hypothesis" (Settle and Fisher 1972). The close association of the species with the prairie habitat and the inferred young age of the species were cited as evidence for this assertion. More recently, a phylogenetic treatment of the Silphium genus reconstructed *S. integrifolium* to be most closely related to the common prairie cup plant *S. perfoliatum*, and *S. wasiotense* (Clevinger and Panero 2000). Medley (1989) used populations endemic to the dry-mesic, mixed hardwood forests of eastern Kentucky to describe *S. wasiotense* and estimated the species to be closely related to other congeners found on limestone and sandstone soils throughout the unglaciated Cumberland Plateau (*S. brachiatum* and *S. mohrii*). The present study represents the first application of population genetic data to the topic of identifying the geographic orgin of *S. integrifolium*. Our results contradict the "prairie origin hypothesis" and suggest that the geographic origin of *S. integrifolium* is in the American southeast.

We hypothesize that *S. integrifolium* has expanded its range from unglaciated regions near present day Mississippi or Texas to cover the American prairie *i.e.* "refugium origin hypothesis". The evidence for our refugium origin hypothesis is multifold. First, there is a consistent correlation between latitude and diversity. Mississippi is the most genetically diverse population sampled. Texas, in the southern region, is also very diverse. However the level of divergence found between the two populations is within the range of variation found within Mississippi, but not Texas. From the sequence divergence data, this pattern suggests that the Texas population originated from Mississippi plants, however more sampling is required to better understand the progenitor taxon in this split. Results from demographic modeling using three-dimensional joint site frequency spectra generally support the refugium origin hypothesis (Figure 5), however confidence intervals around exact parameter estimates are broad (Table 3). We hypothesize that after the split between Mississippi and Texas populations, both eastern and southern lineages expanded northward, reducing genetic variation with northward progression. The expansion of Oklahoma populations into Kansas and Nebraska resulted in a large reduction in genetic diversity. This contraction appears to be most recent and results in western samples still having considerable amounts of the ancestry layer that is dominant in southern samples (Figure 2A, 2B).

The overall negative values of Tajima's D in all populations, excluding Illinois, suggest a relatively recent population expansion for the entire species. We interpret the patterns of Tajima's D across populations to help us identify the most recent population expansion. Western and southern region populations have the largest negative values of Tajima's D,

indicating a large excess of rare variants possibly due to a very recent population expansion after bottlenecks. The smaller negative Tajima's D values found in other eastern populations may be the result of a more modest and/or a more ancient expansion, after which genetic variation has started to accumulate at sites with initially rare variants.

The results of our phylogenetic analyses support the refugium origin hypothesis. We observe a series of nested relationships within the `SVD Quartets` phylogeny (Fig. 2C). Western samples form a monophyletic clade that is nested within a larger clade also containing southern samples. The larger clade containing all southern and western samples is sister to an eastern sample clade. Based on pubescence and number of ray florets of our specimens (unpublished data) we suggest that the eastern clade corresponds to the subspecies *S. integrifolium var. integrifolium* and the other clade corresponds to the subspecies *S. integrifolium var. laeve* (`http://www.efloras.org`). The high levels of observed variation in Mississippi are consistent with samples being placed throughout the eastern region clade and outside of the larger *S. integrifolium* clade. The results of our `TWISST` analyses suggest that the general topological pattern we observe in our `SVD Quartets` phylogeny has support across the majority of transcriptome markers. The southern and western regions are found to be sister taxa to each other more often than either is to the eastern region taxon. This pattern is consistent with our demographic inference results that western and southern regions have diverged most recently within this species, and that their common ancestor split from an eastern region population (Fig 5). Southern and western regions are found to be sister to the eastern region in equal proportions. This pattern is expected under the refugium origin hypothesis.

The two competing hypotheses are not necessarily mutually exclusive. The prairie origin hypothesis (Settle and Fisher 1970, 1972) coupled with the historical range of the prairie (Kurtz 2013) implicitly suggested to many that the central prairie was the center of *S. integrifolium* diversity. At its largest expanse, the historical prairie included a small amount of habitat in present day Mississippi that was geographically separated from the larger contiguous prairie (Kurtz 2013). We can not rule out that ancestral *S. integrifolium* originated in the small, restricted prairie habitat of Mississippi. However, an origin in the geographic center of the current prairie is not consistent with the genetic variation observed. The best remaining hypotheses, given our observed patterns of genetic variation, would be an origin in the southeastern edge of its current range. A more thorough sampling of Mississippi and Texas *S. integrifolium* and wide geographic sampling of *S. wasiotense*, and *S. perfoliatum* are recommended to further investigate the geographic origin of *S. integrifolium* and interspecific relationships.

Potentially adaptive candidate genes

Outlier analyses based on BayeScan indicate that several dozens of the genes with the highest average divergence have SNPs that are statistical outliers, falling outside the distribution of most transcriptome markers. Several of these genes show interesting annotations. Together, these candidate loci may have been involved in adaptation to the drier environment with more extreme temperatures experienced during *S. integrifolium*'s

hypothesized expansion out of the American southeast.

The SNPs with extremely low false discovery rate ($q < 0.001$) have patterns of genetic variation highly unlikely to have occurred by chance. The best explanation for these patterns is that they are very tightly linked to genetic variation under selection during local adaptation, and indeed may be in genes important for adaptation. These genes encode proteins involved in transcriptional regulation (Transcriptional corepressor SEUSS), secondary compound metabolism, disease tolerance, and abiotic stress responses (Delta(8)-fatty-acid desaturase, aquaporin TP1), all of which could have been important for adapting to local conditions. In order to better understand the local environment facets that were potential drivers of adaptation, we looked for climatic variables that were correlated with each selected protein across the landscape. The large number of mitochondrial and chloroplast targeted proteins suggests a possible role for cytonuclear interactions. The presence of several genes involved in cytokinin responses may be explained by altered root-shoot ratios being important in developing the extreme drought tolerance for which this species is known. In addition to being candidate genes that may underlie local adaptation, the high divergence at many of these proteins make them interesting to examine further as targets for breeding, particularly in cases where they may alter phenotypes important for domestication. As an example, the variation found in 13S globulin seed storage protein 2 is potentially important because in other species this protein family is a major driver of seed nutritional content.

Including diversity into breeding program

Perhaps motivated by the prairie origin hypothesis, the overwhelming majority of germplasm involved in *S. integrifolium* breeding programs derive from collections from the western region (Vilela et al. 2018). From those limited collections, significant heritable domestication improvements have been made in the accumulation of above ground biomass and seed yield (Vilela et al. 2018). Our population genetic results suggest that past domestication improvements have been achieved with a limited genetic variation resource. The western region, while containing considerable genetic variation, is depauperate of variation relative to the eastern and southern regions. Evidence for population recovery from a recent genetic bottleneck (e.g., low Tajimas D values, Fig. 6) could imply that potentially useful variation may have been lost during adaptation to this drought prone climate.

The elevated genetic variation across the rest of the species range and the increased divergence between regions is evidence that useful variation that may be beneficial to domestication efforts can be further incorporated into the breeding program. Characterizing the potential primary gene pool of a crop can help breeders identify useful parents in at least two ways. First, demonstrating trait or allele association with climatic, soil or geographic variables could enable breeders to predict which wild populations might produce individuals with extreme values for traits of interest. Recent studies have found that the domestication-targeted traits, seed size and seed oil profile, in wild populations have clear geographic patterns (Reinert et al. 2019). Second, information about the

distribution of genetic variation within and between populations and phylogenetic relationships between populations may inform the sampling strategy for a wild germplasm collection campaign designed to capture the most genetic variation in the least number of accessions that need to be subsequently characterized and maintained. The kind of genetic variation thus preserved ex situ may require careful screening (*e.g.* for resistance to a particular pathogen) to be valuable to breeders, or may be held in reserve for future breeders seeking new genetic variation to adapt crops to evolving pests or climate change. This information should also help inform in situ conservation strategies, and in this case it argues for high priority to be placed on conserving rare native prairies in the southern United States (Campbell and Seymour 2012) as these appear to be the center of origin of at least one *Silphium* species and perhaps a similar pattern will be found for other native grassland species.

Acknowledgements

The authors thank (1) John Holmquist, who found and collected many of the wild populations and helped with RNA extractions, (2) other members of the public, botanical gardens, and prairie conservation organizations who have donated wild *Silphium* germplasm to the The Land Institute's collection over the years, and (3) the financial support of the Perennial Agriculture Project and The Land Institute's donors.

Supporting information

Additional Supporting Information may be found online in the supporting information section at the end of the article.

Literature Cited

Andreasen, K. and B. G. Baldwin, 2001. Unequal evolutionary rates between annual and perennial lineages of checker mallows (Sidalcea, Malvaceae): evidence from 18s–26s rDNA internal and external transcribed spacers. Molecular Biology and Evolution 18:936–944.

Bai, C., W. S. Alverson, A. Follansbee, and D. M. Waller, 2012. New reports of nuclear DNA content for 407 vascular plant taxa from the United States. Annals of botany 110:1623–1629.

Boutet, E., D. Lieberherr, M. Tognolli, M. Schneider, P. Bansal, A. J. Bridge, S. Poux, L. Bougueleret, and I. Xenarios, 2016. UniProtKB/Swiss-Prot, the manually annotated section of the UniProt KnowledgeBase: how to use the entry view. Pp. 23–54, *in* Plant Bioinformatics. Springer.

Bradburd, G. S., G. M. Coop, and P. L. Ralph, 2018. Inferring continuous and discrete population genetic structure across space. Genetics 210:33–52.

Bryant, D. M., K. Johnson, T. DiTommaso, T. Tickle, M. B. Couger, D. Payzin-Dogru, T. J. Lee, N. D. Leigh, T.-H. Kuo, F. G. Davis, et al., 2017. A tissue-mapped axolotl de novo transcriptome enables identification of limb regeneration factors. Cell reports 18:762–776.

Bushnell, B., 2018. Bbtools: A suite of fast, multithreaded bioinformatics tools designed for 493 analysis of DNA and RNA sequence data. Joint Genome Institute: https://jgi. doe. gov/494 data-and-tools/bbtools 495.

Butters, J., B. J. Spiesman, E. G. Murrell, and T. K. Kim, 2019. Living on the edge: Determining the potential of multipurpose perennial crops as habitat for native pollinators and natural enemies. *in* Entomology.

Campbell, J. and W. Seymour, 2012. The flora of Pulliam Prairie, Chickasaw County, Mississippi: a significant remnant of native vegetation in the Black Belt region. J. Mississippi Acad. Sci., in press.[For correct versions of all 3 papers, see following web address.]¡ http://bluegrasswoodland. com/Mississippi_Black_Belt. html .

Chen, M., J. E. Markham, and E. B. Cahoon, 2012. Sphingolipid $\delta 8$ unsaturation is important for glucosylceramide biosynthesis and low-temperature performance in Arabidopsis. The plant journal 69:769–781.

Chifman, J. and L. Kubatko, 2014. Quartet inference from snp data under the coalescent model. Bioinformatics 30:3317–3324.

Clevinger, J. A. and J. L. Panero, 2000. Phylogenetic analysis of Silphium and subtribe Engelmanniinae (Asteraceae: Heliantheae) based on ITS and ETS sequence data.

American Journal of Botany 87:565–572. URL
http://www.jstor.org/stable/2656600.

Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E.
Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, et al., 2011. The variant call format
and VCFtools. Bioinformatics 27:2156–2158.

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut,
M. Chaisson, and T. R. Gingeras, 2013. STAR: ultrafast universal RNA-seq aligner.
Bioinformatics 29:15–21.

Emms, D. M. and S. Kelly, 2015. Orthofinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome biology
16:157.

Fiedler, A. K. and D. Landis, 2007. Attractiveness of Michigan native plants to arthropod
natural enemies and herbivores. Environmental entomology 36:751–765.

Foll, M., 2012. Bayescan v2. 1 user manual. Ecology 20:1450–1462.

Foll, M. and O. Gaggiotti, 2008. A genome-scan method to identify selected loci
appropriate for both dominant and codominant markers: a Bayesian perspective.
Genetics 180:977–993.

Fox, D. L. and P. L. Koch, 2003. Tertiary history of C4 biomass in the Great Plains, USA.
Geology 31:809–812.

———, 2004. Carbon and oxygen isotopic variability in Neogene paleosol carbonates:
constraints on the evolution of the C4-grasslands of the Great Plains, USA.
Palaeogeography, Palaeoclimatology, Palaeoecology 207:305–329.

Garrison, E. and G. Marth, 2012. Haplotype-based variant detection from short-read
sequencing. arXiv preprint arXiv:1207.3907 .

Günther, T. and G. Coop, 2013. Robust identification of local adaptation from allele
frequencies. Genetics 195:205–220.

Haas, B. and A. Papanicolaou, 2017. Transdecoder https://transdecoder. github. io.

Haas, B. J., A. Papanicolaou, M. Yassour, M. Grabherr, P. D. Blood, J. Bowden, M. B.
Couger, D. Eccles, B. Li, M. Lieber, et al., 2013. De novo transcript sequence
reconstruction from RNA-seq using the Trinity platform for reference generation and
analysis. Nature protocols 8:1494.

Isaacs, R., J. Tuell, A. Fiedler, M. Gardiner, and D. Landis, 2009. Maximizing
arthropod-mediated ecosystem services in agricultural landscapes: the role of native
plants. Frontiers in Ecology and the Environment 7:196–203.

Jouganous, J., W. Long, A. P. Ragsdale, and S. Gravel, 2017. Inferring the joint demographic history of multiple populations: beyond the diffusion approximation. Genetics 206:1549–1567.

Knapp, A. K., J. M. Briggs, S. L. Collins, S. R. Archer, M. S. BRET-HARTE, B. E. Ewers, D. P. Peters, D. R. Young, G. R. Shaver, E. Pendall, et al., 2008. Shrub encroachment in North American grasslands: shifts in growth form dominance rapidly alters control of ecosystem carbon inputs. Global Change Biology 14:615–623.

Kurtz, C., 2013. A practical guide to prairie reconstruction. University of Iowa Press.

Langmead, B. and S. L. Salzberg, 2012. Fast gapped-read alignment with Bowtie 2. Nature methods 9:357.

Lauenroth, W. K., 1979. Grassland primary production: North American grasslands in perspective. Pp. 3–24, *in* Perspectives in grassland ecology. Springer.

Leopold, A., 1968. A Sand County almanac, and sketches here and there. Oxford University Press, USA.

Li, B. and C. N. Dewey, 2011. RSEM: accurate transcript quantification from RNA-seq data with or without a reference genome. BMC bioinformatics 12:323.

Liu, A. and J. M. Burke, 2006. Patterns of nucleotide diversity in wild and cultivated sunflower. Genetics 173:321–330.

Lü, S., H. Zhao, D. L. Des Marais, E. P. Parsons, X. Wen, X. Xu, D. K. Bangarusamy, G. Wang, O. Rowland, T. Juenger, et al., 2012. Arabidopsis eceriferum9 involvement in cuticle formation and maintenance of plant water status. Plant physiology 159:930–944.

Madden, T., 2013. The BLAST sequence analysis tool. *in* The NCBI Handbook [Internet]. 2nd edition. National Center for Biotechnology Information (US).

Martin, S. H. and S. M. Van Belleghem, 2017. Exploring evolutionary relationships across the genome using topology weighting. Genetics 206:429–438.

Matthews, W. J., 1988. North American prairie streams as systems for ecological study. Journal of the North American Benthological Society 7:387–409.

Maurel, C., J. Reizer, J. I. Schroeder, and M. J. Chrispeels, 1993. The vacuolar membrane protein gamma-TIP creates water specific channels in Xenopus oocytes. The EMBO Journal 12:2241–2247.

Medley, M., 1989. Silphium wasiotensis (Asteraceae), a new species from the Appalachian plateaus in eastern Kentucky. SIDA, Contributions to Botany 13:285–291.

Nei, M., 1972. Genetic distance between populations. The American Naturalist 106:283–292.

———, 1987. Molecular evolutionary genetics. Columbia university press.

Paradis, E., 2010. pegas: an R package for population genetics with an integrated–modular approach. Bioinformatics 26:419–420.

Pembleton, L., N. Cogan, and J. Forster, 2013. StAMPP: an R package for calculation of genetic differentiation and structure of mixed-ploidy level populations. Molecular Ecology Resources 13:946–952.

Picard Tools, 2018. URL `http://broadinstitute.github.io/picard/`.

Pickrell, J. K. and J. K. Pritchard, 2012. Inference of population splits and mixtures from genome-wide allele frequency data. PLoS genetics 8:e1002967.

Pritchard, J. K., M. Stephens, and P. Donnelly, 2000. Inference of population structure using multilocus genotype data. Genetics 155:945–959.

Reich, D., K. Thangaraj, N. Patterson, A. L. Price, and L. Singh, 2009. Reconstructing Indian population history. Nature 461:489–494.

Reinert, S., D. L. Van Tassel, B. Schlautman, N. C. Kane, and B. S. Hulke, 2019. Assessment of the biogeographical variation of seed size and seed oil traits in wild Silphium integrifolium Michx. genotypes. Plant Genetic Resources Pp. 1–10.

Samson, F. B., F. L. Knopf, and W. R. Ostlie, 2004. Great Plains ecosystems: past, present, and future. Wildlife Society Bulletin 32:6–15.

Sayers, E. W., T. Barrett, D. A. Benson, E. Bolton, S. H. Bryant, K. Canese, V. Chetvernin, D. M. Church, M. DiCuccio, S. Federhen, et al., 2009. Database resources of the national center for biotechnology information. Nucleic acids research 38:D5–D16.

Scythe, 2014. A Bayesian adapter trimmer. URL `https://github.com/vsbuffalo/scythe`.

Settle, W. J. and T. R. Fisher, 1970. The varieties of Silphium integrifolium. Rhodora 72:536–543.

———, 1972. A biosystematic study of Silphium integrifolium Michaux (compositae). The Ohio Journal of Science 72:254–265.

Simão, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov, 2015. Busco: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics 31:3210–3212.

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30:1312–1313.

Stanford, G., 1990. Silphium perfoliatum (cup-plant) as a new forage. *in* Proceedings of the Twelfth North American Prairie Conference, Cedar Falls, IA, Pp. 33–37.

Sterck, L., S. Rombauts, K. Vandepoele, P. Rouze, and Y. Van de Peer, 2007. How many genes are there in plants (and why are they there)? Current opinion in plant biology 10:199–203.

Swofford, D. L., 2000. Paup 4.0.

Ustak, S. and J. Munoz, 2018. Cup-plant potential for biogas production compared to reference maize in relation to the balance needs of nutrients and some microelements for their cultivation. Journal of environmental management 228:260–266.

Van Tassel, D. L., K. A. Albrecht, J. D. Bever, A. A. Boe, Y. Brandvain, T. E. Crews, M. Gansberger, P. Gerstberger, L. González-Paleo, B. S. Hulke, et al., 2017. Accelerating silphium domestication: An opportunity to develop new crop ideotypes and breeding strategies informed by multiple disciplines. Crop Science 57:1274–1284.

Vilela, A., L. González-Paleo, K. Turner, K. Peterson, D. Ravetta, T. Crews, and D. Van Tassel, 2018. Progress and bottlenecks in the early domestication of the perennial oilseed Silphium integrifolium, a sunflower substitute. Sustainability 10:638.

Wang, C., Z. A. Szpiech, J. H. Degnan, M. Jakobsson, T. J. Pemberton, J. A. Hardy, A. B. Singleton, and N. A. Rosenberg, 2010. Comparing spatial maps of human population-genetic variation using Procrustes analysis. Statistical applications in genetics and molecular biology 9.

Wang, T., A. Hamann, D. Spittlehouse, and C. Carroll, 2016. Locally downscaled and spatially customizable climate data for historical and future periods for North America. PLoS One 11:e0156720.

Weaver, J. E., 1954. North American Prairie. Johnsen Publishing Company, USA.

Weaver, J. E., L. Stoddart, and W. Noll, 1935. Response of the prairie to the great drought of 1934. Ecology 16:612–629.

Weir, B. S. and C. C. Cockerham, 1984. Estimating F-statistics for the analysis of population structure. evolution 38:1358–1370.

Wever, C., M. Höller, L. Becker, A. Biertümpfel, J. Köhler, D. van Inghelandt, P. Westhoff, R. Pude, and E. Pestsova, 2019. Towards high-biomass yielding bioenergy crop Silphium perfoliatum L.: phenotypic and genotypic evaluation of five cultivated populations. Biomass and Bioenergy 124:102–113.

Williams, J. W., B. N. Shuman, T. Webb III, P. J. Bartlein, and P. L. Leduc, 2004. Late-Quaternary vegetation dynamics in North America: scaling from taxa to biomes. Ecological Monographs 74:309–334.

Wilson, G. W. and D. C. Hartnett, 1998. Interspecific variation in plant responses to mycorrhizal colonization in tallgrass prairie. American journal of botany 85:1732–1738.

Table 1: Analysis of molecular variance (AMOVA) based on hierarchical grouping stratification among regions, among populations within regions, and within populations. $** = p \leq 0.01$, $*** = p \leq 0.0001$

Sources of variation	df	SS	Variance components	Percent of variance
Regions	2	0.44905	0.00796**	29.4869%
Populations	6	0.23872	0.00375***	13.9031%
Within Populations	58	0.88744	0.01530	56.610%
Total	66	1.57522	0.027028	100.00%

Table 2: Results from f_3 tests for introgression to test the hypothesis that the Illinois population is admixed. Significantly negative test statistics indicate that IL population has history of a non tree-like branching process *i.e.* past introgression between populations P1 and P2.

Focal	P1	P2	f3	std. error	Z–score	p value
IL	WI	MS	−0.0012	0.0010	−1.1329	1.286e−01
IL	WI	AR	−0.0004	0.0011	−0.3531	3.620e−01
IL	WI	MO	−0.0004	0.0010	−0.3724	3.548e−01
IL	WI	TX	−0.0031	0.0011	−2.7611	2.881e−03
IL	WI	OK	−0.0042	0.0012	−3.5755	1.748e−04
IL	WI	KS	−0.0045	0.0010	−4.3112	8.118e−06
IL	WI	NE	−0.0044	0.0011	−4.1234	1.867e−05

Topology	AIC	loglik	s0	s1	nu12	nu0	nu1	nu2	m0_12	m0_2	m0_1	m2_1	T1	T2
(East, (South, West))	23316.94	-11646.47	0.614 (0.503)	0.837 (0.022)	0.753 (1.865)	0.596 (1.06)	0.323 (0.378)	0.476 (0.079)	1.507 (11.426)	1.557 (0.38)	1.833 (4.908)	1.816 (0.867)	1.848 (3.91)	0.818 (2.712)
(South, (East, West))	23324.14	-11650.07	0.489 (0.092)	0.805 (0.211)	0.308 (0.202)	0.543 (0.097)	0.347 (0.258)	0.744 (0.662)	0.573 (0.74)	1.326 (1.459)	1.982 (0.036)	1.386 (0.841)	1.735 (0.094)	0.609 (0.027)
(West, (South, East))	23344.46	-11660.23	0.3 (1.054)	0.516 (0.005)	0.513 (0.332)	0.991 (1.011)	0.688 (0.333)	0.507 (0.233)	1.303 (3.305)	1.397 (0.041)	1.654 (0.651)	1.925 (1.128)	1.403 (6.349)	0.544 (0.027)

Figure 1: Results from a principal components analysis of *S. integrifolium* genotypes. The two major axes of variation distinguish samples (top panel). A Procrustes transformation of PC coordinates onto latitude and longitude coordinates of *S. integrifolium* genotypes displays relationship between geography and genetic variation (bottom panel). Abbreviated population text locations are the Procrustes transformed principle component coordinates. Lines point from transformed coordinates to geographic sampling locations.

Figure 2: *Silphium integrifolium* samples comprise three genetic groups. **Top panels:** Inferred ancestry results of *Silphium integrifolium* samples from a `conStruct` model that includes spatial information and K = 3 (Bradburd et al. 2018). **(A)**: Average amounts of inferred ancestry from three genetic layers for each population are displayed as pie charts. **(B)**: Proportion of inferred ancestry of each genetic layer for all sampled plants. **Bottom panels:** Patterns of phylogenetically reconstructed relationships among samples. **(A)**SVD Quartets reconstruction of phylogenetic relationships between *S. integrifolium* samples. Bootstrap node support values are not shown, however every node has 100% bootstrap support. **(B)** Summary of `TWISST` support for the `SVD Quartets` inferred phylogeny topology. The region of the sister tip for each focal taxon was recorded for each phylogeny subsampled in `TWISST` analysis across all marker contigs. The excess of southern and western tips being found as sister supports their hypothesized most recent split.

Figure 3: Relationships between geographic distance and genetic differentiation. Geographic distance was measured as Euclidean distance between sample midpoints within each population. Differences in color reference within and between region comparisons; colored letters (E, S, W) represent within region comparisons and gray letter pairs (E/S, E/W, S/W) represent between region comparisons.

Figure 4: Distribution of nucleotide diversity at four fold degenerate sites. **(A)** Mean nucleotide diversity between samples from each population. As a general pattern, diversity decreases with latitude. **(B)** Pairwise nucleotide diversity among samples within regions (π) and **(C)** among samples between geographic regions (D_{xy}).

Figure 5: Summarized results of best fitting demographic inference model (symmetric migration rates) for each lineage branching scenario. Width of population segment is equal to population size relative to ancestral population size. Migration rate estimates are reported in each figure. Estimated model parameters with standard deviations and model fit statistics are listed in Table 3. Time and migration rates are in genetic units $(2N_e)$.

Figure 6: Distribution of Tajima's D across contigs within populations. Colors correspond to different regions.