Changes in local mineral homeostasis facilitate the formation of benign and malignant testicular microcalcifications

Ida Marie Boisen[1,2], Nadia Krarup Knudsen[1], John Erik Nielsen[3], Ireen Kooij[1], Mathilde Louise Bagger[1], Jovana Kaludjerovic[4], Peter J. O'Shaughnessy[5], Peter W. Andrews[6], Noriko Ide[4], Birgitte Grønkær Toft[7], Anders Juul[3], Arnela Mehmedbašić[1], Anne Jørgensen[1], Lee B. Smith[8], Richard W. Norman[9], Ewa Rajpert-De Meyts[3], Beate Lanske[4], Martin Blomberg Jensen[1]*

1: Division of Translational Endocrinology, Department of Endocrinology and Internal Medicine, University Hospital Copenhagen, Herlev-Gentofte, DK-2730 Herlev, Denmark

2: Group of Skeletal, Mineral, and Gonadal Endocrinology, Department of Growth and Reproduction, Rigshospitalet, University of Copenhagen, DK-2100 Copenhagen, Denmark

3: Department of Growth and Reproduction, Rigshospitalet, University of Copenhagen, DK-2100 Copenhagen, Denmark

4: Division of Bone and Mineral Research, Harvard School of Dental Medicine/Harvard Medical School, Harvard University, Boston, MA 02108, US

5: College of Medical, Veterinary and Life Sciences, Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, G61 1QH, Glasgow, UK

6: Centre for Stem Cell Biology, Department of Biomedical Science, University of Sheffield, Western Bank, Sheffield S10 2TN, UK

7: Department of Pathology, Rigshospitalet, DK-2100 Copenhagen, Denmark

8: MRC Centre for Reproductive Health, University of Edinburgh, The Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh, EH16 4TJ, UK

9: Department of Urology, Dalhousie University, Halifax NS B3H 2Y9, Nova Scotia, Canada

Keywords: FGF23, microcalcifications, testicular germ cell cancer, reproduction, testis

Abstract: 249 words

Word count: 4814

Figures: 8

Tables: 0

*Address for correspondence

Martin Blomberg Jensen

Martin.blomberg.jensen@regionh.dk

All authors declare no conflicts of interests

Abstract

Testicular microcalcifications consist of hydroxyapatite and have been associated with an increased risk of testicular germ cell tumors (TGCTs) but may also be the result of benign causes such as loss-of-function variants in the phosphate-transporter gene *SLC34A2*. Here, we show that fibroblast growth factor 23 (FGF23), a regulator of phosphate homeostasis, is expressed in testicular germ cell neoplasia *in situ* (GCNIS), embryonal carcinoma (EC), and human embryonic stem cells. FGF23 is not glycosylated in TGCTs and therefore cleaved into a C-terminal fragment which competitively antagonizes full-length FGF23. Here, *Fgf23* knockout mice presented with marked calcifications in the epididymis, spermatogenic arrest, and focally germ cells expressing the osteoblast marker bone gamma-carboxyglutamate protein (BGLAP). Moreover, the frequent testicular microcalcifications in mice with no functional androgen receptor and lack of circulating gonadotropins is associated with lower *Slc34a2* and higher *Slc34a1/Bglap* expression compared with wild-type mice. In accordance, human testicular specimens with microcalcifications also have lower *SLC34A2* and a subpopulation of germ cells express SLC34A1, BGLAP, and RUNX2 highlighting aberrant local phosphate handling and expression of bone-specific proteins. Mineral disturbance *in vitro* using calcium or phosphate treatment induced deposition of calcium-phosphate in a spermatogonial cell line and this effect was fully rescued by the mineralization-inhibitor pyrophosphate. In conclusion, testicular microcalcifications may arise secondary to local alterations in mineral homeostasis, which in combination with impaired Sertoli cell function and reduced levels of mineralization-inhibitors due to high alkaline phosphatase activity in GCNIS and TGCTs, facilitate osteogenic-like differentiation of testicular cells and deposition of hydroxyapatite.

Introduction

Testicular microlithiasis in infertile men is associated with an increased risk of testicular germ cell tumors (TGCTs)[1-4]. However, both ultrasonic and histological microcalcifications are also frequent findings in the testis of asymptomatic men and the etiology remains largely unknown in both malignant and benign cases. The histological microcalcifications have been shown by Raman spectroscopy to consist of hydroxyapatite[5,6], which is normally found exclusively in the skeleton and requires the presence of cells with osteoblast-like characteristics[7,8]. Extra-skeletal mineralization is often caused by an imbalance in promoters and inhibitors of biomineralization[7,9-11]. Tissue mineralization is tightly controlled by small integrin-binding ligand N-linked glycoproteins (SIBLINGs) such as osteopontin (OPN) and dentin matrix acidic phosphoprotein 1 (DMP1) that regulate mineralization in concert with osteoblast-specific factors such as bone gamma-carboxyglutamic acid-containing protein (BGLAP) and runt-related transcription factor 2 (RUNX2). Pyrophosphate (PPi) generated by ectonucleotide pyrophosphatase/phosphodiesterase 1 (ENPP1) is an essential suppressor of hydroxyapatite deposition and alkaline phosphatase (ALP) inactivates PPi[12]. ALP abundance is thus a central factor for local phosphate homeostasis, and during osteogenic differentiation increased ALP activity is a hallmark of the bone-like characteristics of the cells in other extra-skeletal tissue such as blood vessels. In this paper, ALP is used as an acronym for all four isozymes of ALP as all can be found in TGCTs[13]. In addition, accumulating evidence suggests that fibroblast growth factor 23 (FGF23) and its receptor Klotho, the phosphate transport proteins solute carrier family SLC34A-1, -2, and -3, and vitamin D metabolizing enzymes are essential for maintaining stable mineral homeostasis and that imbalance in their activity may lead to extra-skeletal calcifications[14-18]. Interestingly, loss-of-function variants in the *GALNT3* gene and hence a lack of degradation-protective glycosylation of FGF23 leads to systemic hyperphosphatemia and severe testicular microcalcifications[19-21]. Intratesticular phosphate levels seem to be essential in the formation of testicular microlithiasis as loss-of-function variants in the main testicular phosphate transporter *SLC34A2* also lead to severe testicular microcalcifications without affecting systemic phosphate levels[22]. Testicular microlithiasis is also found in patients with loss-of-function variants in the gene encoding ATP binding cassette subfamily C member 6 (*ABCC6*), which results in abnormal PPi metabolism and thus less inhibition of mineralization[23]. These studies indicate that particularly local and potentially systemic changes in phosphate homeostasis induce testicular microcalcifications. Therefore, we hypothesized that the etiology of testicular microcalcifications involves disturbances in intratesticular mineral homeostasis.

TGCTs originate from a common progenitor germ cell neoplasia *in situ* (GCNIS), which is characterized by the expression of pluripotency factors such as octamer-binding transcription factor 4 (OCT4), NANOG, and high alkaline phosphatase activity[13,24-27]. GCNIS cells undergo malignant transformation and form either a seminoma that retains germ cell characteristics or a non-seminoma that may contain embryonal carcinoma (EC) resembling human embryonic stem cells (hESCs), teratoma, yolk sac tumor, choriocarcinoma, or a combination of these[28]. During the malignant transformation of GCNIS into invasive TGCTs duplication of chromosome 12p (often as isochromosome 12p) is the most consistent chromosomal aberration, and may be involved in the dedifferentiation from GCNIS to EC[28]. In healthy individuals, FGF23

is produced exclusively in osteocytes, and the location of the gene on chromosome 12p may be important for its presence and role in TGCTs. Full-length FGF23 increases phosphate excretion by decreasing the luminal expression of SLC34A1 in the kidney and by lowering the concentration of active vitamin D via increased expression of the inactivating enzyme CYP24A1 and reduced expression of the activating enzyme CYP27B1[29]. We have previously shown that the local metabolism of vitamin D disappeared during the malignant transformation of GCNIS to EC and that FGF23 was expressed in an EC-derived cell line[30]. O-glycosylation by GALNT3 protects FGF23 from rapid degradation, and full-length FGF23 activates a receptor complex consisting of Klotho heterodimerization with FGF receptor 1 (FGFR1). However, the cleaved C-terminal fragment (cFGF23) can bind and serve as a competitive antagonist for intact FGF23 (iFGF23), and iFGF23 is also capable of inducing Klotho-independent effects[31]. Loss-of-function variants in the genes *FGF23*, *GALNT3*, or *Klotho* lead to tumoral calcinosis partly through osteogenic-like differentiation of mesenchymal-derived cells due to altered mineral homeostasis and vitamin D metabolism[32]. Here, we investigated whether ectopic FGF23 production and abundant ALP activity in TGCTs alter testicular mineral homeostasis, which alone or in combination with impaired Sertoli cell function may facilitate osteogenic-like differentiation and thus be responsible for the frequent deposition of hydroxyapatite in TGCTs and dysgenetic testes. To address this question, we investigated the expression of mineral regulators and bone factors in human tissues containing TGCT-associated calcifications, *Fgf23* knockout (*Fgf23*$^{-/-}$) mice, hypogonadal (no gonadotropins) mice, androgen-insensitive mice with microcalcifications, mice with microcalcifications due to Sertoli cell ablation, mice xenografted with a TGCT-derived cell line, mice treated with a high phosphate diet, mineral deposition in a mouse spermatogonial cell line (GC1), and an *ex vivo* mouse testis culture model.

Results

Testicular microcalcifications are associated with the presence of renal phosphate transporters and bone markers in human testis

Ultrasonic testicular microlithiasis is often observed adjacent to TGCTs but can also be found in testes without TGCTs or GCNIS (Fig. 1A). Ultrasonic testicular microlithiasis is not always associated with histological microcalcifications but intratesticular mineral depositions can be confirmed by von Kossa or alizarin red staining (Fig. 1A), and Raman spectroscopy has previously demonstrated that testicular microcalcifications consist of hydroxyapatite[5]. In GCNIS-containing tubules adjacent to microcalcifications we found that the renal phosphate transporter SLC34A1 and FGF23 were ectopically expressed. In contrast, Klotho was expressed in the cytoplasm of the germ cells in normal tubules, with or without adjacent microcalcifications, but not in GCNIS. FGFR1 expression was found in both GCNIS and normal tubules (Fig. 1B). The expression of the bone marker BGLAP was present in tissue with GCNIS and adjacent to the microcalcifications but not the normal tubules with full spermatogenesis (Fig. 1B and Suppl. Table 1). The microcalcifications vary in size from small calcifications focally in the center of seminiferous tubules to occupying most of the intraluminal area (Fig. 1C). In very rare severe cases there was intratesticular bone formation (Fig. 1D). We investigated the presence of several bone factors in the cells adjacent to the microcalcifications. The early osteoblast marker RUNX2 was not expressed in testes with normal spermatogenesis. However, occasionally, when microcalcifications or bone formation were detected, a small fraction of the germ cells in the seminiferous tubules (ST) had cytoplasmic staining of RUNX2 supporting a bone-like phenotype of some germ cells (Fig. 1D, Suppl. Fig. 1, and Suppl. Table 1). The shift in expression of the phosphate transporters from SLC34A2 to SLC34A1 and the presence of bone factors in testis with microcalcifications were also evident at the transcriptional level. *SLC34A1* was expressed at a very low level in normal testis but was highly expressed in 25% of the specimens containing GCNIS. In contrast, the phosphate transporter *SLC34A2* was abundantly expressed in the germ cells from normal testis but the expression was lower in specimens containing GCNIS (Fig. 1E).

FGF23 is produced by human testicular germ cell tumors and embryonic stem cells

FGF23, a potent regulator of phosphate homeostasis, is expressed in the EC-derived cell line NTera2[30], and the *FGF23* gene is located on chromosome 12p which is duplicated in TGCTs and could be associated with the formation of testicular microcalcifications. Thus, expression of *FGF23* was investigated in human testicular specimens containing 'normal testis' (adjacent tissue to TGCT/GCNIS without presence of malignant cells), Sertoli-cell-only pattern, GCNIS, seminoma, and EC with or without teratoma components. The transcriptional level of *FGF23* was highest in EC components, significantly higher compared with normal testis, and was positively associated with the expression of the pluripotency factors *NANOG* and *OCT4* (R=0.66, P=0.049, and R=0.61, P=0.068, respectively). Two EC specimens with polyembryoma components (the most dedifferentiated structure resembling early embryonic formation) had several fold higher levels of *FGF23* (Fig. 2A-B). A higher expression of *FGF23* in GCNIS compared with normal testis was also confirmed in microdissected GCNIS cells,

while fetal gonocytes and human embryonic stem cells (hESCs) also expressed *FGF23* (Fig. 2C). The high expression of *FGF23* in hESC and EC was validated by a ~30-fold higher expression in different strains of hESC cells and the EC-derived cell line NTera2 compared with normal testis (Fig. 2D). In contrast, normal testicular tissue had a high expression of the receptor Klotho (*KL*), whereas the expression was low in GCNIS cells and hESC (Fig. 2C). IHC confirmed the marked expression of FGF23 solely in OCT4-positive GCNIS cells that also expressed FGFR1, while Klotho was exclusively expressed in normal germ cells and not found in any of the OCT4-positive cancer cells including seminoma and EC (Fig. 2E and Suppl. Table 1). FGF23 was also expressed in the EC cells characterized by concomitant OCT4 expression but was virtually absent in seminoma cells despite their marked OCT4 expression suggesting that the chromosome 12p duplication (occurring in both seminoma and EC) alone is not sufficient to maintain FGF23 production. Western blot confirmed the presence of FGF23 with two bands corresponding to the sizes of cFGF23 and iFGF23 in GCNIS and EC. cFGF23 was also expressed in NTera2 cells, and neither cFGF23 nor iFGF23 were detected in normal testis or the seminoma-derived cell line TCam2 (Fig. 2F). cFGF23 and iFGF23 were expressed in an anaplastic seminoma sample, which is considered a rare intermediate between seminoma and EC due to the high proliferative index and morphological resemblance with EC (Fig. 2F and Suppl. table 1). Production and release of FGF23 in GCNIS and EC cells were examined by measuring the total FGF23 (iFGF23 + cFGF23) and cFGF23 concentrations in seminal fluid from healthy men and testicular tumor patients. Healthy men had undetectable iFGF23 levels in the seminal fluid, and a few (8/19) men had detectable cFGF23. A few men with GCNIS (2/7) or EC (2/10) had detectable levels of iFGF23 in the seminal fluid while total FGF23 were detected in the majority (9/10) of men with EC and their total FGF23 concentration in seminal fluid was significantly higher compared with healthy men (Fig. 2G).

Testicular phosphate homeostasis in NTera2 xenograft mice is altered by FGF23-independent actions

In NTera2 xenograft tumors, FGF23 was expressed in only 5-10% of tumor cells, which indicates that the phenotype of the NTera2 cells changes *in vivo*. This may be caused in part by somatic differentiation of the FGF23-negative fraction comprising 90-95% of the tumors, as also illustrated by the similar expression pattern of OCT4 within the tumor (Fig. 3A). Human iFGF23 in serum from nude mice with tumors was 4-fold higher compared with control nude mice (Fig. 3B) and thus much lower than the EC tumors samples. The presence of iFGF23 in serum and the undetectable levels of iFGF23 in NTera2 cells grown *in vitro* suggests that NTera2 cells xenografted into nude mice may undergo somatic differentiation in response to host stimuli and facilitate GALNT3-mediated glycosylation of iFGF23 to prevent its cleavage (Fig. 2F). Interestingly, NTera2 tumor-bearing mice had significantly higher *Slc34a1* and *Bglap* expression in testis compared with nude mice without tumors but no change in *Slc34a2* suggesting that the effect induced by the tumor was cFGF23 or FGF23 independent (Fig. 3C). Expression of the phosphate transporter *Slc34a1* in the testis was positively correlated with the tumor size (R^2=0.53, P=0.012) (Fig. 3D). Exogenous treatment with iFGF23 or cFGF23 in human testis specimens cultured *ex vivo* for 24 h did not change the expression of phosphate transporters or *BGLAP* (Fig. 3E), indicating that FGF23 alone is not responsible for the change in expression of these genes but may be caused by other factors released or induced by the tumors.

Mineralization in testis and epididymis of *Fgf23*[-/-] mice and in mice treated with a high phosphate diet

Fgf23[-/-] mice had elevated serum concentrations of calcium and phosphate, smaller testes, and often spermatogenic arrest at the spermatocyte stage compared with control mice (Fig. 4A and data not shown). Testicular evaluation showed no deposition of hydroxyapatite, but DMP1, OPN, and BGLAP were expressed in the cytoplasm of some germ cells in up to 15% of the tubules in *Fgf23*[-/-] mice and more rarely in WT mice (Fig. 4B-C). To distinguish the effects of FGF23 loss from high phosphate alone, WT mice received a high phosphate diet, which showed that short-term high serum phosphate did not induce testicular microlithiasis or induce any testicular changes in the expression of phosphate transporters or *Bgalp* (Fig. 4D). A more severe phenotype was observed in cauda epididymis with marked mineralization in 25% of the *Fgf23*[-/-] mice and none of the WT mice. DMP1, OPN, and BGLAP were highly expressed in the epididymis of *Fgf23*[-/-] mice (Fig. 4E-F), while DMP1 and BGLAP were not detected in the epididymal lumen of WT mice. OPN was expressed in the head of sperm in both WT and *Fgf23*[-/-] mice, although it was expressed exclusively in the luminal part in the calcified section of epididymis of *Fgf23*[-/-] mice.

Testicular microcalcifications in hypogonadal, androgen receptor knockout, and Sertoli cell-ablated mice

Testicular microcalcifications are not exclusively found in mice with knockout of phosphate transport regulators but also in ~30% of hypogonadal (*hpg*) mice that lack circulating gonadotropins (follicle-stimulating hormone and luteinizing hormone). The proportion of mice with testicular microcalcifications increased to ~94% in mice with concomitant global loss of the androgen receptor (*AR*) (*hpg*.ARKO)[33] (schematically illustrated in Fig. 5A). Sertoli cell-specific ablation of *AR* in *hpg* mice (*hpg*.SCARKO) did not augment the effect on microcalcifications in the *hpg* mice, indicating a role of AR in peritubular cells (the only other cell type in the testis that express AR) or AR-expressing cells outside the testis for the formation of microcalcifications. Thus, suggesting that somatic cells may be important for the bone-like trans-differentiation in mice. Noteworthy, a shift in the expression of phosphate transporters was observed in the testis of *hpg*, *hpg*.SCARKO, and *hpg*.ARKO mice which had higher expression of *Slc34a1* and lower expression of *Slc20a1* and *Slc34a2* compared with WT mice. Thus, supporting the hypothesis that local phosphate homeostasis is involved in the formation of microcalcifications in accordance with the observations in human tissues with microcalcifications. There were no significant differences in *Slc20a1, Slc34a2*, and *Slc34a1* expression between *hpg, hpg*.SCARKO, or *hpg*.ARKO mice despite the higher proportion of mice with microcalcifications in the *hpg*.ARKO group. However*, Bglap* was detected in testis from all *hpg*.ARKO and most *hpg*.SCARKO mice suggesting that *Bglap* is expressed in mice with disturbed Sertoli cell maturation with the highest expression found in *hpg*.ARKO mice which had a 94% prevalence of microcalcifications (Fig. 5B). Ablation of Sertoli cells at postnatal day 18 induced substantial microcalcifications in the adult mice (80 days old) as shown by alizarin red and von Kossa staining (Fig. 5C). The microcalcifications persisted when assessed in mice 1-year of age (data not shown). The timing of ablation seemed to be important because Sertoli cell ablation in adulthood (after puberty) did not cause microcalcifications when evaluated after 90 days but was detectable one year later (data not shown). This shows that immature prepubertal cells are more prone to form microcalcifications. The formation of microcalcifications depends

201 on the presence of germ cells with stem cell potential as ablation of Sertoli cells during embryonic development, which

202 causes loss of germ cells, did not result in testicular microcalcifications.

Alkaline phosphatases in TGCTs and effects of calcium, phosphate, and pyrophosphate on mineralization in a
spermatogonial cell line

ALP degrades the potent inhibitor of mineralization PPi and is highly expressed in the cells of mineralized tissue. Clinically,

ALP expression is used as a marker for GCNIS and seminoma. Alkaline phosphatase activity was markedly higher in GCNIS

tubules compared with normal seminiferous tubules as determined by NBT/BCIP staining (Fig. 6A). All four types of ALPs

were expressed in normal testis and the expression in GCNIS with microlithiasis and seminoma was not significantly higher,

although some tumor specimens have marked expression of all four ALP isozymes. This shows that alkaline phosphatase

activity is higher in GCNIS/TGCTs and typically mediated by a combination of different isozymes rather than a single type

of ALP (Fig. 6B). To determine if mineralization can be induced in germ cells, the mouse spermatogonial cell line GC1 was

treated with increasing concentrations of calcium and phosphate, and mineral deposition was subsequently visualized by

alizarin red. Mineralization was already observed after co-treatment with calcium and phosphate for two days, but also

after seven days of treatment with calcium or phosphate alone (Fig. 6C). With a low serum phosphate concentration (0.9

mM), 3 mM calcium was required to obtain rapid mineralization after two days, whereas even 4 mM phosphate did not

induce mineralization in a normal calcium concentration (1.8 mM) after two or four days. However, an increment in

calcium to 2.0 mM enabled five of the six phosphate doses to induce rapid mineralization of the GC1 cells. This indicates

that low calcium levels are protective in the high intratesticular phosphate environment. Longer exposure (four and seven

days) augmented the response in a dose-dependent manner. The addition of PPi, an inhibitor of mineralization, reduced

mineralization, whereas co-treatment with PPi and pyrophosphatase (PPA1) that catalyzes the hydrolysis of

pyrophosphate (PPi) to inorganic phosphate (Pi) reintroduced mineralization (Fig. 6D). To assess the regulation of

phosphatase activity in germ cells, a human cell line TCam2 was used as a positive control. TCam2 is derived from a

seminoma and retains high expression of ectopic alkaline phosphatase, as assessed by NBT/BCIP staining, which was

critically dependent on an alkaline environment with marked phosphatase activity at pH 9, but not at pH 7 (Fig. 6E).

Staining with Fast Blue RR revealed that alkaline phosphatase activity was increased in the presence of osteogenic medium

compared with control medium, indicating that germ cells may increase their alkaline phosphatase activity in the presence

of factors that facilitate bone differentiation (Fig. 6F). However, an *ex vivo* model using wild-type mouse testicles (Fig. 7A)

cultured for 14 days with osteogenic medium -/+ phosphate did not change the expression of the bone markers *Bglap*,

Runx2, or *Alpl* (Fig. 7B). However, occasionally one or few tubules had expression of RUNX2, which was observed more

often in specimens treated with osteogenic medium than in specimens treated with vehicle (Fig. 7C). Few of the specimens

only had Sertoli cells in the tubules due to loss of germ cells and were therefore excluded from the analysis. Runx2 protein

expression was not observed in those tissue pieces.

We propose that the formation of testicular microcalcifications occurs because of one or more of the following events:

disturbed local phosphate homeostasis, decreased function of the Sertoli cells, reduction of mineralization inhibitors, or

aberrant germ cell function. These changes alone or in combination can drive osteogenic-like differentiation of germ cells

resulting in intratubular deposition of microcalcifications (Fig. 8).

Discussion

This translational study demonstrates that testicular microcalcifications, both benign and malignant, occur secondary to changes in gonadal phosphate homeostasis and are often accompanied by a subsequent osteogenic-like differentiation of germ cells or testicular somatic cells. Microcalcifications are more prevalent in testicular biopsies with GCNIS than without[34], which highlights that GCNIS or TGCTs may release factors involved in the development of microlithiasis, but the accompanied impaired Sertoli cell function appears to be of greater importance than the presence of malignant germ cells. In this manuscript, we show that impaired Sertoli cell function in both mice and humans increases the risk of testicular microlithiasis and this association seems to be more persistent than the malignant linkage.

The systemic master regulator of phosphate, FGF23, is highly expressed in GCNIS, EC, and hESC but not in classical seminoma. This suggests that the expression of FGF23 is linked with the transformation of GCNIS to the invasive EC and is not a result of duplication of chromosome 12p which is also prevalent in classical seminoma. The marked expression in EC and hESCs highlights that FGF23 is an early embryonic signal, which is further supported by the strong correlation between *FGF23* and pluripotency factors *OCT4* and *NANOG* in EC and the high *FGF23* expression in EC with polyembryoma formation. Moreover, it is in accordance with previous reports identifying *FGF23* transcripts in hESCs and during early embryonic life[35,36].

Phosphate levels in the testis are 3-fold higher than in serum[37,38]. In vascular and soft tissue calcification, FGF23-Klotho activity has been suggested to regulate ion metabolism and subsequently induce the presence of bone-like cells in the vessels[39,40]. The observed hydroxyapatite in the cauda epididymis of *Fgf23*[-/-] mice shows that loss of FGF23 signaling causes deposition of hydroxyapatite. As EC and GCNIS have no GALNT3 expression[41], FGF23 will rapidly be cleaved into cFGF23 which is in accordance with the detection of mainly cFGF23 (and not iFGF23) in seminal fluid from GCNIS and EC patients. High intratesticular cFGF23 may bind the Klotho/FGFR1 receptor and antagonize the effect of iFGF23[31] thereby inducing a phenotype resembling *Fgf23*[-/-] mice with increased expression of *Bglap*. However, in our human testis *ex vivo* model neither cFGF23 nor iFGF23 induced changes in phosphate transporters or *BGLAP* suggesting that short-term exposure (24 h) to high cFGF23 does not induce changes in phosphate transporters, bone factors, or microcalcifications. The epididymal phenotype of *Fgf23*[-/-] mice does not appear to be mediated via systemic hyperphosphatemia as a high phosphate dietary exposure for weeks did not result in testicular or epididymal microcalcifications or changes in testicular phosphate transporter expression. Instead, we have previously shown that germ cell-specific deletion of Klotho in mice leads to aberrant mineral homeostasis, particularly calcium transport through TRPV5[42], which may be important as low calcium appears to be protective for inducing microcalcifications in a high phosphate environment. Local testicular mineral levels depend mainly on the presence and activity of specific transporters and sensors of calcium and phosphate in the reproductive organs, rather than on systemic concentrations[43]. This may also explain why patients with loss-of-function variants in the predominant testicular phosphate transporter *SLC34A2* present with testicular microcalcifications[22].

Moreover, patients with loss-of-function variants in *GALNT3*, which lead to the early degradation of intact FGF23 to its C-
terminal form, present with global calcifications, severe testicular microlithiasis, and microcalcifications[19,20].

Fgf23[-/-] mice exhibit hypogonadism[44] and spermatogenic arrest, which resembles the condition of some men with
testicular dysgenesis syndrome that occasionally have testicular microcalcifications despite no malignancy[2,45]. Benign
microcalcifications may be facilitated by low levels of androgens or gonadotropins as such hormonal imbalances can
disrupt the Sertoli cell function. As a result, the germ cells may remain in their prepubertal stem cell state, rather than
maturing properly leaving them susceptible to external stimuli that could trigger abnormal differentiation. Changes in the
local testicular phosphate concentration (which is 3-fold higher than serum) could exert such a stimuli[37,38,46,47]. This
hypothesis was supported by the presence of microcalcifications in *hpg* mice with concomitant global *AR* ablation as these
mice have immature Sertoli cells and are unable to complete spermatogenesis[33]. Moreover, these mice had altered
testicular phosphate homeostasis characterized by a non-significant tendency towards lower *Slc34a2,* but significantly
higher *Slc34a1* expression compared with WT mice. However, both *hpg* and *hpg* mice with Sertoli cell-specific *AR* deletion
(*hpg.SCARKO*) had similar changes in phosphate transporters and only 30-36% frequency of microlithiasis, suggesting that
additional mechanisms are contributing to the formation of microlithiasis. The bone marker *Bglap* was detected
exclusively in all the *hpg* mice with concomitant global *AR* deletion (*hpg.ARKO*) but not in *hpg* mice with intact AR, which
supports that some testicular cells undergo osteogenic trans-differentiation, and this may facilitate or accompany more
abundant deposition of hydroxyapatite. We can not show the cell-of-origin, but we have previously shown that vitamin D
induces an osteogenic-like differentiation of NTera2 cells which start to express bone-specific proteins such as BGLAP[30].
However, the osteogenic-like differentiation in *hpg* mice occurs in non-malignant prepubertal/non-maturated testicular
cells although we cannot clarify the cell-of-origin.

Peritubular cells may be a good candidate because they are mesenchymal-derived and such fibroblast-like cells can
undergo osteogenic-like differentiation during vascular calcification at other sites of extra-skeletal mineralization[48].
However, human Sertoli cells can also form large-cell calcifying tumors[49] indicating that multiple gonadal cell types have
the potential to become bone-like. In mice, Sertoli cell-ablation before puberty caused extensive intratubular
microcalcifications[50], whereas Sertoli cell-ablation in adulthood did not result in the formation of microcalcifications when
assessed three months later despite a greater burden of immediate intratubular cell death. This finding is important
because it demonstrates that microcalcifications are not caused by massive intratubular cell death in mice but are critically
dependent on germ cells with stem cell potential. Thus, indicating that Sertoli cell maturation and function are essential
for protection against osteogenic-like differentiation of the germ cells or peritubular cells and that benign
microcalcifications are more likely to occur as a consequence of prepubertal Sertoli cell disturbance. The spermatogonial
stem cells may also be the cell type that undergoes osteogenic-like differentiation as they can spontaneously develop into
all three germ layers when isolated from adult mice, form teratomas in nude mice, and participate in the formation of
multiple organs when injected into blastocytes[51]. RUNX2 is an essential transcription factor for osteoblast development,

and although germ cells express a different isoform of RUNX2 (lacking exon 4-8), it is transcribed from the same promoter as in bone[52,53]. RUNX2 was undetectable in healthy seminiferous tubules using an antibody targeting exon 5, but was detected in GCNIS, hyalinized/calcified tissues, and the otherwise seemingly healthy seminiferous tubules adjacent to bone in the testis, which could indicate that the bone-specific RUNX2 isoform may also be expressed in the testis when microcalcifications occur. Also, the mature bone marker BGLAP was expressed in a fraction of cells that morphologically resemble germ cells. Unfortunately, there was no cell-tracing marker in the mouse models so the required tracing experiments to support that these bone-like cells originated from the germ cells could not be conducted. Instead, a spermatogonial mouse cell line was exposed to calcium and phosphate at concentrations similar to those found in the testis or the epididymis. After 2-7 days of exposure, the spermatogonia started to deposit minerals illustrating that they developed some osteoblast-like characteristics, but they did not undergo complete osteogenic differentiation. A similar experiment using an adult mouse testis culture model without exposure to gonadotropins or androgens showed that treatment with an osteogenic cocktail for 14 days did not induce *Runx2*, *Bgalp*, or *Alpl* but a focal expression of the RUNX2 (using the antibody targeting exon 5) was found in specimens treated with both vehicle and the osteogenic cocktail, indicating that RUNX2 expression was switched to the bone-specific isoform[54] during *ex vivo* culturing. Interestingly, RUNX2 was more frequently found in specimens treated with the osteogenic cocktail and phosphate.

One of the most potent inhibitors of mineralization is PPi, which can be degraded by alkaline phosphatase or pyrophosphatase[55]. PPi prevented the mineralization induced by calcium and phosphate in the GC1 cell line, which was reintroduced in the presence of pyrophosphatase-degrading PPi. This highlights the role of PPi as an inhibitor of spermatogonial mineralization. Alkaline phosphatases convert PPi to Pi and are expressed by primordial and malignant germ cells[13,27]. Four genes encode alkaline phosphatases in humans (five in mice), and all are expressed in malignant germ cells, including the tissue non-specific alkaline phosphatase (TNAP)[13]. TNAP is required for normal bone formation and regulates vascular matrix mineralization by inactivation of the mineralization inhibitors PPi and OPN[56]. We show here that TCam2 cells have high alkaline phosphatase activity in alkaline pH, suggesting that mineralization is promoted by malignant cells. This implies that men with GCNIS and TGCTs are prone to microcalcifications due to lower PPi levels caused by increased alkaline phosphatase activity. Several other inhibitors and promoters of mineralization could be involved, but the marked epididymal mineralization observed in *Fgf23*[-/-] mice, despite the presence of SIBLING proteins that inhibit hydroxyapatite formation and the extensive testicular microcalcifications in men with loss-of-function variants in *SLC34A2*[22], suggest that a change of just one component can be enough to promote the formation of microcalcifications. Bone formation in the testis is commonly observed in teratomas but can occur without malignancy. Mature bone tissue formation has also been reported in the testis after trauma or bleeding without signs of GCNIS or invasive TGCTs[57]. Here, we also show intratesticular bone in a specimen despite no detection of GCNIS or malignant cells. A comprehensive analysis of germ cells adjacent to the bone tissue revealed focally bone-specific RUNX2 positive germ cells but none of them expressed the other investigated bone markers or deposition of hydroxyapatite, which indicates that the mature

bone may have been formed in the interstitial compartment through a different mechanism related to the trauma rather
than alterations in mineral homeostasis leading to testicular microcalcifications.

In conclusion, we show that the formation of testicular microcalcifications occurs as a result of both malignant and benign
etiology, which includes one or more of the following events: aberrant function of Sertoli cells, disturbed local phosphate
homeostasis, change in mineralization inhibitors, and aberrant germ cell function. These changes alone or in combination
can drive osteogenic-like differentiation of germ cells resulting in intratubular deposition of microcalcifications consisting
of hydroxyapatite. Thus, this study suggests that microcalcifications alone should not be considered a marker for
malignancy in the testis.

Materials and Methods

Human tissue samples

Patients were recruited from the Department of Growth and Reproduction, Rigshospitalet, Denmark in accordance with the Helsinki Declaration after approval from the local ethics committee (Permit No. H-15000931, H-17004362, KF 01 2006-3472). Adult testis samples were obtained from orchiectomy specimens performed due to TGCTs. The tissue surrounding the tumor contained tubules with either GCNIS cells or normal/impaired spermatogenesis. Each sample was divided into fragments, which were either snap-frozen and stored at -80°C for RNA extraction or fixed overnight at 4°C in formalin or paraformaldehyde and were subsequently embedded in paraffin. An experienced pathologist evaluated all samples and IHC markers were used to examine the histological subtypes of the TGCT. Fetal gonads were collected after elective terminations for non-medical reasons and gestational age was estimated by ultrasound at the Department of Gynecology (Rigshospitalet/Hvidovre Hospital, Denmark) after approval by the local ethics committee (H-1-2012-007).

Microarray

For this study, the data were extracted from microarray analysis, which was performed and published in full previously[58]. Briefly, snap-frozen tissues were fixed in 75% ethanol and analyzed for AP-2γ (sc-12762, Santa Cruz) by IHC to identify gonocytes and GCNIS and a slide was stained for alkaline phosphatase activity detectable in GCNIS and fetal germ cells. Before microdissection, slides were stained with nitroblue tetrazolium (NBT)-5-bromo-4-chloro-3-indolyl phosphate (BCIP). RNA was purified using Ambion RNAqueous micro kit (Applied Biosystem/Ambion). The quality of RNA was evaluated using Bioanalyzer Picokit (Agilent Technologies). RNA samples were amplified in two rounds using the MessageAMP™ II Kit (Applied Biosystem/Ambion). RNA was reverse transcribed using 50 ng/µL random hexamer primers. Analyses were conducted on three hESCs, microdissected fetal gonocytes, microdissected GCNIS, and tissue containing different percentages of GCNIS and normal testis samples. Agilent Whole Human Genome Microarray 4x44K chips, Design Number 014850 (Agilent Technologies) were used for arrays. Hybridization and scanning of one-color arrays were performed as described by the manufacturer and analyzed using Agilent Feature Extraction software (version 9.1.3.1).

Quantitative RT-PCR (qRT-PCR)

RNA was extracted from 13 tissue specimens with classical seminoma, 10 ECs where some included components of teratoma, and 30 GCNIS samples (GCNIS adjacent to EC, seminoma, mixed tumor, and six GCNIS samples without an overt tumor). RNA from normal testis was purified from three orchiectomy specimens with seminiferous tubules containing varying degrees of full and partial spermatogenesis but with no GCNIS or tumor. Two normal testis RNA samples were purchased from different companies (Applied Biosystems/Ambion and Clontech). RNA and cDNA preparation and qRT-PCR were conducted as described previously[59]. qRT-PCR was performed using primers listed in Suppl. Table 3. The representative bands from each primer combination were sequenced for verification (Eurofins MWG GmbH). qRT-PCR analyses were performed twice in duplicates on two different plates using a Stratagene Mx300P cycler with SYBR Green QPCR Master Mix (Stratagene). Changes in gene expression were determined with the comparative CT method using

376 glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) (ALP expression in human normal testis, GCNIS with microlithiasis,
377 and seminoma) or *β2-microglobulin* (*β2M*) (all other gene expression analyses) as control genes. One sample in the
378 *hpg*.ARKO group was lost during analyses, and *Slc34a1* was therefore not determined in that mouse.

Western blot

Frozen tissues prepared for western blot included adult normal testis, testis with GCNIS, homogeneous anaplastic
seminoma, EC samples, and cell lysates from NTera2 and TCam2 cells. The tissues were homogenized in RIPA lysis buffer
with protease inhibitor cocktail set lll (Calbiochem). After centrifugation, supernatants were collected and diluted in SDS
Sample buffer and heated for 5 min at 95˚C. Proteins were separated on an 8-16% gradient gel (Pierce) and transferred to
a PVDF membrane (Bio-Rad). The membrane was blocked in 5% milk for 1 h and incubated overnight with primary antibody
(diluted 1:200, #sc-16849). ß-actin was used as loading control (diluted 1:200, #sc-47778). The membranes were further
incubated for 1 h with an HRP-conjugated secondary antibody. Membranes were washed in TBST, and protein signals were
detected using 1-step NBT/BCIP (Pierce).

Immunohistochemistry

IHC staining was performed as described previously[60]. Briefly, antigen retrieval was conducted by microwaving for 15 min
in TEG or CIT buffer before incubation with 2% non-immune goat serum (Zymed Histostain kit) or 0.5% milk powder diluted
in TBS. Subsequently, slides were incubated with primary antibodies overnight at 4˚C (Suppl. Table 2) followed by a
secondary antibody, and a peroxidase-conjugated streptavidin complex (Zymed Histostain kit) was used as a tertiary layer.
Visualization was performed with amino ethyl carbazole. Between incubation steps, the slides were washed with TBS.
Counterstaining was performed with hematoxylin eosin (HE)-staining. All experiments were performed with control
staining without primary antibody. Staining was classified according to an arbitrary semiquantitative reference scale
depending on the intensity of staining and the proportion of cells stained: +++, strong staining in nearly all cells; ++,
moderate staining in most of the cells; +, weak staining or a low percentage of cells stained; +/-, very weak staining in
single cells; none, no positive cells detected.

NTera2 xenografts and treatments

All animal experiments were performed in compliance with the Danish Animal Experiments Inspectorate (license number
2011/561-1956) and conducted as previously described[30]. Mice were housed and interventions were conducted at
Pipeline Biotech in a pathogen-free area. NTera2 cells were grown under standard conditions in 175 cm^2 flasks and media
were changed every 48 hours. Matrigel (BD Biosciences) was diluted in Dulbecco's modified Eagle's medium (DMEM)
before being mixed 1:1 with NTera2 cells. NTera2 cells (5x10^6 cells) were injected subcutaneously into the flanks of male
nude mice 6-8 weeks-old NMRI mice (Fox1^{nu}, Taconic Europe). The NTera2 cells were grown on the mice for 28-56 days.
Tumor volumes were calculated from two tumor diameter measurements using a Vernier caliper: tumor volume = L x W
x ½W. If a tumor diameter of 12 mm was reached the animals were sacrificed. At the end of the study, mice were

euthanized according to the animal welfare euthanasia statement. Tumors, kidneys, and testes were harvested and

weighed, and one half was fixed in formalin and embedded in paraffin for IHC, and the other half was snap-frozen.

Hypogonadal mice with androgen receptor ablation globally or specifically in the Sertoli cells and Sertoli cell-ablated

mice

The *hpg* mice were bred in Glasgow and all procedures were carried out under UK Home Office License and with the

approval of a local ethical review committee. *hpg*.SCARKO and *hpg*.ARKO mice were generated as previously described[33,61].

SCARKO and ARKO mice were generated by crossing mice carrying floxed *Ar* (*Ar*[fl]) with mice expressing Cre regulated by

the Sertoli-specific promoter of AMH (*Amh-Cre*) or the ubiquitous promoter PGK-1. *hpg*.SCARKO mice were generated by

crossing female *Ar*[fl/fl] mice heterozygous for the GnRH deletion (*hpg*/+) (C3HE/HeH-101/H) with male *hpg*/+ mice

expressing *Amh-Cre*. The *hpg*.ARKO was generated the same way using a *Pgk1-Cre* instead of *Amh-Cre*. The mice were

euthanized at 9-11 weeks of age by cervical dislocation. Sertoli cell ablated mice were generated as previously described[50].

Briefly, *Amh-Cre* mice were bred with diphtheria-toxin receptor (iDTR) mice, and the offspring were injected

subcutaneously with a single acute dose (100 ng in 50 μl) of diphtheria-toxin or with 50 μl sterile water (vehicle) at

postnatal day 18. The mice were euthanized on postnatal day 80.

Fgf23[-/-] **mice**

Fgf23[-/-] *mice* were generated as previously described[62]. Mice were bred in Boston and all studies were approved by the

Institutional Animal Care and Use Committee of Harvard. The mice were euthanized at 8 weeks of age by using carbon

dioxide followed by cervical dislocation.

Biochemical Analyses

Human iFGF23 and total FGF23 (iFGF23 + cFGF23) were measured in duplicates in batched assays using assays from

Immutopics (Immutopics, #60-6100, #60-6600) according to the manufacturer's instructions. The seminal fluid FGF23

measurements in healthy men have been published previously[42]. Total serum calcium and phosphorus levels were

determined using Stanbio LiquiColor Kits (Stanbio Laboratory).

In vitro **cultures**

hESC lines were cultured in Sheffield as described previously[63]. H7 cells with or without genomic aberrations were sorted

using a fluorescence-activated cell sorter (FACS) and only cells expressing the pluripotency marker SSEA3

(undifferentiated cells) were analyzed. GC1 cells (ATCC® CRL-2053™) were grown under standard conditions at 37˚C at

5% CO_2 in DMEM supplemented with penicillin (100 U/ml) and streptomycin (100 mg/ml) (Gibco), 10% FBS (Gibco,

#11573397), and L-glutamine (Gibco, #25030-024) (2 mM final conc.) was also added to the media. To investigate

calcification in the GC1 cells, the cells were grown in 24-well plates and treated with increasing concentrations of $CaCl_2$

and $NaH_2PO_4 \cdot Na_2HPO_4$ alone or in combination with PPi (Merck, #P8010) and/or phosphatase (Prospec, #ENZ-241). The

medium was changed every second day. Human and mouse testes were cultured *ex vivo* using a hanging drop culture

approach[64] or culture on agarose gel pieces, respectively. In human testis specimens, the effects of 50 ng/ml iFGF23 or

200 ng/ml cFGF23 were investigated after being added to culture media with 0.1% BSA for 24 h. In the mouse *ex vivo*

cultures, testis from adult 8 weeks old mice were dissected and treated with osteogenic medium in a gel-based tissue

model. Briefly, Agarose gels (1.5%) were cut into 8x8x8 mm^3 cube, placed in 4-well plates and soaked in culture media

(DMEM-F12, 1X penicillin/streptomycin, 1X insulin, transferrin and selenium supplement (ITSS), 10% fetal bovine serum

(FBS)) for a minimum of 24 hours before setup of tissue cultures. Mouse testis was dissected into 1.5 mm^3 pieces and

placed in opposite corners on each gel. The tissues were cultured for 14 days at 34 °C and 5% CO_2 in 325 µl media with

osteogenic medium containing 50 µg/ml Ascorbic acid (Merck, #A4544), 10 mM β-glycerol phosphate (Merck, #G9422),

and either -/+ 2-, 4-, or 8-mM phosphate or vehicle. The medium was changed every second day. At least three replicate

tissue pieces from the same mouse and treatment group were fixed in formalin followed by paraffin embedding for IHC

and histological analyses. Tissue was evaluated for morphology and technical quality based on HE-staining.

von Kossa, alizarin red, BCIP/NBT, and fast blue RR staining

For von Kossa staining of tissue sections, the paraffin sections were first deparaffinized and rinsed in ddH$_2$O. Sections were

then incubated with 1% silver nitrate solution under ultraviolet light for 20 min. Thereafter, the sections were rinsed in

ddH$_2$O and incubated with 5% sodium thiosulfate for 5 min and rinsed in ddH$_2$O (and sometimes counterstained with

Mayer's hematoxylin) and dehydrated through graded alcohol and cleared in xylene. For alizarin red staining of tissue

sections, deparaffinized sections were stained with 2% alizarin red solution (pH 4.1-4.3) for 5 minutes. The sections were

then dehydrated in acetone and acetone-xylene (1:1) and cleared in xylene. For alizarin red staining of GC1 cells, the cells

were washed in PBS and fixed in 10% formalin for 10 min. Thereafter, the cells were washed in ddH$_2$O and incubated with

0.5% alizarin red (pH 4.1-4.3) for 20 min protected from light with light shaking. Cells were washed 2-3 times with ddH$_2$O

and left to dry. Cryosections (10 µm) were treated with BCIP/NBT mixture for 90 sec. The BCIP/NBT mixture consisted of

45 µl BCIP stock solution and 35 µl NBT stock solutions in 10 ml water. Stock solutions: 50 mg/ml 5-bromo-4-chloro-3-

indolyl phosphate (BCIP, Sigma #B8503) in 100% dimethylformamide (Sigma #D4254) and 75 mg/ml nitroblue tetrazolium

(NBT, Sigma #N6876) in 70% v/v dimethylformamide/distilled water. The reaction was stopped in water (1 min). BCIP/NBT

staining of cell lines was performed the same way but in citrate buffer (pH=7) or relevation buffer (pH=9.2-9.5). For fast

blue RR staining, the cells were washed in PBS and then fixed with 3.7% formalin for 10 min. Then, the cells were rinsed

in PBS and stained for 15 min at RT lightly shaking and protected from light. The cells were stained for alkaline phosphatase

activity with Napthol AS-MX, 144 n,n-dimethylformamide and Fast Blue RR salt (Sigma #F0500).

Statistics

Data were analyzed using GraphPad Prism v. 8 (GraphPad Software). Gaussian distribution of all numerical variables was

evaluated by QQ plots. Unpaired Student's t-test was used between two groups of mice. One-way ANOVA followed by

Dunnett's multiple comparisons test was used to test for differences between gene expression in human normal testis

and tumors, *ex vivo* cultures of human and mouse testes, and between WT and the different *hpg* mouse models. Kruskal-

Wallis test with Dunn's multiple comparisons test was used to test for differences between levels of iFGF23 and total
FGF23 (iFGF23 + cFGF23) in seminal fluid from healthy men and tumor patients. For correlation analyses, residual plots
were evaluated to ensure the validity of the correlation analyses. As a result, linear regression analyses were used to
investigate the link between *FGF23* and the pluripotency genes *OCT4* and *NANOG* in EC, and between tumor size and
Slc34a1 expression in kidney and testis of xenografted mice. Statistical significance was determined at the following levels:
not significant (ns) $p>0.05$, *$p<0.05$, **$p<0.01$, *** $p<0.001$, and **** $p<0.0001$.

Acknowledgments

566 We thank Giulio Spagnoli for the MAGE antibody. We gratefully acknowledge several urologists and pathologists of the
567 Greater Copenhagen area hospitals for their help with collecting the tissue samples. We thank Betina F. Nielsen, Bonnie
568 Håkansson, Ana R. Nielsen, and Brian V. Hansen for their skillful technical assistance. We appreciate the efforts of Carsten
569 L. Buus and Klaus Kristensen from Pipeline Biotech, who performed the xenografting. We also thank all the patients and
570 donors who took part in this study.

Funding

572 Danish Cancer Society, Forskningsrådet for sundhed og sygdom, Novo Nordisk Foundation, Aase og Ejnar Danielsens fond,
573 Hørslev Fonden, Dagmar Wilhelms fond, and Ib Henriksens fond.

References

1. Tan, I. B. *et al.* Testicular microlithiasis predicts concurrent testicular germ cell tumors and intratubular germ cell neoplasia of unclassified type in adults. *Cancer* **116**, 4520–4532 (2010).

2. Pedersen, M. R., Rafaelsen, S. R., Møller, H., Vedsted, P. & Osther, P. J. Testicular microlithiasis and testicular cancer: review of the literature. *Int. Urol. Nephrol.* **48**, 1079–1086 (2016).

3. Leblanc, L. *et al.* Testicular microlithiasis and testicular tumor: a review of the literature. *Basic Clin. Androl.* **28**, 8 (2018).

4. Barbonetti, A. *et al.* Testicular Cancer in Infertile Men With and Without Testicular Microlithiasis: A Systematic Review and Meta-Analysis of Case-Control Studies. *Front. Endocrinol. (Lausanne).* **10**, 1–8 (2019).

5. De Jong, B. W. D. *et al.* Raman spectroscopic analysis identifies testicular microlithiasis as intratubular hydroxyapatite. *J. Urol.* **171**, 92–96 (2004).

6. Smith, G. D., Steele, I., Barnes, R. B. & Levine, L. A. Identification of seminiferous tubule aberrations and a low incidence of testicular microliths associated with the development of azoospermia. *Fertil. Steril.* **72**, 467–471 (1999).

7. Kirsch, T. Determinants of Pathologic Mineralization. *Curr. Opin. Rheumatol.* **18**, 174–180 (2006).

8. Boström, K., Watson, K. E., Stanford, W. P. & Demer, L. L. Atherosclerotic calcification: Relation to developmental osteogenesis. *Am. J. Cardiol.* **75**, 88B-91B (1995).

9. Rowe, P. S. N. Regulation of Bone−Renal Mineral and Energy Metabolism: The PHEX, FGF23, DMP1, MEPE ASARM Pathway. *Crit. Rev. Eukaryot. Gene Expr.* **22**, 61–86 (2012).

10. Martin, A. & Quarles, L. D. Evidence for FGF23 Involvement in a Bone-Kidney Axis Regulating Bone Mineralization and Systemic Phosphate and Vitamin D Homeostasis. in *Advances in Experimental Medicine and Biology* (ed. Kuro-o, M.) vol. 728 65–83 (Springer US, 2012).

11. Shanahan, C. M., Crouthamel, M. H., Kapustin, A. & Giachelli, C. M. Arterial Calcification in Chronic Kidney Disease: Key Roles for Calcium and Phosphate. *Circ. Res.* **109**, 697–711 (2011).

12. Hessle, L. *et al.* Tissue-nonspecific alkaline phosphatase and plasma cell membrane glycoprotein-1 are central antagonistic regulators of bone mineralization. *Proc. Natl. Acad. Sci.* **99**, 9445–9449 (2002).

13. Hofmann, M. C., Jeltsch, W., Brecher, J. & Walt, H. Alkaline Phosphatase Isozymes in Human Testicular Germ Cell Tumors, Their Precancerous Stage, and Three Related Cell Lines. *Cancer Res.* **49**, 4696–4700 (1989).

14. Yuan, Q. *et al.* Increased Osteopontin Contributes to Inhibition of Bone Mineralization in FGF23-Deficient Mice. *J. Bone Miner. Res.* **29**, 693–704 (2014).

15. Lieben, L. *et al.* Normocalcemia is maintained in mice under conditions of calcium malabsorption by vitamin D−induced inhibition of bone mineralization. *J. Clin. Invest.* **122**, 1803–1815 (2012).

16. Golub, E. E. Role of matrix vesicles in biomineralization. *Biochim. Biophys. Acta - Gen. Subj.* **1790**, 1592–1598 (2009).

17. Komori, T. *et al.* Targeted Disruption of Cbfa1 Results in a Complete Lack of Bone Formation owing to Maturational Arrest of Osteoblasts. *Cell* **89**, 755–764 (1997).

18. Ducy, P. *et al.* Increased bone formation in osteocalcin-deficient mice. *Nature* **382**, 448–452 (1996).

19. Garringer, H. J. *et al.* Two novel GALNT3 mutations in familial tumoral calcinosis. *Am. J. Med. Genet. Part A* **143A**,
2390–2396 (2007).

20. Campagnoli, M. F. *et al.* Familial tumoral calcinosis and testicular microlithiasis associated with a new mutation of
GALNT3 in a white family. *J. Clin. Pathol.* **59**, 440–442 (2006).

21. Kato, K. *et al.* Polypeptide GalNAc-transferase T3 and Familial Tumoral Calcinosis. *J. Biol. Chem.* **281**, 18370–
18377 (2006).

22. Corut, A. *et al.* Mutations in SLC34A2 cause pulmonary alveolar microlithiasis and are possibly associated with
testicular microlithiasis. *Am. J. Hum. Genet.* **79**, 650–656 (2006).

23. Vanakker, O. M. *et al.* Visceral and testicular calcifications as part of the phenotype in pseudoxanthoma
elasticum: ultrasound findings in Belgian patients and healthy carriers. *Br. J. Radiol.* **79**, 221–225 (2006).

24. Skakkebaek, N. E. Possible carcinoma-in-situ of the testis. *Lancet* **300**, 516–517 (1972).

25. Rajpert-De Meyts, E. *et al.* Developmental expression of POU5F1 (OCT-3/4) in normal and dysgenetic human
gonads. *Hum. Reprod.* **19**, 1338–1344 (2004).

26. Hart, A. H. *et al.* The pluripotency homeobox gene NANOG is expressed in human germ cell tumors. *Cancer* **104**,
2092–2098 (2005).

27. Jacobsen, G. K. & Norgaard-Pedersen, B. Placental alkaline phosphatase in testicular germ cell tumours and in
carcinoma-in-situ of the testis. *Acta Pathol. Microbiol. Immunol. Scand. - Sect. A Pathol.* **92**, 323–329 (1984).

28. Rodriguez, S. *et al.* Expression profile of genes from 12p in testicular germ cell tumors of adolescents and adults
associated with i(12p) and amplification at 12p11.2–p12.1. *Oncogene* **22**, 1880–1891 (2003).

29. Shimada, T. *et al.* FGF-23 Is a Potent Regulator of Vitamin D Metabolism and Phosphate Homeostasis. *J. Bone
Miner. Res.* **19**, 429–435 (2003).

30. Blomberg Jensen, M. *et al.* Vitamin D Metabolism and Effects on Pluripotency Genes and Cell Differentiation in
Testicular Germ Cell Tumors In Vitro and In Vivo. *Neoplasia* **14**, 952–962 (2012).

31. Goetz, R. *et al.* Isolated C-terminal tail of FGF23 alleviates hypophosphatemia by inhibiting FGF23-FGFR-Klotho
complex formation. *Proc. Natl. Acad. Sci.* **107**, 407–412 (2010).

32. Farrow, E. G., Imel, E. A. & White, K. E. Hyperphosphatemic familial tumoral calcinosis (FGF23, GALNT3 and
αKlotho). *Best Pract. Res. Clin. Rheumatol.* **25**, 735–747 (2011).

33. O'Shaughnessy, P. J., Monteiro, A., Verhoeven, G., De Gendt, K. & Abel, M. H. Occurrence of testicular
microlithiasis in androgen insensitive hypogonadal mice. *Reprod. Biol. Endocrinol.* **7**, 88 (2009).

34. Kang, J., Rajpert-De Meyts, E., Giwercman, A. & Skakkebæk, N. E. The association of testicular carcinoma 'in situ'
with intratubular microcalcifications. *J. Urol. Pathol.* **2**, 235–242 (1994).

35. Cormier, S., Leroy, C., Delezoide, A.-L. & Silve, C. Expression of fibroblast growth factors 18 and 23 during human
embryonic and fetal development. *Gene Expr. Patterns* **5**, 569–573 (2005).

36. Wei, C. L. *et al.* Transcriptome Profiling of Human and Murine ESCs Identifies Divergent Paths Required to
Maintain the Stem Cell State. *Stem Cells* **23**, 166–185 (2005).

37. Jenkins, A. D., Lechene, C. P. & Howards, S. S. Concentrations of Seven Elements in the Intraluminal Fluids of the
Rat Seminiferous Tubules, Rete Testis, and Epididymis. *Biol. Reprod.* **23**, 981–987 (1980).

38. Jenkins, A. D., Lechene, C. P. & Howards, S. S. The Effect of Estrogen Administration In Vivo on the Elemental
Composition of the Intraluminal Fluids of the Seminiferous Tubules, Rete Testis, and Epididymis of the Rat. *J.*
Androl. **4**, 272–275 (1983).

39. Memon, F. *et al.* Does Fgf23–klotho activity influence vascular and soft tissue calcification through regulating
mineral ion metabolism? *Kidney Int.* **74**, 566–570 (2008).

40. Leszczynska, A. & Murphy, J. M. Vascular Calcification: Is it rather a Stem/Progenitor Cells Driven Phenomenon?
Front. Bioeng. Biotechnol. **6**, 1–8 (2018).

41. Rajpert-De Meyts, E. *et al.* Changes in the profile of simple mucin-type O-glycans and polypeptide GalNAc-
transferases in human testis and testicular neoplasms are associated with germ cell maturation and tumour
differentiation. *Virchows Arch.* **451**, 805–814 (2007).

42. Bøllehuus Hansen, L. *et al.* Influence of FGF23 and Klotho on male reproduction: Systemic vs direct effects. *FASEB*
J. **34**, 12436–12449 (2020).

43. Boisen, I. M. *et al.* Calcium transport in male reproduction is possibly influenced by vitamin D and CaSR. *J.*
Endocrinol. **251**, 207–222 (2021).

44. DeLuca, S. *et al.* Amelioration of the premature ageing-like features of Fgf-23 knockout mice by genetically
restoring the systemic actions of FGF-23. *J. Pathol.* **216**, 345–355 (2008).

45. Rebourcet, D. *et al.* Sertoli Cells Maintain Leydig Cell Number and Peritubular Myoid Cell Activity in the Adult
Mouse Testis. *PLoS One* **9**, e105687 (2014).

46. Sharpe, R. M., McKinnell, C., Kivlin, C. & Fisher, J. S. Proliferation and functional maturation of Sertoli cells, and
their relevance to disorders of testis function in adulthood. *Reproduction* **125**, 769–784 (2003).

47. O'Shaughnessy, P. J., Verhoeven, G., De Gendt, K., Monteiro, A. & Abel, M. H. Direct action through the sertoli
cells is essential for androgen stimulation of spermatogenesis. *Endocrinology* **151**, 2343–2348 (2010).

48. Ronchetti, I., Boraldi, F., Annovi, G., Cianciulli, P. & Quaglino, D. Fibroblast involvement in soft connective tissue
calcification. *Front. Genet.* **4**, 1–16 (2013).

49. Gourgari, E., Saloustros, E. & Stratakis, C. A. Large-cell calcifying Sertoli cell tumors of the testes in pediatrics.
Curr. Opin. Pediatr. **24**, 518–522 (2012).

50. Rebourcet, D. *et al.* Sertoli cells control peritubular myoid cell fate and support adult leydig cell development in
the prepubertal testis. *Dev.* **141**, 2139–2149 (2014).

51. Guan, K. *et al.* Pluripotency of spermatogonial stem cells from adult mouse testis. *Nature* **440**, 1199–1203 (2006).

52. Ogawa, S. *et al.* Cbfa1, an Essential Transcription Factor for Bone Formation, Is Expressed in Testis from the Same
Promoter Used in Bone. *DNA Res.* **7**, 181–185 (2000).

53. Jeong, J. *et al.* Expression of Runx2 transcription factor in non-skeletal tissues, sperm and brain. *J. Cell. Physiol.*
217, 511–517 (2008).

54. Kanto, S., Grynberg, M., Kaneko, Y., Fujita, J. & Satake, M. A variant of Runx2 that differs from the bone isoform
in its splicing is expressed in spermatogenic cells. *PeerJ* **4**, e1862 (2016).

55. Orriss, I. R. Extracellular pyrophosphate: The body's "water softener". *Bone* **134**, (2020).

56. Lomashvili, K. A., Cobbs, S., Hennigar, R. A., Hardcastle, K. I. & O'Neill, W. C. Phosphate-Induced Vascular
Calcification: Role of Pyrophosphate and Osteopontin. *J. Am. Soc. Nephrol.* **15**, 1392–1401 (2004).

57. Yoneda, F., Kagawa, S. & Kurokawa, K. Dystrophic Calcifying Nodule with Osteoid Metaplasia of the Testis. *Br. J.*
Urol. **51**, 413 (1979).

58. Sonne, S. B. *et al.* Analysis of Gene Expression Profiles of Microdissected Cell Populations Indicates that Testicular
Carcinoma In situ Is an Arrested Gonocyte. *Cancer Res.* **69**, 5241–5250 (2009).

59. Blomberg Jensen, M. *et al.* Expression of the vitamin D receptor, 25-hydroxylases, 1α-hydroxylase and 24-
hydroxylase in the human kidney and renal clear cell cancer. *J. Steroid Biochem. Mol. Biol.* **121**, 376–382 (2010).

60. Blomberg Jensen, M. *et al.* Vitamin D receptor and vitamin D metabolizing enzymes are expressed in the human
male reproductive tract. *Hum. Reprod.* **25**, 1303–1311 (2010).

61. O'Shaughnessy, P. J., Monteiro, A., Verhoeven, G., De Gendt, K. & Abel, M. H. Effect of FSH on testicular
morphology and spermatogenesis in gonadotrophin-deficient hypogonadal mice lacking androgen receptors.
Reproduction **139**, 177–184 (2010).

62. Sitara, D. *et al.* Homozygous ablation of fibroblast growth factor-23 results in hyperphosphatemia and impaired
skeletogenesis, and reverses hypophosphatemia in Phex-deficient mice. *Matrix Biol.* **23**, 421–432 (2004).

63. Andrews, P. W., Goodfellow, P. N., Shevinsky, L. H., Bronson, D. L. & Knowles, B. B. Cell-surface antigens of a
clonal human embryonal carcinoma cell line: Morphological and antigenic differentiation in culture. *Int. J. Cancer*
29, 523–531 (1982).

64. Jørgensen, A. *et al.* Hanging drop cultures of human testis and testis cancer samples: a model used to investigate
activin treatment effects in a preserved niche. *Br. J. Cancer* **110**, 2604–14 (2014).

Figure 1

Deposition of hydroxyapatite is linked with the induction of bone markers and changes in the expression of

phosphate transporters. (**A**) Ultrasonic detection of testicular microlithiasis adjacent to a tumor (left) or in an otherwise

healthy testis (right). Arrows indicate testicular microlithiasis. Staining of calcifications by von Kossa and alizarin red in

human testis. (**B**) IHC staining of proteins potentially involved in the formation of testicular microcalcifications adjacent

to microcalcifications in human testis. OCT4: marker of GCNIS cells. Control: without primary antibody. Scale bars

correspond to 125 μm (left) and 25 μm (right). (**C**) Mild (left) and moderate (right) microcalcifications in human testis

stained with von Kossa. Scale bars correspond to 50 μm. (**D**) Severe bone formation in testis from a man with no known

history of local trauma or other predisposing factors of ectopic ossification. IHC staining with RUNX2. Scale bars

correspond to 1.25 mm (upper), 50 μm (lower left), and 100 μm (lower right). (**E**) Expression levels of the bone marker

(*BGLAP*) and phosphate transporters (*SLC34A1* and *SLC34A2*) in NT (normal testis), SCO (Sertoli-cell-only) pattern, and

GCNIS (germ cell neoplasia *in situ*). Data are presented as mean ± SEM. *$p < 0.05$.

Figure 2

Ectopic expression of FGF23 in human testicular germ cell tumors and embryonic stem cells. (**A**) Expression levels of

the pluripotency factors *NANOG* and *OCT4*, and *FGF23* in normal testis (NT), Sertoli-cell-only (SCO) pattern, GCNIS,

seminoma (SEM), and EC +/- teratoma (T). (**B**) Correlations between gene expression of *FGF23* and *OCT4* or *NANOG* in EC

+/- T. (**C**) Microarray showing expression levels of *FGF23* and *KL* in microdissected fetal gonocytes and GCNIS compared

with testicular tissue samples with varying percentages of GCNIS, hESC, and NT. (**D**) Expression level of *FGF23* in NT,

hESC, and NTera2 cells (n=1-5). (**E**) IHC staining of GCNIS with adjacent NT, EC, and SEM. Scale bars correspond to 250

μm (upper panels) and 100 μm (lower panels). (**F**) Western blot of iFGF23 and cFGF23 in NT, GCNIS, anaplastic SEM, EC,

NTera2 cells, and TCam2 cells. (**G**) Levels of seminal iFGF23 or total FGF23 (iFGF23 + cFGF23) in healthy men or men with

GCNIS or EC. Parentheses on the x-axis indicate sample size. Data are presented as mean ± SEM. *$p < 0.05$, **$p < 0.01$,

****$p < 0.0001$. Correlations indicated by R^2.

Figure 3

Change in expression of phosphate transporters and a bone marker in testis of FGF23-producing NTera2 cell tumors in

xenografted mice. (**A**) OCT4 and FGF23 expression in the tumor on the flank of the nude mice. The scale bar

corresponds to 250 μm. (**B**) Serum levels of human iFGF23 in control or NTera2 cell inoculated nude mice (n=5-6). (**C**)

Expression level of the phosphate transporters *Slc34a1* and *Slc34a2* and the bone marker *Bglap* in testis of control or

NTera2 cell xenografted nude mice (n=5-6). (**D**) Correlations between tumor size and expression of *Slc34a1* in testis of

the vehicle and NTera2 cell xenografted nude mice (n=11). (**E**) Effects of iFGF23 (50 ng/ml) or cFGF23 (200 ng/ml) after

24 h in human *ex vivo* cultures of testis tissue without GCNIS. Data are presented as mean ± SEM. *$p < 0.05$, **$p < 0.01$.

Correlation is indicated by R^2.

Figure 4

Mineralization in reproductive organs of *Fgf23* knockout mice. (**A**) Serum levels of calcium and Pi in WT and *Fgf23*[-/-]

mice. (**B**) Table of bone marker expression observed in premeiotic germ cells in WT and *Fgf23*[-/-] mice. (**C**) IHC staining of

bone markers and vitamin D processing enzymes in WT and *Fgf23*[-/-] mice. Scale bar corresponds to 100 μm. (**D**)

Expression level of the calcification-regulators *Abcc6* and *Enpp1*, the phosphate transporters *Slc34a2* and *Slc34a1*, and

the bone marker *Bglap* in testis of mice that received normal or high phosphate diet (n=8-10). (**E**) IHC staining of bone

markers and staining of minerals with von Kossa and alizarin red in epididymis of WT and *Fgf23*[-/-] mice. The scale bar

corresponds to 250 μm. (**F**) Table of bone marker expression observed in the epididymis of WT and *Fgf23*[-/-] mice. Data

are presented as mean ± SEM. *p<0.05, ***p<0.001, and ****p<0.0001.

Figure 5

Testicular microcalcifications in hypogonadal and Sertoli cells ablated mice. (**A**) Schematic representation of the

prevalence of microlithiasis in ARKO, SCARKO, *hpg*, *hpg*.SCARKO, and *hpg*.ARKO mice models. (**B**) Expression levels of

Slc20a1, *Slc34a2*, *Slc34a1*, and *Bglap* in testis of *hpg*, *hpg*.SCARKO, and *hpg*.ARKO mice models (n=2-4). Data are

presented as mean ± SEM. *p<0.05. (**C**) Microcalcifications in Sertoli cell-ablated mice. Mineral stainings with alizarin red

or von Kossa. Scale bars correspond to 500 μm (upper pictures) and 50 μm (lower pictures).

Figure 6

Alkaline phosphatase activity in normal testis and GCNIS and calcium and phosphate induced mineralization *in vitro*.

(**A**) Staining of alkaline phosphatase activity with BCIP/NBT in normal testis (NT) with adjacent tubules containing GCNIS.

Scale bars correspond to 1 mm (left picture) and 100 μm (right pictures). (**B**) Expression levels of the four alkaline

phosphatases in normal testis, GCNIS with microlithiasis, and seminoma. (**C**) Mineralization stained with alizarin red in

GC1 cells treated with increasing concentrations of calcium and Pi for 2, 4, or 7 days. (**D**) Mineralization stained with

alizarin red in GC1 cells with calcium and/or phosphate in the presence of the mineralization-inhibitor pyrophosphate

(PPi) and/or pyrophosphatase (PPA1) that catalyzes the hydrolysis of pyrophosphate (PPi) to inorganic phosphate (Pi) for

4 days. (**E**) Staining of alkaline phosphatase activity with BCIP/NBT in GC1 or TCam2 cells in pH=7 or pH=9.2-9.5. (**F**) Fast

blue RR staining of alkaline phosphatase activity in GC1 cells cultured in control or osteogenic medium for 14 days.

Figure 7

(**A**) Schematic representation of the mouse testis *ex vivo* model. (**B**) Expression levels of *Bgalp*, *Runx2*, and *Alpl* in *ex vivo*

mouse testis tissue culture specimens treated with osteogenic medium -/+ phosphate. (**C**) Table of the prevalence of

focal expression of RUNX2 in mouse testis *ex vivo* cultures with vehicle and osteogenic treatment for 14 days.

Representative RUNX2 staining in the *ex vivo* culture treated with osteogenic media and 8 mM phosphate. Scale bars

correspond to 50 μm (left picture) and 25 μm (right pictures).

Figure 8

Overview of events that can contribute to the formation of testicular microlithiasis. We suggest that the formation of

testicular microcalcifications occurs as a consequence of one or more of the following events: decreased function of the

Sertoli cells, disturbed local phosphate homeostasis, reduction of mineralization inhibitors, or aberrant germ cell

function. These changes alone or in combination can drive osteogenic-like differentiation of germ cells resulting in

intratubular deposition of microcalcifications consisting of hydroxyapatite.

Supplementary table 1

Immunohistochemical expression of selected proteins in human specimens

	n	FGF23	KLOTHO	FGFR1	BGLAP	RUNX2	OPN
Normal testis	7-15	÷	++ to +++	++ to +++	÷	÷	+
Germ cell carcinoma *in situ*	6-18	+++	÷ to +/-	++ to +++	+	+++	+
Hyalinized/calcified testis	5-14	÷ to +	+/- to +	++	+++	+++	+++
Classical seminoma	4-15	÷ to +	÷	++	÷	÷	÷
Anaplastic seminoma	2	++	+/-	++	N.A.	N.A.	N.A.
Embryonal carcinoma	4-13	++ to +++	÷ to +/-	++	+/- to ++	+/- to +	+/- to +
Teratoma	3-6	÷ to +	+/- to ++	++	+/- to ++	+/- to +	+/- to +
Fetal testis	3	÷ to +/-	÷	++	N.A.	N.A.	N.A.

÷   Not expressed
+/-   Barely detectable in few cells
+   Expressed in few cells
++   Markedly expressed in some cells or expressed in most cells
+++   High expression in most cells
N.A.   Not available

Supplementary table 2

List of primary antibodies used for IHC staining.

Antibody	Catalog no.	Manufacturer	Dilution	Buffer
BGLAP	sc-30044	Santa Cruz biotechnology	1:50 / 1:100	CIT
CYP24A1	sc-66851	Santa Cruz biotechnology	1:500	CIT
CYP27B1	sc-67261	Santa Cruz biotechnology	1:100	CIT
DMP1	sc-73633	Santa Cruz biotechnology	1:100	TEG
FGF23	sc-16849	Santa Cruz biotechnology	1:75 / 1:200	TEG
FGFR1	sc-33928	Santa Cruz biotechnology	1:500	TEG
Klotho	sc-22220	Santa Cruz biotechnology	1:100 / 1:200	TEG
MAGE	Gift from Prof. Spagnoli	Non-commercial	1:500	TEG
NPT2a	sc-33928	Santa Cruz biotechnology	1:200	TEG
OCT4	sc-5279	Santa Cruz biotechnology	1:50 / 1:100	TEG
OPN	sc-73631	Santa Cruz biotechnology	1:100	CIT
ALP	M7191	Abcam	1:100	CIT
RUNX2	sc-10758	Santa Cruz biotechnology	1:200 / 1:400	TEG
SOX9	AB5535	Merck Millipore	1:400	TEG

Supplementary table 3

List of primers used for qRT-PCR

Gene	Forward primer	Reverse primer
	Human	
ALPI	TACACGTCCATCCTGTACGG	CTCGCTCTCATTCACGTCTGG
ALPL	ATGGGATGGGTGTCTCCACA	CCACGAAGGGGAACTTGTC
ALPP	CACCTGTGGGACTTGAGGAC	TTTGGCTCTCGACCAGTGTC
ALPP2	GGCATCATCCCAGTTGAGGAG	GCACATGCTTGTCTACACTGTAT
B2M	ATCCAATCCAAATGCGGCATC	AGTATGCCTGCCGTGTGAAC
BGLAP	CACTCCTCGCCCTATTGGC	GTGGTCAGCCAACTCGTCA
FGF23	CACCTGCAGATCCACAAGAA	TAATCACCACAAAGCCAGCA
GAPDH	TCAACGACCACTTTGTCAAGC	GGTGGTCCAGGGGTCTTACTC
NANOG	TGATTTGTGGGCCTGAAGAAAA	GAGGCATCTCAGCAGAAGACA
OCT4	GACTCCTCGGTCCCTTTCC	CAAAAACCCTGGCACAAACT
SLC34A1	CCCTCAGGTCCTACACAGGAT	GGAGCAGACGAAGAGGTAGAG
SLC34A2	CTGAGGCACCTGTAACCAAGA	GGAGCACACGAAAAAGTAGAGA
	Mouse	
Abcc6	GCATCGTTCAGGCTCGAGTG	CGAGGAGCGCCTGGAGTTAC
Alpl	CACGGCCATCCTATATGGTAA	GGGCCTGGTAGTTGTTGTGA
B2m	CGAGACATGTGATCAAGCATCA	TATTGCTCAGCTATCTAGGATAT
Bglap	CTGACCTCACAGATCCCAAGC	TGGTCTGATAGCTCGTCACAAG
Cyp24a1	ACACTGGCAGAGTACCACAAG	TCCAGGGTTTGATCTCTAGCC
Cyp27b1	GCACAGTTTACGTTGCCGAC	CGTTAGCAATCCGCAAGCA
Enpp1	AAGCGCTTACACTTCGCTAAAAG	TGATGGATTCAACGCAAGTTG
Runx2	AGAGTCAGATTACAGATCCCAGG	TGGCTCTTCTTACTGAGAGAGG
Slc20a1	TCATCATTGCATTTGTCTTGGCA	CGGATGGTTTCGCTCACTTTG
Slc34a1	GACAATGCCATCCTATCCAACC	GAGCCCATGATGATCGGAATG
Slc34a2	GACTGAACTGCTACCCTCCTA	ACCATTTGATCCCAGTGTCCT

Supplementary figure 1

Deposition of hydroxyapatite and bone markers. IHC staining of proteins potentially involved in the formation of testicular microcalcifications adjacent to microcalcifications in human testis. Scale bar corresponds to 100 µm.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

A

Serum calcium (mg/dl) — WT, *Fgf23*⁻/⁻ ***

Serum Pi (mg/dl) — WT, *Fgf23*⁻/⁻ ****

B

Premeiotic germ cells

Protein	WT	*Fgf23*⁻/⁻
DMP1	1/4	4/4
OPN	2/5	2/2
BGLAP	1/6	4/4

C

	DMP1	OPN	BGLAP
WT			
Fgf23⁻/⁻			

D

Relative mRNA expression — Normal diet, High phosphate diet

Abcc6, *Enpp1* *, *Slc34a2*, *Slc34a1* ND, *Bglap* ND

E

	DMP1	OPN	BGLAP	von Kossa	Alizarin red
WT					
Fgf23⁻/⁻					

F

Epididymis

Protein	WT	*Fgf23*⁻/⁻
DMP1	0/4	4/5
OPN	0/6	4/5
BGLAP	0/6	4/5

Fig. 5

A

Androgen Receptor KO
ARKO

Androgen Receptor KO in Sertoli cells
SCARKO

No circulating gonadotrophins

hpg

hpg.SCARKO

hpg.ARKO

Microlithiasis: No   No   36%   30%   94%

B

Slc20a1

Slc34a2

Slc34a1

Bglap

C

Alizarin red   von Kossa

Fig. 6

A

NT

GCNIS

B

Relative mRNA expression

ALPP   ALPP2   ALPI   ALPL

☐ Normal testis
☐ GCNIS with microlithiasis
■ Seminoma

C

D2   D4   D7

Pi (mM)
0.9 1.5 2.0 2.5 3.0 4.0

Calcium (mM)
1.8
2.0
2.5
3.0

D

Calcium (3 mM)	-	+	+	+
Pi (4 mM)	-	+	+	+
PPi (2.5 μM)	-	-	+	+
PPA1 (1 μg/ml)	-	-	-	+

E

pH=7   pH=9.2-9.5

GC1 cells

TCam2 cells

F

Control medium

Osteogenic medium

Fig. 7

A

Testes

Pieces ~ 1.5 mm³

B

C

Treatment	RUNX2
Vehicle	2/6
Osteogenic differentiation media	2/2
Osteogenic differentiation media + 2 mM phosphate	0/4
Osteogenic differentiation media + 4 mM phosphate	4/4
Osteogenic differentiation media + 8 mM phosphate	4/4

Fig. 8

Decreased function of support cells
- Androgen receptor mutations
- Large calcifying Sertoli cell tumor
- Testicular dysgenesis syndrome

Disturbed local phosphate homeostasis
- Mutations in *SLC34A2*
- Mutations in *GALNT3*

Testicular microlithiasis

Reduced mineralization inhibition
- Alkaline phosphatase
- Pyrophosphate
- ENPP1
- ABCC6
- OPN/DMP1

Aberrant germ cell function
- Osteogenic-like differentiation
- Gonadoblastoma
- Heat
- Trauma

Supplementary figure 1