Performance in even a simple perceptual task depends on mouse secondary visual areas

Hannah C Goldbach[1*], **Bradley Akitake**[1*], **Caitlin E Leedy**[1], **Mark H Histed**[1]

[1]Unit on Neural Computation and Behavior, National Institute of Mental Health Intramural Program, National Institutes of Health, Bethesda, MD 20892

*For correspondence:
mark.histed@nih.gov (M.H.H.)

*These authors contributed equally to this work

Abstract   Primary visual cortex (V1) in the mouse projects to numerous brain areas, including several secondary visual areas, frontal cortex, and basal ganglia. While it has been demonstrated that optogenetic silencing of V1 strongly impairs visually-guided behavior, it is not known which downstream areas are required for visual behaviors. Here we trained mice to perform a contrast-increment change detection task, for which substantial stimulus information is present in V1. Optogenetic silencing of visual responses in secondary visual areas revealed that their activity is required for even this simple visual task. *In vivo* electrophysiology showed that, although inhibiting secondary visual areas could produce some feedback effects in V1, the principal effect was profound suppression at the location of the optogenetic light. The results show that pathways through secondary visual areas are necessary for even simple visual behaviors.

Introduction

Several kinds of neural computations are required to perform sensorimotor tasks. For example, it has been thought that some cerebral cortical neurons represent aspects of the sensory world in neural activity, and other neurons decode that representation into a form useful for motor action. An important step in understanding brain function is determining where representations are present, and where computations like decoding occur. These questions depend on knowing which areas are involved during sensorimotor behaviors. Here, we ask whether secondary visual areas are involved in a simple visual behavior, or whether such behavior can be performed by operating on ("decoding") sensory representations in V1 or other areas directly, bypassing secondary areas.

The rodent visual system includes several different cortical visual areas, which all respond to visual stimuli, but emphasize different aspects of representation. The borders and locations of mouse visual areas were initially identified through reversals in retinotopy (*Dräger, 1975*; *Wagor et al., 1980*) and cytoarchitecture (*Caviness, 1975*). Later, injections of anterograde tracers into V1 showed that V1 projects to several secondary visual areas, or higher visual areas, of the mouse brain (*Wang and Burkhalter, 2007*). More recently, the boundaries of cortical visual areas in mice have been examined through mapping retinotopy via hemodynamic intrinsic imaging (*Schuett et al., 2002*; *Kalatsky and Stryker, 2003*; *Garrett et al., 2014*; *Juavinett et al., 2017*) and calcium imaging (*Andermann et al., 2011*; *Murakami et al., 2017*; *Zhuang et al., 2017*). Imaging studies have identified as many as twelve individual, retinotopically mapped areas (*Zhuang et al., 2017*) with differing strengths of connectivity to V1. The most heavily studied secondary visual areas, which receive the largest proportions of projections from V1, are the lateral and medial output pathway areas: LM (lateromedial), AL (anterolateral), RL (retrolateral), and PM (posteriomedial) (*Froudarakis et al., 2019*). While more than half of V1 projections target these four secondary visual areas (*Han*

et al., 2018), V1 also projects to a number of other brain areas including frontal cortex and basal ganglia (*Han et al., 2018*; *Khibnik et al., 2014*), both of which are involved in decision-making and action selection.

We sought to determine whether activity in higher visual areas was essential for a simple visual task. Alternatively, it could be that such activity is not always used, and for some tasks, information flows out of V1 via projections that bypass secondary areas (such as V1's projections to frontal cortex or basal ganglia). As an example, consider an analogous question in a larger species. In the primate visual object ("where") pathway, V1 carries representations of small visual edges (*Hubel and Wiesel, 1968*), while areas in inferotemporal (IT) cortex represent complex objects such as faces (*Tsao, 2014*). Identifying the orientation of contours and edges could be carried out by a neural computation that reads out, or decodes, V1 directly, bypassing secondary visual areas such as those in IT cortex. If V1 is read out directly, one might expect that suppression of IT cortical areas without affecting V1 would impact tasks that require face perception, but leave tasks that involve orientation discrimination unaffected. On the other hand, some data (e.g. on unconscious vs. conscious perception, *Leopold, 2012*) supports the idea that macaque V1 is simply a conduit for information to secondary visual areas, with no behaviorally relevant direct readout. In the mouse, we are able to investigate which cortical areas are involved in a given behavior using transgenic lines expressing optogenetic proteins, which allow the inactivation of several different cortical visual areas just by moving an optogenetic light spot. To study how secondary visual areas contribute to mouse perceptual behavior, we chose to study a basic task in which V1 contains substantial information. We trained animals to report changes in stimulus contrast.

Neural activity in V1 carries substantial information about visual stimulus contrast (*Albrecht and Hamilton, 1982*; *Sclar et al., 1990*; *Vaiceliunaite et al., 2013*). Neurons in mouse V1 are tuned for oriented visual stimuli, with a tuning curve that reaches a peak at the cell's preferred orientation. The same neurons simultaneously encode contrast: as stimulus contrast is increased, single V1 neurons increase their firing rates monotonically (*Busse et al., 2011*; *Glickfeld et al., 2013*; *Khastkhodaei et al., 2016*). As expected from single neurons' responses, the V1 neural population response varies monotonically with stimulus contrast, and thus the average or summed V1 response does not completely saturate, even up to 100% contrast (*Glickfeld et al., 2013*; *Histed, 2018*). These features of the population response suggest that mouse V1 activity can be simply (linearly) decoded to allow animals to detect a change in contrast of a visual stimulus. While the structure of population neural variability could in principle make linear decoding difficult, linear decoders that do not take trial-to-trial variability into account perform well on decoding orientation from mouse V1 (*Berens et al., 2012*; *Rumyantsev et al., 2020*; *Stringer et al., 2019*; *Kafashan et al., 2020*). This suggests that linear decoders would also be able to extract stimulus contrast from the V1 population. Thus, a contrast change detection task does not seem to require any information in more complex representations beyond V1. If the mouse brain had circuitry to decode V1 directly, secondary visual areas might not be involved in the task and inhibiting them would not impact animals' behavior.

To study whether secondary visual areas are involved in contrast change detection, mice were trained on a contrast-increment detection task in which they report the presence of a flashed Gabor by releasing a lever. This task was previously found to require V1; that is, suppressing V1 optogenetically produced retinotopically-specific impairments in contrast-detection performance (*Glickfeld et al., 2013*; *Jin and Glickfeld, 2020*). Through optogenetically stimulating all inhibitory neural classes (via the *VGAT-ChR2-EFYP* transgenic mouse line; *Zhao et al., 2011*) we first confirm this finding: inhibition of V1 negatively affects performance in this task. We then inhibit secondary visual areas, and find that suppression of lateral areas (e.g. LM/AL, and RL) and medial areas (PM) also negatively affect performance. Via electrophysiology, we find that inhibition of secondary visual areas primarily acts to suppress secondary area visual responses, rather than suppressing V1 responses via feedback. Thus, these data show that secondary visual areas are involved in a task that, in principle, could have been performed by the mouse reading out V1 directly. We also

find that the effect size caused by secondary area suppression can be large, as suppressing lateral
visual areas (LM/AL and RL) produces behavioral effects of similar or even higher magnitude than
V1 at the same optogenetic light intensity. In contrast, suppression of PM, the largest medial area,
only weakly affected the animals' performance. Paralleling these behavioral findings, any feedback
effects on V1 neural activity were negligible when PM was suppressed, but could be seen in LM at
high light intensity, arguing that LM and V1 activity is linked to some extent. Finally, the behav-
ioral effects in all areas are primarily due to sensitivity (d') and not due to changes in false alarm
rates. In sum, lateral secondary visual areas (LM/AL and RL) form an interconnected network with
V1 that contributes to performance in even simple visually-guided behaviors, where substantial
information is available in, and linearly decodable from, V1.

Results

We studied mice of both sexes expressing ChR2 in all inhibitory neurons (*VGAT-ChR2-EYFP* mouse
line, *Zhao et al., 2011*). We implanted a cranial window and identified visual cortical areas using
hemodynamic intrinsic imaging (Materials and Methods, *Figure 1*, *Figure 1–Figure Supplement 1*).
We cemented a fiber-optic cannula over the cranial window to create a light spot targeted to the
area of visual cortex we wished to inhibit, and trained these animals on a contrast-increment
change detection task (Materials and Methods, *Figure 1A*). Behaving animals were head-fixed in
front of an LED monitor and trained to release a lever in response to the onset (increase in con-
trast, from a neutral gray background) of a Gabor patch for a liquid reward. We varied the contrast
change magnitude in order to obtain psychometric curves, which measure the contrast needed for
animals to perform the task at a set percentage correct — their perceptual threshold (*e.g. Figure 1D*,
open circles; below we discuss effects also on perceptual sensitivity, or d').

Once mice were trained to threshold, we introduced a pulsed blue LED train (470 nm; on for
200 ms, off for 1000 ms) throughout each trial. On 50% of trials (ON trials), the LED pulse occurred
with the visual stimulus. (Due to latency between visual stimulus onset and first spikes in the
cortex, we turned the light pulse on 55 ms after visual stimulus onset; 8-12 ms before the visual
response occurred in V1 *Figure 5–Figure Supplement 1*.) On the remaining trials (OFF trials), the
visual stimulus appeared while the light was off (*Figure 1A*, 820 ms after the last light pulse was
turned off).

To limit the information the animals could gain about correct response (*i.e.* that they could
gain about visual stimulus onset time) from the optogenetic light pulses, the light pulse train was
present on each trial, and the pulse train phase was randomized by varying the first pulse onset
time after the trial began (Materials and Methods). On each trial, the visual stimulus onset time
was chosen randomly (according to a geometric distribution), the light pulse train phase offset
was chosen randomly (uniform distribution), and then the visual stimulus onset time was adjusted
downward to be during the previous light pulse (ON trials) or after the previous light pulse offset
(OFF trials). Thus, each trial, whether ON or OFF, contained a similar pulse train, and we randomly
intermixed ON and OFF trials. These factors made it difficult for animals to use information about
the light pulses to identify the type of trial and change their strategy (such as their perceptual cri-
terion) from trial to trial. While some small amount of information about the visual stimulus time
might in theory be available, because the visual stimulus was locked to discrete times relative to
pulse onset or offset (Materials and Methods), we observed no evidence animals used light pulse
information to guide responses. We found (details below): (1) the optogenetic stimulation system-
atically decreased, not increased, animals' performance, (2) control-site stimulation did not change
performance, and (3) sensitivity (d') analyses yielded qualitatively the same results as hit-rate mea-
sures. Therefore, this behavioral change detection paradigm allows us to examine effects on visual
perception due to inhibition of different mouse visual areas. Finally, while cognitive factors like at-
tention and motivation can change over experimental sessions from one day to the next, many
such factors should act similarly on both trial types. Therefore, to control for changes in attention,
motivation, or perceptual criterion choice (*Macmillan and Creelman, 2004*) from day to day, we

made our comparisons always between ON and OFF trials within each daily experimental session
(e.g. gray vs blue curves, *Figure 1C,F*).

V1 inhibition confirmed to impair behavior

We first sought to validate our methods in primary visual cortex. Based on previous work (*Glick-*
feld et al., 2013), we expected that inhibition of V1 during a contrast-increment change detection
task would impair the animals' behavior. Indeed, we found that inhibiting V1 during behavior,
via a retinotopically-aligned spot of light over V1, produced an intensity-dependent impairment
in behavior (increase in psychometric threshold; *Figure 1C*). In order to obtain a measure of how
effective the optogenetic inhibition was at impairing behavior, for each stimulation light spot we fit
a piecewise-linear function to the threshold increases as a function of optogenetic light intensity.
This yielded an estimate of the intensity required to produce a 100% increase in threshold (yielded
the 100% crossing point; *Figure 1D*, *Figure 1–Figure Supplement 2*), and allowed us to compare the
behavioral impact of inhibition across light spots *Figure 1I-K*. False alarm (early response) rates
were only changed a small amount by stimulation (for this V1 spot: mean ± s.d. early rate increase
(OFF - ON): +7.4% ± 3.7%; overall in V1: +7.7 ± 4.7%.) Note that false alarm rates vary between
light ON and OFF trials, but cannot vary across contrasts, because by definition false alarms must
occur before stimulus presentation — before any stimulus is accessible to perception (Materials
and Methods). Since false alarm rates changed only a small amount, we expected that effects mea-
sured with sensitivity (d') would be similar to those measured with hit rates. To test this, we also
computed d' thresholds in an analogous manner to hit rate thresholds. We fit a Naka-Rushton
function to d' values at each contrast level (Materials and Methods, *Herrmann et al., 2012*), and
did so for light ON and light OFF trials to find two thresholds for d'. Sensitivity threshold changes
were strongly correlated with hit rate threshold changes (*Figure 1E*). We also show below that the
pattern of effects with inhibition of different visual areas are similar for hit rate and for sensitivity.
Because early rate changes were not large, and thresholds measured with sensitivity and with hit
rate were related (for all spots in V1, Mann-Whitney $U = 21.5$, $N_1 = N_2 = 8$, $p = 0.15$ two-tailed), here
and below we primarily calculate threshold changes on hit rate.

Inhibiting control areas, away from visual areas, would be expected to produce little effect in
this task. Indeed, inhibiting control areas in the same manner as we inhibited V1 produced no
effect on threshold (*Figure 1F*). While the light OFF threshold in panel F varies from the light OFF
threshold in panel C, this was not a major effect across our data: light OFF threshold for all V1 sites
was mean 12.5%, 95% CI 11.3-13.6%; for all control sites, mean 9.8%, 95% CI 8.7-11%, and there
was no systematic relationship between threshold change and light OFF threshold: across all V1
sites, slope of linear regression $t = -1.2$, N = 8, $p = 0.23$. To determine the minimum intensity that
would be necessary to produce a 100% increase in threshold (*Figure 1G*), we fit the same piecewise-
linear function (*Figure 1–Figure Supplement 2* for slope calculations) to this data across all collected
intensities. For these control spots, the data put a lower bound on the intensity required to produce
a behavioral effect, but the upper confidence interval for these plots extends upward indefinitely,
as we found no intensity high enough for these spots to cause strong behavioral effects and thus
no upper bound. As with V1 inhibition, changes in false alarm rates were low (summarized in
Figure 4–Figure Supplement 1), and thus, with little hit rate or false alarm rate change, sensitivity
(d') changes were also low (*Figure 1H*).

To measure the spatial falloff of optogenetic inhibition, we used electrophysiological recording
combined with optogenetic stimulation *in vivo* (*Figure 1L*). For inhibition light spots with a half-max
radius of 0.36 ± 0.07 mm (N = 5 ephys. spots; behavioral spot sizes 0.42 ± 0.13 mm, N = 30), we
found that the half max radius (via Gaussian fit, *Figure 1–Figure Supplement 3*) of visual response
suppression was 0.56 mm (bootstrap 95% CI 0.54-0.59%), confirming that inhibitory effects are
spatially limited by the edge of the LED spot. Note that here we measure the spatial falloff of sup-
pression of visual responses. Cortical baseline (spontaneous) firing rates are more sensitive to op-
togenetic manipulation than evoked visual firing rate changes (*Glickfeld et al., 2013*; *Histed, 2018*),

Figure 1. Optogenetic inhibition of V1 confirmed to affect behavior.

Figure 1. (from previous page): (A) Schematic of the visual detection behavior task. Animals' task was to release a lever when they detected a change in a visual stimulus. Stimuli were Gabor patches which increased in contrast, appearing on a neutral gray screen. Cortical activity was suppressed via optogenetic activation of inhibitory neurons (*VGAT-ChR2-EYFP* mouse line). On each trial, a train of light pulses was applied to the cortex (Materials and Methods). Visual stimuli were delivered either during a light pulse (light ON trials) or when the light pulse was off (light OFF trials). **(B)** Cortical areas were identified using hemodynamic intrinsic imaging of cortical responses to a set of 6 visual stimuli in different positions in visual space (Materials and Methods, *Figure 1–Figure Supplement 1*) **(C)** LED stimulation in V1 decreases animal's performance (single session; rightward shift indicates decreased performance). Filled circles: performance at one contrast. Open circles: threshold (62% point of fit curve). **(D)** Piecewise-linear function fit to all sessions from the spot in panel C, measuring the intensity required to double the psychometric threshold (100% crossing point, pink; mean 0.53, bootstrap 95% CI 0.40-0.66 mW/mm^2). Slopes were fit by pooling across stimulation locations and experimental days; fitting slope separately for each day produced similar results *Figure 1–Figure Supplement 2*. **(E)** Change in d' threshold increase vs change in hit-rate threshold increase for the spot in panel C (Materials and Methods, *Figure 4–Figure Supplement 1*, *Figure 1–Figure Supplement 2*). **(F)** Single day example behavior from LED stimulation in a control area outside of V1, same conventions as panel C. **(G)** Same as panel D, fit to sessions from control spot in panel F **(H)** Same as panel E, fit to sessions from the control spot in panel F; mean 100% crossing point 6.6, bootstrap 95% CI 4.6-8.1 mW/mm^2. **(I)** 100% crossing points for all V1 spots and two control spots, as determined by regression fits in E & G. Ordering of spots is arbitrary; points are offset along x axis to allow display of confidence intervals for each point. **(J)** Contours of all V1 spots and controls, color weighted by 100% crossing point. **(K)** Heatmap of effect size in V1, generated by averaging the 100% crossing point at each pixel. **(L)** Spatial fall-off of inhibition. Full-width at half max of normalized suppression = 0.56 mm (red line, 95% CI 0.54-0.59 mm). These measurements were made at 0.3 mW/mm^2, near the mean of V1 crossing points (panel I). Average behavioral spot radius is 0.42 ± 0.13 mm (black dotted line, mean ± SEM, N = 30). Average electrophysiology spot radius is 0.36 ± 0.07 mm (blue shaded area and dotted line, mean ± SEM, N = 5).

Figure 1–Figure supplement 1. Visual area identification via imaging.
Figure 1–Figure supplement 2. Slope determination for piecewise-linear function.
Figure 1–Figure supplement 3. Spatial fall-off of inhibition is restricted to illuminated areas.

---

and consistent with this, we found that visual response changes are more restricted in spatial ex-
tent than spontaneous (baseline) response change *Figure 1–Figure Supplement 3*. (The restricted
spatial extent of suppression is also supported by the falloff of behavioral effects across the cortex;
Discussion.) To make these measurements of spatial falloff of visual responses, we positioned an
array of four electrodes so three electrodes were in V1 and the last electrode shank was in PM,
which was targeted with optogenetic light. We then plotted the visual response suppression in
units normalized to the level of visual response with no inhibition at that electrode. As discussed
below, we chose this configuration because there is little feedback effect on V1 of stimulating PM,
and, in fact, at this light intensity, little feedback effect of stimulating any secondary visual area.

Our V1 results were consistent across animals and across V1 spots (mean 100% crossing pt
0.46, bootstrap 95% CI 0.33-0.58 mW/mm^2; *Figure 1I*). In total, we tested eight spots within V1,
all of which produced substantially larger effects (*i.e.* lower 100% crossing points) than inhibition
of control spots (N = 4, mean control spot 100% crossing point 5.1, & bootstrap 95% CI 2.3-8.0
mW/mm^2; Mann-Whitney $U = 0.0$, V1 N = 8, control N = 4, $p = 0.0042$ two-tailed) (*Figure 1L*). Notably,
inhibition of an area (control spot 1) placed within V1 but offset from the retinotopic location of the
visual stimulus produced no effect (*Figure 1L*, control spot 1). To visualize the effects of inhibiting
V1 across all our stimulation locations (N = 10; N = 8 V1, N = 2 control), we averaged the 100%
crossing points at each pixel within V1 to generate a heat map (*Figure 1M*). Because this heatmap
representation is influenced by the locations of the optogenetic light spots, it is not intended to be
quantitative, but instead to provide a visual guide to assess the spatial pattern of effects given the
spot locations used (all shown in *Figure 1L* and below, to allow comparison between visual areas.)

Together, this data confirms that inhibiting mouse V1 negatively affects an animal's ability to
perform a simple contrast-increment change detection.

Inhibiting areas lateral to V1 (LM/AL and RL) degrades contrast change detection

218 We next inhibited secondary visual areas, beginning with the areas LM, AL, and RL, which are lat-
219 eral to V1. Our main goal was to determine whether any of the lateral visual areas could impact
220 behavior in this task. Therefore, from here on, we refer to LM and AL together as LM/AL, as our
221 optogenetic light spots could not easily differentiate between the two. AL is small relative to our

Figure 2. Inhibiting lateral areas degrades contrast change detection behavior. **(A)** All sessions from an example spot on the border of AL and LM, showing a large effect (mean 100% crossing pt 0.22, bootstrap 95% CI 0.15-0.31 mW/mm^2). **(B)** All sessions from an example spot within RL, showing a large effect (mean 100% crossing pt 0.33, bootstrap 95% CI 0.23-0.40 mW/mm^2). **(C)** All sessions from a control area between LM and V1, showing no effect even at high intensities (mean 100% crossing pt 9.3, bootstrap 95% CI 3.3-10 mW/mm^2). **(D)** Map of spots within LM/AL/RL and control locations, colored by 100% crossing pt. Crossing points from panels A-C are shown on colorbar (black arrows). (Note slight difference for visual display in colorbar extents compared to *Figure 1J-K*). **(E)** Heatmap of effect size, generated by averaging the 100% crossing point at each pixel; pixels with no data are colored black.

spots' effects (*Figure 1L*) and further, the retinotopic representation of our stimulus is on the bor-
der between these two areas (*Garrett et al., 2014*; *Zhuang et al., 2017*).
Inhibiting these lateral areas, LM/AL and RL, produced large increases in psychometric thresh-
old at low intensities. This resulted in small 100% crossing points (mean 100% crossing point 0.36,
bootstrap 95% CI 0.16-0.59 mW/mm^2) (*Figure 2A*, *Figure 2B*). Inhibition of a control spot between
V1 and LM (same as control spot 1, *Figure 1*) produced no effect on behavior, even at high intensi-
ties (*Figure 2C*). The difference between effects on behavior due to inhibition of lateral areas and
control areas was large (Mann-Whitney $U = 0.0$, lateral N = 6, control N = 4, $p = 0.0071$ two-tailed;
Figure 2E).
The piecewise-linear function we used fixed the slope of each curve. While that did not capture
the exact slope of each optogenetic site (*cf*, *Figure 2A-B*), we used this approach because it seems
likely that true underlying slopes do not vary substantially across sites in these data, as we observed
in the V1 data (*Figure 1–Figure Supplement 2*).
In sum, inhibition of these secondary visual areas, lateral to primary visual cortex, resulted in
substantial degradation of the animals' ability to perform the contrast change detection task.

Inhibiting PM produces smaller effects than inhibiting lateral areas (LM/AL & RL)

In addition to inhibiting areas within LM, AL, and RL, we also inhibited spots medial to V1, within area PM. We found that the average effect of inhibiting spots within PM was to produce little effect on behavior (mean PM 100% crossing point 1.7, bootstrap 95% CI 0.67-2.7 mW/mm^2) (*Figure 3A*, *Figure 3B*). Inhibiting control areas located outside PM had small or undetectable effects on behavior, even at relatively high light intensities (> 5 mW/mm^2, *Figure 3C*). Of five light spots targeted to PM, only one showed an obvious effect, and the effect was of medium size compared to V1 and the lateral areas (*Figure 3D*, *Figure 3E*).

Most PM spots produced small effects on behavior, with the one PM spot that produced a large effect at the most anterior point of PM. That area of PM has a retinotopic representation that is slightly more temporal in azimuth (*Juavinett et al., 2017*; *Zhuang et al., 2017*) than the stimulus we used (25° azimuth). To determine if smaller PM effects relative to LM/AL/RL were due to the retinotopic position of the stimulus, we measured effects of inhibiting anterior PM while animals performed the same task with a stimulus at 45° azimuth (*Figure 3–Figure Supplement 1*). We found similarly small effects as the mean PM effect. The reason for the one larger outlier PM effect is unclear, but across 6 PM spots, using both 25° and 45° azimuth stimuli*Figure 3–Figure Supplement 1*, other PM effects were consistent, leading to a mean PM effect that was small. Thus, even for a more eccentric stimulus that may be better represented in PM, effects of inhibiting PM were smaller than inhibiting LM/AL and RL.

Inhibition of LM/AL/RL produces similar or larger effects on behavior as inhibiting V1

We compared the strength of behavioral effects of inhibition across areas by computing the mean effect (mean 100% crossing point, the measure of how much light is required to produce a fixed degradation in behavior) in each region: lateral areas, PM, and V1. We found that the mean effect was similarly strong (smaller crossing point intensity) in the lateral areas and in V1 (*Figure 4A*; Mann-Whitney $U = 17.5$, lateral N = 6, V1 N = 8, $p = 0.22$ two-tailed), while the mean effect across all PM spots was weaker (larger crossing point intensity) than in V1. The average response heat maps (*Figure 4*; lighter colors are stronger effects on behavior and thus smaller crossing point intensities) show the spatial extent of these effects given the spot positions we tested.

Perceptual behavioral performance can be affected both by changes in sensitivity, the amount of information subjects can use about the stimulus change, or by changes in perceptual criterion, the subject's willingness to respond "yes" (in this case, willingness to release the lever). A more permissive criterion increases hit rate at the cost of a higher false alarm rate. In the V1 data (*Figure 1*), we found that optogenetic inhibition caused little change in false alarm rate, and thus sensitivity (d') changes mirrored the hit rate changes. We asked whether there were similar or different changes in false alarm rates between inhibiting V1 and secondary visual areas. We found no significant difference in early rates between areas (mean ± s.d. early rate increase (OFF - ON) V1: 7.4% ± 4.7%; LM/AL & RL: 9.1% ± 6.0%; PM: 6.6% ± 5.0%; control: 6.7% ± 5.2%) (*Figure 4–Figure Supplement 1*). Given the small effect on early rate, which also varied little between areas, we expected little change in the pattern of our results when measured in terms of sensitivity. Indeed, calculating 100% crossing points in terms of sensitivity (d') had no effect on the observed differences between areas (V1: Mann-Whitney $U = 21.5$, $N_1 = N_2 = 8$, $p = 0.15$ two-tailed; LM/AL and RL: Mann-Whitney $U = 13.5$, $N_1 = N_2 = 6$, $p = 0.26$ two-tailed; PM: Mann-Whitney $U = 9.0$, $N_1 = N_2 = 5$, $p = 0.27$ two-tailed; Control: Mann-Whitney $U = 5.0$, $N_1 = N_2 = 4$, $p = 0.24$, two-tailed) (*Figure 4C*,*Figure 1–Figure Supplement 2*).

When, as here, false alarm rate change is small compared to change in hit rate, changes in absolute criterion (c) are primarily due to changes in hit rate ($c = (z(H) + z(FA))/2$, where H and FA are the hit and false alarm rates, and $z(\cdot)$ is the inverse normal CDF; *Macmillan and Creelman, 2004*). However, relative criterion (c'; $= c/d'$) adjusts for changes in sensitivity. $\Delta c'$ approaches a

Figure 3. Inhibiting medial areas produces weaker effects on behavior. **(A)** All sessions from an example spot within PM, which produced a weak effect (mean 100% crossing pt 1.7, bootstrap 95% CI 1.6-2.0 mW/mm²). Vertical pink dashed line represents average crossing point from LM/AL and RL (0.36 mW/mm²). **(B)** All sessions from an example spot a second spot within PM, which also produced a weak effect (mean 100% crossing pt 3.1, bootstrap 95% CI 2.5-3.5 mW/mm²). **(C)** All sessions from a control area outside PM, showing very little effect even at high intensities (mean 100% crossing pt 6.6, bootstrap 95% CI 4.6-8.1 mW/mm²). **(D)** Map of spots within PM and control locations, colored by 100% crossing pt. Crossing points from panels A-C are shown on colorbar (black arrows). **(E)** Heatmap of effect size, generated by averaging the 100% crossing point at each pixel.

Figure 3–Figure supplement 1. Moving the visual stimulus does not affect PM results

---

286   constant as false alarm rate change goes to zero:

$$\Delta c' = \frac{\Delta c}{\Delta d'} = \frac{z(H_1) + z(FA) - z(H_0) - z(FA)}{2[z(H_1) - z(FA) - z(H_0) + z(FA)]} = \frac{1}{2}$$

Consistent with this, we found that absolute criterion did increase with decreases in hit rate (linear
regression, N = 300 sessions from all visual areas, $t = -22$, $p < 0.001$) but we saw little change in
relative criterion with optogenetic inhibition ($t = -0.87$, N = 300, $p = 0.33$). Thus, the systematic
effects of inhibition on behavior seem to be primarily changes in sensitivity.
Effects of optogenetic inhibition could affect both sensory responses or, in principle, animals'
motor responses. One way to determine whether our optogenetic inhibition affected motor re-
sponses is to vary the motor response the animal uses to report a stimulus change, while keeping
the sensory stimulus the same. We did this by training an animal to perform the task using the
forepaw ipsilateral to the visual stimulus (in contrast to all other behavioral data in this work, where
animals were trained to use their contralateral forepaw). If inhibition affected motor responses,
we would expect to see a different pattern of behavior effects when animals' motor response type
was changed. Instead, we saw similar effects in V1, LM/AL, and RL regardless of whether the ipsilat-
eral or contralateral paw was used to report change (*Figure 4–Figure Supplement 3*). While these

Figure 4. Inhibiting lateral areas produces larger effects than inhibiting PM. **(A)** Average 100% crossing point (± bootstrap 95% confidence interval) across all spots within a given area (V1 / Lateral / Medial / Control). V1 and lateral areas (LM/AL/RL) require very low light intensities to double threshold (mean 0.46, bootstrap 95% CI 0.33-0.58 mW/mm^2 and 0.36, CI 0.16-0.59 mW/mm^2, respectively). Inhibited areas within PM required much higher intensities (mean 1.7, bootstrap 95% CI 0.67-2.7 mW/mm^2) to achieve the same effect, and control areas required very high intensities (mean 5.1, bootstrap 95% CI 2.3-8.0 mW/mm^2). **(B)** Heatmap of all inhibited spots, colored by weighted hit-rate 100% crossing point. Light intensity required to produce effects in lateral secondary areas is similar to or larger than that needed in V1. Mean crossing points from spots within V1, LM/AL & RL (L), PM (M) and control (C) are represented on colorbar with black arrows. **(C)** Heatmap of all inhibited spots, colored by weighted d' 100% crossing point. When 100% crossing point is calculated in terms of d', spots in V1 and lateral areas still require much less intensity to double threshold (means 0.68 and 0.48 mW/mm^2, respectively) than spots in PM and control areas (means 3.0 and 6.1 mW/mm^2, respectively). Mean crossing points are represented on the colorbar using the same notation as **B**.

Figure 4–Figure supplement 1. False-alarm rates do not show systematic variation across areas.

Figure 4–Figure supplement 2. d' 100% crossing point calculations.

Figure 4–Figure supplement 3. Secondary visual area effects are not motor effects

---

data cannot rule out some motor contribution, they are evidence that inhibition of these visual
areas produced an effect primarily on sensory information, not motor execution.

Finally, the visual stimulus location we used for the experiments above (+25° azimuth, 0° ele-
vation) is near the border of area PM, and it might be possible that a stimulus in a more temporal
location (*i.e.* displaced in the horizontal direction away from the animal's nose) could produce a
larger neural response and perhaps reveal larger effects of inhibition. However, when we mea-
sured the effect of inhibition with a visual stimulus at a more lateral azimuth (+45° azimuth, 0°
elevation), we found effects on behavior similarly small as we found before (**Figure 3–Figure Sup-**
plement 1), suggesting that the effect of PM inhibition is not large, even for stimuli at a different,
more eccentric, retinotopic position.

In sum, inhibition of areas lateral to V1 (LM/AL and RL) produces the strongest effects on contrast-
increment change detection, as large or larger than in V1. The fact that some secondary visual areas
(the lateral areas) show these strong effects implies that V1 is not being directly decoded by remote
areas, but instead that information must pass through these secondary visual areas before being
used for contrast change detection.

Optogenetic inhibition acts primarily to suppress secondary visual areas, not on V1 via feedback connections

317 Although we found above that inhibiting secondary visual areas could impact animals' contrast-
318 detection behavior, it could have been possible that the effect of inhibiting secondary visual areas
319 was merely to suppress V1 activity, in a similar way to inhibiting V1 itself. To explore this possibility,
320 we performed a series of electrophysiology experiments, where we inhibited in V1, lateral areas, or

Figure 5. Feedback suppression of V1 during secondary visual area inhibition has little effect on behavior.

Figure 5. (from previous page): (A) Direct optogenetic inhibition of V1, lateral, and medial areas. Plus sign indicates visual stimulus at a mapped retinotopic location in V1. **(B)** Average visual responses in each cortical area with increasing amounts of optogenetic inhibition (0, 0.3, and 3.0 mW/mm^2). Responses shown are baseline corrected to r_{min}, a 25 ms period where optogenetic inhibition has reduced or eliminated spontaneous neural activity but before visually evoked spikes arrive to the cortex. The visual response period is shown as r_{vis}. Black and blue bars show the duration of the visual stimulus and optogenetic inhibitory stimulus respectively. **(C)** Summary of direct inhibition effects on visually evoked responses for all intensities tested. Normalized visual response of 1 is no attenuation (black dotted line). The mean LED intensity for 100% behavior threshold change in V1 is shown with the pink dotted line. **(D)** Inhibition of feedforward V1 connections to secondary visual areas (green arrows). **(E)** Average visual responses measured in lateral areas and PM with increasing optogenetic inhibition applied to V1. Vertical scale bars in panels vary, to more clearly illustrate relative suppression with optogenetic stimulation; quantification is shown in panel F. **(F)** Summary of inhibiting V1 and effects on feedforward visual responses in lateral areas and PM for all intensities tested. **(G)** Feedback suppression of V1 during inhibition of secondary visual areas (red arrows). **(H)** Average visual responses measured in V1 with increasing optogenetic inhibition applied to lateral areas or PM. As in row above, vertical scalebar varies between areas and quantification is shown in panel I. **(I)** Summary of direct V1 inhibition versus feedback suppression for all intensities tested. Yellow shaded region shows the difference between direct and feedback suppression at behaviorally relevant intensities. **(J)** Visual responses in V1 are similarly affected by direct inhibition of V1 at a low intensity (0.1 mW/mm^2) or lateral area feedback suppression at a higher intensity (0.3 mW/mm^2) (mean ± SEM: direct, N = 10; feedback, N = 26; t-test, $t = 0.06$, $p = 0.95$). In contrast, the behavioral effects of inhibiting the lateral areas are not reflective of V1 activity. (mean ± SEM: V1 inh. at 0.1 mW/mm^2, N = 8; lat. inh. at 0.3 mW/mm^2, N = 6; Mann-Whitney $U = 12.0$, $p = 0.06$, one-sided; with 4/6 lat. spots sig greater than V1, Mann-Whitney $U = 0.0$, one-sided, $p < 0.05$ multiple comparisons corrected (Bonferroni)).

Figure 5–Figure supplement 1. Cortical responses to visual stimulus and response timing.

Figure 5–Figure supplement 2. Identification of inhibitory and excitatory units.

---

medial areas while simultaneously recording at the location where optogenetic light was delivered
(Materials and Methods; *Figure 5A*).

We first examined how visual responses were suppressed at the site where optogenetic light
was delivered. We found that optogenetic stimulation of inhibitory neurons reduced or abolished
visual responses at the site of light stimulation in recordings from V1, lateral, and medial secondary
visual areas (units recorded from AL/LM border or in PM; *Figure 5B*). We found that complete sup-
pression directly under the LED could be achieved at low LED intensities (in V1, 85 ± 6% suppression
at 0.3 mW/mm^2), similar to the intensities that produce behavioral effects (in V1, 0.46 mW/mm^2 to
produce a 100% threshold change, *Figure 5C*).

We next examined visual responses to contrast-increments in the different visual cortical areas.
Peak visual responses were similar in size between V1 and lateral areas, and both were significantly
larger than peak responses in PM (*Figure 5–Figure Supplement 1A and B*). We measured visual
stimulus latency (time to half max response, sigmoidal fit between r_{min} to r_{vis}) in V1 to be 65 ms
(bootstrap 95% CI 63-67 ms). There was a small delay in response in the lateral areas relative to V1
(time to half max response = 73 ms, bootstrap 95% CI 70-77 ms) but a notably larger delay for PM
(time to half-max response = 90 ms, bootstrap 95% CI 87-93 ms, *Figure 5–Figure Supplement 1C*).

Since mouse V1 operates as an inhibition stabilized network (*Sanzeni et al., 2020*), at lower light
intensities many inhibitory neurons should decrease their firing while excitatory cells also decrease
their firing, but at high light intensities some inhibitory neurons (especially the narrow-waveform
inhibitory cells, *Sanzeni et al., 2020*) increase their firing. To examine differences in baseline firing
rate change due to stimulation, we sorted units by waveform width (*Figure 5–Figure Supplement 2A*
and B), finding a bimodal distribution of spike waveforms (*Figure 5–Figure Supplement 2C*), as in
Sanzeni et al. (*2020*). For simplicity in the analyses below, we dropped narrow-waveform units
(likely the narrow-spiking basket cells *Speed et al., 2019*; *Sanzeni et al., 2020*), thereby eliminating
units that increased their baseline rates to inhibitory stimulation. For analysis of visual responses,
we focused on wide-waveform units (excitatory and also likely non-basket inhibitory cells). (Addi-
tionally, we saw that all areas - V1, lateral areas, and PM - displayed some signatures of inhibition
stabilized networks, supporting the idea that many visual cortical areas operate in the inhibition-
stabilized regime even without visual input. *Figure 5–Figure Supplement 2D-L*.)

As V1 is the primary recipient of visual input to the cortex and then projects to other visual
areas, we expected that suppressing V1 would reduce responses in other visual areas by blocking
feedforward flow of visual information. Indeed, inhibiting V1 while recording downstream visual re-

sponses in either lateral areas or PM (*Figure 5D*) showed that inhibiting V1 strongly decreased visual
responses in both secondary areas (*Figure 5E*). We saw this effect at LED intensities comparable
to those that produced 100% changes in behavioral threshold, and the feedforward suppression
was even stronger at higher intensities (*Figure 5F*). These measurements support the idea that the
majority of the visual information seen in secondary visual areas passes through V1.

We next examined whether our behavioral effects in secondary areas originated solely from
local suppression of visual responses or if they could have arisen from feedback suppression of V1.
That is, we studied whether inhibition of secondary visual areas produced behavioral deficits due to
suppression of responses in the secondary area itself, or whether inhibition could have potentially
produced behavioral deficits merely by acting on the V1 responses via feedback connections. To
study this, we recorded in V1 and simultaneously inhibited lateral areas or PM while awake animals
were passively viewing the visual stimulus used for behavior (*Figure 5G*). We saw that optogenetic
suppression of lateral areas did partially reduce V1 responses. At light intensities that produced
behavioral effects (0.36 mW/mm^2), inhibiting LM/AL produced moderate (31 ± 4%) suppression
in V1. However, the same light stimulation nearly eliminated visual responses in LM/AL. Thus,
the principal effect of stimulation of LM/AL was to suppress LM/AL, with a secondary effect of
suppression in V1.

In contrast to the feedback effects on LM/AL, we found that suppression of medial areas had lit-
tle to no feedback effect on V1 visual responses, even at the highest intensities we used (*Figure 5H,*
right).

Finally, to directly compare the effects on both areas (V1 and LM/AL) of optogenetic suppression
of LM/AL, we examined V1 and LM/AL stimulation intensities that produced a matching magnitude
of V1 suppression. Delivering 0.1 mW/mm^2 to V1 and 0.3 mW/mm^2 to LM/AL, while in both cases
recording in V1, produced similar levels of suppression on the visual responses of V1 neurons
(*Figure 5J, left*). We were then able to examine in our behavioral data how the two types of stimu-
lation (0.1 mW/mm^2 to V1; 0.3 mW/mm^2 to LM/AL), which affected V1 responses similarly, affected
animals' behavior. We found that LM/AL stimulation, and not V1 stimulation, produced large degra-
dation of animals' contrast change detection behavior (*Figure 5J, right*). These observations argue
that when lateral secondary areas LM/AL are suppressed, it is not the change in V1 responses that
drives the behavioral effect (since stimulating V1 itself at an intensity necessary to produce the
same V1 change has no effect on behavior). Instead, likely it is the dramatic suppression of LM/AL
activity (88 ± 12%) at this light intensity that leads to the behavioral degradation. In sum, though in-
hibition of lateral areas does suppress activity in V1, feedback suppression alone could not explain
the observed behavioral results.

Discussion

This work examines the effect of inhibiting secondary visual areas in *VGAT-ChR2-EFYP* mice per-
forming a contrast-increment change detection task. Neurons in V1, lateral, and medial secondary
visual areas change their firing rate in response to changes in stimulus contrast, but only suppress-
ing V1 and the lateral areas produced a large impact on animals' performance in the task. Inhibiting
areas lateral to V1 sometimes caused deficits larger than those seen when inhibiting V1 directly.
Behavioral effects were due to changes in sensitivity (d′) without large changes in false alarm rates.
When inhibiting the secondary areas, light intensities that produced behavioral changes produced
the largest changes in neural firing in the secondary areas themselves, not in V1 via feedback sup-
pression. Taken together, our results provide evidence that while substantial information about
the contrast changes is present in V1, secondary areas AL/LM and RL form an essential part of the
neural circuit that controls this behavior.

399 Behavioral changes are consistent with sensory, not motor representations, and are spa-
400 tially specific

We performed two control experiments to determine whether inhibition affected sensory signals
rather than motor planning or execution, and to determine if PM effects were impacted by retino-
topic maps. First, if our inhibition affected motor performance, we would have expected that such
an effect would vary based on whether animals were using the contralateral or ipsilateral forepaw.
Instead, we found that training animals to perform the task with their ipsilateral forepaw produced
similar results as when animals used their contralateral forepaw (*Figure 4–Figure Supplement 3*).
Second, we examined the possibility that our observed PM effects were weaker because PM may
contain a weaker representation of the nasal than the temporal visual field. To rule this out, we
measured the effect of inhibiting PM during the same task but performed with a stimulus further
into the temporal visual field (45° azimuth relative to midline), where previous studies have found
robust retinotopic representations in PM. Using this stimulus, we found similar magnitude effects
as with our 25° eccentricity stimulus (*Figure 3–Figure Supplement 1*, arguing that the weaker effects
in PM relative to lateral areas were not due to the eccentricity of the visual stimulus.

Our results examine the effect of inhibiting the cortex at a variety of different locations, and
rely on several observations to establish spatial specificity. First, our key finding is that inhibiting
lateral visual areas, LM and/or AL as well as RL, produces similar-sized, or larger, effects on behav-
ior as inhibiting V1. The cortical distance involved between the lateral areas and V1 (approximately
2 mm) and the fact that several control spots with equally distant spacing (as well as PM loca-
tions) produced little or no effect, is itself evidence that our effects are due to inhibition at the site
where the light was positioned. Second, the spatial specificity observed in behavioral experiments
is confirmed by our neurophysiological recordings. We observed that the evoked visual response
showed a similar spatial falloff as the light intensity patterns (spots) that we used, with the half-
maximum suppression approximating the half-maximum light intensity for our average spot sizes
(*Figure 1L*). Inhibition affected neurons' baseline or spontaneous activity over a larger area than the
area over which evoked visual responses were affected (*Figure 1–Figure Supplement 3*), consistent
with balanced-state spiking models of cortical function, where small shifts of excitatory-inhibitory
balance can cause large changes in spontaneous state while neurons continue to produce evoked
responses to input (e.g. *Vogels and Abbott, 2009*). Another factor that may have changed our inten-
sity thresholds, but did not affect the pattern of results across areas, is variability in effect size due
to stimulation spot size. Even if there is some association of effect size with spot size, considering
a subset of spots with similar sizes confirms our effects (e.g. *Figure 2D*, compare "A", "B", and the
control spots shown in dashed lines; also see *Figure 3D*, regions marked "B", "C", etc.). In one case,
we used a small spot to separate AL/LM from V1 (*Figure 2D*, spot "C"), and inhibiting there produced
no discernible effect on behavior up to the highest power that we used (*Figure 2C*), though the spot
area was not much smaller than other LM/AL spot areas, which produced effects at much lower
light intensities (*Figure 2D*). Even if this spot were ignored, our principal finding that inhibition of
secondary visual areas affects this behavior is unchanged. Finally, the heatmaps we use to display
the spatial extent of behavioral effects (e.g. *Figure 4B-C*) are limited to showing effects only at the
locations where we sited stimulation spots, and thus they are principally useful not for quantifica-
tion of the potential effect at every pixel location, but instead are useful to visualize average effects
across areas.

442 Roles of mouse cortical visual areas in contrast perception behaviors

These findings show not only that lateral areas LM/AL/RL play a role in this behavior, but also
that medial areas like PM do not play a major role. (Some visual cortex parcellations include an
area, anteromedial or AM, between PM and RL, *Garrett et al., 2014*; *Zhuang et al., 2017*, but our
conclusions do not significantly depend on whether our medial regions are part of an anterior
section of PM, or a separate area AM.) Both lateral and medial areas have previously been shown
to encode different features of visual stimuli, such as spatial and temporal frequencies (*Andermann*

et al., 2011; *Murakami et al., 2017*), and thus might play object identification or localization roles, as in the 'what' and 'where' pathways of the primate visual system (*Mishkin and Ungerleider, 1982*). In this case, both lateral and medial areas might be representing different features of the visual world, and other types of behavioral decision-making might expose the role of PM (see *Jin and Glickfeld, 2020*). On the other hand, *Siegle et al.* (*2019*) constructed a hierarchy of visual areas based on electrophysiology data (LM->RL->LP->AL->PM->AM), and predicted that medial areas (PM/AM) were higher areas within the hierarchy, which serve to amplify or decode change-related signals. Our data argue against that hypothesis. If PM/AM were important for processing signals for decisions about contrast changes, inhibiting them should also serve to impair behavior, and we found only weak mean effects.

A recent study (*Jin and Glickfeld, 2020*) lends further support to the idea that PM is not crucial for contrast-perception or change detection tasks. Jin and colleagues found, as we did, that lateral areas are involved in contrast change detection tasks, and medial area inhibition has a smaller effect. Our electrophysiological results go beyond these behavioral results, and show that the effect of inhibiting secondary visual areas is not caused by effects on V1 but substantially caused by changes in the visual signals in the secondary areas. However, our results on perceptual sensitivity and criterion did differ from theirs; while *Jin and Glickfeld, 2020* found little or no change in sensitivity due to inhibiting PM, as we did, they found an effect on false alarm rates when PM was inhibited. Because subjects have some ability to choose their own perceptual criterion in perceptual behaviors (*Macmillan and Creelman, 2004*), it seems possible that this difference in false alarm rate is an effect of behavioral training. In sum, the lack of sensitivity change when PM is suppressed argues that PM is not part of the circuit that controls behavior in this contrast perception task, while V1, LM/AL and RL do play important roles in this behavior.

Brain area interactions during perceptual behaviors

Our results shed light on the inter-area circuits used in a perceptual behavioral task. Recent work has shown that mouse premotor cortex is involved in downstream processing of sensory signals (*Wu et al., 2020*; *Zatka-Haas et al., 2020*), but it has been unclear whether secondary visual areas are involved in all or most sensory tasks that involve V1. Our data argue that lateral secondary areas are not bypassed during behavior, and therefore suggests the secondary areas' activity is decoded into a choice or decision representation via those areas' projections to downstream regions. Evidence for choice or decision signals has been found in areas including frontal and premotor cortex and the striatum, but not in either V1 or secondary areas (*Steinmetz et al., 2019*), arguing that the decoding happens outside both V1 and the secondary areas. However, there are several caveats to this sequential interpretation of information flow from V1 to the secondary areas and then on to decoding areas. First, it is possible but unlikely that suppressing activity in cortical areas, as we did, acts on behavior by changing the activity of another area that is truly performing the neural computations for the task. For that to be true, it would require that the visual responses in each of these cortical visual areas are all epiphenomenal. A more likely alternative hypothesis to the sequential interpretation is that multiple brain areas are read out together in parallel. Deficits in visual detection similar to those we observed are also seen with inhibition of the mouse superior colliculus (SC) (*Wang et al., 2020*). Our observations and the SC results suggest that cortical circuits and the colliculus both carry information about visual tasks, and since optogenetic inhibition in each of these studies does not totally abolish the behavior, the areas may each contribute information in near-threshold tasks such as this. This might imply that when animals are performing near perceptual threshold, they use all available representations (e.g. multiple cortical areas, the SC, and potentially others) together to perform the behavior. If it were true that animals do use, or decode, all available representations in sensory areas, this would have implications not just for the circuits used during perceptual tasks, but also for the total amount of information available to the brain about particular stimuli, as calculations of this sort have previously focused only on V1 (*Kafashan et al., 2020*; *Rumyantsev et al., 2020*; *Stringer et al., 2019*).

499 How patterns of neural activity are processed across multiple brain regions to control behavior
500 is a central question about brain function. The role of secondary visual areas in sensory behaviors
501 shown by these results suggests that the mouse visual system, like the visual systems of other
502 species, contains a set of cortical areas that work in concert during perceptual behavior.

503 ## Materials and Methods

Key Resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Chemical compound	Tamoxifen	Sigma-Aldrich	T5648-5G	
Chemical compound	PEDOT; poly(3,4-ethylenedioxythiophene)	Sigma-Aldrich	687553	
Chemical compound	PSS; poly(styrenesulfonate)	Sigma-Aldrich	243051	
Genetic reagent	VGAT-ChR2-EYFP	The Jackson Laboratory	RRID:IMSR_JAX:014548	
Genetic reagent	Ai148	The Jackson Laboratory	RRID:IMSR_Jax:030328	
Genetic reagent	Cux2-CreERT2	MMRRC	RRID:MMRRC_032779-MU	
Software	MWorks	The MWorks Project		mworks.github.io
Software	Offline Sorter	Plexon	RRID:SCR_000012	
Other	C&B Metabond	Parkell	S380	
Other	Kwik-sil	World Precision Instruments	KWIK-SIL	

Animals
All experimental procedures were approved by the NIH Institutional Animal Care and Use Commit-
tee (IACUC) and complied with Public Health Service policy on the humane care and use of labo-
ratory animals. Twelve *VGAT-ChR2-EYFP* mice (ChR2 targeted at the *Slc32a1* locus) were used for
behavioral and electrophysiology experiments, and three tamoxifen-treated *Ai148;Cux2-CreERT2*
mice (GCaMP6f targeted at the *Igs7* locus and CreERT2 targeted at the *Cux2* locus, respectively)
were used for imaging experiments. (Jax stock #s 01458, 030328; MMRRC # 032779-MU; seven
females and eight males; singly housed on a reverse light / dark cycle).

Cranial window implantation
Mice were given intraperitoneal dexamethasone (3.2 mg/kg) and anesthetized with isoflurane (1.0-
3.0% in 100% O_2 at 1 L/min). Using aseptic technique, a titanium headpost was affixed using C&B
Metabond (Parkell) and a 5 mm diameter craniotomy was made, centered over V1 (-3.1 mm ML,
+1.5 mm AP from lambda). A 5 mm cranial window was cemented into the craniotomy, providing
chronic access to visual cortex and surrounding higher visual areas.Post-surgery mice were given
sub-cutaneous 72-hour slow-release buprenorphine (0.50 mg/kg) and recovered on a heating pad.

Hemodynamic imaging & visual area map fitting
To determine the location of V1 and surrounding visual areas, we delivered small visual stimuli to
head-fixed animals at different retinotopic positions and measured hemodynamic-related changes

in absorption by measuring reflected 530 nm light (*Schuett et al., 2002*). Light was delivered with
a 530 nm fiber coupled LED (M350F2, Thorlabs) passed through a 530 nm-center bandpass filter
(FB530-10, Thorlabs). Images were collected on a Zeiss Discovery stereo microscope with a 1X wide-
field objective through a green long-pass emission filter onto a Retiga R3 CCD camera (QImaging,
Inc., captured at 2 Hz with 4x4 binning). We presented upward-drifting square wave gratings (2
Hz, 0.1 cycles/degree) masked with a raised cosine window (10° diameter) at different retinotopic
locations for 5 seconds with 10 seconds of mean luminance between each trial. Stimuli were pre-
sented in random order at six positions in the right monocular field of view. The hemodynamic
response to each stimulus was calculated as the change in reflectance of the cortical surface be-
tween the baseline period and a response window starting 2-3 s after stimulus onset. This window
corresponds to the previously reported timecourse of visually evoked hemodynamic responses in
mouse V1 (*Heimel et al., 2007*). Visual area maps were fitted to the cortex based on centroid of
each stimulus response.

Widefield GCaMP Imaging

To validate the placement of the hemodynamic map fits, we applied our hemodynamic imaging
of V1 and map fitting procedure in animals in which we also performed GCaMP imaging of V1
and secondary visual areas. Animals (*Ai148;CuX2-CreERT2*; given tamoxifen at P18-22) expressing
GCaMP6f in L2/3 cortical excitatory cells were head fixed while a Gabor stimulus (same stimulus
used in behavior: +25° azimuth, 0° elevation, 12° in diameter, spatial frequency 0.1 cycle/degree)
was shown in their right monocular visual field. We recorded GCaMP fluorescence changes through
the 5 mm cranial implant over the contralateral visual cortex. Green fluorescence images were
captured using an eGFP filter, at a frame rate of 10 Hz (1 stimulus frame followed by 32 post-
stimulus frames, 4x4 binning) for 150 stimulus presentations. A 10 uL water reward was given
randomly (10% probability) to keep animals awake and alert.

Behavioral Task

Mice were head-fixed in sound-attenuating custom boxes, and trained to hold a lever and release
it when a Gabor stimulus increased its contrast (*Histed et al., 2012*). Mice were required to a lever
for a maximum of 4.1 s. At a random time point during this period, a small Gabor patch (+25°
azimuth, 0° elevation, 12° in diameter, spatial frequency 0.1 cycle/degree) was presented for 100
ms and animals had 550 ms to report the stimulus by releasing the lever. Possible trial outcomes
included a correct release within the reaction time window (hit), a failure to release the lever in
response to the stimulus (miss), or a false alarm trial where the animal released the lever before
the stimulus had appeared (early). Lever releases occurring in the first 100 ms following the stim-
ulus presentation were counted as earlies or false alarms, as the animal could not have possibly
perceived the stimulus and reacted in 100 ms. Trials resulting in hits were rewarded with 1.5-3 uL
liquid reward. We began training with a full-screen, 100% contrast stimulus. As animals learned
to perform in this task, the size and contrast of the stimulus were decreased and location of the
stimulus was slowly moved to the desired azimuth and elevation. As performance stabilized, we
also increased the number of contrast levels to cover a range of contrasts or difficulties in order
to determine the animals psychometric threshold (the contrast at which the animal was getting
63% of trials correct). Sessions lasted until the animal reached its daily water supplement or until
the animal stopped on its own, whichever came first (mean session length = 384 ± 89 trials). Be-
cause head-fixed mice make rare eye movements and SC inhibition does not systematically change
those movements (*Wang et al., 2020*), mice have no visual streak and small difference in ganglion
cell density across the retina, the secondary areas are small enough that our spots covered a large
part of their retinotopic maps, and the few eye movements made by head-fixed mice are thought
to be compensatory movements that occur with attempted head rotation (*Meyer et al., 2020*), we
did not record animals' eye movements.

Optogenetic Inhibition

Fiber optic cannulae (400 um diameter 0.39 NA, Thorlabs CFMLC14L02) were dental-cemented above specific areas as determined by hemodynamic intrinsic imaging. Spot size was determined as the full-width half-maximum (FWHM) of the illuminated area (spot area mean ± SEM, 0.42 ± 0.13 mm^2, N=30). We used a fiber-coupled LED light source (M470F3, ThorLabs) to deliver illumination with a peak wavelength of 470 nm to the brain. Once animals were performing well on the visual task and we had a full psychometric curve, we introduced a pulsed LED (on for 200 ms, off for 1000 ms) to each trial. The LED pulse train began with the start of the trial and pulsed 0-5 times before the visual stimulus was presented. On 50% of trials, the visual stimulus was presented 55 ms before the last LED pulse in the train (ON trials). On the remaining 50% of trials, the visual stimulus was presented 180 ms before the last LED pulse (OFF trials). Visual onset times were distributed according to a geometric distribution, whose mean was adjusted to balance the desire for a flat hazard function (to discourage waiting for later stimulus times; often reflected in false alarm rates < 5% of trials) and to keep false alarms less than approximately 50% of trials. Once a visual onset time was randomly chosen, it was adjusted by lowering it to either 55 ms after the previous light pulse onset (ON trials) or 820 ms after the previous light pulse offset (OFF trials).

Behavior Analysis

All analyses were performed with Python. Threshold increases were determined for each behavioral session by fitting a Weibull cumulative density function (hit rates) or Naka-Rushton function (d'; *Herrmann et al., 2012*) across contrast levels. The upper asymptote (lapse rate) and threshold (63% point between the upper and lower asymptotes, Weibull; 50% point, Naka-Rushton) were fit for each trial type with a single slope for LED ON and OFF trials. Only sessions with low lapse rates (<20%) and sufficient trials at each contrast (average 10 trials per contrast level, per trial type) were included for further analysis (539 sessions total; 166 excluded, 373 sessions included). To determine the 100% crossing point for each spot, or the intensity required to double the psychometric threshold, a piecewise-linear function was fit to all percent increases obtained from various LED intensities. Fits were initially done on V1 data with variable slopes. The slopes of spots with low slope CI ratios (< 9 upper:lower CI, *Figure 1–Figure Supplement 2B*) were averaged and the regressions were re-fit using this determined slope (268.7). Piecewise-linear functions were fit in the same manner described above (*Figure 4–Figure Supplement 2*), first with variable slopes to V1 data and then again with a fixed average slope (261.2), in order to determine a d' 100% crossing point for each spot. Heatmaps were generated by averaging the 100% crossing point at each pixel. Pixels with no data are black.

Electrophysiological recordings

For recording experiments, we first affixed a 3D printed ring to the cranial window to retain fluid. With the ring in place, we removed the cranial window and flushed the craniotomy site with sterile normal saline to remove debris. Kwik-Sil silicone adhesive (World Precision Instruments) was used to seal the craniotomy between recording days. Optogenetic light was delivered using the same optical fiber and cannula used for behavioral experiments. The cannula distance from the dura was adjusted to provide a light spot with a full width at half maximum intensity of 0.36 ± 0.07 mm (mean ± SEM, N=5, measured with a USB digital macro documentation microscope, Opti-TekScope). We targeted a multisite silicon probe electrode (NeuroNexus; 32-site model 4x8-100-200-177; 4 shank, 8 sites/shank; sites electrochemically coated with PEDOT:PSS [poly(3,4-ethylenedioxythiophene): poly(styrenesulfonate)]) to the center of the LED spot using a micromanipulator (MPC-200, Sutter Instruments). With the fiber optic cannula and electrode in place, we removed the saline buffer using a sterile absorbent triangle (Electron Microscopy Sciences, Inc.) and allowed the dura to dry for 5 minutes. After insertion, we waited 30-60 minutes without moving the probes to reduce slow drift and provide more stable recordings. We isolated single and multiunit threshold crossings (3 times RMS noise) by amplifying the site signals filtered between 750 Hz and 7.5 Khz (Cerebus, Blackrock

619 microsystems). The visual stimulus (90% contrast Gabor, 100 ms duration, same as during behav-
620 ioral experiments) was presented on every trial, paired with a 200 ms LED pulse at varying intensity
621 (180 reps each intensity). To keep animals awake and alert, animals were water-scheduled and a
622 1 uL water reward was randomly provided on 5% of the stimulus trials; we verified animals were
623 licking in response to rewards during the experiments.

Electrophysiology Analysis

Spike waveforms were extracted and sorted using Offline Sorter (Plexon, Inc.). Single units were
identified using waveform clusters that showed separation from noise and unimodal width distri-
butions. Single units had SNR (*Kelly et al., 2007*; *Histed, 2018*; *Sanzeni et al., 2020*) of 3.50 ± 0.34
(mean ± std, N = 221) with a multi unit SNR of 3.08 ± 0.27 (mean ± std, N = 69). Multi units were
excluded from further analysis. In addition to examining the overall average spike rates of the re-
sponses, we calculated mean activity in three response windows: a baseline (r_{base}) window 40 ms
in duration ending 10 ms prior to the delivery of the optogenetic stimulus, an inhibitory minimum
(r_{min}) window of 25 ms starting 25 ms after optogenetic stimulation that ends just before spikes
arrive to V1 (~40 ms, *Sanzeni et al., 2020*), and a maximum visual response (r_{vis}) window of 50 ms
starting 65 ms post visual stimulus for V1 and lateral areas. The max window was shifted 25 ms
later for PM due to delay in the midpoint of the visual response (*Figure 1–Figure Supplement 1*,
Figure 5–Figure Supplement 1). Visually responsive units were identified as having an r_{vis} 10% or
greater over r_{base} with no inhibition (0 mW/mm^2 optogenetic stim). Response traces were base-
line corrected to r_{min}, and the normalized change in spikes rates calculated as r_{vis} - r_{min} / r_{vis} at 0
mW/mm^2.

Acknowledgments

We are grateful to Victoria Scott for assistance with animal husbandry and to Patrick Wright for technical advice on hemodynamic imaging. We thank Richard Krauzlis, Bruno Averbeck, and members of the Histed lab for their comments on the manuscript.

References

Albrecht DG, Hamilton DB. Striate cortex of monkey and cat: contrast response function. Journal of Neurophysiology. 1982; 48(1):217–237. https://doi.org/10.1152/jn.1982.48.1.217, doi: 10.1152/jn.1982.48.1.217, pMID: 7119846.

Andermann ML, Kerlin AM, Roumis DK, Glickfeld LL, Reid RC. Functional Specialization of Mouse Higher Visual Cortical Areas. Neuron. 2011 Dec; 72(6):1025–1039. https://linkinghub.elsevier.com/retrieve/pii/S0896627311010129, doi: 10.1016/j.neuron.2011.11.013.

Berens P, Ecker AS, Cotton RJ, Ma WJ, Bethge M, Tolias AS. A Fast and Simple Population Code for Orientation in Primate V1. Journal of Neuroscience. 2012 Aug; 32(31):10618–10626. http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.1335-12.2012, doi: 10.1523/JNEUROSCI.1335-12.2012.

Busse L, Ayaz A, Dhruv NT, Katzner S, Saleem AB, Schölvinck ML, Zaharia AD, Carandini M. The detection of visual contrast in the behaving mouse. The Journal of neuroscience : the official journal of the Society for Neuroscience. 2011 August; 31(31):11351—11361. https://europepmc.org/articles/PMC6623377, doi: 10.1523/jneurosci.6689-10.2011.

Caviness VS. Architectonic map of neocortex of the normal mouse. The Journal of Comparative Neurology. 1975 Nov; 164(2):247–263. http://doi.wiley.com/10.1002/cne.901640207, doi: 10.1002/cne.901640207.

Dräger UC. Receptive fields of single cells and topography in mouse visual cortex: MOUSE VISUAL CORTEX. Journal of Comparative Neurology. 1975 Apr; 160(3):269–289. http://doi.wiley.com/10.1002/cne.901600302, doi: 10.1002/cne.901600302.

Froudarakis E, Fahey PG, Reimer J, Smirnakis SM, Tehovnik EJ, Tolias AS. The Visual Cortex in Context. Annual Review of Vision Science. 2019 Sep; 5(1):317–339. https://www.annualreviews.org/doi/10.1146/annurev-vision-091517-034407, doi: 10.1146/annurev-vision-091517-034407.

Garrett ME, Nauhaus I, Marshel JH, Callaway EM. Topography and Areal Organization of Mouse Visual Cortex. Journal of Neuroscience. 2014 Sep; 34(37):12587–12600. http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.1124-14.2014, doi: 10.1523/JNEUROSCI.1124-14.2014.

Glickfeld LL, Histed MH, Maunsell JHR. Mouse Primary Visual Cortex Is Used to Detect Both Orientation and Contrast Changes. Journal of Neuroscience. 2013 Dec; 33(50):19416–19422. http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.3560-13.2013, doi: 10.1523/JNEUROSCI.3560-13.2013.

Han Y, Kebschull JM, Campbell RAA, Cowan D, Imhof F, Zador AM, Mrsic-Flogel TD. The logic of single-cell projections from visual cortex. Nature. 2018 Apr; 556(7699):51–56. http://www.nature.com/articles/nature26159, doi: 10.1038/nature26159.

Heimel JA, Hartman RJ, Hermans JM, Levelt CN. Screening mouse vision with intrinsic signal optical imaging. European Journal of Neuroscience. 2007; 25(3):795–804. https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1460-9568.2007.05333.x, doi: 10.1111/j.1460-9568.2007.05333.x.

Herrmann K, Heeger DJ, Carrasco M. Feature-based attention enhances performance by increasing response gain. Vision Research. 2012 Dec; 74:10–20. https://linkinghub.elsevier.com/retrieve/pii/S0042698912001381, doi: 10.1016/j.visres.2012.04.016.

Histed MH. Feedforward Inhibition Allows Input Summation to Vary in Recurrent Cortical Networks. eneuro. 2018 Jan; 5(1):ENEURO.0356–17.2018. http://eneuro.org/lookup/doi/10.1523/ENEURO.0356-17.2018, doi: 10.1523/ENEURO.0356-17.2018.

Histed MH, Carvalho LA, Maunsell JHR. Psychophysical measurement of contrast sensitivity in the behaving mouse. Journal of Neurophysiology. 2012 Feb; 107(3):758–765. https://www.physiology.org/doi/10.1152/jn.00609.2011, doi: 10.1152/jn.00609.2011.

Hubel DH, Wiesel TN. Receptive fields and functional architecture of monkey striate cortex. The Journal of Physiology. 1968 Mar; 195(1):215–243. http://doi.wiley.com/10.1113/jphysiol.1968.sp008455, doi: 10.1113/jphysiol.1968.sp008455.

Jin M, Glickfeld LL. Mouse higher visual areas provide both distributed and discrete contributions to visually guided behaviors. bioRxiv. 2020 Mar; http://biorxiv.org/lookup/doi/10.1101/2020.03.21.001446, doi: 10.1101/2020.03.21.001446.

Juavinett AL, Nauhaus I, Garrett ME, Zhuang J, Callaway EM. Automated identification of mouse visual areas with intrinsic signal imaging. Nature Protocols. 2017 Jan; 12(1):32–43. http://www.nature.com/articles/nprot.2016.158, doi: 10.1038/nprot.2016.158.

Kafashan M, Jaffe A, Chettih SN, Nogueira R, Arandia-Romero I, Harvey CD, Moreno-Bote R, Drugowitsch J. Scaling of information in large neural populations reveals signatures of information-limiting correlations. bioRxiv. 2020; https://www.biorxiv.org/content/early/2020/01/11/2020.01.10.902171, doi: 10.1101/2020.01.10.902171.

Kalatsky VA, Stryker MP. New paradigm for optical imaging: temporally encoded maps of intrinsic signal. Neuron. 2003 May; 38(4):529–545. doi: 10.1016/s0896-6273(03)00286-1, place: United States.

Kelly RC, Smith MA, Samonds JM, Kohn A, Bonds AB, Movshon JA, Sing Lee T. Comparison of Recordings from Microelectrode Arrays and Single Electrodes in the Visual Cortex. Journal of Neuroscience. 2007 Jan; 27(2):261–264. http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.4906-06.2007, doi: 10.1523/JNEUROSCI.4906-06.2007.

Khastkhodaei Z, Jurjut O, Katzner S, Busse L. Mice Can Use Second-Order, Contrast-Modulated Stimuli to Guide Visual Perception. Journal of Neuroscience. 2016; 36(16):4457–4469. https://www.jneurosci.org/content/36/16/4457, doi: 10.1523/JNEUROSCI.4595-15.2016.

Khibnik LA, Tritsch NX, Sabatini BL. A Direct Projection from Mouse Primary Visual Cortex to Dorsomedial Striatum. PLoS ONE. 2014 Aug; 9(8):e104501. https://dx.plos.org/10.1371/journal.pone.0104501, doi: 10.1371/journal.pone.0104501.

Leopold DA. Primary Visual Cortex: Awareness and Blindsight. Annual Review of Neuroscience. 2012 Jul; 35(1):91–109. http://www.annualreviews.org/doi/10.1146/annurev-neuro-062111-150356, doi: 10.1146/annurev-neuro-062111-150356.

Macmillan NA, Creelman CD. Detection Theory: A User's Guide. Psychology Press; 2004. Google-Books-ID: P094AgAAQBAJ.

Meyer AF, O'Keefe J, Poort J. Two Distinct Types of Eye-Head Coupling in Freely Moving Mice. Current Biology. 2020; http://www.sciencedirect.com/science/article/pii/S096098222030556X, doi: https://doi.org/10.1016/j.cub.2020.04.042.

Mishkin M, Ungerleider LG. Contribution of striate inputs to the visuospatial functions of parieto-preoccipital cortex in monkeys. Behavioural Brain Research. 1982 Sep; 6(1):57–77. https://linkinghub.elsevier.com/retrieve/pii/016643288290081X, doi: 10.1016/0166-4328(82)90081-X.

Murakami T, Matsui T, Ohki K. Functional Segregation and Development of Mouse Higher Visual Areas. The Journal of Neuroscience. 2017 Sep; 37(39):9424–9437. http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.0731-17.2017, doi: 10.1523/JNEUROSCI.0731-17.2017.

Rumyantsev OI, Lecoq JA, Hernandez O, Zhang Y, Savall J, Chrapkiewicz R, Li J, Zeng H, Ganguli S, Schnitzer MJ. Fundamental bounds on the fidelity of sensory cortical coding. Nature. 2020 Apr; 580(7801):100–105. http://www.nature.com/articles/s41586-020-2130-2, doi: 10.1038/s41586-020-2130-2.

Sanzeni A, Akitake B, Goldbach HC, Leedy CE, Brunel N, Histed MH. Inhibition stabilization is a widespread property of cortical networks. eLife, in press. 2020; https://www.biorxiv.org/content/early/2019/06/02/656710, doi: 10.1101/656710.

Schuett S, Bonhoeffer T, Hübener M. Mapping Retinotopic Structure in Mouse Visual Cortex with Optical Imaging. The Journal of Neuroscience. 2002 Aug; 22(15):6549–6559. http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.22-15-06549.2002, doi: 10.1523/JNEUROSCI.22-15-06549.2002.

Sclar G, Maunsell JHR, Lennie P. Coding of image contrast in central visual pathways of the macaque monkey. Vision Research. 1990; 30(1):1 – 10. http://www.sciencedirect.com/science/article/pii/0042698990901233, doi: https://doi.org/10.1016/0042-6989(90)90123-3.

Siegle JH, Jia X, Durand S, Gale S, Bennett C, Graddis N, Heller G, Ramirez TK, Choi H, Luviano JA, Groblewski PA, Ahmed R, Arkhipov A, Bernard A, Billeh YN, Brown D, Buice MA, Cain N, Caldejon S, Casal L, et al. A survey of spiking activity reveals a functional hierarchy of mouse corticothalamic visual areas. bioRxiv. 2019 Oct; http://biorxiv.org/lookup/doi/10.1101/805010, doi: 10.1101/805010.

Speed A, Rosario JD, Burgess CP, Haider B. Cortical State Fluctuations across Layers of V1 during Visual Spa-
tial Perception. Cell Reports. 2019 Mar; 26(11):2868–2874.e3. https://www.cell.com/cell-reports/abstract/
S2211-1247(19)30216-5, doi: 10.1016/j.celrep.2019.02.045, publisher: Elsevier.

Steinmetz NA, Zatka-Haas P, Carandini M, Harris KD. Distributed coding of choice, action and engagement
across the mouse brain. Nature. 2019 Dec; 576(7786):266–273. https://doi.org/10.1038/s41586-019-1787-x,
doi: 10.1038/s41586-019-1787-x.

Stringer C, Michaelos M, Pachitariu M. High precision coding in visual cortex. bioRxiv. 2019 Jun; http://biorxiv.
org/lookup/doi/10.1101/679324, doi: 10.1101/679324.

Tsao D. The Macaque Face Patch System: A Window into Object Representation. Cold Spring Harbor
Symposia on Quantitative Biology. 2014 Jan; 79:109–114. http://symposium.cshlp.org/content/79/109, doi:
10.1101/sqb.2014.79.024950, publisher: Cold Spring Harbor Laboratory Press.

Vaiceliunaite A, Erisken S, Franzen F, Katzner S, Busse L. Spatial integration in mouse primary visual cortex.
Journal of Neurophysiology. 2013 Aug; 110(4):964–972. https://www.physiology.org/doi/10.1152/jn.00138.
2013, doi: 10.1152/jn.00138.2013.

Vogels TP, Abbott LF. Gating multiple signals through detailed balance of excitation and inhibition in spiking
networks. Nature Neuroscience. 2009 Apr; 12(4):483–491. http://www.nature.com/articles/nn.2276, doi:
10.1038/nn.2276.

Wagor E, Mangini NJ, Pearlman AL. Retinotopic organization of striate and extrastriate visual cortex in the
mouse. The Journal of Comparative Neurology. 1980 Sep; 193(1):187–202. http://doi.wiley.com/10.1002/cne.
901930113, doi: 10.1002/cne.901930113.

Wang L, McAlonan K, Goldstein S, Gerfen CR, Krauzlis RJ. A Causal Role for Mouse Superior Colliculus in Visual
Perceptual Decision-Making. Journal of Neuroscience. 2020; 40(19):3768–3782. https://www.jneurosci.org/
content/40/19/3768, doi: 10.1523/JNEUROSCI.2642-19.2020.

Wang Q, Burkhalter A. Area map of mouse visual cortex. The Journal of Comparative Neurology. 2007 May;
502(3):339–357. http://doi.wiley.com/10.1002/cne.21286, doi: 10.1002/cne.21286.

Wu Z, Litwin-Kumar A, Shamash P, Taylor A, Axel R, Shadlen MN. Context-Dependent Decision Making in
a Premotor Circuit. Neuron. 2020 Apr; 106(2):316–328.e6. https://linkinghub.elsevier.com/retrieve/pii/
S0896627320300611, doi: 10.1016/j.neuron.2020.01.034.

Zatka-Haas P, Steinmetz NA, Carandini M, Harris KD. A perceptual decision requires sensory but not action cod-
ing in mouse cortex. bioRxiv. 2020 Dec; http://biorxiv.org/lookup/doi/10.1101/501627, doi: 10.1101/501627.

Zhao S, Ting JT, Atallah HE, Qiu L, Tan J, Gloss B, Augustine GJ, Deisseroth K, Luo M, Graybiel AM, Feng
G. Cell type–specific channelrhodopsin-2 transgenic mice for optogenetic dissection of neural circuitry
function. Nature Methods. 2011 Sep; 8(9):745–752. http://www.nature.com/articles/nmeth.1668, doi:
10.1038/nmeth.1668.

Zhuang J, Ng L, Williams D, Valley M, Li Y, Garrett M, Waters J. An extended retinotopic map of mouse cortex.
eLife. 2017 Jan; 6:e18372. https://elifesciences.org/articles/18372, doi: 10.7554/eLife.18372.

777 **Figure 1–Figure supplement 1.** Intrinsic hemodynamic and GCaMP imaging to fit and test visual area maps **(A)** Schematic of hemodynamic intrinsic imaging. Upward drifting Gabor stimuli were presented to head fixed animals randomly at 6 locations in their right visual field (Materials and Methods). Each stimulus was shown to the animal 30 times. The cortical surface was evenly illuminated using a fiber coupled 530 nm LED and imaged at 2 Hz through a bandpass green emissions filter. Example behavior visual stimulus location is shown at the center of the hemodynamic map (yellow circle at 25° azimuth (az), 0° elevation (el), red dotted lines). **(B)** Averaged response map and time course of cortical reflectance change at all 6 stimulus locations. Increased cortical blood flow due to the visual stimulus results in a drop in reflectance (Δref./ref.) The period of maximal change in Δref./ref. (%) occurs 2 s after stimulus presentation (*Heimel et al., 2007*). Colored dot and boundary indicates the centroid and 75% contour. **(C)** Scaled and rotated retinotopic map of mouse V1 and secondary visual areas positioned using the 6 hemodynamic response locations in a *Ai148 x Cux2-CreERT2* animal which expresses GCaMP6f in cortical excitatory neurons. Elevation and azimuth contours from *Zhuang et al.* (*2017*). **(D)** Averaged GCaMP response to a behavior visual stimulus presented for 100 ms with 5 s of 50% grey screen between presentations. GCaMP6f fluorescence increases ($\Delta F/F$) are prominent and well delineated by the area map in V1, and the lateral and medial areas. Colored dot and boundary indicates the centroid and 50% contour. The time courses of the GCaMP6f responses shown at right are much faster than hemodynamic responses. When imaged at 2 Hz the PM response is delayed compared to V1 or the lateral areas, consistent with electrophysiology results (*Figure 5–Figure Supplement 1*.

Figure 1–Figure supplement 2. Slope determination for piecewise-linear functions were fit via least-squares regression. All spots from all areas were first fit with a variable-slope piecewise-linear function. We averaged the variable slopes of spots with low slope CI ratios to determine the fixed slope used across all spots from all areas to determine 100% crossing points (266.7 % threshold change / mW/mm^2). **(A)** Example spot with variable slope (mean 100% crossing point 0.46, bootstrap 95% CI 0.38-0.81 mW/mm^2). **(B)** Slopes of all spots with low slope CIs (< 9, within blue circle) were averaged to produce a single fixed slope. **(C)** Same example spot as (A), with fixed slope (mean 100% crossing point 0.53, bootstrap 95% CI 0.40-0.66 mW/mm^2). **(D)** Fixing the slope did not dramatically affect the 100% crossing points. Spots included here are from all visual areas.

Figure 1–Figure supplement 3. Spatial fall-of of optogenetic inhibition is restricted to illuminated areas.. We used 4 shank electrodes with wide 400 um shank spacing to measure cortical activity directly under the LED and out to a distance of 1.2 mm. **(A)** Spontaneous baseline neural activity was highly sensitive to optogenetic inhibition. The level of suppression shown is normalized to the highest intensity tested (3.0 mW/mm^2) for which baseline activity is abolished (orange dotted line at 1). **(B)** Visually evoked activity is comparatively resistant to inhibition. The level of suppression shown is normalized to control neural activity during 0 mW/mm^2 (no LED stim) trials (black dotted line at 0).

Figure 3–Figure supplement 1. Moving the visual stimulus does not affect PM results. The retinotopic map in area PM is such that stimuli at 0° elevation and 25° azimuth are at the very anterior edge of PM, while a stimulus at the same elevation but 45° azimuth is in the anterior part of PM but well within the PM boundary (*Zhuang et al., 2017*). Therefore, we trained an animal to perform the task with the stimulus at 45° azimuth. The results were similar to those with a 25° azimuth stimulus: little change in threshold when PM was inhibited, confirming that the small PM effects we found were not due to the retinotopic position of the stimulus. **(A)** Spot outlines and averaged heatmap **(B)** of hit-rate threshold changes with stimulus at 45° azimuth and at 25° azimuth. Also included is V1 data from two animals; note the V1 retinotopic representation of 25° and 45° azimuth is separated by ≈ 400 μm (*Garrett et al., 2014*; *Zhuang et al., 2017*), smaller than the V1 spots shown in (A-B), and so both V1 spots are expected to suppress responses for stimuli at both azimuths. **(C)** Moving the stimulus to a different retinotopic location and inhibiting locations within PM produced in each case a high 100% crossing point (*i.e.* optogenetic inhibition still produced little effect on behavior.) Data: mean ± bootstrapped 95% CI. Note that one PM inhibition location (with 25° azimuth stimulus) produced an effect on behavior similar to the V1 effects, but the other locations all showed high crossing points, similar to the effects of control stimulation (e.g. **Figure 3E**).

Figure 4–Figure supplement 1. False-alarm rates do not show systematic variation across areas. **(A-C)** Changes in false-alarm rates between ON and OFF trial types show little correlation with light intensity. Therefore, it would be inappropriate to predict a crossing point in the same manner as we did for hit-rate and d'. False-alarm rate analysis below is done using average false-alarm rates across all intensities. **(D)** We saw little significant difference across areas in average false-alarm rate changes (orange). To rule out that different light intensities used in different areas could contribute to changes, we computed the average intensity used in each area (blue). False alarm rate changes range between 5-10% in all four areas, with a small trend for smaller FA changes in medial and control areas. Higher average intensities were used in medial areas and control areas, as sensitivity changes were not seen at low intensities. **(E)** Average false-alarm rate map generated by averaging the false-alarm rate increase at each pixel. False-alarm rates were generally small (0-10%). Pixels with no data are black. For consequences of false alarm rate (and hit rate) variation on perceptual performance, see comparison of hit-rate and d' thresholds (Results, **Figure 1**, **Figure 4** and *Figure 4–Figure Supplement 2*).

Figure 4–Figure supplement 2. Predicting 100% crossing point using d' values does not alter behavioral effects. **(A-C)** Piecewise-linear functions were fit to d' data in the same manner described in *Figure 1–Figure Supplement 2* to predict the intensity necessary to double sensitivity (the d' 100% crossing point). **(D)** Plotting behavioral effects in terms of sensitivity (d') does not affect our main conclusion that inhibiting lateral areas (LM/AL/RL) produce effects similar to those seen in V1, while inhibiting PM produces effects similar to those seen in control areas.

783 **Figure 4–Figure supplement 3.** Changing the motor response from contralateral to ipsilateral paw produced a similar pattern of behavioral changes. Thus, observed behavioral deficits are primarily sensory, not motor, effects. **(A-B)** In all other data reported in this work, animals were trained to use the paw contralateral to the inhibited hemisphere to press and release the lever. To determine how effects would change when the motor response modality was changed, we trained one animal to use its ipsilateral paw and inhibited areas matched to those inhibited in an second animal trained to use its contralateral paw. Inhibiting similar areas in the two animals (yellow dashed lines surround spots from animal using its ipsilateral paw, blue dashed lines: contralateral paw) produced similar results. **(C)** Crossing points (with 95% CIs for the piecewise-linear functions) for the data shown in A-B. There is a trend for the ipsilateral-paw animal to have a mean crossing point larger (*i.e.* require more power to achieve a similar change in threshold) than the contralateral-paw animal for areas V1, RL, LM/AL. However, in none of these cases do the thresholds do not rise to the level of the control data. Therefore, while we cannot rule out that there may be some motor contribution to the inhibition-induced behavioral changes, changing the animals' motor response shows that the effects on performance are primarily non-motor (sensory) effects.

Figure 5–Figure supplement 1. Characteristics of V1 and secondary visual area cortical responses to visual stimulus and response timing. **(A)** Schematic of electrophysiology recordings in V1, lateral, and medial areas. Plus sign indicates visual stimulus at a mapped retinotopic location in V1. **(B)** Mean change in spike rate recorded in indicated cortical area in response to 100 ms Gabor behavior visual stim. (mean ± SEM: V1, N = 14; lateral, N = 10; medial, N = 16; t-tests: V1-lateral, n.s, $t = 0.80$, $p = 0.49$; V1-medial, *, $t = 5.13$, $p < 0.0001$; lateral-medial, *, $t = 6.57$, $p < 0.0001$). **(C)** Plot of average unit responses (visual stimulus presentation at time = 0) for indicated cortical area with sigmodial fit (solid red lines). Black bar indicates visual stimulus. Time point for half-maximal response shown for each fit with red dot.

Figure 5–Figure supplement 2. V1, lateral areas, and PM display characteristics of inhibition stabilization. **(A)** Direct optogenetic inhibition of V1, lateral areas, and PM to probe for ISN effects. Plus sign indicates visual stimulus at a mapped retinotopic location in V1. **(B)** Schematic of 4 shank probes and electrode placements superimposed on cortical area map. Individual shanks are shown as dots with color indicating cortical area for analysis. **(C)** Spike width histogram of all units across all areas recorded shows expected bi-modal distribution corresponding to wide and narrow units (E and I units respectively, N = 221). We used 450 ms as a cutoff between wide and narrow units (red dotted line). **(D)** Spike width histogram of all V1 units (N = 77). **(E)** Response of wide (E) visually responsive V1 units (normalized to r_{base}, N = 10). **(F)** Peak inhibitory effect profiles for wide and narrow waveforms (r_{min} - r_{base}) in V1 with increasing amounts of optogenetic stimulus (wide, N = 10; narrow, N = 6). **(G)** Spike width histogram of all lateral units (N = 71). **(H)** Response of wide visually responsive lateral units (normalized to r_{base}, N = 10). **(I)** Peak inhibitory effect profiles for wide and narrow waveforms in lateral areas with increasing amounts of optogenetic stimulus (wide, N = 10; narrow, N = 5). **(J)** Spike width histogram of all medial units (N = 73). **(K)** Response of wide visually responsive medial units (normalized to r_{base}, N = 16). **(L)** Peak inhibitory effect profiles for wide and narrow waveforms in PM with increasing amounts of optogenetic stimulus (wide, N = 16; narrow, N = 7).